Chloroplast DNA methylation in the kelp *Saccharina latissima* is determined by origin and influenced by cultivation

Authors: Lydia Scheschonk*[1], Anne M. Leiros Nilsen*[2], Kai Bischof[1], Alexander O. Jueterbock[2]

*shared first authorship

[1] University of Bremen, Marine Botany & MARUM, 28359 Bremen, Germany

[2] Algal and Microbial Biotechnology division, Faculty of Biosciences and Aquaculture, Nord University, Universitetsalleen 11, 8026 Bodø, Norway

Contact: alexander.juterbock@nord.no

Abstract

Chloroplast DNA is methylated in the kelp *Saccharina japonica*, in contrast to most plants. Its function is yet largely unexplored. We detected methylation in the chloroplast DNA of the congener *Saccharina latissima*, a non-model macroalgal species of high ecological (wild populations) and economical (wild and cultured populations) importance in the North Atlantic. To the functional relevance of chloroplast DNA methylation, we compared for the first time methylation patterns between wild and cultured kelp from different climatic regions (High-Arctic (79 °N) and temperate (53 °N), laboratory samples at 5 °C, 10 °C and 15 °C). Our results suggest genome-wide differences in methylated sites, and methylation level, between the climatic regions. At gene level, our data found functions related to photosynthesis to be the predominant affected case only for differential methylation between origins, but not between growth conditions. Here, sample origin led to significant differences between cultivated and wild samples due to differential methylation of genes related to DNA replication in the Spitsbergen samples. Both findings indicate that origin and cultivation

strongly affected the chloroplast methylome, but differently. Similar methylomes for samples

from the same origin – independent from whether they grow in the wild or in the lab – suggest

that origin-specific methylation marks on the chloroplast genome are inherited. However, the

capacity for rapid adaptation (to cultivation conditions) could be shown for *Saccharina*

latissima during this study. Given that DNA-methylation affects gene expression, our study

suggests that lab-cultivation alters epigenetically determined kelp chloroplast characteristics

at least to the same degree as ecotypic differentiation does.

Keywords:

Chloroplast, organelle genome methylation, epigenetics, non-model organism, cultivation,

aquaculture, marine macrophyte, marine algae

Introduction

Primary production, by which photoautotrophic organisms convert oxidised inorganic carbon

into organic carbon via photosynthesis, is the one crucial process that ensures the existence

of nearly all food webs, both terrestrial and marine. Since chloroplasts are presumed to have

developed from a free-living cyano-bacteria to an endosymbiontic organelle (Nishimura 2010),

they feature their own, albeit reduced, genome. Due to the reduced nature of their genome,

the organelles are semi-autonomous, and depend on extra-organellar products for their

internal processes (Wang et al. 2020). Epigenetic mechanisms, including histone modification,

non-coding RNA, and DNA methylation (Boquete et al. 2021), have been shown to play vital

roles in many cellular processes, such as regulation of gene expression, mRNA processing,

timing of DNA replication, determination of chromatin structure, or silencing of transposons

(Finnegan et al. 1998; Zhang et al. 2018). DNA methylation is the most stable epigenetic

mechanism that can persist for multiple generations (Lämke and Bäurle 2017), thus is

presumed to be most relevant for epigenetic adaptation. In contrast, histone modifications

and ncRNAs are often only stable for several days or weeks (Lämke and Bäurle, 2017).

However, the function of chloroplast genome methylation is yet unclear, but intensely

researched in the few species of mostly agricultural use that show chloroplast methylation,

such as tomato, rice, or the unicellular green alga Chlamydomonas (Kobayashi et al. 1990;

Nishimura et al. 1999; Muniandy et al. 2019). In marine primary producers, the first

chloroplast genome methylation has recently been documented in the brown macroalga

Saccharina japonica (Teng et al. 2021). Its functional relevance remains unknown.

Macroalgae are the predominant source of coastal marine primary production (Field et al.

1998; Filbee-Dexter 2020). Nearly exclusively restricted to the shorelines of the continents,

macroalgae act as foundation species of marine rocky shore ecosystems, in addition to

forming the base of food webs, and hence, have high and diverse ecological value (Bartsch et

al. 2008; Teagle et al. 2017; Duffy et al. 2019). Their economic value is increasing exponentially

as macroalgae cultivation gains momentum outside of Asia as an environmentally sustainable

new blue economy with a multitude of applications (Cai et al. 2021). In 2019, marine

macroalgae accounted for ~30% of the global production of marine aquaculture commodities

(120 million t), worth US$ 14.7 billion (Cai et al. 2021). However, in recent years macroalgae

distributions have shifted northward in response to increasing (summer) water temperatures

(Krause-Jensen et al. 2020; Bringloe et al. 2020). Heat waves affect both wild populations and

aquaculture yields. In kelp, the respective thermal tolerance of gameto- and sporophyte

stages seems to be the strongest factor defining their global distribution limits (Monteiro et

al. 2019; Liesner et al. 2020). Temperature is a controlling factor in many biological processes

such as photosynthetic carbon acquisition. Enzymatic activity of RubisCO, the carbon

capturing enzyme in photosynthesis, increases by a factor of 7 between 5°C and 15°C (Cen and

Sage 2005), and has been shown to be especially vulnerable to high temperatures. As the

chloroplast genome carries several photosynthesis-relevant enzymes, investigating its

methylation at different temperatures is particularly relevant when considering methylome-

based temperature adaptation in accordance to the eco-genotype approach (Scheschonk et

al. 2022). While epigenetic mechanisms have been extensively researched in terrestrial plants,

knowledge about kelp epigenetics still is limited to a handful of studies (Cock et al. 2010; Fan

et al. 2020a; Scheschonk et al. 2022), with a single study on chloroplast DNA methylation (Teng

et al. 2021). In many primary producers, the chloroplast genome can be used as control to

estimate the rate of false positive methylation calls, as the chloroplast genome commonly has

been presumed to be un-methylated, despite data suggesting otherwise (*Saccharina japonica*,

Teng et al 2021; Chlamydomonas reinhardtii, Nishimura 2010; Nishimura et al. 2002;

Nishimura et al. 1999; Nishiyama et al. 2004; Nishiyama et al. 2002; Umen and Goodenough

2001). Epigenetic mechanisms hold the key to rapid local adaptation, and hence the capacity

for species survival and ecosystem stabilisation despite swift alterations in biotic and abiotic

factors. Especially in the face of the soaring changes in local and global ecosystems due to the

anthropogenic climate crisis, gaining knowledge in this field is crucial. Our study is the first

attempt to determine the influence of temperature, origin, and growth condition (wild *vs*

cultivated) on the methylome of chloroplasts of the kelp *Saccharina latissima* (sugar kelp). In

the northern hemisphere, this boreal-temperate marine brown alga, a congener species to *S.*

japonica with a large latitudinal distribution range, is among the macroalgal species of both

high ecological and economic value (Bartsch et al. 2008; Cai et al. 2021). In this fundamental

research study, we tested the following hypotheses:

1) *Saccharina latissima* features chloroplast genome methylations, as has been reported for

S. japonica (Teng et al 2021).

2) The chloroplast methylome mirrors the nuclear methylome (Scheschonk et al. 2022) in

showing strongest epigenetic differentiation between origins, with modifications specific to

temperature and lab cultivation.

Results

Methylation levels and methylated sites

According to *in-silico* digestion, there were 69 potential MethylRAD sites for CG methylation,

and 201 for CHG methylation in the chloroplast genome of *Saccharina latissima*. 20.7% and

24.8% of these sites were methylated, respectively (Table 1).

Table *1:* Potentially methylated sites for each sequence context, the mean number of methylated sites
per sample and standard deviations for the means. Methylation level is the mean percentage of
potential methylation sites methylated in samples, shown in the table with standard deviations (SD).

Sequence Context	Potential Sites (in-silico)	Methylated sites (mean +/- SD)	Methylation level (mean +/- SD)
CG	69	14.3 ± 1.5	20.7% ± 2.2%
CHG	201	49.8 ± 3.0	24.8% ± 1.5%

The number of methylated CG sites was significantly higher in Helgoland than in Spitsbergen

samples both in laboratory samples ($p < 0.001$, Fig. 1 A) and in laboratory and field samples

combined ($p < 0.0001$, Fig. 1 B), but no significant differences were found between treatment

temperatures or between field and laboratory conditions for sites in CHG contexts (Suppl. Fig.

1 C, D).

Methylation levels normalized to reads per million (RPM) showed no significant differences in

CG contexts between any groups (Suppl. Fig. 1 A, B), but the Spitsbergen population showed

higher methylation levels than the Helgoland population in the laboratory samples (p < 0.05,

Fig. 1 C). There were no significant differences between field and lab conditions in CHG context

(Fig. 1 D).

Figure *1*: **Significant differences in methylated sites and methylation level between conditions.**
Methylated CG sites (A, B), and methylation levels as reads per million (RPM) for CHG sites (C, D)
between populations and temperatures in laboratory samples (A, C) and compared between lab and
field samples (B, D). ns: not significant. Numbers at the bottom of the bars indicate sample size.
Significance codes after p-value correction (Benjamini-Hochberg) for multiple comparisons: '< 0.05': *,
'< 0.01': **, '< 0.001': ***, '< 0.0001': ****.

 Principal Component Analysis

Laboratory and field samples formed distinct groups along the first dimension with 95%

confidence ellipses not overlapping when both origins were combined (Fig. 2 A). The first two

principal components equally contributed about 14 % of the total sample variation, which was

similar between the comparisons of 'lab vs field' (Fig. 2 A) and 'Helgoland vs Spitsbergen' (Fig.

2 B). However, for the origins (Fig. 2 B), the populations of Helgoland and Spitsbergen formed

distinct groups along the second dimension when lab and field samples were combined (Fig.

2 B), while when field and laboratory samples were separately analysed for origin, Helgoland

and Spitsbergen samples again split along the first axes (Fig. 2 C, D).

Temperatures in the laboratory samples (5°C, 10°C, 15°C) did not separate into distinct groups

(Fig. 2 E).

Figure *2*: **Principal Component Analysis for A: all samples, grouped by laboratory or field samples. B:**
laboratory samples grouped by origin. C: all samples, grouped by origin. D: field samples grouped by
origin. E: laboratory samples of both origins grouped by temperature treatment. Ellipses represent
95% confidence intervals around the group medians.

Differentially methylated sites and GO terms
Comparing the origins (Helgoland vs Spitsbergen, Fig. 3), differentially methylated sites were

as often higher methylated in Spitsbergen samples as in Helgoland samples (Suppl.Tab.1-3).

Of the eight genes with significant methylation differences between the origins (Fig. 3), only

three genes (psaB_(1), atpA, yfc3) were identical to the genes with significant methylation

differences between the origins at a respective temperature within laboratory samples (Table

2, see colour coding suppl. Table 5+7, sheets 'ST5 Origin' and 'ST7 Temperature').

Figure *3:* **VennEuler diagram showing the numbers of differentially methylated sites between**
Helgoland and Spitsbergen when analysing field samples (blue), lab samples (orange), and all
(lab+field, grey) samples (suppl. Tab. 1-3). The labels give the product of the genes that were
differentially methylated in the respective partitions. _(1) and _(2) depict a gene that showed
differential methylation at two different sites. For the exact site position in the respective gene, see
suppl. Table 5 ('ST5 Origins').

Of the 69 genes that were analysed comparing growth conditions (Fig. 4), 16 genes were

differentially methylated between laboratory and field samples, two of which were also

differentially methylated between origins (Fig. 3, Table 2). Many of those genes were

differentially methylated at two different positions (suppl. Table 6, sheet 'ST6 Cultivation').

Figure *4*: **VennEuler diagram showing the numbers of differentially methylated sites between lab**
and field samples when analysing Helgoland samples (yellow), Spitsbergen samples (green), and the
samples from both origins together (**grey**, suppl. Tab.4)**.** The labels give the product of the genes that
were differentially methylated in the respective partitions, with one gene without annotation record
at the ncbi database. _(1) and _(2) refer to a gene that showed differential methylation at two different
positions. For the exact position in the respective gene, see suppl. Table 6 ('ST6 Cultivation').

Two genes were differentially methylated between temperature treatments (Table 2), with

'acetolactate synthase large subunit' (QKE47546.1) at two comparisons: 5°C to 10 °C (padj <

0.0001), and 5 °C to 15°C (padj < 0.0017). All other significant differentially methylated genes

were found between origins at different temperatures, but only partially overlapped with

those obtained when analysing 'all samples' (Fig. 3, see colour codes in suppl. Tables 5 + 7

(sheets 'ST5 Origins' + 'ST7 Temperature')).

Table *2*: **All conditions that were found to be differentially methylated by the DESeq2 analysis when**
analysing the different rearing temperatures of the laboratory samples. Most significances were
detected between the origins at respective temperature (Helgoland vs Spitsbergen @ T5/10/15), but

only 3 of those were identical to those detected when all lab samples were being analysed as a group
(Fig. 3, Suppl. Table 7 ('ST7 Temperature')). However, genes QKE47546.1 and QKE47431.1 (bold letters)
showed significantly different methylation when analysing for temperature difference as factor (origin,
T vs T).

Comparison	Gene Product	Gene ID
5° vs 10° Helgoland	**acetolactate synthase large subunit**	**QKE47546.1**
5° vs 15° Helgoland	**acetolactate synthase large subunit**	**QKE47546.1**
5° vs 10° Spitsbergen	**photochlorophyllide reductase subunit B**	**QKE47431.1**
5° Helgoland vs Spitsbergen	photosystem I P700 apoprotein A2 _(1)	QKE47459.1
	ATP synthase CF1 alpha chain	QKE47418.1
	acetolactate synthase large subunit	QKE47546.1
	photochlorophyllide reductase subunit B	QKE47431.1
10° Helgoland vs Spitsbergen	photosystem I P700 apoprotein A2 _(1)	QKE47459.1
	ATP synthase CF1 alpha chain	QKE47546.1
	photosystem I assembly protein	QKE47473.1
	photochlorophyllide reductase subunit B	QKE47431.1
15° Helgoland vs Spitsbergen	photosystem I P700 apoprotein A2 _(1)	QKE47459.1
	ATP synthase CF1 alpha chain	QKE47546.1

193 None of the extracted GOterms were significantly enriched according to the 'topGO'

194 enrichment analysis.

Analysis of Epigenetic Distance

In both comparisons ('cultivation', 'origin'), the epigenetic distances between the samples

within respective groups were significantly different between groups (Fig. 5). For 'cultivation',

distances were significantly higher in laboratory samples than in field samples ($p < 0.0001$,

median field 153, lab 210), while for 'origin' the samples obtained from Spitsbergen showed

significantly higher epi-distances than Helgoland ($p < 0.0001$, median Helgoland 171,

Spitsbergen 190; suppl. Table 8).

Fig. 5: **Epigenetic distance (Euclidean distance) for 'cultivation', Laboratory (orange) and Field (blue) samples (A), and for 'origin', Helgoland (yellow) and Spitsbergen (green) samples (B).** The small black dots depict the pairwise distances between the individual samples of each group (from the PCA, Fig. 2). The light grey dots are outliers detected by the boxplot script, but were included during epi-distance analyses. The boxes represent the interquartile range (IQR) of the respective group, and the horizontal line within the boxes gives the median Euclidean distance.

Discussion

Like recently reported for the kelp *Saccharina japonica* (Teng et al 2021), we detected

methylations in the chloroplast DNA of its congener *S. latissima* (Table 1). The chloroplast

methylome significantly differed between origins (Fig. 3, Table 2, Fig. 5 B), and growth

conditions (Fig. 4, Fig. 5 A), but temperature had a lesser impact than expected (Table 2).

Differentially methylated sites between origins (Spitsbergen or Helgoland) were located in

genes with functions related to photosynthesis. In contrast, differentially methylated sites

between growth conditions (laboratory or field) were almost exclusively located in genes

related to DNA replication and cell division (suppl. Table 5 + 6, sheets 'ST5 Origin' + 'ST6

Cultivation'; Fig. 3, Fig. 4). This suggests that the chloroplast methylome is origin-specific,

indicating an epigenetic memory, while showing capacity of rapid adaptation to shifts in

growth conditions via the epigenetic mechanism of cytosine methylation.

Sample origin

Population origin had a strong impact on the methylome according to PCA clusters (Fig. 2 B,

C, D), which was reflected in methylation levels, and methylated sites which significantly

differed between origins (Fig 1 A, C). Photosystem I (PSI) was the component most affected

by differences between the origins, as genes coding for varying aspects crucial to PSI

significantly differed at a total of four sites in three genes (Fig. 3, Table 2), with the strongest

effect on PSI P700 chlorophyll a apoprotein A2. In addition, the gene coding for apoprotein A2

in addition was one of two genes differentially methylated between the origins in all analysed

temperature conditions of the laboratory samples (Table 2, suppl. Table 7 sheet 'ST7

Temperature'). Therefore, the difference can be assumed to be solely accounted for by origin.

While methylation levels were significantly different only between the origins (Fig. 1), the

differences appear to be independent of the rearing condition (laboratory or wild), as most of

the differentially methylated genes between origins overlapped in field and lab samples (Fig.

3). Furthermore, the PCA indicated that origin plays a more pronounced role when analysing

laboratory and field samples separately (Fig. 2 C, D), while it is less pronounced when both

growth conditions were combined (Fig. 2 B). Hence, part of the methylome seems to be origin-

specific even after years of lab-cultivation. The differentially methylated genes were

exclusively related to functions of photosynthesis (Fig. 3), which likely can be explained by the

latitudinal differences in the light regimes. Light conditions in the High-Arctic are strongly

seasonal, with absence of photosynthetically active radiation (PAR) for at least three months

during the arctic winter (polar night), and continuous PAR during the arctic summer (polar

day). In boreal-temperate regions, PAR is available throughout the year. Whether or not the

differences observed here are of genetic nature cannot be assessed by our setup, but the

prevalence of the significant difference, detectable in the epigenetic signature that prevailed

through cultivation between origins, is a strong indicator for the eco-genotype approach

(Scheschonk et al 2022). This presumes that adaptation to a specific location is not solely a

matter of genetic nature (fixed only in the DNA), but that epigenetic mechanisms, especially

inheritable ones such as cytosine methylation, play an important role in local adaptation.

Considering methylation and gene expression were shown to be negatively correlated in the

S. japonica nuclear and chloroplast genes (Fan et al. 2020a; Teng et al. 2021) the (double-)

methylation of the gene coding for apoprotein A2 (Fig. 3, suppl. Table 5 sheet 'ST5 Origin'), for

example, which is one of the crucial components of PSI, can be presumed to have a direct

impact on the photosynthetic capacity which according to our results is likely to differ

between origins. Like with the nuclear genome (Scheschonk et al 2022) this has crucial

implications for cultivation processes as well as restoration efforts, as it again shows the

importance of suitable origin when considering local growth condition.

Cultivation effects
PCA showed that cultivation affected the methylome as strongly as origin (2 A, C). While lab

and field samples showed no significant differences in genome-wide methylation level and

number of methylated sites (Fig. 1), they displayed significantly different methylation levels

at 23.19 % of the sites (Fig. 4, suppl. Table 6 (sheet 'ST6 Cultivation'). Most (12 out of 16) of

the sites differentially methylated between rearing conditions had higher methylation levels

in the lab than in field samples (Suppl. Tab. 1). This suggests that those genes were higher

expressed in the field samples (Fan et al. 2020a; Teng et al. 2021). Unexpectedly, epigenetic

distance was higher in lab samples, even though they experience more uniform environmental

conditions than field samples. When translated to transcription, a higher epigenetic distance

is likely to result in more variable phenotypes. This might suggest that the kelp methylome

diversifies rather under stable lab conditions than under variable field conditions, at least

when said cultivation conditions were not previously encountered. Potential explanatory

factors cover the whole process of cultivation, from obtaining the sori, and securing the

zoospores, or handling during the fertilisation process, to culture conditions at any stage.

The lab cultures, which were grown from gametophytes that had been kept in culture for

about four years prior to the experiment (from 2014/2015 to 2018), experienced stable abiotic

factors such as nutrient availability, water current, light/dark cycle, and (before the

experimental treatments) constant temperature, and were protected from biotic stressors

such as grazing, pathogens, and competitors. Thus, cultivated / hatchery kelp might not have

activated those genes required to acclimate to environmental stressors encountered in the

wild due to (gametophyte) culture conditions, which might affect their performance when

deployed at sea for aquaculture or restoration purposes.

Nevertheless, origin had the strongest influence on the differences in cultivated samples at

gene level, as lab-field differences in methylation could only be detected in the Spitsbergen

population (Fig. 4, Table 2).

Another factor that might have impacted the methylomes of lab cultivated and wild kelp is

the age of the sporophytes. The field sporophytes were adult, while the lab samples were

juveniles with a size of approximately 12-15 cm. In plants, cytosine methylation levels tend to

increase with age, but these differences are most pronounced in meristematic areas

(Dubrovina and Kiselev 2016), which were avoided in the kelp samples. Because meristematic

regions were omitted, and there was no pattern consistent with older individuals (field

samples) having higher methylation levels in the chloroplast genome, the likelihood that the

results were affected by sporophyte age differences is low. Temperature as factor could be

shown to have an origin-specific impact on the methylome during cultivation (Table 2; see

next paragraph).   However, which step, or factor, or combination thereof, during the

cultivation process is likely to have caused the increase in epigenetic distance will have to be

assessed in future studies. The difference in methylation between origins that persisted under

laboratory conditions might be due to epigenetic inheritance of a molecular memory of the

environment.

Whether or not genetic differences can result in the epigenetic differences that we observed

has yet to be determined. In plants and kelp, methylation can be present in CG, CHG, and CHH

sequence contexts, where H is any of the bases A, C, or T (Richards and Pigliucci 2020).

Cytosine methylation can be heritable across several generations (Lämke and Bäurle 2017),

hence is a likely candidate for rapid non-genetic but heritable local adaptation. The inheritance

of epigenetic traits induced by environmental factors has been repeatedly reported (Foust et

al. 2016; Gugger et al. 2016; Keller et al. 2016). This suggests that methylation differences can

remain stable across generations in cultivation, indicating that a transgenerational epigenetic

memory likely is present in the *S. latissima* chloroplast genome.

Temperature and the methylome
The temperature treatments did not form distinct clusters in the PCA. The cultivated *S.*

*latissima*s' chloroplast methylome, unlike the nuclear methylome, did not differ in

methylation levels and numbers of methylated sites in response to the rearing temperatures

of 5 °C, 10 °C and 15 °C. Significant differences solely caused by temperature were detected

in 2.9 % of the genes (Table 2). However, when analysing the origins at temperature level

(Table 2), most differences between origins found at discrete temperatures were in other

genes or at other gene sites than those detected for the origin analysis. This again suggests

the inheritance of an epigenetic memory in accordance with the eco-genotype approach

(Scheschonk et al. 2022), as the methylomes reacted origin-specific at discrete temperatures,

and it cannot be ruled out whether these epigenetic differences have a genetic base.

Increasing methylation levels have been shown to be positively correlated with heat stress

(Boyko et al., 2010; Correia et al., 2013). Heat stress is expected to excite a reaction in the

chloroplast due to the high temperature sensitivity of photosynthesis, but 15 °C, the highest

cultivation temperature, cannot be considered as 'heat', neither for chloroplast enzymes, nor

for *S. latissima* as organism. This temperature might have elicited a stress response in the

Spitsbergen samples, due to its presumed lack of epigenetic memory for this temperature. If

this were the case, however, the origins' response to 15°C during cultivation should have been

much stronger (more sites/ significant difference in methylation level) than to the lower

temperatures. However, 15°C was actually the temperature that had the least response in

terms of differential methylation (Table 2), hence this should not be considered the reason

here. A possible explanation for the overall low amount of (differentially) methylated sites

might be that the temperature exposure of the laboratory samples was not stressful for the

physiological processes regulated by the chloroplast genome. However, a reason for the

similarity of chloroplast DNA methylation at the different temperatures might be that

methylation sites affected by these temperatures are probably predominantly located in the

nucleus. In *Saccharina latissima*, it has been shown that temperature treatments affect the

nuclear methylome, and that differences in the methylome exist between populations from

different latitudes, suggesting that epigenetic modifications contribute to local adaptation

(Scheschonk et al. 2022). In addition, significant differences in the nuclear methylome were

found between wild and cultivated individuals (Scheschonk et al. 2022). Earlier research on *S.*

latissima suggests that short-term acclimation to different growth temperatures can lead to

changes in heat tolerance and affect photosynthetic performance (Davison et al. 1991).

Temperatures above approximately 20 °C generally result in poor growth, but at temperatures

below 15°C *S. latissima* shows a high capacity for acclimation (Andersen et al. 2013; Diehl and

Bischof 2021). Site-specific differences in circannual temperature may have contributed to the

different methylation patterns of populations from the high Arctic (Spitsbergen) and

temperate (Helgoland) regions. Here, heat stress may have occurred in the Helgoland

population during previous generations, as summer water temperatures can approach the

upper thermal tolerance of *S. latissima* (Bolton and Lüning 1982; Lüning 1984; Andersen et al.

2013). However, of the genes differentially methylated between origins, most were found at

discrete temperatures (5/10/15 °C), while not detectable when comparing the origins over all

lab samples.

Methylation levels compared with the nuclear genome
The methylation levels of sites in the chloroplast genome were ca. 100 times higher than levels

in the nuclear genome as documented by Scheschonk et al. (2022). In the nuclear genome,

the mean methylation level was 0.21% ± 0.13% SD (Scheschonk et al. 2022), while in the

chloroplast, the levels were 20.7% ± 2.2% for CH contexts and 24.8% ± 1.5% for CHG contexts

(Table 1). The higher methylation level in the chloroplast can be explained by the low numbers

of potential methylation sites in the chloroplast genome (Table 1). In comparison, in

Saccharina japonica the methylation level was reported as 1.4% in the nuclear genome (Fan

et al. 2020a) and in chloroplasts 17.1% at CG sites and 58.1% at CHG sites (Teng et al. 2021) of

sporophytes. Thus, the pattern of higher methylation in the chloroplast than the nuclear

genome is repeated in both *S. japonica* and *S. latissima*, with some dissimilarities that could

be caused by different sequencing methods and the inclusion of gametophytes in the studies

of *S. japonica* (Fan et al. 2020a; Teng et al. 2021). An example from land plants is in *Sequoia*

sempervirens, which has more similar methylation levels in its nuclear genome (7.0% - 8.6%)

and chloroplast genome (7.1% -7.6%), so in this plant the pattern of higher methylation in the

chloroplast is not present (Huang et al. 2012).

In the *S. latissima* nuclear genome, methylation was more site-specific in the population from

lower latitudes: the level of CG methylation was consistently higher, but the number of

methylated sites was lower in temperate Helgoland than in Arctic Spitsbergen (Scheschonk et

al. 2022). Such pattern of methylation becoming more site-specific has also been observed in

cyanobacteria in response to stress exposure (Hu et al. 2018). In contrast, in this study, the

chloroplast genome showed more methylated sites in Helgoland than in Spitsbergen and

higher methylation levels of CHG sites in Spitsbergen. Thus, the population specific pattern in

the nuclear methylome was not repeated in the chloroplast methylome.

In the chloroplast genome, there were no differences in methylated sites or methylation levels

between temperature treatments or between laboratory and field samples. In contrast, in the

nuclear genome, a response in methylation levels and numbers of methylated sites to warmer

temperature treatments was observed, especially in the Helgoland samples (Scheschonk et al.

2022). This difference suggests that the chloroplast genome responds differently to rising

temperatures than the nuclear genome, even though photosynthesis is highly sensitive to

temperature (Cen and Sage 2005).

However, the lack of statistically significant differences in the chloroplast genome as

compared with the nuclear genome may partly be explained by the smaller number of genes.

The chloroplast genome has a size of 130 kb and contains 139 protein-coding genes, three

rRNA genes and 29 tRNA genes (Fan et al. 2020b), while the nuclear genome has a size of 545

Mb and contains 18,733 protein-coding genes (Ye et al. 2015).

In the nuclear genome, most of the differentially methylated sites between lab and field

samples and between populations were in non-genic repetitive contexts (transposable

elements). In contrast, in the chloroplast genome, the differentially methylated sites were in

coding regions of genes, except for very few sites in a promoter region (Suppl. Tab. 1-4). The

higher number of methylated sites in coding regions in the chloroplast are likely related to

differences between chloroplast and nuclear genomes: chloroplast genomes are more

condensed and have far fewer noncoding regions than nuclear DNA (Clegg et al. 1994).

Comparison with the congener species Saccharina japonica
The number of methylated sites found in the chloroplast genome of *S. japonica* (4894 CG and

5807 CHG sites, Teng et al. 2021) was >20 times higher than in chloroplasts of *S. latissima*

(Suppl.Fig.7; 69 CG sites and 201 CHG sites, Table 1) identified in this study.

The low number of methylation sites is an expected outcome of the MethylRAD method. The

restriction enzyme used for digestion targets CCGG and CCHGG sites, which reduces the

number of potentially methylated sites compared to CG/CHG, and in addition targets the

specific bases within the degenerate adaptor endings 5'-NNNT-3' and 5'-NNNC-3', which only

constitute about one-eighth of all CG and CHG sites (Wang et al. 2015), while it does not target

CHH sites. CHH contexts were found to be the highest proportion in *S. japonica* based on

whole-genome bisulfite sequencing ('WGBS'; Fan et al. 2020a; Teng et al. 2021). Hence, the

methylation levels of the *S. latissima* chloroplast genome are expected to be much higher than

the mean number of covered sites we obtained, and are likely to be similar to that of *S.*

japonica when analysed with the same sequencing method (WGBS). However, as we found

both origin and cultivation to significantly influence differences in the methylome within the

same species, this should be taken into account for inter-species study comparisons as well.

Further, the life stage of the analysed samples plays a major role for methylation levels. In

both the nuclear, and chloroplast genome *Saccharina japonica* gametophytes were shown to

feature more methylation than sporophytes, and there is evidence suggesting that

methylation plays a role in regulating the kelp's life cycle (Fan et al. 2020a; Teng et al. 2021;

Yang et al. 2021).

Conclusion

Our results showed that sample origin and growth condition can strongly influence epigenetic mechanisms, while temperature had a minor effect. As methylation differences in genes are likely to trigger gene expression differences (Richards and Pigliucci 2020), this would directly relate to eco-phenotype. The scale-up in cultivation of *Saccharina latissima* during the last decade is likely to continue, and cultivation sites overall will likely shift more northward, in accordance to the preferences of wild populations. The capacity of fast adaptation to changed environmental factors, facilitated by epigenetic mechanisms, could prove particularly useful in the face of rapid anthropogenic climate change. However, due to the high ecological and economic value of this kelp, the implication of differences in methylation due to cultivation relative to origin, and especially the possibility of chloroplast methylation influencing photosynthetic capacity need to be further assessed. As most genes that were differentially methylated in the chloroplast genome between origins contribute to aspects of photosynthesis, an urgent question arising from our findings is whether differences in chloroplast methylation impact photosynthetic capacity. Photosynthetic capacity is directly linked to primary production, and the capability to build up and maintain biomass. This would have implications for both cultivated and wild populations. In cultivars, the importance of matching hatchery and cultivation conditions to origin has been implied by our results. Furthermore, due to the significant differences observed in the methylomes of cultivated *S. latissima*, it might be possible to detect an epigenetic 'fingerprint' of cultivated specimen in the future, to detect escapees to the wild. Regarding the current trend in northward shift, the further north populations of *Saccharina latissima* settle, the more pressure the months of darkness during polar night will pose (Scheschonk et al. 2019). Photosynthetic pigments were shown to be functional throughout the prolonged darkness of the Arctic winter (Scheschonk et al 2019), indicating the chloroplasts of High Arctic specimen retained photosynthetic

capacity during polar night. In which context methylation of chloroplast DNA facilitated by

differences in the eco-phenotypes might have played a role in this context will be of interest,

especially regarding populations adapted to light conditions at more southern distributional

habitats, shifting northwards. In *Saccharina latissima*, phenotypic variation among

populations is often driven by phenotypic plasticity, and the species shows high plasticity in

response to changing abiotic factors (Bolton & Lüning 1982; Diehl and Bischof 2021). So far,

research on differences between origins in *S. latissima* has focused on genetic differentiation,

with significant genetic variation between populations (Møller Nielsen et al. 2016; Paulino et

al. 2016; Guzinski et al. 2020). Epigenetic differences have largely been unexplored as an

alternative source of molecular variation between *S. latissima* populations. The term 'eco-

phenotypes' has recently been introduced to include both epigenetic and genetic mechanisms

into the concept of heritable molecular variation that contributes to adaptation to local

environments and intraspecific variation between origins (Scheschonk et al. 2022). The

differences we observed between origins within the laboratory and the field samples indicate

that the eco-phenotype approach should be applied to the organellar chloroplast genome as

well. Furthermore, our data suggests that the influence of cultivation seems to be at least

partially origin-specific, and that the influence of cultivation and origin on the methylome are

interconnected. Epigenetic modifications have been found to be under some degree of

genetic control, but can also be triggered by environmental factors such as temperature

changes (Boyko et al. 2010; Correia et al. 2013). Hence, the role of genetic differences

controlling the epigenetic responses reported here needs to be assessed in further detail in

future studies.

Materials and Methods

Laboratory cultures, field samples, and 'DNA- Extraction and MethlylRAD sequencing are described in more detail in Scheschonk et al (2022). Changes and specific adjustments to the present study were necessary during the bioinformatics and statistics part, and are detailed below ('Bioinformatics and Statistics').

Laboratory cultures

As described in more detail in Scheschonk et al (2022), four cultures per origin (S1-S4, H1-H4; technical replicates) were obtained from mono-parentally fertilised clonal gametophyte cultures of *Saccharina latissima* (Linnaeus) C.E. Lane, C. Mayes, Druehl & G.W. Saunders from Helgoland, German Bight (Nordstrand, 54°11'18.9"N 7°54'14.1"E; HG1, HG2, HG3, HG4 in culture since 2014) and Spitsbergen, Svalbard (Hansneset, Kongsfjorden, 78°59'26.0"N 11°58'42.3"E; SG1, SG2, SG3, SG4 in culture since 2015; see table Suppl. Table 9) were kept at each of the three temperatures (5 °C, 10 °C, and 15 °C) under white light at 15 μmol m-2 s-1 (18 : 6 hours light : dark) from as early as initiation of fertilization. Only sporophytes with a fresh weight exceeding 120 mg (without stipe) were taken for DNA extraction.

Field Samples

Adult sporophytes were sampled from Helgoland, German Bight (54°11'18.9"N 7°54'14.1"E), and from Spitsbergen, Svalbard (78°59'26.0"N 11°58'42.3"E) at the same locations from which the gametophytes for the laboratory grown sporophytes originated. From each sporophyte, 4 discs (ø=3 cm) were cut from the fronds, omitting the meristematic region, 'midsection', and reproductive tissues. Samples from lab and field were dried in cellulose bags stored in silica. In total 21 samples from the laboratory cultures, and 20 samples from the wild were analysed. In contrast to the field samples, our laboratory samples were uniparentally fertilised. However, given that chloroplast genes are transmitted solely from the maternal parent (Kuroiwa 1991, in Nishimura 2010), the chloroplast genome inheritance is identical in

uniparentally and bi-parentally fertilised individuals. This circumstance enables a direct

comparison between the chloroplast methylomes of our cultured and wild samples.

Bioinformatics and Statistics of MethylRAD sequencing

Clean-up of raw data

DNA was extracted and MethylRAD sequencing libraries were prepared as described in detail

in Scheschonk et al (2022). The sequences were de-multiplexed by sample and quality

trimmed with TrimGalore! v 0.4.1

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Bases with Phred-

score <20 (--quality 20) were eliminated, the adapter sequence was removed (--stringency 3),

and the terminal 2bp (both ends) were removed to eliminate any artefacts at the ligation

position (--clip_R1 2 \ --three_prime_clip_R1 2). Reads were checked for fragment (over-)

representation, base AT CG content bias, and read length with FastQC v0.11.8 before and after

quality trimming.

In-silico digestion

For an overview of potential methylation sites, and as a backbone to map sequences to the

reference chloroplast genome for the *S. latissima* chloroplast (Fan et al. 2020b) was digested

in silico using the custom python script InSilicoTypeIIbDigestion_corrected.py

(http://marinetics.org/2017/04/11/REdigestions.html) and settings for simulating the

restriction enzyme FspEI. The *in silico* digestion found 270 recognition sites, where 69 were

CCGG recognition sites, 61 CCTGG sites, 77 CCTGT sites, and 63 CCAGG sites.

Mapping and annotation

The sequenced reads were mapped to the *in silico* digested reference genome of the *S.*

latissima chloroplast, combined with the *in silico* digested genome of the closely related

species *Saccharina japonica* to account for reads belonging to the nuclear genome and avoid

false mappings to the chloroplast genome for those reads. Reads were mapped using Burrows-

Wheeler Aligner (BWA Version: 0.7.17-r1188) (Li and Durbin 2010). Duplicate mappings were

excluded from further analyses by filtering out reads with a mapping quality <10 using

samtools (v 1.9). A count table was created listing the number of reads that mapped back to

each methylation site. For this, htseq-count (v 0.7.2) was called on each sample alignment

with *in silico* digested fragment in gff3 format as count features. An additional table was made

with counts normalized to 'reads per million (RPM) = (Reads mapped per site ×10^6)/(Total

number of mapped reads)' to account for varying sequencing depths.

Only methylated sites that were covered in at least one library with coverage >2 were kept in

count tables and RPM tables. The sites were annotated for their sequence context (CG or

CHG), genomic region, gene ID numbers, and gene functions based on the gff3 annotation file

for the *S. latissima* chloroplast genome from NCBI (Accession: MT151382.1) (Fan et al. 2020b).

Statistics
Principal Component Analysis
Differences in the chloroplast methylome were investigated with principal component

analysis (PCA) on RPM tables using the R package 'FactomineR' (Lê et al. 2008). Since lab and

field samples separated clearly into distinct groups in the PCA, their methylomes were

compared first for all samples, and then separated into origins (Helgoland and Spitsbergen).

Within the lab and field samples, and for all samples, we compared origins (Helgoland and

Spitsbergen). Within the lab samples, we compared between the treatment temperatures of

5°C, 10°C, and 15°C.

Outlier detection
The outlier function in the R package 'FactoInvestigate' (Thuleau & Husson, 2020) detected

two outliers, FH1 and FH9 (Helgoland field samples) in the PCA. The outliers showed few reads

mapping to the genome, few methylated sites, and high RPM values for some of the

methylated sites (Suppl. Fig. 2). Thus, FH1 and FH9 samples were excluded from all follow-up

analyses. No outliers were found in the Spitsbergen samples. As both outliers belonged to the

same group (Helgoland field samples), outlier removal led to a reduction in variance explained

by the first two principal components from 69.9% to 28.9% (Suppl. Figs. 2, 4, 5).

Methylation levels

We tested for differences in DNA methylation levels as RPM and for differences in the

numbers of methylated sites between populations, between treatment temperatures, and

between lab and field samples, using Wilcoxon rank sum tests. P-values were corrected with

the Benjamini-Hochberg method (Benjamini and Hochberg 1995) to control for the false

discovery rate, and results with adjusted p values padj <0.05 were considered significant. The

tests were performed using the R package 'rstatix' v0.7.0 (Kassambara, 2021).

We tested for correlation between the numbers of quality trimmed reads, and the numbers

of quality-trimmed reads that mapped back to the *Saccharina latissima* chloroplast genome

(Suppl. Fig. 4C, 5C, 4D, 5D).

Differential methylation analysis

We tested for significant differential methylation with the R package 'DESeq2' (Love et al.

2014) between 'Helgoland' *vs* 'Spitsbergen' in all samples ('All', or 'both locations'), and in only

field samples and only lab samples. Further, between 'field' *vs* 'lab' in all samples, and in only

Helgoland samples, and only Spitsbergen samples. We also compared pairwise between

temperature treatments: both origins combined (Helgoland + Spitsbergen 5 °C *vs* 10 °C; 5 °C

vs 15 °C; 10 °C *vs* 15 °C), separately for Helgoland and Spitsbergen (5 °C *vs* 10 °C; 5 °C *vs* 15 °C;

10 °C *vs* 15 °C), and for each temperature (Helgoland vs Spitsbergen @ 5 °C; 10 °C; 15 °C).

'DESeq2' normalised raw methylation counts before testing for differential methylation, and

corrected p-values with the Benjamini-Hochberg method for controlling false discovery rates

(Benjamini & Hochberg, 1995). Methylated sites with adjusted p-value padj < 0.05 were

considered significantly differentially methylated.

GO term analysis

Gene ontology (GO) terms for biological processes, cellular components, and molecular functions were extracted from the database https://www.uniprot.org/ using Gene ID numbers (Suppl. Tab. 1-4). The R package 'topGO' (Alexa & Rahnenführer, 2009) was used to test for enrichment of the GO terms.

Gene names and functions

Names and functions of differentially methylated genes were extracted from the chloroplast genome of *Saccharina latissima* published on NCBI (strain y-c14; https://www.ncbi.nlm.nih.gov/nuccore/MT151382.1?report=graph).

Analysis of epigenetic distance

Groups of the aspects 'cultivation' (lab vs field), and 'origin' (Helgoland vs Spitsbergen) were analysed for differences in epigenetic distances among the samples of each group. The Euclidean distances from the PCA including all samples were used for this. The Shapiro-Wilk indicated non-normal distribution for groups analysed for cultivation (p-value lab $p < 0.0001$, field $p < 0.0001$) and for origin (Helgoland $p < 0.0001$, Spitsbergen $p < 0.0001$), hence both tests for epigenetic distance were carried out applying the non-parametric Wilcox-rank-sum test (suppl. Table 8).

Data availability

Raw sequencer data of this study are available at NCBI SRA BioProject PRJNA809008.

Acknowledgements

We thank S. Niedzwiedz (University of Bremen, Germany) for securing the field samples, and G. Hoarau and H. Reiss (Nord University, Bodø, Norway) for providing lab space and sequencer. Thank you, M. Kopp (Nord University, Bodø, Norway), for the support during the sequencing procedure, and for preparing some of the libraries. Thanks to I. Bartsch and A.

590   Wagner (Alfred Wegener Institute for Marine and Polar Research, Bremerhaven, Germany)

591   for providing and hatching the sporophyte cultures.

Funding

593   LS acknowledges funding by Deutscher Akademischer Auslandsdienst (DAAD; research travel

594   grant 'BremenIdeaOut' 2019). AN acknowledges funding by Nord University (Master Thesis

595   and transitional stipend). AJ acknowledges funding by Nord University (research talents

596   grant). Parts of this work have been submitted in partial fulfilment of the requirements for

597   Master of Science, at Nord University, Bodo (AN), and Dr. rer. nat, University of Bremen (LS).

Author Contributions

599   LS designed and conducted the experiment, LS and MK extracted and sequenced the samples.

600   AN, LS and AJ analysed and interpreted the data. LS and AN wrote the manuscript. All authors

601   read and commented on previous versions of the manuscript, and approved the final version.

References

Alexa, A., & Rahnenführer, J. (2009). Gene set enrichment analysis with topGO. Bioconductor

Improv, 27, 1-26.

Andersen GS, Pedersen MF, Nielsen SL (2013) Temperature acclimation and heat tolerance of

photosynthesis in Norwegian *Saccharina latissima* (Laminariales, Phaeophyceae). Journal of

Phycology 49:689–700. doi: 10.1111/jpy.12077

Bartsch I, Wiencke C, Bischof K, et al (2008) The genus Laminaria *sensu lato*: recent insights

and developments. Eur J Phycol 43:1–86. doi: 10.1080/09670260701711376

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. Journal of the Royal Statistical Society B 57:289–300.

Bolton JJ, Lüning K (1982) Optimal growth and maximal survival temperatures of Atlantic

Laminaria species (Phaeophyta) in culture. Marine Biology 66:89–94. doi:

10.1007/BF00397259

Boquete MT, Muyle A, Alonso C (2021) Plant epigenetics: phenotypic and functional diversity

beyond the DNA sequence. American Journal of Botany 108:553–558. doi: 10.1002/ajb2.1645

Boyko A, Blevins T, Yao Y, et al (2010) Transgenerational adaptation of *Arabidopsis* to stress

requires DNA methylation and the function of dicer-like proteins. PLoS ONE. doi:

10.1371/journal.pone.0009514

Bringloe TT, Verbruggen H, Saunders GW (2020) Unique biodiversity in Arctic marine forests

is shaped by diverse recolonization pathways and far northern glacial refugia. Proceedings of

the National Academy of Sciences 117:22590–22596. doi: 10.1073/pnas.2002753117

Cai J, Lovatelli A, Aguilar-Manjarrez J, et al (2021) Seaweeds and microalgae: an overview for

unlocking their potential in global aquaculture development. FAO Fisheries and Aquaculture,

Rome

Cen Y, Sage RF (2005) The Regulation of Rubisco Activity in Response to Variation in

Temperature and Atmospheric CO_2 Partial Pressure in Sweet Potato 1 [w]. Plant physiology

139:979–990. doi: 10.1104/pp.105.066233.1981

Clegg MT, Gaut BS, Learn GH, Morton BR (1994) Rates and patterns of chloroplast DNA

evolution. Proceedings of the National Academy of Sciences of the United States of America

91:6795–6801. doi: 10.1073/pnas.91.15.6795

Cock JM, Sterck L, Rouzé P, et al (2010) The *Ectocarpus* genome and the independent evolution

of multicellularity in brown algae. Nature 465:617–621. doi: 10.1038/nature09016

Correia B, Valledor L, Meijón M, et al (2013) Is the Interplay between Epigenetic Markers

Related to the Acclimation of Cork Oak Plants to High Temperatures? PLoS ONE. doi:

10.1371/journal.pone.0053543

Davison IR, Greene RM, Podolak EJ (1991) Temperature acclimation of respiration and

photosynthesis in the brown alga *Laminaria saccharina*. Marine Biology 110:449–454. doi:

10.1007/BF01344363

Diehl N, Bischof K (2021) Coping with a changing arctic: Mechanisms of acclimation in the

brown seaweed *Saccharina latissima* from Spitsbergen. Marine Ecology Progress Series

657:43–57. doi: 10.3354/meps13532

Dubrovina AS, Kiselev K V. (2016) Age-associated alterations in the somatic mutation and DNA

methylation levels in plants. Plant Biology 18:185–196. doi: 10.1111/plb.12375

Duffy JE, Benedetti-Cecchi L, Trinanes J, et al (2019) Toward a coordinated global observing

system for seagrasses and marine macroalgae. Frontiers in Marine Science. doi:

10.3389/fmars.2019.00317

Fan X, Han W, Teng L, et al (2020a) Single-base methylome profiling of the giant kelp

Saccharina japonica reveals significant differences in DNA methylation to microalgae and

plants. New Phytologist 225:234–249. doi: 10.1111/nph.16125

Fan X, Xie W, Wang Y, et al (2020b) The complete chloroplast genome of *Saccharina latissima*.

Mitochondrial DNA Part B: Resources 5:3499–3500. doi: 10.1080/23802359.2020.1825999

Field CB, Behrenfeld MJ, Randerson JT (1998) Primary Production of the Biosphere: Integrating

Terrestrial and Oceanic Components. 281:237–241.

Filbee-Dexter K (2020) Ocean Forests Hold Unique Solutions to Our Current Environmental

Crisis. One Earth 2:398–401. doi: 10.1016/j.oneear.2020.05.004

Finnegan E., Genger RK, Peacock WJ, Dennis ES (1998) DNA methylation in plants. Annual

Review of Plant Physiology and Plant Molecular Biology 49:223–247. doi:

10.1146/annurev.arplant.49.1.223

Foust CM, Preite V, Schrey AW, et al (2016) Genetic and epigenetic differences associated with

environmental gradients in replicate populations of two salt marsh perennials. Molecular

Ecology 25:1639–1652. doi: 10.1111/mec.13522

Gugger PF, Fitz-Gibbon S, Pellegrini M, Sork VL (2016) Species-wide patterns of DNA

methylation variation in *Quercus lobata* and their association with climate gradients.

Molecular Ecology 25:1665–1680. doi: 10.1111/mec.13563

Guzinski J, Ruggeri P, Ballenghien M, et al (2020) Seascape genomics of the sugar kelp

Saccharina latissima along the north eastern Atlantic latitudinal gradient. Genes 11:1–28. doi:

10.3390/genes11121503

Hu L, Xiao P, Jiang Y, et al (2018) Transgenerational epigenetic inheritance under

environmental stress by genome-wide DNA methylation profiling in cyanobacterium.

Frontiers in Microbiology 9:1–11. doi: 10.3389/fmicb.2018.01479

Huang LC, Hsiao LJ, Pu SY, et al (2012) DNA methylation and genome rearrangement

characteristics of phase change in cultured shoots of *Sequoia sempervirens*. Physiologia

Plantarum 145:360–368. doi: 10.1111/j.1399-3054.2012.01606.x

Kassambara, A. (2021). *rstatix*: Pipe-friendly framework for basic statistical tests

(R package version 0.7. 0) https://cran.rproject.org/web/packages/rstatix/index.html

Keller TE, Lasky JR, Yi S V. (2016) The multivariate association between genomewide DNA

methylation and climate across the range of *Arabidopsis thaliana*. Molecular Ecology 25:1823–

1837. doi: 10.1111/mec.13573

Kobayashi H, Ngernprasirtsiri J, Akazawa T (1990) Transcriptional regulation and DNA

methylation in plastids during transitional conversion of chloroplasts chromoplasts. The EMBO

Journal 9:307–313. doi: 10.1002/j.1460-2075.1990.tb08112.x

Krause-Jensen D, Archambault P, Assis J, et al (2020) Imprint of Climate Change on Pan-Arctic

Marine Vegetation. Frontiers in Marine Science 7:1–28. doi: 10.3389/fmars.2020.617324

Kuroiwa T (1991) The Replication, Differentiation, and Inheritance of Plastids with Emphasis

on the Concept of Organelle Nuclei. International Review of Cytology 128:1–62. doi:

10.1016/S0074-7696(08)60496-9

Lämke J, Bäurle I (2017) Epigenetic and chromatin-based mechanisms in environmental stress

adaptation and stress memory in plants. Genome Biology 18:1–11. doi: 10.1186/s13059-017-

1263-6

Le S, Josse J, Husson F (2008) FactoMineR: An R Package for Multivariate Analysis. Journal of

Statistical Software 25:1–18. doi: 10.18637/jss.v025.i01

Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler transform.

Bioinformatics 26:589–595. doi: 10.1093/bioinformatics/btp698

Liesner D, Fouqueau L, Valero M, et al (2020) Heat stress responses and population genetics

of the kelp *Laminaria digitata* (Phaeophyceae) across latitudes reveal differentiation among

North Atlantic populations. Ecology and Evolution 10:9144–9177. doi: 10.1002/ece3.6569

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biology 15:550. doi: 10.1186/s13059-014-0550-8

Lüning K (1984) Temperature tolerance and biogeography of seaweeds: The marine algal flora

of Helgoland (North Sea) as an example. Helgoländer Meeresuntersuchungen 38:305–317.

doi: 10.1007/BF01997486

Møller Nielsen M, Paulino C, Neiva J, et al (2016) Genetic diversity of *Saccharina latissima*

(Phaeophyceae) along a salinity gradient in the North Sea-Baltic Sea transition zone. Journal

of phycology 52:523–531. doi: 10.1111/jpy.12428

Monteiro C, Heinrich S, Bartsch I, et al (2019) Temperature Modulates Sex-Biased Gene

Expression in the Gametophytes of the Kelp *Saccharina latissima*. Frontiers in Marine Science

6:1–12. doi: 10.3389/fmars.2019.00769

Muniandy K, Tan MH, Song K, et al (2019) Comparative sequence and methylation analysis of

chloroplast and amyloplast genomes from rice. Plant Molecular Biology 100:33–46. doi:

10.1007/s11103-019-00841-x

Nishimura Y (2010) Uniparental inheritance of cpDNA and the genetic control of sexual

differentiation in *Chlamydomonas reinhardtii*. Journal of Plant Research 123:149–162. doi:

10.1007/s10265-009-0292-y

Nishimura Y, Misumi O, Kato K, et al (2002) An mt+ gamete-specific nuclease that targets mt-

chloroplasts during sexual reproduction in *C. reinhardtii*. Genes and Development 16:1116–

1128. doi: 10.1101/gad.979902

Nishimura Y, Misumi O, Matsunaga S, et al (1999) The active digestion of uniparental

chloroplast DNA in a single zygote of *Chlamydomonas reinhardtii* is revealed by using the

optical tweezer. Proceedings of the National Academy of Sciences of the United States of

America 96:12577–12582. doi: 10.1073/pnas.96.22.12577

Nishiyama R, Ito M, Yamaguchi Y, et al (2002) A chloroplast-resident DNA methyltransferase

is responsible for hypermethylation of chloroplast genes in *Chlamydomonas* maternal

gametes. Proceedings of the National Academy of Sciences of the United States of America

99:5925–5930. doi: 10.1073/pnas.082120199

Nishiyama R, Wada Y, Mibu M, et al (2004) Role of a nonselective *de Novo* DNA

methyltransferase in maternal inheritance of chloroplast genes in the green alga,

Chlamydomonas reinhardtii. Genetics 168:809–816. doi: 10.1534/genetics.104.030775

Paulino C, Neiva J, Coelho NC, et al (2016) Characterization of 12 polymorphic microsatellite

markers in the sugar kelp *Saccharina latissima*. Journal of Applied Phycology 28:3071–3074.

doi: 10.1007/s10811-016-0811-x

Richards CL, Pigliucci M (2020) Epigenetic Inheritance. A Decade into the Extended

Evolutionary Synthesis. Paradigmi XXXVIII:463–494. doi: 10.30460/99624

Scheschonk L, Becker S, Hehemann JH, et al (2019) Arctic kelp eco-physiology during the polar

night in the face of global warming: A crucial role for laminarin. Marine Ecology Progress Series

611:59–74. doi: 10.3354/meps12860

Scheschonk L, Bischof K, Kopp M, Jueterbock A (2022) Differences by Origin in Methylome

Suggest Eco-phenotypes in the Kelp *Saccharina latissima*. Evolutionary Applications 1–17. doi:

10.1111/eva.13382

Teagle H, Hawkins SJ, Moore PJ, Smale DA (2017) The role of kelp species as biogenic habitat

formers in coastal marine ecosystems. Journal of Experimental Marine Biology and Ecology

492:81–98. doi: 10.1016/j.jembe.2017.01.017

Teng L, Han W, Fan X, et al (2021) Integrative analysis of chloroplast DNA methylation in a

marine alga—*Saccharina japonica*. Plant Molecular Biology 105:611–623. doi:

10.1007/s11103-020-01113-9

Thuleau, S., & Husson, F. (2020). *FactoInvestigate*: automatic description of factorial analysis.

[R package] version 1. https://CRAN.R-project.org/package=FactoInvestigate

Umen JG, Goodenough UW (2001) Chloroplast DNA methylation and inheritance in

Chlamydomonas. Genes and Development 15:2585–2597. doi: 10.1101/gad.906701

Wang L, Leister D, Kleine T (2020) Chloroplast development and genomes uncoupled signaling

are independent of the RNA-directed DNA methylation pathway. Scientific Reports 10:1–12.

doi: 10.1038/s41598-020-71907-w

Wang S, Lv J, Zhang L, et al (2015) MethylRAD: A simple and scalable method for genome-wide

DNA methylation profiling using methylation-dependent restriction enzymes. Open Biology.

doi: 10.1098/rsob.150130

Yang X, Wang X, Yao J, Duan D (2021) Genome-Wide Mapping of Cytosine Methylation

Revealed Dynamic DNA Methylation Patterns Associated with Sporophyte Development of

Saccharina japonica. International Journal of Molecular Sciences. doi: 10.3390/ijms22189877

Ye N, Zhang X, Miao M, et al (2015) Saccharina genomes provide novel insight into kelp

biology. Nature Communications. doi: 10.1038/ncomms7986

Zhang H, Lang Z, Zhu J (2018) Dynamics and function of DNA methylation in plants. Nature

Reviews Molecular Cell Biology. doi: 10.1038/s41580-018-0016-z

Figure Legends

Figure *1*: **Significant differences in methylated sites and methylation level between conditions.**
Methylated CG sites (A, B), and methylation levels as reads per million (RPM) for CHG sites (C, D)
between populations and temperatures in laboratory samples (A, C) and compared between lab and
field samples (B, D). ns: not significant. Numbers at the bottom of the bars indicate sample size.
Significance codes after p-value correction (Benjamini-Hochberg) for multiple comparisons: '< 0.05': *,
'< 0.01': **, '< 0.001': ***, '< 0.0001': ****.

Figure *2*: **Principal Component Analysis for A: all samples, grouped by laboratory or field samples. B:**
laboratory samples grouped by origin. C: all samples, grouped by origin. D: field samples grouped by
origin. E: laboratory samples of both origins grouped by temperature treatment. Ellipses represent
95% confidence intervals around the group medians.

Figure *3:* **VennEuler diagram showing the numbers of differentially methylated sites between**
Helgoland and Spitsbergen when analysing field samples (blue), lab samples (orange), and all
(lab+field, grey) samples (suppl. Tab. 1-3). The labels give the product of the genes that were
differentially methylated in the respective partitions. _(1) and _(2) depict a gene that showed
differential methylation at two different sites. For the exact site position in the respective gene, see
suppl. Table 5 ('ST5 Origins').

Figure *4*: **VennEuler diagram showing the numbers of differentially methylated sites between lab**
and field samples when analysing Helgoland samples (yellow), Spitsbergen samples (green), and the
samples from both origins together (**grey**, suppl. Tab.4)**.** The labels give the product of the genes that
were differentially methylated in the respective partitions, with one gene without annotation record
at the ncbi database. _(1) and _(2) refer to a gene that showed differential methylation at two different
positions. For the exact position in the respective gene, see suppl. Table 6 ('ST6 Cultivation').

Fig. 5: **Epigenetic distance (Euclidean distance) for 'cultivation', Laboratory (orange) and Field (blue)**
samples (A), and for 'origin', Helgoland (yellow) and Spitsbergen (green) samples (B). The small black
dots depict the pairwise distances between the individual samples of each group (from the PCA, Fig.
2). The light grey dots are outliers detected by the boxplot script, but were included during epi-distance
analyses. The boxes represent the interquartile range (IQR) of the respective group, and the horizontal
line within the boxes gives the median Euclidean distance.

Supplementary

 Suppl. Figures

 Suppl. Fig. *1*: D**ifferences in methylated sites and methylation level between conditions.** Methylation
 levels as reads per million (RPM) for CHG sites (A, B), and methylated CG sites (A, B) between
 populations and temperatures in laboratory samples (A, C) and compared between lab and field
 samples (B, D). ns: not significant. Numbers at the bottom of the bars indicate sample size.

Suppl. Fig. *2*: **Scree plots showing the amount of variance explained by each dimension**. On the left
including all samples, on the right when removing two outliers.

Suppl.Fig. *3:* **Overview of the chloroplast methylome in *S. latissima* (MT151382) and *S. Japonica***

(NC_0198523.1). *S. japonica* data from Teng et al. 2021. Numbers on the outside of the circle represent

locations within the genomes. Numbers on y-axes show the methylation levels at each site. The inner

circle shows CHG methylation, and the outer circle shows CG methylation. Numbers along the outer

circles refer to megabases along the chloroplast genomes.

Suppl.Fig. *4*: **Sequencing output and methylation characteristics for each sample from Helgoland.**
Plots were used for outlier detection for PCA. A: Number of methylated sites (with more than two
reads) for each sample. B: Boxplot showing reads per million (RPM) for each sample, with each point
representing one methylated site. C: The number of sequenced and quality-trimmed reads for each
sample. D: Numbers of quality-trimmed reads that mapped back to the chloroplast genome of
Saccharina latissima.

Suppl.Fig. *5*: **Sequencing output and methylation characteristics for each sample from Spitsbergen.**
Plots were used for outlier detection for PCA. A: Number of methylated sites (with more than two
reads) for each sample. B: Boxplot showing reads per million (RPM) for each sample, with each point
representing one methylated site. C: The number of sequenced and quality-trimmed reads for each
sample. D: Numbers of quality-trimmed reads that mapped back to the chloroplast genome of
Saccharina latissima.

Suppl. Tab. 1

Sites differentially methylated (p < 0.05) between Helgoland and Spitsbergen populations when including all samples in the comparison. Positive Log2FC values indicate increased methylation in Spitsbergen compared to Helgoland, while negative values indicate higher methylation in Helgoland samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
21922-21953	1.04	0.196	2.66E-06	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CHG
68544-68575	0.933	0.302	0.0274	QKE47473.1	ycf3	gene	photosystem I assembly protein	CHG
39358-39389	0.456	0.159	0.0475	QKE47440.1	groEL	gene	60 kDa chaperonin	CHG
51728-51760	-4.00	0.903	0.000164	QKE47458.1	psaA	promoter	photosystem I P700 apoprotein A1	CG
9045-9077	-4.17	0.77	2.05E-06	QKE47418.1	atpA	gene	ATP synthase CF1 alpha chain	CG
55344-55376	-4.96	0.644	9.14E-13	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CG

Suppl. Tab. 2

Sites with differential DNA methylation (p < 0.05) between Helgoland and Spitsbergen populations when comparing only field samples. Positive Log2FC values indicate increased methylation in Spitsbergen compared to Helgoland, while negative values indicate higher methylation in Helgoland samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
21922-21953	0.867	0.256	0.0117	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CHG
39358-39389	0.776	0.222	0.0104	QKE47440.1	groEL	gene	60 kDa chaperonin	CHG
75619-75650	0.727	0.206	0.0104	QKE47479.1	cbbX	gene	cfxQ-like protein	CHG
55567-55598	-0.528	0.173	0.0259	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CHG
51728-51760	-3.15	0.994	0.0209	QKE47458.1	psaA	promoter	photosystem I P700 apoprotein A1	CG
55344-55376	-4.49	1.21	0.0104	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CG

Suppl. Tab. 3

Sites with differential DNA methylation (p < 0.05) between samples originating from Helgoland or Spitsbergen when comparing only laboratory samples. Positive Log2FC values indicate increased methylation in Spitsbergen compared to Helgoland, while negative values indicate higher methylation in Helgoland samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
68544-68575	1.43	0.457	0.023	QKE47473.1	ycf3	gene	photosystem I assembly protein	CHG
21922-21953	1.18	0.285	0.000824	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CHG
9045-9077	-5.24	0.701	2.71E-12	QKE47418.1	atpA	gene	ATP synthase CF1 alpha chain	CG
51728-51760	-5.24	1.43	0.00438	QKE47458.1	psaA	promoter	photosystem I P700 apoprotein A1	CG
55344-55376	-5.68	0.639	4.27E-17	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CG

Suppl. Tab. 4

Sites differentially methylated (p < 0.05) between laboratory and field samples. Positive Log2FC values indicate higher methylation in laboratory samples compared to field samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
88361-88392	2.27	0.479	0.000137	QKE47493.1	rpl3	gene	50S ribosomal protein L3	CHG
15235-15266	1.16	0.282	0.000659	QKE47426.1	rpoC2	gene	DNA-directed RNA polymerase beta' chain	CHG
88373-88404	0.987	0.298	0.00741	QKE47493.1	rpl3	gene	50S ribosomal protein L3	CHG
17309-17340	0.952	0.31	0.0124	QKE47426.1	rpoC2	gene	DNA-directed RNA polymerase beta' chain	CHG
64927-64959	0.752	0.173	0.000294	QKE47469.1	petA	gene	cytochrome f	CG
34231-34262	0.598	0.198	0.0134	QKE47436.1	secA	promoter	preprotein translocase subunit A	CHG
102711-102742	0.591	0.172	0.00654	QKE47520.1	tufA	gene	translation elongation factor Tu	CHG
124685-124717	0.561	0.181	0.0123	QKE47544.1	ycf39	gene	NmrA domain-containing protein	CG
46721-46753	0.551	0.195	0.0237	QKE47451.1	ycf42	gene	putative peroxiredoxin ycf42	CG
32980-33011	0.514	0.143	0.00416	QKE47436.1	secA	gene	preprotein translocase subunit A	CHG
102611-102643	0.49	0.175	0.0241	QKE47520.1	tufA	gene	translation elongation factor Tu	CG
22542-22574	0.439	0.161	0.0264	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CG
75988-76019	-0.435	0.158	0.0253	QKE47479.1	cbbX	gene	cfxQ-like protein	CHG
104220-104251	-0.567	0.173	0.00741	QKE47523.1	ftsH	gene	ATP-dependent zinc metalloprotease FtsH	CHG
127333-127364	-0.641	0.191	0.00741	QKE47546.1	ilvB	gene	acetolactate synthase large subunit	CHG
95574-95605	-0.728	0.16	0.000183	QKE47509.1	rps5	gene	30S ribosomal protein S5	CHG

Suppl.Table *5* see excel file 'SupplTab5_6_7_Fig3_4_Tab2' sheet 'ST5 Origins'
Suppl.Table *6* see excel file 'SupplTab5_6_7_Fig3_4_Tab2' sheet 'ST6 Cultivation'
Suppl.Table *7* see excel file 'SupplTab5_6_7_Fig3_4_Tab2' sheet 'ST7 Temperature'

Suppl.Table *8:* Statistics results of the epigenetic distance analysis.					
Epidistance field/lab test results				**Euclidean distance**	
		statistic	10499		**Mean dist.**
		p	0.0000000173	*Field*	153
Wilcoxon rank sum test		*p.signif*	****	*Lab*	219
	Field	*statistic (W)*	0.945		
		p	0.00000106		
	Lab	*statistic (W)*	0.932		
Shapiro-Wilks test		*p*	0.0000000264		
Epidistance Helgoland/Spitsbergen test results				**Euclidean distance**	
		statistic	10330		**Mean dist.**
		p	0.00000000231	*Helgoland*	171
Wilcoxon rank sum test		*p.signif*	****	*Spitsbergen*	190
	Helgoland	*statistic (W)*	0.965		
		p	0.0000305		
	Spitsbergen	*statistic (W)*	0.957		
Shapiro-Wilks test		*p*	0.0000172		

871 Suppl.Table 9: Fertilisation scheme of the laboratory male (♂) and female (♀) gametophyte cultures
872 from Helgoland (HG1-4) and Spitsbergen (SG1-4), resulting in the cultures (Helgoland (H1-4) and
873 Spitsbergen (S1-4) raised under laboratory conditions.

Gametophyte Cultures	HG2 ♀ X HG2 ♂	HG3 ♀ X HG3 ♂	HG4 ♀ X HG4 ♂	SG1 ♀ X SG1 ♂	SG2 ♀ X SG2 ♂	SG3 ♀ X SG3 ♂	SG34 ♀ X SG4 ♂
Sporophyte Culture	H2	H3	H4	S1	S2	S3	S4

875