Chloroplast DNA methylation in the kelp *Saccharina latissima* is determined by origin and influenced by cultivation

Authors: Lydia Scheschonk*[1], Anne M.L. Nilsen*[2], Kai Bischof[1], Alexander Jueterbock[2]

*shared first authorship

[1] University of Bremen, Marine Botany & MARUM, 28359 Bremen, Germany

[2] Algal and Microbial Biotechnology division, Faculty of Biosciences and Aquaculture, Nord University, Universitetsalleen 11, 8026 Bodø, Norway

Contact:

during revision: lyd_sch@uni-bremen.de

after acceptance: alexander.juterbock@nord.no

Abstract

DNA cytosine methylation is an important epigenetic mechanism in genomic DNA, but absent in the chloroplast DNA of most land plants. We detected methylation in the chloroplast DNA of the kelp *Saccharina latissima*, a non-model macroalgal species of high ecological (wild populations) and economic importance. Since the functional role of the chloroplast methylome is yet largely unknown, we compared for the first time the chloroplast DNA cytosine methylation between wild and cultured kelp from different climatic origins (High-Arctic (79 °N) and temperate (54 °N), laboratory samples at 5 °C, 10 °C and 15 °C). Our results suggest genome-wide differences in methylated sites, and methylation level, between the origins, and a strong effect of cultivation. Functions related to photosynthesis showed differential methylation only between origins. Significant differences in methylation between cultivated and wild samples of genes related to transcription and translation were unique to

the high-arctic samples. Both findings indicate that origin and cultivation strongly, but

differently affect the chloroplast methylome. Similar methylomes for samples from the same

origin – independent from whether they grow in the wild or in the lab – suggest that origin-

specific methylation marks on the chloroplast genome are inherited. Given that DNA-

methylation affects gene expression, our study suggests that lab-cultivation alters

epigenetically determined kelp chloroplast characteristics at least to the same degree as

ecotypic differentiation does, but likely on a much shorter timescale. This indicates the

capacity for rapid non-genetic adaptation in the kelp *Saccharina latissima*.

Keywords:

Chloroplast, organelle genome methylation, epigenetics, non-model organism, cultivation,

aquaculture, marine macrophyte, marine algae

Introduction

Primary production, by which photoautotrophic organisms convert oxidised inorganic carbon

into organic carbon via photosynthesis, is the one crucial process that ensures the existence

of nearly all food webs, both terrestrial and marine. Since chloroplasts are presumed to have

developed from a free-living cyano-bacteria to an endosymbiontic organelle (Nishimura 2010),

they feature their own, albeit reduced, genome. Due to the reduced nature of their genome,

the organelles are semi-autonomous, and depend on extra-organellar products for their

internal processes (Wang et al. 2020). Epigenetic mechanisms include histone modification,

non-coding RNA, and DNA methylation (Boquete et al. 2021). They have been shown to play

vital roles in many cellular processes, such as regulation of gene expression, mRNA processing,

timing of DNA replication, determination of chromatin structure, or silencing of transposons

(Finnegan et al. 1998; Zhang et al. 2018). DNA cytosine methylation is the most stable

epigenetic mechanism that can persist for multiple generations (Lämke and Bäurle 2017), thus

is presumed to be most relevant for epigenetic adaptation. In contrast, histone modifications

and ncRNAs are often only stable for several days or weeks (Lämke and Bäurle, 2017).

However, the function of the chloroplast genome methylation is yet unclear, but intensely

researched in the few species of mostly agricultural use that show chloroplast methylation,

such as tomato and rice, or the unicellular green alga Chlamydomonas (Kobayashi et al. 1990;

Nishimura et al. 1999; Muniandy et al. 2019). In marine primary producers, the first

chloroplast genome methylation has recently been documented in the brown macroalga

Saccharina japonica (Teng et al. 2021). Its functional relevance remains unknown.

Macroalgae are the predominant source of coastal marine primary production (Field et al.

1998; Filbee-Dexter 2020). Nearly exclusively restricted to the shorelines of the continents,

macroalgae act as foundation species of marine rocky shore ecosystems, in addition to

forming the base of food webs, and hence, have high and diverse ecological value (Bartsch et

al. 2008; Teagle et al. 2017; Duffy et al. 2019). Their economic value is increasing exponentially

as macroalgae cultivation gains momentum outside of Asia as an environmentally sustainable

new blue economy with a multitude of applications (Cai et al. 2021). In 2019, marine

macroalgae accounted for ~30% of the global production of marine aquaculture commodities

(120 million t), worth US$ 14.7 billion (Cai et al. 2021). However, in recent years macroalgae

distributions have shifted northward in response to increasing (summer) water temperatures

(Krause-Jensen et al. 2020; Bringloe et al. 2020). Heat waves affect both wild populations and

aquaculture yields. In kelp, the respective thermal tolerance of gameto- and sporophyte

stages seems to be the strongest factor defining their global distribution limits (Monteiro et

al. 2019; Liesner et al. 2020). Temperature is a controlling factor in many biological processes

such as photosynthetic carbon acquisition. Enzymatic activity of RubisCO, the carbon

capturing enzyme in photosynthesis, increases by a factor of 7 between 5°C and 15°C (Cen and

Sage 2005), and has been shown to be especially vulnerable to high temperatures. As the chloroplast genome carries several photosynthesis-relevant enzymes, investigating its methylation at different temperatures is particularly relevant when considering methylome-based temperature adaptation in accordance to the eco-genotype approach (Scheschonk et al. 2022). While epigenetic mechanisms have been extensively researched in terrestrial plants, knowledge about kelp epigenetics still is limited to a handful of studies (Cock et al. 2010; Fan et al. 2020a, b; Scheschonk et al. 2022), with a single study on chloroplast DNA methylation (Teng et al. 2021). In primary producers, the chloroplast genome has been presumed to be un-methylated, despite data suggesting otherwise (*Saccharina japonica*, Teng et al 2021; *Chlamydomonas reinhardtii*, Nishimura 2010;). Epigenetic mechanisms hold the key to rapid local adaptation, and hence the capacity for species survival and ecosystem stabilisation despite swift alterations in biotic and abiotic factors. Especially in the face of the staggering changes in local and global ecosystems due to the anthropogenic climate crisis, gaining knowledge in the field of eco-evolutionary dynamics is crucial. Our study is the first attempt to determine the influence of temperature, origin, and growth condition (wild *vs* cultivated) on the methylome of chloroplasts of the kelp *Saccharina latissima* (sugar kelp). In the northern hemisphere, this boreal-temperate marine brown alga, a congener species to *S. japonica* with a large latitudinal distribution range, is among the macroalgal species of both high ecological and economic value (Bartsch et al. 2008; Cai et al. 2021).

In this fundamental research study, we tested the following hypotheses:

1) *Saccharina latissima* features chloroplast genome methylations, as has been reported for *S. japonica* (Teng et al 2021).

98 2) The chloroplast methylome mirrors the nuclear methylome (Scheschonk et al. 2022) in

99 showing epigenetic differentiation between origins that remains under cultivation, suggesting

100 a trans-generational epigenetic memory.

101 3) The chloroplast methylome changes in response to temperature and lab cultivation, thus

102 providing potential for rapid non-genetic adaptation to environmental change.

Materials and Methods

104 Laboratory cultures, field samples, and 'DNA- Extraction and MethlylRAD sequencing are

105 described in more detail in Scheschonk et al (2022). Changes and specific adjustments to the

106 present study were necessary during the bioinformatics and statistics part, and are detailed

107 below ('Bioinformatics and Statistics').

Laboratory cultures

As described in more detail in Scheschonk et al (2022), four cultures per origin (S1-S4, H1-H4;

technical replicates) were obtained from mono-parentally fertilised clonal gametophyte

cultures of *Saccharina latissima* (Linnaeus) C.E. Lane, C. Mayes, Druehl & G.W. Saunders from

Helgoland, German Bight (Nordstrand, 54°11'18.9"N 7°54'14.1"E; HG1, HG2, HG3, HG4 in

culture since 2014) and Spitsbergen, Svalbard (Hansneset, Kongsfjorden, 78°59'26.0"N

11°58'42.3"E; SG1, SG2, SG3, SG4 in culture since 2015; see table Suppl. Table 1) were kept at

each of the three temperatures (5 °C, 10 °C, and 15 °C) under white light at 15 µmol m-2 s-1

(18 : 6 hours light : dark) from as early as initiation of fertilization. Only sporophytes with a

fresh weight exceeding 120 mg (without stipe) were taken for DNA extraction.

Field Samples

Adult sporophytes were sampled from Helgoland, German Bight (54°11'18.9"N 7°54'14.1"E),

and from Spitsbergen, Svalbard (78°59'26.0"N 11°58'42.3"E) at the same locations from which

the gametophytes for the laboratory grown sporophytes originated. From each sporophyte, 4

discs (∅=3 cm) were cut from the fronds, omitting the meristematic region, 'midsection', and

reproductive tissues. Samples from lab and field were dried in cellulose bags stored in silica.

In total 21 samples from the laboratory cultures, and 20 samples from the wild were analysed.

In contrast to the field samples, our laboratory samples were uniparentally fertilised.

However, given that chloroplast genes are transmitted solely from the maternal parent

(matrilineally; Kuroiwa 1991, in Nishimura 2010), the chloroplast genome inheritance is

identical in uniparentally and bi-parentally fertilised individuals. This circumstance enables a

direct comparison between the chloroplast methylomes of the cultured and wild samples.

Bioinformatics and Statistics of MethylRAD sequencing
131

Clean-up of raw data
DNA was extracted and MethylRAD sequencing libraries were prepared as described in detail

in Scheschonk et al (2022). The sequences were de-multiplexed by sample and quality

trimmed   with   TrimGalore!   v   0.4.1

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Bases with Phred-

score <20 (--quality 20) were eliminated, the adapter sequence was removed (--stringency 3),

and the terminal 2bp (both ends) were removed to eliminate any artefacts at the ligation

position (--clip_R1 2 \ --three_prime_clip_R1 2). Reads were checked for fragment (over-)

representation, base AT CG content bias, and read length with FastQC v0.11.8 before and after

quality trimming.

In-silico digestion
For an overview of potential methylation sites, and as a backbone to map sequences to the

reference chloroplast genome for the *S. latissima* chloroplast (Fan et al. 2020b) was digested

in silico using the custom python script InSilicoTypeIIbDigestion_corrected.py

(http://marinetics.org/2017/04/11/REdigestions.html) and settings for simulating the

restriction enzyme FspEI. The *in silico* digestion found 270 recognition sites, where 69 were

CCGG recognition sites, 61 CCTGG sites, 77 CCTGT sites, and 63 CCAGG sites.

Mapping and annotation

The sequenced reads were mapped to the *in silico* digested reference genome of the

S. latissima chloroplast, combined with the *in silico* digested genome of the closely related

species *Saccharina japonica* to account for reads belonging to the nuclear genome and avoid

false mappings to the chloroplast genome for those reads. Reads were mapped using Burrows-

Wheeler Aligner (BWA Version: 0.7.17-r1188) (Li and Durbin 2010). Duplicate mappings were

excluded from further analyses by filtering out reads with a mapping quality <10 using

samtools (v 1.9). A count table was created listing the number of reads that mapped back to

each methylation site. For this, htseq-count (v 0.7.2) was called on each sample alignment

with *in silico* digested fragment in gff3 format as count features. An additional table was made

with counts normalized to 'reads per million (RPM) = (Reads mapped per site $\times 10^6$)/(Total

number of mapped reads)' to account for varying sequencing depths.

Only methylated sites that were covered in at least one library with coverage >2 were kept in

count tables and RPM tables. The sites were annotated for their sequence context (CG or

CHG), genomic region, gene ID numbers, and gene functions based on the gff3 annotation file

for the *S. latissima* chloroplast genome from NCBI (Accession: MT151382.1) (Fan et al. 2020b).

Statistics

Principal Component Analysis

Differences in the chloroplast methylome were investigated with principal component

analysis (PCA) on RPM tables using the R package 'FactomineR' (Lê et al. 2008). Since lab and

field samples separated clearly into distinct groups in the PCA, their methylomes were

compared first for all samples, and then separated into origins (Helgoland and Spitsbergen).

Within the lab and field samples, and for all samples, we compared origins (Helgoland and

Spitsbergen). Within the lab samples, we compared between the treatment temperatures of

5°C, 10°C, and 15°C.

The outlier function in the R package 'FactoInvestigate' (Thuleau & Husson, 2020) detected

two outliers, FH1 and FH9 (Helgoland field samples) in the PCA. The outliers showed few reads

mapping to the genome, few methylated sites, and high RPM values for some of the

methylated sites (Suppl. Fig. 1). Thus, FH1 and FH9 samples were excluded from all follow-up

analyses. No outliers were found in the Spitsbergen samples. As both outliers belonged to the

same group (Helgoland field samples), outlier removal led to a reduction in variance explained

by the first two principal components from 69.9% to 28.9% (Suppl. Figs. 1-3).

We tested for differences in DNA methylation levels as RPM and for differences in the

numbers of methylated sites between populations, between treatment temperatures, and

between lab and field samples, using Wilcoxon rank sum tests. P-values were corrected with

the Benjamini-Hochberg method (Benjamini and Hochberg 1995) to control for the false

discovery rate, and results with adjusted p values padj <0.05 were considered significant. The

tests were performed using the R package 'rstatix' v0.7.0 (Kassambara, 2021).

We tested for correlation between the numbers of quality trimmed reads, and the numbers

of quality-trimmed reads that mapped back to the *Saccharina latissima* chloroplast genome

(Suppl. Fig. 2C, 3C, 2D, 3D).

We tested for significant differential methylation with the R package 'DESeq2' (Love et al.

2014) between 'Helgoland' *vs* 'Spitsbergen' in all samples ('All', or 'both locations'), and in only

field samples and only lab samples. Further, between 'field' *vs* 'lab' in all samples, and in only

Helgoland samples, and only Spitsbergen samples. We also compared pairwise between

temperature treatments: both origins combined (Helgoland + Spitsbergen 5 °C *vs* 10 °C; 5 °C

vs 15 °C; 10 °C *vs* 15 °C), separately for Helgoland and Spitsbergen (5 °C *vs* 10 °C; 5 °C *vs* 15 °C;

10 °C *vs* 15 °C), and for each temperature (Helgoland vs Spitsbergen @ 5 °C; 10 °C; 15 °C).

'DESeq2' normalised raw methylation counts before testing for differential methylation, and

corrected p-values with the Benjamini-Hochberg method for controlling false discovery rates

(Benjamini & Hochberg, 1995). Methylated sites with adjusted p-value padj < 0.05 were

considered significantly differentially methylated.

GO term analysis

Gene ontology (GO) terms for biological processes, cellular components, and molecular

functions were extracted from the database https://www.uniprot.org/ using Gene ID numbers

(Suppl. Table 2-5). The R package 'topGO' (Alexa & Rahnenführer, 2009) was used to test for

enrichment of the GO terms.

Gene names and functions

Names and functions of differentially methylated genes were extracted from the chloroplast

genome of *Saccharina latissima* published on NCBI (strain y-c14;

https://www.ncbi.nlm.nih.gov/nuccore/MT151382.1?report=graph).

Analysis of epigenetic distance

Epigenetic distances among lab and field samples (cultivation) or among samples from

Helgoland vs Spitsbeergen (origin) were estimated as euclidean distances among samples in

the PCA . T. The Shapiro-Wilk indicated non-normal distribution for groups analysed for

cultivation (p-value lab p < 0.0001, field p < 0.0001) and for origin (Helgoland p < 0.0001,

Spitsbergen p < 0.0001). Hence, we tested for differences in epigenetic distances among the

cultivation or origin categories using non-parametric Wilcox-rank-sum test (suppl. Table 6).

Results

Methylation levels and methylated sites

In-silico digestion identified 69 MethylRAD sites for CG methylation, and 201 for CHG

methylation in the chloroplast genome of *Saccharina latissima*. 20.7% and 24.8% of these sites

were methylated, respectively (Table 1).

Table *1:* Potentially methylated sites for each sequence context, the mean number of methylated sites per sample and standard deviations for the means. Methylation level is the mean percentage of potential methylation sites methylated in samples, shown in the table with standard deviations (SD).

Sequence Context	Potential Sites (in-silico)	Methylated sites (mean +/- SD)	Methylation level (mean +/- SD)
CG	69	14.3 ± 1.5	20.7% ± 2.2%
CHG	201	49.8 ± 3.0	24.8% ± 1.5%

The number of methylated CG sites was significantly higher in Helgoland than in Spitsbergen

samples both in laboratory samples ($p < 0.001$, Fig. 1 A) and in laboratory and field samples

combined ($p < 0.0001$, Fig. 1 B), but no significant differences were found between treatment

temperatures or between field and laboratory conditions for sites in CHG contexts (Suppl. Fig.

3 C, D).

Methylation levels normalized to reads per million (RPM) showed no significant differences in

CG contexts between any groups (Suppl. Fig. 4 A, B), but the Spitsbergen population showed

higher methylation levels than the Helgoland population in the laboratory samples ($p < 0.05$,

Fig. 1 C). There were no significant differences between field and lab conditions in CHG context

(Fig. 1 D).

Figure *1*: **Significant differences in methylated sites and methylation level between conditions.**
Methylated CG sites (A, B), and methylation levels as reads per million (RPM) for CHG sites (C, D)
between populations and temperatures in laboratory samples (A, C) and compared between lab and
field samples (B, D). ns: not significant. Numbers at the bottom of the bars indicate sample size.
Significance codes after p-value correction (Benjamini-Hochberg) for multiple comparisons: '< 0.05': *,
'< 0.01': **, '< 0.001': ***, '< 0.0001': ****.

 Principal Component Analysis

Laboratory and field samples formed distinct groups along the first principal component with

95% confidence ellipses not overlapping when both origins were combined (Fig. 2 A). The first

two principal components equally contributed about 14 % of the total sample variation, which

was similar between the comparisons of 'lab vs field' (Fig. 2 A) and 'Helgoland vs Spitsbergen'

(Fig. 2 B). However, for the origins (Fig. 2 B), the populations of Helgoland and Spitsbergen

formed distinct groups along the second principal component when lab and field samples

were combined (Fig. 2 B), while when field and laboratory samples were separately analysed

for origin, Helgoland and Spitsbergen samples again split along the first component (Fig. 2 C,

D). Temperatures in the laboratory samples (5°C, 10°C, 15°C) did not separate into distinct

groups (Fig. 2 E).

Figure *2*: **Principal Component Analysis for A: all samples, grouped by laboratory or field samples. B:**
laboratory samples grouped by origin. C: all samples, grouped by origin. D: field samples grouped by
origin. E: laboratory samples of both origins grouped by temperature treatment. Ellipses represent
95% confidence intervals around the group medians.

Differentially methylated sites and GO terms

Comparing the origins (Helgoland vs Spitsbergen, Fig. 3), differentially methylated sites were

as often higher methylated in Spitsbergen samples as in Helgoland samples (Suppl.Tab.1-3).

Of the eight genes with significant methylation differences between the origins (Fig. 3), only

three genes (psaB_(1), atpA, yfc3) were identical to the genes with significant methylation

differences between the origins at a respective temperature within laboratory samples (Table

2, see colour coding suppl. Table 7+8, sheets 'ST7 Origin' and 'ST8 Temperature').

Figure *3:* **VennEuler diagram showing the numbers of differentially methylated sites between**
Helgoland and Spitsbergen when analysing field samples (blue), lab samples (orange), and all
(lab+field, grey) samples (Suppl. Table 2-4). The labels give the product of the genes that were
differentially methylated in the respective partitions. (_1) and (_2) depict a gene that showed
differential methylation at two different sites. For the exact site position in the respective gene, see
suppl. Table 7 ('ST7 Origins').

Of the 69 genes that were analysed comparing growth conditions (Fig. 4), 16 genes were

differentially methylated between laboratory and field samples, two of which were also

differentially methylated between origins (Fig. 3, Table 2). Many of those genes were

differentially methylated at two different positions (suppl. Table 9, sheet 'ST9 Cultivation').

Figure *4*: **VennEuler diagram showing the numbers of differentially methylated sites between lab**
and field samples when analysing Helgoland samples (yellow), Spitsbergen samples (green), and the
samples from both origins together (**grey**, Suppl. Table 5)**.** The labels give the product of the genes
that were differentially methylated in the respective partitions, with one gene without annotation
record at the ncbi database. (_1) and (_2) refer to a gene that showed differential methylation at two
different positions. For the exact position in the respective gene, see suppl. Table 9 ('ST9 Cultivation').

Two genes were differentially methylated between temperature treatments (Table 2), with

'acetolactate synthase large subunit' (QKE47546.1) at two comparisons: 5°C to 10 °C (padj <

0.0001), and 5 °C to 15°C (padj < 0.0017). All other significant differentially methylated genes

were found between origins at different temperatures, but only partially overlapped with

those obtained when analysing 'all samples' (Fig. 3, see colour codes in suppl. Tables 7 + 8

(sheets 'ST7 Origins' + 'ST8 Temperature')).

Table *2*: **All conditions that were found to be differentially methylated by the DESeq2 analysis when**
analysing the different rearing temperatures of the laboratory samples. Most significances were
detected between the origins at respective temperature (Helgoland vs Spitsbergen @ T5/10/15), but

only 3 of those were identical to those detected when all lab samples were being analysed as a group
(Fig. 3, Suppl. Table 8 ('ST8 Temperature')). However, genes QKE47546.1 and QKE47431.1 (bold letters)
showed significantly different methylation when analysing for temperature difference as factor (origin,
T vs T).

Comparison	Gene Product	Gene ID
5° vs 10° Helgoland	**acetolactate synthase large subunit**	**QKE47546.1**
5° vs 15° Helgoland	**acetolactate synthase large subunit**	**QKE47546.1**
5° vs 10° Spitsbergen	**photochlorophyllide reductase subunit B**	**QKE47431.1**
5° Helgoland vs Spitsbergen	photosystem I P700 apoprotein A2 _(1)	QKE47459.1
	ATP synthase CF1 alpha chain	QKE47418.1
	acetolactate synthase large subunit	QKE47546.1
	photochlorophyllide reductase subunit B	QKE47431.1
10° Helgoland vs Spitsbergen	photosystem I P700 apoprotein A2 _(1)	QKE47459.1
	ATP synthase CF1 alpha chain	QKE47546.1
	photosystem I assembly protein	QKE47473.1
	photochlorophyllide reductase subunit B	QKE47431.1
15° Helgoland vs Spitsbergen	photosystem I P700 apoprotein A2 _(1)	QKE47459.1
	ATP synthase CF1 alpha chain	QKE47546.1

None of the extracted GOterms were significantly enriched according to the 'topGO'

enrichment analysis.

Analysis of Epigenetic Distance
In both comparisons ('cultivation', 'origin'), the epigenetic distances between the samples

within respective groups were significantly different between groups (Fig. 5). For 'cultivation',

distances were significantly higher in laboratory samples than in field samples ($p < 0.0001$,

median field 153, lab 210), while for 'origin' the samples obtained from Spitsbergen showed

significantly higher epi-distances than Helgoland ($p < 0.0001$, median Helgoland 171,

Spitsbergen 190; suppl. Table 6).

Fig. 5: **Epigenetic distance (Euclidean distance) for 'cultivation', Laboratory (orange) and Field (blue) samples (A), and for 'origin', Helgoland (yellow) and Spitsbergen (green) samples (B).** The small black dots depict the pairwise distances between the individual samples of each group (from the PCA, Fig. 2 A, B). The light grey dots are outliers detected by the boxplot script, but were included during epi-distance analyses. The boxes represent the interquartile range (IQR) of the respective group, and the horizontal line within the boxes gives the median Euclidean distance.

Discussion

Regarding Hypothesis 1, we could confirm a methylome in the chloroplast DNA of *Saccharina latissima* (Table 1). As with the nuclear genome (Scheschonk et al. 2022) the chloroplast methylome significantly differed between origins (Fig. 3, Table 2, Fig. 5 B), and growth conditions (Fig. 4, Fig. 5 A), suggesting a trans-generational epigenetic memory (Hypothesis 2). However, growth temperature had a smaller impact than expected (Table 2; Hypothesis 3). Differentially methylated sites between origins (Spitsbergen or Helgoland) were located in genes with functions related to photosynthesis or ATP synthesis. However, differentially methylated sites between growth conditions (laboratory or field) were predominantly located in genes related to translation and transcription (suppl. Table 7 + 9, sheets 'ST7 Origin' + 'ST9

Cultivation'; Fig. 3, Fig. 4). This suggests that the chloroplast methylome is origin-specific,

indicating an epigenetic memory, while showing capacity of rapid adaptation to shifts in

growth conditions via the epigenetic mechanism of cytosine methylation.

Sample origin

Population origin had a strong impact on the methylome according to PCA clusters (Fig. 2 B,

C, D), which was reflected in methylation levels, as well as in the number of methylated sites

that significantly differed between origins (Fig 1 A, C). Photosystem I (PSI) was the component

most affected by differences between the origins, as genes coding for varying aspects crucial

to PSI significantly differed at a total of four sites in three genes (Fig. 3, Table 2), with the

strongest effect on PSI P700 chlorophyll a apoprotein A2. The same gene was one of two genes

differentially methylated between the origins in all analysed temperature conditions of the

laboratory samples (Table 2, suppl. Table 8 sheet 'ST8 Temperature'). Therefore, the

difference can be assumed to be solely accounted for by origin. Part of the methylome was

observed to be origin-specific even after years of gametophyte lab-cultivation. Methylation

levels were significantly different between the origins (Fig. 1), independent from the rearing

condition (laboratory or wild). As most of the differentially methylated genes between origins

overlapped in field and lab samples (Fig. 3), this indicates the prevalence of a

transgenerational epigenetic memory, which is further backed by PCA results. PCA in addition

indicated the influence of cultivation on transgenerational memory, as origin played a more

pronounced role when analysing laboratory and field samples separately (Fig. 2 C, D) *vs*

combined (Fig. 2 B),. The difference was less pronounced in the laboratory than the field

samples (Fig. 2 C *vs* D). The differentially methylated genes were exclusively related to

functions of photosynthesis (Fig. 3), which likely can be explained by the latitudinal differences

in the light regimes between origins. Light conditions in the High-Arctic are strongly seasonal,

with absence of photosynthetically active radiation (PAR) for at least three months during the

arctic winter (polar night), and continuous PAR during the arctic summer (polar day). In boreal-

temperate regions, PAR is available throughout the year. Whether or not the differences

observed here are genetically induced cannot be assessed by our setup, but the prevalence of

the significant difference, detectable in the epigenetic signature that prevailed between

origins through cultivation, is a strong indicator for the eco-genotype approach (Scheschonk

et al 2022). This presumes that adaptation to a specific location is not solely a matter of genetic

nature (fixed only in the DNA), but that epigenetic mechanisms, especially inheritable ones

such as cytosine methylation, play an important role in local adaptation and eco-evolutionary

dynamics of a species (Richards and Pigliucci 2020; Jablonka and Lamb 2015) that are partly

independent from underlying genetic variation. Considering methylation and gene expression

were shown to be negatively correlated in the *S. japonica* nuclear and chloroplast genes (Fan

et al. 2020a; Teng et al. 2021) the (double-) methylation of the gene coding for apoprotein A2

(Fig. 3, suppl. Table 7 sheet 'ST7 Origin'), for example, which is one of the crucial components

of PSI, can be presumed to have a direct impact on the photosynthetic capacity which,

according to our results, is likely to differ between origins. Like with the nuclear genome

(Scheschonk et al 2022) this has crucial implications for cultivation processes as well as

restoration efforts, as it shows the importance of suitable origin when considering local

growth condition.

Cultivation effects

PCA showed that cultivation affected the methylome to the same degree as origin (2 A, C).

While lab and field samples showed no significant differences in genome-wide methylation

level and number of methylated sites (Fig. 1), they displayed significantly different

methylation levels for 23.19 % of the methylation sites (Fig. 4, suppl. Table 9 (sheet 'ST9

Cultivation'). Most (12 out of 16) of the sites differentially methylated between rearing

conditions had higher methylation levels in the lab than in field samples (Suppl. Table 2). This

suggests those genes were higher expressed in the field samples (Fan et al. 2020a; Teng et al.

2021), possibly due to a higher variety of genes required in the more variable field conditions.

Unexpectedly, epigenetic distance was higher in lab samples, even though they experience

more uniform environmental conditions than field samples. When translated to transcription,

a higher epigenetic distance is likely to result in more variable phenotypes. This might suggest

that the kelp methylome diversifies rather under stable lab conditions than under variable

field conditions, at least when the given cultivation conditions were not previously

encountered. Potential explanatory factors cover the whole process of cultivation, from

obtaining the sori, and securing the zoospores, or handling during the fertilisation process, to

culture conditions at any stage.

However, origin as factor had the strongest influence on the differences in cultivated samples

at gene level, while lab-field differences in methylation could only be detected in the

Spitsbergen population (Fig. 4, Table 2). A factor that might as well explain differences in the

methylomes between lab cultivated and wild kelp is the age of the sporophytes. The field

sporophytes were adult, while the lab samples were juveniles with a size of approximately 12-

15 cm. In plants, cytosine methylation levels tend to increase with age, but these differences

are most pronounced in meristematic areas (Dubrovina and Kiselev 2016), which were

avoided in the kelp samples. Because meristematic regions were omitted, and older

individuals (field samples) did not show higher methylation levels in the chloroplast genome,

the likelihood that the results were affected by sporophyte age differences is low.

Temperature as factor could be shown to have an origin-specific impact on the methylome

during cultivation (Table 2; see next paragraph). However, which step, or factor, or

combination thereof, during the cultivation process is likely to have caused the increase in

epigenetic distance will have to be assessed in future studies.

Whether or not genotypic differences can result in the epigenetic differences that we

observed has yet to be determined. In plants and algae, DNA cytosine methylation can be

present in CG, CHG, and CHH sequence contexts, where H is any of the bases A, C, or T

(Richards and Pigliucci 2020). Cytosine methylation can be heritable across several

generations (Lämke and Bäurle 2017; Anastasiadi et al 2021), hence is a likely candidate for

rapid non-genetic but heritable local adaptation. The inheritance of epigenetic traits induced

by environmental factors has been repeatedly reported (Foust et al. 2016; Gugger et al. 2016;

Keller et al. 2016). The difference in methylation between origins that persisted under

laboratory conditions indicates epigenetic inheritance of a molecular memory of the

environment. It suggests that methylation differences can remain stable across generations

in cultivation, indicating that a transgenerational epigenetic memory likely is present in the

S. latissima chloroplast genome.

Temperature and the methylome

The temperature treatments did not form distinct clusters in the PCA. Unlike the nuclear

methylome, the cultivated *S. latissima*s' chloroplast methylome did not differ in methylation

levels and numbers of methylated sites in response to the rearing temperatures of 5 °C, 10 °C

and 15 °C. Significant differential methylation solely caused by temperature were detected in

2.9 % of the genes (Table 2). However, when analysing the origins at temperature level (Table

2), most differences between origins found at discrete temperatures were in other genes than

those detected for the origin analysis. This again suggests the inheritance of an epigenetic

memory in accordance with the eco-genotype approach (Scheschonk et al. 2022), as the

methylomes reacted origin-specific at discrete temperatures. However, it cannot be ruled out

whether these epigenetic differences have a genotypic base.

437
The methylation levels of sites in the chloroplast genome were ca. 100 times higher than levels

in the nuclear genome as documented by Scheschonk et al. (2022). In the nuclear genome,

the mean methylation level was 0.21% ± 0.13% SD (Scheschonk et al. 2022), while in the

chloroplast, the levels were 20.7% ± 2.2% for CH contexts and 24.8% ± 1.5% for CHG contexts

(Table 1). The higher methylation level in the chloroplast can be explained by the low numbers

of potential methylation sites in the chloroplast genome (Table 1). The pattern of higher

methylation in the chloroplast than the nuclear genome is repeated in both *S. japonica* and

S. latissima, with some dissimilarities that could be caused by different sequencing methods

and the inclusion of gametophytes in the studies of *S. japonica* (Fan et al. 2020a; Teng et al.

2021).

Photosynthesis is highly sensitive to temperature (Cen and Sage 2005). However, the main

temperature response, in terms of methylation, is observable in the nuclear, but not in the

chloroplast genome. The lack of statistically significant differences in the chloroplast genome

as compared with the nuclear genome might partly be explained by the smaller number of

genes. The chloroplast genome has a size of 130 kb and contains 139 protein-coding genes,

three rRNA genes and 29 tRNA genes (Fan et al. 2020b), while the nuclear genome has a size

of 545 Mb and contains 18,733 protein-coding genes (Ye et al. 2015). However, no significant

differences were observed in neither methylation levels nor methylated sites between

temperature treatments, despite the ~ 100 x higher methylation level of the chloroplast. In

contrast, in the nuclear genome, a response in methylation levels and numbers of methylated

sites to warmer temperature treatments was observed, especially in the Helgoland samples

(Scheschonk et al. 2022). Possible the lack of statistical significance indicates locations in the

methylome where the methylation is highly conserved, and hence crucial on a functional level

regardless of origin or external (new) impact. Aother possible explanation for the overall low

amount of (differentially) methylated sites might be that the temperature exposure of the

laboratory samples was not stressful for the physiological processes regulated by the

chloroplast genome. The difference between nuclear and chloroplast methylome suggests

that between 5 °C and 15 °C the modulation of temperature responses is solely regulated in

the nuclear genome. In *Saccharina latissima*, it has been shown that temperature treatments

affect the nuclear methylome, and that differences in the methylome exist between

populations from different latitudes, suggesting that epigenetic modifications contribute to

local adaptation (Scheschonk et al. 2022). In addition, significant differences in the nuclear

methylome were found between wild and cultivated individuals (Scheschonk et al. 2022).

Earlier research on *S. latissima* suggests that short-term acclimation to different growth

temperatures can lead to changes in heat tolerance, and affect photosynthetic performance

(Davison et al. 1991), which supports the temperature priming hypothesis suggested for the

nuclear methylome. However, most of the differentially methylated genes were found

between origins at discrete temperatures (5/10/15 °C), but were not detected when

comparing the origins over all lab samples.

In the nuclear genome, most of the differentially methylated sites between lab and field

samples and between populations were in non-genic repetitive contexts (transposable

elements). In contrast, in the chloroplast genome, the differentially methylated sites were in

coding regions of genes, except for very few sites in a promoter region (Suppl. Table 2-5). This

is not surprising, given that chloroplast genomes are more condensed and have far fewer

noncoding regions than nuclear DNA (Clegg et al. 1994).genomes.

 Conclusion

The chloroplast genome mainly contributes to processes of photosynthesis. Our results

showed that sample origin and cultivation can strongly influence epigenetic mechanisms in

the chloroplast, while temperature had a minor effect. Eco-phenotypes are likely developed

in the chloroplast, as methylation differences in genes are prone to trigger differences in gene

expression (Richards and Pigliucci 2020, Anastasiadi et al 2021). The scale-up in cultivation of

Saccharina latissima during the last decade is likely to continue, and cultivation sites overall

will likely shift northward, in accordance to the preferences of wild populations. The capacity

of fast adaptation to changed environmental factors, facilitated by epigenetic mechanisms,

could prove particularly useful in the face of rapid anthropogenic climate change. However,

due to the high ecological and economic value of this kelp, the implication of changes in

methylation due to cultivation, and especially the possibility of differential chloroplast

methylation influencing photosynthetic capacity, need to be further assessed. Photosynthetic

capacity is directly linked to primary production, and the capability to build up and maintain

biomass. Processes impacting the chloroplast methylome would have implications for both

cultivated and wild populations. In cultivars, the importance of matching hatchery and

cultivation conditions to origin has been implied by our results. Furthermore, due to the

significant differences observed in the methylomes of cultivated *S. latissima*, it might be

possible to detect an epigenetic 'fingerprint' of cultivated specimen in the future, to detect

escapees to the wild. Regarding the current trend in northward shift, the further north

populations of *Saccharina latissima* settle, the more pressure the months of darkness during

polar night will pose (Scheschonk et al. 2019). Photosynthetic pigments were shown to be

functional throughout the prolonged darkness of the Arctic winter (Scheschonk et al 2019),

indicating the chloroplasts of High Arctic specimen retained photosynthetic capacity during

polar night. In which context differential methylation of chloroplast DNA facilitated by

differences in the eco-phenotypes plays a role in this context will be of interest, especially

regarding populations adapted to light conditions at more southern distributional habitats,

shifting northwards. In *Saccharina latissima*, phenotypic variation among populations is often

driven by phenotypic plasticity, and the species shows high plasticity in response to changing

abiotic factors (Bolton & Lüning 1982; Diehl and Bischof 2021). So far, research on differences

between origins in *S. latissima* has focused on genetic differentiation, with significant genetic

variation between populations (Møller Nielsen et al. 2016; Paulino et al. 2016; Guzinski et al.

2020). Epigenetic differences have largely been unexplored as an alternative source of

molecular variation between *S. latissima* populations. The differences we observed between

origins within the laboratory and the field samples indicate that the eco-phenotype approach

(Scheschonk et al 2021) is applicable to the organellar chloroplast genome as well.

Furthermore, our data suggests that the influence of cultivation seems to be at least partially

origin-specific, or that the influence of cultivation and origin on the methylome are

interconnected. Epigenetic modifications have been found to be under some degree of

genetic control, but can also be triggered by environmental factors such as temperature

changes (Boyko et al. 2010; Correia et al. 2013). Hence, the role of genetic differences

controlling the epigenetic responses reported here needs to be assessed in further detail in

future studies.

Data availability

529   Raw sequencing data of this study are available at NCBI SRA BioProject PRJNA809008.

Acknowledgements

We thank S. Niedzwiedz (University of Bremen, Germany) for securing the field samples, and G. Hoarau and H. Reiss (Nord University, Bodø, Norway) for providing lab space and sequencer. Thank you, M. Kopp (Nord University, Bodø, Norway), for the support during the sequencing procedure, and for preparing some of the libraries. Thanks to I. Bartsch and A. Wagner (Alfred Wegener Institute for Marine and Polar Research, Bremerhaven, Germany) for providing and hatching the sporophyte cultures.

Funding

LS acknowledges funding by Deutscher Akademischer Auslandsdienst (DAAD; research travel grant 'BremenIdeaOut' 2019). AN acknowledges funding by Nord University (Master Thesis and transitional stipend). AJ acknowledges funding by Nord University (research talents grant). Parts of this work have been submitted in partial fulfilment of the requirements for Master of Science, at Nord University, Bodo (AN), and Dr. rer. nat, at University of Bremen (LS).

Author Contributions

LS designed and conducted the experiment. AN, LS and AJ analysed and interpreted the data. LS and AN wrote the manuscript. All authors read and commented on previous versions of the manuscript, and approved the final version.

References

Alexa, A., & Rahnenführer, J. (2009). Gene set enrichment analysis with topGO. Bioconductor Improv, 27, 1-26.

Anastasiadi D, Venney CJ, Bernatchez L, Wellenreuther M (2021) Epigenetic inheritance and reproductive mode in plants and animals. Trends in Ecology and Evolution 36:1124–1140. doi: 10.1016/j.tree.2021.08.006

Andersen GS, Pedersen MF, Nielsen SL (2013) Temperature acclimation and heat tolerance of

photosynthesis in Norwegian *Saccharina latissima* (Laminariales, Phaeophyceae). Journal of

Phycology 49:689–700. doi: 10.1111/jpy.12077

Bartsch I, Wiencke C, Bischof K, et al (2008) The genus Laminaria *sensu lato*: recent insights

and developments. Eur J Phycol 43:1–86. doi: 10.1080/09670260701711376

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. Journal of the Royal Statistical Society B 57:289–300.

Bolton JJ, Lüning K (1982) Optimal growth and maximal survival temperatures of Atlantic

Laminaria species (Phaeophyta) in culture. Marine Biology 66:89–94. doi:

10.1007/BF00397259

Boquete MT, Muyle A, Alonso C (2021) Plant epigenetics: phenotypic and functional diversity

beyond the DNA sequence. American Journal of Botany 108:553–558. doi: 10.1002/ajb2.1645

Boyko A, Blevins T, Yao Y, et al (2010) Transgenerational adaptation of *Arabidopsis* to stress

requires DNA methylation and the function of dicer-like proteins. PLoS ONE. doi:

10.1371/journal.pone.0009514

Bringloe TT, Verbruggen H, Saunders GW (2020) Unique biodiversity in Arctic marine forests

is shaped by diverse recolonization pathways and far northern glacial refugia. Proceedings of

the National Academy of Sciences 117:22590–22596. doi: 10.1073/pnas.2002753117

Cai J, Lovatelli A, Aguilar-Manjarrez J, et al (2021) Seaweeds and microalgae: an overview for

unlocking their potential in global aquaculture development. FAO Fisheries and Aquaculture,

Rome

Cen Y, Sage RF (2005) The Regulation of Rubisco Activity in Response to Variation in
Temperature and Atmospheric CO 2 Partial Pressure in Sweet Potato 1 [w]. Plant physiology
139:979–990. doi: 10.1104/pp.105.066233.1981

Clegg MT, Gaut BS, Learn GH, Morton BR (1994) Rates and patterns of chloroplast DNA
evolution. Proceedings of the National Academy of Sciences of the United States of America
91:6795–6801. doi: 10.1073/pnas.91.15.6795

Cock JM, Sterck L, Rouzé P, et al (2010) The *Ectocarpus* genome and the independent evolution
of multicellularity in brown algae. Nature 465:617–621. doi: 10.1038/nature09016

Correia B, Valledor L, Meijón M, et al (2013) Is the Interplay between Epigenetic Markers
Related to the Acclimation of Cork Oak Plants to High Temperatures? PLoS ONE. doi:
10.1371/journal.pone.0053543

Davison IR, Greene RM, Podolak EJ (1991) Temperature acclimation of respiration and
photosynthesis in the brown alga *Laminaria saccharina*. Marine Biology 110:449–454. doi:
10.1007/BF01344363

Diehl N, Bischof K (2021) Coping with a changing arctic: Mechanisms of acclimation in the
brown seaweed *Saccharina latissima* from Spitsbergen. Marine Ecology Progress Series
657:43–57. doi: 10.3354/meps13532

Dubrovina AS, Kiselev K V. (2016) Age-associated alterations in the somatic mutation and DNA
methylation levels in plants. Plant Biology 18:185–196. doi: 10.1111/plb.12375

Duffy JE, Benedetti-Cecchi L, Trinanes J, et al (2019) Toward a coordinated global observing
system for seagrasses and marine macroalgae. Frontiers in Marine Science. doi:
10.3389/fmars.2019.00317

Fan X, Han W, Teng L, et al (2020a) Single-base methylome profiling of the giant kelp

Saccharina japonica reveals significant differences in DNA methylation to microalgae and

plants. New Phytologist 225:234–249. doi: 10.1111/nph.16125

Fan X, Xie W, Wang Y, et al (2020b) The complete chloroplast genome of *Saccharina latissima*.

Mitochondrial DNA Part B: Resources 5:3499–3500. doi: 10.1080/23802359.2020.1825999

Field CB, Behrenfeld MJ, Randerson JT (1998) Primary Production of the Biosphere: Integrating

Terrestrial and Oceanic Components. 281:237–241.

Filbee-Dexter K (2020) Ocean Forests Hold Unique Solutions to Our Current Environmental

Crisis. One Earth 2:398–401. doi: 10.1016/j.oneear.2020.05.004

Finnegan E., Genger RK, Peacock WJ, Dennis ES (1998) DNA methylation in plants. Annual

Review   of   Plant   Physiology   and   Plant   Molecular   Biology   49:223–247.   doi:

10.1146/annurev.arplant.49.1.223

Foust CM, Preite V, Schrey AW, et al (2016) Genetic and epigenetic differences associated with

environmental gradients in replicate populations of two salt marsh perennials. Molecular

Ecology 25:1639–1652. doi: 10.1111/mec.13522

Gugger PF, Fitz-Gibbon S, Pellegrini M, Sork VL (2016) Species-wide patterns of DNA

methylation variation in *Quercus lobata* and their association with climate gradients.

Molecular Ecology 25:1665–1680. doi: 10.1111/mec.13563

Guzinski J, Ruggeri P, Ballenghien M, et al (2020) Seascape genomics of the sugar kelp

Saccharina latissima along the north eastern Atlantic latitudinal gradient. Genes 11:1–28. doi:

10.3390/genes11121503

Hu L, Xiao P, Jiang Y, et al (2018) Transgenerational epigenetic inheritance under

environmental stress by genome-wide DNA methylation profiling in cyanobacterium.

Frontiers in Microbiology 9:1–11. doi: 10.3389/fmicb.2018.01479

Huang LC, Hsiao LJ, Pu SY, et al (2012) DNA methylation and genome rearrangement

characteristics of phase change in cultured shoots of *Sequoia sempervirens*. Physiologia

Plantarum 145:360–368. doi: 10.1111/j.1399-3054.2012.01606.x

Jablonka E and Lamb MJ (2015) The inheritance of acquired epigenetic variations.

International Journal of Epidemiology 44, 1094–1103

Kassambara, A. (2021). *rstatix*: Pipe-friendly framework for basic statistical tests

(R package version 0.7. 0) https://cran.rproject.org/web/packages/rstatix/index.html

Keller TE, Lasky JR, Yi S V. (2016) The multivariate association between genomewide DNA

methylation and climate across the range of *Arabidopsis thaliana*. Molecular Ecology 25:1823–

1837. doi: 10.1111/mec.13573

Kobayashi H, Ngernprasirtsiri J, Akazawa T (1990) Transcriptional regulation and DNA

methylation in plastids during transitional conversion of chloroplasts chromoplasts. The EMBO

Journal 9:307–313. doi: 10.1002/j.1460-2075.1990.tb08112.x

Krause-Jensen D, Archambault P, Assis J, et al (2020) Imprint of Climate Change on Pan-Arctic

Marine Vegetation. Frontiers in Marine Science 7:1–28. doi: 10.3389/fmars.2020.617324

Kuroiwa T (1991) The Replication, Differentiation, and Inheritance of Plastids with Emphasis

on the Concept of Organelle Nuclei. International Review of Cytology 128:1–62. doi:

10.1016/S0074-7696(08)60496-9

Lämke J, Bäurle I (2017) Epigenetic and chromatin-based mechanisms in environmental stress

adaptation and stress memory in plants. Genome Biology 18:1–11. doi: 10.1186/s13059-017-

1263-6

Le S, Josse J, Husson F (2008) FactoMineR: An R Package for Multivariate Analysis. Journal of

Statistical Software 25:1–18. doi: 10.18637/jss.v025.i01

Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler transform.

Bioinformatics 26:589–595. doi: 10.1093/bioinformatics/btp698

Liesner D, Fouqueau L, Valero M, et al (2020) Heat stress responses and population genetics

of the kelp *Laminaria digitata* (Phaeophyceae) across latitudes reveal differentiation among

North Atlantic populations. Ecology and Evolution 10:9144–9177. doi: 10.1002/ece3.6569

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biology 15:550. doi: 10.1186/s13059-014-0550-8

Lüning K (1984) Temperature tolerance and biogeography of seaweeds: The marine algal flora

of Helgoland (North Sea) as an example. Helgoländer Meeresuntersuchungen 38:305–317.

doi: 10.1007/BF01997486

Møller Nielsen M, Paulino C, Neiva J, et al (2016) Genetic diversity of *Saccharina latissima*

(Phaeophyceae) along a salinity gradient in the North Sea-Baltic Sea transition zone. Journal

of phycology 52:523–531. doi: 10.1111/jpy.12428

Monteiro C, Heinrich S, Bartsch I, et al (2019) Temperature Modulates Sex-Biased Gene

Expression in the Gametophytes of the Kelp *Saccharina latissima*. Frontiers in Marine Science

6:1–12. doi: 10.3389/fmars.2019.00769

Muniandy K, Tan MH, Song K, et al (2019) Comparative sequence and methylation analysis of

chloroplast and amyloplast genomes from rice. Plant Molecular Biology 100:33–46. doi:

10.1007/s11103-019-00841-x

Nishimura Y (2010) Uniparental inheritance of cpDNA and the genetic control of sexual

differentiation in *Chlamydomonas reinhardtii*. Journal of Plant Research 123:149–162. doi:

10.1007/s10265-009-0292-y

Nishimura Y, Misumi O, Kato K, et al (2002) An mt+ gamete-specific nuclease that targets mt-

chloroplasts during sexual reproduction in *C. reinhardtii*. Genes and Development 16:1116–

1128. doi: 10.1101/gad.979902

Nishimura Y, Misumi O, Matsunaga S, et al (1999) The active digestion of uniparental

chloroplast DNA in a single zygote of *Chlamydomonas reinhardtii* is revealed by using the

optical tweezer. Proceedings of the National Academy of Sciences of the United States of

America 96:12577–12582. doi: 10.1073/pnas.96.22.12577

Nishiyama R, Ito M, Yamaguchi Y, et al (2002) A chloroplast-resident DNA methyltransferase

is responsible for hypermethylation of chloroplast genes in *Chlamydomonas* maternal

gametes. Proceedings of the National Academy of Sciences of the United States of America

99:5925–5930. doi: 10.1073/pnas.082120199

Nishiyama R, Wada Y, Mibu M, et al (2004) Role of a nonselective *de Novo* DNA

methyltransferase in maternal inheritance of chloroplast genes in the green alga,

Chlamydomonas reinhardtii. Genetics 168:809–816. doi: 10.1534/genetics.104.030775

Paulino C, Neiva J, Coelho NC, et al (2016) Characterization of 12 polymorphic microsatellite

markers in the sugar kelp *Saccharina latissima*. Journal of Applied Phycology 28:3071–3074.

doi: 10.1007/s10811-016-0811-x

Richards CL, Pigliucci M (2020) Epigenetic Inheritance. A Decade into the Extended

Evolutionary Synthesis. Paradigmi XXXVIII:463–494. doi: 10.30460/99624

Scheschonk L, Becker S, Hehemann JH, et al (2019) Arctic kelp eco-physiology during the polar

night in the face of global warming: A crucial role for laminarin. Marine Ecology Progress Series

611:59–74. doi: 10.3354/meps12860

Scheschonk L, Bischof K, Kopp M, Jueterbock A (2022) Differences by Origin in Methylome

Suggest Eco-phenotypes in the Kelp *Saccharina latissima*. Evolutionary Applications 1–17. doi:

10.1111/eva.13382

Teagle H, Hawkins SJ, Moore PJ, Smale DA (2017) The role of kelp species as biogenic habitat

formers in coastal marine ecosystems. Journal of Experimental Marine Biology and Ecology

492:81–98. doi: 10.1016/j.jembe.2017.01.017

Teng L, Han W, Fan X, et al (2021) Integrative analysis of chloroplast DNA methylation in a

marine alga—*Saccharina japonica*. Plant Molecular Biology 105:611–623. doi:

10.1007/s11103-020-01113-9

Thuleau, S., & Husson, F. (2020). *FactoInvestigate*: automatic description of factorial analysis.

[R package] version 1. https://CRAN.R-project.org/package=FactoInvestigate

Umen JG, Goodenough UW (2001) Chloroplast DNA methylation and inheritance in

Chlamydomonas. Genes and Development 15:2585–2597. doi: 10.1101/gad.906701

Wang L, Leister D, Kleine T (2020) Chloroplast development and genomes uncoupled signaling

are independent of the RNA-directed DNA methylation pathway. Scientific Reports 10:1–12.

doi: 10.1038/s41598-020-71907-w

Wang S, Lv J, Zhang L, et al (2015) MethylRAD: A simple and scalable method for genome-wide

DNA methylation profiling using methylation-dependent restriction enzymes. Open Biology.

doi: 10.1098/rsob.150130

Yang X, Wang X, Yao J, Duan D (2021) Genome-Wide Mapping of Cytosine Methylation

Revealed Dynamic DNA Methylation Patterns Associated with Sporophyte Development of

Saccharina japonica. International Journal of Molecular Sciences. doi: 10.3390/ijms22189877

Ye N, Zhang X, Miao M, et al (2015) Saccharina genomes provide novel insight into kelp

biology. Nature Communications. doi: 10.1038/ncomms7986

Zhang H, Lang Z, Zhu J (2018) Dynamics and function of DNA methylation in plants. Nature

Reviews Molecular Cell Biology. doi: 10.1038/s41580-018-0016-z

Figure Legends

Figure *1*: **Significant differences in methylated sites and methylation level between conditions.**
Methylated CG sites (A, B), and methylation levels as reads per million (RPM) for CHG sites (C, D)
between populations and temperatures in laboratory samples (A, C) and compared between lab and
field samples (B, D). ns: not significant. Numbers at the bottom of the bars indicate sample size.
Significance codes after p-value correction (Benjamini-Hochberg) for multiple comparisons: '< 0.05': *,
'< 0.01': **, '< 0.001': ***, '< 0.0001': ****.

Figure *2*: **Principal Component Analysis for A: all samples, grouped by laboratory or field samples. B:**
laboratory samples grouped by origin. C: all samples, grouped by origin. D: field samples grouped by
origin. E: laboratory samples of both origins grouped by temperature treatment. Ellipses represent
95% confidence intervals around the group medians.

Figure *3:* **VennEuler diagram showing the numbers of differentially methylated sites between**
Helgoland and Spitsbergen when analysing field samples (blue), lab samples (orange), and all
(lab+field, grey) samples (Suppl. Table 2-4). The labels give the product of the genes that were
differentially methylated in the respective partitions. (_1) and (_2) depict a gene that showed
differential methylation at two different sites. For the exact site position in the respective gene, see
suppl. Table 7 ('ST7 Origins').

Figure *4*: **VennEuler diagram showing the numbers of differentially methylated sites between lab**
and field samples when analysing Helgoland samples (yellow), Spitsbergen samples (green), and the

samples from both origins together (**grey**, Suppl. Table 5)**.** The labels give the product of the genes
that were differentially methylated in the respective partitions, with one gene without annotation
record at the ncbi database. (_1) and (_2) refer to a gene that showed differential methylation at two
different positions. For the exact position in the respective gene, see suppl. Table 9 ('ST9 Cultivation').

Fig. 5: **Epigenetic distance (Euclidean distance) for 'cultivation', Laboratory (orange) and Field (blue)**
samples (A), and for 'origin', Helgoland (yellow) and Spitsbergen (green) samples (B). The small black
dots depict the pairwise distances between the individual samples of each group (from the PCA, Fig. 2
A,B). The light grey dots are outliers detected by the boxplot script, but were included during epi-
distance analyses. The boxes represent the interquartile range (IQR) of the respective group, and the
horizontal line within the boxes gives the median Euclidean distance.

Supplementary

Suppl. Figures

 Suppl. Fig. *1*: **Scree plots showing the amount of variance explained by each dimension**. On the left
 including all samples, on the right when removing two outliers.

 Suppl.Fig. *2*: **Sequencing output and methylation characteristics for each sample from Helgoland.**
 Plots were used for outlier detection for PCA. A: Number of methylated sites (with more than two
 reads) for each sample. B: Boxplot showing reads per million (RPM) for each sample, with each point
 representing one methylated site. C: The number of sequenced and quality-trimmed reads for each
 sample. D: Numbers of quality-trimmed reads that mapped back to the chloroplast genome of
 Saccharina latissima.

 Suppl.Fig. *3*: **Sequencing output and methylation characteristics for each sample from Spitsbergen.**
 Plots were used for outlier detection for PCA. A: Number of methylated sites (with more than two
 reads) for each sample. B: Boxplot showing reads per million (RPM) for each sample, with each point
 representing one methylated site. C: The number of sequenced and quality-trimmed reads for each
 sample. D: Numbers of quality-trimmed reads that mapped back to the chloroplast genome of
 Saccharina latissima.

Suppl. Fig. *4*: **Differences in methylated sites and methylation level between conditions.** Methylation levels as reads per million (RPM) for CHG sites (A, B), and methylated CG sites (A, B) between populations and temperatures in laboratory samples (A, C) and compared between lab and field samples (B, D). ns: not significant. Numbers at the bottom of the bars indicate sample size.

780   Suppl.Fig. *5:* **Overview of the chloroplast methylome in *S. latissima* (MT151382) and *S. Japonica***
781   **(NC_0198523.1).** *S. japonica* data from Teng et al. 2021. Numbers on the outside of the circle represent
782   locations within the genomes. Numbers on y-axes show the methylation levels at each site. The inner
783   circle shows CHG methylation, and the outer circle shows CG methylation. Numbers along the outer
784   circles refer to megabases along the chloroplast genomes.

Suppl. Table 1: Fertilisation scheme of the laboratory male (♂) and female (♀) gametophyte cultures from Helgoland (HG1-4) and Spitsbergen (SG1-4), resulting in the cultures (Helgoland (H1-4) and Spitsbergen (S1-4) raised under laboratory conditions.

Gametophyte Cultures	HG2 ♀ X HG2 ♂	HG3 ♀ X HG3 ♂	HG4 ♀ X HG4 ♂	SG1 ♀ X SG1 ♂	SG2 ♀ X SG2 ♂	SG3 ♀ X SG3 ♂	SG34 ♀ X SG4 ♂
Sporophyte Culture	H2	H3	H4	S1	S2	S3	S4

Suppl. Table 2 Sites differentially methylated (p < 0.05) between Helgoland and Spitsbergen populations when including all samples in the comparison. Positive Log2FC values indicate increased methylation in Spitsbergen compared to Helgoland, while negative values indicate higher methylation in Helgoland samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
21922-21953	1.04	0.196	2.66E-06	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CHG
68544-68575	0.933	0.302	0.0274	QKE47473.1	ycf3	gene	photosystem I assembly protein	CHG
39358-39389	0.456	0.159	0.0475	QKE47440.1	groEL	gene	60 kDa chaperonin	CHG
51728-51760	-4.00	0.903	0.000164	QKE47458.1	psaA	promoter	photosystem I P700 apoprotein A1	CG
9045-9077	-4.17	0.77	2.05E-06	QKE47418.1	atpA	gene	ATP synthase CF1 alpha chain	CG
55344-55376	-4.96	0.644	9.14E-13	QKE47459.1	psaB	gene	photosystem I P700 apoprotein	CG

| | | | | | | | | | A2 | |

Suppl. Table 3

Sites with differential DNA methylation (p < 0.05) between Helgoland and Spitsbergen populations when comparing only field samples. Positive Log2FC values indicate increased methylation in Spitsbergen compared to Helgoland, while negative values indicate higher methylation in Helgoland samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
21922-21953	0.867	0.256	0.0117	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CHG
39358-39389	0.776	0.222	0.0104	QKE47440.1	groEL	gene	60 kDa chaperonin	CHG
75619-75650	0.727	0.206	0.0104	QKE47479.1	cbbX	gene	cfxQ-like protein	CHG
55567-55598	-0.528	0.173	0.0259	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CHG
51728-51760	-3.15	0.994	0.0209	QKE47458.1	psaA	promoter	photosystem I P700 apoprotein A1	CG
55344-55376	-4.49	1.21	0.0104	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CG

Suppl. Table 4
Sites with differential DNA methylation (p < 0.05) between samples originating from Helgoland or Spitsbergen when comparing only laboratory samples. Positive Log2FC values indicate increased methylation in Spitsbergen compared to Helgoland, while negative values indicate higher methylation in Helgoland samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
68544-68575	1.43	0.457	0.023	QKE47473.1	ycf3	gene	photosystem I assembly protein	CHG
21922-21953	1.18	0.285	0.000824	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CHG
9045-9077	-5.24	0.701	2.71E-12	QKE47418.1	atpA	gene	ATP synthase CF1 alpha chain	CG
51728-51760	-5.24	1.43	0.00438	QKE47458.1	psaA	promoter	photosystem I P700 apoprotein A1	CG
55344-55376	-5.68	0.639	4.27E-17	QKE47459.1	psaB	gene	photosystem I P700 apoprotein A2	CG

Suppl. Table 5

Sites differentially methylated (p < 0.05) between laboratory and field samples. Positive Log2FC values indicate higher methylation in laboratory samples compared to field samples.

Location	Log2FC	Log2FC std error	Adjusted p value	Gene ID	Gene name	Region	Product	Sequence context
88361-88392	2.27	0.479	0.000137	QKE47493.1	rpl3	gene	50S ribosomal protein L3	CHG
15235-15266	1.16	0.282	0.000659	QKE47426.1	rpoC2	gene	DNA-directed RNA polymerase beta' chain	CHG
88373-88404	0.987	0.298	0.00741	QKE47493.1	rpl3	gene	50S ribosomal protein L3	CHG
17309-17340	0.952	0.31	0.0124	QKE47426.1	rpoC2	gene	DNA-directed RNA polymerase beta' chain	CHG
64927-64959	0.752	0.173	0.000294	QKE47469.1	petA	gene	cytochrome f	CG
34231-34262	0.598	0.198	0.0134	QKE47436.1	secA	promoter	preprotein translocase subunit A	CHG
102711-102742	0.591	0.172	0.00654	QKE47520.1	tufA	gene	translation elongation factor Tu	CHG
124685-124717	0.561	0.181	0.0123	QKE47544.1	ycf39	gene	NmrA domain-containing protein	CG
46721-46753	0.551	0.195	0.0237	QKE47451.1	ycf42	gene	putative peroxiredoxin ycf42	CG
32980-33011	0.514	0.143	0.00416	QKE47436.1	secA	gene	preprotein translocase subunit A	CHG
102611-102643	0.49	0.175	0.0241	QKE47520.1	tufA	gene	translation elongation factor Tu	CG
22542-22574	0.439	0.161	0.0264	QKE47428.1	rpoB	gene	DNA-directed RNA polymerase beta chain	CG
75988-76019	-0.435	0.158	0.0253	QKE47479.1	cbbX	gene	cfxQ-like protein	CHG
104220-104251	-0.567	0.173	0.00741	QKE47523.1	ftsH	gene	ATP-dependent zinc metalloprotease FtsH	CHG
127333-127364	-0.641	0.191	0.00741	QKE47546.1	ilvB	gene	acetolactate synthase large subunit	CHG
95574-95605	-0.728	0.16	0.000183	QKE47509.1	rps5	gene	30S ribosomal protein S5	CHG

Suppl.Table *6:* Statistics results of the epigenetic distance analysis.					
Epidistance field/lab test results				**Euclidean distance**	
		statistic	10499		**Mean dist.**
		p	0.0000000173	*Field*	153
Wilcoxon rank sum test		*p.signif*	****	*Lab*	219
	Field	*statistic (W)*	0.945		
		p	0.00000106		
	Lab	*statistic (W)*	0.932		
Shapiro-Wilks test		*p*	0.0000000264		
Epidistance Helgoland/Spitsbergen test results				**Euclidean distance**	
		statistic	10330		**Mean dist.**
		p	0.00000000231	*Helgoland*	171
Wilcoxon rank sum test		*p.signif*	****	*Spitsbergen*	190
	Helgoland	*statistic (W)*	0.965		
		p	0.0000305		
	Spitsberge n	*statistic (W)*	0.957		
Shapiro-Wilks test		*p*	0.0000172		

Suppl.Table *7* see excel file 'SupplTab7_8_9_Fig3_4_Tab2' sheet 'ST7 Origins'
Suppl.Table *8* see excel file ' SupplTab7_8_9_Fig3_4_Tab2' sheet 'ST8 Temperature'
Suppl.Table *9* see excel file ' SupplTab7_8_9_Fig3_4_Tab2' sheet 'ST9 Cultivation'

804