Specific interaction between Group B *Streptococcus* CC17 hypervirulent clone and phagocytes

Anne-Sophie Bourrel[1,2,×], Amandine Picart[1,×], Jose-Carlos Fernandez[1], Constantin Hays[1,2], Virginie

Mignon[4], Bruno Saubaméa[4], Claire Poyart[1,2,3], Agnès Fouet[1], Asmaa Tazi[1,2,3], Julie Guignot[1*]

[1]Université Paris Cité, Institut Cochin, INSERM, U1016, CNRS, UMR8104, F-75014 Paris, France;

[2]Hôpitaux Universitaires Paris Centre, Cochin, Assistance Publique Hôpitaux de Paris, France; [3]Centre

National de Référence des Streptocoques, France ; [4]Plateforme PICMO, US25 INSERM, UAR3612 CNRS,

Faculté de Pharmacie, Université Paris Cité, Paris, France.

[×] These authors contributed equally to this work

* To whom correspondence should be addressed: julie.guignot@inserm.fr

ORCID number: 0000-0001-8890-4028. Tel: + 33 1 40 51 64 13; Fax: + 33 1 40 51 64 54.

Institut Cochin, 22 rue Méchain 75014 Paris, France.

ORCID number A. Fouet, 0000-0001-7715-9157;

CONFLICT OF INTEREST:

The authors have declared that no conflict of interest exists.

Keywords: CC17, *Streptococcus agalactiae*, Group B *Streptococcus*, phagocytosis, adhesion, THP-1.

Running title: Hypervirulent Group B Streptococcus and phagocytes

Abbreviations: CC17 (clonal complex 17), GBS (group B *Streptococcus*).

ABSTRACT

Streptococcus agalactiae also named Group B *Streptococcus* (GBS) is the most significant pathogen

causing invasive infections, such as bacteremia and meningitis, in neonates. Worldwide

epidemiological studies have shown that a particular clonal complex (CC) of capsular serotype III, the

CC17, is strongly associated with meningitis in neonates and is therefore designated as the

hypervirulent clone. Macrophages are a permissive niche for intracellular bacteria of all GBS clones. In

this study we deciphered the specific interaction of GBS CC17 strains with macrophages. Our study

revealed that CC17 strains are phagocytosed at a higher rate than GBS non-CC17 strains by human

monocytes and macrophages both in cellular models and primary cells. CC17 enhanced phagocytosis

is due to an initial enhanced-attachment step to macrophages mediated by the CC17 specific surface

protein HvgA and the PI-2b pilus (Spb1). We showed that two different inhibitors of scavenger

receptors (fucoidan and poly(I)) specifically inhibited CC17 adhesion and phagocytosis while not

affecting those of non-CC17 strains. Once phagocytosed, both CC17 and non-CC17 strains remained in

a LAMP-1 positive vacuole that ultimately fuses with lysosomes where they can survive at similar rates.

Finally, both strains displayed a basal egress which occurs independently from actin and microtubule

networks. Our findings provide new insights into the interplay between the hypervirulent GBS CC17

and major players of the host's innate immune response. This enhanced adhesion leading to higher

phagocytosis could reflect a peculiar capacity of the CC17 lineage to subvert the host immune

defenses, establish a niche for persistence or disseminate.

INTRODUCTION

The innate immune system plays a crucial role during host's interaction with pathogens, particularly in neonates who exhibit a naive adaptive immune system and are more vulnerable to invasive infections (1). Innate immunity largely relies on professional phagocytic cells, which constitute the first line of defense against pathogens. The large repertoire of phagocytic surface receptors that mediate the binding and internalisation of the pathogen is divided in opsonic and non-opsonic receptors. Opsonic receptors include the Fc receptors and complement receptors, which recognize antibody- or complement-opsonized pathogens, but also some non-opsonized pathogens (2, 3). Non-opsonic receptors are constituted by a wide array of family receptors including Toll like receptors (TLRs), integrins, C-type lectins or scavenger receptors that directly interact with pathogen-associated molecular patterns (PAMPs). Receptor recognition triggers intracellular signaling cascades leading to actin polymerization- which generates membrane protrusions required for bacterial engulfment. Signal transduction during phagocytosis is specific to the cell type and to the receptor(s) involved and can impact the intracellular fate of the microorganism (4). Once internalized, the pathogen resides in a phagosome, which rapidly undergoes maturation and acquires numerous proteins such as the late endosome/lysosome-associated membrane proteins LAMP-1 and LAMP-2. Ultimately, the phagosome fuses with lysosomes to become a phagolysosome with microbicidal features including a very acidic lumen (pH ~ 4.5), lysosomal proteases, and reactive oxygen species (ROS). This in turn leads to microbial killing and production of pro-inflammatory mediators that orchestrate the initiation of adaptive immunity.

Streptococcus agalactiae, also named Group B *Streptococcus* (GBS), is a capsulated chain-forming Gram-positive coccus commonly found in the gut and the lower urogenital tract of approximately 20% healthy adults (5). Although rarely pathogenic in adults, GBS is the leading cause of invasive infections in neonates (6). Two neonatal GBS-associated syndromes, referred to as "early-onset disease" (EOD) and "late-onset disease" (LOD), have been described, depending on the age at onset. EOD often manifests by respiratory distress that rapidly progresses into bacteremia and less frequently meningitis. In contrast, LOD manifests as a bacteremia highly associated with meningitis (20-50% cases) (7, 8). Based on the capsular polysaccharide composition, GBS has been classified into 10 distinct serotypes (Ia, Ib, and II–IX). Serotype III is the most frequently identified serotype in young infants (61.5%) (9). More specifically, within serotype III, multilocus sequence typing (MLST) has identified a hypervirulent lineage termed clonal complex 17 (CC17) which accounts for a substantial proportion (25%) of EOD cases and the vast majority (70%) of LOD cases worldwide, whereas it is isolated in about only 13% of non-pregnancy-related adult infections and colonization (7, 10, 11). GBS CC17 is a highly successful invasive clone in the neonate strongly associated with neonatal meningitis (~70%) in both

EOD and LOD (12-14). CC17 strains harbor a number of specific genes including the genes encoding

two surface proteins named Srr2 and HvgA that are involved in the crossing of the intestinal and blood

brain barriers (8, 15-17). Although not totally exclusive to the CC17 lineage, the pilus island PI-2b

encoding the Spb1 adhesin is found in the vast majority of the CC17 strains, while rarely present in

other human lineages (15, 18).

GBS has developed several mechanisms to resist immune detection, phagocytosis and microbial

degradation (19). The capsular polysaccharide (CPS) is one of the major virulence factors of GBS

allowing evasion from the host immune system, notably by preventing host complement deposition

on the bacterial surface (20, 21). Although capsule limits phagocytosis of complement-opsonized GBS,

opsonized-GBS can be phagocytosed by Fc and complement receptors respectively (19, 22). However,

there are large gaps in our understanding of the phagocytosis of non-opsonized GBS. The receptor(s)

and signaling pathway(s) involved in non-opsonized GBS phagocytosis are unknown. Interestingly, non-

opsonized GBS uptake leads to a better survival within phagocytes (19, 23-25). The persistence of GBS

in macrophages relies on numerous bacterial factors (19, 26). The expression of some virulence factors

associated to the CC17 lineage such as Spb1 have been described to increase phagocytosis (27-29).

However, the capacity of the CC17 lineage to exhibit enhanced phagocytosis and survival in

macrophages compared to those of other GBS lineages is still disputed (25, 30-32).

In the present study, we used a collection of GBS isolates representing a range of capsular types and

CCs and showed that CC17 strains are more phagocytosed than non-CC17 isolates. Our data support

that the enhanced phagocytosis of CC17 strains is solely due to an increased initial adhesion to

phagocytes. Once internalized, GBS CC17 and non-CC17 do not differentially affect the phagolysosome

maturation nor their ability to survive within phagocytes and can be released in the extracellular

compartement.

RESULTS

GBS CC17 strains are more phagocytosed than non-CC17 strains by macrophages and monocytes

The literature shows conflicting results on GBS lineages capacity to be phagocytosed (25, 30-32).

Therefore, we performed a phagocytosis assay using THP-1 differentiated macrophages to compare

the level of phagocytosis of a relevant collection of GBS isolates responsible for human invasive

infections including 27 CC17 strains and 40 non–CC17 strains (Table S2, Fig. 1A). The CC17 strains were

more phagocytosed than the non-CC17 strains (2.3-fold increase). To further characterize the lineage

differences and the phagocytosis process, we next used two well-characterized and fully sequenced

GBS isolates, one CC17 strain (BM110) and one CC23 strain (NEM316). They displayed a difference in

phagocytosis level comparable to that of CC17 versus non-CC17 strains in THP-1 macrophages (2.5-

fold increase) (Fig. S1). Using these two strains, we confirmed the higher phagocytosis of GBS CC17 in

primary human monocytes derived macrophages (hMDM) (Fig. 1B), in non-differentiated THP-1

monocytes (Fig. 1C) and in primary human monocytes (Fig. 1D).

Since the capsule is a major bacterial factor involved in GBS phagocytosis, we analyzed its contribution

to the enhanced phagocytosis of GBS CC17 strains. First, we stratified the results of the phagocytosis

assay performed with the clinical GBS strains presented in Fig. 1A according to their capsular type.

CC17 strains being almost exclusively of capsular type III and more rarely of type IV, we distinguished

within CC17 and non-CC17 isolates those of capsular type III, IV and all the other types (Fig. 2A). The

CC17 strains higher phagocytosis was independent from the capsular type. To further test the influence

of the capsular type, we analyzed the phagocytosis level of BM110 and NEM316 mutants, inactivated

for capsule expression. As expected, an increased phagocytosis of the capsule deficient strains was

observed with both mutant strains (Fig. 2B). Furthermore, the higher phagocytosis of the CC17 isolate

compared to the non-CC17 isolate could also be observed with the corresponding non-capsulated

mutants (Fig. 2B).

Altogether, these results show that CC17 strains are more phagocytosed than non-CC17 strains in a

capsule independent manner.

HvgA and Spb1 promote phagocytosis of CC17 strains

This observation prompted us to search for CC17 specific virulence factors that may account for this

enhanced phagocytosis. To this end, we tested the phagocytosis level of BM110 inactivated in genes

encoding CC17-specific (Srr2, HvgA) or -highly associated (Spb1) surface proteins (Fig. 3A) (15). Both

ΔhvgA and Δspb1 mutants exhibited reduced phagocytosis in THP-1 macrophages (65% and 54%

reduction, respectively) compared to the wild-type (WT) strain. In contrast, there was no difference

between the phagocytosis of the Δ*ssr2* and the WT strain. The role of Spb1 in GBS phagocytosis has

already been described (27, 28, 33), but not that of HvgA. To confirm the direct involvement of the

HvgA surface protein in phagocytosis, we used a close relative non-pathogenic *Lactococcus lactis*, in

which an expression vector encoding HvgA and enabling its presentation at the bacterial surface was

introduced (16). The heterologous expression of HvgA in *L. lactis* led to a 3-fold increase in

phagocytosis, as compared to *L. lactis* carrying an empty vector (Fig. 3B). The involvement of HvgA in

CC17 phagocytosis was also observed using hMDM (Fig. 3C).

The signal transduction leading to phagocytosis is similar between CC17 and non-CC17 strains

Cell membrane protrusion formation leading to bacterial engulfment results from cytoskeleton

rearrangements, which are mediated by specific signaling cascades induced by bacterial binding to cell

surface receptors (34). Numerous signaling molecules regulate the cytoskeleton, including the small

Rho GTPases Rho, Rac and CDC42 known to allow the formation of actin stress fibers, lamellipodial

protrusions, and filopodia, respectively (35). To better characterize the molecular pathways involved

in GBS engulfment, we performed phagocytosis assays using a wide array of inhibitors targeting

different signaling molecules (Table S3, Fig. 4A). BM110 phagocytosis was inhibited by cytochalasin D,

ZCL278, staurosporine, LY29002 and PP2 indicating that CC17 phagocytosis required actin, the small

GTPase CDC42, as well as the different kinases, PKC, Pi3K and Src, respectively. In contrast,

microtubules, the small GTPase Rac and the kinases Syk and Rock were not involved in CC17

phagocytosis, as indicated by the absence of effect of nocodazole, NSC23766, BAY61-3606 and Y27632

respectively (Fig. 4A). Interestingly, we found that NEM316, BM110Δ*hvgA* or BM110Δ*spb1*

phagocytosis involved the same signaling proteins than BM110 (Fig. 4B, 4C and 4D respectively).

To gain further insight into these findings, we characterized the early contact between GBS and

macrophages, which occurs prior to phagocytosis by Scanning Electron Microscopy (SEM) (Fig. 4E).

SEM revealed that numerous BM110 chains were adherent to macrophages in contrast to only a few

NEM316 chains (Fig. 4E, left panel). In addition, the uptake of both BM110 and NEM316 was

accompanied by the formation of filopodia-like cellular protrusions that were observed at 15 min

(Fig.4E, middle panel) and lamellipodia-like structure at 30 min (Fig. 4E, right panel).

Taken together, our results show that CC17 and non-CC17 are both captured by filopodia and that

bacterial uptake involves similar signaling proteins. Our results also suggest that enhanced

phagocytosis of CC17 strains may be due, at least in part, to an increased adhesion.

CC17 enhanced phagocytosis is due to initial higher adhesion

To test a potential adhesion difference between CC17 and non-CC17 strains, we analyzed the adhesion

of GBS strains to THP-1 macrophage. To prevent bacterial engulfment, adhesion was assessed either

at 4°C or at 37°C in the presence of cytochalasin D, an inhibitor of actin polymerization (Fig. S2A). Data

showed that bacterial adhesion to phagocytes was similar for all strains independently of the method

used to block phagocytosis. In both cases, adhesion of the BM110 strain was higher than that of

NEM316. We next assessed adhesion at 4°C and observed a higher adhesion of CC17 clinical strains

than of non-CC17 strains (1.9-fold increase, Fig. 5A, Table S2) to THP-1 macrophages. The enhanced

adhesion was also observed with strain BM110 compared to NEM316 on THP-1 macrophages (Fig.

S2A), THP-1 monocytes (Fig. 5B) and hMDM (Fig. 5C). As observed for phagocytosis, the increased

adhesion of CC17 isolates was not related to the capsular serotype (Fig. S2B) and was still observed in

the absence of capsule expression (Fig. 5D). Both *ΔhvgA and Δspb1* mutants exhibited reduced

adhesion on THP-1 macrophages (23% and 48% reduction, respectively) compared to the WT strain

(Fig. 5E). The heterologous expression of HvgA in *L. lactis,* led to a 2-fold increase in adhesion to THP-

1 macrophages, relative to *L. lactis* carrying an empty vector (Fig. 5F). The involvement of HvgA in CC17

adhesion was also observed using hMDM (Fig. S2C). These results indicate that CC17 strains have a

higher level of adhesion compared to non-CC17 strains. Furthermore, the increased adhesion of CC17

strains is mediated by both Spb1 and HvgA.

To determine whether CC17 higher phagocytosis was solely due to the increased adhesion or to a more

complex phenomenon involving both increased adhesion and uptake, we compared the ratio of

phagocytosed bacteria to the initial number of adherent bacteria. BM110 and its derivative mutant

strains and NEM316 strains showed similar ratios of phagocytosed bacteria (Fig. 5G). Next, we infected

THP-1 macrophages with the strain BM110 at a lower multiplicity of infection than NEM316 (MOI 2

and MOI 10, respectively) to obtain similar numbers of adherent bacteria (Fig. 5H). In that case, similar

numbers of phagocytosed bacteria were obtained with both strains.

Taken together, our results show that the enhanced phagocytosis of BM110 strain is solely due to its

initial increased adhesion to macrophages.

CC17 uptake by macrophages is inhibited by scavenger receptor inhibitors

Phagocytes express multiple receptors involved in bacterial adhesion and uptake. To identify the

precise nature of the macrophage receptor(s) involved in GBS recognition and uptake, GBS CC17

phagocytosis was evaluated in cells pre-treated with various blocking molecules targeting putative

receptor(s) (Table S3, Fig. 6A). Whereas the inhibitors targeting FCγR, complement receptors, integrins

and lectin receptors did not affect GBS CC17 phagocytosis, fucoidan and poly(I), two different

scavenger receptor inhibitors, significantly reduced it by 60% and 32%, respectively. Importantly, the

inactive analog of poly(I), poly(C), used as a negative control, had no effect on CC17 phagocytosis (Fig.

6A). Interestingly, these phenotypes were specific to BM110 since no reduction of NEM316

phagocytosis was observed using these two scavenger receptor inhibitors (Fig. 6B). Importantly,

fucoidan was also found to reduce BM110 adhesion to THP-1 macrophages whereas it did not reduce

NEM316 adhesion (Fig. 6C). These results suggest that some scavenger receptors contribute to a

significant proportion of the GBS CC17 adhesion that lead to its uptake into macrophages.

Scavenger receptors constitute a large family of membrane-associated pattern recognition receptors

(PRRs) that act as receptors mediating direct non-opsonic adhesion and uptake of microbes and/or

their products (36). To determine which scavenger receptor(s) contribute(s) to the adhesion/uptake

of GBS CC17 in THP-1 macrophages, we characterized by RT-PCR their expression profile in THP-1

macrophages. Given that the enhanced phagocytosis of GBS CC17 is conserved in hMDM and

monocytes (Fig. 1B, C, D), we extended the analysis of scavenger receptors expression profile to these

cell types (Fig. 6D). Using the recent classification of scavenger receptors (37), we identified that, out

of the 25 different scavenger receptors tested, 13 were expressed by these three cellular models (Fig.

6D). Next, we silenced the expression of each of these 13 individual scavenger receptors by siRNA to

address their involvement in GBS CC17 uptake. Efficient knockdown levels were confirmed on two

individual genes expressed by THP-1 macrophages for which the knockdown efficiency was 78 and 98

% (Fig. S3A). As an additional control, the silencing of the complement receptor impaired the

phagocytosis of complemented-red blood cells (96% decreased) (Fig. S3B), demonstrating that siRNA

in THP-1 macrophages was efficient. However, when individual scavenger receptors were silenced, no

significant reduction in BM110 uptake was observed (Fig. 6E). Last, in agreement with the phagocytosis

assays using phagocytosis receptor inhibitors (Fig. 6A), BM110 phagocytosis was not decreased in

THP-1 macrophages silenced for FCyR, complement receptors, α5 or β3 integrin expression, Plaur or

CD44 (Fig. S3C) or TLRs (Fig. S3D).

CC17 and non-CC17 strains display similar capacity to survive within macrophages

After phagocytosis, the phagocytic vacuole containing bacteria undergoes maturation events leading

to the formation of a phagolysosome required for efficient bacterial degradation. GBS is known to

survive within macrophages for up to 24 hours (24, 25, 27, 31, 32). In order to gain insight into the

physiological relevance of our study, we compared the intracellular fate of BM110 and NEM316 strains

in THP-1 macrophages. The maturation of phagosomes containing BM110-GFP and NEM316-GFP

(Table S1) was followed by fluorescence microscopy at various time-points after phagocytosis (t=0,

t=2h30 and t=5h). BM110 bacteria colocalizing with LAMP-1 and with the Lysotracker (a fluorescent

dye labeling acidic organelles such as phagolysosome) were quantified after immunofluorescence

staining (Fig. 7A and C). No statistical differences were observed in the kinetic of phagosome

maturation between BM110 and NEM316 whatever the time-point (Fig. 7B and D), indicating that the

delivery of BM110 and NEM316 to late endosomes or lysosomes is similar.

We next compared the survival ability of 10 CC17 and 17 non-CC17 strains clinical isolates by

quantifying intracellular viable bacteria 2h30 after phagocytosis (Fig. 7E). Survival results were

normalized for each GBS strain to the number of initially phagocytosed bacteria. No significant

difference was observed in the survival ability of CC17 and non-CC17 strains (Fig. 7E), whatever the

capsular type (Fig. S4A). The survival rate of BM110 and NEM316 strains was also similar 24 h after

phagocytosis (Fig. S4B). Consistently, HvgA and Spb1 were not involved in CC17 survival within THP-1

macrophages (Fig. 7F) or hMDM (Fig. 7G).

Taken together, these results revealed that the intracellular fate of GBS CC17 and non-CC17 after

adhesion and phagocytosis is similar.

CC17 and non-CC17 strains can escape macrophages by a passive phenomenon

We next analyzed the capacity of GBS strains to escape THP-1 macrophages. After phagocytosis and

antibiotic treatment to eliminate extracellular bacteria, cells were incubated in antibiotic free medium

for short periods of time (45 min and 1h30) to allow potential bacterial escape from infected cells

without any bacterial multiplication within the extracellular media. GBS egress normalized to the

percentage of initial phagocytosis showed that BM110 and NEM316 strains displayed similar capacity

to escape from macrophages (Fig. 8A). GBS egress from THP-1 macrophages was neither inhibited by

cytochalasin D nor by nocodazole (Fig. 8B), indicating that it is independent of actin cytoskeleton and

microtubules, respectively.

In conclusion, both strains of GBS can escape from the intracellular compartment to be released into

the extracellular compartment at similar rates. Furthermore, bacterial escape is a passive

phenomenon, suggesting that escape does not result from the recycling of the bacteria-containing

vacuole to the plasma membrane.

DISCUSSION

The innate immune response is critical for pathogen clearance, especially in neonates in whom the adaptive immune system is not yet fully mature and the specific antibodies required for adaptive immune response are usually lacking. Phagocytic cells populate all tissues and organs of the body where they play an important role in pathogen clearance.

The ability of GBS lineages to interact specifically with phagocytes, particularly their ability to be phagocytosed, remains controversial. In a study using 35 isolates from various GBS lineages, no significant differences were observed in the phagocytosis of GBS strains stratified by capsular serotype or MLST (30). In contrast, a study performed with a more discrete number of GBS isolates (15 GBS strains including 4 CC17 strains) found that CC17 were phagocytosed by macrophages at significantly higher rates than other GBS isolates (32). Our study, performed with a higher number of invasive GBS clinical isolates (27 CC17 isolates and 40 non-CC17 isolates) demonstrates that GBS CC17 strains are more phagocytosed than non-CC17 strains. These conflicting results may be attributed to the use, in the first study, of heat-inactivation of GBS, that could have denatured bacterial surface proteins involved in phagocytosis (30). In agreement with this hypothesis, we demonstrated that the higher phagocytosis of CC17 strains does not depend on the polysaccharidic capsule, a major GBS virulence factor which protects the bacteria from phagocytosis but, rather on the expression of two surface proteins, Spb1, the backbone protein of the PI-2b pilus, and the HvgA adhesin, that promote a higher level of phagocytosis. In contrast, we did not observe any role for the Srr2 surface protein in GBS CC17 uptake by macrophages. In agreement with this result, we and others have found that blocking the known Srr2-receptors, α5β1 and αvβ3 integrins, by mimetic peptides (RGD and/or RGfV) or by silencing α5 or β3 integrin expression does not affect CC17 phagocytosis (this study and (23)). Together, these data are consistent and indicate that the Srr2 surface protein is not involved in the uptake of GBS by phagocytes.

Importantly, we found that the higher phagocytosis mediated by Spb1 and HvgA was solely the result of an increase of the initial adhesion to phagocytes. Both proteins have previously been described to play a role in adherence to host intestinal epithelial and cerebral endothelial cells (16, 28, 38, 39). Interestingly, in these two cell types, CC17 strains were found to have an approximately 2.5-fold increase in adhesion compared to non-CC17 strains. The increased ability to adhere to both intestinal epithelial and cerebral endothelial cells was found to have physiological functionality, as CC17 strains have a significantly greater ability to colonise the gut and cross the blood-brain barrier (16). Although it is not possible to address *in vivo* whether a 2.3-fold increase in phagocytosis for CC17 strains contributes to the hypervirulence of CC17 strains, our previous data support this hypothesis. After initial adherence to phagocytes, both CC17 and non-CC17 where captured by filopodia and internalized

within phagocytes in a process that involved similar signalling molecules. Filopodia formation is known

to require the Cdc42 GTPase activity leading to actin polymerization (35). Accordingly, we found that

GBS phagocytosis was strongly dependent on actin and Cdc42.

While CC17 had a higher adhesion rate, it shared, with non-CC17 strains, a common signaling pathway

leading to phagocytosis. This may reflect the recognition of similar cellular receptors - mediating

bacterial adhesion and subsequent internalisation of CC17 and non-CC17 strains - but with different

affinities. That both strain types interact with the same molecules but with different affinity has

previously been described for GBS interaction with fibrinogen which exhibits a 10-fold higher affinity

for CC17 strains than for non-CC17 strains (40). Alternatively, it may reflect an internalisation process

in a two-step mechanism involving a *sensus stricto* adhesion receptor followed by engulfment

mediated by a phagocytic receptor. In this scenario, CC17 or non-CC17 strains would recognise

different adhesion receptors, while the internalisation process would involve a common phagocytic

receptor. Two-step phagocytosis mechanisms have previously been described involving cross-talk

between TLRs and scavenger receptors (41, 42).

The identity of the cellular receptor(s) involved in the non-opsonized phagocytosis of GBS has been

carried out by many teams (22, 23, 43). Despite these efforts, such receptor(s) are still unknown. Our

study indicates that FcγRs, complement receptors and CD14 are not involved in the phagocytosis of

non-opsonized GBS, as described previously (22, 23, 43). Accordingly, GBS phagocytosis is independent

of Syk and Rock kinases that are known to be involved in FCγR and complement receptor signalling,

respectively (44). Interestingly, fucoidan and poly(I), two broad scavenger receptor inhibitors reduced

the phagocytosis of the BM110 CC17 strain, while having no significant effect on the adhesion and

phagocytosis of a non-CC17 strain. Scavenger receptors are a diverse family of proteins that recognize

polyanionic ligands and can act as PRR involved in the recognition of various microbial ligands as well

as in their phagocytosis (45). The polynucleotide ligand poly(I) acts by efficient competitive inhibition

of the binding of other polyanionic ligands to the cationic collagenous binding domain of some

scavenger receptors. Similarly, fucoidan easily interacts with positively charged molecules because of

its high negative charge at neutral pH (46, 47). These molecules have been successfully used to inhibit

adhesion and/or phagocytosis of several pathogens (48, 49). The inhibitory effect of fucoidan and

poly(I) on CC17 adhesion and phagocytosis led us to conclude that scavenger receptor(s) might be

involved in the adhesion and/or uptake of non-opsonized CC17 bacteria by macrophages. Since

scavenger receptors have a wide distribution, being expressed by phagocytes as well as by epithelial

and endothelial cells (37), it is tempting to speculate that Spb1 and HvgA promote adhesion to host

cells by recognizing the same receptors on phagocytes, epithelial and endothelial cells. Interestingly, it

has been recently described that the scavenger receptor SSC4D, expressed by phagocytes and

epithelial cells, interacts with BM110 strain (50). SSCD4 is a secreted scavenger receptor making this molecule unlikely to be involved in the initial adherence of CC17 bacteria to the macrophage surface. In addition, silencing SSC4D expression does not affect CC17 phagocytosis (not shown) confirming its absence of involvement in this phenotype. Despite our efforts, the siRNA screen performed to identify scavenger receptor(s) involved in GBS uptake was unsuccessful. Several hypotheses could explain this failure. Because CC17 increased phagocytosis is a phenotype conserved between macrophages and monocytes from cell lines and primary cells, we hypothesized that the receptor(s) involved in CC17 adhesion/phagocytosis is also conserved in these different cell types. However, we cannot exclude the possibility that different scavenger receptors in each cellular model allow CC17 uptake, or that several scavenger receptors have a cooperative effect. Also, the list of scavenger receptors is constantly growing and we might have missed the scavenger receptor(s) involved in GBS CC17 adhesion/phagocytosis. Finally, we cannot exclude a non-specific or an indirect effect of fucoidan and poly(I) on the signaling of others PRRs, as described previously (51).

Numerous pathogens have evolved sophisticated strategies to avoid capture and/or killing by phagocytes. GBS efficiently survive within phagocytes by expressing multiple bacterial factors involved in acid stress response and ROS resistance, including the orange carotenoid pigment hemolysin, the glutathione or the superoxide dismutase (SodA) (26, 52). The capacity of different GBS lineages to survive within macrophages has been controversial (25, 27, 31, 32). The discrepancies between the different studies likely come from either the low number of bacterial strains tested or to the calculation of survival rate (25, 27, 31, 32). Indeed, depending on whether the intracellular survival ability is related to the number of phagocytosed bacteria (intracellular bacteria at T0) or to the initial bacterial inoculum different results will be obtained. As CC17 strains display higher adhesion level leading to higher initial load of intracellular CC17 this might *in fine* lead to a higher recovering number of intracellular bacteria at later time points. By calculating the survival rate in relation to the number of phagocytosed bacteria, we found that CC17 and non-CC17 strains display similar intracellular fate with similar survival capacity. In agreement with this result, we found that Sbp1 and HvgA proteins do not contribute to intracellular survival. We also showed that after internalisation, bacteria can be released into the external compartment. As observed for bacterial survival, we found that bacterial egress of CC17 and non-CC17 strains occurs at a similar rate when normalised to the initially phagocytised bacteria. However, due to enhanced phagocytosis of CC17 strains, this results in a greater number of bacteria being released into the extracellular compartment. Interestingly, we found that bacterial egress occurs even when actin or microtubule networks are disrupted. Since most vacuolar movements involve the cytoskeleton, this suggests that GBS release into the extracellular compartment is a passive

phenomenon that may be due to spontaneous macrophage lysis. Further studies are needed to
confirm this hypothesis.

The hypervirulent CC17 clone is strongly associated with LOD (60% to 80% of the cases) for which the
gastrointestinal tract is likely to be the portal of entry. In the gut, CC17 strains can transcytose *via* M
cells and are found associated with CD11c - a marker for macrophages and dendritic cells - positive
cells in the Peyer's patches (8). Interestingly, an increased proportion of CD11c positive cells have been
found in the mesenteric lymph node of CC17-infected mice (8). This suggests that CD11c positive cells
may act as Trojan horses, to promote the dissemination of CC17 bacteria in the gut. In addition, the
higher CC17 intracellular load could also contribute to the establishment of a survival niche and
bacterial persistence in tissues.

In conclusion, the specific features associated with the interaction between the CC17 lineage and the
phagocytes likely contribute to its particular virulence and to the pathophysiology of infection. Indeed,
the increased CC17 bacterial loads in macrophages may confer a selective advantage to promote better
persistence and dissemination.

MATERIALS AND METHODS

Bacterial strains

Bacterial strains used in this study are listed in Table S1. GBS BM110 (CC17) and NEM316 (CC23) are well characterized capsular serotype III strains isolated from blood cultures of neonates (16). BM110 and NEM316 were used as CC17 and non-CC17 reference strains, respectively. Other GBS clinical strains isolated from invasive infections (*i.e.* from blood culture, cerebrospinal fluid or other normally sterile sites) were obtained from the French National Reference Center for Streptococci (http://www.cnr-strep.fr). Capsular genotyping and MLST was performed for all GBS strains (Table S2). All bacterial strains were cultured in Todd Hewitt (TH) broth or agar (Difco Laboratories, Detroit, MI) at 37°C, except for *L. lactis* which was grown at 30°C. When necessary, antibiotics were used to maintain plasmids at the following concentrations, spectinomycin 100 μg.mL^{-1}, erythromycin 10 μg.mL^{-1}.

THP-1 cell line and culture conditions

THP-1 (TIB-202 from ATCC) cell line is a spontaneous immortalized monocyte-like cell line, derived from the peripheral blood of a childhood case of acute monocytic leukemia. THP-1 cells were cultured at 37°C under a 5% CO_2 atmosphere in RPMI1640 + Glutamax (Gibco), 10% Fetal Calf Serum (FCS, Gibco) and 0.05 mM β-mercapto-ethanol. Cells were passaged twice per week and kept at a cell concentration below 1×10^6 cells /mL. THP-1 cell line was routinely tested for mycoplasma contamination (Mycoalert mycoplasma detection kit, LONZA). To differentiate monocytic THP-1 into adherent macrophages, cells were seeded in 24-well plates at 3×10^5 cells per well in culture media containing 200 nM phorbol 12-myristate 13-acetate (PMA) during 3 days, then media was replaced by fresh RPMI1640 complete media without PMA. Cells were incubated for an additional 5 days without any medium change to allow complete differentiation into macrophages as described previously (53).

Isolation and culture of primary human Peripheral Blood Mononuclear Cells (PBMCs) and Monocytes-Derived Macrophages (hMDMs)

Blood of adult healthy donors was used (Etablissement Français du Sang, Ile de France, Site Saint-Antoine-le Crozatier) with the appropriate ethics prior approval as stated in the EFS/ INSERM agreement #18EFS030, ensuring that all donors gave a written informed consent and providing anonymized samples. Primary human monocytes were negatively selected using the EasySep Human Monocyte enrichment kit (StemCell®) following the manufacturer's instructions. For hMDM, primary human PBMCs were isolated by density gradient sedimentation in Histopaque® (Sigma) according to the manufacturer's recommendations. Monocytes were then selected by adhesion to dishes for 2 h at 37°C in FCS-free medium (RPMI 1640 medium supplemented with 0.1 mg.ml^{-1} gentamicin). Monocytes

402   were differentiated into hMDMs for at least 7 days in complete medium containing 10 ng.mL^{-1} of

403   recombinant human macrophage colony-stimulating factor (rhM-CSF; R&D systems).

Phagocytosis and survival assay in monocytes and macrophages

Monocytes and macrophages from THP-1 cell line and primary cells were used to study the

phagocytosis and the intracellular survival of GBS strains. All experiments were carried out with GBS

strains grown to mid-log phase and washed twice with sterile Phosphate buffer saline (PBS) before

infection. After addition of bacteria, a quick centrifugation (5 min, 200 × g) was performed to

synchronize infection. Phagocytes were infected for 45 min at 37°C and under a 5% CO_2 atmosphere

in RPMI 1640 medium with GBS strains, using a multiplicity of infection (MOI) of 10 bacteria per cell

except when otherwise specified. After infection, phagocytes were washed 3 times with PBS.

Remaining extracellular bacteria were killed using RPMI 1640 medium supplemented with Penicillin G

/ Streptomycin (100 units.mL^{-1} / 0.1 mg.mL^{-1}) and Gentamicin (0.2 mg.mL^{-1}) at 37°C and 5% CO_2. After

antibiotic treatment, cells were washed twice and either lysed directly by osmotic choc using sterile

ice-cold H_2O to evaluate phagocytosis (T0) or further incubated in RPMI1640 medium to follow

intracellular survival before being lysed (2h30 or 24h). Serial dilutions of the lysates were plated onto

TH agar for colony forming units (CFU) counting. The percentage of phagocytosis was calculated as

follows: (CFU phagocytosis / CFU in inoculum) × 100. The percentage of survival was calculated as

follows: (CFU survival / CFU phagocytosis) × 100. When specified, results were expressed as normalized

to the control strain (BM110). Assays were performed in triplicate and were repeated at least three

times.

Adhesion assay onto monocytes and macrophages

For adhesion assays, infection was performed as described for phagocytosis but at 4°C and with ice-

cold RPMI1640 medium to avoid engulfment of bacteria and phagocytosis. In some experiments,

adhesion was also tested at 37°C using cells that had been pre-treated for 1 hour with cytochalasin D.

After 3 washes with ice-cold PBS to remove non-adherent bacteria, phagocytes were directly lysed by

osmotic shock with ice-cold H_2O. Serial dilutions of lysates were then plated onto TH agar for CFU

counting. The percentage of adhesion was determined by calculating the ratio between the CFUs of

adherent bacteria and the CFUs from the inoculum × 100. When specified, results were expressed as

normalized to the control strain. Assays were performed in triplicate and were repeated at least three

times.

Phagocytosis assay in the presence of inhibitors/reagents

Selected compounds were tested for their ability to block GBS phagocytosis. All inhibitors and reagents

used in this study are listed in Table S3. THP-1 macrophages were pretreated with inhibitors/reagents

for 1h prior infection with BM110 or NEM316 strains. Inhibitors/reagents were left during the time of

infection at the same concentration. Treatments did not affect bacterial or cell viability as assessed by

Lactate dehydrogenase release in the culture supernatant (LDH cytotoxicity kit®, Promega, according

to the manufacturer's instructions; data not shown). The percentage of phagocytosis was calculated

as follows: (CFU phagocytosis / CFU in inoculum) × 100 and results were normalized to the untreated

condition.

Evaluation of bacterial egress from macrophages

To assess bacterial egress from macrophages, cells were infected as described previously, and then

treated with antibiotics to kill extracellular bacteria for 45 min. Cells culture medium was next replaced

by medium without any antibiotic. Bacterial egress was quantified in the supernatant of infected

macrophages after 45 min or after 1h30 following the replacement of the cell culture medium. The

percent of bacterial egress was calculated as follows: (CFU released bacteria / CFU phagocytosed) ×

100.

RNA extraction, cDNA synthesis and RT-PCR

RNAs were extracted from non-infected THP-1 macrophages, monocytes and hMDM using the

NucleoSpin® RNA kit (Macherey-Nagel®) according to the manufacturer's instructions. RNA

concentration was determined using a NanoDrop 2000 spectrophotometer. Then, 1 µg RNA was

reverse-transcribed into cDNA using the SuperScript® IV Reverse Transcription Kit (Invitrogen),

following the manufacturer's instructions. PCR were carried out with Green Taq DNA polymerase

(Promega) on cDNA using primers described in Table S4. Primers were designed in order to amplify all

variants of the targeted receptor(s). Amplification products were resolved in 1% (w/v) agarose gel.

siRNA transfection

To silence the expression of putative phagocytic receptors, pools of three or four siRNA duplexes

(Dharmacon) were used. The silencer select-negative control-1 siRNA (Ambion) was used as negative

control. THP-1 were seeded in twenty-four-well plates for 24 h in the presence of 200 nM of PMA then

transfected with 50 nM siRNA using Lipofectamine RNAIMAX (Invitrogen) according to the

manufacturer's instructions and incubated for a further 3 days. Efficiency of knockdown was assessed

by Western Blot analysis on 2 independent targets.

Immunofluorescent staining and image analysis

To follow intracellular fate of GBS strains, THP-1 macrophages were infected for 45 minutes as

described for phagocytosis then washed extensively and fixed at different time points post-

phagocytosis (0, 2h30 or 5h) on glass coverslips with 4% paraformaldehyde (PFA) for 10 min then

washed with PBS. Following fixation, samples were permeabilized and saturated with PBS containing

0.2% saponin and 10% serum for 1h at room temperature, then incubated for at least 1h at room

temperature with the primary anti-LAMP1 antibody (DSHB clone H4A3). After 3 washes with PBS,

samples were incubated for 1h at room temperature with conjugated-secondary antibodies (Jackson

ImmunoResearch Laboratories).

To enable detection of acidic compartments (phagolysosome), THP-1 macrophages were incubated 30

min before the end of infection with Lysotracker red DND-99 (Invitrogen) at a final concentration of 75

nM, fixed with 4% PFA for 10 min, then washed with PBS.

After 3 washes in PBS, coverslips were mounted in DAKO fluorescent mounting media (DAKO). Image

acquisitions were realized using confocal laser scanning microscopy (Leica DMI6000) coupled to a

spinning disk confocal head (YokogawaCSU-X1M1), controlled by the Metamorph 7.7.5 software

(Cochin Institute microscopy facility, IMAG'IC). Images were processed using the ImageJ software. At

least 300 cells were analyzed for each sample in four independent experiments.

Scanning Electron Microscopy

All incubations were performed at room temperature except when otherwise indicated. Macrophages

grown on glass coverslips were infected as described previously and then fixed in 0.1M sodium

cacodylate buffer, pH 7.4 containing 2.5% glutaraldehyde and 1% PFA for 60 min. After washing in

cacodylate buffer (2x10 min), cells were fixed in 1% OsO_4 diluted in cacodylate buffer for 45 min at 4°C.

After washing in cacodylate buffer (2x10 min), samples were dehydrated in an ascending series of

ethanol (30%, 50%, 70%, 95%, 100%, 100%, 100% - 10 min each), followed by hexamethyldisilazane

/ethanol (1/1) for 10 min and hexamethyldisilazane for 10 min. After overnight air drying, each

coverslip was placed on a double-sided sticky tape on the top of an aluminum stub and sputter coated

with Au/Pd. Images were acquired using a Jeol LV6510 (Jeol, Croissy-sur-Seine, France).

Statistical analysis

All assays were performed in at least in three independent experiments, each experiment was carried

out in triplicate. Data represent mean ± SEM or median and statistical analyses were performed using

GraphPad Prism 10.0 (GraphPad Software, San Diego, California). Significance levels were set at *$p \leq$

0.05; **$p \leq 0.01$; ***$p \leq 0.001$; **** $p \leq 0.0001$.

AUTHOR CONTRIBUTIONS

JG designed the research. ASB, AP, JCF, CH and JG performed the experiments. ASB, AP, JG analyzed the data. BS and VM performed electron microscopy experiments. AF, AT, CP provided intellectual input and guidance. JG wrote the manuscript and ASB, AF, AT contributed to finalizing the manuscript. All authors discussed results and reviewed the final version of the manuscript.

ACKNOWLEDGMENTS

We are grateful to Glen Ulett for the gift of the GFP plasmid. We thank Cristina Candeias for technical assistance, Shaynoor Dramsi for BM110Δ*spb1* and NEM316Δ*cps* mutant strains and the Imag'IC core facility of the Cochin Institute. The H4A3 antibody developed by August, J.T. / Hildreth, J.E.K. was obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242.

FINANCIAL SUPPORT

This work was supported by Agence Nationale de la Recherche (ANR) (Grant StrepB2brain, ANR-17-CE15-0026-01). ASB was a doctoral fellow funded by Université Paris Cité.

REFERENCES

1. Marodi L. 2006. Neonatal innate immunity to infectious agents. Infect Immun 74:1999-2006.
2. Mittal R, Sukumaran SK, Selvaraj SK, Wooster DG, Babu MM, Schreiber AD, Verbeek JS, Prasadarao NV. 2010. Fcgamma receptor I alpha chain (CD64) expression in macrophages is critical for the onset of meningitis by Escherichia coli K1. PLoS Pathog 6:e1001203.
3. Orrskog S, Rounioja S, Spadafina T, Gallotta M, Norman M, Hentrich K, Falker S, Ygberg-Eriksson S, Hasenberg M, Johansson B, Uotila LM, Gahmberg CG, Barocchi M, Gunzer M, Normark S, Henriques-Normark B. 2013. Pilus adhesin RrgA interacts with complement receptor 3, thereby affecting macrophage function and systemic pneumococcal disease. MBio 4:e00535-12.
4. Rosales C, Uribe-Querol E. 2017. Phagocytosis: A Fundamental Process in Immunity. Biomed Res Int 2017:9042851.
5. Patras KA, Nizet V. 2018. Group B Streptococcal Maternal Colonization and Neonatal Disease: Molecular Mechanisms and Preventative Approaches. Front Pediatr 6:27.
6. Seale AC, Bianchi-Jassir F, Russell NJ, Kohli-Lynch M, Tann CJ, Hall J, Madrid L, Blencowe H, Cousens S, Baker CJ, Bartlett L, Cutland C, Gravett MG, Heath PT, Ip M, Le Doare K, Madhi SA, Rubens CE, Saha SK, Schrag SJ, Sobanjo-Ter Meulen A, Vekemans J, Lawn JE. 2017. Estimates of the Burden of Group B Streptococcal Disease Worldwide for Pregnant Women, Stillbirths, and Children. Clin Infect Dis 65:S200-S219.
7. Tazi A, Plainvert C, Anselem O, Ballon M, Marcou V, Seco A, El Alaoui F, Joubrel C, El Helali N, Falloukh E, Frigo A, Raymond J, Trieu-Cuot P, Branger C, Le Monnier A, Azria E, Ancel PY, Jarreau PH, Mandelbrot L, Goffinet F, Poyart C. 2019. Risk Factors for Infant Colonization by Hypervirulent CC17 Group B *Streptococcus*: Toward the Understanding of Late-onset Disease. Clin Infect Dis doi:ciz033 [pii] 10.1093/cid/ciz033 5427769 [pii].
8. Hays C, Touak G, Bouaboud A, Fouet A, Guignot J, Poyart C, Tazi A. 2019. Perinatal hormones favor CC17 group B *Streptococcus* intestinal translocation through M cells and hypervirulence in neonates. Elife 8.
9. Madrid L, Seale AC, Kohli-Lynch M, Edmond KM, Lawn JE, Heath PT, Madhi SA, Baker CJ, Bartlett L, Cutland C, Gravett MG, Ip M, Le Doare K, Rubens CE, Saha SK, Sobanjo-Ter Meulen A, Vekemans J, Schrag S, Infant GBSDIG. 2017. Infant Group B Streptococcal Disease Incidence and Serotypes Worldwide: Systematic Review and Meta-analyses. Clin Infect Dis 65:S160-S172.
10. Joubrel C, Tazi A, Six A, Dmytruk N, Touak G, Bidet P, Raymond J, Trieu Cuot P, Fouet A, Kerneis S, Poyart C. 2015. Group B *Streptococcus* neonatal invasive infections, France 2007-2012. Clin Microbiol Infect 21:910-6.
11. Vuillemin X, Hays C, Plainvert C, Dmytruk N, Louis M, Touak G, Saint-Pierre B, Adoux L, Letourneur F, Frigo A, Poyart C, Tazi A. 2021. Invasive group B *Streptococcus* infections in non-pregnant adults: a retrospective study, France, 2007-2019. Clin Microbiol Infect 27:129 e1-129 e4.
12. Manning SD, Springman AC, Lehotzky E, Lewis MA, Whittam TS, Davies HD. 2009. Multilocus sequence types associated with neonatal group B streptococcal sepsis and meningitis in Canada. J Clin Microbiol 47:1143-8.
13. Teatero S, McGeer A, Low DE, Li A, Demczuk W, Martin I, Fittipaldi N. 2014. Characterization of invasive group B *Streptococcus* strains from the greater Toronto area, Canada. J Clin Microbiol 52:1441-7.
14. van Kassel M, de Boer G, Teeri S, Jamrozy D, Bentley S, Brouwer M, Van der Ende A, Van de Beek D, Bijlsma M. 2021. Molecular epidemiology and mortality of group B streptococcal meningitis and infant sepsis in the Netherlands: a 30-year nationwide surveillance study. The lancet microbe 2.
15. Brochet M, Couve E, Zouine M, Vallaeys T, Rusniok C, Lamy MC, Buchrieser C, Trieu-Cuot P, Kunst F, Poyart C, Glaser P. 2006. Genomic diversity and evolution within the species *Streptococcus agalactiae*. Microbes Infect 8:1227-43.

16. Tazi A, Disson O, Bellais S, Bouaboud A, Dmytruk N, Dramsi S, Mistou MY, Khun H, Mechler C,
Tardieux I, Trieu-Cuot P, Lecuit M, Poyart C. 2010. The surface protein HvgA mediates group B
Streptococcus hypervirulence and meningeal tropism in neonates. J Exp Med 207:2313-22.
17. Deshayes de Cambronne R, Fouet A, Picart A, Bourrel AS, Anjou C, Bouvier G, Candeias C,
Bouaboud A, Costa L, Boulay AC, Cohen-Salmon M, Plu I, Rambaud C, Faurobert E, Albiges-Rizo
C, Tazi A, Poyart C, Guignot J. 2021. CC17 group B *Streptococcus* exploits integrins for neonatal
meningitis development. J Clin Invest 131.
18. Springman AC, Lacher DW, Wu G, Milton N, Whittam TS, Davies HD, Manning SD. 2009.
Selection, recombination, and virulence gene diversity among group B streptococcal
genotypes. J Bacteriol 191:5419-27.
19. Korir ML, Manning SD, Davies HD. 2017. Intrinsic Maturational Neonatal Immune Deficiencies
and Susceptibility to Group B *Streptococcus* Infection. Clin Microbiol Rev 30:973-989.
20. Chang YC, Nizet V. 2020. Siglecs at the Host-Pathogen Interface. Adv Exp Med Biol 1204:197-
214.
21. Marques MB, Kasper DL, Pangburn MK, Wessels MR. 1992. Prevention of C3 deposition by
capsular polysaccharide is a virulence mechanism of type III group B streptococci. Infect
Immun 60:3986-93.
22. Henneke P, Takeuchi O, Malley R, Lien E, Ingalls RR, Freeman MW, Mayadas T, Nizet V, Akira
S, Kasper DL, Golenbock DT. 2002. Cellular activation, phagocytosis, and bactericidal activity
against group B *Streptococcus* involve parallel myeloid differentiation factor 88-dependent
and independent signaling pathways. J Immunol 169:3970-7.
23. Valentin-Weigand P, Benkel P, Rohde M, Chhatwal GS. 1996. Entry and intracellular survival of
group B streptococci in J774 macrophages. Infect Immun 64:2467-73.
24. Cornacchione P, Scaringi L, Fettucciari K, Rosati E, Sabatini R, Orefici G, von Hunolstein C,
Modesti A, Modica A, Minelli F, Marconi P. 1998. Group B streptococci persist inside
macrophages. Immunology 93:86-95.
25. Cumley NJ, Smith LM, Anthony M, May RC. 2012. The CovS/CovR acid response regulator is
required for intracellular survival of group B *Streptococcus* in macrophages. Infect Immun
80:1650-61.
26. Sullivan MJ, Prince D, Goh KGK, Katupitiya L, Gosling D, Crowley MR, Crossman DK, Ulett GC.
2023. Dual RNA sequencing of group B *Streptococcus*-infected human monocytes reveals new
insights into host-pathogen interactions and bacterial evasion of phagocytosis. Sci Rep
13:2137.
27. Chattopadhyay D, Carey AJ, Caliot E, Webb RI, Layton JR, Wang Y, Bohnsack JF, Adderson EE,
Ulett GC. 2011. Phylogenetic lineage and pilus protein Spb1/SAN1518 affect opsonin-
independent phagocytosis and intracellular survival of Group B *Streptococcus*. Microbes Infect
13:369-82.
28. Perichon B, Guignot J, Szili N, Gao C, Poyart C, Trieu-Cuot P, Dramsi S. 2019. Insights into
Streptococcus agalactiae PI-2b pilus biosynthesis and role in adherence to host cells. Microbes
Infect 21:99-103.
29. Six A, Bellais S, Bouaboud A, Fouet A, Gabriel C, Tazi A, Dramsi S, Trieu-Cuot P, Poyart C. 2015.
Srr2, a multifaceted adhesin expressed by ST-17 hypervirulent Group B *Streptococcus* involved
in binding to both fibrinogen and plasminogen. Mol Microbiol 97:1209-22.
30. Rogers LM, Gaddy JA, Manning SD, Aronoff DM. 2018. Variation in Macrophage Phagocytosis
of *Streptococcus agalactiae* Does Not Reflect Bacterial Capsular Serotype, Multilocus
Sequence Type, or Association with Invasive Infection. Pathog Immun 3:63-71.
31. Korir ML, Laut C, Rogers LM, Plemmons JA, Aronoff DM, Manning SD. 2017. Differing
mechanisms of surviving phagosomal stress among group B *Streptococcus* strains of varying
genotypes. Virulence 8:924-937.
32. Flaherty RA, Aronoff DM, Gaddy JA, Petroff MG, Manning SD. 2021. Distinct Group B
Streptococcus Sequence and Capsule Types Differentially Impact Macrophage Stress and
Inflammatory Signaling Responses. Infect Immun 89.

33. Perichon B, Szili N, du Merle L, Rosinski-Chupin I, Gominet M, Bellais S, Poyart C, Trieu-Cuot P,
Dramsi S. 2017. Regulation of PI-2b Pilus Expression in Hypervirulent *Streptococcus agalactiae*
ST-17 BM110. PLoS One 12:e0169840.
34. Uribe-Querol E, Rosales C. 2020. Phagocytosis: Our Current Understanding of a Universal
Biological Process. Front Immunol 11:1066.
35. Krugmann S, Jordens I, Gevaert K, Driessens M, Vandekerckhove J, Hall A. 2001. Cdc42 induces
filopodia by promoting the formation of an IRSp53:Mena complex. Curr Biol 11:1645-55.
36. Pombinho R, Sousa S, Cabanes D. 2018. Scavenger Receptors: Promiscuous Players during
Microbial Pathogenesis. Crit Rev Microbiol 44:685-700.
37. PrabhuDas MR, Baldwin CL, Bollyky PL, Bowdish DME, Drickamer K, Febbraio M, Herz J, Kobzik
L, Krieger M, Loike J, McVicker B, Means TK, Moestrup SK, Post SR, Sawamura T, Silverstein S,
Speth RC, Telfer JC, Thiele GM, Wang XY, Wright SD, El Khoury J. 2017. A Consensus Definitive
Classification of Scavenger Receptors and Their Roles in Health and Disease. J Immunol
198:3775-3789.
38. Adderson EE, Takahashi S, Wang Y, Armstrong J, Miller DV, Bohnsack JF. 2003. Subtractive
hybridization identifies a novel predicted protein mediating epithelial cell invasion by virulent
serotype III group B *Streptococcus agalactiae*. Infect Immun 71:6857-63.
39. Sharma P, Lata H, Arya DK, Kashyap AK, Kumar H, Dua M, Ali A, Johri AK. 2013. Role of pilus
proteins in adherence and invasion of *Streptococcus agalactiae* to the lung and cervical
epithelial cells. J Biol Chem 288:4023-34.
40. Seo HS, Minasov G, Seepersaud R, Doran KS, Dubrovska I, Shuvalova L, Anderson WF, Iverson
TM, Sullam PM. 2013. Characterization of fibrinogen binding by glycoproteins Srr1 and Srr2 of
Streptococcus agalactiae. J Biol Chem 288:35982-96.
41. Doyle SE, O'Connell RM, Miranda GA, Vaidya SA, Chow EK, Liu PT, Suzuki S, Suzuki N, Modlin
RL, Yeh WC, Lane TF, Cheng G. 2004. Toll-like receptors induce a phagocytic gene program
through p38. J Exp Med 199:81-90.
42. Amiel E, Alonso A, Uematsu S, Akira S, Poynter ME, Berwin B. 2009. Pivotal Advance: Toll-like
receptor regulation of scavenger receptor-A-mediated phagocytosis. J Leukoc Biol 85:595-605.
43. Albanyan EA, Vallejo JG, Smith CW, Edwards MS. 2000. Nonopsonic binding of type III Group B
Streptococci to human neutrophils induces interleukin-8 release mediated by the p38
mitogen-activated protein kinase pathway. Infect Immun 68:2053-60.
44. Lee HJ, Woo Y, Hahn TW, Jung YM, Jung YJ. 2020. Formation and Maturation of the
Phagosome: A Key Mechanism in Innate Immunity against Intracellular Bacterial Infection.
Microorganisms 8.
45. Taban Q, Mumtaz PT, Masoodi KZ, Haq E, Ahmad SM. 2022. Scavenger receptors in host
defense: from functional aspects to mode of action. Cell Commun Signal 20:2.
46. Lin Z, Tan X, Zhang Y, Li F, Luo P, Liu H. 2020. Molecular Targets and Related Biologic Activities
of Fucoidan: A Review. Mar Drugs 18.
47. Sigola LB, Fuentes AL, Millis LM, Vapenik J, Murira A. 2016. Effects of Toll-like receptor ligands
on RAW 264.7 macrophage morphology and zymosan phagocytosis. Tissue Cell 48:389-96.
48. Baba M, Snoeck R, Pauwels R, de Clercq E. 1988. Sulfated polysaccharides are potent and
selective inhibitors of various enveloped viruses, including herpes simplex virus,
cytomegalovirus, vesicular stomatitis virus, and human immunodeficiency virus. Antimicrob
Agents Chemother 32:1742-5.
49. Thelen T, Hao Y, Medeiros AI, Curtis JL, Serezani CH, Kobzik L, Harris LH, Aronoff DM. 2010. The
class A scavenger receptor, macrophage receptor with collagenous structure, is the major
phagocytic receptor for *Clostridium sordellii* expressed by human decidual macrophages. J
Immunol 185:4328-35.
50. Cardoso MS, Santos RF, Almeida S, Sa M, Perez-Cabezas B, Oliveira L, Tavares J, Carmo AM.
2021. Physical Interactions With Bacteria and Protozoan Parasites Establish the Scavenger
Receptor SSC4D as a Broad-Spectrum Pattern Recognition Receptor. Front Immunol
12:760770.

51.   Dutot M, Grassin-Delyle S, Salvator H, Brollo M, Rat P, Fagon R, Naline E, Devillier P. 2019. A
667   marine-sourced fucoidan solution inhibits Toll-like-receptor-3-induced cytokine release by
668   human bronchial epithelial cells. Int J Biol Macromol 130:429-436.
669   52.   Shabayek S, Spellerberg B. 2017. Acid Stress Response Mechanisms of Group B Streptococci.
670   Front Cell Infect Microbiol 7:395.
671   53.   Daigneault M, Preston JA, Marriott HM, Whyte MK, Dockrell DH. 2010. The identification of
672   markers of macrophage differentiation in PMA-stimulated THP-1 cells and monocyte-derived
673   macrophages. PLoS One 5:e8668.

Figure 1: CC17 GBS strains are more phagocytosed by monocytes and macrophages than non-CC17 GBS strains. (A, B, C, D) Phagocytosis level of GBS strains was assessed by CFU count after infection at MOI 10 followed by antibiotic treatment to kill extracellular bacteria using (A) THP-1 macrophages (B) hMDM primary cells, (C) THP-1 monocytes or (D) human primary monocytes. (A) Phagocytes were infected with CC17 and non-CC17 GBS clinical isolates from invasive infections, each dot representing a clinical strain. Triangles correspond to BM110 strain (CC17) and NEM316 (non-CC17). Results are expressed as the percentage of phagocytosis relative to BM110 strain phagocytosis, with horizontal lines indicating median value. (B, C, D) Phagocytes were infected with GBS strains BM110 and NEM316. Results are expressed as the percentage of phagocytosed bacteria normalized to (B, C) the initial inoculum or (D) to BM110 strain phagocytosis.

686   Statistical analysis: data shown are (A) median or (B, C, D) mean ± SEM of at least four independent

687   experiments. (A, B, C) t test or (D) Mann Whitney test, were performed with *, $p < 0.05$; **, $p < 0.01$;

688   ****, $p < 0.0001$.

Figure 2: GBS capsule is not involved in the hyper phagocytosis of CC17 strains. (A, B) Phagocytosis level of GBS strains was assessed by CFU count after infection at MOI 10, followed by antibiotic treatment to kill extracellular bacteria. (A) THP-1 macrophages were infected with CC17 and non-CC17 GBS clinical isolates from invasive infections, each dot representing a clinical strain. Triangles correspond to BM110 strain (CC17) and NEM316 (non-CC17). Results are shown according to capsular type (CPS III, CPS IV, or other CPSs) and are expressed as the percentage of phagocytosis relative to BM110 strain phagocytosis, with horizontal lines indicating median value. (B) THP-1 macrophages were infected with GBS strains BM1110 and NEM316 and their respective mutant strains impaired for

700   capsule expression (Δ*cps*). Results are expressed as the percentage of phagocytosed bacteria

701   normalized to the initial inoculum.

702   Statistical analysis: data shown are (A) median or (B) mean ± SEM of at least four independent

703   experiments. (A, B) Two-way ANOVA, was performed with ns, non-significant; **, $p < 0.01$; ****, $p <$

704   0.0001.

Figure 3: The HvgA surface protein and Spb1 pilus protein are involved in CC17 hyper-phagocytosis.

(A, B, C) Phagocytosis level of GBS strains was assessed by CFU count after infection at MOI 10 followed by antibiotic treatment to kill extracellular bacteria using (A, B) THP-1 macrophages (C) hMDM primary cells. (A, C) Macrophages were infected with BM110 and its derivative mutant strains (Δ*srr2*, Δ*hvgA* or Δ*spb1*), or (B) with *Lactococcus lactis* strain carrying an empty vector (+vector) or a vector expressing the HvgA protein (+HvgA). (A, C) Results are expressed as the relative levels to BM110 strain, or (B) to the control *Lactococcus lactis* strain +vector.

Statistical analysis: data shown are mean ± SEM of at least four independent experiments. (A) Kruskall Wallis test with Dunn's multiple comparisons or (B, C) t test, were performed with ns, non-significant; *, $p < 0.05$; ***, $p < 0.001$; ****, $p < 0.0001$.

Figure 4: Signal transduction leading to phagocytosis is similar for CC17 and non-CC17 strains. (A, B, C, D) THP-1 macrophages were pre-treated for one-hour prior infection with cytochalasin D, ZCL178, staurosporin, LY29002, PP2, nocodazole, NSC23766, BAY61-3606 or Y27632 to inhibit actin polymerization, CDC42, protein kinase C, PI3kinase, Src kinase, microtubule polymerization, Rac, Syc and Rock, respectively. (A) Phagocytosis level of BM110 (B) NEM316 (C) Δ*spb1* or (D) Δ*hvgA* GBS strains

were assessed by CFU count after infection at MOI 10 followed by antibiotic treatment to kill

extracellular bacteria. Results are expressed as the relative levels to untreated condition. (E) Scanning

electron microscopy (SEM) micrographs of THP-1 macrophages showing (left panel, scale bar 5 μm)

bacterial adherence to cell surface and (middle panel and right panels, scale bar 1 μm) bacterial capture

by filopodia after 15 min or lamellipodia after30 min of infection.

Statistical analysis: data shown are mean ± SEM of at least four independent experiments. One-Way

ANOVA with Dunnett multiple comparisons, were performed with ns, non-significant; **, $p < 0.01$; ***,

$p < 0.001$, ****, $p < 0.0001$.

Figure 5: CC17 GBS strains are more adherent to phagocytes than non-CC17 strains. (A, B, C, D, E, F)
Adhesion level of GBS strains was assessed by CFU count after infection at MOI 10 at 4°C to avoid
bacterial engulfment using (A, D, E, F) THP-1 macrophages (B) THP-1 monocytes (C) hMDM primary
cells. (A) Phagocytes were infected with CC17 and non-CC17 GBS clinical isolates from invasive

infections, each dot representing a clinical strain. Triangles correspond to BM110 strain (CC17) and

NEM316 (non-CC17). Results are expressed as the percentage of adhesion relative to BM110 strain

adhesion, with horizontal lines indicating median value. (B, C, D, E, G, H) Phagocytes were infected with

GBS strains or (F) with *Lactococcus lactis* strain carrying an empty vector (+vector) or a vector

expressing HvgA protein (+HvgA). Results were normalized to (B, C, D) the initial inoculum (E) BM110

strain or (F) *L. lactis* control strain (+vector). (G,H) (G) Adhesion and phagocytosis were assessed by

CFU count after infection at MOI 10. Results were expressed as the ratio between CFU of phagocytosed

bacteria and adherent bacteria. (H) Adhesion and phagocytosis were assessed by CFU count after

infection at MOI 10 or 2. Results were normalized to the initial inoculum.

Statistical analysis: data shown are mean ± SEM of at least four independent experiments. (A, B, C, D,

F) t Test (E) Kruskall Wallis test with Dunn's multiple comparisons or (G, H) One-Way ANOVA with

Tukey's multiple comparisons test, were performed with ns, non-significant; *, $p < 0.05$; **, $p < 0.01$;

***, $p < 0.001$; ****, $p < 0.0001$.

Figure 6: Scavenger receptor inhibitors lead to a decreased phagocytosis level of CC17 GBS in macrophages. (A, B, E) Phagocytosis level in THP-1 macrophages was assessed by CFU count after infection at MOI 10 with (A, E) BM110 or (B) NEM 316 GBS strains followed by 1h antibiotic treatment to kill extracellular bacteria. (A) Macrophages were pre-treated 1h before infection to block FCγ receptors (Fc Block™), complement receptor 3 (CR3 blocking antibody), lectin receptors (mannose, mannan, laminarin), integrins receptors (RGD and RGDfV mimetic peptides), or (A, B) scavenger receptors inhibitors (Fucoidan and poly(I) or inactive analog poly(C). (A, B) Results are expressed as the percentage of phagocytosis relative to the untreated control condition except for poly (I) that was

relative to its inactive analogue poly(C). (C) Adhesion level in untreated or fucoidan treated-THP-1

macrophages was assessed by CFU count after infection with BM110 and NEM316 at 4°C to avoid

bacterial engulfment. Results are expressed as the percentage of phagocytosis relative to the

untreated control condition (D) Expression of scavenger receptors genes was assessed by RT-PCR using

THP-1 macrophages (macro), THP-1 monocyte (mono) and hMDM (hMDM) cDNA. Actin and VE-

cadherin gene expression was used as positive and negative controls of expression, respectively.

Genomic DNA (gDNA) was used to validate primers efficiency. (E) Involvement of scavenger receptors

on BM110 phagocytosis was assessed by silencing scavenger receptors expression by si-RNA. Results

are expressed as the percentage of phagocytosis relative to the scramble control condition.

Statistical analysis: data shown are mean ± SEM of at least four independent experiments. (A, B, E)

Kruskall Wallis test with Dunn's multiple comparisons or (C) t test were performed, with ns, non-

significant; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Figure 7: CC17 and non-CC17 GBS strains exhibit similar intracellular fate. (A, B, C, D) Intracellular fate of BM110 and NEM316 strains was followed by microscopy in THP-1 macrophages. (A, C) Representative confocal microscopy images of THP-1 macrophages infected with BM110-GFP (green) and labelled with (A) anti-LAMP1 antibody (red) or (C) Lysotracker® (red). Boxed areas correspond to magnification of insets. Dotted lines indicate cell perimeters. Scale bar: 5 μm. (B, D) Quantification of BM110-GFP and NEM316-GFP associated with (B) LAMP1 or (D) Lysotracker® was performed at 0, 2h30

or 5h post-phagocytosis. Results are expressed as the percentage of bacteria positive for (B) LAMP1 or

(D) Lysotracker® staining. (E, F, G) Percentage of bacterial survival was assessed by CFU count 2h30

after phagocytosis in (E, F) THP-1 macrophage or (G) hMDM. (E) Macrophages were infected with CC17

and non-CC17 GBS clinical isolates from invasive infections, each dot representing a clinical strain. For

each strain, results are expressed as the percentage of survival relative to corresponding the

corresponding level of phagocytosis, with horizontal lines indicating median value. Triangles

correspond to BM110 strain (CC17) and NEM316 (non-CC17) (F, G) THP-1 macrophages were infected

with the BM110 strain and its derivative mutant strains (Δ*hvgA* or Δ*spb1*)*.* Results are expressed as the

percentage of viable intracellular bacteria normalized to the initial intracellular load (phagocytosis).

Statistical analysis: data shown are mean ± SEM of at least four independent experiments. (B, D) Two-

way ANOVA with Tuckey multiple comparison, (E) t Test or (F) Kruskall Wallis test with Dunn's multiple

comparisons or (G) One-Way ANOVA tests were performed with ns, non-significant.

Fig 8: GBS strains can escape macrophages independently of actin and microtubules. (A, B) THP-1
were infected with GBS strains at MOI 10 then treated with antibiotic to kill extracellular bacteria.
Bacterial egress was measured by CFU count in cell supernatant devoid of antibiotic after (A) 45 min
or 1h30 or (B) 45 min post-antibiotic removal. Results are expressed as the percentage of viable
bacteria released in the cell supernatants normalized to the number of phagocytosed bacteria.
Bacterial egress was measured for (A) BM110 or NEM316 strains in the absence of treatment or (B)
BM110 in the presence of actin (cytochalasin D) or microtubules (nocodazole) depolymerizing agents
added after phagocytosis.

Statistical analysis: data shown are mean ± SEM of at least four independent experiments. (A) Two-
way ANOVA with Dunnett's multiple comparison (B) one-Way ANOVA with Dunnett's multiple
comparisons tests were performed with ns, non-significant.