An Autoimmune Transcriptional Circuit Driving Foxp3[+] Regulatory T cell Dysfunction

Tomokazu S. Sumida[1,2,8], Matthew R. Lincoln[1,2,8], Liang He[2,3,4,8], Yongjin Park[2,3, 8], Mineto Ota[6],

Helen A. Stillwell[1], Greta A. Leissa[1], Keishi Fujio[6], Alexander M. Kulminski[4], Charles B. Epstein[2],

Bradley E. Bernstein[2,7], Manolis Kellis[2,3], David A. Hafler[1,2, †]

[1]Departments of Neurology and Immunobiology, Yale School of Medicine, New Haven, CT, USA

[2]Broad Institute of MIT and Harvard, Cambridge, MA, USA

[3]Computer Science and Artificial Intelligence Laboratory, MIT, Cambridge, MA, USA

[4]Biodemography of Aging Research Unit, Social Science Research Institute, Duke University, Durham, NC, USA

[6]Department of Allergy and Rheumatology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

[7]Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA, USA.

[8]These authors contributed equally

[†] Correspondence to David Hafler (david.hafler@yale.edu)

Present Address:

M.R.L.: Division of Neurology, Department of Medicine, University of Toronto, and Keenan Research Centre for Biomedical Science of St. Michael's Hospital, Toronto, ON, Canada

L.H.: Department of Public Health, Faculty of Health Sciences, University of Southern Denmark, Odense M, Denmark

Y.P.: Department of Pathology and Laboratory Medicine, Department of Statistics, University of British Columbia, Vancouver, BC, Canada

Abstract

Autoimmune diseases, among the most common disorders of young adults, are mediated by genetic and environmental factors. While CD4$^+$Foxp3$^+$ regulatory T cells (Tregs) play a central role in preventing autoimmunity, the molecular mechanism underlying their dysfunction is unknown. Here, we performed comprehensive transcriptomic and epigenomic profiling of Tregs in the autoimmune disease multiple sclerosis (MS) to identify central transcriptional programs regulating human autoimmunity. We discovered that upregulation of a primate-specific short *PRDM1* isoform (*PRDM1-S*) induces *SGK1* independent from evolutionally conserved long *PRDM1*, leading to destabilization of Foxp3 and Treg dysfunction. This aberrant *PRDM1-S/SGK1* axis is shared among other autoimmune diseases. Furthermore, by chromatin landscape profiling in MS Tregs we identified a *PRDM1-S* specific *cis*-regulatory element associated with enriched binding of AP-1/IRF transcription factors. Our study identifies evolutionally emerged *PRDM1-S* and epigenetic priming of AP-1/IRF as key drivers of pathogenic Treg programs leading to human autoimmune disease.

(146 words)

Introduction

Human autoimmune diseases constitute a leading cause of death among young adults with

an increasing incidence in recent years. Multiple sclerosis (MS) is a canonical, genetically mediated

autoimmune disease induced by environmental factors where genetic perturbation of *cis*-regulatory

elements in pathogenic immune cells leads to immune dysregulation and generation of

autoreactive T cells and antibodies[1-3]. Among immune cells, CD4[+] T cells play a central role in both

mediating and regulating autoimmunity. As CD4[+] T cells display a large degree of functional

diversity, interrogation of total CD4[+] T cell populations has not to date identified causal

transcriptional changes[4,5]. Thus, we hypothesized that interrogation of CD4[+] T cell subpopulations

is required to elucidate the pathophysiological characteristics of CD4[+] T cells in autoimmune

disorders, allowing identification of central transcriptional factors associated with loss of immune

regulation

Human CD4[+] Foxp3[+] regulatory T cells (Tregs) play a central role in the maintenance of

immune homeostasis and prevention of autoimmunity[6-8]. We first demonstrated that Tregs from

patients with autoimmune disease exhibit a dysfunctional phenotype[9-11] and this has been

subsequently found among multiple autoimmune disorders. Recent evidence has shown that

environmental factors, such as vitamin D and fatty acids affect Treg phenotype and function. In

addition, high salt intake has been epidemiologically linked with autoimmune diseases[12] and higher

physiologic salt concentrations induce proinflammatory Th17 cells mediated by SGK1[13] and

modulate the stability of Tregs[14,15], resembling the phenotype observed in autoimmune diseases

including MS[16]. These environmental factors are known to act, in part, through epigenetic changes

that can be identified by examining the histone and methylation landscape of cells[17]. A recent

investigation of the genetic architecture of 20 autoimmune diseases and MS showed that causal

variants are enriched in regulatory elements that are active in immune cells[3]. Moreover, many of

these genetic loci are common among multiple autoimmune diseases, implicating the shared

immunomodulatory mechanism in human autoimmunity[18,19]. These findings, together with

increases in the incidence of autoimmune diseases over the past three decades that cannot be

explained by genetic factors alone, points to both genetic and epigenetic factors as key mediators

of risk for autoimmunity. While these lines of evidence strongly suggest that both genetic and

epigenetic alterations might disturb Treg homeostasis, the underlying mechanism in Treg

dysfunction in human autoimmune diseases has not been elucidated.

We interrogated the phenotypic and functional characteristics of human CD4[+] T cells,

focusing primarily on memory Treg (mTreg) through comprehensive transcriptomic and epigenetic

profiling. We adopted both bulk and single-cell RNA-seq (scRNA-seq) for transcriptomic profiling

and Assay for Transposase-Accessible Chromatin using sequencing (ATAC-seq) for probing

epigenetic regulation to understand the molecular mechanisms that drive dysfunctional programs

in MS Tregs. Here, we reveal that *PRDM1*, which encodes Blimp1, is upregulated in both mTreg

and mTconv from patients with MS with a more significant increase in mTreg. This transcriptional

signature of Tregs is shared across different autoimmune diseases, suggesting the common

transcriptional signature driving dysfunctional Tregs in human autoimmunity. Specifically, an

alternative short isoform of *PRDM1 (PRDM1-S)*, which codes for protein in primates but not in

rodents, is primarily elevated in MS mTreg compared to the evolutionary conserved long *PRDM1*

isoform (*PRDM1-L*). Gene overexpression experiments in primary human Tregs, together with bulk

and single-cell RNA-seq transcriptional profiling, demonstrate a unique link between the *PRDM1-*

S and *SGK1* that accounts for Treg dysfunction observed in MS. Moreover, while genome-wide

chromatin accessibility in mTreg remains comparable between MS and healthy controls, the

transcription factor (TF) footprints and motifs within accessible chromatin regions are different with

significantly enriched binding of AP-1 and IRF family TFs observed in MS mTreg, suggesting

rewiring of regulatory circuits. Finally, CRISPR activation (CRISPRa)-based exploration of *cis*-

regulatory elements for the *PRDM1-S* identified a *cis*-regulatory element harboring AP-1 and IRF4

composite motif. These findings suggest a possible novel regulatory program by which Tregs

become dysfunctional in humans that is shared across multiple autoimmune diseases. Our

multimodal datasets of human CD4[+] T cells provide a rich resource for understanding the loss of

immune regulation in autoimmune diseases and suggest that the primate specific short *PRDM1*

isoform is a critical, targetable transcriptional regulator in human autoimmunity.

Results

Deep transcriptomic analysis of memory Treg and Tconv highlight PRDM1 upregulation in

MS (Figure 1)

We sought to identify CD4[+] T cell transcriptional differences between patients with MS and

healthy controls. As previous studies had not identified significant differences in bulk transcriptional

profiles of whole CD4[+] T cells between MS subjects and healthy controls [4,5], we divided CD4[+] T

cells into four major subpopulations based on two categories; Tconv vs. Treg, and naïve vs.

memory (Figure S1A), where each population is hypothesized to be involved in MS pathogenesis.

For example, mTconv contains pathogenic CD4[+] T cells in patients with MS, such as myelin-

reactive T cells, which display the signatures of Th1 and/or Th17 cells[20-23]. Moreover, mTregs with

reduced suppressive function are implicated in MS pathophysiology[24,25]. To provide a
comprehensive transcriptomic catalogue of memory CD4$^+$ T cell subpopulations in patients with
relapsing-remitting MS (RRMS), we performed RNA-seq on *ex vivo* mTreg and mTconv isolated
from the peripheral blood of untreated RRMS patients free of steroid treatment and healthy control
subjects as a discovery cohort (Figure 1A, S1A). The control subjects were matched by age, sex,
and ethnicity (clinical characteristics are described in Supplementary Table 1). Among a total of 90
RNA-seq samples (48 mTconv and 42 mTreg samples) in the discovery cohort (HC; n = 21, MS; n
= 30), 84 samples (HC; n = 20, MS; n = 26) passed quality control (Method Details). We identified
21 and 243 differentially expressed genes (DEGs, defined as |log2FC| >0.6, FDR<0.1) between
MS and healthy subjects for mTreg and mTconv, respectively (Figure 1B). Several DEGs were up-
or down-regulated in the same direction in both mTreg and mTconv: *PRDM1*, *BCL3*, and *PIM3*
were upregulated genes, and *ID3*, *TOB2*, and *LBH* were downregulated genes in MS (Figure 1C,
S1B). *PRDM1* was identified as one of the top genes significantly upregulated in MS in both mTreg
and mTconv (Figure S1C). Reduced expression of *ID3* in both cell types in MS reflects the negative
regulation of *ID3* by *PRDM1* consistent with the known function of *ID3* in maintaining Foxp3
expression in Treg[26] (Figure 1D). We confirmed that the frequency of Foxp3$^+$ Tregs within CD4$^+$ T
cells were not changed in patients with MS (Figure S1D), consistent with previous studies that
quantitative loss of Foxp3$^+$ Tregs is unlikely to be the primary feature of disturbed peripheral T cell
tolerance in patients with MS or other autoimmune diseases.

We validated several of the top DEGs (*PRDM1, BCL3, RHBDD2, TOB2, LBH*) in mTreg by
using qPCR with an independent cohort of patients with MS (n = 16) and healthy controls (n = 23)
(Figure 1A and E, Figure S1E, patient characteristics are described in Supplementary Table 1). We
further confirmed the up regulation of *PRDM1* in MS mTreg and mTconv at the protein level by
intracellular staining of Blimp1 using flow cytometry (Figure 1E). We then asked whether this Treg
transcriptional signature observed in MS is shared across different autoimmune diseases using
ImmuNexUT data[27] where transcriptomic profiles of multiple peripheral immune cells, including
mTreg and mTconv, were explored at population scale across multiple autoimmune diseases.
Notably, the transcriptional signature that we observed in MS Treg was also observed in Treg
among most of the autoimmune diseases we analyzed (total 12 diseases: 10 diseases from
ImmuNexUT, T1D from P. Gao *et al.*[28], and MS from this study) (Figure 1F, S1F). Inverse regulation
of *PRDM1* and *ID3* observed in MS was highlighted in Tregs from patients with systemic lupus
erythematosus (SLE), Idiopathic inflammatory myopathy (IIM), and ANCA-associated vasculitis

(AAV) (Figure S1G). Our bulk RNA-seq transcriptional profiling of memory Tconv and Treg

highlighted *PRDM1* as a key regulatory factor in dysfunctional Tregs in autoimmune diseases.

Figure 1

Figure 1 *Deep transcriptomic analysis of memory Treg and Tconv highlight PRDM1 upregulation in MS.* **(A)** Schematic of study design. Memory Tconv (mTconv) or memory Tregs (mTreg) were FACS sorted from peripheral blood CD4⁺ T cells from patients with MS and healthy control subjects (HC). Bulk RNA-seq was performed on the discovery cohort and DEGs were identified (HC: n=20, MS: n=26). The selected DEGs were validated by an independent validation cohort (HC: n=23, MS: n=16). **(B)** Volcano plots showing DEGs for mTreg and mTconv between MS and HC. **(C)** Overlapped DEGs between mTreg and mTconv. **(D)** qPCR validation for *PRDM1* expression in both discovery and validation cohorts. **(E)** Protein validation for Blimp1 expression using flow cytometry (HC; n=12, MS; n=9). **(F)** Heatmap depicting expression patterns of selected six genes in mTreg/Fr2 eTreg across 12 autoimmune diseases (data were extracted from M. Ota *et al.* and P. Gao *et al.*). P*<0.05, P**<0.01; Statistical significance computed by unpaired t-test **(D, E)**.

Single-cell dual omics analysis reveals elevated PRDM1 in Th17-like Treg in MS. (Figure 2)

To gain a deeper understanding of cellular heterogeneity and novel cell types underlying

disease mechanisms, we designed and performed single-cell RNA-sequencing (scRNA-seq) to

profile CD4⁺ T cells. To overcome the sparsity of scRNA-seq data[29,30], we used Cellular Indexing

of Transcriptomes and Epitopes by sequencing (CITE-seq)[31,32], and profiled 44 surface protein

markers (Supplementary Table 2) and mRNA expression simultaneously in total CD4⁺ T cells

enriched with Treg cells. Additionally, to avoid experimental batch effects between controls and

MS, we adopted hashing technology to pool cells into a single run of the 10x Genomics platform.

Across five experimental batches, we obtained a comparable number of Treg and Tconv cells

(Figure S2A). Demographic backgrounds (age, sex, and ethnicity) are controlled in each batch of

10x Genomics processing between controls and MS, and a total of five paired MS and control

samples were included (Supplementary Table 1).

We established cell identities based on surface marker proteins, eliciting prior knowledge

of cell-type-specific signatures. We first built a data matrix of 25,267 features (proteins and genes)

and 36,983 cells, combining five batches of CITE-seq profiles. We conducted a basic quality control

procedure to remove low-quality cells (Figure S2B) followed by batch normalization to control

batch-specific bias (Methods). We annotated cell type identities in semi-supervised training guided

by combinations of surface markers (Figure 2A), clearly demonstrating distinctive cell populations

in both the protein and transcriptomic space as can be seen in the gene-expression-based

clustering patterns in the UMAP visualization (Figure 2B). We confirmed that cell type annotation

results were not affected by batch labels or disease groups (Figure S2C). Our experimental design

also consistently enriched rare Treg populations and provided sufficient cells to study the variation

within Tregs (Figure S2D). The markers for Treg vs Tconv (*FOXP3, IKZF2*, *IL2RA/*CD25, and

*IL7R/*CD127), and memory vs naive (CD45RA, CD45RO) clearly distinguished four CD4$^+$ T cell

subpopulations (Figure 2C, Figure S2E).

For differentially expressed gene[33] analysis between MS and control subjects at the pseudo-

bulk level[34], we quantified subject-level gene expression profiles after adjusting contributions of

putative confounding factors, which stem from unmeasured technical covariates but may create

spurious associations with disease status and cell types (see Methods for details)[35]. We found that

single-cell-based analysis identified 90 up-regulated genes, and 16 down-regulated genes in MS

mTreg (Figure S2F). We compared the DEG effect size calculated in the scRNA-seq data (y-axis)

with the bulk DEG results (x-axis) (Figure 2D and E), which confirmed that a substantial number of

DEGs found in the bulk RNA-seq analysis are replicated in the matched subpopulations (mTreg

and mTconv) with a statistically significant correlation in both mTreg and mTconv (mTreg; r = 0.28,

p=2^{-178}, mTconv; r = 0.25, p=2^{-149}). For example, upregulation of *PRDM1* and downregulation of

LBH were demonstrated in MS mTreg (Figure 2D). *PRDM1* was also upregulated in mTconv, which

further validated the importance of *PRDM1* in the MS T cell signature (Figure 2D and E, Figure

S2G). *DDIT4*, which suppresses mTOR function, was upregulated in mTreg as well as mTconv

and downregulation of *TRAF3IP3* in MS mTreg, supporting the dysfunctional Treg property and

skewed Th17 signature in MS[36-39]. Of note, expression of CD45RO and *FOXP3* was not

significantly altered between MS and control subjects (Figure S2G), indicating our differential
analysis based on dual omics single-cell analysis was not biased by skewed memory differentiation
or significant loss of *FOXP3* gene expression in MS mTreg.

To elucidate the plasticity of the memory CD4⁺ T cell population, we further determined the T
helper cell subtypes (Th1, Th2, Th17, Th1/17) in mTreg and mTconv by using key marker
expression with CITE-seq (Figure 2F, G, Figure S3A-E, see also Methods). We then analyzed the
gene expression changes between MS and controls at each mTreg subtype and found that *PRDM1*
was significantly upregulated in Th17-like mTreg (mTreg17) (Figure 2H). Upregulation of CD226

Figure 2

Figure 2 *Single-cell dual omics analysis reveals elevated PRDM1 in Th17-like Treg in MS.* **(A)** Surface protein guided CD4⁺ T cells subtype annotation. Four CD4⁺ T cell subtypes were distinguished by CD25, CD127, CD45RO, and CD45RA protein expression. **(B)** UMAP based on gene expression for CD4⁺ T cells demonstrating decent overlap with protein-based subtype annotation. **(C)** *FOXP3*, *IKZF2*, *PRDM1*, and *BACH2* expressions on UMAP (all cells passed QC are plotted). **(D)** Combined differential analysis for bulk- and scRNA-seq in mTreg. Representative differentially expressed genes with pseudo-bulk analysis with scRNA-seq are shown. Gene expression changes in MS relative to control with indicated genes are computed at the single-cell level (see detailed in Methods). The size of dots is scaled proportionally to the average number of mRNA reads quantified within each batch and condition; the y-axis (disease effect) shows average gene expression after adjusted by confounding factors by matching; the error bars capture one standard deviation for the disease effects in Bayesian inference. Experimental batches for paired HC and MS samples are color coded (#1-5). **(E)** Combined differential analysis for bulk- and scRNA-seq in mTconv. **(F)** Sub-cell type analysis based on CITE-seq. Surface CXCR3 (CD183) and CCR6 (CD196) protein expressions of four subtypes are shown (log10 scale). **(G)** Heatmaps showing the marker genes and proteins to define subtypes for each mTconv and mTreg. **(H)** The changes of *PRDM1* expression at subtype-level analysis in mTreg between MS and control subjects.

and *CD6* in mTreg17 and CD278 (ICOS) and *JUNB* in Th17 indicates the skewed pathogenic Th17-

like signature in MS memory T cells (Figure S3F)[40-43]. Given the causative role of Th17 in MS

pathophysiology, our results at single-cell resolution further dissected the features of MS mTreg

and highlighted the potent role of *PRDM1* in skewing mTreg and mTconv toward the pathogenic

Th17-like signature in the context of MS.

Elevated alternative short PRDM1 isoform in MS mTreg. (Figure 3)

Our unbiased transcriptomic profiling of MS mTreg using bulk and single-cell RNA-seq

identified *PRDM1* as a top candidate transcription factor accounting for dysfunctional Treg

properties in MS. *PRDM1* encodes the Blimp1 protein which functions as a zinc finger

transcriptional repressor initially identified as a protein that binds to the promoter of *IFNB1* and

suppresses its expression[44]. The development and function of a variety of immune cells are also

under the control of Blimp1. CD4⁺ T cell-specific *PRDM1* deletion exaggerates the proinflammatory

reaction in multiple murine models of autoimmune diseases, including EAE[45,46]; in contrast,

previous studies demonstrated Blimp1 as an essential factor driving inflammatory programs in

Th17 differentiation[47]. This contradictory evidence suggests the context-dependent roles of *PRDM1*

in CD4⁺ T cells in mice. Recently, the role of *PRDM1* in Treg cells was studied by using Treg-

specific *PRDM1* knockout mice in the context of autoimmunity. Loss of their suppressive capacity

in *PRDM1*-deficient Treg indicates that *PRDM1* plays a critical role in maintaining Treg

homeostasis in tissue and positively regulates its suppressive function[48-50]. Thus, our result

showing upregulation of *PRDM1* in MS mTreg fundamentally contradicted observations in mice.

In humans, it has been appreciated that *PRDM1* has two major isoforms: the original full-
length isoform and another short isoform generated by alternative promoter usage[51]. The short
PRDM1 isoform (*PRDM1-S;* encoding a short form of Blimp1, Blimp1-S) arose during dry-nosed
primate evolution, and thus is not coded in the mouse genome. Blimp1-S lacks the N-terminal
region of Blimp1, which results in missing a part of the PR domain that is important in mediating
interaction with co-factors of Blimp1 (Figure 3A). Thus, Blimp1-S is implicated as a dominant
negative form against the full-length Blimp1[51,52]. Chromatin accessibility analysis at the *PRDM1*
locus showed that the *PRDM1-S* promoter region is significantly accessible in human Treg and our
previous HiDRA analysis[53] revealed stronger activity at the promoter of *PRDM1-S* than that of full-
length *PRDM1* isoform (*PRDM1-L*), which is consistent with DNase I hypersensitive site (DHS)
data of human primary total Treg[54] and T cells[55] (Figure 3B). In addition, bulk RNA-seq with nine
different peripheral blood human immune cells confirmed that both short and long *PRDM1* isoforms
were expressed in a cell-type-dependent manner. More specifically, monocytes and B cells mainly
express *PRDM1-L*. In contrast, NK cells dominantly express *PRDM1-S*, and CD4[+] T cells (including
Treg) and CD8[+] T cells express more *PRDM1-S* than *PRDM1-L*, especially in the memory
population, suggesting cell type-specific roles for each isoform (Figure 3C, S4A). This cell-type
specific expression patterns of *PRDM1-S* and *-L* were further validated at protein level by western
blot (Figure 3D, S4B). There are three bands detected between 80-110 kD consistent with previous
studies[56], though the band sizes are slightly higher in our data. To confirm the band size of Blimp1-
S and Blimp1 protein, we overexpressed the open reading frame of Blimp1-S or Blimp1 in 293T
cells by lentiviral transduction and found that lower two bands are derived from Blimp1-S while the
higher molecular weight band is from Blimp1 (Figure S4B). We then analyzed our mTreg bulk RNA-
seq at transcript level and found that *PRDM1-S* is significantly increased in MS mTregs compared
to healthy controls, though there is a moderate difference for *PRDM1-L* expression between MS
and control subjects (Figure 3E). This alteration was further validated by qPCR in both discovery
and validation cohorts (Figure S4C). Notably, this upregulation of *PRDM1-S* expression was also
observed in SLE Tregs from ImmuNexUT data[27] (Figure S4D). These data suggest that *PRDM1-S*
is a key regulator of Treg function in autoimmune diseases. These lines of data prompted us to
hypothesize that the aberrant induction of *PRDM1-S* may confer dysfunctional properties to MS
Tregs. Given that Blimp1-S can serve as a dominant negative isoform against conventional Blimp1,
we first examined whether the ratio of *PRDM1-S* and *PRDM1-L* is altered between MS vs HC.
However, there was no significant difference observed in this ratio with either bulk RNA-seq or
qPCR data, suggesting that the balance between *PRDM1-S* and *PRDM1-L* was not significantly
disrupted in MS Tregs (Figure S4E). Although *PRDM1-S* level was significantly upregulated in MS

Tregs, *PRDM1-L* levels were not changed or slightly increased in MS Treg (Figure 3E, S4C), thus

PRDM1-S mediated effects in MS Tregs could be independent from its dominant negative function

against *PRDM1-L*.

In order to identify genes that are differentially correlated between *PRDM1-S* and *PRDM1-L*,

we further analyzed the co-expression pattern of immune-related genes with *PRDM1-S* and

PRDM1-L in mTreg from HC and MS (Figure 3F); genes associated with effector and tissue

resident Treg signature (i.e. *BATF, CCR8, ICOS, CD69, IL1RL1, AREG, IRF4*) were co-expressed

with both *PRDM1-L* and *PRDM1-S,* particularly in MS Tregs, which reflects the skewing feature of

MS mTreg towards effector/tissue resident properties. Differentially expressed genets in MS

mTregs, such as *BCL3,* were positively correlated with both *PRDM1* isoforms in MS but negatively

correlated in healthy controls. Of note, these positive correlations in MS are stronger for *PRDM1-*

S compared to *PRDM1-L*. This type of correlation pattern (positive correlation for MS but negative

for control Tregs and higher correlation with *PRDM1-S* than *PRDM1-L*: Box 1 and 2 in Figure 3F)

was observed for the following genes; *IRF1*, which acts as a negative regulator for Foxp3

expression[57]; *BATF* and *FOSL2*, which belong to AP-1 family, with crucial functions for Treg

differentiation and maintenance[58,59]; *SGK1*, which is known to disrupt Treg homeostasis[14,15] and

Figure 3

Figure 3 *Elevated alternative short PRDM1 isoform in MS mTreg.* **(A)** Schematic of *PRDM1* short and long isoforms. PR; PR domain, Pro/Ser; Proline/serine rich region, ZnF; five C2H2 zink fingers. **(B)** ATAC-seq (mTreg and nTreg), DHS (ENCODE primary Treg, Roadmap primary T cells, ENCODE monocytes, Roadmap Primary B cells), and HiDRA peaks at *PRDM1* locus. **(C)** *PRDM1-S* and *PRDM1-L* isoform expression across 9 different immune cell types in peripheral blood assessed by bulk RNA-seq (n=6). **(D)** Western blot analysis of Blimp1 expression from 8 different immune cell types in peripheral blood. Conventional Blimp1 and alternative Blimp1-S are distinguished by different size. **(E)** *PRDM1-S* and *PRDM1-L* gene expression assessed by bulk RNA-seq in mTreg between MS and HC. P**<0.01; Statistical significance computed by one way ANOVA with Dunn's multiple comparisons tests. **(F)** Heatmaps depicting the Spearman's correlation for curated immune related genes with *PRDM1-S* or *PRDM1-L* in HC or MS.

260   play pathogenic roles in EAE[60]. We also noted that suppressive molecules enhancing Treg function

261   (i.e. *IKZF4, TIGIT, LAG3, ID3*) are negatively correlated with *PRDM1-S* in MS but not in healthy

262   controls (Box 3 in Figure 3F), supporting the association between *PRDM1-S* and dysfunctional

263   Treg properties in MS. Taken together, we identified *PRDM1-S* as a significantly upregulated

264   alternative isoform that partially accounts for the dysfunctional gene signature of MS mTreg.

Short PRDM1 induces SGK1 and Treg dysfunction. (Figure 4)

Next, we hypothesized that *PRDM1-S* has a unique feature independent of interfering from

PRDM1-L function. To test this hypothesis, we transduced *PRDM1-S* or *PRDM1-L* into primary

human Tregs by using a lentivirus-based overexpression (OE) system. Isolated human primary

Tregs were infected with lentivirus encoding *PRDM1-S* or *PRDM1-L* with GFP reporter and GFP

positive transduced cells were sorted by FACS after four days of culture. Overexpression for each

transcript was confirmed by qPCR (Figure S5A). Since only three amino acids are unique to

PRDM1-S (Blimp1-S) coding sequence compared to *PRDM1*-L (Blimp1), overexpression of

Blimp1-S open reading frame cannot be detected by qPCR, because qPCR primers are targeted

to unique sequence of 5' UTR region of *PRDM1-S*. Instead, we confirmed upregulation of total

PRDM1 but not *PRDM1-L* with Blimp1-S OE by qPCR (Figure S5A). Protein level induction of

Blimp1-S with this system was confirmed by western blot (Figure S4B). Bulk RNA-seq was

performed on sorted GFP positive cells and highlighted 100 genes exhibiting nominal evidence of

differential expression ($|log2FC| > 0.6$, P value < 0.05) between *PRDM1-S* overexpression and

GFP control (Figure 4A). We found that *SGK1* was one of the major upregulated genes induced by

PRDM1-S but not by *PRDM1-L* overexpression (Figure 4A, S5B). We validated this observation by

qPCR in human primary Treg cells and Jurkat T cells (Figure 4B). Moreover, *SGK1* was

upregulated in our scRNA-seq analyses, particularly in the Th17-like Treg subset together with

PRDM1, consistent with the role of SGK1 skewing pathogenic Th17 like signature in Tregs[60] (Figure

4C, S5C). This *PRDM1-SGK1* axis was a common feature among other autoimmune diseases in

a published dataset[27], where both *PRDM1* and *SGK1* were significantly increased in Tregs from

patients with SLE and ANCA-associated vasculitis (Figure S1E, S5D). Finally, we examined the
impact of *PRDM1-S* on Treg function by *in vitro* Treg suppression assay and found that Tregs with
PRDM1-S overexpression (OE) exhibited lower suppressive function than control GFP OE Treg,
strongly indicating that aberrant *PRDM1-S* expression causes Treg dysfunction (Figure 4D).
PRDM1-S OE decreased the level of both full-length Foxp3 and the exon 2-containing suppressive
isoform of Foxp3 protein, further confirming the impaired suppressive function on Tregs with
PRDM1-S OE (Figure 4E, S5E). These data support the unique role of *PRDM1-S* as a positive
regulator for *SGK1* in human Treg cells, especially in the subset of potentially pathogenic Th17-
like Treg, where *SGK1* plays a proinflammatory role under high sodium conditions and is
responsible for pathogenic features in a murine MS model via suppressing Foxp3 expression and
stablization[14,15]. Thus, these data suggest that the *PRDM1-S/SGK1* axis underlies Treg dysfunction
in MS.

Figure 4

Figure 4 *Short PRDM1 induces SGK1 and Treg dysfunction.* (A) Volcano plot showing statistical significance and fold change for genes differentially expressed by *PRDM1-S* overexpression in primary mTreg. **(B)** *SGK1* expression was assessed by qPCR for overexpression of *PRDM1-S* and *PRDM1-L* with primary human mTregs (left) and Jurkat T cells (right). P***<0.001, P****<0.0001; Statistical significance computed by one way ANOVA with Dunn's multiple comparisons tests. **(C)** Pseudo-bulk analysis of *SGK1, PRDM1, RORC, IRF4 and BATF* expression in scRNA-seq at each of four major mTreg subtypes. **(D)** *In vitro* Treg suppression assay with human primary Tregs. T effector cell proliferation was assessed after 5 days of co-culture with Tregs transduced with GFP control vector vs *PRDM1-S* OE vector. P**<0.01; Statistical significance computed by two-way repeated measures ANOVA. **(E)** Flow cytometry analysis for Foxp3 in primary Treg cells by overexpression of *PRDM1-S* compared to GFP control (n=10). P**<0.01; Statistical significance computed by paired t test.

Comprehensive analysis for chromatin accessibility reveals AP-1 and IRF enrichment in MS mTreg. (Figure 5)

To further elucidate the regulatory mechanisms underlying the dysfunctional properties of MS mTregs, we performed Assay for Transposase Accessible Chromatin sequencing (ATAC-seq) to identify chromatin-accessible signatures and epigenetic regulatory elements. We first determined the change of mTreg chromatin accessibility between MS and healthy controls. To our surprise, there was no significant difference in genome-wide chromatin accessibility between MS and healthy controls in mTreg (FDR<0.05), suggesting that global chromatin accessibility is not per se the major factor regulating gene expression in MS mTreg. We hypothesized that differential binding of regulatory TFs may account for the changes in mTreg gene expression between MS and control subjects. To identify TFs that potentially drive the observed gene expression signature in MS mTreg, we analyzed the enrichment of TF motifs and TF footprints within accessible regions between MS and healthy controls in mTreg (Figure 5A). We observed an enrichment of AP-1 and IRFs TF motifs that are important for CD4[+] T cell activation and differentiation in MS mTreg (Figure 5B).

To gain deeper insight into captured accessible regulatory elements, we performed a differential footprint analysis on ATAC-seq peaks by using HINT (Hmm-based IdeNtification of Transcription factor footprints)[61] and TOBIAS[62]. Consistent with our motif analysis, footprint analysis demonstrated the enrichment of AP-1 family TFs and IRFs in mTregs from MS compared to control subjects[63-66] (Figure 5C). AP-1 transcriptional activity is negatively regulated by direct interaction with Foxp3[67] and AP-1 is postulated to serve as a pioneer factor at Treg-specific regulatory elements where Foxp3 subsequently replaces it to establish Treg-specific enhancer architecture and DNA methylation [68,69]. Our observation of AP-1 enrichment in dysfunctional Tregs in MS could thus stem from the lower or impaired activity of Foxp3 in MS mTreg and reflect a more effector Tconv-like mTreg function in patients with MS[70,71].

IRF family TFs share a common DNA binding sequence (IRF binding element). Of interest, IRF-1 and IRF-2, but not IRF-4 nor IRF-8, are known to compete with evolutionally conserved *PRDM1-L*[72,73]. IRF-1 plays a critical role in Treg differentiation and maintenance[57,74] and it was of interest that *IRF1* was more co-expressed with *PRDM1-S* as compared to *PRDM1-L* in MS mTreg (Figure 3F). In addition, the enrichment of IRF-1 TF motif and footprint in MS mTreg (Figure 5B, C) could reflect the disrupted *PRDM1-L*-mediated gene regulation in MS mTreg. These results suggest that *PRDM1-L* mediated gene regulation could be disrupted in the context of MS (Figure S6A). To test this hypothesis, we first defined the set of genes that are specifically regulated by *PRDM1-L* by performing *PRDM1-L*-specific gene knockdown in human primary Tregs. A total of

1566 differentially expressed genes (defined as |log2FC| >1, FDR<0.05; 753 upregulated and 813

downregulated genes) were identified and defined as "*PRDM1-L* signature genes" (Figure S6A and

B). We performed single-cell level correlation analysis with our scRNA-seq data using a scalable

negative binomial mixed model, NEBULA[75]. We then investigated putative pathogenic downstream

effects of *PRDM1* expressions in single-cell analysis, examining whether the *PRDM1-L* signature

genes were perturbed by *PRDM1* genes between the mTreg cell groups derived from the healthy

and MS samples. We first ranked genes according to the correlation with single-cell *PRDM1*

expression levels in two independent cell groups derived from the healthy and MS samples,

respectively (thus, two lists of genes--one for the healthy and the MS). At a p-value threshold (the

x-axes of Figure S6C), we investigated whether the number of genes in the top lists were from the

PRDM1-L signature genes. So as not to be biased by a fixed p-value threshold, we conducted

bootstrapped enrichment tests for each p-value threshold. We found that the *PRDM1*-L signature

genes were significantly depleted in the top list of the genes derived from the MS mTreg group,

suggesting that the interactions between *PRDM1* and the target genes were more frequently

disrupted in the disease cells. We further explored the isoform level transcriptional co-regulation

by using our bulk RNA-seq data. Approximately half of positively correlated *PRDM1-L* signature

genes in Tregs from healthy controls lost their positive correlation in MS Tregs (Figure S6D,

Supplementary Table 3). Given that *PRDM1-L* plays a crucial role in maintaining Treg function [48-

50], we next assessed the correlation between *PRDM1-L* and Treg signature genes[76,77]. As observed

with *PRDM1-L* signature genes, approximately half of the positively correlated Treg signature

genes in healthy controls lost their positive correlation in MS Tregs (Figure S6E). Taken together,

our ATAC-seq results revealed that while genome-wide chromatin accessibility was not significantly

altered, differential binding of TFs (AP-1 and IRFs) to regulatory elements could serve as key

upstream regulatory factors and drive dysfunctional Treg gene programs by possibly disturbing

PRDM1-L mediated gene regulation in patients with MS.

Figure 5

Figure 5 *AP-1 and IRF TF bindings are enriched in MS mTreg.* **(A)** Schematic of ATAC-seq experiments for mTreg from MS (n=26) and healthy control (n=21). **(B)** TF motif enrichment analysis in mTreg between MS and HC by HINT. IRF and AP-1 motifs are significantly enriched in MS mTreg. **(C)** TF footprint enrichment analysis in mTreg between MS and HC by HINT and TOBIAS. IRF and AP-1 footprints are significantly enriched in MS mTreg.

Identification of active enhancer for Short PRDM1 in human T cells. (Figure 6)

We sought to determine the regulatory mechanisms that induce upregulation of *PRDM1-S* as

opposed to *PRDM1-L* in human mTreg. We reasoned that the enriched binding of AP-1 and IRFs

exert its regulatory function through binding to *cis*-regulatory elements for *PRDM1*, especially

PRDM1-S. To identify the *cis*-regulatory elements for *PRDM1-S*, we prioritized the accessible

chromatin regions surrounding the *PRDM1* locus (+/-0.5MB window around the transcriptional start

site of *PRDM1*) and examined the association between *PRDM1* expression and chromatin

accessibility in our human primary T cell ATAC-seq and RNA-seq data. Twenty significant

accessible regions (p<0.001) were identified as potential regulatory elements associated with

PRDM1 expression (Figure 6A). The majority of these peaks overlapped with H3K27ac ChIP-seq

signals for primary human Tregs[55], nominating them as potential enhancers. To functionally

validate these accessible chromatin regions, we adopted the CRISPR activation (CRISPRa)

system in Jurkat T cells that stably expresses catalytically inactive Cas9 fused to the transcriptional

activator VP64 (dCas9-VP64) [78] (Figure 6A).

First, we confirmed that *PRDM1-S* and *PRDM1-L* are independently regulated through

different promoter activity by our CRISPRa method with sgRNAs targeting each promoter region,

though there may be interactions between the transcription start site (TSS) of *PRDM1-L* and

PRDM1-S (Figure S7A). Next, we designed three sgRNAs for each accessible region and

generated sgRNA expressing lentiviral particles for twenty candidate regulatory elements (coded

as #1 to #20) (Supplementary Table 4, Figure 6A). dCas9-VP64 expressing Jurkat T cells were

infected by lentivirus encoding each sgRNA, then the double positive cells for GFP (dCas9-VP64)

and RFP (sgRNA) were sorted by FACS (Figure 6A). We observed that sgRNAs targeting the #2

peak region (-339,554 bp upstream of the *PRDM1-L* TSS) mediated a unique induction of *PRDM1-*

S but not *PRDM1-L* compared to control sgRNAs (Figure 6B). This #2 peak region is reported as

one of the "double elite" regulatory elements for *PRDM1* in the GeneHancer dataset[79], which

reflects a higher likelihood of prediction accuracy for both enhancer and target gene (Figure 6B;

top). The enhancer function of the #2 peak *cis*-regulatory element was further validated by

independent experiments with not only CRISPRa but also CRISPRi (Figure 6C, S7B).

To further clarify the function of this region as a *cis*-regulatory element at *ex vivo* human

primary Tregs, we sought to decode the histone modification. The main obstacle in the investigation

of chromatin state by using conventional ChIP-seq technique is the limited number of *ex vivo*

primary Tregs available for determining multiple histone marks from the same sample. Recent

development of multiplexed, indexed T7 ChIP-seq (Mint-ChIP)[80] allows us to identify both active

and repressive epigenetic marks by using histone modification-specific antibodies with limited cell

numbers. We performed Mint-ChIP on human primary Tregs in collaboration with the ENCODE

project. Histone modifications (H3K27ac, H3K4me1, and H3K4me3) distinguishing active

enhancers were assessed on *ex vivo* human primary Treg from MS and control subjects. The #2

peak region was marked by H3K4me1 and H3K27ac but without H3K4me3, suggesting the #2

peak region functions as an active enhancer (Figure 6D; top). Importantly, this #2 peak region

overlaps with AP-1 family and IRF4 ChIP-seq peaks and contains AP-1/IRF composite motif[81],

suggesting this enhancer element regulates *PRDM1-S* via AP-1 and IRF4 binding (Figure 6D;

bottom).

Given that BATF and IRF4 are known to bind in a cooperative fashion on the AP-1/IRF

composite motif facilitating Th17 differentiation [63,82], we examined the role of IRF4 and BATF on

PRDM1-S expression in Tregs by performing IRF4 and BATF knockdown experiments. Surprisingly,

PRDM1-S was upregulated by knocking down *IRF4* or *BATF* while in contrast, *PRDM1-L* was

downregulated by *IRF4* KD (Figure S7C). These data indicate that *IRF4* differentially regulates

PRDM1-S and *PRDM1-L* in human primary Tregs. Of note, *IRF4* or BATF KD did not affect *FOXP3*

expression, suggesting that the loss of IRF4 and BATF in human Tregs can induce *PRDM1-S* and

SGK1 expression without significant reduction of *FOXP3* expression, further indicating that the

dysfunctional *PRDM1-S/SGK1* axis observed in MS Tregs is counter regulated by the core effector

Treg regulator IRF4 and BATF. These data also suggest that the *cis*-regulatory element (#2 peak

region identified as upstream of *PRDM1-S*, Figure 6) serves as a negative regulatory element for

PRDM1-S expression via IRF4/BATF binding. Indeed, CRISPRi targeting on this #2 peak region

suppress *PRDM1-S* expression (Figure 6C). Thus, these results highlight IRF4/BATF as potential

upstream transcription factors negatively regulating *PRDM1-S and* indicating that a newly identified

cis-regulatory element that contains the AP-1/IRF composite motif may account for aberrant

PRDM1-S induction in MS mTreg.

Figure 6

Figure 6 *Active enhancer element for short PRDM1 with AP-1 and IRF bindings.* **(A)** Schematic experimental overview. 1. Identification of candidate *cis*-regulatory elements regulating *PRDM1* expression from ATAC-seq peaks. 2. CRISPRa based examination of *PRDM1-S* specific *cis*-regulatory elements. **(B)** CRISPRa validation for top 20 *PRDM1* associated regulatory elements. Top: Top 20 accessible chromatin elements that are associated with *PRDM1* expression are highlighted in red. Potential interactions of regulatory elements with *PRDM1* gene are analyzed by GeneHancer database and shown on the top. Middle and bottom; CRISPRa-induced expression of short *PRDM1* (middle) and Long *PRDM1* (bottom) were assessed by qPCR. Detailed information for all 20 regions is shown in Supplementary Table 4. **(C)** Validation CRISPRa and CRISPRi experiment for #2 peak sgRNA-1 (n=4). **(D)** Top: H3K27ac, H3K4me1, and H3K4me3 MINT-ChIP signal on the #2 peak region in mTreg from HC and MS. Four replicates of HC and two replicates of MS are merged into one representative track respectively. Middle: Footprint analysis on #2 peak region with TOBIAS footprint score. Bottom: AP-1 and IRF ChIP-seq signals identified on #2 peak region in ENCODE data are shown. AP-1/IRF composite motif identified in #2 peak region is highlighted. P*<0.05, P**<0.01, P****<0.0001; Statistical significance computed by one way ANOVA with Dunn's multiple comparisons tests.

Discussion

Disruption of peripheral CD4$^+$ T cell homeostasis is a central component driving pathogenesis

of autoimmune disease where autoreactive T cells lose tolerance to self-antigen by both intrinsic

and extrinsic mechanisms. Treg-mediated surveillance is central for controlling activation of

autoreactive CD4$^+$ T cells and dysfunctional Tregs are a hallmark of MS and other autoimmune

diseases [7,8]. In addition, recent analysis of genome-wide association studies emphasizes the

substantial contribution of CD4$^+$ T cells, including Tregs, as potentially causal mediators of

autoimmune disease [3]. Although several phenotypic changes have been identified in dysfunctional

Tregs, the underlying molecular mechanisms leading to breakdown of Treg suppressive function

in patients with autoimmune diseases are unknown. Here, by using MS as a model for studying the

molecular mechanisms of human Treg dysfunction, we examined transcriptional and epigenetic

alteration in human Tregs, identifying a previously unknown role of an alternative short *PRDM1*

isoform in dysfunctional Treg. *SGK1* was identified as a target of short *PRDM1*, which has been

reported to confer the pathogenic function of Treg in both human and mouse[14,15,60]. Moreover, both

PRDM1 and *SGK1* were upregulated in Tregs from different autoimmune diseases such as SLE

and ANCA-associated vasculitis[27], suggesting that the *PRDM1/SGK1* axis could serve as a

common feature of Treg dysfunction in the context of human autoimmunity. Finally, exploration of

epigenetic changes in MS Treg revealed an active enhancer element that induces short *PRDM1*

transcription, which was validated by CRISPRa/i experiments. AP-1 family, especially BATF, and

IRF4 directly bound to this regulatory element, implicating the role of these TFs in contributing to

PRDM1-S mediated Treg dysfunction in autoimmune disease. Thus, this study provides a novel

mechanistic insight of dysfunctional Tregs in the context of MS and potential therapeutic targets to

reverse Treg dysfunction in autoimmune diseases.

Previous studies exploring transcriptional alterations that occur in CD4$^+$ T cells in patients

with MS as compared to control subjects did not identify significant differences[4,5]. We hypothesized

that it was critical to examine CD4$^+$ T cell subpopulations. Thus, we further segregated CD4$^+$ T

cells into four subpopulations and performed transcriptional characterization using bulk RNA-seq

with deeper sequencing. This allowed us to detect significant differences in gene expression

between patients with MS and control subjects at both the gene and transcript level. Our bulk RNA-

seq based findings in the discovery cohort were further confirmed by two different means; (1)

utilization of a validation cohort with the same method as the discovery cohort, and (2) utilization

of CITE-seq to determine the subpopulations and assess the difference within the third cohort,

leading to highly reproducible findings. One caveat is that we were unable to quantitate *PRDM1*

isoform expression at single cell resolution in our 10x genomics dataset. Full length mRNA-
capturing scRNA-seq has the potential to differentiate between *PRDM1* isoforms; thus, this
technique could be applied to elucidate further insight at single-cell resolution[83]. This approach will
also enable characterization of further differences in T cell transcriptomics between MS and healthy
subjects.

Our transcriptomic analysis of *PRDM1* isoforms identified expression patterns between
PRDM1-S and *PRDM1-L* that are highly cell type specific. Memory T cells and NK cells expressed
higher levels of *PRDM1-S* as compared to *PRDM1-L,* while monocytes/DCs, and B cells
preferentially express *PRDM1-L.* Of note, *PRDM1* expression across all B cell linage (naïve,
unswitched memory, switched memory, double negative B cells, and plasmablasts) is strictly
limited to *PRDM1-L* over *PRDM1-S* (ImmuNexUT dataset). We also observed that the balance
between *PRDM1-S* and *-L* is dynamic and changes after T cell activation and cytokine milieu *in*
vitro. Thus, our findings demonstrate a tight linkage between *PRDM1* isoforms and immune
pathways related to AP-1/IRFs in autoimmune diseases and highlight a fundamental role in
primates for *PRDM1* isoform switching in regulating immune responses.

Epigenetic alterations in MS Tregs were assessed by using bulk ATAC-seq. Conventional
characterization with differential analysis of chromatin accessibility did not reveal significant
alterations of chromatin accessibility between patients with MS and healthy donors at a genome-
wide level. We then used our ATAC-seq data to elucidate TF footprint enrichment and found that
AP-1 and IRF family TFs are significantly enriched in MS mTreg compared to that of healthy
controls. It was of interest that neither AP-1 nor IRFs were detected as differentially expressed
genes by transcriptomic analysis, indicating that TF activity cannot be inferred by RNA expression.
Conventional TF ChIP-seq is thought to be the best method to provide direct evidence of TF activity;
however, it is technically challenging to achieve with primary human *ex vivo* mTregs due to their
low frequency in circulating blood. Thus, here we employed ATAC-seq footprint analysis to
overcome these issues for epigenetic profiling in human Tregs and provided novel evidence of
enriched AP-1 and IRFs bindings in MS mTregs that contributing to the Treg transcriptional
signature. These data agree with previous studies highlighting the important role of AP-1 in
establishing epigenetic state and linking genetic susceptibility to T cell activation[84]. Of note, recent
studies identified Treg specific eQTL effects with MS associated CD28 susceptible locus[85], which
was also confirmed with ImmuNexUT and DICE data (data not shown). Given that CD28 signaling
is central in the activation of AP-1 function in T cells, our findings of AP-1 enrichment in MS mTregs
potentially suggest that the genomic susceptibility of MS could be mediated by high activity of the

AP-1 family in mTregs, resulting in increased *PRDM1-S*. Further studies focused on the functional
properties of the MS-associated SNP at the CD28 locus are warranted.

A fundamental question relates to the elucidation of molecular interactions between
environment triggers and gene transcription driven by allelic variation associated with disease risk
that lead to autoimmune disease. We have previously shown that high Na^+ induces SGK-1 and
subsequent inactivation of Foxo1 leading to dysfunctional Tregs. This SGK1-Foxo pathway plays
a role in driving pathogenic Th17 cells especially under high salt environment[22]. Other recent
studies in humans have demonstrated higher Na^+ tissue levels in a subset of patients with MS[86].
Of note, gene set enrichment analysis (GSEA) demonstrated a dysfunctional Foxo1 and Foxo3 KO
Treg signature that was enriched in MS and SLE mTreg (Figure S8), consistent with our previous
studies showing impaired function of Foxo1 in dysfunctional Tregs in MS[15,87,88]. Thus, we
hypothesize that molecular interactions leading to dysfunctional Tregs in autoimmunity can be
driven in part through SGK-1 mediated by high sodium concentration as an environmental factor.
However, it should be pointed out that this could be just one of many environmental factors that
drive autoimmune disease.

In summary, our data uncover fundamental molecular mechanisms by which Treg dysfunction
is triggered in patients with MS and potentially other autoimmune diseases. Identification of the
primate-specific alternative short *PRDM1* isoform induction in MS Tregs highlights the importance
of studying human tissues in addition to mouse models in obtaining insight into disease
pathogenesis. Enhancement of the *PRDM1/SGK1* axis in mTreg was observed in the other
autoimmune diseases, suggesting shared mechanisms among dysfunctional Tregs in the context
of autoimmunity. Furthermore, our findings link epigenetic priming of AP-1 and IRF TF binding in
MS Tregs to short *PRDM1* induction. Finally, we believe our rich data of both transcriptome and
epigenome profiles on human memory CD4[+] T cells and Tregs will be a useful tool to explore further
insights into pathogenic mechanisms of dysfunctional T cells in autoimmune diseases.

Methods

Study subjects and ethics statement

Peripheral blood was drawn from people with MS and healthy controls who were recruited as part

of an Institutional Review Board (IRB)-approved study at Yale University, and written consent was

obtained. All experiments conformed to the principles set out in the WMA Declaration of Helsinki

and the Department of Health and Human Services Belmont Report.

Human T cell isolation and culture

Peripheral blood mononuclear cells (PBMCs) were prepared from whole blood by Ficoll gradient

centrifugation (Lymphoprep, STEMCELL Technologies) and used directly for total CD4+ T cell

enrichment by negative magnetic selection using Easysep magnetic separation kits (STEMCELL

Technologies). Cell suspension was stained with anti-CD4 (RPA-T4), anti-CD25 (clone 2A3), anti-

CD45RO (UCHL1), anti-CD45RA (HI100) and anti-CD127 (hIL-7R-M21, all from BD Biosciences)

for 30 minutes at 4°C. Naïve Tconv (CD4+/CD25neg/CD127+/CD45ROneg/CD45RA+), Naive

Treg (CD4+/CD25hi/CD127neg/CD45ROneg/CD45RA+), Memory Tconv

(CD4+/CD127+/CD45RO+/CD45RAneg), and Memory Treg

(CD4+/CD25hi/CD127neg/CD45RO+/CD45RAneg) were sorted on a FACSAria (BD Biosciences).

Sorted cells were plated in 96-well round-bottom plates (Corning) and cultured in RPMI 1640

medium supplemented with 5% Human serum, 2 nM L-glutamine, 5 mM HEPES, and 100 U/ml

penicillin, 100 μg/ml streptomycin, 0.5 mM sodium pyruvate, 0.05 mM nonessential amino acids,

and 5% human AB serum (Gemini Bio-Products). Cells were seeded (30,000-50,000/well) into

wells pre-coated with anti-human CD3 (2 μg/ml, clone UCHT1, BD Biosciences) along with soluble

anti-human CD28 (1 μg/ml, clone 28.2, BD Biosciences) in the presence or absence of human IL-

2 (50 U/ml).

Lentiviral transduction

<u>*Lentiviral production*</u>

Lentiviral plasmids encoding shRNA for gene knockdown for *PRDM1-L* or open reading frame of

overexpression for *PRDM1-S* and *PRDM1-L* were obtained from Sigma-Aldrich (MISSION shRNA)

and Horizon Discovery Biosciences (Precision LentiORF), respectively. dCas9-VP64-2A-GFP

(Addgene 61422) and pHR-SFFV-dCas9-BFP-KRAB (addgene 46911) were used for generating

Jurkat T cell lines for CRISPRa and CRISPRi, respectively. EF1a-RFP-H1-gRNA vector

(CASLV502PA-R from System bioscience) was modified to introduce BsaI cut site and single

sgRNAs were cloned into it by using Golden Gate Assembly kit (BsmBI-v2, New England Biolabs

#E1602). All single sgRNAs used in this study are listed in Supplementary Table 5. Each plasmid

was transformed into One Shot Stbl3 chemically competent cells (Invitrogen) and purified by

ZymoPURE plasmid Maxiprep kit (Zymo research). Lentiviral pseudoparticles were obtained after

plasmid transfection of 293T cells using TurboFectin 8.0 Transfection Reagent (Origene). The

medium was replaced after 6-12 h with fresh media with 1X Viral boost (Alstem). The lentivirus

containing media was harvested 72 h after transfection and concentrated 80 times using Lenti

Concentrator (Origene). LV particles were then resuspended in RPMI 1640 media without serum

and stored at -80°C before use. Virus titer was determined by using Jurkat T cells and Lenti-X

GoStix Plus (Takara Clontech).

Lentiviral transduction

FACS-sorted Tregs were plated at 50,000 cells/well in round bottom 96 well plates pre-coated with

anti-human CD3 (2 μg/ml, clone UCHT1, BD Biosciences) and soluble anti-human CD28 (1 μg/ml,

clone 28.2, BD Biosciences), in the presence of human IL-2 (50 U/ml). After 24 h, cells were

transferred into Retronectin coated 96 well plates and 25-50 μl of lenti particles were added to each

well, then spinfected with high-speed centrifugation (1000 g) for 1.5 hour at 32 °C. Immediately

after centrifugation, cells are placed back to the culture. On day 5, cells are harvested and GFP

positive cells are sorted by FACSAria or analyzed by Fortessa. Gene knockdown effect by *PRDM1-*

L shRNA is shown in Figure S6B.

Jurkat T cells were plated at 50,000 cells/well in round bottom 96 well plates and 25 μl of lenti

particles were added to each well, and spinfected as well as above. On day 3-5, cells were scaled

up to 12 well plates and followed by the second scale-up at day 7-9 into 6 well plates. Cells were

stimulated with PMA and Ionomycin (50 ng/ml and 250 ng/ml respectively) for four hours and

GFP$^+$/RFP$^+$ double positive cells were sorted directly into RNA lysis buffer by FACS Aria.

Suppression assay

CD4$^+$CD25$^+$CD127^{neg} Treg cells were sorted on a FACS Aria (BD Biosciences). Treg cells were

transduced with lentiviral particles containing *PRDM1-S* ORF or GFP control. GFP positive cells

were sorted by FACS at day 5, and freshly sorted T effector cells were labeled with cell trace

violet dye and then co-cultured with GFP$^+$ T$_{reg}$ cells (1 x 10^4) at different ratio with human Treg

inspector beads at 2:1 bead-to-cell ratio. The proliferation of T effector cells was determined at

day 4 on a BD Fortessa instrument (BD Bioscience).

Flow cytometry analysis

Cells were stained with LIVE/DEAD Fixable Near-IR Dead Cell Stain kit (Invitrogen) and surface

antibodies for 30 min at 4°C. For intracellular cytokine staining, cells were fixed with BD Cytofix™

Fixation Buffer (BD Biosciences) for 10 min at RT, then washed with PBS. Intracellular staining

was performed in Foxp3 permeabilization buffer (Thermo Fisher) for 30 min at 4°C. The following

antibodies were used: anti-Foxp3 (clone 150D, Biolegend, and clone PCH101, Thermo Fisher),

anti-Blimp1 (clone 3H2-EB, Thermo Fisher). All antibody information is listed in Supplementary

Table 6. Cells were acquired on a BD Fortessa flow cytometer and data was analyzed with FlowJo

software v10 (Threestar).

Immunoblotting

Cells were lysed with RIPA buffer containing EDTA-free cOmplete protease inhibitor cocktail

(Roche) and Halt phosphatase inhibitor cocktail (Thermo Fisher Scientific). Extracted protein was

quantified with a BCA kit (Thermo Scientific). 0.8-1 μg of protein extract was loaded in each lane,

followed by separation by 7.5% SDS-PAGE and transfer to a nitrocellulose membrane. After 1 hour

of blocking with 2.5 % bovine serum albumin (BSA) containing 1X Tris-Buffered Saline, 0.1%

Tween20 Detergent (TBST), the blotted membranes were then incubated overnight with primary

antibodies (anti-Blimp1 (C14A4, 1:1,000, Cell Signaling Technologies or C-7, 1:200, Santa Cruz

Biotechnology), anti-β-actin (A5441, 1:10,000, Sigma-Aldrich)) in TBST with 2.5% BSA. Primary

antibodies were detected by the secondary antibody horseradish peroxidase (HRP)–conjugated

anti-rabbit or mouse (Cell Signaling Technology), and SuperSignal West Femto Maximum

Sensitivity ECL Substrate (Pierce) was used as a chemiluminescent substrate for detecting HRP.

The images were obtained with a ChemiDoc Imaging system (Bio-Rad).

Bulk RNA-seq and ATAC-seq library preparation and sequencing

Bulk RNA-seq: FACS sorted cells (5,000 cells) were subjected to cDNA synthesis using SMART-

Seq v4 Ultra Low Input RNA Kit for sequencing (Takara/Clontech). Barcoded libraries were

generated by the Nextera XT DNA Library Preparation kit (Illumina) and sequenced with a 2x100

bp paired-end protocol on the HiSeq 4000 Sequencing System (Illumina).

Bulk ATAC-seq: We adopted FAST-ATAC for our FACS sorted CD4$^+$ T cells (5,000 cells) (Corces

et al., 2016). Cells were pelleted by centrifugation at 500 g for 7 min at 4°C, then resuspended with

50 μl of transposase mixture (25 μL of 2x TD buffer (Illumina), 2.5 μL of TDE1 (Illumina), 0.5 μL of

1% digitonin (Thermo Fisher), 22 μL of nuclease-free water). Transposition reactions were

incubated at 37°C for 30 minutes in a thermal shaker with agitation at 300 RPM. Transposed DNA

was purified using a MinElute Reaction Cleanup kit (QIAgen) and purified DNA was eluted in 20

µL elution buffer. Transposed fragments were amplified and purified as described previously [89] with

modified primers [90]. Libraries were quantified using qPCR (KAPA Library Quantification Kit) prior

to sequencing. All Fast-ATAC libraries were sequenced using a 2x100 bp paired-end protocol on

the HiSeq 4000 Sequencing System (Illumina).

Mint-ChIP library preparation and sequencing

We used 800,000-1,000,000 cryopreserved sorted human primary Tregs for Mint-ChIP profiling.

Cells were thawed and immediately pelleted by centrifugation at 400 g for 7 min at 4°C, then

resuspended with 100 µL of ice cold PBS. Approximately 100,000 cells per antibody are lysed in

detergent and chromatin is digested with micrococcal nuclease. Double stranded adapters (that

contain both a promoter for transcription by T7 RNA polymerase and a demultiplexing adapter) are

ligated to the chromatin. Chromatin is mixed overnight with antibodies recognizing histone

modifications, and immune complexes are captured using protein A / protein G magnetic bead

mixtures. In addition, some chromatin is set aside overnight to enable preparation of an antibody

free, input control library. Immobilized immune complexes are washed, and immunoprecipitated

DNA is eluted using proteinase K. Recovered DNA is purified with SPRI beads and subject to T7

RNA synthesis, thus creating an RNA copy of the immunoprecipitated DNA. RNA is copied back

to cDNA using a random primer containing a 5' extension, enabling subsequent PCR amplification

with Illumina indexed sequencing primers. PCR products are purified and mixed together to enable

multiplex Illumina sequencing. DNA is sequenced using a paired end protocol; in Mint-ChIP3, the

first 8 bases of Illumina Read2 serve as an inline barcode enabling demultiplexing of the chromatin

using the ligated barcoded adapter. The detailed protocol and primer/adaptor sequences are

described in dx.doi.org/10.17504/protocols.io.wbefaje.

Mint-ChIP data processing

We processed the Mint-ChIP FASTQ files of each sample using the ENCODE3 ChIP-seq pipeline

provided by Anshul Kundaje (https://github.com/ENCODE-DCC/chip-seq-pipeline2) with the

following parameters "chip.pipeline_type=histone, chip.aligner=bowtie2, chip.true_rep_only=true,

chip.paired_end=true, chip.ctl_paired_end=true, chip.always_use_pooled_ctl=false" specified in

its json file. We used the default value for all the other parameters. Briefly, the pipeline first mapped

the reads to the hg19 human reference genome using bowtie2. The aligned reads were filtered and

duplicated reads were removed. The peak calling was then performed using MACS2 with a control

sample for each individual. Sample quality was assessed with a cross-correlation plot.

Bulk RNA-seq analysis

After sequencing, adapter sequences and poor-quality bases (quality score < 3) were trimmed

with Trimmomatic. Remaining bases were trimmed if their average quality score in a 4 bp sliding

window fell below 5. FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) was

used to obtain quality control metrics before and after trimming. Remaining reads were aligned to

the GRCh38 human genome assembly with STAR 2.5.2 [91]. We used Picard

(https://github.com/broadinstitute/picard) to remove optical duplicates and to compile alignment

summary statistics and RNA-seq summary statistics. After alignment, reads were quantitated to

gene level with RSEM [92] using the GENCODE annotation [93].

We conducted our initial quality control assessment on the entire dataset, including memory and

naïve Tconv and Treg cells obtained from MS patients and healthy controls. A subset of these

were stimulated with IL-2 as described above, with the remainder collected in the *ex vivo* state.

We used principal component analysis to identify potential sample swaps. We considered genes

that were quantitated >1 count per million (cpm) in ≥ 15 samples, normalizing expression values

by the trimmed mean of M-values as implemented in edgeR [94]. We used limma [95] with the voom

transformation [96] to identify differentially expressed genes (DEGs) within mTconv and mTreg cell

populations separately. We used RUV-seq [97] to account for batch effect and other sources of

systematic variation; we included RUV parameters along with a sex covariate in our final model.

DEG list for *ex vivo* mTreg and mTconv between MS vs HC are shown in Supplementary Table

7.

Co-expression analysis in bulk RNA-seq and scRNA-seq

The co-expression between two genes was computed as the Spearman correlation coefficient of

normalized gene expression. The normalized gene expression was calculated by dividing the raw

count by the library size of the sample and its scaling factor obtained from the TMM normalization.

The co-expression analysis of *PRDM1* in the scRNA-seq data was conducted for genes that had

the average count per cell >0.005 in the mTreg and mTconv. The co-expression was measured

by the log(fold-change) between *PRDM1* and the gene of interest in a negative binomial mixed

model with the subjects as random effects implemented in NEBULA [75]. We used normalized

expression of *PRDM1* (the raw count divided by the library size of the cell) as the explanatory

variable and included in the model the proportion of reads from ribosomal protein genes and

mitochondrial genes as covariates. To assess the differential co-expression, the model was fitted

for 4896 and 4676 cells from the five MS patients and five healthy controls separately.

Bulk ATAC-seq analysis

The official pipeline of the Encyclopedia of DNA Elements [54] consortium
(https://github.com/kundajelab/atac_dnase_pipelines) {*kundajelab/atac_dnase_pipelines: 0.3.0*}
was adopted to preprocess the ATAC-seq raw data. The preprocessing started with the paired-end
ATAC-seq fastq files of each subject. More specifically, reads were trimmed for adapters using
cutadapt [98] and mapped to the human reference genome (hg19) using bowtie2 [99]. The output raw
bam files were filtered, deduped, and converted to single-ended Tn5-shifted tagalign files, which
were then used as the input for peak calling. The deduped bam files were used in the downstream
motif and footprint analyses described in a later section.

We first called sample-specific narrow peaks (FDR<0.01) using MACS2 [100] from the tagalign file of
each of the samples, separately, with the command "macs2 callpeak -f BED -g hs -q 0.01 --nomodel
--shift -75 --extsize 150 -B --SPMR --keep-dup all --call-summits". We calculated the fraction of
reads in called peak regions (FRiP score) for each sample. We then called group-specific narrow
peaks (FDR<0.01) by providing MACS2 with the tagalign files of all high-quality samples
(FRiP>0.1) within each of the eight groups (healthy/MS mTconv, mTreg, naive Treg (nTreg), and
naive Tconv (nTconv)) using the command "macs2 callpeak -t <tagalign files of the high-quality
samples within the group> -f BED -g hs -q 0.01 --nomodel --shift -75 --extsize 150 -B --SPMR --
keep-dup all --call-summits". To obtain a unified set of peak regions across all groups, we merged
the group-specific peak regions that were overlapping or had maximum distance <100bp using
BEDTools [101] with the command "bedtools merge -i -d 100".

Given the unified set of peak regions, the number of reads overlapping a peak was called for each
subject using BEDTools. This raw count matrix was used for the downstream analyses. In the
analysis of differential accessibility, we first filtered out peaks with low counts and normalized the
counts of each subject using TMM [94]. Then the *voom* function in the limma package [95] was used to
perform the differential analysis between the cases and controls within each cell type with FRiP,
sex and ethnicity as covariates.

Correlation analysis between RNA-seq and ATAC-seq data
The analysis of the correlation between the accessibility of the adjacent open chromatin regions
and the *PRDM1* expression was performed using a linear regression model in which the normalized
PRDM1 expression (raw count divided by the total library size and the scaling factor) was the
dependent variable and the adjusted ATAC-seq peak height was the explanatory variable. The
adjusted ATAC-seq peak height was the residual obtained by fitting a weighted linear model using
limma-voom for the peak with FRiP as the covariate. We included 106 samples from all the cell

populations and both groups and these samples had both RNA-seq and ATAC-seq measurements.
We interrogated 72 ATAC-seq peaks for which *PRDM1* was annotated as the nearest gene by
Homer. These peaks span from ~350k bp upstream to ~100k bp downstream of the transcription
start site of the long isoform of *PRDM1*.

ATAC-seq footprint analysis
To conduct differential footprint analysis between the disease groups and cell types, we generated
group-level bam files by merging the deduped bam files within each of the eight groups. A total of
47 high-quality ATAC-seq samples with FRiP>0.1 were included (MS group: 26 mTregs, control
group: 21 mTregs). The differential footprint analysis was performed using HINT v0.12.3 [102] and
TOBIAS v0.10.1 [62]. In the analysis using HINT, we first called footprints using the command "rgt-
hint footprinting --atac-seq --paired-end" with the group-specific bam files and peaks as the input
files. We then identified predicted binding sites using the command "rgt-motifanalysis
matching" for the 579 JASPAR (2018) core motifs for vertebrates. Finally, we identified
footprints showing differential binding activity between cell types or disease groups using
the command "rgt-hint differential" based on the results from the previous two steps. The
same bam files and input files were used in the analysis adopting TOBIAS, in which we first ran
"TOBIAS ATACorrect" to obtain bias-corrected signals and then ran "TOBIAS FootprintScores" to
obtain footprint scores. Finally, we performed the differential footprint analysis with the command
"TOBIAS BINDetect".

Single-cell RNA-seq using 10x Genomics platform
CD4$^+$ T cells and CD25^{hi} CD4$^+$ T cells were negatively isolated from PBMCs separately by using
Easysep human CD4$^+$ T cell isolation kits and EasySep Human CD4$^+$CD127^{low}CD25$^+$ Regulatory
T Cell Isolation Kit (STEMCELL Technologies), respectively. To avoid batch effects between
healthy and MS samples, and to increase the numbers of Tregs to analyze, we used hashing
technology (Biolegend) to pool samples in a single run of the 10x Genomics platform. Each MS
sample was processed with a paired healthy control subject matched for age, sex, and ethnicity. A
total of five healthy and MS sample pairs were analyzed. 100,000 cells for each cell type were
subjected to Total-seq C and Hashtag antibody staining. 2 ug per 1 million cells for Total-seq C
and 1 ug per 1 million cells for Hashing antibodies were used for the staining. Cells were washed
three times with PBS containing 2% FBS and four hashed samples (total CD4$^+$ T cells and
CD25^{hi}CD4$^+$ T cells from each of HC and MS) were pooled into one sample. The cellular
concentration was adjusted to 1,000/μL and loaded into the 10x Genomics instrument aiming to

recover 10,000 cells for library preparation and sequencing. Generated libraries were then

sequenced on the NovaSeq (Illumina) with a target of 50,000 reads/cell (2 × 150 paired-end reads).

Dual omics single-cell analysis

Cell type assignment by protein surface markers

We define a latent indicator variable z_{jk} to mark the assignment of a cell j to a cell type k and

estimate the posterior probability of $z_{jk}=1$ by the stochastic expectation maximization (EM)

algorithm. We assume that a normalized vector for each cell x_j follows von Mises-Fisher (vMF)

distribution [103] with cell type k-specific mean vector μ_k and shared concentration parameter κ:

$P(x_j|z_{jk}=1) \sim \exp(\kappa\ \theta_k^\top x_j)$. We modified the existing EM algorithm [104] and enforced the sparsity of

the mean vector μ based on the prior knowledge of the cell-type-specific activity of marker

proteins/genes/features. Simply, we allow μ_{gk} to take non-zero values if and only if a feature g is a

known marker for the cell type k.

To improve the quality of inference, we also take advantage of "negative" marker protein labels and
build an "adversarial" model for each cell type and contrast with the likelihood of the corresponding
"positive" model.
● nTconv: CD3+, CD4+, CD8-, CD25-/CD127+, CD45RA+/CD45RO-
● mTconv: CD3+, CD4+, CD8-, CD25-/CD127+, CD45RA-/CD45RO+
● nTreg: CD3+, CD4+, CD8-, CD25+/CD127-, CD45RA+/CD45RO-
● mTreg: CD3+, CD4+, CD8-, CD25+/CD127-, CD45RA-/CD45RO+

To further dissect the cell types within mTreg cells, we sorted single-cell CITE-seq vectors based
on the following definitions:
● Treg1: CD183+ / CD194- / CD196-
● Treg1/17: CD183+ / CD194- / CD196+
● Treg17: CD183- / CD194+ / CD196+
● Treg2: CD183- / CD194+ / CD196-
759
Batch correction and visualization of scRNA-seq data

We used the top 100 principal components of the log-transformed scRNA-seq data matrix to

characterize intercellular similarity and clustering patterns across ~45k cells and ~15k genes. We

managed to adjust discrepancies across five different batches using the batch-balancing k-nearest

neighborhood method [105] followed by adjustment of principal component, subtracting out the mean

difference between batches [106].

Single-cell differential expression analysis

We compared cell-type-specific gene expression profiles between the MS and HC subjects by

estimating unbiased subject-level pseudo-bulk profiles for each gene using CoCoA-diff [35]. CoCoA-

diff can improve the statistical power in case-control scRNA-seq study while adjusting for unwanted

confounding effects existing across individuals. We first estimated latent factors, which may

confound gene and cell-type-specific expressions with the disease labels. We established a

controlled baseline for each cell derived from the MS subjects by imputing counterfactual gene

expression values based on the 100 cells found in the HC cells in the top 10 PC space (BBKNN-

based weighted average). Likewise, we imputed counterfactual values for the HC cells using the

MS cells. More precisely, for each gene g and cell j, we have observed Y_{gj}[16] and counterfactual

(imputed) $Y_{gj}^{(HC)}$ if a cell j were derived from the MS; we have observed $Y_{gj}^{(HC)}$ and counterfactual

$Y_{gj}^{(MS)}$ if a cell j were from the HC. We then aggregate both factual (observed) and counterfactual

cell profiles within each subject and cell type to estimate causal effects by comparing the average

disease effect (ADE), average disease effect in the disease subject [107], and average disease effect

in the control subject (ADC). Denoting the subject-level aggregate profiles λ_{gi}[16] and $\lambda_{gi}^{(HC)}$ (for gene

g and subject i), we define ADE of a gene g as: $ADE(g) = \Sigma_{i=1..10} \log[\lambda_{gi}^{16} / \lambda_{gi}^{(HC)}] / 10$, $ADC(g) = \Sigma_i$

$_{in\ 5\ HC\ subjects} \log[\lambda_{gi}^{16} / \lambda_{gi}^{(HC)}] / 5$, $ADD(g) = \Sigma_{i\ in\ 5\ MS\ subjects} \log[\lambda_{gi}^{16} / \lambda_{gi}^{(HC)}] / 5$. We implemented

Bayesian inference methods in a C++ program that calculates gene-level statistics, including

posterior mean and standard error, efficiently handling ten thousand genes and hundred thousand

cells [35].

Data and code availability

The accession number for the whole transcriptome sequencing data, dual omics single-cell

sequencing data, and processed data reported in this paper will be provided in GEO/SRA.

Acknowledgements

We thank patients and families, members of the Yale MS clinic; Lesley Devine and Chao Wang for

technical assistance with flow cytometry; Mei Zhong and the staff of the Yale Stem Cell Center for

technical assistance with bulk RNA-seq and ATAC-seq sample preparation and sequencing.

Funding

This work was supported by Race to Erase MS Young investigator award to T.S.S.; a Career

Transition Fellowship from the Consortium of MS Centers and the National MS Society to M.R.L.;

NIH grants (P01 AI073748, U24 AI11867, R01 AI22220, UM 1HG009390, P01 AI039671, P50

CA121974, R01 CA227473) to D.A.H. NIA grants (R01 AG047310, R01 AG061853, R01

AG065477, and R01 AG070488) to A.M.K.

Author contributions

Conceptualization: T.S.S., M.R.L., L.H., Y.P., M.K., and D.A.H. Recruitment of study participants:
T.S.S., M.R.L., and D.A.H. Specimen collection: T.S.S., M.R.L., and H.A.S. Human primary cell
experiments: T.S.S., H.A.S., and G.A.L. Mint-ChIP experiment and analysis: T.S.S., M.R.L., L.H.,
C.B.E. and B.E.B. Transcriptomic and eQTL analysis of ImmuNexUT data: M.O., K.F. Data
analysis: T.S.S., M.R.L., L.H., and Y.P. Visualization: T.S.S., M.R.L., L.H., and Y.P. Writing review
& editing: T.S.S., M.R.L., L.H., Y.P., A.M.K., M.K., and D.A.H. wrote the manuscript, with input from
all co-authors. Supervision: M.K., and D.A.H.

Declaration of interests

D.A.H. has received research funding from Bristol-Myers Squibb, Novartis, Sanofi, and Genentech.
He has been a consultant for Bayer Pharmaceuticals, Bristol Myers Squibb, Compass Therapeutics,
EMD Serono, Genentech, Juno therapeutics, Novartis Pharmaceuticals, Proclara Biosciences,
Sage Therapeutics, and Sanofi Genzyme. B.E.B. declares outside interests in Fulcrum
Therapeutics, Arsenal Biosciences, HiFiBio, Cell Signaling Technologies and Chroma Medicine.
815

Supplementary Figures

Figure S1

Figure S1 Transcriptomic analysis in mTreg and mTconv with MS patients, related to Figure 1. (A) Experimental workflow for isolating four major CD4+ T cell subpopulations by FACS. **(B)** Top individual DEGs expression at subject level. **(C)** Correlation of *PRDM1* expression in mTreg and mTconv. **(D)** Flow cytometry-based assessment of the frequency of FOXP3+ T cells in HC vs MS. (n=34 (HC), 35 (MS)) **(E)** qPCR validation for top four DEGs in mTreg. Data for the discovery cohort (left side) and validation cohort (right side) are shown for each HC (yellow) and MS (purple). **(F)** *PRDM1* expression in Fr2 eTreg and Fr1 nTreg across 10 and 6 autoimmune diseases are shown respectively (data were extracted from M. Ota *et al.*). **(G)** *PRDM1 and ID3* expression in Fr2 eTreg from HC, SLE, IIM, and AAV are shown respectively (data were extracted from M. Ota *et al.*) P*<0.05, P**<0.01, P***<0.001, P****<0.0001; Statistical significance computed by one way ANOVA with Dunn's multiple comparisons tests.

Figure S2

Figure S2 Single-cell dual omics analysis with CD4⁺ T cells in MS, related to Figure 2. (A) Experimental workflow for dual omics single-cell analysis of HC and MS CD4⁺ T cells. Age, sex, and ethnicity matched HC and MS subject are processed at the same time as one experimental batch. Total five experimental batches were included in this study. **(B)** Histograms showing numbers of genes detected per cell (left) and frequency of mitochondrial genes per cell (right). Cells before and after quality control selection are highlighted in purple and light gray respectively. **(C)** Gene expression UMAP of all cells color coded for experimental five batches (#1-5) (top) and disease condition (bottom). **(D)** Annotated cell numbers of CD4⁺ T cell subpopulation within total CD4⁺ T cells (left), CD4⁺CD25⁺⁺ T cells (middle), and combined total cells (right). Cell numbers for each batch and summary of five batches from HC and MS are shown. **(E)** Representative gene and protein expressions UMAP. **(F)** Numbers of upregulated and downregulated DEGs in each CD4⁺ T cell subpopulation are shown. **(F)** Representative differential gene analysis in each CD4⁺ T cell subpopulation is depicted (See also Methods).

Figure S3 Single cell dual omics analysis with CD4⁺ T cells sub cell types in MS, related to Figure 2. (A) Surface protein guided mTreg and mTconv subtype annotation. CD196, CD183, and CD194 expressions were shown in four subtypes for each mTreg and mTconv. **(B)** Heatmaps showing the marker genes and proteins to define subtypes for each mTconv and mTreg at individual subject level. **(C)** Key marker expressions (CD196, CD183, CD194, *CXCR3* and *GATA3*) for each subtype in mTreg and mTconv. **(D)** UMAP based on CITE-seq based protein expressions for the subtypes in mTreg and mTconv. **(E)** Numbers of upregulated and downregulated DEGs in each subtype for mTreg (top) and mTconv (bottom). **(F)** Representative differential gene and surface protein analysis in each subtype in mTreg and mTconv are depicted (See also Methods).

Figure S4

Figure S4 Elevated alternative short *PRDM1* isoform in MS mTreg, related to Figure 3. (A) Representative bulk RNA-seq coverage tracks at *PRDM1* locus from nine different immune cell types in peripheral blood. Unique exonic regions for *PRDM1-L* and *PRDM1-S* are highlighted in orange and blue respectively. **(B)** Western blot analysis of Blimp1 expression from 8 different immune cell types in peripheral blood and Blimp1-S or Blimp-1 overexpressed (OE) 293T cells by anti-Blimp1 Ab (C-7) (top) and anti-Blimp1 Ab (C14A4) (bottom). **(C)** qPCR validation of short and long *PRDM1* isoform expression between HC and MS from discovery cohort (left box plot) and validation cohort (right box plot). P*<0.05, P**<0.01; Statistical significance computed by unpaired t test. **(D)** Short and long *PRDM1* isoform expression between HC and SLE from ImmuNexUT data. **(E)** Ratio of *PRDM1-S* vs *PRDM1-L* in bulk RNA-seq and qPCR data. P*<0.05, P**<0.01; Statistical significance computed by Mann-Whitney test.

Figure S5

Figure S5 Short *PRDM1* induces *SGK1* and Treg instability, related to Figure 4. (A) qPCR validation of total *PRDM1* and long *PRDM1* expression by short and long *PRDM1* isoform overexpression in mTreg. P*<0.05, P**<0.01; Statistical significance computed by paired t test. **(B)** Volcano plot showing statistical significance and fold change for genes differentially expressed by long *PRDM1* overexpression in primary mTregs. **(C)** *SGK1* expression assessed by scRNA-seq in four main CD4+ T cell subpopulations (top) and mTreg sub cell-types (bottom). ADE: average disease effect between disease cells and matched healthy cells across all the individuals, both MS and HC. ADC: average disease effect only measured within the healthy control group. ADD: average disease effect only measured within the disease group. **(D)** *SGK1* expression in Fr2 eTreg and Fr1 nTreg across 10 and 6 autoimmune diseases respectively (data were extracted from *M. Ota et al.*). P*<0.05, P**<0.01, P***<0.001, P****<0.0001; Statistical significance computed by one way ANOVA with Dunn's multiple comparisons tests. **(E)** Flow cytometry analysis for the specific splicing isoforms of FOXP3 containing exon 2 (FOXP3-Ex2) in primary Treg cells by overexpression of *PRDM1-S* compared to GFP control (n=5). P**<0.01; Statistical significance computed by paired t test.

Figure S6

Figure S6 Disrupted *PRDM1-L mediated gene regulation in MS mTregs*, related to Figure 5. (A) Schematic of how *PRDM1-L* mediated gene regulation is disrupted by enriched TF binding of IRF1/2. **(B)** Normalized gene expression for *PRDM1-S* and *PRDM1-L* on human primary Tregs with and without *PRDM1-L* specific gene knockdown by lentiviral shRNA transduction. Lentiviral transduced GFP⁺ cells were sorted by FACS at day 5. mRNAs were isolated and bulk RNA-seq was performed. **(C)** Quantile-quantile plot showing the co-expression correlation between *PRDM1-L* signature genes with *PRDM1* at a single-cell level in mTreg for MS and HC. Correlation between *PRDM1-L* signature genes and *PRDM1* expression is stronger in HC compared to MS. Wilcoxon's test p-value as summary values between MS vs HC: p-value = 0.005763. **(D, E)** Heatmaps depicting the Spearman's correlation between *PRDM1-L* expression and *PRDM1-L* signature genes **(D)** and Treg signature genes **(E)**.

Figure S7

Figure S7 Identification of active enhancer for short *PRDM1* in human T cells, related to Figure 6. (A) Schematic of CRISPRa experiment for short and long *PRDM1* induction with targeting each promoter element (left) and qPCR quantification of short and long *PRDM1* expression (right). **(B)** Validation CRISPRa and CRISPRi experiment for #2 peak independent from Figure 6B (n=4). Peak #2 sgRNA-1 and -3 were validated as functional *cis*-regulatory elements for *PRDM1-S*. **(C)** Lentiviral shRNA-based gene knockdown for IRF4 and BATF in human primary Tregs. Human primary Tregs are isolated by FACS and stimulated with anti-CD3/CD28 antibodies. Lenti particles were transduced at day 1 and GFP+ cells were sorted by FACS at day 4-5. Expressions of each gene assessed by qPCR are shown. P*<0.05, P**<0.01, P***<0.001, P****<0.0001; Statistical significance computed by one way ANOVA with Dunn's multiple comparisons tests.

Figure S8

Figure S8 Dysfunctional Foxo1 and Foxo3 KO Treg signatures in MS and SLE mTregs. Dot plots showing gene set enrichment analysis (GSEA) for Foxo signaling on Tregs in MS (left) and SLE (right) Tregs. Gene sets generated by Foxo1/3 KO Tregs and Foxo1 constitutive active (CA) Tregs are used. NES; Normalized enrichment score.

References

1   Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in
regulatory DNA. *Science* **337**, 1190-1195 (2012).
https://doi.org:10.1126/science.1222794

2   Yoshida, H. *et al.* The cis-Regulatory Atlas of the Mouse Immune System. *Cell* **176**, 897-
912 e820 (2019). https://doi.org:10.1016/j.cell.2018.12.036

3   Farh, K. K. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease
variants. *Nature* **518**, 337-343 (2015). https://doi.org:10.1038/nature13835

4   Hellberg, S. *et al.* Dynamic Response Genes in CD4+ T Cells Reveal a Network of
Interactive Proteins that Classifies Disease Activity in Multiple Sclerosis. *Cell Rep* **16**,
2928-2939 (2016). https://doi.org:10.1016/j.celrep.2016.08.036

5   Brorson, I. S. *et al.* No differential gene expression for CD4(+) T cells of MS patients and
healthy controls. *Mult Scler J Exp Transl Clin* **5**, 2055217319856903 (2019).
https://doi.org:10.1177/2055217319856903

6   Cvetanovich, G. L. & Hafler, D. A. Human regulatory T cells in autoimmune diseases.
Curr Opin Immunol **22**, 753-760 (2010). https://doi.org:10.1016/j.coi.2010.08.012

7   Miyara, M. *et al.* Human FoxP3+ regulatory T cells in systemic autoimmune diseases.
Autoimmun Rev **10**, 744-755 (2011). https://doi.org:10.1016/j.autrev.2011.05.004

8   Dominguez-Villar, M. & Hafler, D. A. Regulatory T cells in autoimmune disease. *Nat
Immunol* **19**, 665-673 (2018). https://doi.org:10.1038/s41590-018-0120-4

9   Viglietta, V., Baecher-Allan, C., Weiner, H. L. & Hafler, D. A. Loss of functional
suppression by CD4+CD25+ regulatory T cells in patients with multiple sclerosis. *J Exp
Med* **199**, 971-979 (2004). https://doi.org:10.1084/jem.20031579

10   Dominguez-Villar, M., Baecher-Allan, C. M. & Hafler, D. A. Identification of T helper type
1-like, Foxp3+ regulatory T cells in human autoimmune disease. *Nat Med* **17**, 673-675
(2011). https://doi.org:10.1038/nm.2389

11   McClymont, S. A. *et al.* Plasticity of human regulatory T cells in healthy subjects and
patients with type 1 diabetes. *J Immunol* **186**, 3918-3926 (2011).
https://doi.org:10.4049/jimmunol.1003099

12   Farez, M. F., Fiol, M. P., Gaitan, M. I., Quintana, F. J. & Correale, J. Sodium intake is
associated with increased disease activity in multiple sclerosis. *J Neurol Neurosurg
Psychiatry* **86**, 26-31 (2015). https://doi.org:10.1136/jnnp-2014-307928

13   Kleinewietfeld, M. *et al.* Sodium chloride drives autoimmune disease by the induction of
pathogenic TH17 cells. *Nature* **496**, 518-522 (2013). https://doi.org:10.1038/nature11868

14   Hernandez, A. L. *et al.* Sodium chloride inhibits the suppressive function of FOXP3+
regulatory T cells. *J Clin Invest* **125**, 4212-4222 (2015). https://doi.org:10.1172/JCI81151

15   Sumida, T. *et al.* Activated beta-catenin in Foxp3(+) regulatory T cells links inflammatory
environments to autoimmunity. *Nat Immunol* **19**, 1391-1402 (2018).
https://doi.org:10.1038/s41590-018-0236-6

16   Lappalainen, T. *et al.* Transcriptome and genome sequencing uncovers functional
variation in humans. *Nature* **501**, 506-511 (2013). https://doi.org:10.1038/nature12531

17   Wu, H. *et al.* High salt promotes autoimmunity by TET2-induced DNA demethylation and
driving the differentiation of Tfh cells. *Sci Rep* **6**, 28065 (2016).
https://doi.org:10.1038/srep28065

18   Cotsapas, C. *et al.* Pervasive sharing of genetic effects in autoimmune disease. *PLoS
Genet* **7**, e1002254 (2011). https://doi.org:10.1371/journal.pgen.1002254

19   Lincoln, M. R. *et al.* Joint analysis reveals shared autoimmune disease associations and
identifies common mechanisms. *medRxiv*, 2021.2005.2013.21257044 (2021).
https://doi.org:10.1101/2021.05.13.21257044

20 Panitch, H. S., Hirsch, R. L., Haley, A. S. & Johnson, K. P. Exacerbations of multiple
sclerosis in patients treated with gamma interferon. *Lancet* **1**, 893-895 (1987).
https://doi.org:10.1016/s0140-6736(87)92863-7
21 Lee, Y. *et al.* Induction and molecular signature of pathogenic TH17 cells. *Nat Immunol*
13, 991-999 (2012). https://doi.org:10.1038/ni.2416
22 Wu, C. *et al.* Induction of pathogenic TH17 cells by inducible salt-sensing kinase SGK1.
Nature **496**, 513-517 (2013). https://doi.org:10.1038/nature11984
23 Cao, Y. *et al.* Functional inflammatory profiles distinguish myelin-reactive T cells from
patients with multiple sclerosis. *Sci Transl Med* **7**, 287ra274 (2015).
https://doi.org:10.1126/scitranslmed.aaa8038
24 Schwarz, A. *et al.* Fine-tuning of regulatory T cell function: the role of calcium signals
and naive regulatory T cells for regulatory T cell deficiency in multiple sclerosis. *J*
Immunol **190**, 4965-4970 (2013). https://doi.org:10.4049/jimmunol.1203224
25 Ciccocioppo, F. *et al.* The Characterization of Regulatory T-Cell Profiles in Alzheimer's
Disease and Multiple Sclerosis. *Sci Rep* **9**, 8788 (2019). https://doi.org:10.1038/s41598-
019-45433-3
26 Rauch, K. S. *et al.* Id3 Maintains Foxp3 Expression in Regulatory T Cells by Controlling
a Transcriptional Network of E47, Spi-B, and SOCS3. *Cell Rep* **17**, 2827-2836 (2016).
https://doi.org:10.1016/j.celrep.2016.11.045
27 Ota, M. *et al.* Dynamic landscape of immune cell-specific gene regulation in immune-
mediated diseases. *Cell* **184**, 3006-3021 e3017 (2021).
https://doi.org:10.1016/j.cell.2021.03.056
28 Gao, P. *et al.* Risk variants disrupting enhancers of TH1 and TREG cells in type 1
diabetes. *Proc Natl Acad Sci U S A* **116**, 7581-7590 (2019).
https://doi.org:10.1073/pnas.1815336116
29 Grun, D., Kester, L. & van Oudenaarden, A. Validation of noise models for single-cell
transcriptomics. *Nat Methods* **11**, 637-640 (2014). https://doi.org:10.1038/nmeth.2930
30 Kolodziejczyk, A. A., Kim, J. K., Svensson, V., Marioni, J. C. & Teichmann, S. A. The
technology and biology of single-cell RNA sequencing. *Mol Cell* **58**, 610-620 (2015).
https://doi.org:10.1016/j.molcel.2015.04.005
31 Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single
cells. *Nat Methods* **14**, 865-868 (2017). https://doi.org:10.1038/nmeth.4380
32 Stoeckius, M. *et al.* Cell Hashing with barcoded antibodies enables multiplexing and
doublet detection for single cell genomics. *Genome Biol* **19**, 224 (2018).
https://doi.org:10.1186/s13059-018-1603-1
33 International Multiple Sclerosis Genetics, C. *et al.* Analysis of immune-related loci
identifies 48 new susceptibility variants for multiple sclerosis. *Nat Genet* **45**, 1353-1360
(2013). https://doi.org:10.1038/ng.2770
34 Soneson, C. & Robinson, M. D. Bias, robustness and scalability in single-cell differential
expression analysis. *Nat Methods* **15**, 255-261 (2018).
https://doi.org:10.1038/nmeth.4612
35 Park, Y. P. & Kellis, M. CoCoA-diff: counterfactual inference for single-cell gene
expression analysis. *Genome Biol* **22**, 228 (2021). https://doi.org:10.1186/s13059-021-
02438-4
36 Chapman, N. M. *et al.* mTOR coordinates transcriptional programs and mitochondrial
metabolism of activated Treg subsets to protect tissue homeostasis. *Nat Commun* **9**,
2095 (2018). https://doi.org:10.1038/s41467-018-04392-5
37 Zhang, F., Liu, G., Li, D., Wei, C. & Hao, J. DDIT4 and Associated lncDDIT4 Modulate
Th17 Differentiation through the DDIT4/TSC/mTOR Pathway. *J Immunol* **200**, 1618-
1626 (2018). https://doi.org:10.4049/jimmunol.1601689

929	38	Loo, C. S. *et al.* A Genome-wide CRISPR Screen Reveals a Role for the Non-canonical
930		Nucleosome-Remodeling BAF Complex in Foxp3 Expression and Regulatory T Cell
931		Function. *Immunity* **53**, 143-157 e148 (2020).
932		https://doi.org:10.1016/j.immuni.2020.06.011
933	39	Yu, X. *et al.* Metabolic control of regulatory T cell stability and function by TRAF3IP3 at
934		the lysosome. *J Exp Med* **215**, 2463-2476 (2018). https://doi.org:10.1084/jem.20180397
935	40	Paulos, C. M. *et al.* The inducible costimulator (ICOS) is critical for the development of
936		human T(H)17 cells. *Sci Transl Med* **2**, 55ra78 (2010).
937		https://doi.org:10.1126/scitranslmed.3000448
938	41	Hasan, Z. *et al.* JunB is essential for IL-23-dependent pathogenicity of Th17 cells. *Nat*
939		*Commun* **8**, 15628 (2017). https://doi.org:10.1038/ncomms15628
940	42	Ma, C. *et al.* Critical Role of CD6highCD4+ T Cells in Driving Th1/Th17 Cell Immune
941		Responses and Mucosal Inflammation in IBD. *J Crohns Colitis* **13**, 510-524 (2019).
942		https://doi.org:10.1093/ecco-jcc/jjy179
943	43	Wang, N. *et al.* CD226 Attenuates Treg Proliferation via Akt and Erk Signaling in an EAE
944		Model. *Front Immunol* **11**, 1883 (2020). https://doi.org:10.3389/fimmu.2020.01883
945	44	Keller, A. D. & Maniatis, T. Identification and characterization of a novel repressor of
946		beta-interferon gene expression. *Genes Dev* **5**, 868-879 (1991).
947		https://doi.org:10.1101/gad.5.5.868
948	45	Martins, G. A. *et al.* Transcriptional repressor Blimp-1 regulates T cell homeostasis and
949		function. *Nat Immunol* **7**, 457-465 (2006). https://doi.org:10.1038/ni1320
950	46	Lin, M. H. *et al.* T cell-specific BLIMP-1 deficiency exacerbates experimental
951		autoimmune encephalomyelitis in nonobese diabetic mice by increasing Th1 and Th17
952		cells. *Clin Immunol* **151**, 101-113 (2014). https://doi.org:10.1016/j.clim.2014.02.006
953	47	Jain, R. *et al.* Interleukin-23-Induced Transcription Factor Blimp-1 Promotes
954		Pathogenicity of T Helper 17 Cells. *Immunity* **44**, 131-142 (2016).
955		https://doi.org:10.1016/j.immuni.2015.11.009
956	48	Bankoti, R. *et al.* Differential regulation of Effector and Regulatory T cell function by
957		Blimp1. *Sci Rep* **7**, 12078 (2017). https://doi.org:10.1038/s41598-017-12171-3
958	49	Ogawa, C. *et al.* Blimp-1 Functions as a Molecular Switch to Prevent Inflammatory
959		Activity in Foxp3(+)RORgammat(+) Regulatory T Cells. *Cell Rep* **25**, 19-28 e15 (2018).
960		https://doi.org:10.1016/j.celrep.2018.09.016
961	50	Garg, G. *et al.* Blimp1 Prevents Methylation of Foxp3 and Loss of Regulatory T Cell
962		Identity at Sites of Inflammation. *Cell Rep* **26**, 1854-1868 e1855 (2019).
963		https://doi.org:10.1016/j.celrep.2019.01.070
964	51	Gyory, I., Fejer, G., Ghosh, N., Seto, E. & Wright, K. L. Identification of a functionally
965		impaired positive regulatory domain I binding factor 1 transcription repressor in myeloma
966		cell lines. *J Immunol* **170**, 3125-3133 (2003).
967		https://doi.org:10.4049/jimmunol.170.6.3125
968	52	Boi, M., Zucca, E., Inghirami, G. & Bertoni, F. PRDM1/BLIMP1: a tumor suppressor
969		gene in B and T cell lymphomas. *Leuk Lymphoma* **56**, 1223-1228 (2015).
970		https://doi.org:10.3109/10428194.2014.953155
971	53	Wang, X. *et al.* High-resolution genome-wide functional dissection of transcriptional
972		regulatory regions and nucleotides in human. *Nat Commun* **9**, 5380 (2018).
973		https://doi.org:10.1038/s41467-018-07746-1
974	54	Consortium, E. P. An integrated encyclopedia of DNA elements in the human genome.
975		*Nature* **489**, 57-74 (2012). https://doi.org:10.1038/nature11247
976	55	Roadmap Epigenomics, C. *et al.* Integrative analysis of 111 reference human
977		epigenomes. *Nature* **518**, 317-330 (2015). https://doi.org:10.1038/nature14248
978	56	Smith, M. A. *et al.* PRDM1/Blimp-1 controls effector cytokine production in human NK
979		cells. *J Immunol* **185**, 6058-6067 (2010). https://doi.org:10.4049/jimmunol.1001682

57   Fragale, A. *et al.* IFN regulatory factor-1 negatively regulates CD4+ CD25+ regulatory T
cell differentiation by repressing Foxp3 expression. *J Immunol* **181**, 1673-1682 (2008).
https://doi.org:10.4049/jimmunol.181.3.1673
58   Hayatsu, N. *et al.* Analyses of a Mutant Foxp3 Allele Reveal BATF as a Critical
Transcription Factor in the Differentiation and Accumulation of Tissue Regulatory T
Cells. *Immunity* **47**, 268-283 e269 (2017). https://doi.org:10.1016/j.immuni.2017.07.008
59   Renoux, F. *et al.* The AP1 Transcription Factor Fosl2 Promotes Systemic Autoimmunity
and Inflammation by Repressing Treg Development. *Cell Rep* **31**, 107826 (2020).
https://doi.org:10.1016/j.celrep.2020.107826
60   Wu, C. *et al.* SGK1 Governs the Reciprocal Development of Th17 and Regulatory T
Cells. *Cell Rep* **22**, 653-665 (2018). https://doi.org:10.1016/j.celrep.2017.12.068
61   Li, Z. *et al.* Identification of transcription factor binding sites using ATAC-seq. *Genome*
Biology **20**, 45 (2019). https://doi.org:10.1186/s13059-019-1642-2
62   Bentsen, M. *et al.* ATAC-seq footprinting unravels kinetics of transcription factor binding
during zygotic genome activation. *Nat Commun* **11**, 4267 (2020).
https://doi.org:10.1038/s41467-020-18035-1
63   Li, P. *et al.* BATF–JUN is critical for IRF4-mediated transcription in T cells. *Nature* **490**,
543-546 (2012). https://doi.org:10.1038/nature11530
64   Murphy, T. L., Tussiwand, R. & Murphy, K. M. Specificity through cooperation: BATF-IRF
interactions control immune-regulatory networks. *Nat Rev Immunol* **13**, 499-509 (2013).
https://doi.org:10.1038/nri3470
65   Gate, R. E. *et al.* Genetic determinants of co-accessible chromatin regions in activated T
cells across humans. *Nat Genet* **50**, 1140-1150 (2018). https://doi.org:10.1038/s41588-
018-0156-2
66   Pham, D. *et al.* Batf Pioneers the Reorganization of Chromatin in Developing Effector T
Cells via Ets1-Dependent Recruitment of Ctcf. *Cell Rep* **29**, 1203-1220 e1207 (2019).
https://doi.org:10.1016/j.celrep.2019.09.064
67   Lee, S. M., Gao, B. & Fang, D. FoxP3 maintains Treg unresponsiveness by selectively
inhibiting the promoter DNA-binding activity of AP-1. *Blood* **111**, 3599-3606 (2008).
https://doi.org:10.1182/blood-2007-09-115014
68   Samstein, R. M. *et al.* Foxp3 exploits a pre-existent enhancer landscape for regulatory T
cell lineage specification. *Cell* **151**, 153-166 (2012).
https://doi.org:10.1016/j.cell.2012.06.053
69   Lam, A. J. *et al.* Optimized CRISPR-mediated gene knockin reveals FOXP3-
independent maintenance of human Treg identity. *Cell Rep* **36**, 109494 (2021).
https://doi.org:10.1016/j.celrep.2021.109494
70   Costantino, C. M., Baecher-Allan, C. & Hafler, D. A. Multiple sclerosis and regulatory T
cells. *J Clin Immunol* **28**, 697-706 (2008). https://doi.org:10.1007/s10875-008-9236-x
71   Sambucci, M. *et al.* FoxP3 isoforms and PD-1 expression by T regulatory cells in
multiple sclerosis. *Scientific Reports* **8**, 3674 (2018). https://doi.org:10.1038/s41598-018-
21861-5
72   Xi, H. & Blanck, G. The IRF-2 DNA binding domain facilitates the activation of the class
II transactivator (CIITA) type IV promoter by IRF-1. *Mol Immunol* **39**, 677-684 (2003).
https://doi.org:10.1016/s0161-5890(02)00214-6
73   Mould, A. W., Morgan, M. A., Nelson, A. C., Bikoff, E. K. & Robertson, E. J.
Blimp1/Prdm1 Functions in Opposition to Irf1 to Maintain Neonatal Tolerance during
Postnatal Intestinal Maturation. *PLoS Genet* **11**, e1005375 (2015).
https://doi.org:10.1371/journal.pgen.1005375
74   Lal, G. *et al.* Distinct inflammatory signals have physiologically divergent effects on
epigenetic regulation of Foxp3 expression and Treg function. *Am J Transplant* **11**, 203-
214 (2011). https://doi.org:10.1111/j.1600-6143.2010.03389.x

He, L. *et al.* NEBULA is a fast negative binomial mixed model for differential or co-expression analysis of large-scale multi-subject single-cell data. *Commun Biol* **4**, 629 (2021). https://doi.org:10.1038/s42003-021-02146-6

Hill, J. A. *et al.* Foxp3 transcription-factor-dependent and -independent regulation of the regulatory T cell transcriptional signature. *Immunity* **27**, 786-800 (2007). https://doi.org:10.1016/j.immuni.2007.09.010

Ferraro, A. *et al.* Interindividual variation in human T regulatory cells. *Proceedings of the National Academy of Sciences* **111**, E1111-E1120 (2014). https://doi.org:10.1073/pnas.1401343111

Konermann, S. *et al.* Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* **517**, 583-588 (2015). https://doi.org:10.1038/nature14136

Fishilevich, S. *et al.* GeneHancer: genome-wide integration of enhancers and target genes in GeneCards. *Database (Oxford)* **2017** (2017). https://doi.org:10.1093/database/bax028

van Galen, P. *et al.* A Multiplexed System for Quantitative Comparisons of Chromatin Landscapes. *Mol Cell* **61**, 170-180 (2016). https://doi.org:10.1016/j.molcel.2015.11.003

Glasmacher, E. *et al.* A genomic regulatory element that directs assembly and function of immune-specific AP-1-IRF complexes. *Science* **338**, 975-980 (2012). https://doi.org:10.1126/science.1228309

Ciofani, M. *et al.* A validated regulatory network for Th17 cell specification. *Cell* **151**, 289-303 (2012). https://doi.org:10.1016/j.cell.2012.09.016

Li, J., Comeau, H. Y., Zhang, Z. & Ren, X. Landscape of transcript isoforms in single T cells infiltrating in non-small-cell lung cancer. *J Genet Genomics* **47**, 373-388 (2020). https://doi.org:10.1016/j.jgg.2020.06.006

Yukawa, M. *et al.* AP-1 activity induced by co-stimulation is required for chromatin opening during T cell activation. *J Exp Med* **217** (2020). https://doi.org:10.1084/jem.20182009

Bossini-Castillo, L. *et al.* Immune disease variants modulate gene expression in regulatory CD4(+) T cells. *Cell Genom* **2**, None (2022). https://doi.org:10.1016/j.xgen.2022.100117

Huhn, K. *et al.* Skin sodium is increased in male patients with multiple sclerosis and related animal models. *Proc Natl Acad Sci U S A* **118** (2021). https://doi.org:10.1073/pnas.2102549118

Lowther, D. E. *et al.* PD-1 marks dysfunctional regulatory T cells in malignant gliomas. *JCI Insight* **1** (2016). https://doi.org:10.1172/jci.insight.85935

Lucca, L. E. *et al.* TIGIT signaling restores suppressor function of Th1 Tregs. *JCI Insight* **4** (2019). https://doi.org:10.1172/jci.insight.124427

Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, 21 29 21-21 29 29 (2015). https://doi.org:10.1002/0471142727.mb2129s109

Buenrostro, J. D. *et al.* Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* **523**, 486-490 (2015). https://doi.org:10.1038/nature14590

Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013). https://doi.org:10.1093/bioinformatics/bts635

Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011). https://doi.org:10.1186/1471-2105-12-323

Harrow, J. *et al.* GENCODE: the reference human genome annotation for The ENCODE Project. *Genome Res* **22**, 1760-1774 (2012). https://doi.org:10.1101/gr.135350.111

94 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140 (2010). https://doi.org:10.1093/bioinformatics/btp616

95 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
https://doi.org:10.1093/nar/gkv007

96 Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome Biology* **15**, R29 (2014).
https://doi.org:10.1186/gb-2014-15-2-r29

97 Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data using factor
analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).
https://doi.org:10.1038/nbt.2931

98 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *2011* **17**, 3 (2011). https://doi.org:10.14806/ej.17.1.200

99 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357-359 (2012). https://doi.org:10.1038/nmeth.1923

100 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137
(2008). https://doi.org:10.1186/gb-2008-9-9-r137

101 Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841-842 (2010).
https://doi.org:10.1093/bioinformatics/btq033

102 Gusmao, E. G., Allhoff, M., Zenke, M. & Costa, I. G. Analysis of computational
footprinting methods for DNase sequencing experiments. *Nat Methods* **13**, 303-309
(2016). https://doi.org:10.1038/nmeth.3772

103 Fisher, R. A. Dispersion on a sphere. *Proceedings of the Royal Society of London.*
Series A. Mathematical and Physical Sciences **217**, 295 - 305 (1953).

104 Banerjee, A., Dhillon, I. S., Ghosh, J. & Sra, S. Clustering on the Unit Hypersphere using
von Mises-Fisher Distributions. *J. Mach. Learn. Res.* **6**, 1345-1382 (2005).

105 Polański, K. *et al.* BBKNN: fast batch alignment of single cell transcriptomes.
Bioinformatics **36**, 964-965 (2019). https://doi.org:10.1093/bioinformatics/btz625

106 Hie, B., Bryson, B. & Berger, B. Efficient integration of heterogeneous single-cell
transcriptomes using Scanorama. *Nature Biotechnology* **37**, 685-691 (2019).
https://doi.org:10.1038/s41587-019-0113-3

107 Unterman, A. *et al.* Single-cell multi-omics reveals dyssynchrony of the innate and
adaptive immune system in progressive COVID-19. *Nat Commun* **13**, 440 (2022).
https://doi.org:10.1038/s41467-021-27716-4

1117