Like mother, like daughter? Phenotypic plasticity, environmental covariation, and

heritability of size in a parthenogenetic wasp

Abstract

Dinocampus coccinellae (Hymenoptera:Braconidae, Euphorinae) is a solitary,

generalist Braconid parasitoid wasp that reproduces through thelytokous

parthenogenesis, an asexual process in which diploid daughters emerge from

unfertilized eggs, and uses over fifty diverse species of coccinellid ladybeetles

worldwide as hosts. Here we utilized a common garden and reciprocal transplant

experiment using parthenogenetic lines of *D. coccinellae* presented with three

different host ladybeetle species of varying sizes, across multiple generations to

investigate heritability, plasticity, and environmental covariation of body size. We

expected positively correlated parent-offspring parasitoid regressions, indicative of

heritable size variation, from unilineal (parent and offspring reared on same host

species) lines, since these restrict environmental variation in phenotypes. In

contrast, because multilineal (parent and offspring reared on different host species)

lines would induce phenotypic plasticity of clones reared in varying environments,

we expected negatively correlated parent-offspring parasitoid regressions. Contrary

to expectations, our results indicate (1) little heritable variation in body size, (2)

strong independence of offspring size on the host environment, (3) a consistent

signal of size-host tradeoff wherein small mothers produced larger offspring, and

vice versa, independent of host environment. Our study offers support for a

constrained fecundity advantage model of Cope's Law, wherein *D. coccinellae*

maintains phenotypic plasticity in body size despite parthenogenetic reproduction.

24
Introduction

Size of an organism is a complex and often plastic trait that is correlated with key adaptive traits

such as reproductive success (Bosch and Vicens 2005, Berger et al., 2012), fecundity (Honek

1993), response to varying environments and hosts (Chown and Gaston 2010), developmental

rates (Davidowitz et al., 2003), survival (Callier and Nijhout 2013), and greater depredation

success (Oliveira et al., 2019). At the same time, larger bodied organisms face challenges such

as increased resource need, and strong evolutionary constraints on reproductive tradeoffs

(Blanckenhorn 2000, Shine 1988), which set "thresholds" on size. Theory therefore predicts that

a fecundity advantage for body size only occurs in the presence of energy availability (Shine

1988). The evolution of organismal size has been studied extensively over speciation

timescales (reviewed in Hone and Benton 2005), often pointing to multiple independent

transitions to larger body size (termed as Cope's Rule) across diverse animal taxa, indicating

that there is no one definitive "pathway" or evolutionary strategy for size among species.

Several lines of evidence instead support that plasticity of body size evolves at

microevolutionary scales (Maurer et al., 1992), with standing genetic variation providing the

basis for adaptability of body size plasticity (Gotanda et al., 2015). The positive correlations

between body size of host and adult offspring are well documented in solitary parasitoid wasps,

as parasitoid offspring reared on larger species and sizes of host develop larger parasitoid

offspring than their parent (Mackauer and Chau, 2001; Arakawa et al., 2004; Wang et al., 2008).

As tradeoffs are ever present across phenotypes, the plasticity of parasitoid body size occurs as

a tradeoff between accessing wider ranges of hosts for generalist parasitoids, while at the

expense of producing varying sizes of adult parasitoid offspring (Henry et al., 2006).

Parthenogenetic wasps provide an ideal natural experimental system to test hypotheses of

plasticity of body size, considering their clonal mode of reproduction that maintains genetic

variation, specifically utilizing a combination of common-garden and reciprocal transplant

experiments to control for genetic and environmental variation.

The parasitoid wasp, *Dinocampus coccinellae* (Hymenoptera: Braconidae), is a generalist that is

capable of successfully parasitizing over fifty species of predatory ladybeetles (Coleoptera:

Coccinellidae, subfamily Coccinellinae) across a global distribution (Balduf, 1926; Ceryngier et

al., 2018, Fei et al 2023). *D. coccinellae* primarily displays solitary behavior, and is only known

to asexually reproduce through thelytoky, a mode of parthenogenesis in which females emerge

from unfertilized eggs; with males rarely observed in this species (Slobodchikoff and Daly, 1971;

Wright, 1979; Heimpel and De Boer, 2008; Ceryngier et al., 2018). Briefly, thelytoky is a

parthenogenetic mode of reproduction in which diploid female adults develop from unfertilized

egg clones (Beukeboom et al., 2007; Heimpel and Jetske, 2008; Slobodchikoff and Daly, 1971).

There are genetic forms of thelytoky in which no crossing over occurs (apomictic thelytoky or

premeiotic doubling) or where the fusion of sister or non-sister recombinant chromosomes form

diploid eggs (automictic thelytoky) (Heimpel and Jetske, 2008), regardless restricting genomic

variation from parent to offspring.

Characteristic to the Euphorinae subfamily of Hymenoptera, a parasitoid larva of *D. coccinellae*

consumes the adipose tissue of a parasitized adult ladybeetle as a koinobiont endoparasitoid,

(Balduf, 1926; Orr et al., 1992; Ceryngier et al., 2018), although it has been documented to

oviposit within host ladybeetle larvae and pupae (Obrycki et al., 1985). Across the diverse range

of host ladybeetles, *D. coccinellae* has been reported to preferentially oviposit in coccinellids

which are more mobile, larger, adult, female hosts (Davis *et al*., 2006; Obrycki, 1989). Once an

adult *D. coccinellae* locates a sufficient adult ladybeetle, they arch their stinger under the beetle

and thrust into the abdomen of the host, injecting clonal daughter egg(s) along with

accompanying venom enzymes and the RNA-virus, the *Dinocampus coccinellae* Paralysis Virus

(DcPV) (Balduf, 1926; Orr *et al.,* 1992; Dheilly *et al*., 2015). This is yet another unique facet of

the *D. Coccinellae*, as their life cycle involves an endosymbiotic relationship established with

DcPV, an RNA virus in the Iflaviridae family (Dheilly et al., 2015). In concert with host behavior

modifications mediated by this virus, *D. coccinellae* then use their captive adult host as a

bodyguard to the advantage of the next generation. After approximately a week following

oviposition within a host beetle, the larva emerges from its egg into the fat body of the host's

abdomen, where it undergoes four larval instar stages of development (Balduf, 1926). Multiple

eggs may be deposited within the same host, which is referred to as superparasitism, which has

been documented in several field studies (summarized by Ceryngier et al. 2012). When this

occurs, the first larva to emerge crushes the others with its mandibles (Balduf, 1926). In these

cases of superparasitism, the survivor then cannibalizes its host-mate(s) as its first meal;

otherwise, the larva feeds on adipose tissue and ovaries of coccinellid host throughout

development (15-20 days) (Balduf, 1926). Tetratocytes, which originate from the parasitoid's

egg, aid in providing an initial food source, in addition to the host itself (Okuda *et al.*, 1995).

Following pupation in an external cocoon, the daughter wasp emerges as an adult with fully

developed eggs, with some of these females leaving a varying percentage of their hosts alive

(Orr et al., 1992). The intricate behavioral relationship between an adult *D. coccinellae* wasp

and its host ladybeetle have been described, with successful parasitization, measured as the

percentage of emerged daughter wasps as a proxy for fecundity, varying between different host

species (Orr et al., 1992). However, little is known about fitness of the emergent

parthenogenetic daughter wasps.

A previous study by Vansant et al. (2019), found positive relationships between host and

emergent daughter *D. coccinellae* morphology (e.g. dry mass, wing length, abdominal length),

which has been found to be the case in a variety of parasitoid wasps and their hosts (Brandl and

Vidal, 1987; Mackauer and Chau, 2001; Harvey et al., 2006; Henry et al., 2006; Symonds and

Elgar, 2013). The developmental environmental conditions, including resources that a

developing parasitoid can uptake from its host, substantially determines the body size

phenotype of the emerging parasitoid. Additive genetic effects on the phenotype are expected to

be heritable, and the relative importance of the development environment and additive genetic

effects in determining the phenotype of *D. coccinellae* are not known. However, as *D.*

coccinellae reproduces via thelytoky with little to no genetic recombination, this brings into

question the balance between heritability or phenotypic plasticity of body size as a proxy for

individual fitness.

To test and quantify the heritability and phenotypic plasticity of body size, we established

multigenerational experiments with unilineal and multilineal reciprocal crosses of three host

ladybeetle species exposed to parthenogenetic lines of *D. coccinellae*. Given that total

phenotypic variation of a trait is composed of genetic and environmental variation, a unilineal

line consisting of rearing wasps and resultant clonal daughters on a single host beetle species

background would have minimal variation in size due to environmental effects. Restricting the

environmental variability to a single host species would produce no relationship between parent

and offspring trait values if size is entirely phenotypically plastic, but would produce a positive

relationship if offspring trait values were due to additive genetic variation. Alternately, multilineal

lines, with daughters raised on hosts that were a different size from their mothers, would

produce negative relationships between parent and offspring trait values if size is entirely

phenotypically plastic. Additive genetic effects may cause a residual positive correlation

between mother and daughter pairs after host size has been accounted for in multilineal lines.

The relative importance of additive genetic effects and developmental environment could then

be assessed by combining unilineal and multilineal lines in a single analysis.

Materials and Methods

Experimental setup

D. coccinellae wasps used to start the lineages for reciprocal host-transplants were

obtained from field collections in Kentucky of parasitized adult *Coccinella septempunctata* (*C.*

septempunctata or C7 - JJO personal comm.) and from *Hippodamia convergens* (*H.*

convergens or H. con) from an insectary in San Marcos, CA. Parthenogenetic lines of *D.*

coccinellae were then maintained for at least 4 generations on laboratory populations of three

species of lady beetles – C7, H. con, and *Coleomegilla maculata* (*C. maculata* or C. mac) which

were obtained from field sites in Kentucky (JJO personal comm.). These beetle populations

were maintained on an *ad libitum* diet of *Acyrthosiphon pisum* (pea aphids), which in turn were

maintained on fava bean plants (*Vicia faba*) in insect tents in the California State University San

Marcos (CSUSM) greenhouse in San Marcos, CA until March 2020. Following the COVID-19

outbreak, all insect tents, subsequent crosses, and experimentation were performed (socially

distanced and masked) in AT's garage in Oceanside, CA. Despite temporary relocation of the

experimental setup, all experimental conditions were maintained constant to minimize random

effects, including daily variations in temperature and diurnal cycles. The unlineal and multilineal

experiments were conducted in a single location, with the same species of three hosts, and

therefore approximately evenly distributed across treatments.

In each experimental setup, one adult *D. coccinellae* wasp ('mother') was placed into a paper

soup cup along with four individual ladybeetle hosts, moth (*Ephestia*) eggs for hosts to feed on

and a honey-water soaked cotton ball for both the wasp and beetles to drink from; only one

wasp was introduced per each experiential cup setup and was sealed using a mesh sheet and

an open-face lid. After the mother wasp oviposited into her hosts and died (which always

occurred before her daughters egressed from their cocoons), she was paired with the host that

she emerged from. The remaining four host ladybeetles were fed and tended to until the initial

appearance of the cocoon spun by the larval *D. coccinellae* parasitoid. Once finished

developing in her cocoon, an adult *D. coccinellae* 'daughter' egressed from her cocoon. This

'daughter' *D. coccinellae* was then placed in another experimental cup setup as the next

'mother' *D. coccinellae* with another four individual ladybeetles of the next type of host

coccinellid species, *Ephestia* (moth) eggs for hosts to feed on and honey-water for both host

and wasp to drink. In every introduction cup, the life history data recorded were: wasp

introduction date, parent removal and collection date, cocoon date (if noticed), daughter

eclosion date, and host mortality rate. Mothers were picked at random to oviposit in unilineal or

multilineal conditions.

Parasitized beetles were reared until *D. coccinellae* larva egression from the infected

host as a cocoon woven between the host legs (Vansant et al., 2019). 92 wasp-host pairs and

40 mother-daughter pairs were collected for morphological observations. An expanded

polystyrene foam stage and ruler (mm) was assembled to standardize and scale the

photographed parasitoid-host pairs. Using a Nikon dissection microscope, adult *D. coccinellae*

wasps were photographed in the lateral position, and the corresponding ladybeetle host was

photographed from the dorsal, lateral, and ventral positions. These images were uploaded into

ImageJ (NIH) to obtain the following morphometric measurements in mm for the wasp: head

length, head depth, thorax length, thorax depth, abdomen length and wing length (Figure 1);

and for the host beetle: dorsal body length and depth; lateral body depth, elytron chord length

and pronotum length; and ventral pronotum width, and abdominal length and width (Figure 2);

based on body segments measured in Vansant et al., 2019. Morphometric measurements were

repeated independently by four individuals and averaged to control for observational bias. Each

parent and offspring wasps were then paired for regression analysis, in addition to pairing host

beetle and emergent wasp measurements.

Statistical analysis

All statistical analyses and visualization were performed in R v.4.2.1 (R core Team),

with additional libraries noted below.

Size distributions of all wasp and host ladybeetle morphological variables were

visualized as box plots, grouped by the host species from which the wasp eclosed. Differences

between host species morphologies were tested using one-way ANOVA followed by Tukey HSD

post-hoc procedures for each host morphological variable. Additionally, due to the similarity in

sizes of *C. maculata* and *H. convergens* hosts, we grouped them into a "Small" group and

compared them to the "Large" *C. septempunctata*. Similarly, box plots, and one-way ANOVA

were performed across *D. coccinellae* morphological measurements, with host size as a factor.

The multivariate relationship between host morphology and morphology of *D. coccinellae* that

developed in the host was assessed using canonical correlation analysis (CCA). The body

segment measurements for *D. coccinellae* and their hosts were used (Figure 2). Statistical

significance was assessed using an F approximation of Wilks' Lambda (Rao's F), using the

CCP package in R (CCP version 1.2, Menzel 2022).

Heritabilities of wasp morphological variables were measured with parent/offspring regression.

Crosses were assigned to either the unilineal group (when parent and offspring host species

were the same) or the multilineal group (when parent and offspring host species differed), and

type of cross was included as a factor in the parent/offspring regressions. .odels were fitted

using sum contrasts, such that the slope for the main effect of parent morphology represented

the overall parent/offspring relationship across the two different types of cross. The interaction

of cross type with parent host morphology tested for differences in slope of parent/offspring

relationships between unilineal and multilineal crosses. When a significant interaction occurred,

indicating that unilineal and multilineal slopes differed from one another, we used the *emmeans*

package (Lenth 2022) to test the slopes for difference from zero. A second set of models with

parent host species replacing cross type as a factor in the models, used a similar approach, but

differences in parent/offspring slope between the three host species were tested using post-hoc

procedures. Parent morphology was not orthogonal to the categorical predictor in either sets of

linear models. To determine whether the confounded variation between morphology and the

grouping variable (i.e. cross type or parent host species) was responsible for significant

relationships between parent and offspring morphology, two alternative models were fitted for

each morphological variable with parent morphology entered either before or after the factor,

and then the two alternative orders of entry were tested with Type I (sequential sums of

205 squares) ANOVA. Cases in which a significant parent/offspring regression became non-

206 significant after accounting for the factor were noted (Supplemental Data File).

207   To better understand the association between mother and daughter morphology, CCA

208 was conducted relating mother and daughter morphological variables (mass was not included

209 because of missing values, which would have further reduced the sample size). Initially, the

210 canonical correlation between mother and daughter morphology was assessed without

211 accounting for host type. Then, to determine how much of the canonical correlation between

212 mother and daughter was due to either mother or daughter host environment, partial CCA was

213 conducted using the residuals from a MANOVA that included mother's host, daughter's host, or

214 the combination of both mother's and daughter's host as predictors. Partial CCA that controlled

215 for mother's host in mother's morphology and daughter's host in daughter's morphology, and for

216 mother's host in mother's morphology and combinations of mother's and daughter's hosts in

217 daughter's morphology were also used. Large reductions in canonical correlation between

218 mother and daughter when host was accounted for would indicate that the correlation was

219 principally due to host-mediated effects (e.g. developmental environment, mother's investment

220 decisions at oviposition), whereas stable patterns of correlation after host effects had been

221 accounted for would be consistent with factors driven by the mother's state, independent of the

222 host she developed in or oviposited on. All partial CCA analyses were conducted with the *yacca*

223 package (version 1.4-2, Butts 2022), which included a measure of statistical redundancy

224 between the mother's and daughter's morphological variables. Statistical significance of

225 canonical correlations was assessed using an F approximation of Wilks' Lambda (Rao's F),

226 using the CCP package in R (CCP version 1.2, Menzel 2022).

228 **Results**

229   To establish that the three separate ladybeetle species do indeed provide parasitoid *D.*

230 *coccinellae* larva with significantly different environmental conditions to develop within, we

generated box plots and ran one-way ANOVA between the ladybeetle species across each

body segment (Figure 3a-i). *C. maculata* and *H. convergens* were much more similar in size

than either were to *C. septempunctata*, and some morphological variables were not significantly

different between them (body length, abdominal length, mass). But, once *C. maculata* and *H.*

convergens were grouped together under the 'Small' label and *C. septempunctata* being the

'Large' label (Figure 3a-i), all differences were statistically significant.

237   *D. coccinellae* grown on different host sizes only differ in abdominal length, (Figure 4).

238   The two strongest correlations between host and *D. coccinellae* morphological variables

are between ladybeetle abdominal width and *D. coccinellae* head depth (r = 0.51), and between

ladybeetle abdominal width (V) and *D. coccinellae* thorax length (r = 0.45). The multivariate

correlation between host and *D. coccinellae* morphology is much higher; the first canonical

correlation coefficient R = 0.83 was statistically significant (p = 0.014), but the second through

sixth were not (p > 0.05). The first canonical correlation axis represents a positive relationship of

host abdominal width (loading = -0.94) with *D. coccinellae* size across every variable but head

length (loadings ranging from -0.21 for thorax depth to -0.56 for head depth; Figure 5).

246   Narrow sense heritability (h^2) across each body segment measurement was captured by

the slope of the line of the main effect of the parent morphological variable parent-offspring

linear models (Figures 6a-g and 7a-f). Additionally, the parent-offspring regressions in Figures

6a through 6g distinguish between unilineal and multilineal crosses, whose slopes are

represented by the parent morphology by cross type interaction. Four out of seven unilineal

parent-offspring regressions indicate a slight positive slope, but for some variables they are

negative (wing length, thorax length, head depth), and none of the unilineal slopes are

significantly different from 0 (n = 15). Alternately, regression slopes for all multilineal parent-

offspring pairs uniformly display a slight positive relationship [total n = 25], and in the one

variable (wing length) that had a significant difference in slopes between cross types only the

positive multilineal slope significantly differed from zero (Fig. 6a). The overall parent/offspring

relationship, combined across cross type, was significant for thorax depth (Fig. 6c) and

abdomen length (Fig. 6d), but the slopes did not differ by type of cross for these two variables.

Using parent host species to group the data yielded significant parent/offspring

relationships only for thorax depth (Fig. 7c) and abdomen length (Fig. 7d) as before, but

confounding between parent host species and parent morphology made these relationships

dependent on the order that parent morphology was entered into the model; when they were

entered first they were significant, but when they were entered second they were not. Parent

hosts only differed in their slopes for head depth (Fig. 7e), with *C. maculata* and *C.*

septempunctata differing from one another, and neither differing from *H. convergens*. The

negative slope for *C. maculata* was significantly different from zero, as was the positive slope for

C. septempunctata.

For every CCA of parent and offspring morphology only the first canonical axis was statistically

significant (Rao's F approximation, p-values ranging from 0.002 to 0.02). All of the models

yielded a qualitatively consistent pattern (Figure 8), in which the parent loadings were primarily

(for the "No partial" model that did not account for any host effects) or entirely positive and

offspring loadings were negative except for wing length. The canonical correlation coefficients

were also very consistent, ranging from a low of 0.76 for the "No partial" to a high of 0.8 for the

model in which the combinations of mother and daughter host were accounted for in both the

mother and daughter morphologies. The contrast between mother and daughter loadings

increased in any model that the mother's or daughter's host was accounted for, with all of the

mother's loadings becoming positive for those models.

Discussion

The unique life history strategies of *D. coccinellae* - thelytokous parthenogenesis,

solitary behavior, and the ability to successfully oviposit in an uncharacteristically large range of

host lady beetle species that span a wide spectrum of body sizes and shapes (Balduf, 1926;

Ceryngier et al., 2012, 2018; Wright, 1979) - present a great opportunity to understand the

dynamics of phenotypic microevolution of size. This parasitoid attacks a group of predatory

beetles that are widely used in biological control; our study highlights the importance of

examining the genetic bases of ecological interactions underlying parasitoid-host relationships

(Fei et al 2023, Rodrigues et al 2022, Sentis et al 2022).

Specifically, the diversity in host coccinellid morphology offers *D. coccinellae* (1) different host-

parasitoid conflicts (Orr et al., 1992), (2) different environmental niches for their larvae to

develop in, and (3) varying amounts of adipose tissue to feed upon. Therefore, we would predict

that phenotypic plasticity in *D. coccinellae*'s ability to successfully parasitize its hosts offers the

species a selective advantage at microevolutionary scales, while an occasional sexual

reproductive cycle with a male (Shaw et al., 1999) offers an "escape" from Muller's ratchet (i.e.

irreversible accumulation of deleterious variants towards extinction). It has been well

documented that variation in parasitoid wasp morphology is strongly associated with variation in

body size and morphology of host species (Belshaw *et al.,* 2003; Symonds and Elgar, 2013).

Furthermore, previous research indicates that the environmental variation in host lady beetle

body size strongly influences the body size phenotype of each emergent *D. coccinellae*, with

each next clonal generation being capable of significant size changes relative to the parent

(Vansant *et al.*, 2019).

In this study, we utilize a common-garden, reciprocal transplant experiment over multiple

generations to investigate the variation in body size morphology of emergent *D. coccinellae*

conditioned on (1) the clonal parent, and (2) its host. Our study clearly points to the dependence

of body size morphology and phenotypes of emergent *D. coccinellae* on its clonal parent, further

bolstering the idea of a plastic response to maintain size variation in the species at

microevolutionary scales. As *D. coccinellae* reproduces through thelytoky, the process of

asexual reproduction in which diploid female parasitoids are born from unfertilized eggs, it can

reasonably be expected that body size morphometric traits would exhibit strong correlation as

estimated using parent-offspring regressions, as there is no source of additional genetic

variation to affect relatedness through sexual reproduction and recombination or dominance

(Slobodchikoff and Daly, 1971; Heimpel and De Boer, 2008). Yet, anything but a strong

relationship is observed in our results. Across both the unilineal and multilineal parent-offspring

regressions, most of the relationships return non-significant linear slopes, which imply that there

is no difference from regression slopes of zero, indicating that there is extremely low heritable

variation of size (power analysis estimated that we had an 80% chance of detecting slopes of

0.41 or larger in parent/offspring regressions). This was an interesting finding, as we considered

thelytokous parthenogenesis to be such a strong constrictor on genetic variation, that the

significant shift in body size would have been expected to be at least partially evident in body

size plasticity. This experiment also points to how low heritability could emerge from intense

selection (here artificial). It is possible that an adult *D. coccinellae* can feasibly jump to a

different species of host ladybeetle than that of their mother, given the available distribution of

phenotypic variation in body size across one generation. Yet, repeated host shifts in a rapid

succession of a few generations may introduce too intense an artificial selection pressure for

this trait plasticity to endure, limiting the variation in body size variation of the following *D.*

coccinellae generations. Therefore, as a result of negligible additive genetic variance in body

size morphometric traits, we would also predict that there is little trait variability for natural

selection to act on/work with, thereby minimizing the trait's ability to evolve. This is further

complemented by the lack of significant differences in body size morphometric traits in

emergent *D. coccinellae* among all host types as observed in our experiment. Further analyses

of our controlled reciprocal transplant experiments, to quantify the fecundity of *D. coccinellae*

females, perhaps differentially across different hosts would help predict the fitness

consequences of natural selection on plastic size in these predominantly asexual species.

Additionally, of potential interest then is the differential efficacy of parasitization of small

D. coccinellae on smaller versus larger coccinellid hosts. It has been predicted that host

manipulation via "bodyguard" behavior (Maure et al., 2011, Maure et al., 2013) to protect *D.*

coccinellae pupae from predators is presumably under selection for larger hosts, to possibly

repel larger predatory species, e.g., crickets or carabid beetles. This hypothesis can also be

tested by studying the fecundity, survival, duration of "bodyguard" behavior, and parasitization

rates of emergent *D. coccinellae* across different Coccinellid hosts, while controlling for host

size. It has also been noted that the sex of the coccinellid host, and prey availability in the field

could also influence variability in size of adults (Belnavis, 1988), which were not controlled in our

study.

Multivariate comparison of mother and daughter morphology yielded evidence that

mothers produce offspring that differ from them, independent of the host species. Mother's

loadings on the first canonical correlation axis were positive, while their daughter's morphology

had negative loadings, except for their wing lengths. This suggests that across all

microevolutionary scenarios in our experiment, large mothers produce small daughters with long

wings, while small mothers produce large daughters with short wings. A small body size with

long wings is consistent with better dispersal ability (summarized in Johannson et al., 2009),

and it is possible that large females are preferentially producing daughters that will disperse

greater distances. Smaller mothers that produce large daughters with short wings may be

maximizing the survival probability of their daughters at the expense of their potential dispersal

distances. This pattern of increase in size of koinobionts such as Braconid wasps has also been

previously reported to be correlated with increased longevity and fecundity (Boivin 2010). For

example, *Trissolcus mitsukurii* (Hymenoptera: Scelionidae) which were reared on larger hosts

had higher fecundity than individual parasitoids which were reared on smaller hosts (Arakawa et

al., 2004). Moreover, it was observed that the number of eggs a female *Spathius agrili*

(Hymenoptera: Braconidae) oviposits is positively correlated with host size, in addition to other

explanatory factors (Wang et al., 2008). Since this pattern is independent of both mother and

daughter host species, it is likely that it is mediated by the mother's state. Our observations

therefore offer partial support for Darwin's fecundity-advantage model, but limited by energy

availability from the host environment (Shine 1988). The mechanism for producing these

changes is unknown, but facultative changes in the size of eggs, modified fecundity on energy

availability, or epigenetic regulation of gene expression are some combination of the above are

possible. The phenotypic variance caused by mothers producing daughters who are genetically

identical but are morphologically different would further reduce the narrow sense heritability of

traits, and could explain negative slope estimates for some of the traits.

367   Finally, our results also bring into question the micro- and macroevolutionary

consequences of the evolution and maintenance of thelytokous parthenogenesis from ancestral

arrhenotoky in these species. A recent study on the *D. coccinellae* genome by Sethuraman et

al., 2022 pointed to an early divergence, accelerated rates of genome evolution via manifold

duplications and gene loss along the *D. coccinellae* lineage. Significant duplication events were

reported in transposase activity and stress response gene families, while significant gene losses

were reported among olfactory/odorant receptors and viral-coevolution genes. We surmise that

these duplication (and loss) events contribute to standing genomic variation in *D. coccinellae*

that permit plasticity of size despite parthenogenetic reproduction and alternating reproductive

trade-offs depending on host availability and host-associated energy limitations, independent of

maternal genetics.

Data Availability

All high definition images utilized in assessment of size in this study have been made available

via FigShare (https://doi.org/10.6084/m9.figshare.21685181.v1). All statistical analyses and

morphological data used in this study are accessible via R Markdown at:

https://wkristan.github.io/toval_etal_supplement.zip

References

Arakawa, R., Miura, M., & Fujita, M. (2004). Effects of host species on the body size, fecundity,

and longevity of Trissolcus mitsukurii (Hymenoptera: Scelionidae), a solitary egg

parasitoid of stink bugs. *Applied Entomology and Zoology*, *39*(1), 177-181.

Balduf, W.V., 1926. The bionomics of *Dinocampus coccinellae* Schrank. *Annals of the*

Entomological Society of America, *19*(4), pp.465-498.

Belshaw, R., Grafen, A. and Quicke, D.L., 2003. Inferring life history from ovipositor morphology

in parasitoid wasps using phylogenetic regression and discriminant analysis. *Zoological*

Journal of the Linnean Society, *139*(2), pp.213-228.

Berger, D., Olofsson, M., Friberg, M., Karlsson, B., Wiklund, C., and Gotthard, K., 2012.

Intraspecific variation in body size and the rate of reproduction in female insects -

adaptive allometry or biophysical constraint?. *Journal of Animal Ecology, 81(6),* pp.1244-

1258.

Blanckenhorn, W., 2000. The evolution of body size: what keeps organisms small? *The*

Quarterly Review of Biology, 75(4), pp. 385-407.

Boivin, G., 2010. Phenotypic plasticity and fitness in egg parasitoids. *Neotropical Entomology*,

39, pp.457-463.

Bosch, J. and Vicens, N., 2006. Relationship between body size, provisioning rate, longevity

and reproductive success in females of the solitary bee *Osmia cornuta. Behavioral*

Ecology and Sociobiology, 60, pp.26-33.

Brandl, R. and Vidal, S., 1987. Ovipositor length in parasitoids and tentiform leaf mines:

adaptations in eulophids (Hymenoptera: Chalcidoidea). *Biological Journal of the Linnean*

Society, 32(4), pp.351-355.

Butts, C. T. 2022. yacca: Yet Another Canonical Correlation Analysis Package. R package

version 1.4-2, https://CRAN.R-project.org/package=yacca.

Callier, V. and Nijhout, H., 2013. Body size determination in insects: a review and synthesis of

size- and brain- dependent mechanisms. *Biological Reviews, 88(4),* pp. 944-954.

Cerynigier, P., H.E. Roy, and R.L. Poland. 2012. Natural enemies of ladybird beetles. Pgs. 375-

443. In I, Hodek, H.F. van Emden, and A. Honek (eds). Ecology and Behaviour of

the Ladybird Beetles (Coccinellidae). Wiley-Blackwell, New York.

Ceryngier, P., Nedvěd, O., Grez, A.A., Riddick, E.W., Roy, H.E., San Martin, G., Steenberg, T.,

Veselý, P., Zaviezo, T., Zúñiga-Reinoso, Á. and Haelewaters, D., 2018. Predators and

parasitoids of the harlequin ladybird, *Harmonia axyridis*, in its native range and invaded

areas. *Biological Invasions*, *20*(4), pp.1009-1031.

Chown, S. and Gaston, K., 2010. Body size variation in insects: a macroecological perspective.

Biological Reviews, 85(1), pp. 139-169.

Davidowitz, G., D'Amico, L., and Nijhout, H., 2003. Critical weight in the development of insect

body size. *Evolution & Development, 5(2),* pp. 188-197.

Fei, M., Gols, R., and Harvey, J.A. 2023. The biology and ecology of parasitoid wasps of

predatory arthropods. Annu. Rev. Entomol. 68: 109-128.

González, I and Déjean, S (2021). CCA: Canonical Correlation Analysis. R package version

1.2.1, https://CRAN.R-project.org/package=CCA

Gotanda, K., Correa, C., Turcotte, M., Rolshausen, G., and Hendry, A., 2015. Linking

macrotrends and microrates: re-evaluating microevolutionary support for Cope's rule.

Evolution, 69(5), pp.1345-1354.

Harvey, J.A., Vet, L.E., Witjes, L.M. and Bezemer, T.M., 2006. Remarkable similarity in body

mass of a secondary hyperparasitoid *Lysibia nana* and its primary parasitoid host

Cotesia glomerata emerging from cocoons of comparable size. *Archives of Insect*

Biochemistry and Physiology: Published in Collaboration with the Entomological Society

of America, *61*(3), pp.170-183.

Heimpel, G.E. and De Boer, J.G., 2008. Sex determination in the Hymenoptera. *Annu. Rev.*

Entomol., *53*, pp.209-230.

Henry, L.M., Roitberg, B.D. and Gillespie, D.R., 2006. Covariance of phenotypically plastic traits

induces an adaptive shift in host selection behaviour. *Proceedings of the Royal Society*

B: Biological Sciences, *273*(1603), pp.2893-2899.

Hone, D. and Benton, M., 2005. The evolution of large size: how does Cope's Rule work?

Trends in Ecology & Evolution, 20(1), pp. 4-6.

Honĕk, A., 1993. Intraspecific variation in body size and fecundity in insects: a general

relationship. *Oikos, 66(3),* pp.483-492.

Johansson, F., Söderquist, M., and Bokma, F., 2009. Insect wing shape evolution: independent

effects of migratory and mate guarding flight on dragonfly wings. *Biological Journal of*

the Linnean Society, 97(2), pp. 362-372.

Lenth, R.V. 2022. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.7.3, https://CRAN.R-project.org/package=emmeans

Mackauer, M. and Chau, A., 2001. Adaptive self superparasitism in a solitary parasitoid wasp:

the influence of clutch size on offspring size. *Functional Ecology*, *15*(3), pp.335-343.

Maurer, B., Brown, J., and Rusler, R., 1992. The micro and macro in body size evolution.

Evolution, 46(4), pp. 939-953.

Obrycki, J.J. 1989. Parasitization of native and exotic coccinellids by Dinocampus coccinellae

(Hymenoptera: Braconidae). J. Kansas Entomol. Soc. 62: 211-218.

Obrycki, J.J., Tauber, M.J. and Tauber, C.A., 1985. *Perilitus coccinellae* (Hymenoptera:

Braconidae): parasitization and development in relation to host-stage attacked. *Annals of*

the Entomological Society of America, *78*(6), pp.852-854.

Oliveira, T., Monteiro, A., Morales-Silva, T., Maia, L., and Faria, L., 2019. Multitrophic

interactions drive body size variations in seed-feeding insects. *Ecological Entomology,*

45(3), pp. 538-546.

Orr, C.J. & J.J. Obrycki. 1992. Acceptance behavior of Dinocampus coccinellae (Shrank)

(Hymenoptera: Braconidae). Ann. Entomol. Soc. Am. 85: 722-730.

Rodrigues, L.R., M. Montserrat, and S. Magalhaes. Evolution in agricultural systems: Moving

toward the understanding of complexity. Evol. Appl. 15: 1483-1489.

Sentis, A., J.-L. Hemptinne, A. Magro, and Y. Outreman. 2022. Biological control needs

evolutionary perspectives of ecological interactions. Evol. Appl. 15: 1537-1554.

Sethuraman, A., Tovar, A., Welch, W., Dettmers, R., Arce, C., Skaggs, T., Rothenberg, A.,

Saisho, R., Summerhays, B., Cartmill, R. and Grenier, C., 2022. Genome of the

parasitoid wasp Dinocampus coccinellae reveals extensive duplications, accelerated

evolution, and independent origins of thelytokous parthenogeny and solitary behavior.

G3, *12*(3), p.jkac001.

Shaw, M.R., Geoghegan, I.E. and Majerus, M.E.N., 1999. Males of *Dinocampus coccinellae*

(Schrank)(Hym.: Braconidae: Euphorinae). *Entomologist's Record and Journal of*

Variation, *111*, pp.195-196.

Shine, R., 1988. The evolution of large body size in females: a critique of darwin's "Fecundity

Advantage" model. *The American Naturalist, 131(1),* pp. 124-131.

Slobodchikoff, C.N. and Daly, H.V., 1971. Systematic and evolutionary implications of

parthenogenesis in the Hymenoptera. *American Zoologist*, *11*(2), pp.273-282.

Symonds, M.R. and Elgar, M.A., 2013. The evolution of body size, antennal size and host use in

parasitoid wasps (Hymenoptera: Chalcidoidea): a phylogenetic comparative analysis.

PLoS One, *8*(10), p.e78297.

Vansant, H., Vasquez, Y.M., Obrycki, J.J. and Sethuraman, A., 2019. Coccinellid host

morphology dictates morphological diversity of the parasitoid wasp *Dinocampus*

coccinellae. *Biological Control*, *133*, pp.110-116.

Wang, X. Y., Yang, Z. Q., Wu, H., & Gould, J. R. (2008). Effects of host size on the sex ratio,

clutch size, and size of adult Spathius agrili, an ectoparasitoid of emerald ash borer.

Biological Control, *44*(1), 7-12.

Wright, E.J., 1978. Observations on the copulatory behaviour of *Perilitus coccinellae*

(Hymenoptera: Braconidae).

Figures and Tables

Figure 1 Body segment morphometric traits measured from adult *D. coccinellae* parasitoid

wasps, shown in lateral view with a millimeter scale on the stage. These traits were selected

based on the morphometric segments outlined in Vansant *et al*., 2019.

Figure 2 Body segment morphometric traits measured from host lady beetles which the parasitoid *D. coccinellae* egressed from. Shown from dorsal (Fig. 2a), lateral (Fig. 2b), and ventral (Fig. 2c) perspectives, with a millimeter scale on the stage. These traits were selected based on the morphometric segments outlined in Vansant *et al.*, 2019.

Full Text: 2a) "Body Width: Perpendicular to the midpoint of the body length vector", "Body Length: anterior tip of head to the midpoint of the distal elytra tips." 2b) "Elytron Chord Length: from dorsal pronotum – elytron join to distal elytron apex," "Body Depth: From ventral abdomen at base of hind leg to the top of elytron, perpendicular to the elytron chord vector", and "Pronotum Length: From anterior pronotum point to dorsal intersection of pronotum-elytron." 2c) "pronotum Width: Lateral extremes of pronotum," "Abdominal Length: From prosternum to the midpoint of distal tips at abdomen-elytra joint", and "Abdominal Width: Lateral-most points across abdomen, perpendicular to the abdominal length vector."

Figure 3a-3h Boxplots of morphometric variables measured for *Coleomegilla maculata*, *Hippodamia convergens*, and *Coccinella septempunctata* host ladybeetles across the dorsal, lateral, and ventral image viewpoints. P-values from Tukey post-hoc comparisons are indicated above box plots. Significance level for comparisons between small (combined *H. convergens* and *C. maculata*) and large (*C. septempunctata*) ladybeetles are indicated as asterisks on legend title (* < 0.05, ** < 0.01, *** < 0.001).

3a)

3b)

3c)

3d)

3e)

3f)

3g)

3h)

3i)

Figure 4 Boxplots for morphometric variables measured for *D. coccinellae*, separated by which host ladybeetle species the parasitoid egressed from ('Small' and 'Large' host categories are the same for the host ladybeetle analyses). The p-values are comparisons of means by one-way ANOVA.

4a)

4b)

4c)

4d)

4e)

4f)

4g)

Figure 5. Canonical Correlation Analysis depicting the positive correlation between *D. coccinellae* morphological variables and host ladybeetle morphological variables. Variables for hosts are BL = body length, BW = body width, ECL = elytron chord length, BD = body depth, PL = pronotum length, PW = pronotum width, AL = abdominal length, AW = abdominal width. Variables for wasps are WL = wing length, TL = thorax length, TD = thorax depth, AL = abdomen length, HD = head depth, and HL = head length.

Figure 6 Analysis of covariance relating offspring morphology to parent morphology, grouped by type of cross. Unilineal crosses used the same host species for both parent and offspring, and multilineal crosses used different host species for parent and offspring.. The gray line gives the overall relationship between parent and offspring, and its regression equation and coefficient of determination are shown in black (asterisks indicate p < 0.05). Unilineal and multilineal

regression equations and coefficients of determination match the color for their lines and plot symbols (an asterisk on the legend title indicates a significant Type x parent morphology interaction, and an asterisk next to a type label indicates a slope significantly different from zero for that group).

6a)

6b)

6c)

6d)

6e)

6f)

6g)

Figure 7 Parent offspring analysis of covariance models, grouped by parent host. The overall relationship between parent and offspring is shown with a thick gray line, and its regression equation and coefficient of variation are shown in black. Regression equations and coefficients of variation are colored to match lines and plot symbols for host species. Symbols indicate: * = p < 0.05, † = p < 0.05 only when entered first using sequential sums of squares. Letters after host species names in 7e are post-hoc comparison letters, and * indicates slopes that are significantly different from 0.

7a)   7b)

Figure 8. Summary of the results of canonical correlation analysis between mother and daughter morphology. Each model is shown as a row, labeled along the y-axis. Points give variable loadings (correlation between variable and CCA1 axis), and are labeled by variable abbreviation. The canonical correlation coefficient for CCA1 is given to the right of the points.