Estimating Biochemical Concentration in Food Using Untargeted Metabolomics

Michael Sebek[1,+],   Giulia Menichetti[1,2,+],

Albert-László Barabási[1,2,3†]

[1]Network Science Institute and Department of Physics, Northeastern University, Boston, MA,

USA

[2]Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA,

USA

[3]Department of Network and Data Science, Central European University, Budapest, Hungary

[+]These authors contributed equally to this work

Abstract

Untargeted metabolomics can detect hundreds of biochemicals in food, yet without standards, it cannot quantify them. Here we show that we can take advantage of the universal scaling of nutrient concentrations to estimate the concentration of all biochemicals detected by untargeted metabolomics. We validate our method on 20 raw foods, finding an excellent agreement between the predicted and the experimentally observed concentrations.

†Corresponding author e-mail: barabasi@gmail.com

The documented impact of diet on health has prompted national efforts to catalogue the biochemical composition of food[1], like the USDA Standard Reference[2] that reports the concentration of 150 nutrients in 9,759 foods. The data was collected using gravimetry (AOAC934.06), liquid chromatography (AOAC988.15), and fluorimetry (AOAC967.22), experimental methodologies that can detect and quantify the concentration of a biochemical in food. Meanwhile untargeted metabolomics can detect hundreds more compounds within a single experiment, but only relative concentration are inferred, mainly because of the unknown fraction of molecules ionized for each compound[3,4]. Targeted metabolomics circumvents ionization efficiency[5] by using standard curves to determine the concentration, a low-throughput procedure that eliminates the advantages of untargeted metabolomics over established methodologies by increasing the time and cost over current AOAC methods. Untargeted metabolomics needs to move beyond relative concentrations for adoption by food composition catalogues. Here we report a method to predict concentrations from untargeted metabolomics, offering researchers a way to extend their results beyond relative concentrations. This is advantageous for food composition catalogues which can leverage the predictions to report expected concentrations and their potential variance for foods in a high-throughput way.

Here we show that the universal scaling of nutrient concentrations across the food supply[6] allow us to determine nutrient concentration in untargeted metabolomics. We performed metabolomics experiments on 20 raw plant ingredients to minimize the influence of human processing and span the phylogenetic tree as well as edible parts of plants (fruits, leaves, roots) to get a representative diversity of plant composition across the food supply (SI Experimental Design). Focusing on the 295 biochemicals found in at least 10 ingredients, we obtained their nutrient distribution across the food samples using the experimentally measured peak areas. In metabolomics, the peak area for each compound, n, in each food, f, measures only the ionized concentration, $c_{(f,n)}^{ionized}$. The unknown ionization efficiency, IE_n, prevents peak area from directly measuring the total concentration, $c_{(f,n)}$, following $c_{(f,n)}^{ionized}$

$= IE_{(f,n)}c_{(f,n)}$. The primary contributors to $IE_{(f,n)}$ are the instrumental parameters, the

composition of the mobile phases, and the sample matrix (ionization competition between

co-eluting compounds). Since in our experiments all samples were measured using the same

experimental setup, the differing sample matrices provide the largest contribution to $IE_{(f,n)}$

differences. It is known that the set of $c_{(f,n)}$ for all foods follows the universal features of

nutrient concentration[7]: (i) constant standard deviation ($\langle s_n \rangle$), (ii) symmetric distribution,

and (iii) translational invariance. This universality is rooted in the biochemical mechanisms

responsible for the synthesis and consumption of each compound within an organism. There-

fore, $IE_{(f,n)}$ can break the universality if it greatly varies between food. Yet, we find that

(i)-(iii) apply to the peak areas as well, with $\langle s_n \rangle = 1.41 \pm 0.50$. This suggests that

$$IE_{(f,n)} \approx \langle IE_n \rangle \tag{1}$$

for each compound, where $\langle IE_n \rangle$ is the mean ionization efficiency across all foods, conserving

the universality. We also find that the peak areas follow log-normal distributions similar to

the nutrient concentrations in [7] (SI Universal Scaling Law of Metabolomics). Lastly, we

compare the universal scaling observed for nutrients reported by the USDA (Fig 1a) with

the biochemicals obtained by our experiments (Fig 1b). We find that the linear standard

deviation of peak areas for a biochemical in MassSpec (σ_n^{MS}) relates to the mean peak area

for a biochemical in MassSpec (μ_n^{MS}) in the log-space via $\sigma_n^{MS} = e^{\alpha_\sigma^{MS}}(\mu_n^{MS})^{\beta_\sigma^{MS}}$, where α

and β are parameters of the power law fit, in excellent agreement with the USDA.

Next, we test the relationship between the metabolomics variables (σ_n^{MS}, μ_n^{MS}) and their

respective counterparts of the USDA (σ_n^{US}, μ_n^{US}). Focusing on 19 foods and 31 biochemicals

reported in both datasets (SI USDA – Metabolomics Overlap), we find low correlation ($R^2 =$

0.566), confirming that ionization efficiency obscures the proportionality between the peak

areas measured in MassSpec and the concentrations reported in the USDA via $\mu_n^{US} = \langle IE_n \rangle$

μ_n^{MS}. We can, however, estimate the concentration from peak area by leveraging (1) and the

universal features (i)-(iii), followed by both metabolomics and USDA, indicating that the

position of foods within the nutrient distributions between the datasets is largely conserved.

For example, we can see that for pyridoxine (vitamin B6) the relative position of potato

is the same in the metabolomics and the USDA distributions (Fig 1c). This suggests that

we can use the distance between the individual food measurements and the median in the

linear-space to connect the two distributions via

$$\frac{x^{US}_{(f,n)}}{e^{m^{US}_n}} \approx \frac{x^{MS}_{(f,n)}}{e^{m^{MS}_n}}, \tag{2}$$

where $x^{US}_{(f,n)}$ is the concentration of the biochemical for a food from the USDA, m^{US}_n is the

mean log concentration of the nutrient in the USDA, $x^{MS}_{(f,n)}$ is the peak area of the biochemical

in the food from experiments, and m^{MS}_n is the mean log peak area of the biochemical in the

experiments (SI Proportionality Validation).

To assess the validity of (2) we curate 113 high-quality food-biochemical pairs in overlap

between the two datasets by filtering to analytical values with at least 4 measurements (SI

Curated Pairs), then estimate $x^{US}_{(f,n)}$ and compare to the reported value in the USDA by

calculating prediction error,

$$error = \begin{cases} \frac{x^{est}_{(f,n)}}{x^{US}_{(f,n)}} & x^{US}_{(f,n)} < x^{est}_{(f,n)} \\ \frac{x^{US}_{(f,n)}}{x^{est}_{(f,n)}} & x^{US}_{(f,n)} \geq x^{est}_{(f,n)} \end{cases} \tag{3}$$

where $x^{est}_{(f,n)}$ is the estimated concentration as found by Equation (2), $x^{est}_{(f,n)} = \frac{x^{MS}_{(f,n)}}{e^{m^{MS}_n}} e^{m^{US}_n}$.

When the estimated and reported concentrations are equal, the prediction error is 1.0 and so

values closer to 1.0 is desirable. Using (3), we observe a 3.1 mean error and a 2.4 median error

for the $x^{est}_{(f,n)}$ values (Fig 2a, blue line) with 73% of the values below 4.0 error, an outcome

comparable with other untargeted metabolomics estimation methods[8], which report a 2.0-4.0

mean error.

The methodology (2) requires the concentration m^{US}_n of the biochemicals, known only

for nutrients reported in the USDA. To overcome this limitation, we rely on the finding that compound concentrations in bacterial and human cells are correlated with their chemical properties of the compounds[9], allowing us to infer m_n^{US} for nutrients not present in the USDA. We created a XGBoost model to predict $x_{(f,n)}^{US}$, taking as input the chemical properties of biochemicals (molecular weight, logP, logS, hydrogen bonding inventory, number of charged atoms, non-polar surface area) and phylogenetic lineages of foods (class, order, family, genus classifications). Leave-one-out validation of the trained model shows 70% of the $x_{(f,n)}^{US}$ values within 2.0 error and 90% within 4.0 error of the true value ($R^2 = 0.931$, Fig 2b), confirming the model is accurate (SI Gradient Boosting Methodology).

Using XGBoost to estimate m_n^{US}, (2) can estimate $x_{(f,n)}^{US}$ in individual foods using the peak areas, observing a 3.4 mean error (Fig 2a, red line) and 73% of the values below a 4.0 error. XGBoost allows us to determine the concentration of biochemicals not reported in the USDA, but detected in our experiments. For example, while S-allylcysteine in garlic is not reported by USDA, our untargeted experiments allow us to estimate its concentration as 0.158 g/100g. Using FoodMine[10], we found six published measurements for S-allylcysteine in garlic, with an average at 0.115 g/100g, giving a 1.4 error, demonstrating the possibility of using ML-models to extend the estimated concentrations from (2) beyond the USDA. While promising, the accuracy and generalizability of such models require further study (SI Beyond the USDA).

The proposed methodology offers actionable concentration estimates to complement the standard presence/absence information delivered by untargeted metabolomics, helping managing costs and resources of future studies. We find that our method estimates concentration in untargeted metabolomics with a 3.1 mean error, relying on the universal features of nutrient distributions and offers comparable performance to the current structural similarity method (4.3 mean error) without requiring chemical standards and ionization efficiency prediction method (2.1 mean error) without needing rarely measured ionization efficiencies[8]. Here, we rely on publicly available training data, facilitating the seamless integration of

our methodology in the decision-making process of health risk assessments, as seen with established methods[11] considering food-borne compounds.

References

[1] European Food Information Resource. List of Food Composition Databases at EuroFIR. https://www.eurofir.org/food-information/food-composition-databases (2009).

[2] U.S. Department of Agriculture & Agricultural Research Service. FoodData Central. https://fdc.nal.usda.gov (2019).

[3] Purves, R.W., Gabryelski, W. & Li, L. Rapid Commun Mass Spectrom 12, 695–700 (1998).

[4] Müller, C., Schäfer, P., Störtzel, M., Vogt, S. & Weinmann, W. J. Chromatogr. B 773, 47–52 (2002).

[5] Ankney, J.A., Muneer, A. & Chen, X. Annu. Rev. Anal. Chem. 11, 49–77 (2018).

[6] Menichetti, G., Barabási, A-L., & Loscalzo, J. Annu. Rev. Nutr. (2024).

[7] Menichetti, G. & Barabási, A.-L. Nat. Food 3, 375-382 (2022).

[8] Kruve, A., Kiefer, K. & Hollender, J. Anal. Bioanal. Chem. 413, 1549–1559 (2021).

[9] Bar-Even, A., Noor, E., Flamholz, A., Buescher, J.M. & Milo, R. PLoS Comput. Biol. 7, (2011).

[10] Hooton, F., Menichetti, G. & Barabási, A.-L. Sci. Rep. 10, 16191 (2020).

[11] Groff, L.C., et al. Anal. Bioanal. Chem. 414, 4919–4933 (2022).

Acknowledgements

We thank Dr. Roger Giese and Dr. Pushkar M. Kulkarni at the Department of Pharmaceutical Sciences, Northeastern University for preparing the food samples. Experiments were performed by Metabolon Inc., led by Nino Esile, Dr. Brian Ingram, and Nathan Testa. This

work was partially supported by NIH grant 1P01HL132825, American Heart Association grant 151708, ERC grant 810115-DYNASET, and Rockefeller Foundation grant 2019 FOD 026.

Data and Code Availability

The data and codes used to develop the methodology are openly available at our GitHub page at `https://github.com/Barabasi-Lab/Quantifying-Untargeted-Metabolomics`. The raw metabolomics data for the study is available on Metabolights at `https://www.ebi.ac.uk/metabolights/MTBLS3319`.

Author Contributions

A.L.B. and G.M. conceived the research. M.S. led the metabolomics experiments and performed the data analysis. A.L.B. and G.M. advised and guided the research. M.S. wrote the manuscript. A.L.B. and G.M. reviewed and edited the manuscript.

Competing interests

A.L.B. is a scientific founder of Scipher Medicine, Inc., which applies network medicine strategies to personalized drug selection, and Naring, Inc., which applies data science to food and health. All other authors declare no competing interests.

Figure 1: **Estimating Concentrations in Untargeted Metabolomics. (a)** Universal scaling law for nutrients within the USDA subset, relating the standard deviation, σ_n^{US} vertical axis, to the average nutrient concentration, μ_n^{US} horizontal axis, for 94 nutrients in 510 foods. Dashed lines are confidence intervals. Colored dots are compounds we selected to compare their relative positions between a) and b). **(b)** Universal scaling for our untargeted metabolomics experiments, relating the standard deviation, σ_n^{MS}, to the mean peak area, μ_n^{MS}, for 295 biochemicals in 20 foods. **(c)** Nutrient distributions using the 19 foods in overlap between the USDA and our experiments for pyridoxine: experiments (peak area, red) and USDA (g/100g, blue). The position of potato is shown in each distribution (black lines).

Figure 2: **Prediction Error Curves.** The proportion of estimated nutrient concentrations in individual food items below specified error values. **(a)** Error curve plots using the curated 113 biochemical-food pairs for the calculated μ_n^{US} from the USDA (blue) and predicted μ_n^{US} from a XGBoost model (yellow), taking as input the chemical properties of biochemicals (molecular weight, logP, logS, hydrogen bonding inventory, number of charged atoms, non-polar surface area) and phylogenetic lineages of foods (class, order, family, genus classifications). **(b)** Error curve of the Leave-one-out predicted concentrations for the 18,458 biochemical-food pairs in the training data. Inset: the relationship between calculated $x_{(f,n)}^{US}$ from the USDA and predicted $x_{(f,n)}^{US}$ from the XGBoost model. This shows that we can predict the concentrations reported in the USDA with a correlation of $R^2 = 0.931$ between the predicted and real concentrations.