**A Unitary Mechanism Underlies Adaptation to Both Local and Global Environmental Statistics in Time Perception**

Tianhe Wang[1,2*], Yingrui Luo[1], Richard B. Ivry[2], Jonathan S. Tsay[2], Ernst Pöppel[1,3], and Yan Bao[1,3,4*]

1 School of Psychological and Cognitive Sciences, Peking University, Beijing, China;

2 Department of Psychology and Helen Wills Neuroscience Institute, University of California Berkeley, Berkeley, USA;

3 Institute of Medical Psychology, Ludwig Maximilian University, Munich, Germany,

4 Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China

**Co-corresponding authors (*):**

Tianhe Wang (tianhewang@berkeley.edu)

Yan Bao (baoyan@pku.edu.cn)

**Abstract**

Our sense of the passage of time flexibly adapts to the statistical properties of the temporal context. Humans and non-human species exhibit a perceptual bias towards the mean of durations previously observed as well as serial dependence, a perceptual bias towards the duration of recently processed events. Here we asked whether those two phenomena arise from a unitary mechanism or reflect the operation of two distinct systems that adapt separately to the global and local statistics of the environment. We employed a set of duration reproduction tasks in which the target duration was sampled from distributions with different variances and means. The central tendency and serial dependence biases were jointly modulated by the range and the variance of the prior, and these effects were well-captured by a unitary mechanism model in which temporal expectancies are updated after each trial based on perceptual observations. Alternative models that assume separate mechanisms for global and local contextual effects failed to capture the empirical results.

**Teaser**

Time perception of humans is shaped by a common mechanism that is sensitive to short-term and long-term environmental changes.

**Introduction**

The internal representation of temporal information is essential for a wide range of cognitive functions, from anticipating future events to controlling movements (*1–3*). To improve the precision of temporal perception, the timing system flexibly adapts to the statistical properties of the current context (*4*, *5*). For example, when presented with a set of durations in a perception task, participants have a strong tendency to overestimate relatively short durations and underestimate relatively long durations. Thus, the perceived duration is biased toward the mean of the set (*4*, *6–9*), a phenomenon, known as the "central tendency effect." This effect indicates that temporal perception is sensitive to the global temporal context.

It has also been shown that temporal perception can adapt on a rapid timescale. Participants' perception of the current duration is attracted toward the duration of the previous stimulus(*10–12*). That is, the duration is perceived to be longer after a relatively long duration stimulus compared to a relatively short duration stimulus. This phenomenon, known as serial dependence, suggests that the perceptual system is also sensitive to the local statistics of the environment (*13*, *14*).

Central tendency and serial dependence effects have been observed across a wide range of perceptual tasks(*13*, *15–19*), indicating that they reflect general principles of how the perception system adapts to the statistics of the environment. Both phenomena can be explained under a Bayesian framework. On the one hand, the observers appear to construct a relatively stable global prior that reflects the distribution of the stimulus set (*6*, *20*, *21*). Following Bayesian integration, the current perception is biased toward the mean of the global prior, the central tendency effect. On the other hand, the observer also appears to build a temporal expectancy based on the most recent stimulus, inducing a bias in judging the duration of the current stimulus towards recently experienced stimuli, the serial dependence effect (*22*).

Although the central tendency and serial dependence effects describe two ways in which context can influence behavior, it remains unclear whether they reflect the operation of a unitary process or two separate, adaptive processes. From a unitary view, the perceptual system continuously updates the global prior based on new observations, and the trial-by-trial updating of the global prior could influence the subsequent perception, giving rise to serial dependence(*10*, *11*, *23*). Alternatively, there may be two adaptive systems that operate on different timescales in response to environmental statistics, building up global *and* local priors that give rise to central tendency and serial dependence, respectively.

To arbitrate between these hypotheses, we used a temporal reproduction task (*6*, *24–26*) in which

participants reproduce an interval specified by a visual stimulus. Across conditions, we manipulated the

global distribution by sampling the target durations from different temporal distributions. If serial

dependence and central tendency arise from a shared mechanism, the magnitude of the serial

dependence effect would be impacted by the global temporal distribution. Alternatively, if serial

dependence and central tendency arise from distinct mechanisms, the magnitude of the serial

dependence effect would remain invariant across a wide range of global temporal distributions. We

formalized these hypotheses and compared the predictions of these computational models with the

empirical results. By combining our behavioral experiments and model-based analyses, we sought to

unravel the computational mechanisms underlying the influence of context on temporal perception.

**Results**

*Serial dependence in time perception is attractive and non-linear*

The central tendency effect in temporal perception has been shown in many experimental contexts (*4*, *6–*

*9*). In contrast, the serial dependence effect has not been well established in timing. This motivated us to

begin this project by examining serial dependence in a widely employed temporal reproduction task, the

"ready-set-go" task (*6*). Participants observed a pair of visual events that defined a target interval ("Ready"

and "Set") and then made a single button press ("Go"), attempting to produce an interval between the

Set and Go signals that reproduced the target interval (Fig. 1a). The target durations were randomly

sampled from a uniform distribution that ranged from 500 to 900ms.

Figure 1. Trial and task structure. (a) The stimulus sequence used in **Experiment 1** ("ready-set-go" task). Two

100-ms stimuli flashed in sequence, signifying first a "Ready" signal and then a "Set" signal; the target duration

was the interval between the onset times of the "Ready" and "Set" signals. Participants were instructed to

press the space bar to reproduce the temporal interval after the "Set" signal. Performance feedback was

conveyed for 50 ms via the color of the fixation cross (green = correct; red = incorrect). (b) The stimulus

sequence used in **Experiments 2-4**. We used a duration reproduction task including both go and no-Go trials.

A ripple-shaped stimulus was presented for a fixed duration denoting the temporal interval. After a 300 ms

interval, the fixation point became either a "+" sign or "x" sign, signaling either a "go" or "no-go" trial,

respectively. On Go trials, participants reproduced the temporal interval by holding down a key. When the key

was released, the fixation point turned to a grey circle signaling the end of the trial. In the no-Go trial,

participants were asked to withhold any movement and fixate until the "x" switched to a grey circle after

700ms.

The reproduced durations exhibited robust regression towards the mean (Fig. 2a), replicating the central

tendency effect seen in previous studies (*10, 25, 27*). Quantitatively, the slope of the reproduced duration

to the target duration was significantly smaller than one (0.65 ± 0.20; t(11) = -6.08; p<0.001, Fig S1a). To

examine serial dependence, we first calculated a "deviation" index, the difference between the

reproduction of a target duration on a given trial and the individual's mean reproduction to that target

duration across all trials (see Fig. 2a). The serial dependence effect is calculated by the change in the

deviation index as a function of the difference between the previous and the current target durations.

This method minimizes potential artifacts in the serial dependence function that may be induced by

regression to the mean and reproduction biases (*28, 29*).

Figure 2. Serial dependence in duration reproduction. **Experiment 1**: (a) Reproduced durations of a

representative participant. Grey dots represent the reproduced duration of each trial. The blue circle

represents the participant's mean reproduced duration for each target duration. (b) Serial dependence is

evidence in the non-linear relationship between the deviation index, the current reproduced duration minus

the mean reproduced duration, as a function of the temporal difference between the previous target duration

and the current target duration (positive values indicate previous target was longer). Filled dots denote

individual participants. Empty circles denote the mean of all 12 participants. Red solid line represents the best-

fitted Derivative of Gaussian (DoG) model at the group level. (c) Half-height of the best-fitted DoG in

Experiment 1. Error bars represent an estimate of the standard error obtained from jackknife resampling. (d)

Instantaneous slope obtained from the fitted DoG curve for trials n-1, n-2, n-3, and n+1 trials. **Experiment 2**:

(e) Instantaneous slope obtained from the fitted DoG curve with respect to Go and no-Go trials n-1, n-2, and

n+1 (control condition). (f) Half-height of the best-fitted DoG for trials n-1 and n-2.

We found that the deviation is biased towards the previous stimulus (Fig. 2b): When the target duration

on trial n-1 was longer than the stimulus on trial n, the reproduced duration tended to be longer than

average, and vice-versa. This indicates that the reproduction on the current trial is attracted towards the

stimulus duration (or reproduced duration) of the previous trial. Notably, the shape of serial dependence

is non-linear: The attraction effect peaks when the current stimulus differs from the previous stimulus by

approximately 100ms and then falls off when the difference grows larger.

This non-linear function is well captured by a derivative of Gaussian (DoG) curve (*13*, *28*). To quantify how

previous stimuli influence the current perception in experiment 1, we fit a DoG to the function of the

difference between the target duration in trial n and trial 1-back, 2-back, or 3-back, respectively. We used

the instantaneous slope at zero to measure the sign of serial dependence (see Fig. 2b). Positive

instantaneous slopes indicate that the current perception is attracted towards the previous stimulus

duration; negative slopes indicate that the current perception is repelled from the previous stimulus

duration. We found a positive slope when calculating serial dependency based on trial n-1 (0.092 ± 0.094,

$t(11)$ = 3.39, p=0.006; Fig. 2d), but not when the calculation was based on trial n-2 (0.009 ± 0.079, $t(11)$ =

0.39, p=0.70) or n-3 (0.028 ± 0.095, $t(11)$ = 1.00, p=0.34). To better estimate the magnitude of the serial

dependence effect in response to the previous stimuli, we fit a DoG at the group level. The DoG provided

a good fit (Fig. 2b), outperforming the null-model ($\Delta AIC_n$ = -46.3 ± 7.7) and linear-model ($\Delta AIC_l$ = -18.6 ±

4.6) for the trial n-1 function. The half-height of this function is 7.6 ± 0.6ms and the half-width is 95.8 ±

10.1ms (Fig. 2c); thus, at its peak the serial dependence effect is about 8%. When analyzed at the group

level, we also get a significant serial dependence effect from trial n-2 (2.4 ± 0.8 ms, z = 3.00, p=0.001), but

not from trial n-3 (Fig. 2c), indicating there might be a weak attractive effect that is not evident in the

instantaneous slopes calculated at the individual level. We return to this issue below.

*Serial dependence mainly originates from the temporal reproduction*

In Experiment 2, we asked whether the serial dependence effect originates from temporal perception or

temporal reproduction. To test this, we included No-Go trials in which participants were instructed to not

respond. Temporal reproductions following Go trials could reflect biases arising from processes associated

only with perception, only with motor production, or both. In contrast, temporal reproductions following

No-Go trials should only be influenced by a perceptual bias. Importantly, the Go/No-Go signal was only

presented after the target duration had been presented, ensuring that participants encoded the target

duration on both Go and No-Go trials (Fig. 1b).

In trials immediately following Go trials, we replicated the serial dependence effect observed in
experiment 1 (instantaneous slope: 0.114 ± 0.042; Wilcoxon test: z = 2.03, p = 0.021; Fig. 2e) with a half-
height of 7.4 ± 0.9ms (Fig. 2f) and half-width of 111.0 ± 15.6ms (Fig S3a). In contrast, we did not find a
serial dependence effect from the previous stimulus following a No-Go trial, (instantaneous slope: 0.006
± 0.049; Wilcoxon test: z = 0.01, p = 0.97; Fig. 2e; Fig S3). This dissociation suggests that serial dependence
in timing mainly arises from sequential effects associated with temporal reproduction, not temporal
perception. Consistent with this notion, when we examined trial triplets composed of Go-NoGo-Go trials,
we found a serial dependence effect on the second Go trial towards the first Go trial (rather than the
intervening NoGo trial; instantaneous slope, 0.080 ± 0.034, Wilcoxon test: z = 2.25, p=0.012; Fig. 2e-f).
152
153 *Modeling central tendency and serial dependence*
154 Having established robust signatures of the central tendency and serial dependence effects, we now ask
155 whether these two effects are generated by a unitary mechanism or reflect separate mechanisms that are
156 shaped by global and local statistics respectively. To test this, we formalized our intuitions into a series of
157 computational models.
158
To explain the central tendency effect, we begin with a Bayesian-least-square model (*6*) that assumes the
observer's global temporal expectancy is static (i.e., global prior), with bias constrained only by the global
distribution of temporal stimuli (Global-Only model, Fig 3a). The observer's temporal estimation on a
given trial is the Bayesian integration of the global prior with the actual target duration (corresponding to
the likelihood in Bayesian theory). As such, the estimated duration on each trial is biased towards the
global prior, giving rise to the central tendency effect (Fig. 4a). However, given that the Global-Only model
uses a fixed prior, the observer's estimate will not be influenced by the local context (e.g., recently
experienced stimuli); that is, the basic Bayesian model cannot account for the serial dependence effect.
167

Figure 3. Schematics of different models. (a) In all of the models, the target duration (green) is represented with a normal distribution (likelihood) centered at the target duration with perceptual noise, and the global prior (blue) is the distribution of target stimuli. The posterior (yellow) is obtained by multiplying the global prior with likelihood. (b) In the Global-only and two global prior updating models (BU/MGU), the final estimate of the target duration is the mean of the posterior. The reproduced duration is a normal distribution centered on the target duration with motor noise. (c-d) In the BU and MGU models, after making the motor response, the observer updates the prior based on the reproduced duration. For the MGU (c), a scaled normal distribution centered at the reproduced duration was added to the old prior and the new prior was normalized. For the BU (d), the prior is updated with a Kalman filter. (e) For the Local-Only model, the likelihood is integrated with the local prior in a non-linear manner, with the weight on the local prior increasing when the likelihood and local prior are close and decreasing when they are far. (f) In the DP model, the observer is assumed to hold two priors, one global and one local. After the posterior is computed based on the global prior and likelihood, the posterior is further integrated with the local prior to generate a second posterior (posterior-2). (g) In the second step of the DP and Local-Only models, the new local prior is fully determined by the motor response of the

184    current trial. As with the other three models, the reproduced duration comes from a normal distribution
185    centered at the mean of the posterior with motor noise.
186

187
Figure 4. Model simulations. Predicted central tendency (a) and serial dependence (b) effects for the five
models. (c) Predicted relationship between central tendency and serial dependence for the models. Each line
is the prediction of one model and each dot is the prediction with a specific learning rate parameter. The half-
height increases with the learning rate. The horizontal purple line indicates the mean of the empirical data and
the vertical purple bar indicates SE. (d) Illustration of how the MGU model generates a non-linear serial
dependence. When the current and previous durations are close (left), the previous duration can influence the
shape of the prior in the range around the current duration and make the current posterior shift more relative
to the likelihood. In this region, the bias of the posterior increases with the distance between two successive
stimuli. However, when the current and previous duration are distinct (Right), the previous duration cannot
influence the shape of the prior around the likelihood and thus, the attraction effect decreases.
198
We next considered a unitary mechanism Bayesian Interference model that assumes that the observer
integrates the likelihood and a local prior based solely on the local context (Local-Only model, Fig 3g). The

weights of the local prior and likelihood are determined by the difference between distributions
associated with the current sample and recently experienced samples. Specifically, the observer will rely
less on the local prior when it is far away from the likelihood, and more on the local prior when it is close
to the likelihood. The Local-Only model can produce a non-linear serial dependence effect (Fig. 3c right).
However, this model produces a small central tendency effect (Fig. 4a), inconsistent with the behavioral
results (Fig. 4c). Thus, a local prior by itself is not sufficient to explain adaptative behavior in the current
experiments, indicating that the central tendency effect is not a by-product of serial dependence.
208
We then considered models capable of simultaneously generating serial dependence and central
tendency effects. First, we considered a unitary mechanism model in which the global prior is updated
across trials (Bayes-Updating model). Given that serial dependence is driven by temporal reproduction
(Exp 2), we assume that, following Bayes rule, the prior is updated by integrating a Gaussian centered at
the reproduced duration. Since the priors and posteriors are both Gaussians, the Bayesian integration can
be simplified into a linear weighted sum of the means (i.e., a Kalman filter, fig, 3d). This model is
mathematically similar to what has been previously described as an internal reference model (*10*, *11*).
While this model predicts a central tendency effect and an attractive serial dependence effect. the
predicted serial dependence function is near-linear (Fig. 5b), inconsistent with the empirical results of
Exps 1-2.
219
To get a non-linear serial dependence function, the prior can be updated in a non-Bayesian manner. We
assume that the brain creates the prior distribution by summing multiple Gaussian distributions with
different weights (Mixed Gaussian Updating model, MGU). When perceiving a new duration, the prior is
updated by adding a Gaussian centered at the reproduced duration to the old prior, followed by
normalization (Fig. 3c). This kind of computation has previously been suggested to account for how
Purkinje cells in the cerebellum adapt to the prior for representing temporal information (*26*). Since the
global prior is updated locally, the attraction effect from trial n-1 decreases when the duration of the
current stimulus is far from the duration of the n-1 reproduction (Fig. 5c). Thus, this model predicts a
nonlinear serial dependence function and serial dependence, in accord with the empirical results in
experiment 1-2 (Fig. 4b).
230
In contrast to the unitary mechanism models describe above, an alternative way to produce both serial
dependence and central tendency effect is to consider a hybrid model in which the two biases arise from

two distinct processes (Dual Priors model, DP). Specifically, the DP model assumes that the current

stimulus (likelihood) is integrated with a static global prior (fig. 3a), generating serial dependence. The

posterior is then integrated with a local prior (fig. 3f), inducing non-linear serial dependence. As such, the

Dual Priors model can predict both the central tendency and non-linear serial dependence effects

observed in the data. Note, those two effects are generated by two mechanisms that separately adapt to

the global and local temporal context.

In sum, our computational analyses point to two candidate models that can account for the observed

central tendency and serial dependence effects. These two models diverge in that the MGU model

postulates that central tendency and serial dependence effects arise from a unitary mechanism (non-

linear updates to a global prior), whereas the DP model postulates that these two effects arise from

separate mechanisms (a fixed global prior paired with updates to a local prior). We sought to arbitrate

between the MGU and the DP models in the following sections.

*Contextual effect on serial dependence*

A key difference between the MGU and DP models is that the latter assumes the local prior is mainly

determined by the duration on trial n-1; as such, a serial dependence effect from the n-2 trial will be very

weak. In contrast, the MGU predicts a robust serial dependence effect from trial n-2 given that all (recent)

observations are integrated into the prior. In experiment 1, the group level analysis indicated there was a

positive serial dependence effect from the n-2 trial. However, this effect was not observed in the

individual analyses of the instantaneous slope values.

To address this discrepancy, we sought to enhance the serial dependence effect and examine whether it

will be manifest beyond trial n-1. The MGU and DP models both predict that extending the range of target

durations in the test set will enhance the trial n-1 serial dependence effect (Fig 5). By the MGU, the

observer constructs a concentrated prior when the range is limited (Fig 5d right). As such, the prior is

resistant to updating, yielding a weak serial dependence effect. When the range is expanded, the prior

becomes more distributed (Fig 5d left), resulting in a more pronounced local change after each update

and thus a stronger serial dependence effect (Fig 5b-c). The DP also predicts this pattern but for a different

reason. Perceptual noise scales with duration (*9*), a form of Weber's law. Because of this, the likelihood

becomes relatively flat when the range is increased (Fig 5e). This will result in a greater influence of the

previous production and a strengthening of the serial dependence effect (Fig. Sf-g). Importantly, as noted

above, the MGU and DP models generate very different predictions regarding concerning a trial n-2 serial

dependence effect (Fig 6a): The MGU predicts that the serial dependence effect should be observed from

trial n-2, with a half-height slightly attenuated relative to trial n-1, whereas the DP predicts almost no

serial dependence effect from trial n-2.

Figure 5. The range of the target duration distribution influences the serial dependence effect. (a) Distributions of target durations in different contexts (b-c) Simulations of the MGU model predict that the half-height of the serial dependence effect will increase as the range of target duration is increased. (d) In the MGU model, changing the range of the target durations will impact the width of the prior and therefore the serial dependence effect. (e) In the PD model, since the scalar property in time perception such that the ratio between the SD and mean is a constant, the likelihood will become flatter as the target duration increases. (f-g) Simulations of the DP model also predict that the half-height of serial dependence increases substantially as the range of target duration is increased. (h) Serial dependence effects for trial n-1 in the medium (left) and long condition (right). Filled dots represent individual participants. Blank circles represent the average of all participants. The turquoise and blue lines represent the best-fitted DoG in the medium and long conditions, respectively. The red and green dash lines represent the best-fitted DoG in Experiments 1 and 2, respectively. (i) Half-height of the best-fitted DoG for serial dependence effect from trial n-1 in Experiments 1, 2, and 3 (medium and long conditions). Each filled dot represents an estimate from jackknife resampling. Error bars represent standard error.

Given the predictions of the two models, we extended the range of the target durations in experiment 3. We applied two test sets, one ranging from 520-1260ms (Medium range) and the other from 560-1860ms (Large range, Fig. 5a). We compared the results with those obtained in Experiment 1 in which the target durations ranged from 520-880ms (Short range). Extending the range of target duration successfully enhanced the serial dependence effect from trial n-1. The best-fitted DoG of the medium condition ($\Delta AIC_n$= -27.5 ± 4.2; $\Delta AIC_l$ = -10.8 ± 2.2) and long condition ($\Delta AIC_n$ = -20.7 ± 4.4; $\Delta AIC_l$ = -10.0 ± 2.1) had a higher peak and was broader than that for the short condition (Fig. 5h). Correspondingly, the half-height increased as the distribution became wider (Fig. 5i; medium: 11.8 ± 0.9 ms; long: 22.6 ± 2.0 ms; Zs > 4.9, ps < 0.001). Similarly, the half-widths also increased as the distribution became wider (Zs > 10.5, ps < 0.001), indicating an influence from more distant productions on the previous trial when the test set range increased.

The key question in Experiment 3 centers on the trial n-2 data: Will the increase in the magnitude of the serial dependence effect from trial n-1 be accompanied by a stronger serial dependence effect from non-adjacent trials? In the analysis of the individual functions, there was a significant positive instantaneous slope for the trial n-2 data in both the median and long conditions (Fig 6b). Trial n-3 also showed a

tendency for a positive bias although this did not reach significance. Moreover, fitting the DoG at the

group level showed a significant positive half-height for the serial dependence function from trial n-2 (Fig

6c). Thus, the results are consistent with the prediction of the MGU model and fail to support the DP

model (Fig 6a). Moreover, the MGU is consistent with the central tendency effect in all conditions (short,

medium, long), further supporting the idea that the process that produces the serial dependence effect

uses local information to update the prior (Fig 5g).

Figure 6. The serial dependence effect becomes stronger when the range of the test distribution increases. (a)

Simulation of the half-height with the MGU model (left) and the DP model(right). The upper two panels depict

simulations for the long-range condition and the lower two panels depict simulations for the medium range

condition. Simulations of the MGU model can produce robust serial dependence effects from trial n-2, whereas

the DP model fails to predict this effect. (b) Instantaneous slope of the DoG-fitting curve for trials n-1, n-2, and

n-3, and n+1 (control condition) in the medium and long conditions. The p-values are with respect to difference

from zero. (c) Half-height of the best-fitted DoG for the n-1 and n-2 trials in the medium (left) and long (right)

conditions.

*The variance of prior increases serial dependence*

The MGU and DP models also make differential predictions when the variance of the target distribution

is manipulated (Fig 7a). In Experiment 4, we set the mean of the target set distribution to 900 ms (the

medium condition of Experiment 3). In one condition, we created a set with low variance (720-1080ms,

short-900) compared to that used in experiment 3 where the variance was larger (540-1260ms). In the

second condition, we used a bimodal distribution to increase variance. The GMU predicts that the serial

dependence effect will be enhanced as the variance of the target distribution is increased. The logic here

is similar to that described above in terms of the range of the distribution: Because a low variance test

set, by definition, is more concentrated, the effect of trial-by-trial updating of the prior will be smaller

relative to when the variance is high. In contrast, the DP predicts that the variance of the test set will have

little effect on serial dependence (Fig 7b).

Figure 7. Serial dependence effect becomes stronger when the variance of the test distribution increases. (a)

Illustration of the distributions of target durations in experiment 4. The short-900 condition (purple) has the

same mean as the medium condition (turquoise). The bimodal (orange) has the same range as the medium

condition but has larger variability. (b-c) Predicted half-heights of the DoGs for the different conditions by the

MGU and the DP models. (d-e) Serial dependence effect for trial n-1 of the short-900 condition (d) and bimodal

condition (e). Note the scale for the x-axis is different in d and e (and thus, the function for the medium

condition looks different). Each filled dot represents a participant. The open circles indicate the average across

participants. The purple, orange, and turquoise lines represent the best-fitted DoG for the short-900, the

bimodal-medium, and the medium condition, respectively. Each filled dot represents one participant. Error

bars represent standard error. (f) Half-height of the best-fitted DoG for the trial n-1 data in Experiments 1, 2,

and 4. Each filled dot represents an estimate from jackknife resampling. Error bars represent standard error.

Consistent with the prediction of the MGU, the serial dependence effect was modulated by the variance

of the test set. For the trial n-1 data, the best-fitted DoG in the bimodal condition ($\Delta AIC_n$ = -78.6 ± 5.6;

$\Delta AIC_l$ = -18.3 ± 3.6) yielded the highest and broadest serial dependence function of the three conditions

(fig. 5e); the short-900 condition (short-900: $\Delta AIC_n$ = -48.3 ± 7.8) showed the lowest and narrowest

function (fig. 5d). Statistically, the half-heights of the functions increased as the variance of the prior

increased (short-900: 8.7 ± 0.6 ms; medium: 11.8 ± 0.9 ms; bimodal: 18.7 ± 0.8ms, Zs>4.2, ps<0.001). In

the analysis of the individual functions, the serial dependence effect was found from both trial n-1 and

trial n-2 in the bimodal condition, similar to what was found in the medium condition. In the short-900

serial dependence effect was only significant from trial n-1 (Fig S4). In sum, the results provide additional

evidence that serial dependence is enhanced in higher variance test sets.

Taken together, the results from Experiments 3 and 4 are consistent with the predictions of the MGU

model and at odds with the predictions of the DP model. Manipulations of the distributional properties

of the test set indicate that a unitary process gives rise to both the central tendency and serial dependence

effects.

**Discussion**

The internal representation of duration is context dependent. Previous work has identified two sources

of contextual bias, a central tendency bias in which the temporal representation is attracted towards the

mean of a global prior, and a serial dependence effect in which recently experienced durations serve as

attractors on the current representation of a stimulus duration (*6*, *15*, *20*, *23*, *27*). In the current study, we

asked whether the central tendency and serial dependence effects arise from a common mechanism or

separate mechanisms.

We performed a series of temporal reproduction tasks in which we manipulated the distribution of the

target duration. We observed central tendency and serial dependence effects in all of the experiments.

Although many studies have showed central tendency in duration perception(*6*, *20*), the presence of a

short-term contextual effect, serial dependency, is less clear. Moreover, the designs and analyses used in

previous work has precluded the analysis of the whole serial dependence function(*11*, *23*). In the current

study, we consistently observed a non-linear serial dependence effect in temporal reproduction. By using

a Go/No-Go task(*28*), we found that the effect was dependent on movement reproduction: There was an

attraction effect from the previous target duration after a Go trial, but no observable effect after a No-Go

trial.

To determine if long- and short-term contextual effects arise from a single or distinct Bayesian processes,

we compared five computational models. We first examined two unitary models, one with a static global

prior established during the initial phase of the experiment(*6*), and one with a local prior dictated by

recent experience. Each unitary models could not account for both the observed central tendency and

serial dependence effects. To generate both effects, we considered three more complex models. In one

of these, we assumed that there were two separate mechanisms, one based on a global prior and one

based on a local prior. For the other two models, a unitary mechanism included a process by which the

global prior was continuously updated by recent experience. We rejected the unitary model in which the

prior is updated in a Bayesian optimal manner since it generated a near-linear serial dependence function.

In contrast, simulations of the MGU unitary and DP models yield central tendency and non-linear serial

dependence effects.

To further explore the viability of the MGU and DP models, we considered predictions of the two models

when we manipulated either the range or variance of the test stimuli. While both models predict that the

amplitude of the serial dependence effect from the preceding trial will increase with the range of the test

stimuli, only the MGU model predicts that this manipulation will also enhance the serial dependence

effect from earlier trials (e.g., n-2). Moreover, the MGU model predicts that the serial dependence effect

will be enhanced when the variance of the test set is increased, even when the mean duration is fixed,

whereas the DP model predicts that the variance of the prior should have negligible influence on serial

dependence. The results showed that, when extending the range of the test set, the serial dependence

effect was enhanced and evident in terms of the context established from the n-2 as well as n-1 stimulus.

Similarly, the serial dependence effect was enhanced when increasing the variance of the test set.

Together, these results provide strong support for the MGU model, indicating that the global bias and

serial dependence effects arise from a unitary mechanism in which a single global prior is continuously

updated after each trial by the reproduced stimulus duration.

The nonlinearity of the serial dependence function sheds light on how the prior is updated based on recent

experience. Updating could follow a simple Bayesian optimal integration rule with the prior represented

as a Gaussian. While this model generates a serial dependence effect, the shape of the function is

approximately linear (BU, see Fig 4b). To capture a non-linear serial dependence function, we assumed

that the prior is updated by increasing the weight of the Gaussian centered at the recently reproduced

duration. One benefit of representing the prior with a Gaussian mixture model is that it provides a way to

represent the effects of any test set, regardless of its distributional shape. Indeed, previous studies have

also shown that a mixture of Gaussians provides a good fit of the internal prior in duration perception

experiments (*26*, *30*). Here we show that an updating rule based on this assumption can account for the

non-linear serial dependence function as well as how the function will shift when the variance of the test

set is manipulated.

As noted above, the serial dependence effect was contingent on the participant having produced a

response; it was markedly attenuated after No-Go trials. In theory, it is possible that this local effect is

reflective of some sort of motor memory, where the current motor reproduction is directly biased by the

previous motor reproduction, with the perceived duration unbiased. However, we think it more likely that

the serial dependence effect is mediated by perception: A prior of temporal expectation is constructed

from the reproduced duration, and this prior biases the perception of subsequent target stimuli which in

turn, will be reflected in the next reproduced duration. In support of this idea, several studies have shown

that making a movement of a variable duration influences performance on a subsequent duration

comparison task in which no motor reproduction was made (*12*, *31*, *32*).

We note that serial dependence effects have been observed in previous visual perception studies of

orientation and position that do not involve motor responses (*28*, *29*, *33*). Thus, we do not claim that the

absence of a serial dependence effect in the No-Go condition of Exp 2 should be taken to mean that

movement is a necessary prerequisite. As with all null results, caution is warranted. The design used in

Exp 2 may have been insensitive to capture the contribution of a purely perceptual component to the

prior. Indeed, there is evidence from duration comparison tasks without reproduction of an attractive

sequential effect (*11*, *34*). Follow-up experiments are needed to understand the factors that determine

the weight given to perception and production in forming the prior.

We recognize that, in evaluating the five models, we assumed that a global prior was established during

the initial phase of the experiment. Establishing a global prior obviously requires some integration of the

local context as the participant becomes familiar with the stimulus set. We set our initial "training" phase

based on previous studies which have shown that participants are able to generate a relatively accurate

global prior of the temporal context after about 100 trials (*17*). One open question is how the prior

updating rate changes as a function of training. In our dynamic models, we assumed that the rate

remained constant over the course of the experiment. However, it is possible that the rate of updating

weakens as the prior becomes more established. Correspondingly, the serial dependence effect might

become weaker over time. However, this temporal expectation system might be a rigid system that

recalibrates to the environment with an invariant learning rate. This characterization conforms with a

conceptualization of the operation of the cerebellum in sensorimotor adaptation, and the same principles

might apply to duration representation, another function associated with the cerebellum (*35*, *36*). The

data set in the current experiments are insufficient to examine the dynamics of updating, and we see this

as an important issue to be addressed in future studies.

In summary, the current study provides new insights into how context influences our sense of duration.

The perception of the duration of a stimulus is sensitive to both the global distribution of the stimulus set

as well as recent experience. Importantly, by examining the central tendency and serial dependence

effects in a joint manner, we observed that these two forms of bias are best explained by a unitary

mechanism in which a global prior is updated in an iterative manner based on each observation. Given

that central tendency and serial dependence effects are ubiquitous in perception tasks (*6*, *14*, *15*, *19*, *27*,

*33*), the common Bayesian framework developed here may provide a general account of how perceptual

systems adapt to environmental statistics.

**Methods**

Participants

A total of sixty-four students at Peking University were recruited for the four experiments. All participants

were right-handed with normal or corrected-to-normal vision. In Experiment 1, thirteen participants (8

females, mean age = 21.1, SD ± 1.0) were recruited for the two 1-hour sessions. One participant did not

return for the second session. In Experiment 2, twelve participants (8 females, mean age = 24.1, SD ± 4.2)

completed two 1.5-hour sessions. In Experiment 3 and 4, 52 participants (15 females, mean age = 21.1,

SD ± 2.1, 13 for each of the four conditions) completed a 1-hour experiment. Participants received $10/h

as compensation. All experimental protocols were approved by the institutional review board of the

School of Psychological and Cognitive Sciences, Peking University, and carried out according to the

approved guidelines. Written informed consent was obtained from all participants.

Testing was conducted in a dark room and the stimuli were presented on a 27-inch LCD monitor

(resolution of 1,024 × 768), viewed from a distance of 65 cm. The computer used a Windows 8 operating

system with a refresh rate of 100 Hz. The experiment was written in MATLAB (Mathworks, Natick, MA;

Psychophysics Toolbox Brainard, 1997;(*37*)).

Procedure and design

Experiment 1

We used a "ready-set-go" time-reproduction task(*6*) (Fig. 1a) to measure global (central tendency effect)

and local (serial dependence effect) biases. Each trial started with the presentation of a gray fixation point

(0.5 degrees diameter) at the center of the screen. After a random interval ranging from 0.7-1.2 s (drawn

from an exponential distribution), two 100-ms stimuli flashed in sequence, with the first serving as the

"Ready" signal and the second serving as the "Set" signal. The visual stimulus was either a grey ripple-

shaped arc (Exp 1) or circle (Exps 2-4) (see Fig 1) with a radius of around 12 degrees.

The interval between the onset times of the "Ready" and "Set" signals defined the target interval.

Participants were instructed to press the space bar ("Go") to reproduce the target interval, with the onset

of the reproduction interval defined by the "Set" stimulus. After the keypress, performance feedback was

provided for 50 ms via a change in color of the fixation point: Green indicated that the reproduced

duration was within an acceptable window and red indicated that the reproduced duration fell outside

this window. The criterion window was continuously adapted based on the participant's performance

such that green and red appeared with roughly equal probability. The feedback was provided to

encourage the participant to pay attention to the task. It was relatively uninformative (e.g., did not provide

signed information) because we did not want participants to correct their timing based on the feedback.

Each participant completed two sessions, with each session composed of 10 blocks of 200 trials (2000

trials total). A one-minute break was provided every two blocks. The stimulus set consisted of ten target

durations, ranging from 520 to 880 ms (step size of 40 ms). Target duration was randomized within a block

of 100 trials with the constraint that each duration was presented 10 times.

Experiment 2

The goal of Experiment 2 was to evaluate whether the source of the serial dependence effect is

perceptual, motoric, or a combination of both. To test this, we included Go and No-Go trials in a duration

reproduction task.

After a random interval ranging from 0.7-1.2s (following an exponential distribution), a ripple-shaped

stimulus was presented for the target duration. The spatial distribution of brightness was constant, but

the actual shape varied across trials to avoid repetition suppression effects (*38–41*). Crucially, 300 ms after

the offset of the target stimulus, the fixation point changed to either a "+" or "x", with these symbols

indicating that the current trial was a Go or No-Go trial, respectively. On Go trials, participants were

instructed to depress the space bar for a duration that matched the target duration. The fixation point

changed back to a grey circle right after the release of the keypress. On No-Go trials, participants were

asked to fixate without movement until the "x" disappeared. On these trials, the grey circle reappeared

after 700ms, indicating the start of a new trial. Note that no time constraints were imposed on Go trials;

thus, we anticipated there would be few errors of omission (Go trials) or commission (No-Go trials).

However, we assumed that the target duration would be similarly encoded on all trials since the "+" or

"x" did not appear until after the target stimulus.

There were five target durations (540, 620, 700, 780, 860 ms) with each target duration repeated on 360

trials. For each target duration, 60% were Go trials and 40% were No-Go trials. Target duration and

response requirement were randomized within blocks of 180 trials. Each session consisted of five blocks

(1800 total trials across the two sessions), with a one-minute break between every two blocks.

Experiment 3

The goal of Experiment 3 was to assess how the serial dependence effect is impacted by the range of the

stimulus set. To test this, we employed two new stimulus sets: A medium condition (540, 720, 900, 1080,

1260ms) and a long condition (560, 880, 1200, 1520, 1840ms). We compared performance with these sets

to the data from Experiments 1-2 (where the range was shorter, 540-860ms). The procedure in

Experiment 3 was identical to that of Experiment 2, except that only Go trials were included. There were

five blocks of 150/120 trials (medium/long condition), resulting in a total of 750 trials for the medium

condition and 600 trials for the long condition. The difference was imposed to keep all sessions within one

hour. The target duration was selected at random on each trial with the constraint that each condition

occurred an equal number of times within each block.

Experiment 4

The goal of Experiment 4 was to examine how the serial dependence effect is impacted by the variability

of the stimulus set. To test this, we employed two new stimulus sets. For the short-900 condition, the test

set ranged from 720-1080 ms (steps of 40 ms, mean = 900 ms). This group has the same mean as the

medium condition in Experiment 3 but with a shorter range (equal to that used in Exps 1 and 2). In this

way, the variance of the test set is smaller than that of the medium condition. For the bimodal-medium

condition, the test values were the same as that used in the medium condition of Experiment 3, but the

extreme values (540 ms and 1160 ms) were presented three times as often as the other three test

durations (670, 900, 1030ms). Short-900 condition included 10 blocks of 200 trials and the bimodal

condition includes 10 blocks of 135 trials.

Data analysis.

The logic of these experiments is predicated on the assumption that participants are familiar with the

temporal context. Given this, the first block of each experiment (approx. 7-10 minutes of data collection)

was treated as the familiarization phase and not included in the analysis. In addition, reproduced

durations shorter than 0.3 s or longer than 1.5s (2.5s for Experiments 3-4) were considered outliers and

excluded from the analyses (less than 0.1 % of trials).

The central tendency bias, or regression to the mean was quantified as the regression coefficient between

the reproduced durations and the target durations. To analyze the serial dependence effect, we used a

"deviation" index. For each individual, the average reproduced duration was calculated for each target

duration. The deviation was defined as the reproduced duration for a given trial minus the mean

reproduced duration of all trials with that target duration (*28*, *29*). Positive values indicate that the

reproduced duration for the present trial was longer than the average reproduction for that target and

negative values indicate that the reproduced duration was less than the average reproduction.

To quantify the magnitude of the serial dependence effect, a simplified DoG curve was fit to describe the

deviation index as a function of the difference between the current target duration and reference

duration, where the reference could be the target duration of the previous trial (n-1), two trials back (n-

2), etc., as well as following trial (n+1, serving as a baseline):

568
$$y = abcxe^{-(bx)^2},$$

where y is the deviation, x is the relative target duration of the previous trial, $a$ is half the peak-to-trough

amplitude of the derivative-of-Gaussian, $b$ scales the width of the Gaussian derivative, and $c$ is a constant,

$\sqrt{2}/e^{-0.5}$, which scales the curve to make the $a$ parameter equal to the peak amplitude. As a measure of

serial dependence, we report half the peak-to-trough amplitude (half-height) and half the width of the

best-fitted derivative of a Gaussian. A positive value for the $a$ parameter indicates a perceptual bias

toward the target durations of the previous trials. A negative value for the $a$ parameter indicates a

perceptual bias away from the target durations of the previous trials.

We fit the Gaussian derivative at the group and individual level using constrained nonlinear minimization

of the residual sum of squares. Jackknife resampling was applied to estimate the variation of the

parameters for the group-level fit, where each participant was systematically left out from the pooled

sample. The standard deviation of those estimates represented the standard error of the parameter at

the group level. The half-height and half-width of the best-fitted DoG were compared between groups

with a t-test, where the t-value was computed with the mean and the variance of the parameters

estimated from the jackknife resampling procedure. Bonferroni correction was applied for multiple

comparisons. To test the goodness-of-fit of our model, we computed the ΔAIC for the DoG model

compared with either a non-model (y = 0, ΔAIC$_n$) or a linear model (y = $kx$, ΔAIC$_l$). A negative ΔAIC indicates

DoG performed better than the alternative models.

To determine the extent of serial dependency, we fitted individual serial dependence functions in which

the reference could be the target duration of the previous trial (n-1), two trials back (n-2), etc. We also

tested the serial dependence function for the n+1 trial. Given that this the reference stimulus has not

been experienced, there should be no serial dependence effect here, providing a test of whether the

deviation measure is a valid index to analyze serial dependence. Since individual serial dependence

functions do not always show a strong nonlinearity, $a$ parameter (half-height) can become unreasonably

large. Thus, we opted to use the instantaneous slope of the DoG at the inflection point ($abc$) when

estimating the presence of a serial dependence effect at the individual level. Note that this index measures

the sign of the serial dependence independent of whether or not the function is linear.

In Experiment 2, the Go trials were sorted into two groups based on whether the reproduced interval on

that trial was preceded by a Go trial or a No-Go trial. We calculated the average reproduced durations

and the deviation using the same protocol as Experiment 1 and then fit DoG functions separately for the

Go/Go trial sequence and the No-Go/Go trial sequence to measure the serial dependence effect from trial

n-1. For each function, we calculated the instantaneous slope of the DoG at the inflection point. Note that

if the serial dependence effect is dependent on a motor response, there should be no serial dependence

effect when the preceding trial was a No-Go trial. For Go trials preceded by a No-Go trial, we also analyzed

the serial dependence effect from n-2 Go trials. Group-level DoG fitting was performed on n-1 and n-2 Go

trials separately to quantify the amplitude of serial dependence.

One-sample t-tests and paired t-tests were applied at the group level for comparisons. Normality and

equal variance assumptions were assessed prior to the t-tests. The Wilcoxon Sign-rank test was applied

when the normality assumption was violated. Two-tailed P values are reported for all statistic tests, and

the significance level was set as $p < 0.05$. All analyses were performed with MATLAB 2018b (The

MathWorks, Natick, MA).

*Models*

To account for central tendency and serial dependence effects, we implemented five Bayesian models:

Two single process models (Global-only, Local-Only), a model with both a static global prior and and local

prior (Dual-prior), and two unitary models that capture how a global prior is dynamically updated by

recent experience (Bayes-Updating & Mix-Gaussian-Updating).

Global-Only model

This model is based on a Bayesian observer model that uses a Bayes-Least-square as the mapping rule (*6*).

We assume that the observer builds up an internal prior, $\pi(t_s)$ based on the target durations ($t_s$) observed

during an initial exposure phase, and subsequent judgments are made by reference to this static prior.

The likelihood function describes the probability of the perceived duration ($t_p$) given $t_s$.

$$p(t_p|t_s) = N(t_p|t_s, v_p) \qquad\qquad [1]$$

626 where $N(x|m, s)$ represents a normal distribution with mean m and standard deviation s, and $v_p$ scales

627 the perception noise. The posterior, $\pi(t_s|t_p)$, is the product of the prior multiplied by the likelihood

628 function and appropriately normalized.

629
$$\pi(t_s|t_p) = \frac{p(t_p|t_s)\,\pi(t_s)}{\int p(t_p|t_s)\,\pi(t_s)dt_s}$$
[2]

630 The loss function, $l(t_e, t_s)$, was used to convert the posterior into a single estimate, $t_e$, the mean of the

631 posterior in the present situation.

632
$$l(t_e, t_s) = (t_e - t_s)^2$$
[3]

633
$$t_e = \underset{t_e}{\mathrm{argmin}}\left[\int l(t_e, t_s) * \pi(t_s|t_p)\,dt_s\right]$$
[4]

634 The Bayesian observer makes a response based on $t_e$ :

635
$$p(t_r|t_e) = N(t_r \mid t_e, v_m)$$
[5]

where $t_r$ is the reproduced duration, and $v_m$ scales the motor noise. Previous studies (*6*, *9*) have shown

that scalar forms of perceptual and motor noise provide a better fit of the behavior compared to when $v_p$

and $v_m$ are treated as constants. Thus, we set $v_p$ as $n_p * t_s$, and $v_m = n_m * t_e$, where $n_p$ and $n_m$ are

constants.

For this baseline, Global-only model, we simulated the results under the assumption that the prior was

established based on a data set that would be experienced during the first 7-10 minutes of the

experiments, and then remained fixed for the duration of the experiment. We assumed that participants

learn the true distribution of target durations as the prior:

645
$$\pi(t_s) = U(t_s, [500\ ms, 900\ ms])$$
[6]

where $U(x, [y, z])$ represents a uniform distribution ranging from $y$ to $z$.

Local-Only Model

To capture the serial dependence effect, we adapted a Bayesian integration model from a previous study

that examined serial dependence in magnitude estimation (*22*). In the current context, this model

assumes that the observer integrates the previous response ($t_r$) with the current stimulus ($t_s$) when

estimating the duration of the current stimulus:

653
$$t_{e,i} = (1 - W_{i-1})t_{s,i} + W_{i-1}t_{r,i-1}$$
[7]

654 where $t_{e,i}$ and $t_{s,i}$ indicate the $t_e$ and $t_s$ of trial i, and $t_{r,i-1}$ is the reproduced duration of trial i-1. $W_{i-1}$ is

655 the weight the observer assigns to the trial n-1 response when estimating the current duration. Following

656 Bayes rule to integrate the two samples dictating the current percept, the observer specifies $W_{i-1}$ as

$$W_{i-1} = \frac{1/\sigma_{i-1}^2}{1/\sigma_i^2 + 1/\sigma_{i-1}^2} = \frac{\sigma_i^2}{\sigma_i^2 + \sigma_{i-1}^2} \qquad [8]$$

658 where $\sigma$ is the variance of the estimate. The weight is influenced by the distance between the two stimuli,

$$W_{i-1} = \frac{\sigma_i^2}{\sigma_i^2 + \sigma_{i-1}^2 + (t_{s,i} - t_{s,i-1})^2} \qquad [9]$$

660 The variance is assumed to follow a power law.

$$\sigma = Kt_s{}^\alpha \qquad [10]$$

Assuming that time perception follows a scalar rule, $\alpha = 1$. $K$ is a free parameter regulating the behavior

of the model.

Dual Prior model

We combined the Local-Only and Global-Only models to create a Dual Prior model. It contained two

Bayesian processes. For the first integration, the observer integrates the current stimulus ($t_s$) with the

prior $\pi(t_s)$ to get an estimation $t_e'$, following Global-Only model (Eq [4]-[7]). The second integration

estimates $t_{e,i}$ based on $t_{r,i-1}$, $t_{e,i}'$ and $t_{e,i-1}'$ following the Local-Only model. The formulas [11] and [13]

are rewritten as

$$t_{e,i} = (1 - W_{i-1})t_{e,i}' + W_{i-1}t_{r,i-1} \qquad [11]$$

$$W_{i-1} = \frac{\sigma_i^2}{\sigma_i^2 + \sigma_{i-1}^2 + (t_{e,i}' - t_{e,i-1}')^2} \qquad [12]$$

675 Bayes Updating model

676 We also considered two unitary process models in which a global prior is updated in a dynamic manner.

677 For the Bayes Updating model, the prior is updated following Bayes rule after each observation. Since the

678 prior and likelihood are Gaussian, this model can be expressed as a Kalman filter. On each trial, the

679 estimated duration ($t_e$) is a weighted sum of $t_o$ and the duration of the stimulus ($t_s$).

$$t_e = (1 - w) * t_o + w * t_s \qquad [13]$$

where the weight $w$ is determined by the perceptual noise and variance of the prior. As such, as $w$

increases, the weight given to the prior will increase (i.e., attraction to central tendency). Participants

make a motor response based on $t_e$ with Gaussian motor noise (see equation [5]). Given that Experiment

2 showed that serial dependence is primarily induced by the reproduction component rather than the

perceptual component, we assumed the prior is updated according to $t_r$. After the motor response, $t_0$ is

updated based on $t_r$ following another Kalman filter:

$$t'_o = (1 - k) * t_o + k * t_r \tag{14}$$

where $t'_o$ is the new reference point for the next trial, and $k$ represents the learning rate. To address the

scalar property of timing noise (Weber law), $t_r$, $t_s$, $t_e$, and $t_o$ are taken to be the log value of the respective

durations. Note, that an alternative version of this model could have the observer directly use the

posterior as a new prior (what is known as a fully irritative model). However, as with the Local-Only model,

a fully irritative model will largely overestimate serial dependence given a reasonable central tendency

effect.

Mixed-Gaussian Updating model

In a second unitary model, the prior is represented as a Gaussian mixture model. This model is identical

to the Global-Only model (equation [1-5]) except that the prior is updated after each trial to generate a

better estimate of the temporal context. We applied a simple updating rule here, in which the observer

adds a normal distribution centered at $t_r$ with a standard deviation of $v_p$ to the old test set. The posterior

is then appropriately normalized:

$$\pi'(t_s) = \frac{\pi(t_s) + r * N(t_s | t_r, v_p)}{\int [\pi(t_s) + r * N(t_s | t_r, v_p)] dt_s} \tag{15}$$

where $\pi'(t_s)$ is the new test set, and $r$ is the learning rate.

Simulation procedure.

Simulations of each model were conducted to evaluate the results of Experiment 1. The data from 100

pseudo-participants were generated for each simulation. To determine the values of the free parameters

for the simulations, we referred to a previous study that used a similar design to that employed here and

evaluated the results with the BLS model, or what we refer to as the Global-Only model (6). Based on their

results, we set $n_p = 0.10$ and $n_m = 0.06$ for the Global-Only, MGU, and DP models. Other parameters were

determined to make the amplitude of the serial dependence function roughly similar to the behavioral

results. Specifically, we set $w = 0.7$ and $k = 0.3$ for the BU model, the learning rate $r = 0.3$ for the MGU

model, and $K = 0.06$ for the Local-Only and DP models. We designed Experiments 3 and 4 to focus on the

MGU and DP models, the two models that produce both central tendency and non-linear serial

dependence effects. For simulations of the short conditions, the step between each two adjacent stimuli

was the same as what was used in the other experiments (40 ms). For simulations of the medium and the

long conditions, the step size was set to a smaller value (70 ms) than used in the experiments to improve

resolution. The parameters $n_p$ and $n_m$ were fixed at the values used in the previous simulations, while a

series of *r* and *K values were tested*.

Data and code availability

Data for this paper and codes for analyses are available at https://osf.io/2dz7k/

722
**Author Contributions**

T.W. designed research; T.W. and Y.L. performed research; T.W. analyzed data; T.W. prepared figures;

T.W., Y.L., E.P., Y.B., J.S.T, and R.B.I wrote the paper.

**Competing Interest**

RI is a co-founder with equity in Magnetic Tides, Inc.

**Acknowledgments**

This work was supported by the National Natural Science Foundation of China (31771213 and 31371018)

to Yan Bao. We thank Linfeng Han, Jiashu Wang, and Huihui Zhang for valuable discussions.

**Reference**

1. R. B. Ivry, J. E. Schlerf, Dedicated and intrinsic models of time perception. *Trends Cogn. Sci.* **12**, 273–
280 (7/2008).

2. E. Pöppel, A hierarchical model of temporal perception. *Trends Cogn. Sci.* **1**, 56–61 (5/1997).

3. Y. Bao, A. Szymaszek, X. Wang, A. Oron, E. Pöppel, E. Szelag, Temporal order perception of auditory
stimuli is selectively modified by tonal and non-tonal language environments. *Cognition*. **129**, 579–
585 (2013).

4. H. Lejeune, J. H. Wearden, Vierordt's *The Experimental Study of the Time Sense* (1868) and its legacy.
*Eur. J. Cogn. Psychol.* **21**, 941–960 (09/2009).

5. Vierordt: Der zeitsinn nach versuchen - Google Scholar, (available at
https://scholar.google.com/scholar_lookup?hl=en&publication_year=1868&author=K.+Vierordt&ti
tle=+Der+Zeitsinn+nach+Versuchen+).

6. M. Jazayeri, M. N. Shadlen, Temporal context calibrates interval timing. *Nat. Neurosci.* **13**, 1020–
1026 (8/2010).

7. G. C. Ganzenmüller, S. Hiermaier, M. May, Improvements to the prototype micro-brittle linear
elasticity model of Peridynamics. *arXiv [physics.comp-ph]* (2013), (available at
http://arxiv.org/abs/1312.5543).

8. J. H. Wearden, L. A. Jones, Is the Growth of Subjective Time in Humans a Linear or Nonlinear Function
of Real Time? *Q. J. Exp. Psychol.* **60**, 1289–1302 (09/2007).

9. J. H. Wearden, H. Lejeune, Scalar Properties in Human Timing: Conformity and Violations. *Q. J. Exp.*
*Psychol.* **61**, 569–587 (04/2008).

10. K. M. Bausenhart, O. Dyjas, R. Ulrich, Temporal reproductions are influenced by an internal
reference: Explaining the Vierordt effect. *Acta Psychol.* . **147**, 60–67 (03/2014).

11. O. Dyjas, K. M. Bausenhart, R. Ulrich, Trial-by-trial updating of an internal reference in discrimination
tasks: Evidence from effects of stimulus order and trial sequence. *Atten. Percept. Psychophys.* **74**,
1819–1841 (11/2012).

12. K. M. Bausenhart, D. Bratzke, R. Ulrich, Formation and representation of temporal reference
information. *Current Opinion in Behavioral Sciences*. **8**, 46–52 (04/2016).

13. J. Fischer, D. Whitney, Serial dependence in visual perception. *Nat. Neurosci.* **17**, 738–743 (5/2014).

14. M. Manassi, D. Whitney, Illusion of visual stability through active perceptual serial dependence. *Sci.*
*Adv.* **8**, eabk2480 (2022).

15. K. P. Körding, D. M. Wolpert, Bayesian integration in sensorimotor learning. *Nature*. **427**, 244–247
(2004).

16. H. Sohn, M. Jazayeri, Validating model-based Bayesian integration using prior–cost metamers. *of the*
*National Academy of Sciences* (2021) (available at
https://www.pnas.org/content/118/25/e2021531118.short).

17. M. Berniker, M. Voss, K. Kording, Learning Priors for Bayesian Computations in the Nervous System.
*PLoS One*. **5**, e12686 (2010).

18. A. Liberman, J. Fischer, D. Whitney, Serial Dependence in the Perception of Faces. *Curr. Biol.* **24**,
2569–2574 (11/2014).

19. A. Kiyonaga, J. M. Scimeca, D. P. Bliss, D. Whitney, Serial Dependence across Perception, Attention,
and Memory. *Trends Cogn. Sci.* **21**, 493–497 (07/2017).

20. Z. Shi, D. Burr, Predictive coding of multisensory timing. *Current Opinion in Behavioral Sciences*. **8**,
200–206 (04/2016).

21. N. W. Roach, P. V. McGraw, D. J. Whitaker, J. Heron, Generalization of prior information for rapid
Bayesian time estimation. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 412–417 (2017).

22. G. M. Cicchini, G. Anobile, D. C. Burr, Compressive mapping of number to space reflects dynamic
encoding mechanisms, not static logarithmic transform. *Proceedings of the National Academy of*
*Sciences*. **111**, 7867–7872 (2014).

23. S. Glasauer, Z. Shi, Individual beliefs about temporal continuity explain variation of perceptual biases.
*Sci. Rep.* **12**, 10746 (2022).

24. G. T. Finnerty, M. N. Shadlen, M. Jazayeri, A. C. Nobre, D. V. Buonomano, Time in Cortical Circuits.
*Journal of Neuroscience*. **35**, 13912–13916 (2015).

25. M. Jazayeri, M. N. Shadlen, A Neural Mechanism for Sensing and Reproducing a Time Interval. *Curr.*
*Biol.* **25**, 2599–2609 (2015).

26. D. Narain, E. D. Remington, C. I. D. Zeeuw, M. Jazayeri, A cerebellar mechanism for learning prior
distributions of time intervals. *Nat. Commun.* **9**, 469 (12/2018).

27. Y. Murai, Y. Yotsumoto, Timescale- and Sensory Modality-Dependency of the Central Tendency of
Time Perception. *PLoS One*. **11**, e0158921 (2016).

28. M. Manassi, A. Liberman, A. Kosovicheva, K. Zhang, D. Whitney, Serial dependence in position occurs
at the time of perception. *Psychon. Bull. Rev.* **25**, 2245–2253 (12/2018).

29. D. Pascucci, G. Mancuso, E. Santandrea, C. Della Libera, G. Plomp, L. Chelazzi, Laws of concatenated
perception: Vision goes for novelty, decisions for perseverance. *PLoS Biol.* **17**, e3000144 (2019).

30. L. Acerbi, D. M. Wolpert, S. Vijayakumar, Internal Representations of Temporal Statistics and
Feedback Calibrate Motor-Sensory Interval Timing. *PLoS Comput. Biol.* **8**, e1002771 (2012).

31. J. J. Wehrman, J. H. Wearden, P. Sowman, Short-term effects on temporal judgement: Sequential
drivers of interval bisection and reproduction. *Acta Psychol. .* **185**, 87–95 (04/2018).

32. D. Yon, R. Edey, R. B. Ivry, C. Press, Time on your hands: Perceived duration of sensory events is
biased toward concurrent actions. *J. Exp. Psychol. Gen.* **146**, 182–193 (02/2017).

33. M. Fritsche, E. Spaak, F. P. de Lange, A Bayesian and efficient observer model explains concurrent
attractive and repulsive history biases in visual perception. *Elife*. **9** (2020), doi:10.7554/eLife.55389.

34. F. Sierra, D. Poeppel, A. Tavano, "A bias generating temporal distortions in serial perception"
(Neuroscience, 2021), (available at http://biorxiv.org/lookup/doi/10.1101/2021.10.08.463190).

35. A. Breska, R. B. Ivry, "The human cerebellum is essential for modulating perceptual sensitivity based
on temporal expectations" (Neuroscience, 2021), (available at
http://biorxiv.org/lookup/doi/10.1101/2021.01.15.426903).

36. R. B. Ivry, R. M. C. Spencer, The neural representation of time. *Curr. Opin. Neurobiol.* **14**, 225–232
(04/2004).

37. M. Kleiner, D. Brainard, D. Pelli, A. Ingling, R. Murray, C. Broussard, What's new in psychtoolbox-3.
*Perception*. **36**, 1–16 (2007).

38. W. J. Matthews, Stimulus Repetition and the Perception of Time: The Effects of Prior Exposure on
Temporal Discrimination, Judgment, and Production. *PLoS One*. **6**, e19815 (2011).

39. W. J. Matthews, A. I. Gheorghiu, Repetition, expectation, and the perception of time. *Current
Opinion in Behavioral Sciences*. **8**, 110–116 (04/2016).

40. J. J. New, B. J. Scholl, Subjective time dilation: Spatially local, object-based, or a global visual
experience? *J. Vis.* **9**, 4–4 (2009).

41. V. Pariyadath, D. Eagleman, The Effect of Predictability on Subjective Duration. *PLoS One*. **2**, e1264
(2007).

**Supplementary information**

Figure S1. Central tendency of the reproduced duration as predicted by the MGU model in all experiments. (a-e) Reproduced duration is plotted as a function of target duration for Experiments 1, 3, & 4. The shaded area indicates S.E. The median slope ± S.E. is reported on each figure. (f) The predicated slopes for the central tendency of the MGU model provide a good fit to the data. The dots and error bars indicate median slope ± S.E.

Figure S2. Histogram of the difference between the previous target duration ($S_{n-1}$) and the current target

duration ($S_n$). For individual participants, there are few trials with a large difference between stimuli,

especially for experiment 1 and the short-900 condition. As such, the DoG may not provide a good fit

when used to estimate serial dependence curves at the individual level.

Figure S3. Deviation index functions for Experiment 2. (a) A prominent DoG curve can be seen for the n-1

data when trial n-1 was a Go trial. (b) This curve is markedly attenuated when trial n-1 was a no-Go trial.

(c) A small DoG curve is evident for the n-2 trial when n-1 was a No-Go trail and n-2 is a Go trial. The thick

dashed line is the best-fitted DoG curve. Shaded areas indicate standard error.

Figure S4. Instantaneous slope of the fitted DoG functions for trials n-1, n-2, and n-3, and future (n+1)

trials in the short-900 (a) and bimodal conditions (b). A significant serial dependence effect can be

observed from the n-2 trial in the bimodal condition rather than the short-900 condition in experiment 4.

Dots indicate individual data points and error bars represent standard error. The p-values are based on a

test of whether the observed values differ from zero.