A decrease in fatty acid synthesis rescues cells with limited peptidoglycan

synthesis capacity

Short title: Balancing membrane and cell wall synthesis

Jessica R. Willdigg[a] and John D. Helmann[a*]

[a]Department of Microbiology, Cornell University, Ithaca, NY, USA

[*]Address correspondence to John D. Helmann, jdh9@cornell.edu

ORCID: 0000-0002-3832-3249

Competing Interest Statement: The authors declare no conflict of interest

Keywords: cell wall, peptidoglycan, fatty acid synthesis, elongasome, antibiotic

Abstract

Proper synthesis and maintenance of a multilayered cell envelope is critical for bacterial fitness. However, whether genetic mechanisms exist to coordinate synthesis of the membrane and peptidoglycan envelope layers is unclear. In *Bacillus subtilis*, synthesis of peptidoglycan (PG) during cell elongation is mediated by an elongasome complex acting in concert with class A PBPs (aPBPs). We previously described mutant strains limited in their capacity for PG synthesis due to a loss of aPBPs and an inability to upregulate elongasome function. Growth of these PG-limited cells can be restored by suppressor mutations predicted to decrease membrane synthesis. One suppressor mutation leads to an altered function repressor, FapR*, that functions as a super-repressor and leads to decreased transcription of fatty acid synthesis genes. Consistent with fatty acid limitation mitigating cell wall homeostasis defects, inhibition of FAS by cerulenin also restored growth of PG-limited cells. Moreover, cerulenin rescued the growth of wild-type (WT) cells in the presence of otherwise inhibitory concentrations of β-lactams. These results imply that limiting PG synthesis results in impaired growth, in part, due to an imbalance of PG and cell membrane synthesis and that *B. subtilis* lacks a robust genetic mechanism to coordinate the balanced synthesis of these two envelope layers.

Importance

Understanding how a bacterium coordinates cell envelope synthesis is essential to fully
appreciate how bacteria grow, divide, and resist cell envelope stresses such as β-lactam
antibiotics. The balance between the peptidoglycan cell wall and the essential plasma membrane
structure must be maintained for the cell to maintain shape, turgor pressure and resist external
cell envelope threats. Using *Bacillus subtilis*, we show that cells deficient in peptidoglycan
synthesis can be rescued by compensatory mutations which decrease the synthesis of fatty acids.
Further, we show that inhibiting fatty acids synthesis with cerulenin is sufficient to reduce
susceptibility to β-lactams. Understanding the coordination of cell wall and plasma membrane
synthesis may provide insights relevant to antimicrobial treatment.

Introduction

Understanding how a bacterium coordinates the synthesis of its multiple envelope components is essential to fully appreciate how bacteria grow and divide. The balanced synthesis of the peptidoglycan (PG) cell wall, the cell membrane, and any additional envelope layers must be maintained for the cell to maintain shape and resist turgor pressure and external cell envelope threats. Many of our most powerful antibiotics target PG synthesis, and others disrupt membrane synthesis or integrity. Understanding the consequences of antibiotic inhibition, and the processes that ultimately lead to growth cessation or cell death, will facilitate efforts to counter the emerging threat of antibiotic resistance.

The PG cell wall is an important structure for nearly all bacteria, allowing cells to resist internal turgor pressure to prevent lysis as well as conferring shape and rigidity to the cell. Seemingly contrary to its role in maintaining rigidity, the cell wall is a highly dynamic structure (1, 2). The PG sacculus is constantly remodeled to accommodate the insertion of newly synthesized glycan strands during cell growth and division (3). Lipid II, a lipid-linked disaccharide pentapeptide PG precursor, is flipped across the cell membrane, and then the glycopeptide subunits are incorporated into the sacculus by transglycosylases (TG) and penicillin-binding proteins (PBPs) (4). Elongation of rod-shaped bacterial cells results from the balanced action of bifunctional class A PBPs (aPBPs), which contain both TG and transpeptidase (TP) activities, and an MreB-associated elongasome complex. The elongasome complex, which moves circumferentially around the short axis of the cell, includes RodA (TG), class B PBPs (TP), and associated endopeptidases. The aPBPs move independently and diffusely around the cell during PG assembly (5-8). Although these PG synthesis machines do not move in a

coordinated manner, both aPBPs and the elongasome are necessary to maintain normal cell

length and width (9, 10).

Despite recent advances, there is still much to learn about PG synthesis and assembly. *B.*

subtilis has served as the premier model system for the study of envelope synthesis in Gram-

positive bacteria. We have previously characterized several cell envelope stress response (CESR)

pathways that facilitate adaptation to antibiotics that inhibit envelope synthesis or function (11).

For example, the extracytoplasmic function (ECF) sigma factor σ^M is necessary for intrinsic

resistance to PG synthesis inhibitors such as β-lactam antibiotics and moenomycin, which

inhibits the TG activity of class A PBPs (12, 13). In circumstances where the aPBPs are absent

due to mutation or have been inhibited by antibiotics, cells depend on the elongasome for PG

synthesis. Several components of the elongasome, including RodA, are transcriptionally

activated by σ^M in response to cell envelope stress (14).

An initially puzzling study revealed that cells lacking all four aPBPs (Δ4 aPBP) still

synthesize PG (15), which implied the presence of an alternative TG enzyme for assembly of the

glycan strands. Subsequent work revealed that RodA provides TG activity to the elongasome,

and functions together with class B PBPs that provide TP activity (14, 16, 17). Previously, we

used TnSeq to identify genes that become essential in cells deficient for aPBP activity (18). This

led to the elucidation of a regulatory pathway that leads to the σ^I-dependent upregulation of the

LytE endopeptidase and MreBH scaffolding protein (18). This pathway is initiated by the EcsA

membrane protein, which is required for the function of the RasP intramembrane protease (19).

RasP, in turn, cleaves multiple proteins including the anti-sigma factors for the σ^V (20, 21), σ^W

(19, 22), and σ^I stress responses (23, 24). Thus, in cells lacking all four aPBPs (Δ4 aPBP), or

even just the dominant enzyme PBP1 (*ponA*), the loss of *ecsA*, *rasP*, or *sigI* prevents growth on

LB medium (18). These cells fail to grow since they are unable to upregulate elongasome

activity to compensate for the loss of aPBPs. Growth on LB is observed if the medium is

supplemented with high levels of magnesium (Mg) ion (18), as seen also for other mutant strains

with defects in PG synthesis (25). The mechanism of this suppression is unknown but may be

related to a reduction of autolysin activity (25). We refer to the phenotypically similar *ecsA*

ponA, *rasP ponA*, and *sigI ponA* double mutants as PG-limited strains.

Here, we took advantage of the conditional lethality of these PG-limited strains (*ecsA*

ponA and *rasP ponA)* to select for growth in the absence of high magnesium. This selection

revealed a cohort of suppressor mutations in genes important for fatty acid synthesis (FAS). One

of these mutants, *fapR**, encodes a super-repressor that downregulates the expression of FAS

enzymes. Further, we show that inhibiting FAS with cerulenin is sufficient to restore growth of

PG-limited mutant strains. Characterization of the *fapR** suppressor mutation highlights the

benefits of balancing membrane and PG synthesis during cell envelope biogenesis. Moreover,

these genetic findings suggest that, at least in this model system, cells lack robust genetic

mechanisms to coordinate membrane and PG synthesis, and that imbalanced synthesis can

contribute to fitness defects.

Results

Spontaneous mutations in FAS genes compensate for lethal cell wall defects in *B. subtilis*

We selected spontaneous suppressor mutations that restore growth of strains restricted

due to deficiencies in peptidoglycan (PG) synthesis (*ecsA ponA* and *rasP ponA*) in LB medium.

Using this approach, we previously described three spontaneous suppressor mutations affecting

the essential two-component system kinase, WalK, that affects expression of cell wall synthesis

related genes (18). Here, we report that this selection also led to mutations in several genes

related to FAS. In both *ecsA ponA* and *rasP ponA* genetic backgrounds, we recovered suppressor

mutations in genes encoding all four subunits of the essential enzyme acetyl-CoA carboxylase

(ACC), which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, and constitutes the

first committed step of FAS (26) (Table 1). We hypothesize that these mutations encode reduced

function alleles of ACC that restrict the capacity for membrane biosynthesis. We also recovered

both missense and deletion mutations in *fapR*, encoding the fatty acid and phospholipid synthesis

repressor FapR (27). FapR responds to malonyl-CoA as an inducer to coordinate synthesis of

fatty acids and phospholipids (28) .

fapR* is a dominant mutation that reduces expression of FapR-repressed genes

125   We focused on understanding the mechanism by which an in-frame deletion mutation,

here defined as *fapR** (Figure 1A), suppressed the growth defect of PG-limited parent strains

containing a *ponA* mutation in combination with either *ecsA*, *rasP*, or *sigI* (Figure 1B). The

ability of *fapR** to rescue growth of all three double mutant strains is consistent with our

previous data, which place *ecsA*, *rasP*, and *sigI* together in a regulatory pathway necessary for

the upregulation of *lytE* and *mreBH* to support elongasome function (18). In contrast, deletion of

fapR, which removes negative regulation on *fapR*-regulated enzymes involved in FAS, does not

rescue growth of *sigI ponA* (Figure 1C). The *fapR** allele contains a 24-nucleotide deletion in the

thioesterase-like domain overlapping with residue Phe99, known to be necessary for binding of

the malonyl-CoA inducer (Figure 1A) (28). Thus, we hypothesized that FapR* might have

reduced sensitivity to malonyl-CoA.

Further characterization of the *fapR** strain was performed to define its impact on cell

physiology. Compared to WT, *fapR** strains had a significantly slower average doubling time

(WT, 24 ± 0.033 min, *fapR**, 39 ± 0.11 min). The *fapR** mutant did not exhibit altered cell

morphology, and no statistically significant differences in the length-to-width ratios were

detected (Figure S1). Ectopic overexpression of *fapR** (but not *fapR*) resulted in a fitness defect

in a merodiploid strain also containing native *fapR*, suggesting a dominant effect (Figure 1D).

Collectively, these data led us to hypothesize that FapR* is a dominant super-repressor, and that

rescue of strains limited in their capacity for PG synthesis can be mediated by decreased

transcription of FAS enzymes.

We used qRT-PCR to directly probe the impact of FapR* on the expression levels of

selected genes in the FapR regulon. In the *fapR* null strain, expression of *fabF* and *fabHA* genes

was changed little (as expected, *fapR* was not detected), although the data are consistent with a

modest increase. This suggests that FapR may be largely inactive as a repressor in these mid-

logarithmic phase cells. In contrast, we noted a 5-10-fold decrease in the expression levels of

fabF, fabHA, and *fapR* in the *fapR** strain (Figure 2). Mutations that eliminate the FapR-

malonyl-CoA interaction are lethal, presumably because the cell is unable to sufficiently express

essential FAS genes (28). We therefore hypothesized that the *fapR** allele has reduced, but not

eliminated, the binding of malonyl-CoA to FapR.

FapR* is a super-repressor with reduced response to malonyl-CoA

FapR recognizes a 17-base pair operator sequence with a characteristic inverted repeat

sequence (27, 28). Using the FapR operator from the *fabHA-fabF* regulatory region, we used

electrophoretic mobility shift assays (EMSA) to compare the response of purified FapR and

FapR* to malonyl-CoA. When operator DNA was incubated with FapR, nearly all operator-

containing fragments were shifted, whereas the control DNA fragment was unchanged. Upon

titration with increasing concentrations of malonyl-CoA, the unshifted operator DNA band re-

appeared (Figure 3A), suggesting that FapR is released from the DNA in response to its ligand.

Conversely, DNA-binding by FapR* was relatively insensitive to malonyl-CoA (Figure 3B).

With FapR, 360 µM malonyl-CoA was sufficient for release of 50% of operator DNA, whereas

even >1 mM malonyl-CoA caused only a small decrease in the amount of FapR*-DNA complex

(Figure 3C). This supports our hypothesis that FapR* is behaving as a super-repressor to

decrease the expression of FAS enzymes, resulting in rescue of the PG-limited cells.

Addition of a membrane fluidizing agent can rescue PG-limited cells

The above results suggest that the *fapR** allele reduces membrane synthesis. However,

altering the relative expression of genes in the FAS pathway could also affect membrane

properties. One property known to be critical for the movement of the elongasome during PG

synthesis is membrane fluidity (29). This led us to speculate that *fapR** might facilitate

elongasome function by increasing membrane fluidity.

To determine whether increased membrane fluidity might be advantageous for the PG-

limited *sigI ponA* double mutant, we tested the effects of the chemical fluidizing agent benzyl

alcohol (BnOH). Indeed, BnOH rescued growth of *sigI ponA* at 37 ˚C on LB medium. We note

that under these conditions, the *sigI ponA* strain had a high propensity for generating suppressor

mutations (Figure 4A). Whole genome sequencing of selected colonies revealed mutations in

walH, a negative effector of the essential WalK-WalR two-component system (Table S1),

consistent with our previous findings where gain-of-function mutations in WalK rescue PG-

limited cells (18). However, the *rasP ponA* strain could not be rescued by BnOH under the same

growth conditions (Figure 4A). It is possible that BnOH may induce other stress responses

dependent on RasP. A good candidate is the RasP-dependent σ^W stress response, which increases

expression of *fabF* and decreases expression of *fabHA,* thereby decreasing membrane fluidity

(30). However, growth of a *sigI ponA sigW* triple mutant was still rescued by BnOH, although to

a lesser extent than a *sigI ponA* strain. To test the effects of high σ^W activity on PG-limited cells,

we sought to construct a *rasP ponA rsiW* strain. However, we were unable to construct this

strain, which is both PG-limited and predicted to have decreased membrane fluidity (30). Thus,

σ^W-dependent genes may contribute to the ability of BnOH to rescue the *sigI ponA* strain, but

other RasP-dependent pathways may also be important (Figure 4B). Since both BnOH and *fapR**

rescue growth of the *sigI ponA* mutant strain, it is possible that *fapR** affects both membrane

synthesis and fluidity.

FapR* sensitizes cells to conditions that increase membrane fluidity

To test whether *fapR** alters membrane properties we tested sensitivity to chemical

(detergent) and temperature stress. In general, cells with increased membrane fluidity often

display an increased sensitivity to detergents. Indeed, the *fapR** cells were more sensitive to

Triton X-100 than WT cells (Figure 5A). As noted above, detergents induce the σ^W stress

response, which leads to compensatory alterations in FA composition (30). Indeed, in a *sigW*

fapR* double mutant, sensitivity to Triton X-100 appeared slightly higher, albeit not statistically

significant, compared to the *fapR** single (Figure 5A). In a strain sensitized to heat by deletion of

sigI, the additional presence of a *fapR** mutation increased heat sensitivity as measured by the

reduced growth of the *sigI fapR** double mutant at 56 °C (Figure 5B). Consistently, introduction

of *fapR** rescued PG-limited strains (*ecsA ponA*, *rasP ponA*, and *sigI ponA*) at 37 °C, but not at

52 °C (Figure 5C), even though the *sigI ponA fapR** strain grew at 52 °C in the presence of

magnesium (Figure S2). To determine the effects of the *fapR** mutation on the fatty acid

constituents of the cell membrane, we performed fatty acid methyl ester (FAME) analysis.

FapR* cells contained a slightly lower proportion of straight FA chains and a significant increase

in the ratio of C_{17} to C_{15} acyl-chains (Table S2). It is unclear whether these changes in the FA

composition of the membranes directly contribute to the apparent changes in membrane fluidity.

Overall, the increased sensitivity of *fapR** to both detergents and elevated temperature supports

the hypothesis that *fapR** strains have membranes with altered physical properties.

FapR* increases sensitivity to cerulenin, a FAS inhibitor

Our results suggest that *fapR** rescue of PG-limited cells is correlated with reduced

expression of FAS genes. We therefore predicted that *fapR** mutants may have an increased

sensitivity to cerulenin. Cerulenin is a 12-carbon mycotoxin that inhibits FAS by forming

covalent linkages with β-ketoacyl-ACP synthases, including FabF, to block FA elongation (31-

33). Upon inhibition by cerulenin, malonyl-CoA accumulates, triggering derepression of genes in

the FapR regulon. Since FapR* is less responsive to malonyl-CoA (Figure 3), we anticipated that

this compensatory mechanism might be impaired. As predicted, *fapR** mutant strains had greatly

increased sensitivity to cerulenin at concentrations that did not inhibit WT or *fapR* null mutant

strains (Figure 6). This effect was specific to cerulenin and unlikely to reflect general membrane

permeability effects since no change was noted in sensitivity to other common hydrophobic

antibiotics (Figure S3). Interestingly, *fapR** also did not affect sensitivity to triclosan (Figure 6),

an antibiotic which inhibits the second reduction step of FAS and, like cerulenin, normally

triggers induction of the *fapR* regulon (34).

Chemical inhibition of FAS is sufficient to rescue PG-limited *rasP ponA* cells

Since *fapR** rescues growth of PG-limited strains by restricting FAS, we hypothesized

that antibiotics that inhibit FAS might also rescue growth. Remarkably, cerulenin rescued growth

of the PG-limited *rasP ponA* strain (Figure 7). With optimal concentrations of cerulenin (1.65 or

3.3 µg mL^{-1}), growth was comparable to that of the magnesium supplemented medium (Figure

7). Cerulenin can also lead to a reduction in protein synthesis similar to that experienced

following nutrient deprivation (35, 36). However, PG-limited cells could not be rescued by

treatment with sub-MIC levels of the protein synthesis inhibiter chloramphenicol (Figure S4A),

or by reducing growth rate through non-preferred carbon sources (Figure S4B). Previously, cells

lacking (p)ppGpp were found to be more sensitive to cerulenin (37). However, in our strains the

effect of *fapR** appear to be independent of (p)ppGpp synthesis. Specifically, a *sigI* strain is

sensitive to cefuroxime (CEF), which inactivates PBP1, and this sensitivity is suppressed by

fapR* in both *sigI* and *sigI* (p)ppGpp null strains (Figure S4C).

Cerulenin is antagonistic with selected PG synthesis inhibitors

We reasoned that if restriction of FAS by cerulenin rescues strains that are PG-limited

due to mutations (*rasP ponA*), it might also reduce sensitivity to PG-targeting antibiotics. To

determine if cerulenin affects sensitivity to antibiotics that target PG synthesis we used a

checkerboard assay. When WT *B. subtilis* was treated with cerulenin and CEF, we observed a

fractional inhibitory concentration index (FICI) of 2.5 (Table 2), indicating an additive effect.

However, in a *sigI* mutant (which is more CEF sensitive than WT), we observed strong

cerulenin-CEF antagonism (FICI of 16) (Table 2). Further, treatment of WT cells with low doses

of cerulenin inhibits the characteristic CEF-induced lysis evident as a loss in optical density

(Figure S5). We conclude that antibiotics that inhibit FAS (cerulenin) and cell wall synthesis

(CEF) display an antagonistic interaction in the *sigI* mutant. We next tested a variety of

antibiotics targeting different steps in PG synthesis, including fosfomycin (UDP-GlcNAc-enoyl

pyruvate synthesis), D-cycloserine (D-Ala-D-Ala synthesis), bacitracin (undecaprenyl-

pyrophosphate recycling), and nisin and vancomycin (lipid II binding and sequestration). Of

these compounds, only cerulenin and bacitracin displayed an antagonistic relationship (Table 2),

We conclude that inhibition of FAS can reduce the efficacy of antibiotics that affect membrane-

localized steps in PG synthesis.

Discussion

All cells are defined by a membrane, a boundary layer comprised largely of lipids and

proteins. In Bacteria, the cell membrane is rich in phospholipids and houses numerous proteins

involved in transport of nutrients and ions, energy generation, and cell signaling. The membrane

is supported by the peptidoglycan cell wall, a covalently crosslinked meshwork that provides

structural stability and shape to the cell. In many Bacteria, additional layers may also be present

including the lipopolysaccharide-containing outer membrane in Gram-negative bacteria, S-

layers, or mycomembranes. Collectively, the resulting multilayered structure is defined as the

cell envelope.

Considerable progress has been made in defining how synthesis of each individual

envelope layer is regulated. For example, *in B. subtilis,* FAS and phospholipid synthesis are

coordinated in response to malonyl-CoA precursor availability by the FapR repressor. Depletion

of the fatty acyl-phosphate synthase, PlsX, halts both FAS and phospholipid synthesis, which

suggests that *plsX* also plays a key regulatory role in coordinating FAS and phospholipid

synthesis (38, 39). Similarly, PG synthesis is homeostatically regulated by multiple mechanisms

that control the GlmS-dependent partition of fructose 6-phosphate into amino sugar synthesis.

GlmS expression is feedback regulated by a GlcNAc-sensing riboswitch, and UDP-GlcNAc

negatively regulates the activity of GlmR, an activator of GlmS activity (40, 41). In Gram-

negative bacteria, stabilization or degradation of LpxC, the rate-limiting enzyme for

lipopolysaccharide (LPS) biogenesis, fine-tunes LPS synthesis in response to situational demand

(42). For example, phospholipids mislocalized to the outer leaflet of the outer membrane are

degraded by the phospholipase, PldA, and the resulting fatty acyl-CoA molecules inhibit LpxC

degradation and increase LPS synthesis (43). While these mechanisms regulate individual

structures within the cell envelope, it is presently unclear if or how bacteria coordinate the

synthesis of the various cell envelope layers with each other.

Synthesis of the individual cell envelope layers must be uncoupled under some

conditions. For example, L-forms are able to synthesize membranes even in the absence of PG

(44-46), and photosynthetic Bacteria can synthesize abundant internal membranes without a

coordinate increase in PG synthesis (47). In *E. coli*, FAS has been implicated as a main

determinant of cell size (36). Spontaneous suppressor mutations in *fabH* can compensate for

lethal defects in LPS synthesis, and also reduce the rate of cell envelope growth (48). Further, *B.*

subtilis supplemented with exogenous fatty acids will increase in size by up to 10 percent,

suggesting that increased membrane synthesis can allow increased cell wall synthesis (36).

Here, we sought to better understand the stresses associated with a genetically-imposed,

limited capacity for PG synthesis. As shown previously, cells lacking all four aPBPS ($\Delta 4$ aPBP)

or even just the dominant aPBP (*ponA*) are viable on LB medium only because stress responses

increase the synthetic capacity of the elongasome. The activation of σ^M increases expression of

the RodA TG, as well as other PG synthesis enzymes, and a *ponA sigM* double mutant is inviable

(14). Similarly, σ^I is required in a *ponA* null mutant to increase elongasome activity by

increasing expression of elongasome-associated MreBH and LytE proteins (18).

Using forward genetics, we selected for suppressors of PG-limited strains lacking the

major aPBP (*ponA* mutation) and defective in the σ^I pathway (e.g. *ecsA, rasP* or *sigI* mutants).

One mechanism of rescue involves the up-regulation of the essential WalK-WalR two-

component system through either gain-of-function mutations in WalK (18) or mutations

affecting the negative regulator WalH (Table S1). Here, we demonstrate that this growth

restriction can also be rescued by mutations affecting FAS. We recovered missense mutations

that we hypothesize decrease ACC activity. However, efforts to further explore these mutations

were frustrated by our inability to introduce these mutations into WT cells. This hints that an

insufficient capacity for FAS is not well tolerated in cells with normal PG synthesis capacity.

YqhY has been implicated as a possible negative regulator of ACC (49, 50). However, induction

of *yqhY* did not rescue our PG-limited strains, suggesting that simply overexpressing YqhY may

not be sufficient to downregulate ACC activity.

We focused our attention on *fapR**, an allele that decreases expression of FAS genes and

restores growth on LB medium. The genetic rescue we observe is likely due to a decrease in

membrane synthesis. Indeed, reducing FAS activity with cerulenin can also rescue growth

(Figure 7). Our phenotypic analysis suggests that the *fapR** allele may also alter membrane

properties. Membrane fluidity is important for elongasome-dependent PG-synthesis (51-53), and

the *fapR** allele increases sensitivity to both detergents and temperature. These phenotypes are

suggestive of a possible fluidizing effect on the membrane. Further, a membrane-fluidizing agent

can also rescue growth of the PG-limited *sigI ponA* strain (Figure 4). Taken together, the

decrease in expression of FAS genes, possibly in conjunction with altered membrane properties,

enables the *fapR** allele to restore growth of PG-limited cells. There is currently no known

mechanism in *B. subtilis* to coordinate FAS with PG assembly. In the absence of such a

regulatory mechanism, antibiotics can perturb balanced synthesis and thereby create an

opportunity for spontaneous suppressors to develop, such as *fapR**, that re-balance FAS with PG

synthesis.

Collectively, our results suggest that ongoing membrane synthesis in PG-limited cells

contributes to cell death, and that cells lack a robust mechanism to slow membrane production

when PG synthesis is restricted. These findings highlight the importance of maintaining a

balance between membrane and PG synthesis. A more complete understanding of the balance

between FAS and cell wall synthesis can provide further insights relevant to antimicrobial

treatments and the maintenance of cell homeostasis.

Materials and Methods

Growth conditions, bacterial strains, and plasmids

All strains were cultured in lysogeny broth (LB) medium at 37 °C and aerated on an orbital

shaker at 300 RPM. Glycerol stocks were streaked on LB agar plates and grown overnight at 37

°C. When needed, antibiotics were used in growth media at the following concentrations: 100

μg/mL ampicillin; 10 μg/mL chloramphenicol; mls (1 μg/mL erythromycin plus 2.5 μg/mL

lincomycin); 10 µg/mL kanamycin; 100 µg/mL spectinomycin, 10 µg/mL tetracycline. All

synthetic lethal strains were grown in media supplemented with 20 mM $MgSO_4$. Bacterial strains

used in this study are listed in supplemental Table S3. Deletion strains were created by utilizing

the BKK/BKE genomic knockout library available at the Bacillus Genome Stock Center

(BGSC). Gene deletions with kanamycin or erythromycin cassettes were moved into the lab WT

168 strain, taking advantage of the natural competence of *B. subtilis* grown to OD_{600nm} 0.8 in

modified competence (MC) media. Null gene mutations were created by removing the antibiotic

resistance cassette to create a clean in-frame null mutation using pDR244 as previously

described (Koo et al., 2017). Null mutations were confirmed via colony PCR using the

designated check primers as detailed in Table S4. For strains with low levels of natural

competence, SPP1 transfection was used to create mutations as preciously described (54). The

FapR* suppressor mutations were reconstructed via CRISPR-cas9 mutagenesis using the

pJOE8999 as described previously (55). Primer sequences for repair template, guide RNA and

construct confirmation can be found in Table S4. Constructs produced via CRISPR-cas9

mutagenesis were confirmed via Sanger sequencing performed by the Cornell University

Biotechnology Resource Center (BRC). Genes were expressed ectopically at the *amyE* locus

using the pPL82 plasmid (56) and expressed from a P$_{spac(hy)}$ promoter after induction with 1 mM

IPTG.

Plating efficiency

All strains were grown in 5 mL LB supplemented with 20 mM $MgSO_4$ at 37 °C with shaking

until OD_{600nm} ~0.4-0.5. The cells were washed by centrifuging 1 mL of culture at 5000 RPM, and

the pellet was resuspended in 1 mL LB without additional magnesium. Serial dilutions were

performed and 10 µL of culture was spotted onto an LB-agar plate, air dried for 15 minutes and

then incubated overnight at 37 °C. N=3. A representative image is shown.

Growth assay

From a colony, strains were grown in 5 mL LB medium supplemented with 20 mM $MgSO_4$ at 37

°C with shaking until OD_{600nm} ~0.4-0.5. The cells were washed by centrifuging 1 mL of culture

at 5000 RPM, and the pellet was resuspended in 1 mL LB medium to remove traces of $MgSO_4$ or

resuspended in minimal media (15 mM $(NH_4)2SO_4$, 0.8 mM $MgSO_4$ $7H_2O$, 3.4 mM sodium

citrate dihydrate, 2 mM KPO_4, 4.2 mM potassium glutamate, 40 mM morpholinepropanesulfonic

acid (MOPS), pH 7.4, 0.25 mM Tryptophan, 5 µM $FeSO_4$, 5 µM $MnCl_2$, 2 % carbon as

specified) as required. 1 µL of this culture was added to 199 µL of media with the desired drug

concentration into the well of a 100-well Honeycomb plate (Steri). The OD_{600nm} of each well was

measured at 37 °C with shaking in a Bioscreen C growth curve analyzer (Growth Curves USA,

NJ) every 15 minutes up to a maximum time of 24 hours. All data shown is representative plots

and standard deviations from three biological replicates.

Disc diffusion assay

Cultures were grown in LB medium at 37 °C with aeration and synthetic lethal cultures were

supplemented with 20 mM $MgSO_4$. Cells were harvested at ~0.4 OD_{600nm}, and where applicable,

pelleted and resuspended in fresh LB medium to remove traces of $MgSO_4$. 100 µL of the liquid

culture was added to 4 mL of top agar (0.75 % agar) maintained at 50 °C to prevent

solidification. After mixing, top agar and cells were poured onto a 15 mL agar plate (1.5 % agar)

and allowed to solidify. Top agar was allowed to dry for 15 minutes prior to placing a 6 mm

diameter Whatman paper filter disc on the surface of the top agar. The compound of interest was

them immediately applied to the filter disc and the plates were incubated overnight at 37 °C. The

diameter of the zone of inhibition or the diameter of the zone of growth were measured and

recorded. Compounds were used at the following concentrations: 10 µL of a 25 % solution

Triton X-100, 10 µL of a 10 % solution benzyl alcohol (BnOH), 10 µg cerulenin, 5 µg triclosan

unless otherwise specified, 0.8 µg erythromycin, 8 µg chloramphenicol, 10 µg cefuroxime.

Transcript analysis (Real-time RT-PCR)

To determine gene expression via RT-PCR, strains were grown in LB medium to ~OD_{600nm} 0.4.

To extract RNA, 1.5 mL of cells were incubated at 37 °C for 15 minutes in lysozyme-TE (30

mM Tris-HCl, 1 mM EDTA; pH 8.0 with 15 mg/mL lysozyme) to mediate cell lysis. RNA was

purified as per the manufacturer's instructions using the Qiagen RNeasy Kit. The extracted RNA

was DNase treated according to the Turbo DNase kit manufacturer's instructions (Ambion). 2 µg

of RNA was used in a 20 µL reaction to synthesize 100 ng/µL of cDNA using the Applied

Biosystems high-capacity reverse transcription kit. Gene expression was measured using 100 ng

of cDNA and 0.5 µM of gene specific primer (see Table S4) and 1x SYBR (Bio-Rad) in a qPCR

QuantStudio 7 Pro (ThermoFischer Scientific). *gyrA* was used as a house-keeping gene. Gene

expression was normalized to *gyrA* and then expression values ($2^{-\Delta ct}$) were recorded for each

gene.

Purification of FapR and FapR*

E. coli BL21 DE3 cells were freshly transformed with a pET-16b expression vector containing

an IPTG inducible copy of *fapR* or *fapR** with an N-terminal 10x-His tag. Transformants were

grown overnight in LB medium supplemented with ampicillin as specified and 1 % glucose. The

next day 100 µL of the overnight culture was used to inoculate a 3 mL pre-culture of LB medium

with 1 % glucose and ampicillin. The pre-culture was used to inoculate 500 mL of LB medium

with 1 % glucose and ampicillin and allowed to grow until ~0.2 OD_{600nm}. Cultures were induced

with 1 mM IPTG and additional ampicillin and left at 37 °C for 3 hours. Cultures were spun

down in a chilled rotor at 3500 RPM for 20 minutes at 4 °C and pellets were stored at -80 °C

until purification. Cells pellets were resuspended in 5 mL chilled resuspension buffer (20 mM

Tris-HCl, 500 M NaCl, 10 mM Imidazole, 0.5 tablet of cOmplete Mini protease inhibitor

cocktail (Roche), pH 7.4). Cells were lysed via sonication and then spun down at 4 °C at 12,000

RPM for 30 minutes. The supernatant was further processed using a 1 mL HisTrap HP column

(Cytiva), prepared according to the manufacturer's directions, washed with wash buffer (20 mM

Tris-HCl, 500 mM NaCl, 30 mM imidazole, pH 7.4) and eluted with elution buffer (20 mM Tris-

HCl, 500 mM NaCl, 500 mM imidazole, pH 7.4). The concentration of the purified protein was

determined by Bradford assay with a BSA standard (57).

Electrophoretic mobility shift assay (EMSA). Labeled DNA was prepared using FAM

conjugated oligos (IDT) to amplify a portion of the *pfabHAF* region (see Table S4 for primers)

and the resulting amplicon was digested with *ndeI* and purified to produce FAM fragments either

with or without the *fapR* binding consensus sequence. Equimolar amounts of protein and DNA

were added to 10 µL reactions prepared using buffer as detailed in Schujman et al., 2003.

Briefly, the buffer consisted of 20 mM Tris-HCl (pH 8.0), 0.5 mM EDTA, 50 mM NaCl, 1 mM

DTT, 5 % glycerol. The reactions were incubated in the dark for 10 minutes at 37 °C before

addition of malonyl-CoA followed by an additional 10-minute incubation period under the same

conditions. The samples were loaded onto a pre-run 10 % non-denaturing acrylamide gel in 1x

TBE buffer. Electrophoresis was performed at 100 V and the gel was imaged using a Typhoon

FLA 7000 (GE Healthcare Life Sciences) and quantification was performed using Fiji (58).

Fatty acid methyl esterase analysis (FAME)

Freshly streaked strains were grown in LB medium overnight at 37 °C with shaking. The pre-

culture was then used to inoculate 100 mL of LB medium and grown at 37 °C with shaking until

0.4 OD_{600nm}. 20-30 mg of cells were harvested and stored at -80 °C. Sample preparation and

FAME direct analysis was performed by Microbial ID (Newark, DE). Reported values represent

the average and standard deviations of biological triplicates.

Microscopic evaluation

The indicated strains were grown until ~0.4 OD_{600nm} and spotted onto a slide containing a thin

pad prepared from 1% agarose. The cells were allowed to air-dry before application of a cover

slip. The cells were imaged at 100x magnification with immersion oil using an Olympus BX61

microscope and images were captured using a Cooke Sensicam camera. Cell lengths and widths

were determined using the Fiji (58) plugin, MicrobeJ (59). A minimum of 100 cell lengths and

widths were recorded for statistical analysis.

Minimal inhibitory concentration (MIC) determination and fractional inhibitory

concentration index (FICI) calculation

An MIC for each indicated drug was first identified as the drug concentration where a WT strain

did not reach ~0.2 OD_{600nm} following 10 hours of treatment. The MIC of drugs individually are

as follows: 1000 µg/mL bacitracin, 1.28 µg/mL cefuroxime, 5 µg/mL cerulenin, 1 µg/mL

chloramphenicol, 50 µg/mL D-cycloserine, 50 µg/mL fosfomycin, 200 µg/mL nisin, 0.75 µg/mL

triclosan, 0.15 µg/mL vancomycin. The index (FICI) for drug combinations was calculated as

follows with FIC = (A/MIC$_A$) + (B/MIC$_B$) where A and B are the MIC of each antibiotic in

combination and MIC$_A$ and MIC$_B$ are the MIC of each antibiotic individually. The FICI of < 0.5

is considered as synergistic, 0.5-4 as additive and >4 as antagonistic (60).

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health (grant R35GM122461 to JDH).

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript. We acknowledge Heng Zhao for guidance with the reconstruction

of *fapR** suppressor via CRISPR-cas9 mutagenesis. We thank the Angert lab for use of their

light microscope and Yesha Patel, Ankita Sachla, and Tobias Dörr for helpful comments.

References

1.   Egan AJF, Errington J, Vollmer W. 2020. Regulation of peptidoglycan synthesis and remodelling.

Nat Rev Microbiol 18:446-460.

2.   Zhao H, Patel V, Helmann JD, Dörr T. 2017. Don't let sleeping dogmas lie: new views of

peptidoglycan synthesis and its regulation. Mol Microbiol 106:847-860.

3.   Nguyen LT, Gumbart JC, Beeby M, Jensen GJ. 2015. Coarse-grained simulations of bacterial cell

wall growth reveal that local coordination alone can be sufficient to maintain rod shape. Proc

Natl Acad Sci U S A 112:E3689-98.

4.   Barreteau H, Kovac A, Boniface A, Sova M, Gobec S, Blanot D. 2008. Cytoplasmic steps of

480   peptidoglycan biosynthesis. FEMS Microbiol Rev 32:168-207.

481   5.   Claessen D, Emmins R, Hamoen LW, Daniel RA, Errington J, Edwards DH. 2008. Control of the cell

482   elongation-division cycle by shuttling of PBP1 protein in *Bacillus subtilis*. Mol Microbiol 68:1029-

483   46.

484   6.   Garner EC, Bernard R, Wang W, Zhuang X, Rudner DZ, Mitchison T. 2011. Coupled,

485   circumferential motions of the cell wall synthesis machinery and MreB filaments in *B. subtilis*.

486   Science 333:222-5.

487   7.   Domínguez-Escobar J, Chastanet A, Crevenna AH, Fromion V, Wedlich-Söldner R, Carballido-

488   López R. 2011. Processive movement of MreB-associated cell wall biosynthetic complexes in

489   bacteria. Science 333:225-8.

490   8.   Vigouroux A, Cordier B, Aristov A, Alvarez L, Özbaykal G, Chaze T, Oldewurtel ER, Matondo M,

491   Cava F, Bikard D, van Teeffelen S. 2020. Class-A penicillin binding proteins do not contribute to

492   cell shape but repair cell-wall defects. Elife 9.

493   9.   Straume D, Piechowiak KW, Kjos M, Håvarstein LS. 2021. Class A PBPs: It is time to rethink

494   traditional paradigms. Mol Microbiol 116:41-52.

495   10.   Dion MF, Kapoor M, Sun Y, Wilson S, Ryan J, Vigouroux A, van Teeffelen S, Oldenbourg R, Garner

496   EC. 2019. *Bacillus subtilis* cell diameter is determined by the opposing actions of two distinct cell

497   wall synthetic systems. Nat Microbiol 4:1294-1305.

498   11.   Helmann JD. 2016. *Bacillus subtilis* extracytoplasmic function (ECF) sigma factors and defense of

499   the cell envelope. Curr Opin Microbiol 30:122-132.

500   12.   Eiamphungporn W, Helmann JD. 2008. The *Bacillus subtilis* sigma(M) regulon and its

501   contribution to cell envelope stress responses. Mol Microbiol 67:830-48.

502   13.   Mascher T, Hachmann AB, Helmann JD. 2007. Regulatory overlap and functional redundancy

503   among *Bacillus subtilis* extracytoplasmic function sigma factors. J Bacteriol 189:6919-27.

504   14.   Meeske AJ, Riley EP, Robins WP, Uehara T, Mekalanos JJ, Kahne D, Walker S, Kruse AC,

505   Bernhardt TG, Rudner DZ. 2016. SEDS proteins are a widespread family of bacterial cell wall

506   polymerases. Nature 537:634-638.

507   15.   McPherson DC, Popham DL. 2003. Peptidoglycan synthesis in the absence of class A penicillin-

508   binding proteins in *Bacillus subtilis*. J Bacteriol 185:1423-31.

509   16.   Sjodt M, Rohs PDA, Gilman MSA, Erlandson SC, Zheng S, Green AG, Brock KP, Taguchi A, Kahne

510   D, Walker S, Marks DS, Rudner DZ, Bernhardt TG, Kruse AC. 2020. Structural coordination of

511   polymerization and crosslinking by a SEDS-bPBP peptidoglycan synthase complex. Nat Microbiol

512   5:813-820.

513   17.   Emami K, Guyet A, Kawai Y, Devi J, Wu LJ, Allenby N, Daniel RA, Errington J. 2017. RodA as the

514   missing glycosyltransferase in *Bacillus subtilis* and antibiotic discovery for the peptidoglycan

515   polymerase pathway. Nat Microbiol 2:16253.

516   18.   Patel Y, Zhao H, Helmann JD. 2020. A regulatory pathway that selectively up-regulates

517   elongasome function in the absence of class A PBPs. Elife 9.

518   19.   Heinrich J, Lundén T, Kontinen VP, Wiegert T. 2008. The *Bacillus subtilis* ABC transporter EcsAB

519   influences intramembrane proteolysis through RasP. Microbiology (Reading) 154:1989-1997.

520   20.   Hastie JL, Williams KB, Ellermeier CD. 2013. The activity of σ^V, an extracytoplasmic function σ

521   factor of *Bacillus subtilis*, is controlled by regulated proteolysis of the anti-σ factor RsiV. J

522   Bacteriol 195:3135-44.

523   21.   Hastie JL, Williams KB, Sepúlveda C, Houtman JC, Forest KT, Ellermeier CD. 2014. Evidence of a

524   bacterial receptor for lysozyme: binding of lysozyme to the anti-σ factor RsiV controls activation

525   of the ECF σ factor σ^V. PLoS Genet 10:e1004643.

526   22.   Parrell D, Zhang Y, Olenic S, Kroos L. 2017. *Bacillus subtilis* Intramembrane Protease RasP

527   Activity in *Escherichia coli* and In Vitro. J Bacteriol 199.

528   23.   Liu TY, Chu SH, Hu YN, Wang JJ, Shaw GC. 2017. Genetic evidence that multiple proteases are

529   involved in modulation of heat-induced activation of the sigma factor SigI in *Bacillus subtilis*.

530   FEMS Microbiol Lett 364.

531   24.   Zuber U, Drzewiecki K, Hecker M. 2001. Putative sigma factor SigI (YkoZ) of *Bacillus subtilis* is

532   induced by heat shock. J Bacteriol 183:1472-5.

533   25.   Tesson B, Dajkovic A, Keary R, Marlière C, Dupont-Gillain CC, Carballido-López R. 2022.

534   Magnesium rescues the morphology of *Bacillus subtilis mreB* mutants through its inhibitory

535   effect on peptidoglycan hydrolases. Sci Rep 12:1137.

536   26.   Tong L. 2005. Acetyl-coenzyme A carboxylase: crucial metabolic enzyme and attractive target for

537   drug discovery. Cell Mol Life Sci 62:1784-803.

538   27.   Schujman GE, Paoletti L, Grossman AD, de Mendoza D. 2003. FapR, a bacterial transcription

539   factor involved in global regulation of membrane lipid biosynthesis. Dev Cell 4:663-72.

540   28.   Schujman GE, Guerin M, Buschiazzo A, Schaeffer F, Llarrull LI, Reh G, Vila AJ, Alzari PM, de

541   Mendoza D. 2006. Structural basis of lipid biosynthesis regulation in Gram-positive bacteria.

542   Embo J 25:4074-83.

543   29.   Strahl H, Bürmann F, Hamoen LW. 2014. The actin homologue MreB organizes the bacterial cell

544   membrane. Nat Commun 5:3442.

545   30.   Kingston AW, Subramanian C, Rock CO, Helmann JD. 2011. A σ^W-dependent stress response in

546   *Bacillus subtilis* that reduces membrane fluidity. Mol Microbiol 81:69-79.

547   31.   Moche M, Schneider G, Edwards P, Dehesh K, Lindqvist Y. 1999. Structure of the complex

548   between the antibiotic cerulenin and its target, beta-ketoacyl-acyl carrier protein synthase. J

549   Biol Chem 274:6031-4.

550   32.   Campbell JW, Cronan JE, Jr. 2001. Bacterial fatty acid biosynthesis: targets for antibacterial drug

551   discovery. Annu Rev Microbiol 55:305-32.

552   33.   Zhang YM, White SW, Rock CO. 2006. Inhibiting bacterial fatty acid synthesis. J Biol Chem

553   281:17541-4.

554   34.   Hutter B, Schaab C, Albrecht S, Borgmann M, Brunner NA, Freiberg C, Ziegelbauer K, Rock CO,

555   Ivanov I, Loferer H. 2004. Prediction of mechanisms of action of antibacterial compounds by

556   gene expression profiling. Antimicrob Agents Chemother 48:2838-44.

557   35.   Wille W, Eisenstadt E, Willecke K. 1975. Inhibition of de novo fatty acid synthesis by the

558   antibiotic cerulenin in *Bacillus subtilis*: effects on citrate-Mg^{2+} transport and synthesis of

559   macromolecules. Antimicrob Agents Chemother 8:231-7.

560   36.   Vadia S, Tse JL, Lucena R, Yang Z, Kellogg DR, Wang JD, Levin PA. 2017. Fatty Acid Availability

561   Sets Cell Envelope Capacity and Dictates Microbial Cell Size. Curr Biol 27:1757-1767.e5.

562   37.   Pulschen AA, Sastre DE, Machinandiarena F, Crotta Asis A, Albanesi D, de Mendoza D, Gueiros-

563   Filho FJ. 2017. The stringent response plays a key role in *Bacillus subtilis* survival of fatty acid

564   starvation. Mol Microbiol 103:698-712.

565   38.   Paoletti L, Lu YJ, Schujman GE, de Mendoza D, Rock CO. 2007. Coupling of fatty acid and

566   phospholipid synthesis in *Bacillus subtilis*. J Bacteriol 189:5816-24.

567   39.   Machinandiarena F, Nakamatsu L, Schujman GE, de Mendoza D, Albanesi D. 2020. Revisiting the

568   coupling of fatty acid to phospholipid synthesis in bacteria with FapR regulation. Mol Microbiol

569   114:653-663.

570   40.   Winkler WC, Nahvi A, Roth A, Collins JA, Breaker RR. 2004. Control of gene expression by a

571   natural metabolite-responsive ribozyme. Nature 428:281-6.

41.   Patel V, Wu Q, Chandrangsu P, Helmann JD. 2018. A metabolic checkpoint protein GlmR is

important for diverting carbon into peptidoglycan biosynthesis in *Bacillus subtilis*. PLoS Genet

14:e1007689.

42.   Thomanek N, Arends J, Lindemann C, Barkovits K, Meyer HE, Marcus K, Narberhaus F. 2018.

Intricate Crosstalk Between Lipopolysaccharide, Phospholipid and Fatty Acid Metabolism in

Escherichia coli Modulates Proteolysis of LpxC. Front Microbiol 9:3285.

43.   May KL, Silhavy TJ. 2018. The *Escherichia coli* Phospholipase PldA Regulates Outer Membrane

Homeostasis via Lipid Signaling. mBio 9.

44.   Errington J, Mickiewicz K, Kawai Y, Wu LJ. 2016. L-form bacteria, chronic diseases and the origins

of life. Philos Trans R Soc Lond B Biol Sci 371.

45.   Mercier R, Kawai Y, Errington J. 2013. Excess membrane synthesis drives a primitive mode of cell

proliferation. Cell 152:997-1007.

46.   Siddiqui RA, Hoischen C, Holst O, Heinze I, Schlott B, Gumpert J, Diekmann S, Grosse F, Platzer

M. 2006. The analysis of cell division and cell wall synthesis genes reveals mutationally

inactivated *ftsQ* and *mraY* in a protoplast-type L-form of *Escherichia coli*. FEMS Microbiol Lett

258:305-11.

47.   Niederman RA. 2013. Membrane development in purple photosynthetic bacteria in response to

alterations in light intensity and oxygen tension. Photosynth Res 116:333-48.

48.   Yao Z, Davis RM, Kishony R, Kahne D, Ruiz N. 2012. Regulation of cell size in response to nutrient

availability by fatty acid biosynthesis in *Escherichia coli.* Proc Natl Acad Sci U S A 109:E2561-8.

49.   Tödter D, Gunka K, Stülke J. 2017. The Highly Conserved Asp23 Family Protein YqhY Plays a Role

in Lipid Biosynthesis in *Bacillus subtilis*. Front Microbiol 8:883.

50.   Willdigg JR, Helmann JD. 2021. Mini Review: Bacterial Membrane Composition and Its

Modulation in Response to Stress. Frontiers in Molecular Biosciences 8.

51. Kurita K, Kato F, Shiomi D. 2020. Alteration of Membrane Fluidity or Phospholipid Composition

Perturbs Rotation of MreB Complexes in *Escherichia coli*. Front Mol Biosci 7:582660.

52. van Teeffelen S, Wang S, Furchtgott L, Huang KC, Wingreen NS, Shaevitz JW, Gitai Z. 2011. The

bacterial actin MreB rotates, and rotation depends on cell-wall assembly. Proc Natl Acad Sci U S

A 108:15822-7.

53. Zielińska A, Savietto A, de Sousa Borges A, Martinez D, Berbon M, Roelofsen JR, Hartman AM, de

Boer R, Van der Klei IJ, Hirsch AK, Habenstein B, Bramkamp M, Scheffers DJ. 2020. Flotillin-

mediated membrane fluidity controls peptidoglycan synthesis and MreB movement. Elife 9.

54. Kearns DB, Chu F, Branda SS, Kolter R, Losick R. 2005. A master regulator for biofilm formation

by *Bacillus subtilis*. Mol Microbiol 55:739-49.

55. Altenbuchner J. 2016. Editing of the *Bacillus subtilis* Genome by the CRISPR-Cas9 System. Appl

Environ Microbiol 82:5421-7.

56. Quisel JD, Burkholder WF, Grossman AD. 2001. In vivo effects of sporulation kinases on mutant

Spo0A proteins in *Bacillus subtilis*. J Bacteriol 183:6573-8.

57. Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities

of protein utilizing the principle of protein-dye binding. Anal Biochem 72:248-54.

58. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A.

2012. Fiji: an open-source platform for biological-image analysis. Nat Methods 9:676-82.

59. Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput bacterial cell

detection and quantitative analysis. Nat Microbiol 1:16077.

60. Odds FC. 2003. Synergy, antagonism, and what the chequerboard puts between them. J

Antimicrob Chemother 52:1.

Table 1. **Spontaneous suppressor mutations recovered in PG-limited cells.** Mutants of the

ecsA ponA and *rasP ponA* parent strains were selected that were able to grow in LB medium in

the absence of magnesium supplementation. Each suppressor strain contained a single point

mutation identified via whole genome re-sequencing. The genome position was determined via

comparison to the *B. subtills* reference genome NC_000964.3.

Isolate	Strain Background	Gene	Genome Position	Nucleotide Change	Mutation
HB23854	*ecsA ponA*	*fapR*	1662276	A --> G	Δ24 bp deletion in CDC/ I104V
HB23855	*ecsA ponA*	*fapR*	1662301	C --> T	A112V
HB23856	*ecsA ponA*	*accA*	2988324	C --> A	G129W
HB23857	*ecsA ponA*	*accA*	2988009	A --> G	T234R
HB23860	*ecsA ponA*	*accB*	2532222	T --> C	SNP between promoter and RBS
HB23885	*rasP ponA*	*fapR*	1662190	T --> A	I75N
HB23874	*rasP ponA*	*accA*	2988055	G --> T	N218K
HB23875	*rasP ponA*	*accC*	2531202	T --> C	R169G
HB23883	*rasP ponA*	*accC*	2531393	G --> A	P105L
HB23877	*rasP ponA*	*accD*	2989081	G --> A	T162M
HB23879	*rasP ponA*	*accD*	2989654	T --> deletion	deletion in promoter

Table 2. FICI values of cerulenin in combination with selected cell wall synthesis targeting antibiotics. An MIC was defined as having and OD_{600} below 0.2 after 10 hours of treatment. Susceptibility was defined as follows: < 0.5 synergistic, 0.5-4 indifferent, > 5 antagonistic.

Strain	Compound A	Compound B	FIC	Susceptibility Pattern
WT	Bacitracin	Cerulenin	5	Antagonistic
	Bacitracin	Triclosan	1.03	Indifferent
	Bacitracin	Chloramphenicol	1	Indifferent
	Cefuroxime	Cerulenin	2.5	Indifferent
	D-Cycloserine	Cerulenin	2	Indifferent
	Fosfomycin	Cerulenin	3	Indifferent
	Nisin	Cerulenin	2	Indifferent
	Vancomycin	Cerulenin	2	Indifferent
sigI	Cefuroxime	Cerulenin	16	Antagonistic

FIGURE LEGENDS

Figure 1. **FAS suppressor mutations rescue cells with lethal PG synthesis defects.** (A) The

FapR* 24-nucleotide deletion occurs in the thioesterase-like, malonyl-CoA binding domain of

the protein. Asterisks indicate residues important for coordinating the interaction between FapR

and malonyl-CoA. (B) *fapR** rescues viability of *ecsA ponA*, *rasP ponA*, and *sigI ponA* cells in

LB medium in the absence of magnesium supplementation. (C) Deletion of FapR does not rescue

viability of a *sigI ponA* mutant grown in LB medium in the absence of magnesium

supplementation. (D) *fapR** is dominant in a merodiploid strain expressing at locus *fapR*.

Figure 2. *fapR** **decreases the expression of FAS genes.** (A) Gene expression from candidate

genes in the *fapR* regulon (*fabF, fapR and fabHA)* is decreased in a *fapR** mutant. The x-axis

indicates candidate genes detected, and the legend indicates strain genotypes. ND indicates a

lack of *fapR* gene expression in a *fapR* null background. Gene expression values ($2^{-\Delta ct}$) were

normalized to *gyrA* and plotted. A one-way ANOVA with a Tukey test for multiple comparisons

was performed. P-value cutoff (*) < 0.05.

Figure 3. **FapR* exhibits altered interactions with malonyl-CoA.** (A) Gel shift assay where a

FapR-DNA complex was titrated with increasing concentrations of malonyl-CoA. The lower

DNA bands serve both as a loading control and control for protein interaction specificity as it

does not contain the FapR recognized inverted repeat and is therefore not expected to shift upon

addition of FapR. The ratio of protein-DNA was kept constant for all conditions. (B) Gel shift

assay where FapR*-DNA complex was titrated with increasing concentrations of malonyl-CoA

under same conditions as above. (C) The fraction unbound DNA under different concentrations

of malonyl-CoA was determined by calculating the ratio of intensity of the shifted protein-DNA

band to the unshifted control band. All values were normalized to 1. N=3 representative trials.

Figure 4. **Increased membrane fluidity can restore viability to PG-limited cells.** (A) A

representative zone of growth obtained following application of 10% BnoH to a plate inoculated

with the indicated strains. *sigI ponA* and *sigW sigI ponA* have a higher propensity to form

suppressor mutation on LB medium in the absence of high concentrations of magnesium

compared to that of *rasP ponA* as seen by the isolated colonies on the plate. (B) Treatment with

10% BnOH creates a zone of growth when applied to *sigI ponA* cells grown on LB medium in

the absence of high concentrations of magnesium but fails to rescue *rasP ponA*. A one-way

ANOVA with Tukey test for multiple comparisons was performed. P-value > 0.05. The size of

the filter disc (6mm) was subtracted from the zone of growth values.

Figure 5. *fapR** **cells exhibit increased sensitivity to conditions that increase membrane**

fluidity. (A) FapR* cells demonstrate an increased sensitivity to Triton X-100. P-value cutoff

(*) < 0.05. The size of the filter disc (6mm) was subtracted from all zone of inhibition values.

(B) The *fapR** cells sensitized to heat with a *sigI* null deletion experience increased sensitivity to

heat (56 °C). (C) The *fapR** mediated rescue is lost in the *ecsA ponA*, *rasP ponA*, and *sigI ponA*

at increased temperatures.

Figure 6. *fapR** **sensitizes cells to the FAS-inhibiting effects of cerulenin**. (A) *fapR** exhibits

an increased sensitivity to cerulenin compared to WT and *fapR* null cells, but not to triclosan. No

comparison was performed between antibiotic groups. A one-way ANOVA with Tukey test for

multiple comparisons was performed for each antibiotic group. P-value cutoff (***) < 0.0002.

The size of the filter disc (6mm) was subtracted from the zone of inhibition values.

Figure 7. **Chemical inhibition of FAS by cerulenin is sufficient to rescue the *rasP ponA***

strain. (A) Increasing concentrations of cerulenin rescue growth of a *rasP ponA* strain grown in

LB medium in the absence of high concentrations of magnesium.

Figure 1. **FAS suppressor mutations rescue cells with lethal PG synthesis defects.** (A) The

FapR* 24-nucleotide deletion occurs in the thioesterase-like, malonyl-CoA binding domain of

the protein. Asterisks indicate residues important for coordinating the interaction between FapR

and malonyl-CoA. (B) *fapR** rescues viability of *ecsA ponA*, *rasP ponA*, and *sigI ponA* cells in

LB medium in the absence of magnesium supplementation. (C) Deletion of FapR does not rescue

viability of a *sigI ponA* mutant grown in LB medium in the absence of magnesium

supplementation. (D) *fapR** is dominant in a merodiploid strain expressing at locus *fapR*.

Figure 2. ***fapR* decreases the expression of FAS genes.*** (A) Gene expression from candidate

genes in the *fapR* regulon (*fabF, fapR and fabHA)* is decreased in a *fapR** mutant. The x-axis

indicates candidate genes detected, and the legend indicates strain genotypes. ND indicates a

lack of *fapR* gene expression in a *fapR* null background. Gene expression values ($2^{-\Delta ct}$) were

normalized to *gyrA* and plotted. A one-way ANOVA with a Tukey test for multiple comparisons

was performed. P-value cutoff (*) < 0.05.

Figure 3. **FapR* exhibits altered interactions with malonyl-CoA.** (A) Gel shift assay where a

FapR-DNA complex was titrated with increasing concentrations of malonyl-CoA. The lower

DNA bands serve both as a loading control and control for protein interaction specificity as it

does not contain the FapR recognized inverted repeat and is therefore not expected to shift upon

addition of FapR. The ratio of protein-DNA was kept constant for all conditions. (B) Gel shift

assay where FapR*-DNA complex was titrated with increasing concentrations of malonyl-CoA

under same conditions as above. (C) The fraction unbound DNA under different concentrations

of malonyl-CoA was determined by calculating the ratio of intensity of the shifted protein-DNA

band to the unshifted control band. All values were normalized to 1. N=3 representative trials.

Figure 4. **Increased membrane fluidity can restore viability to PG-limited cells.** (A) A

representative zone of growth obtained following application of 10% BnoH to a plate inoculated

with the indicated strains. *sigI ponA* and *sigW sigI ponA* have a higher propensity to form

suppressor mutation on LB medium in the absence of high concentrations of magnesium

compared to that of *rasP ponA* as seen by the isolated colonies on the plate. (B) Treatment with

10% BnOH creates a zone of growth when applied to *sigI ponA* cells grown on LB medium in

the absence of high concentrations of magnesium but fails to rescue *rasP ponA*. A one-way

ANOVA with Tukey test for multiple comparisons was performed. P-value > 0.05. The size of

the filter disc (6mm) was subtracted from the zone of growth values.

Figure 5. ***fapR* cells exhibit increased sensitivity to conditions that increase membrane***

fluidity. (A) FapR* cells demonstrate an increased sensitivity to Triton X-100. P-value cutoff

(*) < 0.05. The size of the filter disc (6mm) was subtracted from all zone of inhibition values.

(B) The *fapR** cells sensitized to heat with a *sigI* null deletion experience increased sensitivity to

heat (56 °C). (C) The *fapR** mediated rescue is lost in the *ecsA ponA*, *rasP ponA*, and *sigI ponA*

at increased temperatures.

Figure 6. ***fapR* sensitizes cells to the FAS-inhibiting effects of cerulenin***. (A) *fapR** exhibits
an increased sensitivity to cerulenin compared to WT and *fapR* null cells, but not to triclosan. No
comparison was performed between antibiotic groups. A one-way ANOVA with Tukey test for
multiple comparisons was performed for each antibiotic group. P-value cutoff (***) < 0.0002.
The size of the filter disc (6mm) was subtracted from the zone of inhibition values.

74   Figure 7. **Chemical inhibition of FAS by cerulenin is sufficient to rescue the *rasP ponA***

75   **strain.** (A) Increasing concentrations of cerulenin rescue growth of a *rasP ponA* strain grown in

76   LB medium in the absence of high concentrations of magnesium.