**Elucidation of Amyloid-Beta's Gambit in Oligomerization:**

**Truncated Aβ fragments of residues Aβ1-23, Aβ1-24 and Aβ1-25 rapidly seed to form SDS-stable, LMW Aβ oligomers that impair synaptic plasticity.**

**Beatriz Gil[1+], Jamie Rose[2+], Davide Demurtas[3], Gian-Filippo Mancini[4], Jessica Sordet-Dessimoz[4], Vincenzo Sorrentino[5], Nikita Rudinskiy[6], Matthew P. Frosch[7], Bradley T. Hyman[6], Marc Moniatte[8], Tara. L. Spires-Jones[2], Caroline E. Herron[1] and Adrien W. Schmid[8*].**

[1]Conway Institute of Biomolecular and Biomedical Research, University College of Dublin (UCD), Ireland.

[2]UK Dementia Research Institute and Centre for Discovery Brain Sciences at The University of Edinburgh, Edinburgh, Scotland.

[3]Interdisciplinary Centre for Electron Microscopy (CIME), EPFL, Switzerland.

[4]Histology Core Facility, Ecole Polytechnique Fédérale de Lausanne (EPFL), Switzerland.

[5] Healthy Longevity Translational Research Programme, Yong Loo Lin School of Medicine & Department of Biochemistry, National University of Singapore.

[6]Harvard Medical School, Mass General Hospital, Massachusetts Boston, USA

[7]Massachusetts General Hospital and Harvard Medical School, Massachusetts General Institute for Neurodegenerative Disease, Charlestown, MA, USA.

[8]Proteomics Core Facility, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland.

[+] Contributed equally

*To whom correspondence should be addressed:

Adrien W. Schmid, PhD.
Life Sciences (SV), PTP-PCF, Station 15, EPFL
1015 Lausanne, Switzerland
Tel: +41 (0)21 693 06 51
E-mail: adrien.schmid@epfl.ch

**Keywords:** Alzheimer's disease; amyloid-beta, oligomers, antibody, immuno-therapy, aducanumab, LTP, synaptic plasticity, diagnostic.

**Author contributions:** AWS conceived the project, designed and directed the study. CEH and BG planned and carried out LTP studies. AWS performed mass spectrometry and biochemical analysis. JR and TS planned and carried out IF and array tomography studies on human brain tissue. GM and JSD carried out IF studies on Tg mouse brain. DD performed electron microscopy. VS carried out *C. elegans* experiments and analysis. MPF performed histological examination of human brain tissue. NR and AWS carried out ELISA measurements. BTH and MPF provided biological samples. AWS and CEH wrote the paper and all authors read and commented on the manuscript.

**Acknowledgments:** This work was funded by the "Enable Grant", kindly awarded by Dr. André Catana from the Technology Transfer Office (TTO) as well as the Catalyze4Life Innovation grant by Prof. Bart Deplancke and Dr. Kostas Kaloulis of the Ecole Polytechnique Fédérale de Lausanne (EPFL). We thank Dr. Adam Swetloff (TTO) for the constructive and strategic discussions on the project development as well as the support by the whole Life Sciences Faculty. The project was partially funded using internal funding of the EPFL. Human brain sample preparation and ELISA development was supported by the ADRC grant P30AG062421.We are grateful to the Roland Bailly Foundation (Geneva, Switzerland) for the funding of the MS instrument. We extend our gratitude to Dr. David Hacker (Protein Production and Structure Core Facility (EPFL) for his help with antibody production. Prof. Carmen Sandi, Laboratory of Behavioural Genetics (EPFL) for kindly providing us with antibodies for the study as well as Dr. Jose Sanchez-Mut, Laboratory of Neuroepigenetics (EPFL, Prof. J. Gräff lab), providing us with the reversed sequence of Aβ42-1 peptide and Prof. Hilal Lashuel (EPFL) for the Aβ40-arc mutant peptide. We thank Dr. Pamela Valdés (LEN, EPFL) for her support with IF imaging. TSJ and JR are funded by the UK Dementia Research Institute which receives its funding from DRI Ltd, funded by the UK Medical Research Council, Alzheimer's Society, and Alzheimer's Research UK, and the confocal used for IHC/AT was funded by Alzheimer's Research UK (ARUK-EG2016A-6) and a Wellcome Trust Institute Strategic Support Gant.

**Conflict of interest:** EPFL holds a patent on specific amyloid-beta peptide fragment signatures
and the use thereof (WO2015150322).

**Abstract:**

In Alzheimer's disease (AD), Amyloid-beta (Aβ) oligomers are considered an appealing
therapeutic- and diagnostic target. However, to date, the molecular mechanisms associated
with the pathological accumulation or structure of Aβ oligomers remains an enigma to the
scientific community. Here we demonstrate the strong seeding properties of unique Aβ
fragment signatures and show that the truncated Aβ peptides of residues Aβ1-23, Aβ1-24 and
Aβ1-25, rapidly seed to form small, SDS-PAGE stable assemblies of ~5kDa to ~14kDa
molecular mass range. Mass spectrometry analysis of SDS-PAGE fractionated and gel
extracted oligomers revealed that the truncated Aβ isoforms of residues 1-23 to 1-25 form
stable entities with low molecular weight (LMW) oligomers, which strongly resemble the
regularly reported Aβ entities of putative dimeric or trimeric assemblies found in human post-
mortem AD and Tg mouse brain extracts. Furthermore, electrophysiological recordings in the
mouse hippocampus indicate that LMW Aβ assemblies formed by fragments Aβ1-23 to
Aβ1-25 significantly impair long-term-potentiation (LTP) in the absence of full-length
Aβ1-42. Extensive antibody screening highlights the important observation, that the LMW Aβ
assemblies formed by these truncated Aβ peptides escape immuno-detection using
conventional, conformation specific antibodies but, more importantly, the clinical antibody
aducanumab. Our novel findings suggest that there are new Aβ target "loopholes" which can
be exploited for the development of therapeutic antibodies with binding properties against
stable target hotspots present in Aβ oligomers. We provide here a first example of a new class
of monoclonal antibody with unique binding properties against LMW Aβ oligomers, in the
absence of binding to large fibrillar Aβ assemblies, or dense amyloid plaques. Our research
supports a novel, unparalleled approach for targeting early, pathological Aβ species during the
insidious phase of AD and prior to the appearance of large oligomeric or protofibrilar
assemblies.

**Introduction:**

There is mounting evidence suggesting, that early therapeutic intervention will be the most

promising therapeutic strategy in Alzheimer's disease (AD) (1). As a consequence significant

effort has been made in the past to gain insight into early, "molecular triggers" associated with

pathological amyloid-beta (Aβ) aggregation, because Aβ oligomers are known to play a

causative role in early synaptic dysfunction  and cognitive decline in patients suffering from

dementia (2). Therefore, Aβ oligomers are widely accepted as an appealing diagnostic and

therapeutic target (3) in Alzheimer's disease (AD). However, structural characterization of Aβ

oligomers remains a challenging endeavour (4), which is mainly due to the labile and

polymorphic nature found with these oligomeric entities. Consequently, it may not seem to be

surprising, that the lack in clinical efficacy reported with earlier developed immuno-therapies

collectively reflect the intrinsic difficulties observed with the characterization of a "unique

pathological target" associated with neurotoxic Aβ oligomers.

The aetiology and neurotoxic properties of different, low molecular weight (LMW) oligomeric

Aβ species (4–9) such as Aβ dimers and trimers (10) or, larger oligomeric assemblies (11),

have been subjected to thorough studies in the past, in order to identify the "holy grail" of Aβ

assemblies as novel therapeutic targets in AD (12). Numerous studies have narrowed down the

search of pathological Aβ species to LMW species, namely dimers (10,13,14) or trimers (11).

More recently, McDonald and colleagues (15) reported that human AD brain extracts contain

small Aβ structures of a 6–7 kDa molecular mass range, of which these Aβ species may form

part of larger, neurotoxic Aβ assemblies. However, earlier attempts to identify specific Aβ

isoforms associated with LWM Aβ assemblies and, more specifically, the ~4kDa and ~7kDa

species, have failed (16).

Aβ peptide mediated neurotoxicity has been the subject of a host of *in vitro* and *in vivo*

electrophysiological studies, indicating that acute exposure of both, Aβ42 (17) and the more

soluble Aβ40 (18,19) or, Aβ dimers isolated from human AD brains and CSF (14), significantly

impair long-term-potentiation (LTP) in the rodent hippocampus. Earlier studies showed that

intra-cerebral injection of Aβ40 in mice resulted in a rapid generation of truncated Aβ species

(20), which resemble Aβ fragment fingerprints typically observed during *in vitro* Aβ

oligomerization studies (12). The significance of truncated Aβ species is further supported  by

Mazzitelli and colleagues (21), who reported that the co-occurrence of the truncated fragment

Aβ1-24 promotes Aβ42 aggregation and impairs Aβ peptide clearance in the mouse brain. In

strong support with this concept, we provided strong analytical evidence on an aggregation

induced auto-cleavage and hence accumulation of a specific Aβ fragment fingerprint, which

can rapidly seed to form oligomeric Aβ assemblies (12). Taken together, these findings suggest

that a gradual accumulation of different truncated Aβ isoforms, of which some bear C-terminal

amidation ($-CONH_2$), could play a central role in the early seeding events of Aβ

oligomerization. These truncated Aβ species may therefore represent novel diagnostic markers

as well as provide unprecedented therapeutic potential for the development of new antibody

based AD therapies.

Using advanced technologies in quantitative and targeted mass spectrometry, together with our

earlier developed neo-epitope antibody tools (12) against specific C-terminal truncated Aβ

species, we were able to shed light on an enigma associated with LMW Aβ oligomers. Here we

provide analytical proof of detection of specific truncated Aβ species present in LMW Aβ

oligomers. This finding allows unprecedented and selective targeting of oligomeric assemblies,

irrespective of oligomer structure or size. This gain in knowledge has led us to develop a novel,

conformation selective mouse monoclonal antibody with unique binding properties against

small oligomeric Aβ entities formed by the truncated Aβ isoforms Aβ1-23, Aβ1-24 and Aβ1-

25, which appear during the early phase of Aβ peptide aggregation (12). In line with our

antibody target design rational for early, LMW Aβ assemblies, we show that this novel antibody

tool displays unique binding properties against LMW Aβ species, in the absence of binding to

full-length Aβ42 monomers or higher order Aβ assemblies, such as amyloid plaques. We

highlight here the strong need for identifying additional, molecular mechanisms associated with

the early events of Aβ peptide aggregation during the long, insidious phase of AD. This in turn

will allow the improvement of antibody-specific target engagement of early, pathological Aβ

species present in the human AD brain and biofluids.

***Results:***

***Low molecular weight oligomers consist of an assembly of different C-terminally truncated***

***Aβ isoforms.***

Numerous earlier reports show that oligomerization promotes the formation of labile Aβ

assemblies that decompose into SDS-PAGE stable Aβ entities of dimeric, trimeric and

tetrameric nature (12,15). Here, we have used a combination of analytical tools to investigate

the structural composition of these typical LMW Aβ assemblies observed during SDS-PAGE

analysis. Preparation of Aβ42 oligomers was carried out following a standardized and widely

applied protocol for the generation of amyloid derived diffusible ligands (ADDL) (22). In an

initial approach, Aβ oligomers were analysed by SDS-PAGE followed by immuno-blotting,

using the conventional antibodies 4G8 and 6E10, together with a selection of our earlier

developed, highly selective neo-epitope antibody tools (Table I & Supplementary Data Fig.

1a&b.) against different C-terminally truncated Aβ species (12).

| No. | Antibody | Target / Epitope | Host | Type | Applications tested | Source |
|---|---|---|---|---|---|---|
| 1 | N3 | Aβ C-term Asp23 | rabbit | polyclonal | IP, WB, DB | EPFL |
| 2 | N4 | Aβ C-term Val24-NH2 | rabbit | poylclonal | IP, WB, DB | EPFL |
| 3 | N5s | Aβ C-term Gly25-OH | rabbit | polyclonal | IP, WB, DB | EPFL |
| 4 | N5NH | Aβ C-term Gly25-NH2 | rabbit | polyclonal | IP, WB, DB | EPFL |
| 5 | N5ns | Aβ C-term 23-25 | rabbit | polyclonal | IP, WB, DB, ELISA, IHC | EPFL |
| 6 | JD1 | Aβ oligomer | mouse | monoclonal IgG1 | IP, DB, ELISA | EPFL |
| 7 | JD1 | Aβ oligomer | mouse | monoclonal IgM | DB, ELISA | EPFL |
| 8 | 6E10 | Aβ 3-8 | mouse | monoclonal | IP, WB, DB, ELISA, IHC | Covance |
| 9 | 4G8 | Aβ 17-24 | mouse | monoclonal | IP, WB, DB | Covance |
| 10 | MM26-2 | Aβ C-term 42 | mouse | monoclonal | WB | Anaspec |
| 11 | A11 | oligomers | rabbit | polyclonal | DB | Thermo-Fisher |
| 12 | OC | Aβ aggregates | rabbit | polyclonal | DB, IHC | Merck |
| 13 | Aducanumab | Aβ oligomers / fibrils | recombinant | human IgG | DB | Creative Biolabs |
| 14 | Neurogranin | - | sheep | polyclonal | WB | R&D Systems |
| 15 | Somatostatin | - | rabbit | polyclonal | WB | Abbexa |
| 16 | Syn-1 | alpha-synuclein | mouse | monoclonal | WB | BD Tansduction Labs |
| 17 | PSD94 | - | mouse | monoclonal | WB | abcam |
| 18 | b-actin | C-term | mouse | monoclonal | WB | Santa Cruz Biotech. |
| 19 | homer 1 | - | rabbit | polyclonal | WB | Synaptic Systems |

190    **Table I.  Summary of the different antibodies used in the study.** Antibodies no. 1 to5 are referred to neo-epitope specific antibodies in the

191    study. IgM and IgG classes of the monoclonal antibody JD1 are shown, whereas the latter was mainly used in the study.

In line with earlier studies we show that, under denaturing SDS-PAGE conditions, large oligomeric assemblies dissociate into LMW species of putative dimeric, trimeric and tetrameric Aβ species (Fig. 1a). Probing the WB membrane with a combination of neo-epitope antibodies N-3, N-4 and N-5ns against Aβ fragments of residues 1-23 (mw: 2775Da), 1-24 (mw: 2875Da) and 1-25 (mw: 2932Da) respectively, allowed identification of C-terminally truncated Aβ species at the migration level corresponding to Aβ42 monomers, dimers, trimers and tetramers. Detection of these truncated Aβ fragments appeared to be significantly increased in Aβ42 tetramers when compared to trimers and dimers, whereas strong antibody signals were also observed at the Aβ42 monomer level (mw: 4512Da).

To confirm the identification of different Aβ species in LMW oligomers, we excised SDS-PAGE gel bands at the migration level of Aβ42 monomers to tetramers (Fig. 1b, inset: gel), followed by a repeated gel dehydration & rehydration process to allow diffusion of the gel trapped Aβ species into the solution. Following gel extraction, the solution containing different Aβ species was cleared and concentrated using C18 stage tips and subjected to MALDI-TOF/TOF MS analysis. MS analysis revealed a high abundance of fragments Aβ1-23 and Aβ1-25 together with full-length Aβ42 in the tetramer gel band (Fig. 1b; top spectrum). Similarly, the C-terminally amidated ($-NH_2$) forms of fragments Aβ1-24 and Aβ1-25 were also identified, however at significantly lower abundance. Overall, the MS signal intensities found for Aβ1-23 and Aβ1-25 were in line with the immuno-blot signal observed at the identical gel migration level of Aβ42 tetramers ($\leq$15kDa) (Fig. 1a, lanes: 5&8). We could also identify similar Aβ fragment signatures in the trimer gel band, however, at significantly lower ratios when compared to tetramers, which corresponds to the decreased immuno-blotting signals found at the corresponding migration level.

221
222
223
224

225 **Figure 1. Truncated Aβ species rapidly seed to form small oligomeric entities of 6-14kDa molecular weight range. a** left; SDS-PAGE
226 and immuno-blotting (4G8) analysis of Aβ42 ADDLs, showing the typical gel migration pattern of Aβ dimers, trimers and tetramers within
227 the molecular mass range of 6-15kDa. Samples were loaded at high (1ug) and low (100ng) concentrations. Centre IB; re-probed membrane
228 with a mix of neo-epitope antibodies (N3, N4 & N5ns) revealing the presence of C-terminal truncated fragments. (IB right: merged scan of
229 both antibodies at different wavelengths: green=800nm (4G8), yellow=680nm (N3, N4 & N5ns). Neo-epitope antibodies bind to specific
230 bands at the migration level of putative Aβ42 tetramers, trimers and dimers. Substantial neo-epitope antibody binding was also observed at
231 the Aβ42 monomer level, which indicates the presence of C-terminally truncated Aβ species present in this band. **b** MALDI-TOF/TOF analysis
232 of gel extracted (by diffusion) Aβ species found in tetramers (top) and trimers (bottom). Aβ1-25 was partially detected as a C-terminal
233 amidated form (inset grey arrow), whereas Aβ1-24 was predominantly detected as the C-terminal amidated form (Aβ1-24-NH$_2$). The strong
234 MS signal intensities detected for Aβ23 and Aβ25, as compared to Aβ42 are in line with the IB signals found at the migration level of <14kDa
235 (a). **c** WB and quantitative MS analysis of high molecular weight (HMW, ~200kDa) Aβ assemblies detected in the insoluble fraction. WB's
236 were probed using either a combination of 6E10&4G (1[st] blot, left) followed by re-probing with a mixture of neo-epitope antibodies (2[nd] blot
237 from left: N3, N4, N5ns & N5NH$_2$) to detect the presence of fragments Aβ 1-23 to 1-25 in these HMW oligomers. Gel bands were excised at
238 the migration level of HMW oligomers and were subjected to LysN digestion, giving rise to the mid region sequences (inset right) of either
239 16-27 (from full-length Aβ1-42 cleavage) or 16-25 (e.g. derived from Aβ1-25 cleavage product). LysN cleavage products were subjected to
240 PRM analysis and the calculated total peptide areas, as a sum of all MS$_2$ b-ion (b2 to b8) transitions (bottom, left), were used to estimate the
241 relative abundance (histogram, right) of the different Aβ isoforms found within HMW oligomers. (Lanes 1-8 are identical Aβ samples but
242 loaded at high (1, 3, 5) and low (2, 4, 6) concentrations, lane 8 = MWM). **d** WB analysis of Aβ40 (left) and Aβ42 (right) ADDLs. ADDLs
243 were individually IP'ed with neo-epitope antibodies against fragments Aβ23, Aβ24 and Aβ25. Formic acid induced dissociation of ADDLs
244 allowed a specific pull-down of these Aβ fragments (lanes: 2,4,6 and 11,13,15), whereas IP of native Aβ40 (lanes: 3,5,7) and Aβ42 (lanes:
245 12,14,16) ADDLs resulted in a pull-down of an oligomeric complex with substantial enrichment of a band centred at the migration level of
246 ≥6kDa, which corresponds to a trimeric form of these Aβ fragments as seen by the migration pattern of a mixture of *in vitro* aggregated Aβ23
247 to Aβ25 (Figure 1e). (lanes: 1 & 10 = native Aβ40 & 42 ADDLs before IP, lanes: 8/9 = MW). **d** TEM imaging of the seeding properties
248 observed with Aβ1-24, showing predominant spherical, oligomeric structures of 20-50nm diameter (arrows) and some clusters of larger
249 aggregates (scale bar = 100nm). below; LC-MS analysis of Aβ24 trimers (WB inset) as seen by the multiply (m/z=1232.4409 M+H[7]) charged
250 ion. Inset right; WB analysis of the gel migration behaviour of individually aggregated fragments of Aβ23, Aβ24 and Aβ25. All Aβ fragments
251 form substantial amounts of trimers centred at the migration level of >6kDa (white box: Aβ24) as confirmed by LC-MS analysis. A shift in
252 migration profile can be observed between dimeric and trimeric Aβ23 and Aβ24 assemblies, which accounts for the difference in molecular
253 mass (200-300Da) found for the Aβ24 species (see table 2 for comparison). **e** WB analysis of Aβ assemblies formed by the mixture of
254 fragments Aβ1-23, 1-24 and 1-25. The rapid seeding of these fragments results in the formation of a substantial population of dimers (≥5kDa)
255 and trimers (≥6kDa), with some tetramers (≤12kDa). Monomers (≤3kDa) are no longer observed and the dimer (≥5kDa) band is slightly
256 shifted when compared to the Aβ42 monomers at 4.5kDa, which accounts for the difference in molecular mass for the dimeric species. (lanes:
257 1/2 = Aβ42 and lanes: 3/4 = Aβ23, 24 & 25 mixture at high and low concentrations). **f** Summary of neo-epitope antibody properties and
258 application (left green bar) for targeted analysis of early fragment fingerprints as a result of Aβ aggregation. The design and development
259 rational for a new conformational specific antibody (JD1) (right green column) with binding specificity against LMW oligomers (early seeds)
260 derived from early Aβ fragment fingerprints of Aβ1-23 to Aβ1-25 is shown.

263 In addition to the above outlined diffusion extraction procedure, an identical copy of the gel

264 bands was subjected to in-gel digestion, using LysN protease followed by LC-MS analysis, to

265 further corroborate the findings by WB and MALDI-TOF MS, as well as to allow identification

266 of additional Aβ isoforms present within these SDS-PAGE-stable LMW oligomers. High-

267 resolution LC-MS analysis revealed the presence of a diversity of N-terminally, as well as C-

268 terminally modified Aβ species across the different gel migration levels (Supplementary Data

269 Fig. 1c-e). Generally, identification of fragments Aβ1-23, Aβ1-24 and Aβ1-25 could be

270 confirmed in the monomer, dimer, trimer and tetramer bands and these fragments were found

271 to be significantly enriched in the monomer and tetramer bands respectively, which is in line

272 with the WB and MS analysis shown above (Fig. 1a). The increased abundance of truncated Aβ

273 species detected at the migration level of 3-5kDa may result from a gel migration induced

274 dissociation of Aβ assemblies, giving rise to the monomeric and dimeric form of Aβ1-23, Aβ1-

275 24 and Aβ1-25 (Table II). Moreover, LC-MS analysis allowed for unbiased identification of

276 the C-terminally amidated fragment Aβ1-24 ($Val_{24}$). In addition to the above listed truncated

277 forms, we were also able to identify C-terminal amidation at positions $His_{13}$, $His_{14}$ and $Asp_{23}$.

278 Modifications associated with the far C-terminal consisted of a repertoire of specific truncations

279 found at position $Gly_{37}$, $Gly_{38}$, $Val_{39}$ as well as $Ile_{41}$ (Supplementary Data Fig. 1e).

| Aβ Peptide | C-term. | Abbreviation | monoisotopic mass (Da) | average mass (Da) | Dimer (Da) | Trimer (Da) | Tetramer (Da) |
|---|---|---|---|---|---|---|---|
| 1-23 | COOH | 23 | 2775,27 | 2776,96 | 5554 | 8331 | 11108 |
| 1-24-NH2 | CONH2 | 24N | 2873,35 | 2875,11 | 5750 | 8625 | 11500 |
| 1-25 | COOH | 25 | 2931,36 | 2933,15 | 5866 | 8799 | 11733 |
| 1-25-NH2 | CONH2 | 25N | 2930,37 | 2932,16 | 5864 | 8796 | 11729 |
| 1-40 | COOH | 40 | 4327,15 | 4329,86 | 8660 | 12990 | 17319 |
| 1-42 | COOH | 42 | 4511,27 | 4514,10 | 9028 | 13542 | 18056 |

284 **Table II. List of Aβ peptide fragments used in the study.** Theoretical masses of Aβ dimers, trimers and tetramers was calculated using the

285 average peptide mass.

289 **Truncated Aβ species may seed to form SDS-PAGE stable entities with large Aβ**

290 **oligomeric assemblies of a 50-200kDa molecular mass range.**

291 We reasoned that the detection of small Aβ assemblies at the migration level of 5-14kDa may

292 result from the dissociation of larger (≥30kDa), metastable Aβ entities. We therefore decided

293 to analyse the insoluble fraction after centrifugation of the ADDLs (soluble fraction)

294 preparation. This insoluble fraction is known to contain mainly large Aβ assemblies of

295 protofibrillar and fibrillar nature (data not shown). WB analysis of this fraction revealed the

296 presence of different, large Aβ assemblies at the migration level of 50kDa to 200kDa

297 (Supplementary Data Fig. 2a). Probing the membrane with antibodies directed against the N-

298 terminal (6E10) and mid-region (4G8) or, the C-terminal ending of Aβ42 (MM26-2) (Table I)

299 allowed the identification of the bulk of Aβ assemblies centred at the LMW migration level of

300 4-14kDa, as observed with the soluble ADDL fraction shown above. However, application of

301 a cocktail of neo-epitope antibodies (N3, N4, N5ns and N5NH2) also strongly indicated the

presence of distinct Aβ assemblies found at the gel migration level of ~200kDa (Fig. 1c) as well

as ~98kD, ≥62kDa and ≥50kDa (Supplementary Data Fig. 2a). We observed particularly strong

binding of 6E10 & 4G8 as well as N3-N5 (mix) to a large Aβ entity found at ~200kDa (Fig. 1c,

top: WB). We therefore decided to carry out in-gel digestion followed by quantitative MS

analysis of these large Aβ entities in order to confirm the presence of the different Aβ isoforms

of Aβ1-23 to Aβ1-25 within these stable assemblies. Our quantitative MS analysis confirmed

the high abundance of the different Aβ isoforms of 1-23, 1-24NH$_2$, 1-25 and 1-25NH$_2$ present

in these large Aβ assemblies (e.g. 200kDa) of which the overall calculated sum of all truncated

Aβ isoforms accounted ≥40%, when compared to full-length Aβ1-42 (Fig. 1c, bottom:

histogram). We also provide quantitative MS data on the different Aβ isoforms detected at the

gel migration level of ≥50kDa, ~98kDa and ~200kDa (Supplementary Data Fig. 2b).

**Dissociation of Aβ oligomers results in the isolation of Aβ species of 6kDa to 14kDa mass**

**range.**

Given the observation that truncated Aβ species form stable complexes with LMW and HMW

oligomers, we were interested in validating if we are able to pull-down Aβ oligomers formed

by Aβ40 or Aβ42 peptides. To address this question, we prepared ADDLs of which one part of

the preparation was subjected to 70% formic acid (FA) denaturing (3hrs at room temp.)

followed by IP using neo-epitope antibodies N-3s, N-4 or N-5ns (Table I & Supplementary

Data Fig. 1a&b). We show that selective IP of FA dissociated ADDLs resulted in a specific

capturing of two Aβ species centred at the gel migration level of 6kDa and <14kDa, whereas

IP of native (not denatured) ADDLs resulted in the pull-down of the typical oligomeric

assemblies found at a migration level of 6-14kDa (Fig. 1d). The selective pull-down and

therefore isolation of the ≥6kDa Aβ species was found to be more straightforward with Aβ40

assemblies (Fig. 1d, left; lanes 2-7), whereas Aβ42 ADDLs appeared to be more resistant to FA

dissociation (Fig. 1d, right; lanes: 11-16).

**Aβ fragments of Aβ1-23, Aβ1-24 and Aβ1-25 rapidly seed to form SDS-stable, LMW assemblies in the absence of full-length Aβ1-42.**

Because our earlier studies strongly indicated that Aβ aggregation is accompanied by the appearance of the typical fragment fingerprint consisting of truncations at residues 1-23, 1-24 and 1-25, we were interested in understanding the seeding properties using a mixture of these C-terminally truncated Aβ species. We show that, *in vitro* aggregation of these truncated Aβ fragments results in the rapid formation of stable assemblies within the 5-15kDa molecular range, of which dimers (≥5kDa) and trimers (≥6kDa) are the most predominant species formed by these Aβ fragments, whereas monomers (theoretical av. mass range: 2776Da to 2933kDa) were no longer detected (Fig. 1e, lanes: 3-4). Consistent with this observation, earlier reports showed a rapid seeding behaviour with fragment Aβ1-28, revealing gel migration profiles that are consistent with stable Aβ1-28 dimers (22). The gel migration profile of the peptide mixture of Aβ1-23 to Aβ1-25 dimers (>5kDa) was significantly shifted when compared to full-length Aβ1-42 monomers (4.5kDa) (Fig. 1e, lanes 1-2), whereas the trimeric form of these truncated Aβ species (theoretical mass: 8-9kDa) was found at a migration level above ≥6kDa, which strongly resembles the migration profile of earlier reported putative Aβ42 dimers (10) or 4-7kDa assemblies (15) isolated from human AD brain extracts.

We then investigated the self-seeding properties of individual truncated Aβ species Aβ1-23, Aβ1-24-NH$_2$ and Aβ1-25 as well as the C-term amidated form of Aβ1-25 (NH$_2$). In vitro aggregation of Aβ1-24 showed that this fragment has rapid self-seeding properties, resulting in the formation of SDS-PAGE stable assemblies in the MW range of dimers to trimers (Fig. 1f, WB inset). Transmission electron-microscopy (TEM) imaging revealed that fragment Aβ1-24 preferentially seeds to form homogenous spherical aggregates with some clusters of small protofibrillar structures. The gel migration profile of Aβ1-24 and Aβ1-25 peptides were similar (Fig. 1f), whereas the gel migration profile appeared to be slightly shifted when compared to fragment Aβ1-23, which may account for the difference in MW between these Aβ species (Aβ23 vs. Aβ24 dimer = 198Da, Aβ24 dimer vs. Aβ25 dimer = 114Da) (Table II).

The true molecular entity of these small oligomeric assemblies observed by SDS-PAGE was further confirmed using high-resolution LC-MS analysis, revealing the presence of highly stable Aβ1-24 trimers (8.6kDa) (Fig. 1f, bottom: MS spectrum) as well as dimers (5.7kDa) or even pentamers (14.3kDa) (Supplementary Data Fig. 2c-d). The MS analysis of stable dimers to pentamers of these truncated Aβ peptides further corroborates the migration pattern observed by denaturing SDS-PAGE analysis (Fig.1f, WB inset), and therefore rules out any analytical artefacts associated with gel migration (23).

***Detection of early, pathophysiological Aβ seeds requires the development of novel antibody***
***tools, with high binding selectivity for small oligomeric assemblies, in the absence of***
***monomer binding.***

The above reported findings clearly highlight the need for the development of additional

antibody molecules with high binding specificity for LMW Aβ oligomers, irrespective of the

presence of the full-length Aβ42 or Aβ40 target. To address this limitation, we set out to develop

a conformational specific antibody; with preferential binding properties against a repertoire of

LMW Aβ assemblies formed by here described truncated Aβ species. We speculated that the

application of such a conformation specific antibody would serve as an additional analytical

tool as compared to our earlier developed neo-epitope antibodies, therefore allowing the

detection of the gradual appearance of small Aβ entities, prior to the accumulation of large,

metastable oligomeric or protofibrillar species (Fig. 1g).

We therefore immunized mice using a synthetic, oligomeric form of fragment Aβ1-24 and

screened clones for antibodies with binding properties against a linear (denatured), and

aggregated (positive target) form of the different C-terminally truncated Aβ fragments of

interest as well as against denatured, full-length Aβ42 and Aβ40 peptide (negative target).

Using a combination of ELISA and DB techniques, we were able to identify six antibody-

secreting clones (data not provided), with different antibody binding properties against

oligomeric Aβ assemblies of interest, of which the properties of one selected monoclonal

antibody (mAB), referred to as **"*JD1*"** is described below.

**The monoclonal antibody JD1 binds LMW Aβ assemblies, which appear during the early**

**time course of Aβ aggregation.**

To characterize the binding specificity of JD1, we initially compared JD1 binding properties

with the two conventional antibodies 4G8 and 6E10, as well as different, conformation specific

antibodies such as the A11 or OC antibody. We found that the intrinsic binding property of

JD1 is not limited to fragment Aβ1-24, since JD1 binding was also observed for the different

aggregated forms of fragments of Aβ1-23 and Aβ1-25-NH$_2$ (Fig. 2a; bottom DB). Immuno-

blotting analysis indicated that JD1 positive signals found by dot blotting are in agreement with

the presence of LMW oligomeric assemblies found at the Aβ peptide migration range of dimers

and trimers (5-14kDa) and this could be also confirmed using aggregated Aβ42 (Fig. 2a, top

WB, lane: 6). While JD1 clearly binds to oligomeric Aβ assemblies using native DB, we do not

observe positive JD1 binding of these LMW species by WB. The reason for this observation

may result from either dissociation of larger ($\geq$ 14kDa) Aβ assemblies or from a SDS-PAGE

induced denaturing of the conformational epitope.

DB analysis of Aβ aggregation kinetics indicated that significant JD1 binding can be observed

following $\geq$8hrs aggregation of the full-length, wild type (wt) Aβ1-42 peptide (Fig. 2b).

Remarkably, aggregation kinetics analysis of the arctic mutant Aβ40 revealed that, JD1 binding

significantly preceded the detection of oligomeric assemblies formed by Aβ40-Arc peptide,

when compared to wt Aβ42 peptide. TEM imaging indicated that selective JD1 binding was

accompanied by the presence of predominantly oligomeric entities found with Aβ40-Arc (Fig.

2b), which is in line with the morphology observed with fragment Aβ1-24 (Fig. 1f). Probing

Aβ42 peptide aggregation with neo-epitope antibodies revealed that the rise in JD1 positive

species is generally associated with a gradual accumulation of C-terminally truncated Aβ

species of residues 1-23 to 1-25 and this was further confirmed by MALDI-TOF MS (Fig. 2c,

bottom: MS spectrum), whereas no antibody binding was observed for the more soluble, wt

Aβ40 peptide during the same time course of peptide aggregation (Fig. 2c; top DB).

**Figure 2. LMW Aβ oligomers escape immuno-detection using conventional, conformation specific antibodies. a** WB (4G8 & 6E10)
and DB (JD1) analysis of the migration profile of JD1 positive species formed by aggregated Aβ1-23 to Aβ1-25 (lanes: 2-4) and full-length
Aβ42 (lane: 6). The presence of Aβ42 (lane 6) species at the migration level of dimers and trimers is revealed by WB analysis. Below; DB
validation of JD1 positive species. JD1 positive signals found for Aβ42 were associated with the presence of bands detected at ≥6kDa and
≤14kDa (lane: 5). **b** Aggregation kinetic analysis of the mutant Aβ40-Arc versus wt Aβ42 peptide using JD1 (DB) and TEM imaging of Aβ40-
Arc oligomers at 8hrs. Aβ entities of spherical (10-50nm) and small protofibrillar morphology can be observed with Aβ40-Arc (scale bar =
100nm). **c** Monitoring of Aβ42 and Aβ40 aggregation kinetics using JD1 and MS. top: DB validation using JD1 (left) and re-probed with a
mixture neo-epitope antibodies (right): positive JD1 signals are observed following 8hrs of Aβ42 aggregation (DB) and this was accompanied
by an appearance of fragments Aβ23, Aβ24 and Aβ25 as probed with N3, N4 and N5ns (right). Bottom: MS (MALDI-TOF-TOF) analysis of
Aβ fragment signatures detected at t=8hrs. Inset right; WB analysis of IP'ed (JD1) Aβ42 (t=8hrs) (lane 2) showing the pull-down of a
predominant band centred at the migration level of ≥6kDa. (lane 1: Aβ42 starting material). **d** DB validation of Aβ40 and Aβ42 as well as
truncated Aβ1-24 aggregation kinetics using JD1 (Lanes: 1-3), the oligomer specific antibody A11 (lanes: 4-6), conformation specific antibody
OC (lanes: 7-9) and Aducanumab (lanes: 10-12) (samples were spotted at two different concentrations: lanes 1-2, 4-5, 7-8,10-11 = 50ng and
lane 3, 6, 9, 12 = 100ng). A clear difference in antibody binding to aggregated Aβ24 can be seen between JD1 and A11 as well as Aducanumab.
**e** Validation of JD1, Aducanumab as well as human plasma, natural IgGs or 6E10 binding specificity against freshly prepared (lanes: 1-4) or
aggregated (lanes: 5-8) forms of Aβ fragments Aβ23, Aβ24 and Aβ25 as well as Aβ42 (lanes: 9-10). JD1 has strong binding affinity for the
aggregated form of the different Aβ fragments; whereas no binding was observed with Aducanumab or plasma IgGs (lanes. 5-8) against the
soluble or aggregated form of these C-term truncated Aβ species.

## Validation of JD1 binding properties against oligomeric Aβ species and its potential applications for screening methods.

We show that detection of JD1 positive species was accompanied by the gradual accumulation of a typical Aβ fragment fingerprint (Aβ23 to Aβ25) as shown by the application of neo-epitope antibodies (Supplementary Data Fig. 3a) and JD1 proved to be useful for screening aggregation inhibitors, such as the green tea extract (EGCG) (Supplementary Data Fig. 3b). In line with the DB screening we show that, IP-MS analysis with JD1 does not result in the pull-down of soluble, Aβ42 monomers at early time points (t ≤ 30min), when compared to prolonged aggregation times (t = ≥2hrs) (Supplementary Data Fig. 3c). We also took advantage of the C. elegans GMC101 worm model to further understand the binding property of JD1 using a living organism. GMC101 worms constantly express the human Aβ42 isoform in muscle cells and develop age-progressive paralysis and significant amyloid deposition which are A11 positive species (24). WB analysis of aged GMC101 worms revealed a significant accumulation of different LMW Aβ species at the molecular weight range of 5-14kDa. IP-MS analysis of GMC101 extracts confirmed the abundance of different C-terminal truncated Aβ species (Supplementary Data Fig. 3d), which reflects the typical Aβ fragment signatures found in Tg2576 mice and human post-mortem brain tissue extracts (25). Furthermore, we show that JD1 also binds oligomeric assemblies formed by synthetic mouse Aβ42 sequence, though, with significantly lower affinity (Supplementary Data Fig. 3e), whereas, no binding was observed

against the reversed human Aβ42-1 sequence (Supplementary Data Fig. 3f) or aggregated

alpha-synuclein protein (data not shown). These observations collectively rule out JD1's

binding property against oligomeric assemblies formed by other amyloidogenic proteins, as

observed with earlier developed conformation specific antibodies (26). Collectively these

observations highlight JD1's unique binding property for predominantly, small oligomeric Aβ

assemblies, in the absence of binding of full-length Aβ1-42 or Aβ1-40 monomers.

**Small Aβ assemblies formed by the N-terminal fragments Aβ1-23 to Aβ1-25 are not**

**detected by conventional, anti-oligomeric antibodies or natural plasma IgG's.**

Due to the strong seeding properties observed with C-terminally truncated Aβ species, we

investigated if these LMW oligomeric assemblies can be detected using conventional, anti-

oligomeric antibody tools. For this purpose, full-length Aβ42 and Aβ40, as well as the C-

terminally amidated fragment Aβ1-24-NH2 were subjected to aggregation kinetics and probed

with the "anti-oligomer" specific antibody A11 and the "conformation-specific" antibody OC

(6,26,27), as well as a biosimilar of the clinical antibody aducanumab (28,29). No visible

antibody binding was observed at day 0 (data not shown), which is in line with earlier reported

antibody screening studies using A11 and OC (30). Following one day of Aβ peptide

aggregation, significant antibody binding was only observed against peptides Aβ42 and Aβ40

using the OC and aducanumab antibodies (Fig. 2d), whereas A11 reached a similar level of

binding significance following two days of aggregation. Generally, positive immuno-probing

signals for Aβ42 preceded the detection of Aβ40 oligomers, which is likely due to the increased

aggregation propensity found with Aβ42. At day two, JD1 binding to aggregated Aβ42 and

Aβ40 was similar when compared to aducanumab, but with a noticeable lag in detection of

Aβ40 at day 1. Intriguingly, our screening studies revealed a striking difference in antibody

binding affinity for the aggregated Aβ1-24 peptide. Detection of oligomeric Aβ1-24 assemblies

was significantly reduced using the OC or aducanumab antibody and absent with A11 (Fig. 2d

& Supplementary Data Fig. 3g). To determine if the intrinsic binding properties of JD1 are

associated with an oligomeric or linear, denatured form of Aβ peptide, we prepared fresh Aβ

peptide standards by dissolving and sonicating the Aβ peptides in HFIP solution, followed by

solution evaporation and solubilization in a 50mM NaOH solution as described earlier (30).

Sample preparation of freshly prepared (t=0) or aggregated (t=1d) Aβ peptide standards where

screened by DB using JD1 and aducanumab, together with a cocktail of neo-epitope antibodies

(N3, N4, N5ns, N5N) as well as 6E10 for the detection of all Aβ assemblies. Our findings

strongly indicate that JD1 has significantly increased binding properties for the aggregated form

of fragments Aβ1-23, Aβ1-24-NH$_2$, Aβ1-25, Aβ1-25NH$_2$ and that detection of these Aβ species

was significantly reduced with aducanumab (Fig. 2e, lanes: 5-8). Application of neo-epitope

antibodies enabled the detection of both, denatured as well as aggregated forms of the Aβ

peptide, indicating that these antibodies bear high C-terminal binding specificity, irrespective

of Aβ peptide structure, which is in agreement with our earlier reported validation studies (12).

Human plasma is known to contain natural IgGs antibodies that are reactive against a broad

range of different amyloidogenic species. Because natural antibodies have been reported to bear

neuroprotective properties against Aβ peptide (31–33), we were interested in understanding if

these plasma IgG's also bear binding properties against the here described LMW Aβ oligomers.

In-spite of the high, natural abundance of conformation-specific antibodies reported in human

plasma, we could not identify plasma IgG selective binding to the small oligomeric Aβ entities

of fragments Aβ1-23 to Aβ1-25. However, significant plasma IgG binding was observed for

aggregated, but not soluble Aβ42 peptide and levels of plasma IgG binding was comparable

with aducanumab (Fig. 2e, lane: 10). Overall, these observations highlight the important finding

that, the small oligomeric Aβ assemblies formed by the truncated fragments Aβ1-23 to Aβ1-

25, may escape conventional immuno-based detection, using the oligomer-specific antibody

A11 or aducanumab.

***Identification of JD1 positive assemblies in human post-mortem AD brains highlights the***

***pathological significance of LMW Aβ assemblies in AD.***

To further provide evidence of the significance of LMW oligomeric Aβ species in AD brains,

we decided to screen biological samples using a combination of immuno-fluorescence (IF) and

array tomography imaging as well as IP-MS and ELISA assays. Having confirmed the presence

of truncated Aβ fingerprints in a set of human brain samples earlier (12), we set out in

specifically identifying JD1 positive species in post-mortem human brains.

IF imaging of human AD brains revealed the presence of dense core plaques, which were

thioflavin S (ThioS) and OC positive, however, we did not observe any JD1 positive staining

of these dense amyloid plaques (Fig. 3a). Therefore, we were interested in identifying the

presence of JD1 positive species at synapses, because one of the early and important effects of

oligomeric Aβ in AD, is the binding of oligomeric Aβ to synapses, resulting in synaptic

dysfunction and gradual loss of synapses. Tissue from human subjects was prepared for high-

resolution ribbon array tomography as reported previously (34), allowing accurate and high-

resolution detection of individual synapses. Similar to our IF analysis, application of the OC

antibody resulted in strong labelling of dense core Aβ plaques as well as individual postsynaptic

punctae, which co-localized with PSD95 protein staining. No JD1 positive staining could be

observed for dense Aβ plaques, which is in line with the IF analysis of aged (≥9months) Tg2576

mouse brains (data not shown). Interestingly, we clearly observed selective JD1staining at

individual postsynaptic punctae, which co-localized with OC and PSD95 staining (Fig. 3b).

This observation indicates that JD1 positive Aβ assemblies may represent early, neurotoxic Aβ

species associated with synapse binding and therefore resulting in gradual neurodegeneration.

**Figure 3. JD1 positive oligomers are detected in the post-synaptic vicinity of AD brains. a** IF staining of amyloid plaques found in a human AD brain, using Thioflavin S (Thio S), JD1 and OC antibody. Significant plaque staining was observed with Thio S, which co-localized with OC staining, whereas no staining of dense core plaques was found using JD1. **b** Array tomography images of Aβ deposits detected in two post-mortem human AD brains using JD1, anti-PSD95 and OC antibody (left to right). JD1 positive species were detected in the post-synaptic vicinity, which co-localize with PSD95 positive staining. Large images are maximum intensity projections of 5 serial sections and insets are single sections confirming colocalization. (for brain tissue details see Supple. Table I). **c** Example of an IP-WB analysis of human cortical brain tissue extracts using the neo-epitope antibody N5ns. Significant pull-down of dimeric and trimeric Aβ species are observed in a majority of AD (lanes 6-14) but not control (lanes 2-4) brains. (lane 1: IgG ctrl, lane 15: MW, Lane 16: aggregated synthetic Aβ25 dimer). Below; Representative summary of an IP-MS analysis of an AD brain sample (lane 14) using a mixture of antibodies N3, N4 and N5ns. Blue bars highlight the sequences of all Aβ species identified (FDR ≤1%) (PEAKS Studio soft.) **d** ELISA analysis of JD1 positive assemblies found in post-mortem human AD (n=7) brains (FA extracts) but not in age matched controls (HC, n=3). (range: 35pg-132pg / mg brain tissue) (median ± SEM of technical triplicates).

**Targeted mass spectrometry analysis of human AD brain extracts reveals the presence of LMW Aβ assemblies, consistent with the detection of fragments Aβ1-23 to Aβ1-25.**

We decided to analyse human post-mortem brains to further support the results of our high-resolution array-tomography and provide insight into the pathophysiological significance of these truncated Aβ isoforms as well as the LMW oligomers. We employed IP-WB and IP-MS analysis of eight AD and three human control (HC) brain extracts (Supplementary Data Table I) and found that IP with N-5ns enabled the isolation of Aβ species centred at a gel migration level of 5-6kDa and ≥6kDa, which corresponds to our above observed dimeric and trimeric species of fragment Aβ1-25 (Fig. 1e). Significant levels of dimeric and trimeric Aβ species were detected in four AD brains (Fig. 3c, lanes: 7, 12 13 & 14), whereas the detection of these Aβ species was significantly lower in three other AD subjects (Fig. 3c, lanes: 8-10). No Aβ was detected in brains from human controls (Fig. 3c, lanes: 2-4) as well as in one AD subject (lane: 6). Generally, IP-WB analysis of these AD brain extracts strongly resembled the migration profile of the above reported FA dissociated ADDLs (Fig. 1d). Moreover, using IP-MS analysis with a neo-epitope antibody cocktail (N3, N4, N5ns), we were able to confirm the overall presence of different Aβ species in AD brain extracts but not human controls (Fig. 4c; bottom inset & Supplementary Data Fig.4a-e). Overall, our IP-MS analysis allowed the unbiased identification of different truncated Aβ species of residues 1-23, 1-24, 1-25, of which fragment 1-24 was predominantly detected as a C-terminal amidated (-CONH2) form. In addition, we were also able to identify the C-terminally amidated fragment of Aβ1-25 (Supplementary Data Fig. 4e) using a recently developed neo-epitope antibody (N-5NH2) (Table I).

Next, we carried out protein extractions from Tg2576 mouse (age: 9months) brain tissue as well as human post-mortem AD and HC brains, using either TBS, formic acid (FA) or SDS extraction buffers, followed by immuno-precipitation with JD1 and downstream targeted MS

analysis. IP-MS analysis of Tg2576 mice and human AD brains provided unbiased

confirmation of the presence of JD1 positive Aβ assemblies in these brain samples

(Supplementary Data Fig. 5a) and detection of these Aβ species was found to be increased using

FA and SDS extraction buffers, when compared to the more soluble TBS fractions, which is an

observation also reported by others (10). However, we cannot rule out that the use of harsh

extraction buffers, such as FA, could eventually result in substantial dissociation of small

oligomeric Aβ assemblies, therefore contributing to a decrease or even loss of the

conformational epitope associated with JD1 binding.

To further corroborate the presence of JD1 positive species in post-mortem human brain, we

next, applied a sandwich ELISA (Supplementary Data Fig. 5b), using JD1 as capture antibody

and a biotinylated form of 6E10 for detection of oligomeric Aβ species in HC (n=3) and AD

(n=7) cortical brain tissue extracts. Our ELISA measurement revealed that, JD1 positive

assemblies can be detected in all of the AD subjects, though, with considerable inter-subject

variability (30-140pg / mg cortical brain extract), which may reflect the general variability in

Aβ pathology found in AD subjects (Fig. 3d). Indeed, a large inter-subject variability of LMW

Aβ assemblies was also observed by IP-WB (Fig. 3c). No JD1 positive species were detected

in brain extracts from control subjects, however, the low number of subjects used here would

not rule out the presence of low amounts of oligomeric Aβ in other, age matched control

subjects or patients suffering from MCI (35).

Collectively, our brain imaging and screening studies strongly indicate that JD1 has preferential

binding properties against small Aβ assemblies and that antibody binding may be gradually lost

during the time course of on-pathway Aβ peptide aggregation. This may also explain the lack

in JD1 binding to dense core plaques as well as the decrease in JD1 binding affinity following

extended (≥2week) Aβ peptide aggregation (Supplementary Data Fig. 3b). Nevertheless, our

findings do not rule out binding of JD1to protofibrillar or fibrillar structures, given that the Aβ

binding epitope is sufficiently conserved and more importantly exposed, in these larger Aβ

assemblies.

***LMW oligomers formed by fragments Aβ1-23 to Aβ1-25 significantly impair synaptic***
***plasticity.***
Earlier reports showed that unlike full-length Aβ42, the truncated form of Aβ1-16 did not have
any inhibitory effects on synaptic plasticity (36), whereas other reports indicate that  the larger
Aβ1-25 isoform,  serves a sphingolipid-binding domain motif and is rapidly internalized by
neuronal cells (37). This rapid peptide internalization may account for the acute neurotoxicity
observed for some Aβ fragments. Therefore, to address the acute neurotoxic properties of
truncated Aβ species, we decided to generate LMW oligomers using a mixture of the typically
observed Aβ fragment fingerprints during aggregation. For this purpose, a mixture of the
synthetic Aβ fragments Aβ1-23, Aβ1-24-NH2, Aβ1-25 and Aβ1-25-NH2 (50ng/ul) was
incubated over-night followed by size exclusion chromatography (SEC) purification of LMW
oligomeric assemblies that are recognized by JD1.
We identified JD1 positive Aβ species in the elution volume (13ml) corresponding to the mass
of a theoretical molecular standard of ≥13.7kDa (Fig. 4a). WB analysis of the SEC fractions
revealed a population of different Aβ assemblies of molecular weights ranging from ≥5kDa
(dimers) to ≥ 6kDa (trimers), though we cannot rule out that these Aβ entities are the result of
the SDS-PAGE analysis induced dissociation of larger Aβ assemblies. MS analysis of the JD1
positive fraction confirmed the identification of a mixture of all four Aβ isoforms present in the
JD1 positive fraction (data not shown).
646

**Figure 4. LMW oligomeric assemblies derived from C-terminally truncated Aβ species significantly inhibit LTP in the mouse**
**hippocampus. a** left; representative DB screening of JD1 positive assemblies formed by the individual fragments Aβ23 to Aβ25 (lanes 2-5)
as well as using a mixture of all fragments (lane 6). A reverse sequence of Aβ25-1 was used to show that aggregation of this sequence does
not generate JD1 positive assemblies (lane 1). Right; SEC (Superdex 75, 10/300) and DB analysis of LMW assemblies formed by the mixture
of Aβ fragments 1-23 to 1-25. JD1 positive assemblies were detected in SEC fraction volume 13ml, which contained dimeric and trimeric
(≥6kDa) assemblies of these fragments (WB inset below), whereas monomers (≤3kDa) eluted in fraction volume 16ml. **b** Effect of application
of LMW Aβ oligomers (SEC purified JD1 positive assemblies) on LTP in hippocampal slices from young wild type mice. Acute application
of Aβ oligomers (500nM) to hippocampal slices significantly depressed LTP amplitude compared to control conditions (P<0.05). (inset right:
top) Sample traces represent fEPSP before (**1**) and 60min after HFS (**2**) in control LTP, and before (**3**) and after HFS (**4**) Aβ-mediated LTP.
(inset right: bottom) Bar chart comparing magnitude of LTP in control (190.6 ±22.94) and Aβ conditions (123.8 ±9.05) (N=5). All data are
presented as Mean ±SEM and analyzed using unpaired Student's t-test. JD1 positive Aβ oligomer treated sliced (open circles) as compared to
PBS controls (filled black circles). **c** Diagram of the analytical workflow applied for mouse hippocampal tissue extraction post-LTP
recordings, protein digestion, isobaric labelling (TMT), off-gel isoelectric focusing, MS analysis and bioinformatics as well as orthogonal hit
validation by targeted, quantitative MS and immuno-blotting. **d** Volcano plot analysis (y-axis of volcano plots:  -Log10 p-value from two-
sided T test corrected by Benjamini-Hochberg procedure) of differentially regulated proteins (x-axis: Log2 fold-change) in mouse brain tissue
(n=3), following treatment with Aβ oligomers versus PBS controls. The dashed lines indicate a threshold of ± Log2 0.58 fold change (vertical
lines) and a 0.05 p-value cut-off threshold (horizontal line). At the given cut-off, somatostatin (SMS) appears to be significantly up-regulated
(FC= Log2 0.63) in Aβ treated brains. A similar trend in FC was observed for neurogranin (NEUG), however at significantly reduced level
of significance (FC= log2 0.58). Proteins were graphed by fold change (FC, difference) and significance (-Log p) using a false discovery rate
of 0.05 and an S0 of 0.1. Protein ID's in red (SMS and NEUG) were identified as significantly up-regulated (ggVolcanoR tool), whereas only
SMS passed the confidence level of P<0.05 using Perseus software (Supplementary Data Figure 9). **e** Representative example of a regression
analysis of (intra-TMT channel) protein correlation (mouse #2) for control versus Aβ treated brain tissue. A strong proteome correlation was
observed for control versus Aβ treated tissues. As a reference, two protein markers with strong correlation between control and Aβ treated
tissies are highlighted (DLG4 (PSD95) and SYUA protein). A clear correlation deviation was observed for the two synaptic markers SMS
and NEUG.

To set out a strategic approach in assessing the acute neurotoxic properties of LMW oligomers,
we decided to incubate mouse hippocampal brain slices with the above described, SEC purified
JD1 positive assemblies, followed by *in vitro* electrophysiological recordings of long-term
potentiation (LTP) in the Schaffer collateral commissural pathway of the CA1 region in mouse
hippocampal slices. Brain slices were incubated with 500nM JD1 positive Aβ assemblies or an
equal volume of PBS buffer (controls), followed by baseline recordings for 20min prior to high
frequency stimulation (HFS) and LTP recording during 60min. LTP measurements were paired
in a way, where control and Aβ treatment was carried out on hippocampal tissues derived from
the same mouse brain.

Hippocampal tissues treated with these small oligomers resulted in a statistically significant
reduction (123.8 ± 9.05, n=5) (p≤0.01) in LTP measured at 1hr post-HFS, when compared to
matched controls (190.6 ±22.94, n=5, ) (Fig. 4b), which concurs with earlier reports showing
an Aβ mediated attenuation of LTP *in vivo* and *in vitro* (5,17,18). IP-MS (JD1) analysis of the
aCSF bath-solution showed that JD1 positive oligomers remained stable during the time course
of LTP recordings (data not provided).

**LMW oligomers induce acute changes in somatostatin and neurogranin levels in the**
**mouse hippocampus.**

Following LTP recordings, brain slices from control and Aβ treated hippocampal tissues were
subjected to protein extraction, followed by downstream deep proteomic analysis to assess any
putative changes in synaptic markers following acute treatment with LMW Aβ assemblies. We
employed a relative quantitative proteomics approach using isobaric labelling with tandem-
mass-tags (TMT) followed by off-gel isoelectric focusing fractionation (IEF) (Fig. 4c), to study
the relative change in brain protein abundance and post-translational modifications (PTM) as a
result of acute Aβ treatment.

Deep proteomic analysis of mouse hippocampal tissues allowed for unbiased identification of
≤5440 protein groups (FDR ≤1%) of which 6% of this proteome is associated with cellular
signalling mechanisms (Fig. 4d-e, Supplementary Data Fig. 6-8). Overall, extended
bioinformatics analysis of the mouse brain proteome did not reveal major changes in protein
abundances in control versus Aβ treated tissues (Fig. 4d-e and Supplementary Data Fig. 9). This
observation is in agreement with the general knowledge that *de novo* protein synthesis is mainly
associated with late-phase LTP, consisting of several hours or days (38). We nevertheless found
a statistically significant fold-change (-Log p = 2.84) of the pre-synaptic protein somatostatin
(SMS) following Aβ treatment (Fig. 4d-e and Supplementary Data Fig. 9a&b). Interestingly,
rapid hippocampal SMS release has been associated with acute stress in animals (39) and
somatostatin is known to play a key role in regulating Aβ metabolism through modulating
proteolytic degradation by neprilysin (40).

Next, a functional protein partnership search was carried out using the STRING database for
the integration of both, known and predicted protein-protein interaction networks (physical or
functional) to identify potential somatostatin interaction networks (Fig. 5a-b and
Supplementary Data Fig. 8a-b). Active somatostatin interaction was found for the protein
neurogranin (Neug / Nrgn), a well-established marker of neurodegeneration (41,42), whereas
neurogranin itself appears to form a network hub with different pre- and post-synaptic markers
(Fig. 5b). Bioinformatics analysis revealed a small fold change (Log2 fold change = 0.58) for
NEUG in Aβ treated tissue, however, this marker failed to show statistical significance (-Log p
= 1.34) (Fig. 4d-e). The small change in NEUG levels following Aβ treatment is an intriguing
observation because neurogranin expression is known to be regulated by synaptic activity (43)
and memory encoding requires a rapid (≤ 15min) *de novo* synthesis of neurogranin in the mouse
hippocampus (44). Moreover, neurogranin is known to play a key role in the regulation of

metaplasticity, by lowering the threshold for LTP induction (45) and the Aβ oligomer induced

inhibitory effect on LTP can be reversed by increasing neurogranin levels (46).

730
731
732
733

734
735

**Figure 5. Acute Aβ neurotoxicity induces rapid changes in somatostatin and neurogranin levels in the mouse hippocampus. a** Protein abundance analysis showing the relative abundance of different pre- and post-synaptic markers highlighted in red. **b** Selected zoom of the protein association network analysis (STRING), centred on neurograning (Neug) protein and showing an interaction with somatostatin (SMS) protein as well as a network with other proteins. (see Supplementary Data Fig. 8 for larger interaction network). **c** Summary of label-free, quantitative MS of the pre- and post-synaptic markers: somatostatin, alpha-synuclein and neurogranin (from top to bottom). Left; $MS^2$ spectra of one proteotypic peptide monitored for SMS, SYUA and NEUG protein. Centre; $MS^2$ fragment areas of selected $MS^2$ transitions. Right; calculated relative protein areas derived from quantitative MS analysis. **d** Orthogonal validation of SYUA levels found in Aβ treated (lanes: 2, 4 & 6) and control (lanes: 1, 3 & 5) brain tissue. (lane 7: MWM) using WB. Below; Summary of actin normalized (ACTB) densitometry analysis of NEUG levels detected in Aβ treated (110.3±10, mean ± SD, n=3) versus control (94.3±23) (p=0.489, t-test, two-tailed) animals. (ns: not significant). **e** Orthogonal validation of NEUG levels found in Aβ treated (lanes: 2, 4 & 6) and control (lanes: 1, 3 & 5) brain tissue. (lane 7: MWM) using WB. Below; Summary of actin normalized (ACTB) densitometry analysis of NEUG levels detected in Aβ treated (152±38, mean ± SD, n=3) versus control (100±16.5) (p=0.082, t-test, two-tailed) animals. (ns: not significant). **f** Summary of neo-epitope (N3, N4 & N5ns) and JD1 antibody binding features, as well as earlier reported binding properties of different clinical antibodies. The diagram depicts a perspective of the current understanding of the pathological accumulation of Aβ assemblies (morphologies) and antibody molecules against these different Aβ entities: oligomers (ACU 193, aducanumab) and protofibrils (lecanemab), fibrils or amyloid plaques (gantenerumab, aducanumab) or modifications (pyro-E3) found within the N-terminal region of Aβ (donanemab). The combination of both, neo-epitope and JD1 antibodies may allow early interception of small oligomeric species prior to the accumulation of larger Aβ assemblies.

To confirm the significance of the differentially regulated levels found for somatostatin and neurogranin, we decided to analyse the abundance of these two synaptic markers using a targeted, quantitative MS approach known as Parallel Reaction Monitoring (PRM), which allows unbiased identification of low abundant species in complex sample matrices. Quantitative MS analysis revealed that, levels of SMS were indeed significantly increased in Aβ treated mouse brains when compared to control brain slices. Similarly, NEUG levels where approximately ≥ 30% increased in Aβ treated brains, whereas levels of the predominantly found presynaptic marker α-synuclein (SYUA) did not differ between both groups (Fig. 5c and Supplementary Data Fig. 9b&c). Orthogonal validation by WB of the pre- and postsynaptic markers of SYUA (Fig. 5d) and NEUG respectively (Fig. 5e), as well as homer1 and post-synaptic density protein (PSD95 / Dlg4) (Supplementary Data Fig. 9c), further corroborate the findings by MS. However, immuno-blot detection of the small, presynaptic protein somatostatin proved to be challenging (Supplementary Data Fig. 9c) and therefore highlights the advantage of implementing targeted and quantitative MS assays, for an unbiased identification and quantification of scarce proteins (Supplementary Data Fig. 10).

**Aβ oligomer mediated changes in mouse hippocampal phospho-proteome.**

It has been well established that Aβ accelerates hyperphosphorylation of tau protein, which in

turn aggravates neuronal tauopathy (47). We therefore sought to identify putative changes in

site-specific protein phosphorylation, as a result of acute Aβ treatment, using a titanium-dioxide

(TiO$_2$) bead enabled phospho-peptide enrichment approach, followed by LC-MS analysis.

Overall, phospho-enrichment analysis allowed the identification of ≥7800 phospho-sites (FDR

≤1%), in control and Aβ treated hippocampal slices. Our target proteins of interest were

primarily focused on well-established markers of synaptic plasticity, as well as proteins that

have previously been reported to be susceptible to phosphorylation as a result of Aβ toxicity

(48). Phospho-proteomic analysis revealed differential levels of site-specific phosphorylation

of the calcium/calmodulin dependent protein kinase II alpha (CAMKIIa) protein, post-synaptic

density protein (PSD95 / Dlg4), synapsin 1 & 2 (SYN1 & SYN2), synaptophysin (SYPH) as

well as tau (TAU) protein in control versus Aβ treated brains (Supplementary Data Table II).

In spite of the here reported MS identification of some differentially modified tau species found

in acute Aβ treated brains, we were not able to confirm earlier reported tau phosphorylation *in*

*vitro* at residues pS202 or pS416 using structurally different Aβ assemblies (49).

Collectively, our deep proteomic and phospho-proteomic analysis may further shed light on the

acute Aβ mediated effects on synaptic plasticity. The observed, differentially regulated synaptic

protein somatostatin and, to some extent neurogranin, is intriguing, and clearly merits a more

thorough investigation in the future but is beyond the scope of the current study.

**Discussion:**

In this study, we provide analytical insight into the composition of LMW Aβ oligomers. We

show that small Aβ assemblies of putative dimeric, trimeric and tetrameric nature consist of a

combination of full-length Aβ42 as well as a repertoire of C-terminally truncated Aβ species,

of which the latter may also independently seed to form SDS-stable, HMW oligomeric

assemblies of 50kDa-200kDa mass range. Our unbiased MS identification of identical Aβ

fragment fingerprints, together with the JD1 positive assemblies detected in post-mortem AD

brains, further highlights the pathological significance of Aβ fragments; Aβ1-23 to Aβ1-25.

Generally, our findings allow to draw strong parallels with earlier reported observations of

typical Aβ fragment fingerprints (12) as well as the detection of a repertoire of different N- and,

C-terminally truncated Aβ species in human CSF samples (36,49,50). The independent report

of specific Aβ fragments found in human post-mortem AD brains as well as Down-syndrome

(DS) subjects (51,52) further corroborates the significance of the findings reported here.

Nevertheless, it remains to be confirmed, if the here reported key findings mirror earlier reports

on the inhibition of LTP, using similar LMW oligomeric assemblies (22) or, soluble ~7kDa Aβ

species isolated from post-mortem AD brains (9,10,15).

We provide compelling analytical proof, that the application of our recently developed neo-

epitope antibodies against different truncated Aβ isoforms allows for a highly selective pull-

down (IP) of SDS-PAGE stable assemblies from human post-mortem AD brains.  These

isolated Aβ assemblies strongly resemble the LMW Aβ species (4-7kDa) typically found

following IP of formic acid induced dissociation of ADDLs. Our findings are in strong

agreement with earlier reports indicating, that LMW Aβ oligomers consist of a repertoire of

truncated Aβ species and full-length Aβ42 or Aβ40 (16,53). However, earlier analytical

attempts in identifying specific Aβ isoforms associated with LWM Aβ assemblies and, more

specifically, the ~4kDa and ~7kDa Aβ species, have failed (16).

The lack in scientific and clinical consensus on the therapeutic efficacy of some clinical

antibodies as well as the ongoing, heated debate on viable, oligomeric Aβ targets (54) (55),

emphasizes the strong need for identifying highly specific Aβ targets for the development of

next generation disease modifying-therapies. Recent studies corroborate the notion, that

therapeutically targetable Aβ seeds exist during the lag phase of protein aggregation, and that

acute administration of aducanumab in young mice led to a significant reduction of Aβ

pathology in the mouse brain (1). Therefore, a successful clinical translation of specific Aβ

target hotspots will allow a therapeutic interception of early Aβ seeds during the long, insidious

lag phase of AD and prior to the accumulation of larger, metastable Aβ assemblies of

protofibrillar or fibrillar nature (Fig. 5f). Improving antibody-binding properties for small Aβ

assemblies may allow a selective targeting of early, pathological Aβ seeds during disease onset

and therefore attenuate or even prevent the progression to severe dementia.  The here reported

monoclonal antibody binding features observed with JD1 represents a first example of a new

class of future antibody molecules.

Our understanding of JD1 binding properties give rise to important Aβ target "loopholes"

associated with earlier developed anti-conformational antibodies. Yet, it remains to be

determined, if other antibody molecules, such as the anti-oligomer ACU 193 (3) or anti-

protofibril antibody lecanemab, currently undergoing clinical validation (56,57), provide

similar binding properties for the here described LMW Aβ assemblies. Our observation on

aducanumab's decreased binding affinity for LMW oligomeric assemblies has not been

reported before, and this finding is somehow surprising, because aducanumab's binding epitope

is known to span the N-terminal region of residues 3-6. Preferential aducanumab binding was

reported for oligomeric and fibrillar Aβ species derived from full-length Aβ42 peptide (29),

whereas oligomers derived from the N-terminally truncated forms of Aβ 5-42 to Aβ11-42 were

no longer detected with aducanumab (58). It is conceivable, that the strong seeding propensity

observed with fragments Aβ1-23, Aβ1-24 and Aβ1-25, as well as the C-terminally amidated

forms of these fragments, involves rapid conformational changes associated with the amyloid

driving segment of residues 16-25 and that JD1 binding is associated with a unique

conformational structure found within this Aβ segment. Discrepancies in aducanumab binding

properties have been previously observed, following immunization of a transgenic mouse

model of AD, resulting in a clear reduction in amyloid plaques but a failure in clearance of

vascular amyloid (CAA) (1,29). Interestingly, it was reported that the deposition of specific Aβ

isoforms appears to differ in subjects with vascular dementia, when compared with AD patients

(51), which is in line with the positive immuno-staining found for fragment Aβ1-25 in CAA

subjects (12).

Significant IHC staining of Aβ oligomers was reported using different conformation specific

antibodies (3) with binding properties against large (≥80kDa) Aβ assemblies (59). The lack in

JD1 staining of amyloid plaques may not appear surprising, given JD1's binding specificity for

LMW Aβ assemblies and the overall scarcity of Aβ oligomers reported in the human brain,

which is an observation also made by others using different anti-oligomer antibodies (3,28,61).

Nevertheless, our high-resolution array tomography clearly shows co-localization of JD1

positive species at post-synaptic terminals, which is in line with earlier reports, using different

conformational specific antibodies (34).

Taken together, the selective binding profile found with JD1 could be a beneficial antibody

feature, by decreasing the risk of dose-limiting side effects associated with passive

immunization, as well as increasing the antibody's bioavailability for rather early, neurotoxic

oligomeric assemblies (61). Overall, the here reported binding properties of JD1 highlights the

importance in further improving and developing novel antibody tools, with high target

selectivity for small, pathological Aβ seeds or oligomeric entities derived thereof. To date, the

development of highly specific anti-oligomeric antibodies remains a strong unmet medical need

for future disease-modifying therapeutics in AD. Improvement of immuno-based target

selectivity for acute, pathological Aβ species will inevitably provide early insight into the

disease pathology. This in turn will serve as an indispensable diagnostic tool for evaluation of

disease onset and progression, prior to the accumulation of clinical targets associated with pyro-

Glu$_3$ modification (Donanemab) (61) or large Aβ assemblies of protofibrilar (lecanemab) (57)

and fibrillar  structure (gantenerumab) (56) (Fig. 5f).

Monitoring changes in CSF and plasma Aβ42/40 ratios, as well as detection of early Aβ seeds,

together with other, clinically well-established biomarkers of neurodegeneration (e.g. p-Tau,

Nfl, Ngrn), may overall improve the diagnostic accuracy for individuals suffering from mild

cognitive impairment, who may benefit from early therapeutic intervention with future disease

modifying therapies. This could help to attenuate the pathological progression to severe

dementia and overall improve the quality of life in patients in the long run.

**References**

1. Uhlmann RE, Rother C, Rasmussen J, Schelle J, Bergmann C, Ullrich Gavilanes EM, et al. Acute targeting of pre-amyloid seeds in transgenic mice reduces Alzheimer-like pathology later in life. Nat Neurosci. 2020 Dec;23(12):1580–8.

2. Selkoe DJ. Soluble oligomers of the amyloid β-protein impair synaptic plasticity and behavior. Behavioural Brain Research. 2008 Sep;192(1):106–13.

3. Viola KL, Bicca MA, Bebenek AM, Kranz DL, Nandwana V, Waters EA, et al. The Therapeutic and Diagnostic Potential of Amyloid β Oligomers Selective Antibodies to Treat Alzheimer's Disease. Front Neurosci. 2022 Jan 3;15:768646.

4. Glabe CG. Structural Classification of Toxic Amyloid Oligomers. Journal of Biological Chemistry. 2008 Oct;283(44):29639–43.

5. Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, et al. Naturally secreted oligomers of amyloid b protein potently inhibit hippocampal long-term potentiation in vivo. 2002;416:5.

6. Liu P, Reed MN, Kotilinek LA, Grant MKO, Forster CL, Qiang W, et al. Quaternary Structure Defines a Large Class of Amyloid-β Oligomers Neutralized by Sequestration. Cell Reports. 2015 Jun;11(11):1760–71.

7. Cerf E, Sarroukh R, Tamamizu-Kato S, Breydo L, Derclaye S, Dufrêne YF, et al. Antiparallel β-sheet: a signature structure of the oligomeric amyloid β-peptide. Biochemical Journal. 2009 Aug 1;421(3):415–23.

8. Gu L, Liu C, Stroud JC, Ngo S, Jiang L, Guo Z. Antiparallel Triple-strand Architecture for Prefibrillar Aβ42 Oligomers. Journal of Biological Chemistry. 2014 Sep;289(39):27300–13.

9. Yang T, Li S, Xu H, Walsh DM, Selkoe DJ. Large Soluble Oligomers of Amyloid β-Protein from Alzheimer Brain Are Far Less Neuroactive Than the Smaller Oligomers to Which They Dissociate. J Neurosci. 2017 Jan 4;37(1):152–63.

10.    Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, et al.
Amyloid-β protein dimers isolated directly from Alzheimer's brains impair synaptic
plasticity and memory. Nat Med. 2008 Aug;14(8):837–42.

11.    Lesné S, Koh MT, Kotilinek L, Kayed R, Glabe CG, Yang A, et al. A specific amyloid-
β protein assembly in the brain impairs memory. Nature. 2006 Mar;440(7082):352–7.

12.    Rudinskiy N, Fuerer C, Demurtas D, Zamorano S, De Piano C, Herrmann AG, et al.
Amyloid-beta oligomerization is associated with the generation of a typical peptide fragment
fingerprint. Alzheimer's & Dementia. 2016 Sep;12(9):996–1013.

13.    O'Nuallain B, Freir DB, Nicoll AJ, Risse E, Ferguson N, Herron CE, et al. Amyloid -
Protein Dimers Rapidly Form Stable Synaptotoxic Protofibrils. Journal of Neuroscience.
2010 Oct 27;30(43):14411–9.

14.    Klyubin I, Betts V, Welzel AT, Blennow K, Zetterberg H, Wallin A, et al. Amyloid
Protein Dimer-Containing Human CSF Disrupts Synaptic Plasticity: Prevention by Systemic
Passive Immunization. Journal of Neuroscience. 2008 Apr 16;28(16):4231–7.

15.    Mc Donald JM, O'Malley TT, Liu W, Mably AJ, Brinkmalm G, Portelius E, et al. The
aqueous phase of Alzheimer's disease brain contains assemblies built from ∼4 and ∼7 kDa
Aβ species. Alzheimer's & Dementia. 2015 Nov;11(11):1286–305.

16.    Brinkmalm G, Hong W, Wang Z, Liu W, O'Malley TT, Sun X, et al. Identification of
neurotoxic cross-linked amyloid-b dimers in the Alzheimer's brain. :17.

17.    Freir DB, Holscher C, Herron CE. Blockade of Long-Term Potentiation by β-Amyloid
Peptides in the CA1 Region of the Rat Hippocampus In Vivo. Journal of Neurophysiology.
2001 Feb 1;85(2):708–13.

18.    Schmid AW, Freir DB, Herron CE. Inhibition of LTP in vivo by beta-amyloid peptide
in different conformational states. Brain Research. 2008 Mar;1197:135–42.

19.    Freir DB, Herron CE. Nicotine Enhances the Depressive Actions of Aβ $_{1-40}$ on Long-

961           Term Potentiation in the Rat Hippocampal CA1 Region In Vivo. Journal of

962           Neurophysiology. 2003 Jun;89(6):2917–22.

963    20.    McIntee FL, Giannoni P, Blais S, Sommer G, Neubert TA, Rostagno A, et al. In vivo

964           Differential Brain Clearance and Catabolism of Monomeric and Oligomeric Alzheimer's Aβ

965           protein. Front Aging Neurosci [Internet]. 2016 Sep 27 [cited 2020 Apr 7];8. Available from:

966           http://journal.frontiersin.org/Article/10.3389/fnagi.2016.00223/abstract

967    21.    Mazzitelli S, Filipello F, Rasile M, Lauranzano E, Starvaggi-Cucuzza C, Tamborini M,

968           et al. Amyloid-β $_{1-24}$ C-terminal truncated fragment promotes amyloid-β $_{1-42}$ aggregate

969           formation in the healthy brain. acta neuropathol commun. 2016 Dec;4(1):110.

970    22.    Lambert MP, Klein WL. Diffusible, nonfibrillar ligands derived from Abeta1-42 are

971           potent central nervous system neurotoxins. Proc Natl Acad Sci USA. 1998 May;

972    23.    Velasco PT, Heffern MC, Sebollela A, Popova IA, Lacor PN, Lee KB, et al. Synapse-

973           Binding Subpopulations of Aβ Oligomers Sensitive to Peptide Assembly Blockers and scFv

974           Antibodies. ACS Chem Neurosci. 2012 Nov 21;3(11):972–81.

975    24.    Sorrentino V, Romani M, Mouchiroud L, Beck JS, Zhang H, D'Amico D, et al.

976           Enhancing mitochondrial proteostasis reduces amyloid-β proteotoxicity. Nature. 2017

977           Dec;552(7684):187–93.

978    25.    Rudinskiy et al. - 2016 - Amyloid-beta oligomerization is associated with th.pdf.

979    26.    Kayed R, Head E, Sarsoza F, Saing T, Cotman CW, Necula M, et al. Fibril specific,

980           conformation dependent antibodies recognize a generic epitope common to amyloid fibrils

981           and fibrillar oligomers that is absent in prefibrillar oligomers. Mol Neurodegeneration. 2007

982           Dec;2(1):18.

27.    Kayed R, Canto I, Breydo L, Rasool S, Lukacsovich T, Wu J, et al. Conformation

dependent monoclonal antibodies distinguish different replicating strains or conformers of

prefibrillar Aβ oligomers. Mol Neurodegeneration. 2010 Dec;5(1):57.

28.    Gibbs E, Silverman JM, Zhao B, Peng X, Wang J, Wellington CL, et al. A Rationally

Designed Humanized Antibody Selective for Amyloid Beta Oligomers in Alzheimer's

Disease. Sci Rep. 2019 Dec;9(1):9870.

29.    Sevigny J, Chiao P, Bussière T, Weinreb PH, Williams L, Maier M, et al. The antibody

aducanumab reduces Aβ plaques in Alzheimer's disease. Nature. 2016 Sep 1;537(7618):50–

6.

30.    Ladiwala ARA, Litt J, Kane RS, Aucoin DS, Smith SO, Ranjan S, et al. Conformational

Differences between Two Amyloid β Oligomers of Similar Size and Dissimilar Toxicity. J

Biol Chem. 2012 Jul 13;287(29):24765–73.

31.    Britschgi M, Olin CE, Johns HT, Takeda-Uchimura Y, LeMieux MC, Rufibach K, et

al. Neuroprotective natural antibodies to assemblies of amyloidogenic peptides decrease with

normal aging and advancing Alzheimer's disease. Proc Natl Acad Sci USA. 2009 Jul

21;106(29):12145–50.

32.    Klaver AC, Finke JM, Digambaranath J, Balasubramaniam M, Loeffler DA. Antibody

concentrations to Aβ1–42 monomer and soluble oligomers in untreated and antibody–

antigen-dissociated    intravenous    immunoglobulin    preparations.    International

Immunopharmacology. 2010 Jan;10(1):115–9.

33.    Magga J, Puli L, Pihlaja R, Kanninen K, Neulamaa S, Malm T, et al. Human intravenous

immunoglobulin provides protection against Ab toxicity by multiple mechanisms in a mouse

model of Alzheimer's disease. 2010;15.

34.      Pickett EK, Koffie RM, Wegmann S, Henstridge CM, Herrmann AG, Colom-Cadena

1007            M, et al. Non-Fibrillar Oligomeric Amyloid-β within Synapses. Stranahan A, editor. JAD.

1008            2016 Aug 3;53(3):787–800.

1009      35.      Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild Cognitive

1010            Impairment: Clinical Characterization and Outcome. Arch Neurol. 1999 Mar 1;56(3):303.

1011      36.      Portelius E, Andreasson U, Ringman JM, Buerger K, Daborg J, Buchhave P, et al.

1012            Distinct cerebrospinal fluid amyloid β peptide signatures in sporadic and PSEN1 A431E-

1013            associated familial Alzheimer's disease. Mol Neurodegeneration. 2010;5(1):2.

1014      37.      Hebbar S, Lee E, Manna M, Steinert S, Kumar GS, Wenk M, et al. A fluorescent

1015            sphingolipid binding domain peptide probe interacts with sphingolipids and cholesterol-

1016            dependent raft domains. Journal of Lipid Research. 2008 May;49(5):1077–89.

1017      38.      Younts TJ, Monday HR, Dudok B, Klein ME, Jordan BA, Katona I, et al. Presynaptic

1018            Protein Synthesis Is Required for Long-Term Plasticity of GABA Release. Neuron. 2016

1019            Oct;92(2):479–92.

1020      39.      Arancibia S, Payet O, Givalois L, Tapia-Arancibia L. Acute stress and dexamethasone

1021            rapidly increase hippocampal somatostatin synthesis and release from the dentate gyrus

1022            hilus. Hippocampus. 2001;11(4):469–77.

1023      40.      Saito T, Iwata N, Tsubuki S, Takaki Y, Takano J, Huang SM, et al. Somatostatin

1024            regulates brain amyloid β peptide Aβ42 through modulation of proteolytic degradation. Nat

1025            Med. 2005 Apr;11(4):434–9.

1026      41.      Hesse R, Hurtado ML, Jackson RJ, Eaton SL, Herrmann AG, Colom-Cadena M, et al.

1027            Comparative profiling of the synaptic proteome from Alzheimer's disease patients with

1028            focus on the APOE genotype. acta neuropathol commun. 2019 Dec;7(1):214.

1029     42.     Wellington H, Paterson RW, Portelius E, Törnqvist U, Magdalinou N, Fox NC, et al.

1030            Increased CSF neurogranin concentration is specific to Alzheimer disease. Neurology. 2016

1031            Mar 1;86(9):829–35.

1032     43.     Garrido-García A, de Andrés R, Jiménez-Pompa A, Soriano P, Sanz-Fuentes D,

1033            Martínez-Blanco E, et al. Neurogranin Expression Is Regulated by Synaptic Activity and

1034            Promotes Synaptogenesis in Cultured Hippocampal Neurons. Mol Neurobiol. 2019

1035            Nov;56(11):7321–37.

1036     44.     Jones KJ, Templet S, Zemoura K, Kuzniewska B, Pena FX, Hwang H, et al. Rapid,

1037            experience-dependent translation of neurogranin enables memory encoding. Proc Natl Acad

1038            Sci USA. 2018 Jun 19;115(25):E5805–14.

1039     45.     Zhong L, Gerges NZ. Neurogranin Regulates Metaplasticity. Front Mol Neurosci. 2020

1040            Jan 24;12:322.

1041     46.     Kaleka KS, Gerges NZ. Neurogranin restores amyloid β-mediated synaptic transmission

1042            and long-term potentiation deficits. Experimental Neurology. 2016 Mar;277:115–23.

1043     47.     Terwel D, Muyllaert D, Dewachter I, Borghgraef P, Croes S, Devijver H, et al. Amyloid

1044            Activates GSK-3β to Aggravate Neuronal Tauopathy in Bigenic Mice. The American

1045            Journal of Pathology. 2008 Mar;172(3):786–98.

1046     48.     Jin M, Shepardson N, Yang T, Chen G, Walsh D, Selkoe DJ. Soluble amyloid β-protein

1047            dimers isolated from Alzheimer cortex directly induce Tau hyperphosphorylation and

1048            neuritic degeneration. Proc Natl Acad Sci USA. 2011 Apr 5;108(14):5819–24.

1049     49.     Amar F, Sherman MA, Rush T, Larson M, Boyle G, Chang L, et al. The amyloid-β

1050            oligomer Aβ*56 induces specific alterations in neuronal signaling that lead to tau

1051            phosphorylation and aggregation. Sci Signal. 2017 May 9;10(478):eaal2021.

50.    Bibl M, Gallus M, Welge V, Esselmann H, Wiltfang J. Aminoterminally Truncated and

Oxidized Amyloid-β Peptides in the Cerebrospinal Fluid of Alzheimer's Disease Patients.

JAD. 2012 Apr 16;29(4):809–16.

51.    Larner AJ. Hypothesis: amyloid NL-peptides truncated at the N-terminus contribute to

the pathogenesis of Alzheimer's disease. Neurobiology of Aging. 1999;5.

52.    Gkanatsiou E, Portelius E, Toomey CE, Blennow K, Zetterberg H, Lashley T, et al. A

distinct brain beta amyloid signature in cerebral amyloid angiopathy compared to

Alzheimer's disease. Neuroscience Letters. 2019 May;701:125–31.

53.    Gkanatsiou E, Sahlin C, Portelius E, Johannesson M, Söderberg L, Fälting J, et al.

Characterization of monomeric and soluble aggregated Aβ in Down's syndrome and

Alzheimer's disease brains. Neuroscience Letters. 2021 May;754:135894.

54.    Hong W, Wang Z, Liu W, O'Malley TT, Jin M, Willem M, et al. Diffusible, highly

bioactive oligomers represent a critical minority of soluble Aβ in Alzheimer's disease brain.

Acta Neuropathol. 2018 Jul;136(1):19–40.

55.    Benilova I, Karran E, De Strooper B. The toxic Aβ oligomer and Alzheimer's disease:

an emperor in need of clothes. Nat Neurosci. 2012 Mar;15(3):349–57.

56.    Ashe. Alzheimer's target still viable but untested.

57.    Bohrmann B, Baumann K, Benz J, Gerber F, Huber W, Knoflach F, et al.

Gantenerumab: A Novel Human Anti-Aβ Antibody Demonstrates Sustained Cerebral

Amyloid-β Binding and Elicits Cell-Mediated Removal of Human Amyloid-β. JAD. 2012

Jan 12;28(1):49–69.

58.    Sehlin D, Englund H, Simu B, Karlsson M, Ingelsson M, Nikolajeff F, et al. Large

Aggregates Are the Major Soluble Aβ Species in AD Brain Fractionated with Density

Gradient Ultracentrifugation. Bush AI, editor. PLoS ONE. 2012 Feb 15;7(2):e32014.

59.    Desai AA, Smith MD, Zhang Y, Makowski EK, Gerson JE, Ionescu E, et al. Rational

1077            affinity maturation of anti-amyloid antibodies with high conformational and sequence

1078            specificity. Journal of Biological Chemistry. 2021 Jan;296:100508.

1079    60.    Sebollela A, Mustata GM, Luo K, Velasco PT, Viola KL, Cline EN, et al. Elucidating

1080            Molecular Mass and Shape of a Neurotoxic Aβ Oligomer. ACS Chem Neurosci. 2014;8.

1081    61.    Bakrania P, Hall G, Bouter Y, Bouter C, Beindorff N, Cowan R, et al. Discovery of a

1082            novel pseudo β-hairpin structure of N-truncated amyloid-β for use as a vaccine against

1083            Alzheimer's disease. Mol Psychiatry [Internet]. 2021 Nov 15 [cited 2022 Apr 6]; Available

1084            from: https://www.nature.com/articles/s41380-021-01385-7

1085    62.    Cynis H, Frost JL, Crehan H, Lemere CA. Immunotherapy targeting pyroglutamate-3

1086            Aβ: prospects and challenges. Mol Neurodegeneration. 2016 Dec;11(1):48.

1087    63.    Metais C, Brennan K, Mably AJ, Scott M, Walsh DM, Herron CE. Simvastatin

1088            Treatment Preserves Synaptic Plasticity in A $_{NL}$ PPswe/PS1dE9 Mice. C Me. :15.

1089    64.    Hughes B, Herron CE. Cannabidiol Reverses Deficits in Hippocampal LTP in a Model

1090            of Alzheimer's Disease. Neurochem Res. 2019 Mar;44(3):703–13.

1091

1092

1093

1094

1095

1096

1097    **Materials and Methods:**

1098

**Aβ peptide preparation.**

Full-length wild type (wt) Aβ peptides Aβ1-40 and Aβ1-42 Aβ (BACHEM, Switzerland) were

dissolved in 1,1,1,3,3,3-Hexafluoro-2-propanol (HFIP) at a concentration of 1mg/ml, followed

by a 10-min sonication to break any preformed aggregates. HFIP solution was evaporated

under a ventilated fume hood by applying a light stream of $N_2$ gas. The HFIP film containing

the Aβ peptide was either directly re-suspended in 100% DMSO and further diluted to 1%

DMSO in a new buffer or stored dry at −80 °C until use. Aβ peptide fragments comprising of

residues: 1-23, 1-24-NH$_2$, 1-25, 1-25-NH$_2$ (purity of ≥ 97%) were purchased from GenicBio

Ltd. (Shanghai, China) and Aβ1-42 and Aβ1-40 were purchased from Bachem

(Schwitzerland), Aβ40-Arc was kindly provided by Prof. Hilal Lashuel (EPFL) and reverse

peptide Aβ42-1 was kindly provided by Prof. Gräff (EPFL). Recombinant alpha-synuclein

protein was a gift from rPeptide (GA, USA).

1111

**In vitro Aβ peptide aggregation studies.**

Aggregation kinetics were carried out as reported by (Ladiwala et al.). Briefly, HFIP dried Aβ

peptide films were solubilized in 50mM NaOH, sonicated and further diluted to 5mM with

PBS (or 30mM Tris, 150mM NaCl) to a final Aβ concentration of 0.05-0.1 mg/ml for truncated

Aβ fragments and full-length peptides of Aβ1-40, 0.1 mg/ml mutant Aβ1-40 Arc, and 0.05

mg/ml Aβ1-42. Aβ preparations were then centrifuged (22,000 x g) for 30min at 4°C and the

supernatant (90-95% volume) was recovered and transferred into a new Eppendorf tube.

Supernatant samples were then incubated at 37°C and left for spontaneous aggregation during

either 0-20 hrs (short-term) or 1-5 days (intermediate). Aggregated sample aliquots were drawn

at different time points and either analyzed immediately using MS and/or DB or snap frozen

in liquid nitrogen and stored at -80°C. Typically, one microliter was spotted onto a

nitrocellulose membrane for dotblotting (corresponding to either 25, 50ng to100ng / spot) or

mixed with alpha-cyano matrix for MALDI-TOF/TOF analysis.

1125

**Digestion of Aβ peptides.**

Proteolytic digestion using LysN (2ng/ul) was performed overnight at 37 °C in 50 mM

ammonium bicarbonate, pH 10 (LysN buffer). For in-gel digestions, peptides were extracted

from gels and concentrated by Speed-Vacuum prior to LC-MS/MS measurements.

Immunoprecipitated samples (IP), were reduced and alkylated followed by in-solution

digestion at 37°C using standard LysN buffer (50ul) and approximately 70ng of LysN. Dried

samples were resuspended in 5% DMSO and 2.5% FA followed by LC-MS/MS, LC-MRM

and PRM analysis.

1134

**Matrix assisted Laser Desorption/Ionization Time-of-flight Mass Spectrometry (MALDI**

**TOF/TOF).**

Aliquots (2 μl) of samples were used for MALDI-TOF/TOF MS (ABI 4800 model, Applied

Biosystems) measurements. Matrix solution of α-cyano-4-hydroxycinnamic acid (7 mg/ml in

ACN/0.1% TFA (1:1, v/v)) was used for sample deposition. The sample (1 μl) was mixed with

1 μl of matrix solution and then 1 μl of this mixture was deposited in duplicates on the target

plate and allowed to air dry. Samples were analyzed in reflectron positive mode.

1142

**1143 Sandwich ELISA for detection of Aβ oligomers**

JD1 antibody was adsorbed to high protein binding Nunc-Immuno MaxiSorp 96-well plates

(Nunc, Roskilde,) at 4°C for 4hrs, followed by a blocking step for 1hr using a 5% BSA in

PBST (Tween 20: 0.1%) solution. Synthetic, oligomeric preparations of Aβ1-24 or Aβ1-25

standards or, human TBS and FA brain extracts were incubated at 4°C over-night to allow for

binding to the capture antibody JD1. Detection of Aβ species was achieved using the mouse

monoclonal HRP-conjugated 6E10 (Biolegend) antibody at a dilution of 1:1000. Signals were

measured at an absorbance at 450nm using a Tecan Infinite F500 reader. All samples were

analyzed in triplicates.

1152

**1153 Mouse hippocampal tissue extraction, TMT labelling and LC-MS analysis.**

Mouse hippocampal proteins from PBS control and Aβ oligomer (JD1 positive) treated brain

slices were extracted with 2% SDS (in PBS) containing protease and phosphatase inhibitors

(Roche, Switzerland) followed by centrifugation at 14000 x g for 15min to precipitate any

insoluble material. Prior to proteolysis, samples were cleared of excess SDS using detergent

removal spin columns (Cat# 87777, Pierce, Switzerland) followed by total protein content

normalization using a standard Bradford protein assay. Samples were reduced and alkylated

followed by overnight digestion at 37°C using LysC / Trypsin proteases (Promega) and isobaric

labelling using tandem mass tags (TMT) as described elsewhere. Briefly, the TMT reagent (0.8

mg, Thermo Fisher Scientific) was dissolved in acetonitrile, and 10 μL of the reagent was

added to 30 μL of peptides along with anhydrous ACN to give a final concentration of 30%

v/v. The peptides were then pooled after incubating at room temperature for 1h followed by

C18 stage tip clearing and drying in a speed vacuum.

For Off-gel fractionation, samples were suspended in a 4M urea buffer containing ampholytes

and subjected to off-gel IEF fractionation using a 24 well strip (pH 3-10) (Agilent, 3100 Off-

gel fractionator). Following IEF, fractions were recovered and samples were cleared with C18

stage tips followed by speed vacuum drying and storage at -20°C until LC-MS analysis.

1170

**LC-MS/MS analysis**

Mass spectrometry (MS) analysis was performed on an Orbitrap Exploris 480 mass

spectrometer (Thermo Fisher Scientific) coupled to a nano-UPLC Dionex pump. For liquid

chromatography (LC) MS/MS analysis, trypsin / LysC digested samples were resuspended in

30-60ul of a mobile phase A solution (2% ACN / water, 0.1% formic acic) and then separated

by reversed-phase chromatography using a Dionex Ultimate 3000 RSLC nanoUPLC system

on a home-made 75 μm ID × 50 cm C18 capillary column (Reprosil-Pur AQ 120 Å, 1.9 μm)

in-line connected with the MS instrument. Peptides were separated by applying a non-linear

150min gradient ranging from 99% solvent A (2% ACN and 0.1% FA) to 90% solvent B (90%

ACN and 0.1%FA) at a flow rate of 250nl/min. Full-scan MS spectra (300−1500 m/z) were

acquired at a resolution of 120'000 at 200 m/z. Data-dependent MS/MS spectra were recorded

followed by HCD (higher-energy collision dissociation) fragmentation on the ten most intense

signals per cycle (2 s), using an isolation window of 1.4 m/z. HCD spectra were acquired at a

resolution of 60'000 using a normalized collision energy of 32 and a maximum injection time

of 100 ms. The automatic gain control (AGC) was set to 100'000 ions. Charge state screening

was enabled such that unassigned and charge states higher than six were rejected. Precursors

intensity threshold was set at 5'000. Precursor masses previously selected for MS/MS

measurement were excluded from further selection for a duration of 20 s, and the mass

exclusion window was set at 10 ppm. Theoretical isotopic envelopes of Aβ peptides were

drawn using the isotope calculator tool in Data Explorer Software (version 4.9). The mass

spectrometry data generated from this study have been deposited to the ProteomeXchange

Consortium (http://www.proteomexchange.org) via PRIDE partner repository with the dataset

identifier: PXD033934 (10.6019/PXD033934).

1194

1195

1196

1197

**Targeted and quantitative MS analyses**

Parallel reaction montitoring (PRM) analyses were performed as reported earlier (Sathe et al)

using an Orbitrap Exploris 480 mass spectrometer (Thermo Fisher Scientific) coupled to a

nano-UPLC Dionex pump. Peptides from selected target proteins, SST, NEUG, SYUA were

monitored using a targeted inclusion list. The parameters used for the PRM analysis were set

as follows: Orbitrap resolution of 35,000, AGC target value of 5 x 10 5, injection time of 200

ms. A precursor target isolation window of 2 m/z was applied and a normalized collision energy

of 35 was employed for MS/MS fragmentation. The PRM method was carried out in an

unscheduled mode and samples were analysed in duplicates. PRM data were analyzed using

the Skyline software package (MacCoss Lab). Acquired raw files were uploaded to Skyline and

peak integration was verified manually.

1209

**MS raw data processing and database searches**

PEAKS Studio X+ Pro (Bioinformatics Solutions Inc.) software was used for data processing.

The raw MS data files were imported into PEAKS Studio software using the following

parameters for the database search: For protein identification the UniProt / Swiss-Prot mus

musculus database (ID: UP000000589) combined with a decoy database was used. For peptide

identification the following settings were used: Enzyme: Trypsin, missed cleavages: 2 precursor

mass tolerance: 10 ppm, fragment mass tolerance: 0.2 Da, minimum charge: 2, maximum

charge: 5, fixed modifications: Carbamidomethyl (C), variable modifications: Oxidation (M),

phosphorylation (STY). False discovery rate (FDR) was calculated based on the target/decoy

database and peptides as well as proteins with FDR threshold of $\leq 1\%$ were chosen as true

positive hits.

MaxQuant searches: Label-free and TMT-labelled mouse brain proteomes were processed by

MaxQuant as reported earlier (Tyanova 2016). Briefly, MS raw files were processed using the

peptide identification algorithm of the Maxquant software (ver. 1.5.3.12) using the Andromega

search engine. TMT correction factors were updated according to the values provided

manufacture. MS/MS searches were performed with the following parameters: Peptides had

to have a minimum length of six amino acids to be considered for identification.

Carbamidomethylation was set as a fixed modification, while oxidation of methionine and

protein N-terminal acetylation were set as variable modifications. The enzyme specificity was

set to trypsin and allowed up to two miscleavages. The UniProt / Swiss-Prot mus musculus

database (ID: UP000000589) combined with a decoy database was used. A false discovery rate

(FDR)    of    $\leq 1\%$    was    applied    at    the    proteins    and    peptide    level.

Quantitative data analysis was performed using Skyline (version 21.1.0.278, MacCoss lab,

Uni. Washington, USA), an open source software tool application for quantitative data

processing and proteomic analysis. All integrated peaks were manually inspected to ensure

correct peak detection and integration.

Bioinformatic analysis was carried out using the Perseus (v.1.6.15.0) software environment and

compared with the bioinformatics output generated with the PEAKS Studio X+ Pro software to

identify any putative discrepancies due to the analytical stringencies or search engine

algorithms used in both workflows. Volcano plots were generated using the -log10 of the p-
value on the y-axis versus the mean fold-difference of the log2 transformed quantitative data
on the x-axis. A paired t-test with 'Benjamini-Hochberg FDR' method was applied for multiple
hypothesis corrections. Significance curves in the volcano plot corresponded to an S0 value of
0.5 and an FDR cut-off of 0.05. Protein interaction networks were analyzed and plotted using
the open source platforms: STRING DB (http://string-db.org) (Jensen et al.) and Cytoscape
(v.3.9.0) software (Shannon et al).

1246

**SDS-PAGE and Immunoblotting.**
Where indicated in the results, samples were mixed using either standard SDS Lämmli sample
buffer and heated at 80°C for 5 min prior to loading onto gels (Novex Nupage 4-12% Bis-Tris
gels, 1mm, 15 well, Invitrogen) and ran with Novex Nupage MES-SDS buffer. PAGE
separated samples were electroblotted using a constant voltage of 30V (90min) onto
nitrocellulose (0.22μm) membranes. Following transfer, membranes were boiled in PBS for
2min in a microwave oven (800W) and then blocked for 1h at room temperature under constant
rocking using Odyssey blocking buffer (Li-COR Biosciences, Bad Homburg, Germany)
diluted 1:3 in PBS. Following blocking, membranes were incubated at 4 °C with constant
rocking overnight using the commercially available mouse monoclonal antibodies 6E10 and
4G8 (0.5μg /ml) (Enzo, Life Sciences, Switzerland). Membranes were washed four times with
PBS-Tween (PBS containing 0.1% Tween 20), followed by incubation with a goat anti-rabbit
or anti-mouse secondary IgG antibody (highly cross-adsorbed) (dilution, 1:5000) conjugated
to Alexa Fluor 680 or 800 and scanned in a Li-COR scanner at a wavelength of 700 nm and
800 nm respectively. Densitometric analysis of WB band intensities was carried out using
image J software v.1.43 (NIH).

1263

1264

1265

**Dot blotting**.
Typically, 1ul samples were spotted onto a nitrocellulose membrane, which corresponded to a
total peptide load of 25ng (Aβ1-40) and (AB1-42) unless otherwise stated in the Fig.ures.
Samples were left to dry for 15min followed by blocking of the membrane (30min) with 5%
BSA in PBS solution. Membrane strips were incubated with primary antibodies overnight at
4°C on a shaker. For JD1 probing, membranes were incubated 2-4hrs at RT or over-night at
4°C followed by revelation with a goat anti-mouse, secondary Alexa-flour conjugated

antibody. Neo-epitope antibodies N3, N4, N5s N5ns and N5NH2 were applied as described
earlier (12). Identical solutions and revelation procedures with secondary antibodies were used
as described for immunoblotting above.

1276

**Immunoprecipitation of Tg mouse and human brain tissue.**

Tg mouse (9-12 months) (Tg2576) and cortical human post-mortem brain tissue samples were
extracted using either TBS, 1% SDS (in TBS) or formic acid (70-90%FA) as reported earlier
Rudinskiy et al. (12) Briefly, tissue samples were homogenized (20 strokes on ice) in TBS with
protease inhibitors (Roche, Switzerland) using a Teflon homogenizer. Samples were then
subjected to centrifugation (150,000g) during 45min and supernatant was recovered as the TBS
soluble fraction. Protein pellets were subjected to an additional extraction using either SDS
(1%) or FA (70%) followed by centrifugation. Typically, pellets were extracted with 70-90%
FA overnight at 4°C (to minimize formylation adducts) followed by centrifugation. SDS
fractions were diluted to ≤0.1% SDS final concentration and FA fractions were neutralized to
pH 7.5 with 5M sodium hydroxide (NaOH) solution prior to IP. All samples were initially
depleted of endogenous IgG's using a mixture of protein A&G agarose beads (Roche AG,
Switzerland). Typically, 3-5ug/ml rabbit polyclonal antibody (N3, N4, N5ns & N5NH2),
5ug/ml of JD1 (IgG) or 3-5ug/ml of 6E10 or 4G8 mouse monoclonal was used for IP overnight
under continues rotation (4rpm/min) at 4°C. Samples were eluted with 40% ACN /$H_2O$ & 0.1%
TFA for downstream MS analysis and dried in a speed vacuum or, boiled in Lämmli buffer for
WB analysis. IP'ed samples were either directly analysed by WB or MALDI-TOF/TOF or
further digested overnight using LysN proteolysis prior to LC-MS/MS analysis as outlined
above.

1296

1297

1298

1299

1300 **Animals**

1301 Hippocampal slices were prepared from 8 to 10 week old c57/black 6 mice (c57BL6), obtained
1302 from Charles River UK. Mice were housed in the biomedical facility, University College
1303 Dublin with a 12 h light/dark cycle with food and water ad libitum. All experiments were
1304 conducted under license from the Dept. of Health, Ireland, Directive 63/2010/EU.

1305

**Hippocampal Slices**

Parasagittal hippocampal slices, were prepared using a vibrotome (Leica VT1000S) as

described previously (62,63) using ice cold cutting solution (mM) NaCl 87; NaHCO$_3$ 25;

Glucose 25; Sucrose 75; KCl 2.5; NaH$_2$PO$_4$ 1.25; CaCl$_2$ 0.5; MgSO$_4$ 7; bubbled with 95%

O$_2$/5% CO$_2$ (carbogen). Slices (400-µm thick) were transferred to a holding chamber

containing aCSF (mM) NaCl 119; NaHCO$_3$ 26.2; Glucose 11 mM; KCl 2.5; NaH$_2$PO$_4$ 1; CaCl

$_2$ 2.5; MgSO$_4$ 1(also used for recording); bubbled with carbogen and were allowed to recover

at room temperature for at least 90 min Slices were transferred to a recording chamber, secured

by means of a harp with fine nylon strings and perfused with recording aCSF at a rate of

5 ml/min maintained at 28–30 °C for the duration of all experiments. A horizontal puller (DMZ

universal puller, Germany) was used to makerecording electrodes from borosilicate capillary

glass (GC150 F-10, Havard apparatus), Electrodes (2–5 MΩ) were filled with recording aCSF.

The Shaffer-collateral pathway was stimulated using a mono-polar electrode (FHC, Bowdoin,

USA) at 0.033 Hz (duration: 100 µs), the return electrode was a silver/silver chloride wire

placed in the recording bath. Extracellular field excitatory post synaptic potentials (fEPSPs)

were recorded in the CA$_1$ *stratum radiatum* and paired stimuli were delivered with an inter-

stimulus interval of 50 ms in order to monitor paired pulse facilitation (PPF). The voltage

signal was filtered at 5 kHz and stored for off-line analysis using a personal computer

interfaced with a Digidata 1440 and pclamp 10. Signals were amplified by a HS2A headstage

(Molecular Devices, USA) connected to an Axoclamp 2B system (Molecular Devices, USA)

and a Brownlee 410 Precision preamplifier. The stimulus voltage was adjusted to evoke a

fEPSP that was 40–50% of the maximal response (maximum fEPSP just prior to the formation

of a spike caused by cell firing) for the duration of all experiments. When a stable baseline had

been recorded for 20 min a Master 8 (AMPI) timer was used to deliver two trains of high

frequency stimuli (HFS), 100 Hz for 1 s, with an inter-train interval of 30 s. Following the

application of HFS, the synaptic response was recorded for a further period of 60 min. All

results are presented as mean ± SEM. The "*n*" numbers quoted refer to the number of slices

used. Control and test experiments in a given section were conducted on the same day on slices

from the same animal.

1335

**Development of monoclonal antibody against LMW Aβ oligomers.**

Briefly, mice were immunized with the linear sequence of Aβ1-24 (CONH2) as well as an

aggregated form of Aβ1-24 peptide. Eight weeks following immunization, sera were tested by

ELISA for antibodies against fresh (linear) and oligomeric Aβ species. Mice testing positive

for anti-Aβ antibodies were then given booster injections of 50 ug of an oligomeric form of

Aβ1-25. IgM type antibodies consisted of the predominantly identified type of anti-Aβ
antibodies found following this protocol. Antibody type switching and grafting from IgM to
IgG was then carried using recombinant technologies. All work was carried out by Eurogentec
SA, (Liege, Belgium).

1345

**Immunofluorescence and array tomography imaging of human post-mortem brain tissue.**
Samples from the inferior temporal gyrus of 4 AD patients was prepared for array tomography
as described previously (12,34). Brain samples were obtained with ethical approval via the
University of Edinburgh Sudden Death Brain Bank and the Alzheimer Scotland Brain and
Tissue Bank. This study was reviewed and approved by the Edinburgh Brain Bank ethics
committee and the Academic and Clinical Central Office for Research and Development, a
joint office of the University of Edinburgh and NHS Lothian (approval number 15-HV-
016). The Edinburgh Brain Bank is a Medical Research Council funded facility with
research ethics committee (REC) approval (16/ES/0084). Briefly, tissue was collected at
autopsy, fixed for 3 hours in 4% paraformaldehyde with 2.5% sucrose, dehydrated through a
series of ethanols, and embedded in LR White resin. Ribbons of 70nm serial sections were cut
with an ultracut microtome (Leica). Ribbons were stained with, primary antibodies OC (Sigma
AB2286, 1:200), JD1 (IgG, 1:200), and PSD95 (Synaptic Systems 124 014; 1:50) and detected
with anti-mouse IgG alexa fluor 594, and anti-rabbit alexa fluor 405, and anti-guinea pig alexa
fluor 647 (all at 1:50). Images from 5 serial sections were acquired on a Leica TSC SP8 confocal
with a 63x oil objective. Individual images were made into stacks, aligned, and segmented with
custom software that can be found at https://github.com/Spires-Jones-Lab For
immunofluorescence staining of 4 μm paraffin sections from formalin fixed tissue, sections
were de-waxed, rehydrated, and stained with amyloid beta antibodies JD1 (1:500) and OC
(1:1000) and secondaries donkey anti-rabbit 647 and donkey anti-mouse 594. Thioflavin S
counterstain was used to label fibrils (0.01%). Image stacks were acquired on a Leica TSC SP8
confocal with a 63x objective.

1368

1369    **Transmission Electron Microscopy (TEM).**
1370    Aβ peptide (5μl) samples were deposited on Formvar-coated 200-mesh copper grids (Electron
1371    Microscopy Sciences, Hatfield, PA). Grids were washed with two drops of double-distilled
1372    $H_2O$ and stained with two drops of freshly prepared 1% (w/v) uranyl formate (Electron
1373    Microscopy Sciences). The samples were analysed at room temperature using a Tecnai Spirit

1374 (FEI, The Netherlands) electron microscope equipped with a LaB6 filament and operated at an

1375 acceleration voltage of 80 kV. Images were recorded with an Eagle (FEI, The Netherlands)

1376 camera 4098 X 4098 pixels. Representative images of the main Aβ structures were taken at

1377 different magnifications ranging from 30kX to 50kX.

1378

1379 **C. elegans strain**

1380 C. elegans overexpressing Aβ42 peptide were developed as reported before (24). Briefly,

1381 c. elegans were cultured at 20 °C on nematode growth medium (NGM) agar plates seeded with

1382 E. coli strain OP50 unless stated otherwise. The GMC101 strain was used in this study and was

1383 provided by the Caenorhabditis Genetics Center (University of Minnesota).

1384