Moderate High Temperature is Beneficial or Detrimental Depending on Carbon Availability in the Green Alga *Chlamydomonas reinhardtii*

Ningning Zhang[1,#], Benedikt Venn[2,#], Catherine E. Bailey[1], Ming Xia[1], Erin M. Mattoon[1,3], Timo Mühlhaus[2], Ru Zhang[1,*]

[1]Donald Danforth Plant Science Center, St. Louis, Missouri 63132, USA;

[2]Computational Systems Biology, RPTU Kaiserslautern, Kaiserslautern, 67663, Germany;

[3]Plant and Microbial Biosciences Program, Division of Biology and Biomedical Sciences, Washington University in Saint Louis, St. Louis, MO 63130, USA;

[4]Bioinformatics and Computational Biology, Saint Louis University, St. Louis, Missouri 63103, USA.

Ningning Zhang: lucy.nnzhang@gmail.com, co-first-author[#]

Benedikt Venn: venn@rptu.de, co-first-author[#]

Ming Xia: celine.xia@wustl.edu, present address[3]

Catherine E. Bailey: cady.e.bailey@gmail.com, present address[4]

Erin M. Mattoon: emattoon@danforthcenter.org

Timo Mühlhaus: timo.muehlhaus@rptu.de

Ru Zhang: rzhang@danforthcenter.org, corresponding author*

Running title: Algal heat responses interact with carbon availability.

Highlight

We revealed the overlooked, dynamic effects of moderate high temperature in algae depending on carbon availability and demonstrated the importance of carbon metabolism in thermotolerance of photosynthetic cells.

Abstract

High temperatures impair plant and algal growth and reduce food and biofuel production,
but the underlying mechanisms remain elusive. The unicellular green
alga *Chlamydomonas reinhardtii* is a superior model to study heat responses in
photosynthetic cells due to its fast growth rate, many similarities in cellular processes to
land plants, simple and sequenced genome, and ample genetic and genomics resources.
Chlamydomonas grows in light by photosynthesis and/or with the externally supplied
organic carbon source, acetate. Most of the published research about Chlamydomonas
heat responses used acetate-containing medium. Understanding how organic carbon
sources affect heat responses is important for the algal industry but understudied. We
cultivated Chlamydomonas wild-type cultures under highly controlled conditions in
photobioreactors at control of 25°C, moderate high temperature of 35°C, or acute high
temperature of 40°C with and without constant acetate supply for 1- or 4-days. Our results
showed that 35°C increased algal growth with constant acetate supply but reduced algal
growth without sufficient acetate. The overlooked and dynamic effects of 35°C could be
explained by induced carbon metabolism, including acetate uptake and assimilation,
glyoxylate cycle, gluconeogenesis pathways, and glycolysis. Acute high temperature at
40°C for more than 2 days was lethal to algal cultures with and without constant acetate
supply. Our research provides insights to understand algal heat responses and help
improve thermotolerance in photosynthetic cells.
50
51 **Key words:** moderate high temperature, acute high temperature, heat responses,
52 *Chlamydomonas reinhardtii*, acetate, glyoxylate cycle, gluconeogenesis, glycolysis,
53 photobioreactors

54
55 **Abbreviations:**
56 Photobioreactors (PBRs), Tris-acetate-phosphate (TAP) medium, Tris-phosphate (TP)
57 medium
58

Introduction
Many biological processes in photosynthetic cells are sensitive to high temperatures
(Mittler *et al.*, 2012; Schroda *et al.*, 2015; Mishra *et al.*, 2021; Guihur *et al.*, 2022). Global

warming increases the intensity, duration, and frequency of high temperatures in the field,

reducing plant growth and food production (Janni *et al.*, 2020). A recent model revealed

high temperature as the primary climatic driver for yield loss in crops from 1981-2017 in

the U.S. (Ortiz-Bobea *et al.*, 2019). Considering the increasing global temperature and

human population, it is imperative to understand how photosynthetic cells respond to high

temperatures and improve plant thermotolerance.

High temperatures in the field have different intensities and durations. For many land

plants and algae grown in temperate regions, moderate high temperatures refer to heat

slightly above the optimal temperature for growth (at or around 35°C), while acute high

temperatures refer to heat at or above 40°C, which significantly reduces or completely

halts growth (Zhang *et al.*, 2022*a*). Moderate high temperatures are often long-lasting and

frequent in nature with mild effects on photosynthetic cells, while acute high temperatures

are often short-term but severely damaging. Most previous heat-stress experiments in

plants and algae used acute high temperatures at or above 40°C (Hemme *et al.*, 2014;

Rütgers *et al.*, 2017; Balfagón *et al.*, 2019; Kim *et al.*, 2020; Ji *et al.*, 2021), likely due to

the rapid onset and easily quantifiable phenotypes as compared to moderate high

temperatures. Although the impact of moderate high temperatures in photosynthetic cells

can be difficult to investigate due to comparatively mild phenotypes, moderate high

temperatures are physiologically relevant stresses in field conditions. Additionally, global

warming can further increase the frequency and duration of moderate high temperatures

in nature. The frequent and long-lasting features of moderate high temperatures could

reduce agricultural yield significantly (Delorge *et al.*, 2014; Anderson *et al.*, 2021).

Arabidopsis seedlings treated with 5-day heat at 35°C had reduced growth and viability

(Song *et al.*, 2021). Furthermore, long-term (4-6 days) moderate high temperatures

resulted in coral bleaching by causing cell death in endosymbiotic algae and their coral

hosts (Dunn *et al.*, 2004) and destabilizing the symbiotic nutrient cycling relationship

between these two organisms (Rädecker *et al.*, 2021). However, the effects of moderate

high temperatures in photosynthetic cells and the interaction of moderate high

temperatures with nutrient status especially carbon availability are underexplored. Algae

have great potential for production of biofuels and bioproducts (Scranton *et al.*, 2015;

Mathimani and Pugazhendhi, 2019). However, how algal cells respond to high
temperatures is under-investigated as compared to land plants (Schroda *et al.*, 2015). To
study the effects of carbon status, heat intensities and duration on heat responses in
photosynthetic cells, we utilized the unicellular green alga *Chlamydomonas reinhardtii*
(Chlamydomonas throughout).

Chlamydomonas is a superior model to study heat responses in photosynthetic cells
(Schroda *et al.*, 2015). Chlamydomonas can grow in light under photoautotrophic
conditions using photosynthesis (Minagawa and Tokutsu, 2015). It can also grow with
supplied organic carbon source (acetate) in light (mixotrophic) or dark (heterotrophic)
conditions (Sasso *et al.*, 2018), providing a platform to study heat responses under
different light and carbon conditions. Additionally, it has a haploid, sequenced, annotated,
small genome (111 Mb, 17,741 protein-encoding genes) with simpler gene families and
fewer gene duplications than land plants (Merchant *et al.*, 2007; Karpowicz *et al.*, 2011).
Furthermore, several well-established gene editing, cloning, high-throughput, and
functional genomic tools are available in Chlamydomonas (Shimogawara *et al.*, 1998; Li
et al., 2016, 2019; Greiner *et al.*, 2017; Crozet *et al.*, 2018; Wang *et al.*, 2019; Dhokane
et al., 2020; Emrich-Mills *et al.*, 2021).

Outdoor algal ponds frequently experience moderate high temperatures around 35°C
during the summer (Mata *et al.*, 2010; Krishnan *et al.*, 2021), but the effects of moderate
high temperatures on algal growth have been overlooked. Previously published algal heat
treatments were often conducted in flasks in pre-warmed water baths at or above 42°C
with sharp temperature switches (Hemme *et al.*, 2014; Rütgers *et al.*, 2017). While
previous research was highly valuable to understanding algal heat responses, high
temperatures in the field or outdoor ponds often increase gradually. Rapid increases to
high temperatures largely reduce algal viability (Zhang *et al.*, 2022*a*); similar effects have
also been reported in land plants (Mittler *et al.*, 2012). Acute high temperatures at or
above 40°C inhibit algal cell division (Mühlhaus *et al.*, 2011*b*; Hemme *et al.*, 2014;
Zachleder *et al.*, 2019; Ivanov *et al.*, 2021), while moderate high temperature at 35°C only
transiently inhibits cell division during the first 4-8 hour (h) heat (Zhang *et al.*, 2022*a*).

Thus, conducting long-term experiments at moderate high temperatures in flasks can

result in overgrown cultures, nutrient depletion, and light limitation, complicating data

interpretation.

Consequently, it is advantageous to investigate algal heat responses under highly-

controlled conditions in photobioreactors (PBRs) (Zhang *et al.*, 2022*a*; Mattoon *et al.*,

2022*a*), which have several evident strengths: (1) precise temperature regulation with a

sterile temperature probe inside each algal culture; (2) controlled heating to mimic the

heating speed in nature; (3) precisely controlled cultivation conditions, including

temperature, light, nutrient, and air agitation, allowing for reproducible experiments; (4)

automatic recording of culture status every minute, including growth conditions and

optical densities, enabling quantitative growth rate measurements; (5) the availability of

turbidostatic control based on defined parameters (e.g., chlorophyll contents) to enable

frequent culture dilutions with constant medium supply; (6) being able to simulate nutrient

depletion by turning off the turbidostatic control to investigate how nutrient availability

affects algal heat responses. Utilization of PBRs for algal cultivation and heat treatments

can largely reduce compounding effects during high temperature treatments and improve

our understanding of algal heat responses.

Recently, we conducted systems-wide analyses in wild-type (WT) Chlamydomonas

cultures during and after 24-h heat of 35°C and 40°C in PBRs under light in acetate-

containing medium (mixotrophic condition) with turbidostatic control and constant nutrient

supply (Zhang *et al.*, 2022*a*). We chose moderate high temperature of 35°C and acute

high temperature of 40°C based on the algal growth rates under different temperatures

and their relevance to physiologically high temperatures experienced in nature (Krishnan

et al., 2021). The 24-h heat duration allowed us to balance time-course sample collection

over a sufficient duration while maintaining culture viability (heat at 40°C for longer time

killed algal cells); our experimental conditions enabled us to investigate algal heat

responses during and after physiologically relevant high temperatures (Zhang *et al.*,

2022*a*). Our results showed that 40°C inhibited algal growth while 35°C increased algal

growth with constant acetate supply. The growth inhibition at 40°C could be explained by

reduced photosynthesis, impaired respiration, and arrested cell cycle, while these cell
parameters had minor changes in algal cultures treated by 35°C. Our proteomics data
indicated that several proteins involved in acetate uptake and assimilation, glyoxylate
cycle, and gluconeogenesis were up-regulated during 35°C (Zhang *et al.*, 2022*a*).
Chlamydomonas uptakes acetate and feeds it into the glyoxylate cycle and
gluconeogenesis for starch biosynthesis (Johnson and Alric, 2012, 2013). The main
function of glyoxylate cycle is assimilation of 2-carbon compounds (e.g., acetate) into
organic acids that can be metabolized by the cell; the glyoxylate cycle is a shunt of the
tricarboxylic acid (TCA) cycle in mitochondria without releasing CO_2, which allows for the
anabolism of simple carbon compounds in gluconeogenesis, a process to make sugars,
e.g., glucose (Johnson and Alric, 2012; Chew *et al.*, 2019; Walker *et al.*, 2021).

We hypothesized that the increased growth in Chlamydomonas at 35°C with constant
acetate supply was attributable to up-regulated acetate metabolism. In our previous
papers, we found out that 35°C increased the growth rates of Chlamydomonas cultures
with constant acetate supply (Zhang *et al.*, 2022*a*), but not in cultures without acetate
(Mattoon *et al.*, 2022*b*), as compared to the control at 25°C with the same medium type.
Therefore, we hypothesized that the effects of moderate high temperature at 35°C was
dynamic depending on the availability of acetate, e.g., constant-acetate, acetate-
depleting, or no-acetate conditions. The majority of published high temperature research
in Chlamydomonas has been conducted in acetate-containing medium (Voß *et al.*, 2010;
Mühlhaus *et al.*, 2011*a*; Hemme *et al.*, 2014; Rütgers *et al.*, 2017; Zhang *et al.*, 2022*a*),
but the interface between acetate supply and Chlamydomonas heat responses is
understudied. To validate our hypothesis and address these unknown questions, we
cultivated the same Chlamydomonas WT strain in PBRs as before but heated the cultures
at 35°C or 40°C with and without constant acetate supply for 1 or 4 days. Our previous
work was conducted at 35°C and 40°C for 24 h with constant acetate supply (Zhang *et*
al., 2022*a*). In this paper, to investigate the effects of carbon availability on algal
thermotolerance, we included 3 new different heat treatments at 35°C and 40°C: (1)
acetate-depleting condition for 24 h, (2) constant-acetate condition for 4 days, and (3) no-
acetate condition for 4 days. The last two conditions utilized turbidostatic control with

constant nutrient supply. Our new results confirmed our hypothesis about the role of
acetate metabolism in algal thermotolerance and revealed the overlooked effects of
moderate high temperature of 35°C on algal growth. Heat of 35°C was beneficial or
detrimental to Chlamydomonas cultures depending on acetate availability. Acute high
temperature of 40°C was lethal to Chlamydomonas cultures, with and without constant
acetate supply. Our research helps fill the knowledge gaps about how external carbon
supply affects algal heat responses, provides new insights to understand algal heat
responses, has potential applications in production of algal biofuel and bioproducts, and
contributes to improved thermotolerance in photosynthetic cells.

Materials and methods

Algal cultivation

The *Chlamydomonas reinhardtii* wild-type strain CC-1690 (also called *21gr*, mating type
plus) (Sager, 1955; Pröschold *et al.*, 2005; Zhang *et al.*, 2022*b*) was purchased from the
Chlamydomonas resource center and used in all experiments. Algal cultures were grown
in Tris-acetate-phosphate (TAP, with acetate) or Tris-phosphate (TP, without acetate)
medium with revised trace elements (Kropat *et al.*, 2011) in 400 mL photobioreactors
(PBRs) (Photon System Instruments, FMT 150/400-RB) as described previously (Zhang
et al., 2022*a*). Each medium type was used in individual PBRs for independent
experiments. Cultures were illuminated with constant 100 μmol photons m^{-2} s^{-1} light (red:
blue, 1:1 ratio), and agitated by bubbling with filtered air at a flow rate of 1 L min^{-1}. Algal
cultures with targeted cell density around 1-2 $\times 10^6$ cells mL^{-1} (~4-6 μg mL^{-1} chlorophyll
content) were maintained at 25°C for 4 days with constant nutrient supply through
turbidostatic control before different temperature treatments. The turbidostatic mode was
controlled by OD_{680} (optical density at 680 nm), which is proportional to chlorophyll
contents (μg mL^{-1}) and was monitored once per min automatically. When OD_{680} increased
to a maximum value slightly above the target value (for target cell density) due to algal
growth, OD_{680} signaled the control computer to turn on a turbidostatic pump to add fresh
medium to dilute the culture until a minimum OD_{680} slightly below the target value was
reached, then the turbidostatic pump was turned off automatically. Because of the small

OD range we used, the PBR cultures had exponential growth between dilution events
through the turbidostatic control with constant nutrient supply (Fig. 1A). The OD_{680} data
during exponential growth phases in between dilution events was log_2 transformed, and
the relative growth rate was calculated using the slope of $log_2(OD_{680})$. The relative growth
rate is the inverse of the algal culture doubling time. For the treatments without constant
nutrient supply (acetate-depleting condition), the turbidostatic pumps were turned off after
PBR cultures reached steady growth and at the start of different temperature treatments
(Fig. 1B).
225
High temperature treatments in PBRs
After algal cultures in PBRs reached steady growth with turbidostatic control at 25°C for
4 days, PBR temperatures were increased to moderate (35°C) or acute (40°C) high
temperatures in different PBRs for the indicated duration with indicated nutrient
conditions. PBR temperatures were increased from 25°C to 35°C or 40°C gradually over
the course of 30 min with controlled heating speeds. PBR cultures grown under constant
25°C with the same nutrient status and treatment duration served as controls. Algal
cultures were harvested from PBRs at different time points during different treatments for
various measurements.
235
236   **Algal biomass quantification**
237   Algal cultures (90 mL) were harvested from PBRs, centrifuged to remove supernatant,
238   flash frozen in liquid nitrogen, then stored in a -80°C freezer until analysis. Algal cell
239   pellets were freeze dried for 24 h over liquid nitrogen. The remaining algal dry biomass
240   were weighed and quantified.
241
242   **Acetate quantification assay**
243   Algal cultures were harvested from PBRs (2 mL) and centrifuged to collect 500 µL top
244   clear supernatant to a new tube. The supernatant was stored in a -80°C freezer until
245   analysis. The acetate content in the supernatant was quantified using the Acetate
246   Colorimetric Assay Kit (Sigma, Cat No. MAK086) according to manufacture instructions.
247

Cell imaging using light microscopy

Chlamydomonas cultures harvested at select time points with different temperature treatments were fixed with 0.2% glutaraldehyde (VWR, Cat No. 76177-346). Algal cells were imaged with a Leica DMI6000 B microscope and a 63x (NA1.4) oil-immersion objective.

Chlorophyll quantification

Chlorophyll contents in algal cells were quantified as previously described (Zhang *et al.*, 2022*a*). PBR cultures of 1 mL were harvested in 1.5-mL tubes with 2.5 µL 2% Tween20 (Sigma, P9416-100ML, to help cell pelleting), and centrifuged at 18,407 g at 4°C. After removing supernatant, cell pellets were stored in a -80°C freezer until quantification. Later, cell pellets were thawed, resuspended in 1 mL of HPLC grade methanol (100%, Sigma, 34860-4L-R), vortexed for 1 min, incubated in the dark at 4°C for 5 min, and centrifuged at 15,000 g at 4°C for 5 min. Supernatant was pipetted out to a two-sided disposable plastic cuvettes (VWR, 97000-586) for chlorophyll (Chl) quantification at 652 nm and 665 nm in a spectrophotometer (IMPLEN Nonophotometer P300) using the following equations: Chl a + Chl b = $22.12*A_{652} + 2.71*A_{665}$ (in µg mL^{-1} algal cultures) (Zhang *et al.*, 2022*a*). Chlorophyll concentrations were also normalized to cell densities (Chl pg cell^{-1}) or mean cell volume (Chl pg µm^{-3}). Cell density and mean cell volume were measured using a Coulter Counter (Multisizer 3, Beckman Counter, Brea, CA). For 35°C with photoautotrophic medium, algal cells formed palmelloids, clumps of non-motile cells, thus the size of palmelloids instead of cell size was measured by the Coulter Counter.

RNA extraction

PBR cultures of 2 mL were pelleted with Tween-20 (0.005%, v/v) by centrifugation at 11,363 x g and 4°C for 2 min, followed by supernatant removal. The cell pellet was flash frozen in liquid nitrogen and then stored in a -80°C freezer until processing. RNA extraction and RT-qPCR analysis were performed as previously described with minor modifications (Zhang *et al.*, 2022*a*). Total RNA was extracted with TRIzol reagent (Thermo Fisher Scientific, Cat No. 15596026), digested on-column with RNase-free

DNase (Qiagen, Cat No. 79256), purified by RNeasy mini-column (Qiagen, Cat No.
74106), and quantified with Qubit™ RNA BR Assay Kit, (Life technology, Cat No.
Q10210).

RT-qPCR analysis

Total 0.4 µg RNA was reverse transcribed with oligo dT primers using SuperScript® III
First-Strand Synthesis System (Life technology, Cat No. 18080-051) following
manufacturer instructions. Quantitative real-time PCR (RT-qPCR) analysis was
performed using a CFX384 Real-Time System (C 1000 Touch Thermal Cycler, Bio-Rad,
Hercules, California) using SensiFAST SYBR No-ROS kit (Bioline, BIO-98020) following
this program: (1) $95^{o}C$ (2 min); (2) 40 cycles of $95^{o}C$ (5 s), $60^{o}C$ (10 s), and $72^{o}C$ (15 s);
(3) final melt curve at 60°C (5 s), followed by continuous temperature ramping from 60°C
to 99°C at a rate of $0.5°C\ s^{-1}$. Melting curves and qPCR products were tested to ensure
there were no primer dimers or unspecific PCR products. All RT-qPCR products were
sequenced for confirmation. Primers and gene IDs for RT-qPCR can be found in
Supplementary Table 1. *CBLP* (*β-subunit-like polypeptide*, Cre06.g278222) (Schloss,
1990; Xie *et al.*, 2013) and *EIF1A* (*Eukaryotic translation initiation factor 1A,*
Cre02.g103550) (Strenkert *et al.*, 2019) had stable expression among all time points, and
were used as reference genes for RT-qPCR normalization. The relative gene expressions
were calculated relative to the pre-heat time point by using the $2^{-\Delta\Delta CT}$ method (Livak and
Schmittgen, 2001; Hellemans *et al.*, 2007; Remans *et al.*, 2014). Three biological
replicates were included for each time point and treatment. RT-qPCR was used to verify
some selected transcripts and compared with the results of RNA-seq for samples
harvested under the acetate-depleting conditions (Supplemental Fig. 2). RNA-seq for
samples harvested under the constant-acetate conditions were verified by RT-qPCR
previously (Zhang *et al.*, 2022*a*).

RNA-seq analysis

Raw sequencing data for the constant-acetate experiments at 35°C and 40°C were
obtained from Zhang *et al.*, 2022*a*. Raw sequencing data for the constant-acetate

experiment at 25°C and acetate-depleting experiments at 25°C, 35°C, and 40°C were

collected in this study. All raw sequencing data were processed using the same pipeline

as described below.

RNA libraries were prepared and sequenced by the Joint Genome Institute (JGI,

Community Science Program) using the NovaSeq platform generating 151-nt paired-end

reads. Plate-based RNA sample preparation was performed on the PerkinElmer Sciclone

NGS robotic liquid handling system using Illumina's TruSeq Stranded mRNA HT sample

prep kit utilizing poly-A selection of mRNA following the protocol outlined by Illumina in

their user guide, with the following conditions: total RNA starting material was 1 µg per

sample and 8 cycles of PCR was used for library amplification. The prepared libraries

were quantified using KAPA Biosystems' next-generation sequencing library qPCR kit

and ran on a Roche LightCycler 480 real-time PCR instrument. Sequencing of the flowcell

was performed on the Illumina NovaSeq sequencer using NovaSeq XP V1.5 reagent kits,

S4 flowcell, following a 2x151 indexed run recipe.

Samples were quality control filtered using the JGI BBDuk and BBMap pipelines (BBDuk,

2022). Samples were quality assessed using FastQC (Andrews) and mapped to the

Chlamydomonas reinhardtii v5.6 genome (Merchant *et al.*, 2007) using HISAT2 version

2.2.0 (Kim *et al.*, 2015). Reads per feature were counted via featureCounts (Liao *et al.*,

2014). The count matrix was combined with existing data (Zhang *et al.*, 2022*a*), resulting

in a 16,403x101 count matrix (Supplemental Dataset 4, 5). The normalization of the count

matrix was conducted using the median of ratios method (Liao *et al.*, 2014). Prior to

imputation of two missing time points in the constant-acetate-25°C condition (see details

below), transcripts were filtered to have nonzero counts in at least 90% of the samples.

Under constant 25°C, acetate, and light with turbidostatic control (constant-acetate-25°C),

the algal growth rate and cell physiology stays constant (Fig. 5). We had RNA-seq data

for 8 time points (pre-heat, 0, 0.5, 1, 2, 24, 26, and 48 h) under constant-acetate-25°C

and these samples were highly correlated with each other (with Pearson correlation ≥

0.993) (Supplemental Fig. 1). RNA samples for the time points of 4 and 8 h under the

constant-acetate-25ºC condition were unavailable; thus RNA-seq data for the 4 and 8 h
under the constant-acetate-25ºC condition were imputed from other extra time points (0,
0.5, 1, 26, 48 h) of the same condition. For the time points of 4 and 8 h under the constant-
acetate-25ºC condition, each transcript triplicates were sampled randomly using the 5
extra time points under the same condition mentioned above. It was ensured that both
missing time points were not imputed with the same count data. The final count matrix
consists of 14,893 transcripts in 90 samples (5 time points measured at 6 conditions as
triplicates). The natural logarithm was determined for all normalized counts before
averaging triplicates. Pearson correlation coefficients were calculated for all sample pairs.
348
The count data were tested for differential expression using DESeq2 v1.38.3 (Love *et al.*,
2014). DESeq2 applies a Benjamini Hochberg correction for multiple testing. The design
matrix was constructed to test for global kinetic differences between the three different
temperatures and both growth media over time. The following tests were performed
(Supplemental Dataset 1, 3, 5, Supplemental Fig. 5): (A) acetate-depleting 25°C vs
acetate-depleting 35°C; (B) acetate-depleting 25°C vs acetate-depleting 40°C; (C)
constant-acetate 25°C vs constant-acetate 35°C; (D) constant-acetate 25°C vs constant-
acetate 40°C; (E) temperature effect: (constant-acetate 35°C and acetate-depleting 35°C)
vs (constant-acetate 40°C and acetate-depleting 40°C); (F) medium effect: (constant-
acetate 35°C and constant-acetate 40°C) vs (acetate-depleting 35°C and acetate-
depleting 40°C); (G) acetate-depleting 25°C vs constant-acetate 25°C.
360
Functional descriptions were determined for each transcript (Merchant *et al.*, 2007;
Usadel *et al.*, 2009; Venn and Mühlhaus, 2022). Transcripts were grouped according to
their functional description. Time courses of normalized transcript counts were visualized
as $\log_2$(fold change). An ANOVA was performed for each transcript at each treatment to
elucidate whether a transcript underwent a relevant change during its time course (Venn
et al., 2022a). Heatmaps (Supplemental Dataset 1) and functional sets figures
(Supplemental Dataset 2) were created using Plotly.NETv4.0.0 (Schneider *et al.*, 2022).
368

Ontology enrichment was performed using extended MapMan annotations (for example

"PS.lightreaction.LHC" became "PS.Lightreaction.LHC", "PS.Lightreaction" and "PS")

(Supplemental Dataset 3). The measured and filtered transcripts served as background.

If a transcript showed significant differential expression in the respective comparison

(FDR < 0.05), it was considered as significant for its annotation. Enrichment p values

were determined using hypergeometric tests. Multiple testing correction was performed

using the Benjamini-Hochberg method (Benjamini and Hochberg, 1995; Venn *et al.*,

2022). Additional ontology enrichment was performed using ChlamyCyc annotations

(v2023-01-04). This pathway ontology combines KEGG, MapMan, and JGI pathway

information (May *et al.*, 2009). The enrichment procedure and statistical analysis were

the same as above.

Results

To investigate how the availability of an organic carbon source, acetate, affected algal

heat responses, we first cultivated WT Chlamydomonas cells (CC-1690, *21gr*) in PBRs

in Tris-acetate-phosphate (TAP, acetate as an organic carbon source) medium at 25^{o}C

with constant nutrient supply through turbidostatic control (providing frequent fresh

medium for culture dilution) (Fig. 1A). After algal cultures reached steady growth rates in

PBRs, the turbidostatic control was turned off and the temperatures of algal cultures were

increased to 35^{o}C or 40^{o}C or stayed at 25^{o}C for 24 h without constant acetate supply,

defined as the acetate-depleting condition. This experiment allowed us to quantify acetate

consumption under different temperatures, which could not be performed in cultures with

turbidostatic control and frequent medium dilution. It also enabled us to investigate algal

heat responses under the acetate-depleting condition, like shaker cultures, which were

used by most previous algal heat experiments in published literatures (Hemme *et al.*,

2014; Rütgers *et al.*, 2017). Under the acetate-depleting condition, algal cultures were

harvested at different time points to analyze cell physiology, biomass, and transcriptome

(Fig. 1B). Algal dry biomass quantification showed that cultures treated with 35^{o}C had

increased biomass at 8-h heat but decreased biomass at 24-h heat as compared to 25^{o}C

(Fig. 1C). Acetate quantification in the supernatant of algal cultures showed that 35^{o}C-

treated cultures had increased acetate consumption and depleted acetate faster than

25°C or 40°C (Fig. 1D).

We hypothesized the transiently increased and then decreased algal biomass in 35°C-

treated cultures was due to increased acetate uptake and usage which led to acetate

starvation at 24 h of the acetate-depleting condition (Fig. 1). Chlamydomonas uptakes

acetate and feeds it into the glyoxylate and gluconeogenesis cycles to produce starch,

which can be used in glycolysis to generate ATP (Johnson and Alric, 2012, 2013) (Fig.

2A). To verify this hypothesis, we performed RNA-seq analysis and investigated key

transcripts involved in acetate uptake and assimilation, glyoxylate, gluconeogenesis, and

glycolysis cycles under the acetate-depleting condition by comparing with the constant-

acetate condition (Fig. 2B-P). Under the acetate-depleting condition, almost all these

transcripts related to acetate uptake and metabolism were up-regulated under 35°C as

compared to 25°C, especially the early heat time points of 2 and 4 h at 35°C. Similar 35°C

effects on these transcripts were held under the constant-acetate condition, although the

up-regulation of these transcripts were often stronger and lasted longer at 35°C under the

constant-acetate condition than the acetate-depleting condition. In contrast, these

transcripts related to acetate uptake and metabolism were down-regulated under 40°C

as compared to 25°C under both the acetate-depleting and constant-acetate conditions,

especially the 2-h heat time point at 40°C. Our RT-qPCR results of selected transcripts

showed similar trends as our RNA-seq data (Supplemental Fig. 2). These results showed

that 35°C promoted acetate metabolism, from acetate uptake and assimilation to

glyoxylate, gluconeogenesis, and glycolysis cycles; however, these acetate metabolism

pathways were reduced under 40°C.

In addition to the acetate metabolism pathways mentioned above, we investigated

transcripts related to other processes that are often affected by high temperatures

(Supplemental Fig. 3, 4). Chlamydomonas has two heat shock transcription factors, HSF1

and HSF2; HSF1 is the master heat regulator but the function of HSF2 is unclear (Schulz-

Raffelt *et al.*, 2007). The transcripts of both *HSF*s were induced by heat, with higher

induction at 40°C than 35°C under the same acetate condition and higher induction under

the acetate-depleting than constant-acetate condition under the same high temperature

(Supplemental Fig. 3A, B). Heat shock proteins (HSPs) are the hallmarks of heat

responses and function as chaperones to help refold denatured proteins, fold nascent

polypeptides, and prevent denatured protein aggregates (Altschuler and Mascarenhas,

1982; Schroda *et al.*, 2015; Haq *et al.*, 2019). Based on molecular mass, HSPs are

grouped into large HSPs (HSP100/90/70/60, >50 kDa) and small HSPs (10-43 kDa) (Jee,

2016; Bourgine and Guihur, 2021). Under the acetate-depleting condition, many

transcripts related to large HSPs, *HSP60/70/90s*, were down-regulated at 24 h of 25°C

and 35°C; such down-regulation of *HSP60/70/90s* was absent at 24 h of 25°C and 35°C

under the constant-acetate condition or 24 h of 40°C under both acetate conditions

(Supplemental Fig. 4A-C). Additionally, transcripts of *HSP100* and most of the small

HSPs did not have such down-regulation at 24 h of 25°C and 35°C under the acetate-

depleting condition (Supplemental Fig. 4D, E). Almost all transcripts related to the *HSPs*

we examined, including *HSP60/70/90/100s* and small *HSPs*, were up-regulated at 2 h of

40°C, with similar induction fold change under either the acetate-depleting or constant-

acetate condition (Supplemental Fig. 4A-E). This suggested the early induction of these

HSPs under 40°C was temperature-dependent (present at 40°C while much less so under

35°C) but acetate-availability-independent. Alternatively, the acetate-availability might not

differ significantly at 2 h of the acetate-depleting and constant-acetate condition (Fig. 1D).

However, the absence of down-regulated *HSP* transcripts at 24 h of 40°C under the

acetate-depleting condition supported that the induced *HSP* transcripts at 40°C was

independent of acetate.

The global transcriptome changes related to light reaction and Calvin-Benson cycle of

photosynthesis suggested more disturbance of these transcripts under the acetate-

depleting than the constant-acetate conditions at the same temperatures, and more

disturbance of these transcripts at 40°C than 35°C under the same acetate conditions

(Supplemental Fig. 4F-M). These photosynthesis-related transcripts behaved similarly at

40°C under both the acetate-depleting and constant-acetate conditions, suggesting 40°C

heat effects were more dominant than the changes of acetate availability.

Chlamydomonas employs a cellular structure, pyrenoid, to concentrate CO_2 for improved

carbon fixation efficiency, called carbon concentrating mechanisms (CCM) (Meyer *et al.*,
2017; Hennacy and Jonikas, 2020). The CCM is especially important under high
temperature conditions, where the solubility of CO_2 decreases in water by approximately
30% from 25°C to 40°C (Dodds *et al.*, 1956). Under the acetate-depleting conditions,
several transcripts related to CCM were up-regulated at 24 h under 25°C and 35°C, but
not under the same temperature of the constant-acetate condition (Supplemental Fig.
4N). This suggested that as acetate became depleted in the medium and cell density
increased, cells relied more on the photosynthetic carbon fixation by up-regulating CCM.
Such induction of the CCM transcripts was much less at 40°C under the acetate-depleting
condition, which may correlate with the compromised pyrenoid structures with decreased
appearance of thylakoid tubules at 40°C in our previous publication (Zhang *et al.*, 2022*a*).
Thylakoid tubules deliver concentrated CO_2 to the pyrenoid and also serve as the diffusion
path of Calvin-Benson Cycle metabolites (Hennacy and Jonikas, 2020). Transcripts
related to photoprotection had larger induction at 24 h under all three temperatures of the
acetate-depleting than the constant-acetate conditions at the same temperature,
indicating compromised photosynthesis and the need for photoprotection when acetate
was depleted (Supplemental Fig. 4O).

The compromised photosynthesis under high temperatures lead to increased
accumulation of reactive oxygen species (ROS) (Pospíšil, 2016; Janni *et al.*, 2020;
Niemeyer *et al.*, 2021). Some transcripts related to ROS scavenging were down-regulated
at 24 h of 25°C and 35°C under the acetate-depleting condition but much less so under
the constant-acetate condition at the same temperatures (Supplemental Fig. 4R-V). This
may be related to the increased cell density and reduced light and oxidative stress at 24
h of 25°C and 35°C under the acetate-depleting condition. Such down-regulation of
transcripts related to ROS scavenging was not present at 24 h of 40°C under the acetate-
depleting condition. This may suggest relatively increased ROS accumulation and ROS
scavenging activity at 40°C than 25°C and 35°C under the acetate-depleting conditions.
Additionally, transcripts related to ROS scavenging had very similar transcriptional
changes at 40°C under either the acetate-depleting or constant-acetate condition,

suggesting 40°C had more dominant effects on the ROS accumulation and ROS
scavenging activities than acetate.

Transcripts related to cell cycle and cell wall had similar transcriptional patterns under
each of the six conditions (Supplemental Fig. 4W, X). Under diurnal cycles, cell wall genes
were up-regulated following the up-regulation of cell cycle genes and cell walls form
around daughter cells during cell division (Zones *et al.*, 2015). These transcripts were
down-regulated at 24 h of 25°C under the acetate-depleting condition, suggesting the
transition to reduced cell division due to the acetate-depletion. The reduction of these
transcripts was larger and more prolonged under 40°C than 35°C, consistent with
complete and transient inhibition of cell cycle under 40°C and 35°C, respectively (Hemme
et al., 2014; Zhang *et al.*, 2022*a*). At 24 h of 40°C and 35°C, these transcripts were more
reduced under the acetate-depleting condition than the constant-acetate condition. It is
worthwhile to note that the changes during 24 h of the acetate-depletion condition were
dynamic and here we just mentioned the changes at 24 h in the end when acetate was
depleted.

Almost all transcripts mentioned above had little changes at 25°C during all time points
of the constant-acetate condition, validating the steady growth and constant cell status
under the turbidostatic control with constant nutrient and light but without heat treatments
in our algal cultures grown in PBRs (Supplemental Fig. 4).

We performed statistical analysis to investigate global transcript kinetic differences
between different conditions (Supplemental Dataset 1, 3). We then applied ontology
enrichments analysis for transcripts that were significantly expressed in a defined
comparison to identify differentially regulated biological processes (Supplemental Fig. 5,
Supplemental Dataset 3). The top enriched function group that differed at 25°C and 35°C
under both acetate-depleting and constant-acetate condition was the nuclear-encoded
ribosomal proteins that are involved in chloroplast protein synthesis (Supplemental Fig.
5A, C). More than half of the ribosomal related transcripts were down-regulated at 4 h
heat of 35°C as compared to 25°C under both acetate conditions (Supplemental Fig. 4Y,

Z), suggesting 35°C may inhibit chloroplast protein synthesis. The down-regulation of

ribosomal related transcripts at 35°C could recover partially under the constant-acetate

condition, but much less so under the acetate-depleting condition. The top enriched

function group that differs between 25°C and 40°C under the acetate-depleting condition

was the mitochondrial electron transport (Supplemental Fig. 5B). Many transcripts related

to mitochondrial electron transport were down-regulated at 2 h heat of 40°C as compared

to 25°C under the acetate-depleting condition, especially these related to ATP synthesis

(Supplemental Fig. 4AA, AB). When comparing 35°C and 40°C conditions independent

of the acetate availability, the top enriched function group was also the mitochondrial

electron transport (Supplemental Fig. 5E). The results suggested that mitochondrial

electron transport was more sensitive to 40°C than 25°C and 35°C. Under the constant-

acetate condition, 40°C affected transcripts related to RNA processing the most as

compared to 25°C (Supplemental Fig. 5D, 4AC, AD). When comparing the acetate-

depleting and constant-acetate conditions independent of temperatures, the top two

enriched groups were phosphate transport and mitogen-activated protein kinase (MAP

kinase) cascades signaling (Supplemental Fig. 5F). MAP kinase signaling pathways have

important roles in plant growth and stress responses and involve protein phosphorylation

and dephosphorylation (Bigeard and Hirt, 2018; Jagodzik *et al.*, 2018; Zhang and Zhang,

2022). The biggest differences in transcripts related to the MAP kinase signaling

pathways between the acetate-depleting and constant-acetate conditions were at the 24

h time point at 25°C and 35°C when the acetate was depleted (Supplemental Fig. 4AF),

suggesting potential roles of the MAP kinase signaling pathways in nutrient sensing and

growth regulation. When comparing the acetate-depleting and constant-acetate

conditions at 25°C, the top enriched functional groups are dominantly related to cell

motility (Supplemental Fig. 5G). More than half of the transcripts related to cell motility at

25°C were down-regulated at 24 h under the acetate-depleting condition but not the

constant-acetate condition (Supplemental Fig. 4AG, AH). Microscopic images showed

that cells had cilia at 24 h of 25°C under the acetate-depleting condition (Supplemental

Fig. 6). Cell motility needs energy and ciliary beating frequency corelates with the

cytosolic ATP concentration in Chlamydomonas (Takano *et al.*, 2021). Thus, the down-

regulation of transcripts related to cell motility may suggest the reduced cellular ATP level

and the transition from motile to non-motile phase when the acetate was depleted at 24
h of the acetate-depleting condition.

Besides MapMan ontology enrichment, we performed ChlamyCyc pathway enrichment
analysis of differentially regulated transcripts compared between two conditions
(Supplemental Dataset 6). The ChlamyCyc annotations (v2023-01-04) and pathway
ontology combine KEGG, MapMan, and JGI pathway information (May et al. 2009). The
KEGG pathway annotation for Chlamydomonas is sparse, therefore it was difficult to
determine significances. As comparison, for the MapMan functional ontology we got
23,500 annotations in the RNA-seq data, for KEGG pathways we only got 803. Both
analyses showed the mitochondrial activity was the top enrichment for the comparisons
of 25^oC versus 40^oC under the acetate-depleting condition and 35^oC versus 40^oC under
either acetate condition (temperature effects) (Supplemental Fig. 5B, E, Supplemental
Dataset 6).

To further understand our transcriptome data and the effects of acetate, we compared
cell physiology at different temperatures between the acetate-depleting and constant-
acetate conditions (Fig. 3). Under the acetate-depleting condition without medium dilution,
cell density increased over time at 25^oC and 35^oC but no changes at 40^oC because of
cell cycle arrest at 40^oC (Fig. 3A). Cell density was lower at 35^oC than 25^oC at 8 h and 24
h time points, suggesting reduced growth rates at 35^oC at the end of the acetate-depleting
condition. Algal cells treated at 35^oC had transiently increased then decreased cell size
while 40^oC-treated algal cells had steadily increased cell size due to heat-inhibited cell
division (Fig. 3B). Under the acetate-depleting condition, chlorophyll content per culture
volume (mL) had no significant differences between 25^oC and 40^oC but was significantly
higher at 8-h of 35^oC than 25^oC (Fig. 3C), consistent with the increased biomass at the
8-h of 35^oC (Fig. 1C). Chlorophyll content normalized to cell density revealed constant
chlorophyll per cell during 25^oC, but transiently and constantly increased cellular
chlorophyll during 35^oC and 40^oC, respectively (Fig. 3D). Chlorophyll content normalized
to cell volume showed the increased cellular chlorophyll during 40^oC and 35^oC could not
be completely explained by increased cell volume (Fig. 3E).

Based on our previously published data, we summarized cell parameters in algal cultures

grown in the same PBRs but under the 24-h constant-acetate condition through

turbidostatic control (Zhang *et al.*, 2022*a*) (Fig. 3F-J). Cell parameters at different time

points during 25°C had little or no changes as compared to the pre-heat time points,

demonstrating the steady algal growth and effectiveness of the turbidostatic control with

constant acetate supply. Under the constant acetate condition, the smaller changes of

cell density and chlorophyll per mL culture as compared to the acetate-depleting

conditions were due to the turbidostatic control and frequent dilution of the culture (Fig.

3F and 3H). The changes of cell volume and chlorophyll content per cell during 35°C and

40°C had similar trends under both the acetate-depleting and constant-acetate condition,

but the increase of these parameters was larger under 40°C in cultures with the constant-

acetate than the acetate-depleting condition (Fig. 3B, D, G, I). This was supported by the

fold changes of cell parameters by comparing the data under the acetate-depleting and

constant-acetate conditions (Fig. 3K-O).

In addition to the cellular parameters mentioned above using time-course harvesting, we

next utilized non-disruptive and automatic methods to quantify algal growth under

different temperatures. The acetate-depleting condition without turbidostatic control

allowed us to have a large range of cell densities under different temperature treatments

to investigate the optical density (OD) vs cell density relationship and validate automatic

culture monitoring under different temperature treatments. OD_{680} (optical density at 680

nm) monitors chlorophyll content per mL culture (Chapman *et al.*, 2015; Xiao *et al.*, 2015;

Young *et al.*, 2022). OD_{750} (optical density at 750 nm) monitors light scattering and is

thought to be proportional to cell density or cell volume (Chioccioli *et al.*, 2014; Young *et

al.*, 2022). Our PBRs can measure OD_{680} and OD_{720}, but not OD_{750}. However, OD_{720}

serves as a proxy for OD_{750} for light scattering. Thus, we investigated the dynamic

changes of OD_{680} and OD_{720} relative to chlorophyll content, cell density, and cell volume

before and during heat treatments under the acetate-depleting condition (Fig. 4). The

change of OD_{680} and OD_{720} mimicked the change of chlorophyll content per mL algal

culture for all three temperature treatments (Fig. 4A, B, 3C). But the change of OD_{720}

(Fig. 4B) mimicked the change of cell density for 25°C and 35°C but not 40°C (Fig. 3A).

Combining all data from different temperature treatments and time points, both OD_{680} and

OD_{720} were linearly proportional to chlorophyll content per mL algal culture (Fig. 4C, D),

but they were much less proportional to cell volume or cell density (Fig. 4E-H). Our results

showed that OD_{680} and OD_{720} can be used to accurately estimate chlorophyll content in

algal cultures with different heat treatments under a wide range of cell densities. Thus,

OD_{680} and OD_{720} can be used to estimate algal relative growth rates based on chlorophyll

accumulation during an exponential growth phase between two dilution events under the

turbidostatic control in PBRs (Zhang *et al.*, 2022*a*) (See methods for details). The relative

growth rates calculated from both OD_{680} and OD_{720} yielded similar results. Because OD_{680}

had larger values and higher signal/noise ratios than OD_{720}, we used OD_{680} to estimate

relative growth rates of PBR cultures over several days under the turbidostatic controls

with and without acetate supply to investigate how acetate affected algal heat responses

at 35°C and 40°C in long-term (Fig. 5-8). This research goal could not be achieved without

turbidostatic control because our acetate-depleting experiment revealed that acetate was

fully depleted in 24 h during all three temperature treatments of 25°C, 35°C and 40°C

without constant acetate supply (Fig. 1D).

To investigate the effects of acetate on long-term heating, after algal cultures reached

steady growth at 25°C in PBRs with turbidostatic control, we conducted all three

temperature treatments with continuous turbidostatic control and constant acetate supply

for 4 days (Fig. 5). Without heat treatments, algal growth rates and cell parameters stayed

constant at 25°C, demonstrating the effectiveness of our turbidostatic control for steady

algal growth (Fig. 5A). At 35°C with constant acetate supply, the relative growth rates

increased first, but the increase was smaller after 2-day heating at 35°C and stabilized

thereafter. At 40°C, the relative growth rates decreased steadily, and the cultures died

after 2-day heating at 40°C, suggesting algal cells could not acclimate to long-term

constant heat at 40°C, even with constant acetate supply. Cells treated with 40°C had

more than 4-fold increase of cell volume as compared to the pre-heat condition, followed

by reduced cellular chlorophyll and cell death (Fig. 5C-F). The turbidostatic mode by

OD_{680} tightly controlled the chlorophyll contents in unit of ug per mL culture in PBRs during

the treatments of 25°C and 35°C (Fig. 5D), but not 40°C due to the cell cycle arrest and

overwhelmingly increased chlorophyll per cell, and eventually cell death under 40°C (Fig.

5B and E).

To further test the effects of acetate on algal heat responses, we conducted experiments

using algal cultures grown in PBRs under the turbidostatic control with constant nutrient

supply of photoautotrophic medium (no acetate) and different temperature treatments for

4 days (Fig. 6, 7). Our microscopic images showed that 35°C-treated cells in

photoautotrophic medium formed palmelloids (Fig. 6). Palmelloids are clumps of non-

motile cells, which often result from a failure of hatching or release from the mother cell

wall under stressful conditions (Harris, 2009). It was reported that Chlamydomonas cells

aggregated to form palmelloids after 3 min acute heat shock at 50°C in acetate-containing

medium (de Carpentier *et al.*, 2022). The formation of palmelloids did not occur in our

constant-acetate-35°C condition (Zhang *et al.*, 2022*a*), suggesting 35°C was more

stressful without acetate than with acetate. Without heat treatments, algal growth rates

and cell parameters stayed constant at 25°C, demonstrating the effectiveness of our

turbidostatic control for steady algal growth under photoautotrophic conditions (Fig. 7A).

Without acetate but with constant medium supply, algal growth had gradually decreased

growth rates during 35°C (Fig. 7A), unlike the increased growth rates during 35°C with

constant acetate supply (Fig. 5A), strongly supporting the role of acetate in algal heat

tolerance. Without acetate, algal cells with 35°C treatment had increased palmelloid

volume (or size) and chlorophyll contents (Fig. 7B-F). The reduction of growth rates during

40°C was comparable with and without acetate (Fig. 5A and 7A).

Our PBRs could also monitor photosystem II (PSII) operating efficiency using chlorophyll

fluorescence measurement automatically (Fig. 8). Heat at 40°C reduces PSII operating

efficiency dramatically, with more reduction without acetate than with acetate during the

first two days of heating (Fig. 8B, C, D). The acetate effects on PSII operating efficiency

disappeared in algal cells treated with 40°C for more than 2 days (Fig. 8D). The reduction

of PSII operating efficiency during 35°C was much smaller than during 40°C, with higher

PSII operating efficiency in 35°C-treated algal cells with acetate than without acetate (Fig.
8D).

Discussion

By performing algal cultivation under highly controlled conditions in PBRs with and without
the constant organic carbon source, acetate, we investigated how acetate availability
affected the growth of Chlamydomonas under moderate (35°C) and acute high
temperatures (40°C) (Fig. 9). The moderate high temperature of 35°C is frequently
experienced by outdoor, open algal ponds in summertime and enclosed outdoor algal
ponds can often heat up to 40°C or above (Krishnan *et al.*, 2021). Thus, both high
temperatures we chose were physiologically relevant for algae in nature. Additionally,
global warming increases the duration of moderate and acute high temperatures in the
field conditions. By using highly controlled turbidostatic conditions in PBRs, our research
not only investigated the short-term (within 24 h) but also long-term (up to 4-days) heat
effects on algal growth with and without acetate supply or under the acetate-depleting
conditions. Our research advanced the understanding of algal heat responses under
different heat intensities, heat durations, and carbon availability.

Heat of 35°C was beneficial or detrimental depending on carbon availability

Under the turbidostatic control with frequent medium dilutions, moderate high
temperature of 35°C increased algal growth rates with constant acetate supply (Fig. 5A)
but decreased algal growth rates without acetate in photoautotrophic medium (Fig. 7A),
confirming the important role of acetate in heat tolerance. The increased heat tolerance
at 35°C with acetate is not strain-dependent because the acetate effects were similar in
two different algal strains: CC-1690 (this study) and CC-5325 (Mattoon *et al.*, 2022*a*). In
medium starting with acetate but without constant acetate supply (the acetate-depleting
condition), algal biomass increased first but then decreased at 35°C compared to 25°C,
which can be explained by increased acetate uptake and usage initially followed by
acetate depletion and starvation by the end of the 24-h 35°C treatments (Fig. 1C, D).

Under the acetate-depleting condition without medium dilution, besides acetate, other
nutrients may also be depleted gradually during 24 h. We checked known transcription
factors and regulators responding to starvation of other nutrients, including *nitrogen*
starvation responsive regulator 1 (NRR1) (Boyle *et al.*, 2012), *sulfur starvation responsive*
regulator 2.1 (SNRK2.1) (González-Ballester *et al.*, 2010), *phosphorus starvation*
response 1 (PSR1) (Bajhaiya *et al.*, 2016), *copper deficiency response regulator 1*
(CRR1) (Sommer *et al.*, 2010), and *iron deficiency response regulator (Mitogen-Activated*
Protein Kinase 3, MAPK3) (Fei *et al.*, 2017). If these nutrients were depleted, the
corresponding transcription factors and regulators would be up-regulated. Most of these
nutrient-deficiency transcription factors and regulators had minimal or no induction at 24
h of the acetate-depleting condition (Supplemental Fig. 3C-G). One exception is the
NRR1, which was up-regulated at 25°C or 35°C but was down-regulated under 40°C at
24 h of the acetate-depleting condition (Supplemental Fig. 3C). The results suggested
minimal or no limitation of sulfur, phosphorus, copper, and iron during the acetate-
depleting condition, but possibly some nitrogen deficiency at 24 h of the acetate-depleting
condition at 25°C and 35°C but not 40°C. One of the responses to nitrogen deficiency is
the reduction of chlorophyll (Schmollinger *et al.,* 2014). During our acetate-depleting
condition, most transcripts involved in chlorophyll biosynthesis were stable (Supplemental
Fig. 4Q). Additionally, there was no change of the chlorophyll per cell or per cell volume
at 24 h of 25°C as compared to the pre-heat time point which did not have acetate-
limitation; these parameters were higher under 35°C and 40°C than 25°C at 24 h of the
acetate-depleting condition (Fig. 3D, E). Collectively, the results indicated that the
nitrogen limitation may have just started at 24 h of the acetate-depleting condition, thus
would not majorly impact cell physiology. Finally, our results from the 4-day constant-
acetate and no-acetate conditions under the turbidostatic control strongly pointed to the
specific effects of acetate on algal heat tolerance but not other nutrients (Fig. 5, 7). Thus,
the acetate effects are the focus of this paper.

Chlamydomonas uptakes acetate and feeds it into the glyoxylate cycle and
gluconeogenesis for starch biosynthesis and starch is broken down through glycolysis to
make cellular energy (Johnson and Alric, 2012, 2013) (Fig. 2A). The dynamic effects of

35°C on algal growth under different acetate conditions can be explained by the 35°C-
induced acetate metabolism, supported by increased acetate consumption and up-
regulated transcripts related to acetate metabolism pathway at 35°C (Fig. 1, 2). Energy
produced from glycolysis can be used for energy-requiring cellular activities to increase
thermotolerance (Olas *et al.*, 2021), e.g., repair pathways related to photosynthesis
(Murata and Nishiyama, 2018). Pyruvate kinase catalyzes the final step of glycolysis by
converting phosphoenolpyruvate and one ADP to pyruvate and one ATP (Baud *et al.*,
2007; Wulfert *et al.*, 2020). Chlamydomonas mutants deficient in pyruvate kinase were
heat-sensitive under 35°C in acetate-containing medium (Mattoon *et al.*, 2022*a*),
supporting the important roles of glycolysis in thermotolerance of 35°C. Although ATP is
primarily produced through mitochondrial oxidative respiration, the ATP production from
glycolysis can be important under stressful conditions where energy availability is limited
(van Dongen *et al.*, 2011). Under low oxygen conditions, plants increase activity of
pyruvate kinases to produce more ATP (van Dongen *et al.*, 2011). Heat-treated barley
leaves utilized glycolysis as an alternative energy source for thermotolerance based on
proteomics analysis (Rollins *et al.*, 2013).

Additionally, acetate may protect photosynthesis from heat-induced photoinhibition.
Acetate is proposed to protect PSII against photoinhibition by replacing the bicarbonate
associated to the non-heme iron at the acceptor side of PSII, changing the environment
of plastoquinone and affecting PSII charge recombination (Roach *et al.*, 2013). Our PSII
efficiency data supported this report because PSII efficiency was higher with acetate than
without acetate during 25°C, 35°C or 40°C treatments (Fig. 8D). Furthermore,
experimental and modeling analysis suggested that acetate promoted cyclic electron flow
(CEF) around photosystem I (PSI); reducing equivalents produced during the acetate
metabolism reduce plastoquinone pools and increase CEF activity (Johnson and Alric,
2012; Lucker and Kramer, 2013; Chapman *et al.*, 2015). CEF cycles photosynthetic
electrons around PSI, producing only ATP but no NADPH and providing extra ATP
needed for photosynthesis and other cellular activities (Munekage *et al.*, 2004; Baker *et*
al., 2007). CEF balances the ATP/NADPH ratio, contributes to the generation of
transthylakoid proton motive force, and protects both PSI and PSII from photo-oxidative

damage (Johnson, 2011; Yamori and Shikanai, 2016). The increased acetate uptake
under 35°C (Fig. 1D) was coupled with induced CEF activity in Chlamydomonas under
the constant-acetate condition (Zhang *et al.*, 2022*a*). Our transcriptome analysis also
showed up-regulation of transcripts related to PSI and CEF at 2 h of 35°C treatments
under either the constant-acetate or acetate-depleting condition (little acetate depletion
at 2 h), consistent with heat-induced CEF in the presence of acetate (Supplemental Fig.
4J, P).

The effects of acetate on the improved heat tolerance under the moderate high
temperature may also be related to the accumulation of thermoprotective metabolites
(osmolytes, including amino acids, sugars, and sugar alcohols) (Guihur *et al.*, 2022) or
compatible solutes (Hemme *et al.*, 2014; Schroda *et al.*, 2015). Compatible solutes
accumulate from the catabolism of large molecules upon the onset of heat to stabilize
membranes and prevent protein unfolding and aggregation (Schroda *et al.*, 2015). In
Chlamydomonas, these compatible solutes include glycerophos-phoglycerol (GPG),
trehalose, alanine, isoleucine, asparagine, glutamine, and guanosine (Hemme *et al.*,
2014). The elevated acetate metabolism at 35°C with constant acetate supply may
increase the accumulation of these thermoprotective compatible solutes. This is an
understudied but exciting area. Future metabolite analysis may provide more information
about this.

With constant acetate supply, 35°C is beneficial, increasing carbon metabolism and
improving thermotolerance for enhanced growth (Fig. 5). If algal cells can be cultivated
under sterile conditions with constant organic carbon supply, 35°C could be used to
promote algal growth and increase biofuel and bioproduct generation under mixotrophic
conditions in light. Culture temperature of 35°C in closed, outdoor algal ponds may not
be difficult to reach with natural sunlight heating in the summertime of the moderate
climate regions and year-round in the tropical regions. If photoautotrophic medium is used
for enclosed outdoor algal cultivation, acetate could be added to improve algal
thermotolerance before the occurrence of moderate high temperature around 35°C. The
improved algal thermotolerance and growth may outweigh the cost of added acetate.

Without constant acetate supply, 35°C is detrimental. Under the acetate-depleting

condition, the 35°C-elevated carbon metabolism without sufficient carbon input may

deplete cellular carbon reserves. This was not sustainable in the long-term and eventually

reduced biomass accumulation (Fig. 1C). With constant acetate supply, heat at 35°C

transiently arrested the cell cycle, which recovered after 8 h at 35°C (Fig. 3G). In

photoautotrophic medium without acetate, 35°C reduced algal growth rates and resulted

in stressed cells aggregated to form palmelloids (Fig. 6, 7). Such effects of 35°C can

particularly compromise the yields of outdoor algal ponds because they usually do not

contain carbon sources but frequently experience moderate high temperatures (Mata *et*

al., 2010; El-Sheekh *et al.,* 2019).

It is reported that the growth rate of another Chlamydomonas WT strain (CC-125) at 35°C

was about 50% of that at 28°C (0.083 vs 0.166) in photoautotrophic medium under

constant light and continuous dilution but with 2% CO_2 (Vítová *et al.*, 2011). The 50%

growth reduction is comparable to the reduced growth of our data at 35°C without acetate

(Fig. 7A). We used air level CO_2, which is 400 ppm or 0.04% CO_2 to mimic the nature

condition (1/50[th] of 2% CO_2 as in Vítová *et al.*, 2011). CO_2 provides the inorganic carbon

source for algal growth. However, the equivalent 50% growth reduction at 35°C as

compared to the control temperature with (Vítová *et al.*, 2011) and without (this study)

high CO_2 suggests CO_2 (inorganic carbon source) is not as effective as acetate (organic

carbon source) to improve thermotolerance in algal cells. Furthermore, different algal

strains with changes in cultivation conditions may have varying moderate high

temperature ranges. For example, another Chlamydomonas WT strain (CC-137) had

optimal growth at 35°C in photoautotrophic medium with 2% CO_2 in day/night cycle with

semi-dilution (one dilution per day at the end of dark phase) (Lien and Knutsen, 1979).

Such semi-dilution may result in a large range of cell densities, which may affect the light

perceived by each cell. Additionally, the day/night cycle may also alleviate photooxidative

stress as photosynthesis only occurs in light. However, the author mentioned that

cytokinesis and sporulation became inhibited at temperature above 36°C (Lien and

Knutsen, 1979), indicating 35°C may be close to the moderate high temperature range

that would reduce algal growth in this strain without acetate under their cultivation
condition. Additionally, the fast-growing marine alga *Picochlorum celery*, which is tolerant
to high salt, high light, and high temperature, had maximal growth rates around 35°C
which was reduced at temperatures around 40°C, with photoautotrophic medium and
unlimited CO_2 supply in outdoor open ponds under semi-continuous, drain and fill mode
of operation (cells were harvested and fresh medium was added every 2-3 days)
(Krishnan *et al.*, 2021), suggesting the moderate high temperature range is strain-specific
and depending on the cultivation condition.

Furthermore, some tissues in land plants are either mixotrophic or heterotrophic, such as
reproductive tissues (e.g., developing green seeds) (Koley *et al.*, 2022) or sink tissues
(e.g., roots). Reproductive tissues are often considered the most heat sensitive in land
plants (Barnabás *et al.*, 2008; Ullah *et al.*, 2022) and are the main harvest of grain crops
such as soybean, wheat, barley, sorghum, and maize. The up-regulated carbon
metabolism under moderate high temperatures may also occur in land plants, especially
in reproductive or sink tissues. Heat-induced glyoxylate cycle, gluconeogenesis, and
glycolysis have been reported in plants (Rollins *et al.*, 2013; Zhang *et al.*, 2013; Aprile *et*
al., 2013). Carbon metabolism and sugar availability were shown to be essential for heat
tolerance in Arabidopsis (Olas *et al.*, 2021). The thermotolerant wheat cultivar had
increased carbohydrate molecules in sink tissues (Mishra *et al.*, 2021). It is also reported
that heat stress impaired sucrose metabolism in leaves and anthers, which were
associated with heat-induced reproductive failures in chickpea (Kaushal *et al.*, 2013). A
better understanding of the interaction between carbon availability and heat tolerance
could be leveraged to protect sink and/or reproductive tissues from heat stresses.

Heat of 40°C was detrimental to algal cells with and without constant acetate supply
Acute high temperatures at or above 40°C inhibit the algal cell cycle (Fig. 3, 5-7)
(Mühlhaus *et al.*, 2011*b*; Hemme *et al.*, 2014; Zachleder *et al.*, 2019; Ivanov *et al.*, 2021),
alter thylakoid membranes, reduce photosynthesis, and damage respiration (Zhang *et al.*,
2022*a*). These cellular damages took place quickly, mostly within 4 h of heat at 40°C with
constant acetate supply (Zhang *et al.*, 2022*a*). Although 40°C-treated cells grew bigger

and had more chlorophyll per cell under the constant-acetate condition than the acetate-
depleting condition in the short-term (24-h heat) (Fig. 3B, G, D, I, L, N), the constant
acetate supply could not prevent ultimate cell death with 40°C-heat longer than 2 days
(Fig. 5). Without acetate, Chlamydomonas cells also died after 2-day heat at 40°C (Fig
7). The death of Chlamydomonas cultures was also reported at 39°C for 33 h in
photoautotrophic medium without acetate using the same WT strain *21gr* (Zachleder *et*
al., 2019; Ivanov *et al.*, 2021). In conclusion, constant heating at 40°C is lethal for
Chlamydomonas cells independent of acetate availability.

At 24-h of the acetate-depleting condition, algal cells exposed to 40°C had very little
division (only 30% increase of cell density, thus 1.3 X) but they still grew and accumulated
biomass by increasing cell volume (3 X) as compared to the pre-heat condition (Fig. 3B).
In comparison, algal cells exposed to 25°C at 24 h of the acetate-depleting condition had
5.5 X increased cell density but reduced cell volume (55% of the pre-heat values, due to
acetate starvation). If we use fold change of cell density multiplied by fold change of cell
volume to estimate biomass, it could explain the comparable amount of dry biomass in
cultures treated with 40°C and 25°C by the end of 24-h treatment under the acetate-
depleting condition (Fig. 1C). However, the inefficient system with reduced carbon input
and little cell division at 40°C was not sustainable, eventually killing all the algal cells by
the end of 2-day heat of 40°C, even with constant acetate supply (Fig. 5). Our results
suggested that the increased thermotolerance by acetate did not apply to long-term heat
at 40°C.

Acetate had very limited effects on the tolerance to 40°C in Chlamydomonas possibly
because 40°C damaged mitochondrial activities, which are important for acetate
metabolism. Most of the reducing power from acetate assimilation is used in mitochondrial
respiration to produce ATP (Johnson and Alric, 2012, 2013). Our previous results showed
that 40°C reduced mitochondrial activities significantly under the 24-h constant-acetate
condition (Zhang *et al.*, 2022*a*). When comparing 40°C vs 25°C under the acetate-
depleting condition and 40°C vs 35°C independent of the acetate condition, the top
enriched function group was related to mitochondrial electron transport (Supplemental

29-46

Fig. 5B, E), suggesting mitochondria were particularly sensitive to 40°C. This was also

supported by the down-regulation of transcripts related to mitochondrial electron

transport, especially those related to F1-ATPase at 2 h of 40°C (Supplemental Fig. 4AA,

AB, Supplemental Dataset 2). The compromised mitochondrial activities may restrict

acetate uptake and assimilation during 40°C heat treatment. Although there were no

significant differences in acetate consumption between 40°C and 25°C under the 24-h

acetate-depleting conditions (Fig. 1D), the differences may be underestimated and

masked by the acetate depletion. Transcripts of many genes involved in carbon

metabolism, e.g., acetate uptake and assimilation, glyoxylate cycle and gluconeogenesis

pathways, and glycolysis were significantly down-regulated during 40°C under either the

acetate-depleting or constant-acetate condition, especially at 2 h heat (Fig. 2), consistent

with the down-regulated transcripts related to mitochondrial electron transport. Heat at

40°C for longer than 1 day would eventually slow down mitochondrial activities and reduce

acetate-uptake. The decreased acetate uptake and assimilation, in addition to reduced

photosynthesis and arrested cell cycle, may result in the ultimate death of the algal culture

with 2-day heat at 40°C (Fig. 3, 5-8).

Under the acetate-depleting condition, without frequent medium dilution, cell densities

increased at 25°C and 35°C during the 24-h experiments (Fig. 3A). With the constant light

source, the increased cell densities resulted in light shading. However, the amount of

chlorophyll per mL culture was comparable at 24 h of 25°C, 35°C, and 40°C under the

acetate-depleting condition, suggesting comparable level of light shading under all three

temperatures (Fig. 3C). Thus, we could cross off the effects of light shading when

comparing the results at the 24 h of 25°C, 35°C, and 40°C under the acetate-depleting

condition. Additionally, the up-regulated transcripts related to photoprotection at 24 h of

all three temperatures under the acetate-depleting condition when light shading was

present indicated compromised or inefficient photosynthesis that was not due to high light

stress (Supplemental Fig. 4O). The up-regulation of transcripts related to photoprotection

at 24 h of 25°C and 35°C was only present under the acetate-depleting condition but not

the constant-acetate condition, which may be due to reduced CO_2 concentrations with

increased cell density without frequent culture dilution, supported by a recent report that

low CO_2 induced photoprotection independent of light (Ruiz-Sola *et al.*, 2023). The up-

regulation of transcripts related to photoprotection at 24 h of 40°C was similar under either

acetate condition, just with higher induction under the acetate-depleting than the

constant-acetate condition, suggesting it was mainly caused by damaged photosynthesis

at 40°C.

At 24-h of the acetate-depleting condition, the transcripts related to large *HSP*s,

HSP60/70/90s, were down-regulated at 25°C and 35°C, but not under 40°C

(Supplemental Fig. 4A-C). The down-regulation of large *HSP*s may be related to the

increased cell density and light shading, thus reduced light stress at 24 h of 25°C and

35°C under the acetate-depleting condition (Fig. 3A). Besides heat stress, transcripts of

large *HSPs* also responded to light and were also up-regulated under high light stresses

(Anderson et al. 2021). Additionally, the down regulation of large *HSP*s at 25°C and 35°C

but not 40°C at 24 h of the acetate-depleting condition may suggest the energy distribution

between growth and heat tolerance. Most large HSPs are ATP-dependent chaperones,

especially HSP90s (Al-Whaibi, 2011; Xu *et al.*, 2012; Guihur *et al.*, 2022). Down-

regulation of these large HSPs when acetate was depleted may be helpful to reserve

energy for cell growth when the temperature stress was none (25°C) or moderate (35°C).

However, under the acute heat of 40°C, thermotolerance has higher priority than growth,

thus the expression of these large *HSPs* were maintained while cell cycle was arrested

to preserve energy (Supplemental Fig. 4A-C,W, X).

As compared to 40°C, the induction of *HSP*s at 35°C was minimal or none under either

acetate condition (Supplemental Fig. 4A-C), suggesting the 35°C-induced

thermotolerance by acetate may not be related to the accumulation of HSPs. This may

suggest some none-overlapping heat sensing and signaling pathways between 35°C and

40°C, because HSPs have important roles in the classical heat shock responses (Guihur

et al., 2022). Several heat sensors have been proposed (Vu *et al.*, 2019), including

membrane fluidity (Martinière *et al.*, 2011), Ca^{2+} channels in the plasma membrane (Gao

et al., 2012; Finka *et al.*, 2012), unfolded proteins (Rütgers *et al.*, 2017; Li *et al.*, 2020),

and phytochromes (Jung *et al.*, 2016; Legris *et al.*, 2016), and some circadian clock

components (e.g. EARLY FLOWERING 3, ELF3) (Jung *et al.*, 2020). Among these, the

heat effects on phytochromes and ELF3 have been reported around the temperature

ranges of 30°C-35°C (Jung *et al.*, 2016, 2020; Legris *et al.*, 2016), providing potential heat

sensor candidates for moderate high temperatures. Future research to investigate the

shared and unique heat sensors under moderate and acuate high temperatures may

provide more information about how plants sense and adapt to different high

temperatures.

High temperatures affect Chlamydomonas lipid metabolism (Schroda *et al.*, 2015). To

overcome heat-induced membrane fluidity and leakiness, Chlamydomonas reduced

unsaturated membrane lipids and increased the production of saturated lipids to restore

normal membrane viscosity (Schroda *et al.*, 2015; Légeret *et al.*, 2016). Saturated fatty

acids are more stable than unsaturated fatty acids under high temperatures. It was

proposed that the diacylglycerols (DAGs) acyltransferase, DGTT1, converted the

chloroplast unsaturated membrane lipids monogalactosyldiacylglycerols (MGDG) to

DAGs then triacylglycerols (TAGs) in Chlamydomonas with 1 h heat at 42°C (Légeret *et*

al., 2016). Our RNA-seq data showed that the transcripts of *DGTT1* were quickly and

largely up-regulated at 40°C independent of acetate availability (Supplemental Figure

3H), consistent with a strategy to reduce unsaturated membrane lipids during high

temperatures. The transcripts of *DGTT1* were also up-regulated at 35°C, but with a

smaller induction than at 40°C. Additionally, a long chain fatty acid CoA ligase, LCS2,

involves in de novo synthesis of saturated fatty acids by incorporating newly synthesized

fatty acids exported from the plastid into saturated TAGs in lipid droplets (Li *et al.*, 2016;

Li-Beisson *et al.*, 2023). Our RNA-seq data showed that the transcripts of *LCS2* were

quickly and significantly up-regulated at 40°C independent of acetate availability

(Supplemental Figure 3I), suggesting increased de novo synthesis of saturated fatty acids

at 40°C. The de novo fatty acids biosynthesis is the only option for increasing the fraction

of lipids with saturated fatty acids (Hemme *et al.*, 2014). Heat at 40°C reduced

photosynthetic activities within 4 h, which increased chloroplast redox pressure and

oxidative stress under high temperatures (Zhang *et al.*, 2022*a*). Biosynthesis of lipids with

saturated fatty acids via the de novo pathway requires reducing power NADPH and

energy ATP (Ohlrogge and Browse, 1995), which may serve as an electron sink and

alleviate heat-induced redox stresses (Hemme *et al.*, 2014). The transcripts of *LCS2* were

also up-regulated at 24 h of 25°C and 35°C under the acetate-depleting condition

(Supplemental Figure 3I), which may be related to the protective role of the de novo lipid

biosynthesis pathway as an electron sink when acetate was depleted and photosynthesis

was inefficient, consistent with the induced transcripts involved in photoprotection under

the same conditions (Supplemental Figure 4O). Furthermore, our MapMan ontology

enrichment analysis showed that the functional group of lipid metabolism and lipid

degradation was significantly enriched in differentially regulated transcripts between 25°C

and 35°C under the acetate-depleting conditions but not under the constant-acetate

condition, suggesting acetate depletion may make the moderate high temperature more

stressful (Supplemental Figure 5A, C). Our work provides rich datasets for researchers

with interest and expertise in lipid metabolism to further investigate how changes of lipids

affect algal heat responses.

Most previous heat treatments in Chlamydomonas used cultures grown with acetate-

containing medium in flasks without turbidostatic control (Hemme *et al.*, 2014; Rütgers *et*

al., 2017). Our results showed that such heat treatment could not last more than 8 h at

35°C or 24 h of 40°C due to acetate-depletion (Fig. 1D). In fact, many previous heat

experiments were conducted for a short-term, e.g., 1 h (Légeret *et al.*, 2016; Rütgers *et*

al., 2017). Using our highly controlled PBR systems, we could not only conduct heat

experiments under the 24-h acetate-depleting condition but also during 4-day heating with

and without acetate under the turbidostatic control with constant nutrient supply to

uniquely reveal the effects of carbon availability, heat duration and intensity on algal heat

responses. Additionally, we used constant light in our experimental conditions to focus

specifically on the effects of high temperatures. In nature, high temperatures are

accompanied by day and night cycles. The interaction among high temperatures, carbon

status, and diurnal cycles will be an exciting area in future research.

1013

1014   In summary, by using the model green alga Chlamydomonas and highly controlled

1015   cultivation systems in PBRs, we revealed how the availability of an organic carbon source,

acetate, interacted with different intensities and durations of high temperatures in photosynthetic cells. Our research revealed the overlooked effects of moderate high temperature of 35°C, which was beneficial with constant carbon supply or detrimental with insufficient or no carbon supply. We also showed that the damaging effects of acute high temperature of 40°C was dominant and independent of carbon availability. Our research not only helped us understand heat responses in photosynthetic cells but also provided insights for high temperature effects on the production of algal biofuels and bioproducts. Our results will also be useful to study how carbon availability affects heat responses in mixotrophic and heterotrophic (reproductive or sink) tissues in land plants.

Supplementary data

Supplemental Table 1: Gene IDs and primers for RT-qPCR analysis in Fig. 2 and Supplemental Fig. 2 and 3.

Supplemental Dataset 1. RNA-seq data and plots of each transcript detected, including mean normalized read count plots, mean normalized read count heat map, $\log_2$(fold change) of mean normalized read counts relative to the pre-heat time point under the same condition, and statistical analysis using False Discovery Rate (FDR).

Supplemental Dataset 2. $\log_2$(fold change, FC) of mean normalized read counts of groups of transcripts based on MapMan function annotation (MapMan folder) or manual curation (Manual_curation folder). Manual curation was based on the gene lists from this paper Zhang *et al.*, 2022*a*). Each line in the plots represent one unique transcript in that function category under one of six conditions. RNA-seq data with normalized read counts were used for this analysis. $\log_2$FC is relative to the pre-heat time point under the same condition. Black vertical lines indicated the start of heat treatments under 35°C or 40°C or the equivalent time point at 25°C. ANOVA test was applied to the time course (five time points) of one transcript for each condition. If a transcript differed significantly during the time course (or underwent a global significant change) of one condition, it is colored purple (ANOVA significant); blue lines had no significant changes during the time course. Different intensities of purple and blue colors were used to distinguish different transcripts.

1047 Each file opens as a web page. Please use the interactive feature of the figures to see

1048 the gene IDs for each panel.

1049

1050 **Supplemental Dataset 3**. MapMan ontology enrichment. Ontology enrichment was

1051 performed for transcripts with significantly differential expression in the indicated

1052 comparisons using the extended MapMan ontology. The p values were calculated using

1053 a hypergeometric test and corrected for multiple testing using the Benjamini-Hochberg

1054 method.

1055

1056 **Supplemental Dataset 4**. Normalized RNA-seq read counts.

1057

1058 **Supplemental Dataset 5**. RNA-seq data summary, including gene IDs, MapMan function

1059 groups, annotation, adjusted False Discovery Rate (FDR) values for comparisons of

1060 different conditions, mean and standard deviation (sd) of normalized read counts.

1061

Supplemental Dataset 6. ChlamyCyc Pathway enrichment analysis of differentially

regulated transcripts compared between two conditions. The ChlamyCyc annotations

(v2023-01-04) and pathway ontology combine KEGG, MapMan, and JGI pathway

information (May *et al.*, 2009). The p values were calculated using a hypergeometric test

(BioFSharp.Stats) and corrected for multiple testing using the Benjamini-Hochberg

method (FSharp.Stats).

1068

1069 **Supplemental Figures 1-6** (see figure file)

1070

Acknowledgements

We would like to thank Dr. Michael Schroda for discussing the initial experimental idea

under the acetate-depleting conditions, Dr. Trent Northen for recommending the acetate

quantification assay, and Dr. Jaruswan Warakanont for discussion about

Chlamydomonas lipid metabolism. We also want to thank the team of the Department of

Energy (DOE) Joint Genome Institute (JGI) Community Science Program (Anna Lipzen,

Christa Pennacchio, Kerrie Barry, Dr. Jeremy Schmutz, and others) for managing and

sequencing our RNA samples.

1079

Author contributions

RZ designed the experiments. CB and MX operated and maintained the algal growth in

photobioreactors. NZ, CB, and MX harvested algal samples at different time points and

quantified biomass. NZ and MX performed acetate quantification assay. NZ performed

RT-qPCR analysis and extracted RNA for RNA-seq. CB and NZ performed light

microscope imaging. CB, MX, and NZ quantified cell density, cell size, and chlorophyll

contents, and recorded OD_{608}/OD_{720} readings. CB and MX performed 4-day heating

experiments and quantified growth rates as well as cell parameters. CB analyzed PSII

data. BV, TM, RZ did RNA-seq analysis. EMM did initial analysis of the acetate pathways.

NZ, BV, CB, RZ, MX, and EMM prepared the figures. BV wrote the RNA-seq method and

related figure legends. RZ wrote the rest of the manuscript. RZ, EMM, BV, and CB revised

the manuscript.

1092

Conflict of interest

1094 No conflict of interest declared.

1095

Funding statement

This work was supported by the start-up funding from the Donald Danforth Plant Science

Center (DDPSC) and the DOE Biological & Environmental Research (BER) grant (DE-

SC0020400) to RZ. EMM was supported by the William H. Danforth Fellowship in Plant

Sciences and the DDPSC start-up funding (to RZ). BV and TM were supported by the

Germany funding TRR 175 D02 (Project No: 270050988). The RNA-seq service support

was from the Department of Energy (DOE) Joint Genome Institute (JGI) Community

Science Program (CSP, 503414) to RZ. The NovaSeq for RNA-seq (proposal:

10.46936/10.25585/60001126) was conducted by the U.S. Department of Energy Joint

Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is

supported by the Office of Science of the U.S. Department of Energy operated under

Contract No. DE-AC02-05CH11231.

1108

Data availability

All data supporting the findings of this study are available within the paper and within its

supplementary materials published online. Raw RNA-seq data was deposited to

Sequence Read Archive (SRA) following the Data Policy of the Joint Genome Institute

(JGI) with Sequencing Project ID: 1377265.

1114

References

Altschuler M, Mascarenhas JP. 1982. Heat shock proteins and effects of heat shock in plants. Plant molecular biology **1**, 103–115.

Al-Whaibi MH. 2011. Plant heat-shock proteins: A mini review. Journal of King Saud University - Science **23**, 139–150.

Anderson CM, Mattoon EM, Zhang N, *et al.* 2021. High light and temperature reduce photosynthetic efficiency through different mechanisms in the C4 model Setaria viridis. Communications Biology **4**, 1092.

Aprile A, Havlickova L, Panna R, *et al.* 2013. Different stress responsive strategies to drought and heat in two durum wheat cultivars with contrasting water use efficiency. BMC genomics **14**, 821.

Bajhaiya AK, Dean AP, Zeef LAH, Webster RE, Pittman JK. 2016. PSR1 Is a Global Transcriptional Regulator of Phosphorus Deficiency Responses and Carbon Storage Metabolism in Chlamydomonas reinhardtii. Plant physiology **170**, 1216–1234.

Baker NR, Harbinson J, Kramer DM. 2007. Determining the limitations and regulation of photosynthetic energy transduction in leaves. Plant, cell & environment **30**, 1107–1125.

Balfagón D, Sengupta S, Gómez-Cadenas A, Fritschi FB, Azad RK, Mittler R, Zandalinas SI. 2019. Jasmonic acid Is required for plant acclimation to a combination of high light and heat stress. Plant physiology **181**, 1668–1682.

Barnabás B, Jäger K, Fehér A. 2008. The effect of drought and heat stress on reproductive processes in cereals. Plant, cell & environment **31**, 11–38.

Baud S, Wuillème S, Dubreucq B, de Almeida A, Vuagnat C, Lepiniec L, Miquel M, Rochat C. 2007. Function of plastidial pyruvate kinases in seeds of Arabidopsis thaliana. The Plant journal: for cell and molecular biology **52**, 405–419.

BBDuk. 2022. BBMap short read aligner, and other bioinformatic tools.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society. Series B, Statistical methodology **57**, 289–300.

Bigeard J, Hirt H. 2018. Nuclear Signaling of Plant MAPKs. Frontiers in plant science **9**, 469.

Bourgine B, Guihur A. 2021. Heat Shock Signaling in Land Plants: From Plasma Membrane Sensing to the Transcription of Small Heat Shock Proteins. Frontiers in plant science **12**, 710801.

Boyle NR, Page MD, Liu B, *et al.* 2012. Three acyltransferases and nitrogen-responsive regulator are implicated in nitrogen starvation-induced triacylglycerol accumulation in Chlamydomonas. The Journal of biological chemistry **287**, 15811–15825.

de Carpentier F, Maes A, Marchand CH, Chung C, Durand C, Crozet P, Lemaire SD, Danon A. 2022. How abiotic stress-induced socialization leads to the formation of massive aggregates in Chlamydomonas. Plant physiology **190**, 1927–1940.

Chapman S, Paget C, Johnson G, Schwartz J-M. 2015. Flux balance analysis reveals acetate metabolism modulates cyclic electron flow and alternative glycolytic pathways in Chlamydomonas reinhardtii. Frontiers in plant science **6**.

Chew SY, Chee WJY, Than LTL. 2019. The glyoxylate cycle and alternative carbon metabolism as metabolic adaptation strategies of Candida glabrata: perspectives from Candida albicans and Saccharomyces cerevisiae. Journal of Biomedical Science **26**.

Chioccioli M, Hankamer B, Ross IL. 2014. Flow Cytometry Pulse Width Data Enables Rapid and Sensitive Estimation of Biomass Dry Weight in the Microalgae Chlamydomonas reinhardtii and Chlorella vulgaris. PLoS ONE **9**, e97269.

Crozet P, Navarro FJ, Willmund F, *et al.* 2018. Birth of a photosynthetic chassis: A MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas reinhardtii*. ACS synthetic biology **7**, 2074–2086.

Delorge I, Janiak M, Carpentier S, Van Dijck P. 2014. Fine tuning of trehalose biosynthesis and hydrolysis as novel tools for the generation of abiotic stress tolerant plants. Frontiers in plant science **5**.

Dhokane D, Bhadra B, Dasgupta S. 2020. CRISPR based targeted genome editing of *Chlamydomonas reinhardtii* using programmed Cas9-gRNA ribonucleoprotein. Molecular biology reports **47**, 8747–8755.

Dodds WS, Stutzman LF, Sollami BJ. 1956. Carbon Dioxide Solubility in Water. Industrial & Engineering Chemistry Chemical & Engineering Data Series **1**, 92–95.

van Dongen JT, Gupta KJ, Ramírez-Aguilar SJ, Araújo WL, Nunes-Nesi A, Fernie AR. 2011. Regulation of respiration in plants: a role for alternative metabolic pathways. Journal of plant physiology **168**, 1434–1443.

Dunn SR, Thomason JC, Le Tissier MDA, Bythell JC. 2004. Heat stress induces different forms of cell death in sea anemones and their endosymbiotic algae depending on temperature and duration. Cell death and differentiation **11**, 1213–1222.

El-Sheekh MM, Gheda SF, El-Sayed AE-KB, Abo Shady AM, El-Sheikh ME, Schagerl M. 2019. Outdoor cultivation of the green microalga Chlorella vulgaris under stress conditions as a feedstock for biofuel. Environmental science and pollution research international **26**, 18520–18532.

Emrich-Mills TZ, Yates G, Barrett J, *et al.* 2021. A recombineering pipeline to clone large and complex genes in Chlamydomonas. The Plant Cell **33**, 1161–1181.

Fei X, Yu J, Li Y, Deng X. 2017. CrMAPK3 regulates the expression of iron-deficiency-responsive genes in Chlamydomonas reinhardtii. BMC biochemistry **18**, 6.

Finka A, Cuendet AFH, Maathuis FJM, Saidi Y, Goloubinoff P. 2012. Plasma membrane cyclic nucleotide gated calcium channels control land plant thermal sensing and acquired thermotolerance. The Plant cell **24**, 3333–3348.

Gao F, Han X, Wu J, Zheng S, Shang Z, Sun D, Zhou R, Li B. 2012. A heat-activated calcium-permeable channel - Arabidopsis cyclic nucleotide-gated ion channel 6 - is involved in heat shock responses. The Plant Journal **70**, 1056–1069.

González-Ballester D, Casero D, Cokus S, Pellegrini M, Merchant SS, Grossman AR. 2010. RNA-seq analysis of sulfur-deprived Chlamydomonas cells reveals aspects of acclimation critical for cell survival. The Plant cell **22**, 2058–2084.

Greiner A, Kelterborn S, Evers H, Kreimer G, Sizova I, Hegemann P. 2017. Targeting of photoreceptor genes in *Chlamydomonas reinhardtii* via zinc-finger nucleases and CRISPR/Cas9. The Plant cell **29**, 2498–2518.

Guihur A, Rebeaud ME, Goloubinoff P. 2022. How do plants feel the heat and survive? Trends in biochemical sciences **47**, 824–838.

Haq S, Khan A, Ali M, Khattak AM, Gai W-X, Zhang H-X, Wei A-M, Gong Z-H. 2019. Heat Shock Proteins: Dynamic Biomolecules to Counter Plant Biotic and Abiotic Stresses. International journal of molecular sciences **20**, 5321.

Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. 2007. QBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. Genome biology **8**, R19.

Hemme D, Veyel D, Mühlhaus T, *et al.* 2014. Systems-wide analysis of acclimation responses to long-term heat stress and recovery in the photosynthetic model organism *Chlamydomonas reinhardtii*. The Plant cell **26**, 4270–4297.

Hennacy JH, Jonikas MC. 2020. Prospects for engineering biophysical CO2 concentrating mechanisms into land Plants to enhance yields. Annual review of plant biology **71**, 461–485.

Ivanov IN, Zachleder V, Vítová M, Barbosa MJ, Bišová K. 2021. Starch production in *Chlamydomonas reinhardtii* through supraoptimal temperature in a pilot-scale photobioreactor. Cells **10**, 1084.

Jagodzik P, Tajdel-Zielinska M, Ciesla A, Marczak M, Ludwikow A. 2018. Mitogen-Activated Protein Kinase Cascades in Plant Hormone Signaling. Frontiers in plant science **9**, 1387.

Janni M, Gullì M, Maestri E, Marmiroli M, Valliyodan B, Nguyen HT, Marmiroli N. 2020. Molecular and genetic bases of heat stress responses in crop plants and breeding for increased resilience and productivity. Journal of experimental botany **71**, 3780–3802.

Jee H. 2016. Size dependent classification of heat shock proteins: a mini-review. Journal of exercise rehabilitation **12**, 255–259.

Ji S, Siegel A, Shan S-O, Grimm B, Wang P. 2021. Chloroplast SRP43 autonomously protects chlorophyll biosynthesis proteins against heat shock. Nature plants **7**, 1420–1432.

Johnson GN. 2011. Physiology of PSI cyclic electron transport in higher plants. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1807**, 384–389.

Johnson X, Alric J. 2012. Interaction between starch breakdown, acetate assimilation, and photosynthetic cyclic electron flow in *Chlamydomonas reinhardtii*. The Journal of biological chemistry **287**, 26445–26452.

Johnson X, Alric J. 2013. Central carbon metabolism and electron transport in *Chlamydomonas reinhardtii*: metabolic constraints for carbon partitioning between oil and starch. Eukaryotic cell **12**, 776–793.

Jung J-H, Barbosa AD, Hutin S, *et al.* 2020. A prion-like domain in ELF3 functions as a thermosensor in Arabidopsis. Nature **585**, 256–260.

Jung J-H, Domijan M, Klose C, *et al.* 2016. Phytochromes function as thermosensors in Arabidopsis. Science **354**, 886–889.

Karpowicz SJ, Prochnik SE, Grossman AR, Merchant SS. 2011. The GreenCut2 resource, a phylogenomically derived inventory of proteins specific to the plant lineage. The Journal of biological chemistry **286**, 21427–21439.

Kaushal N, Awasthi R, Gupta K, Gaur P, Siddique KHM, Nayyar H. 2013. Heat-stress-induced reproductive failures in chickpea (Cicer arietinum) are associated with impaired sucrose metabolism in leaves and anthers. Functional plant biology: FPB **40**, 1334–1349.

Kim S-C, Guo L, Wang X. 2020. Nuclear moonlighting of cytosolic glyceraldehyde-3-phosphate dehydrogenase regulates Arabidopsis response to heat stress. Nature communications **11**, 3439.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory requirements. Nature methods **12**, 357–360.

Koley S, Chu KL, Mukherjee T, Morley SA, Klebanovych A, Czymmek KJ, Allen DK. 2022. Metabolic synergy in Camelina reproductive tissues for seed development. Science advances **8**, eabo7683.

Krishnan A, Likhogrud M, Cano M, Edmundson S, Melanson JB, Huesemann M, McGowen J, Weissman JC, Posewitz MC. 2021. Picochlorum celeri as a model system for robust outdoor algal growth in seawater. Scientific reports **11**, 11649.

Kropat J, Hong-Hermesdorf A, Casero D, Ent P, Castruita M, Pellegrini M, Merchant SS, Malasarn D. 2011. A revised mineral nutrient supplement increases biomass and growth rate in Chlamydomonas reinhardtii. The Plant journal: for cell and molecular biology **66**, 770–780.

Légeret B, Schulz-Raffelt M, Nguyen HM, Auroy P, Beisson F, Peltier G, Blanc G, Li-Beisson Y. 2016. Lipidomic and transcriptomic analyses of *Chlamydomonas reinhardtii* under heat stress unveil a direct route for the conversion of membrane lipids into storage lipids. Plant, cell & environment **39**, 834–847.

Legris M, Klose C, Burgie ES, Rojas CCR, Neme M, Hiltbrunner A, Wigge PA, Schäfer E, Vierstra RD, Casal JJ. 2016. Phytochrome B integrates light and temperature signals in Arabidopsis. Science **354**, 897–900.

Li X, Fauser F, Jinkerson RE, *et al.* 2019. A genome-wide algal mutant library and functional screen identifies genes required for eukaryotic photosynthesis. Nature genetics **51**, 627–635.

Li Z, Tang J, Srivastava R, Bassham DC, Howell SH. 2020. The transcription factor bZIP60 links the unfolded protein response to the heat stress response in maize. The Plant cell **32**, 3559–3575.

Li X, Zhang R, Patena W, *et al.* 2016. An indexed, mapped mutant library enables reverse genetics studies of biological processes in *Chlamydomonas reinhardtii*. The Plant cell **28**, 367–387.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics **30**, 923–930.

Li-Beisson Y, Warakanont J, Riekhof W, Benning C. 2023. Chapter 2 - Chlamydomonas glycerolipid metabolism. In: Grossman AR,, In: Wollman F-A, eds. The Chlamydomonas Sourcebook (Third Edition). London: Academic Press, 51–97.

Lien T, Knutsen G. 1979. Synchronous growth of *Chlamydomonas reinhardtii*(chlorophyceae): a review of optimal conditions. Journal of phycology **15**, 191–200.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods **25**, 402–408.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology **15**, 550.

Lucker B, Kramer DM. 2013. Regulation of cyclic electron flow in *Chlamydomonas reinhardtii* under fluctuating carbon availability. Photosynthesis research **117**, 449–459.

Martinière A, Shvedunova M, Thomson AJW, Evans NH, Penfield S, Runions J, McWatters HG. 2011. Homeostasis of plasma membrane viscosity in fluctuating temperatures. The New phytologist **192**, 328–337.

Mata TM, Martins AA, Caetano NS. 2010. Microalgae for biodiesel production and other applications: A review. Renewable and Sustainable Energy Reviews **14**, 217–232.

Mathimani T, Pugazhendhi A. 2019. Utilization of algae for biofuel, bio-products and bio-remediation. Biocatalysis and agricultural biotechnology **17**, 326–330.

Mattoon EM, McHargue W, Bailey CE, *et al.* 2022*a*. High-throughput Identification of Novel Heat Tolerance Genes via Genome-wide Pooled Mutant Screens in the Model Green Alga Chlamydomonas reinhardtii. bioRxiv, 2022.07.13.499508.

Mattoon EM, McHargue W, Bailey CE, *et al.* 2022*b*. High-throughput identification of novel heat tolerance genes via genome-wide pooled mutant screens in the model green alga Chlamydomonas reinhardtii. Plant, cell & environment.

May P, Christian J-O, Kempa S, Walther D. 2009. ChlamyCyc: an integrative systems biology database and web-portal for Chlamydomonas reinhardtii. BMC genomics **10**, 209.

Merchant SS, Prochnik SE, Vallon O, *et al.* 2007. The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. Science **318**, 245–250.

Meyer MT, Whittaker C, Griffiths H. 2017. The algal pyrenoid: key unanswered questions. Journal of experimental botany **68**, 3739–3749.

Minagawa J, Tokutsu R. 2015. Dynamic regulation of photosynthesis in *Chlamydomonas reinhardtii*. The Plant journal: for cell and molecular biology **82**, 413–428.

Mishra D, Shekhar S, Chakraborty S, Chakraborty N. 2021. High temperature stress responses and wheat: Impacts and alleviation strategies. Environmental and experimental botany **190**, 104589.

Mittler R, Finka A, Goloubinoff P. 2012. How do plants feel the heat? Trends in biochemical sciences **37**, 118–125.

Mühlhaus T, Weiss J, Hemme D, Sommer F, Schroda M. 2011*a*. Quantitative Shotgun Proteomics Using a Uniform 15N-Labeled Standard to Monitor Proteome Dynamics in Time Course Experiments Reveals New Insights into the Heat Stress Response of Chlamydomonas reinhardtii. Molecular & Cellular Proteomics **10**, M110.004739.

Mühlhaus T, Weiss J, Hemme D, Sommer F, Schroda M. 2011*b*. Quantitative shotgun proteomics using a uniform ^{15}N-labeled standard to monitor proteome dynamics in time course experiments reveals new insights into the heat stress response of *Chlamydomonas reinhardtii*. Molecular & cellular proteomics: MCP **10**, M110.004739.

Munekage Y, Hashimoto M, Miyake C, Tomizawa K-I, Endo T, Tasaka M, Shikanai T. 2004. Cyclic electron flow around photosystem I is essential for photosynthesis. Nature **429**, 579–582.

Murata N, Nishiyama Y. 2018. ATP is a driving force in the repair of photosystem II during photoinhibition. Plant, cell & environment **41**, 285–299.

Niemeyer J, Scheuring D, Oestreicher J, Morgan B, Schroda M. 2021. Real-time monitoring of subcellular H2O2 distribution in *Chlamydomonas reinhardtii*. The Plant cell.

Ohlrogge J, Browse J. 1995. Lipid biosynthesis. The Plant cell **7**, 957–970.

Olas JJ, Apelt F, Annunziata MG, John S, Richard SI, Gupta S, Kragler F, Balazadeh S, Mueller-Roeber B. 2021. Primary carbohydrate metabolism genes participate in heat-stress memory at the shoot apical meristem of Arabidopsis thaliana. Molecular plant **14**, 1508–1524.

Ortiz-Bobea A, Wang H, Carrillo CM, Ault TR. 2019. Unpacking the climatic drivers of US agricultural yields. Environmental research letters: ERL [Web site] **14**, 064003.

Pospíšil P. 2016. Production of reactive oxygen species by photosystem II as a response to light and temperature stress. Frontiers in plant science **7**, 1950.

Pröschold T, Harris EH, Coleman AW. 2005. Portrait of a species: *Chlamydomonas reinhardtii*. Genetics **170**, 1601–1610.

Rädecker N, Pogoreutz C, Gegner HM, *et al.* 2021. Heat stress destabilizes symbiotic nutrient cycling in corals. Proceedings of the National Academy of Sciences of the United States of America **118**.

Remans T, Keunen E, Bex GJ, Smeets K, Vangronsveld J, Cuypers A. 2014. Reliable gene expression analysis by reverse transcription-quantitative PCR: reporting and minimizing the uncertainty in data accuracy. The Plant cell **26**, 3829–3837.

Roach T, Sedoud A, Krieger-Liszkay A. 2013. Acetate in mixotrophic growth medium affects photosystem II in Chlamydomonas reinhardtii and protects against photoinhibition. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1827**, 1183–1190.

Rollins JA, Habte E, Templer SE, Colby T, Schmidt J, von Korff M. 2013. Leaf proteome alterations in the context of physiological and morphological responses to drought and heat stress in barley (Hordeum vulgare L.). Journal of experimental botany **64**, 3201–3212.

Ruiz-Sola MÁ, Flori S, Yuan Y, *et al.* 2023. Light-independent regulation of algal photoprotection by CO2 availability. Nature communications **14**, 1–12.

Rütgers M, Muranaka LS, Schulz-Raffelt M, Thoms S, Schurig J, Willmund F, Schroda M. 2017. Not changes in membrane fluidity but proteotoxic stress triggers heat shock protein expression in *Chlamydomonas reinhardtii*. Plant, cell & environment **40**, 2987–3001.

Sager R. 1955. Inheritance in the green alga *Chlamydomonas reinhardtii*. Genetics **40**, 476–489.

Sasso S, Stibor H, Mittag M, Grossman AR. 2018. From molecular manipulation of domesticated Chlamydomonas reinhardtii to survival in nature. eLife **7**, e39233.

Schloss JA. 1990. A *Chlamydomonas* gene encodes a G protein β subunit-like polypeptide. Molecular & general genetics: MGG **221**, 443–452.

Schmollinger S, Mühlhaus T, Boyle NR, *et al.* 2014. Nitrogen-sparing mechanisms in *Chlamydomonas* affect the transcriptome, the proteome, and photosynthetic metabolism. The Plant cell **26**, 1410–1435.

Schneider K, Venn B, Mühlhaus T. 2022. Plotly.net: A fully featured charting library for .NET programming languages. F1000Research **11**, 1094.

Schroda M, Hemme D, Mühlhaus T. 2015. The *Chlamydomonas* heat stress response. The Plant journal: for cell and molecular biology **82**, 466–480.

Schulz-Raffelt M, Lodha M, Schroda M. 2007. Heat shock factor 1 is a key regulator of the stress response in *Chlamydomonas*. The Plant journal: for cell and molecular biology **52**, 286–295.

Scranton MA, Ostrand JT, Fields FJ, Mayfield SP. 2015. *Chlamydomonas* as a model for biofuels and bio-products production. The Plant journal: for cell and molecular biology **82**, 523–531.

Shimogawara K, Fujiwara S, Grossman A, Usuda H. 1998. High-efficiency transformation of *Chlamydomonas reinhardtii* by electroporation. Genetics **148**, 1821–1828.

Sommer F, Kropat J, Malasarn D, Grossoehme NE, Chen X, Giedroc DP, Merchant SS. 2010. The CRR1 nutritional copper sensor in Chlamydomonas contains two distinct metal-responsive domains. The Plant cell **22**, 4098–4113.

Song Z-T, Zhang L-L, Han J-J, Zhou M, Liu J-X. 2021. Histone H3K4 methyltransferases SDG25 and ATX1 maintain heat-stress gene expression during recovery in Arabidopsis. The Plant journal: for cell and molecular biology **105**, 1326–1338.

Strenkert D, Schmollinger S, Gallaher SD, *et al.* 2019. Multiomics resolution of molecular events during a day in the life of *Chlamydomonas*. Proceedings of the National Academy of Sciences of the United States of America **116**, 2374–2383.

Takano W, Hisabori T, Wakabayashi K-I. 2021. Rapid estimation of cytosolic ATP concentration from the ciliary beating frequency in the green alga Chlamydomonas reinhardtii. The Journal of biological chemistry **296**, 100156.

Ullah A, Nadeem F, Nawaz A, Siddique KHM, Farooq M. 2022. Heat stress effects on the reproductive physiology and yield of wheat. Journal of agronomy and crop science **208**, 1–17.

Usadel B, Poree F, Nagel A, Lohse M, Czedik-Eysenberg A, Stitt M. 2009. A guide to using MapMan to visualize and compare Omics data in plants: a case study in the crop species, Maize. Plant, cell & environment **32**, 1211–1229.

Venn B, Mühlhaus T. 2022. CSBiology/OntologyEnrichment: Release 0.0.1 (0.0.1). Zenodo. https://doi.org/10.5281/zenodo.6340412.

Venn B, Weil L, Schneider K, Zimmer D, Mühlhaus T. 2022. fslaborg/FSharp.Stats. Zenodo. https://doi.org/10.5281/zenodo.6337056.

Vítová M, Bišová K, Hlavová M, Kawano S, Zachleder V, Cížková M. 2011. *Chlamydomonas reinhardtii*: duration of its cell cycle and phases at growth rates affected by temperature. Planta **234**, 599–608.

Voß B, Meinecke L, Kurz T, Al-Babili S, Beck CF, Hess WR. 2010. Hemin and Magnesium-Protoporphyrin IX Induce Global Changes in Gene Expression in Chlamydomonas reinhardtii. Plant physiology **155**, 892–905.

Vu LD, Gevaert K, De Smet I. 2019. Feeling the heat: searching for plant thermosensors. Trends in plant science **24**, 210–219.

Walker RP, Chen Z-H, Famiani F. 2021. Gluconeogenesis in Plants: A Key Interface between Organic Acid/Amino Acid/Lipid and Sugar Metabolism. Molecules **26**.

Wang L, Yang L, Wen X, Chen Z, Liang Q, Li J, Wang W. 2019. Rapid and high efficiency transformation of *Chlamydomonas reinhardtii* by square-wave electroporation. Bioscience reports **39**, BSR20181210.

Wulfert S, Schilasky S, Krueger S. 2020. Transcriptional and Biochemical Characterization of Cytosolic Pyruvate Kinases in Arabidopsis thaliana. Plants **9**.

Xiao L, Young EB, Grothjan JJ, Lyon S, Zhang H, He Z. 2015. Wastewater treatment and microbial communities in an integrated photo-bioelectrochemical system affected by different wastewater algal inocula. Algal Research **12**, 446–454.

Xie B, Bishop S, Stessman D, Wright D, Spalding MH, Halverson LJ. 2013. *Chlamydomonas reinhardtii* thermal tolerance enhancement mediated by a mutualistic interaction with vitamin B12-producing bacteria. The ISME journal **7**, 1544–1555.

Xu Z-S, Li Z-Y, Chen Y, Chen M, Li L-C, Ma Y-Z. 2012. Heat shock protein 90 in plants: molecular mechanisms and roles in stress responses. International journal of molecular sciences **13**, 15706–15723.

Yamori W, Shikanai T. 2016. Physiological functions of cyclic electron transport around Photosystem I in sustaining photosynthesis and plant growth. Annual review of plant biology **67**, 81–106.

Young EB, Reed L, Berges JA. 2022. Growth parameters and responses of green algae across a gradient of phototrophic, mixotrophic and heterotrophic conditions. PeerJ **10**, e13776.

Zachleder V, Ivanov I, Vítová M, Bišová K. 2019. Cell cycle arrest by supraoptimal temperature in the alga *Chlamydomonas reinhardtii*. Cells **8**, 1237.

Zhang N, Mattoon EM, McHargue W, *et al.* 2022*a*. Systems-wide analysis revealed shared and unique responses to moderate and acute high temperatures in the green alga Chlamydomonas reinhardtii. Communications biology **5**, 460.

Zhang N, Pazouki L, Nguyen H, *et al.* 2022*b*. Comparative phenotyping of two commonly used Chlamydomonas reinhardtii background strains: CC-1690 (21gr) and CC-5325 (The CLiP mutant library background). Plants **11**, 585.

Zhang X, Rerksiri W, Liu A, Zhou X, Xiong H, Xiang J, Chen X, Xiong X. 2013. Transcriptome profile reveals heat response mechanism at molecular and metabolic levels in rice flag leaf. Gene **530**, 185–192.

Zhang M, Zhang S. 2022. Mitogen-activated protein kinase cascades in plant signaling. Journal of integrative plant biology **64**, 301–341.

Zones JM, Blaby IK, Merchant SS, Umen JG. 2015. High-resolution profiling of a synchronized diurnal transcriptome from *Chlamydomonas reinhardtii* reveals continuous cell and metabolic differentiation. The Plant cell **27**, 2743–2769.

Figures

Fig. 1.

Fig. 1. **Under the acetate-depleting condition, moderate high temperature of 35°C transiently increased but then decreased Chlamydomonas biomass accumulation.** **(A)** Before heat treatments, Chlamydomonas cells (CC-1690, also called *21gr*, wild-type) were grown in photobioreactors (PBRs) in Tris-acetate-phosphate (TAP) medium (acetate as an organic carbon source) with a light intensity of 100 µmol photons m^{-2} s^{-1} and constant bubbling of air. Algal cultures were maintained turbidostatically with frequent medium dilution using a small range of OD$_{680}$, which monitors chlorophyll content. Figure cited and modified from a supplemental figure of our previous paper (Zhang *et al.*, 2022*a*). **(B)** Experimental outline and sampling time points for the acetate-depleting condition. **(C)** Under the acetate-depleting condition, algal dry biomass (in 90 mL cultures) increased with 8-h heat but decreased with 24-h heat of 35°C as compared to the constant 25°C. Mean ± SE, *n*=3-8 biological replicates. **(D)** Heat of 35°C accelerated acetate

consumption. Acetate content in algal supernatant was quantified using the Acetate Colorimetric Assay Kit. The red dashed line marks the start of heat or the corresponding time point at constant 25°C. **(C, D)** Statistical analyses were performed using two-tailed t-test assuming unequal variance; *, p<0.05, the colors of the asterisks match the heated condition as compared to 25°C at the same time points; #, p<0.05, for the comparisons between 35°C and 40°C at the same time points. Not significant, ns.

Fig. 2.

Fig. 2. Moderate high temperature of 35°C up-regulated transcripts related to acetate uptake and assimilation, glyoxylate cycle, gluconeogenesis, and glycolysis pathways. (A) Simplified pathways of acetate metabolism in Chlamydomonas based on

these papers (Johnson and Alric 2013; Durante et al. 2019). Key enzymes are in boxes and were analyzed via RNA-seq in panels B-P. Acetate uptake/assimilation enzymes: GFY3, putative acetate transporter; ACS2/3, acetyl-CoA synthases; ACK2, acetate kinase. Glyoxylate cycle key enzymes: ICL1, isocitrate lyase; MAS1, malate synthase; CIS2, Citrate synthase; MDH1/2/4, NAD-dependent malate dehydrogenases. Gluconeogenesis key enzymes: PCK1, phosphoenolpyruvate carboxykinase; FBP1, fructose-1,6-bisphosphatase. Glycolysis key enzymes: PYK2/3/4, pyruvate kinase 2/3/4. See all gene IDs and annotations in Supplemental Table 1. PEP, phosphoenolpyruvate. Fru-6-P, fructose-6-phosphate. Fru-1,6-P$_2$, fructose-1,6-bisphosphate. **(B-P)** Transcriptional change of the indicated genes under six different conditions, presented as log$_2$(fold change) of mean normalized read counts relative to the pre-heat time point. In each panel, each line represents the transcriptional time course of the indicated gene under one condition as indicated by the legend. The black vertical lines mark the start of heat or the corresponding time point at constant 25^{o}C. Solid lines for the acetate-depleting condition and dashed lines for the constant-acetate condition. Black, brown, red lines for 25^{o}C, 35^{o}C, and 40^{o}C temperature treatments, respectively. See Supplemental Dataset 1 for bigger pictures and more information.

Fig. 3.

Fig. 3. Algal cells subjected to different temperature treatments under the acetate-depleting condition were compared at the physiological level with those under the constant-acetate condition. (A-E) Cell parameters from algal cultures under the acetate-depleting condition. The algal cultivation and heat treatments were the same as in Fig. 1B. **(F-J)** Cell parameters from algal cultures under the constant-acetate condition, with turbidostatic control, data plotted based on the results from Zhang *et al.*, 2022*a*. **(A-**

J) Two panels on the same row share the same y axis. The red dashed lines mark the start of heat or the corresponding time point at constant 25°C. **(K-O)** Fold change of the indicated cell parameters from algal cultures under the acetate-depleting condition (filled bars) and under the constant-acetate condition (empty bars). (**A-O**) Mean ± SE, n = 3-7 biological replicates. Statistical analyses were performed with two-tailed t-test assuming unequal variance; *, p<0.05, for the comparisons between 35°C or 40°C with 25°C at the same time point under the same acetate condition, the colors of the asterisks match the heat condition; #, p<0.05, for the comparisons between 35°C and 40°C at the same time point under the same acetate condition; ^, p<0.05, for the comparisons between the acetate-depleting and constant-acetate conditions at the same time point and under the same temperature. **(D, E, I, J)** Cell parameters at different time points during constant 25°C had little change, not significantly different from the pre-heat time points (p>0.05).

Fig. 4.

Fig. 4. Both OD$_{680}$ and OD$_{720}$ were linearly proportional to chlorophyll contents under different temperatures and can be used to monitor algal growth rates automatically. Black, brown, red lines and symbols represent 25°C, 35°C, and 40°C treatments, respectively. **(A, B)** OD$_{680}$ and OD$_{720}$ automatically monitored algal growth every one min in algal cultures grown in photobioreactors with different temperatures under the acetate-depleting conditions. The algal cultivation and heat treatments were the same as in Fig. 1B. Multiple lines with the same colors represent biological replicates

(n= 4-8) under the same condition. **(C-H)** OD_{680} and OD_{720} were linearly proportional to Chlorophyll (Chl) contents per mL cultures but not cell volume or cell density. Chl contents **(C, D)**, cell volume **(E, F)**, and cell densities **(G, H)** were plotted against OD_{680} or OD_{720} readings in algal cultures with different treatments. Dashed blue lines are linear trendlines of best fit and R-squared values are displayed.

Fig. 5.

Fig. 5. Under the constant-acetate condition with turbidostatic control, moderate (35°C) and acute high temperatures (40°C) increased and reduced algal growth during 4-day heating, respectively. Black, brown, red lines and symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat or the corresponding time point at constant 25°C. **(A)** Chlamydomonas cultures were grown in photobioreactors (PBRs) with turbidostatic control at different temperatures in acetate-containing medium. Algal cultures were first acclimated at 25°C for 4 days before the temperature was switched to 35°C or 40°C or stayed at 25°C for 4 days. Relative growth rates were calculated based on the cycling of OD_{680} caused by the turbidostatic control (see Fig. 1A and method for details). Each temperature treatment had 3 biological replicates in separate PBRs. **(B-F)** Cell parameters were quantified from algal cultures harvested at different time points with different treatments. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing 35°C or 40°C with 25°C at the same time point. *, $p < 0.05$, the colors of the asterisks match the heat condition; #, $p < 0.05$, for the comparisons between 35°C and 40°C at the same time point. Chl, Chlorophyll.

Fig. 6

Fig. 6. Under the no-acetate condition (photoautotrophic medium) with turbidostatic control, 35ºC-treated cells formed palmelloids and 40ºC-treated cells were bleached after 4-day heat treatments. Light microscopic images of Chlamydomonas cells under different conditions. Chlamydomonas cultures were grown in photobioreactors with turbidostatic control at different temperatures in photoautotrophic medium without acetate. Algal cultures were first acclimated at 25ºC for 4 days before the temperature was switched to 35ºC or 40ºC or stayed at 25ºC for 4 days. Images shown are representative results from three biological replicates. Black scale bar on each panel represents 10 µm.

Fig. 7.

Fig. 7. Under the no-acetate condition (photoautotrophic medium) with turbidostatic control, both moderate (35°C) and acute high temperatures (40°C) reduced algal growth during 4-day heating. Black, brown, red lines and symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat or the corresponding time point at constant 25°C. **(A)** Chlamydomonas cultures were grown in photobioreactors (PBRs) with turbidostatic control at different temperatures in photoautotrophic medium without acetate. Algal cultures were first acclimated at 25°C for 4 days before the temperature was switched to 35°C or 40°C or stayed at 25°C for 4 days. Relative growth rates were calculated based on the cycling of OD_{680} caused by the turbidostatic control. Each temperature treatment had at least 3 biological replicates in separate PBRs. **(B-F)** Cell parameters were quantified from algal cultures harvested at different time points with different treatments. Because 35°C-treated cells under photoautotrophic condition formed palmelloids (see Fig 6), the term of "palmelloid" was used in place of "cell" for this condition only. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing 35°C or 40°C with 25°C at the same time point. *, $p < 0.05$, the colors of the asterisks match the heat condition; #, $p < 0.05$, for the comparisons between 35°C and 40°C at the same time point. Chl, Chlorophyll.

Fig. 8

Fig. 8. Under the turbidostatic control and no nutrient limitation, algal cultures had higher photosystem II (PSII) operating efficiency with acetate than without acetate during treatment of 25°C, 35°C, or 40°C, although the acetate effects diminished after 2-day heat of 40°C due to cell death. PSII operating efficiency was measured automatically in photobioreactors under different conditions. Black, brown, red lines and symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat or the corresponding time point at constant 25°C. **(A)** Acetate-depleting condition at 25°C, 35°C, or 40°C for 24 h, in medium started with acetate but no turbidostatic control, thus no constant acetate supply. **(B)** Constant-acetate condition at 25°C, 35°C, or 40°C for 4 days, in medium with acetate and turbidostatic control, thus no nutrient limitation. **(C)** No-acetate condition at 25°C, 35°C, or 40°C for 4 days, in photoautotrophic medium without acetate but with turbidostatic control, thus no nutrient limitation. **(D)** The comparison of PSII operating efficiency in the constant-acetate (data from panel B, filled bars) and no-acetate (data from panel C, empty bars) conditions during the indicated periods at 25°C, 35°C, or 40°C for 4 days. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing the indicated groups. *, $p < 0.05$, comparing 35°C or 40°C with 25°C with the same medium type at the same time; #, $p < 0.05$, for the comparisons between 35°C and 40°C with the same medium type at the same time; ^, $p < 0.05$, for the comparisons between the same temperature treatment at the same time but with and without acetate. The colors of the significant marks match the heat condition, brown for 35°C and bright red color for 40°C related comparisons, and dark red for 35°C and 40°C comparisons.

Fig. 9.

Fig. 9. Acetate availability affected Chlamydomonas heat tolerance under the moderate high temperature (35ºC) but had much less effects under the acute high temperature (40ºC). In medium containing the organic carbon source, acetate, heat of 35ºC accelerated acetate metabolism by upregulating transcripts related to acetate uptake and assimilation, glyoxylate cycle, gluconeogenesis, and glycolysis pathways. With constant acetate supply, 35ºC increased algal growth. Under the acetate-depleting condition, 35ºC increased algal growth transiently followed by decreased growth and biomass accumulation due to acetate starvation. Without acetate supply in photoautotrophic medium, 35ºC reduced algal growth rates. Under either constant-acetate or depleting-acetate condition, heat of 40ºC down-regulated transcripts related to acetate metabolism, but up-regulated transcripts related to heat shock proteins (HSPs) and reactive oxygen species (ROS) scavenging. In medium with or without acetate, 40ºC

decreased algal growth rates, inhibited cell division, damaged photosynthesis (e.g., photosystem II, PSII), reduced chlorophyll contents, and eventually led to cell death. Thus, Chlamydomonas cells could not survive continuous heat of 40°C for longer than 2 days, independent of acetate supply. Brown and red labels and lines represent the effects of 35°C and 40°C, respectively. Purple, blue, and green labels and lines represent the acetate-depleting, constant-acetate, and no-acetate conditions, respectively.

Supplemental Fig. 1.

Supplemental Fig. 1. RNA-seq data from algal samples harvested from photobioreactors (PBRs) under the constant-acetate-25°C with turbidostatic control were highly correlated with each other. The Pearson correlations were generated by comparing all RNA-seq samples to the first sample (marked by *) of the constant-acetate-25°C condition. Under constant acetate, 25°C, and light (constant-acetate-25°C) with turbidostatic control in PBRs, the algal growth rates and cell physiologies stay constant (Fig. 5). We had RNA-seq data for 8 time points under the constant-acetate-25°C condition, and these samples were highly correlated with each other (with Pearson correlation ≥ 0.993). The 0 h of the constant-acetate-25°C condition corresponds to the time when PBRs reach expected high temperatures during heat treatments, which takes about 30 min from 25°C to 35°C or 40°C. All pre-heat samples were harvested from PBRs under the same constant-acetate-25°C condition with turbidostatic control and these RNA-seq samples had Pearson correlation > 0.95.

Because the cell status and transcriptome stay steady under the constant-acetate-25°C, we used the five extra time points under the constant-acetate-25°C condition (with underlines) to impute the 2 h and 4 h time points under the constant-acetate-25°C condition. See method for details.

Supplemental Fig. 2.

Supplemental Fig. 2. RT-qPCR analysis was consistent with RNA-seq results. Fold changes of four selected transcripts involved in the glyoxylate cycle (*ICL1* and *MAS1*) and gluconeogenesis cycle (*PCK1* and *FBP1*) as in Fig. 2A were calculated based on RT-qPCR (**A-D**) or RNA-seq (**E-H**) results. The red dashed lines mark the start of heat or the corresponding time point at constant 25°C. For RT-qPCR results, the relative fold changes were calculated by normalizing the transcript expression to the reference genes *CBLP*, *EIF1A* and the pre-heat level. For RNA-seq results, the relative fold changes were calculated based on normalized RNA-seq read counts relative to the pre-heat level. Values are mean ± SE, n = 3 biological replicates. Glyoxylate cycle key enzymes: ICL1, isocitrate lyase; MAS1, malate synthase. Gluconeogenesis key enzymes: PCK1, phosphoenolpyruvate carboxykinase; FBP1, fructose-1,6-bisphosphatase (See all gene IDs and RT-qPCR primers in Supplemental Table 1).

Supplemental Fig. 3

Supplemental Fig. 3 continued

Supplemental Fig. 3 continued

Supplemental Fig. 3 continued

Supplemental Fig. 3. Log$_2$(fold change) of the indicated transcript relative to pre-heat under different conditions. Log$_2$(fold change) were determined on the mean normalized transcript abundance, relative to the pre-heat time point. Each line represents one condition as indicated by the legend. The black vertical lines mark the start of heat or the corresponding time point at constant 25°C. Solid lines for the acetate-depleting condition and dashed lines for the constant-acetate condition. Black, brown, red lines for 25°C, 35°C, and 40°C temperature treatments, respectively. See all gene IDs in Supplemental Table 1 and Supplemental Dataset 1 for bigger pictures and more information.

Supplemental Fig. 4.

A. Transcripts related to HSP60s (HSP, heat shock protein)

Supplemental Fig. 4. Log$_2$(fold change, FC) of mean normalized read counts of groups of transcripts based on MapMan function annotation or manual curation. Plots are Log$_2$FC of transcripts related to the indicated function groups under six different treatments, as compared to the pre-heat time points. RNA-seq data with normalized read counts were used for this analysis. Black vertical lines mark the start of heat treatments or the corresponding time point at constant 25°C. Each line represents one unique transcript in the indicated function group. ANOVA test was applied to the time course (five time points) of one transcript for each condition. If a transcript differed significantly during the time course (or underwent a global significant change) of one condition, it is colored purple (ANOVA significant); blue lines had no significant changes during the time course. Different intensities of purple and blue colors were used to distinguish different transcripts. The total number of transcripts plotted in each function group is marked on the top left panel. See interactive figures with gene IDs and annotations in Supplemental Dataset 2. **See more panels of Supplemental Fig. 4 below.**

Supplemental Fig. 4. continued

Supplemental Fig. 4. continued

E. Transcripts related to small HSPs

F. Transcripts related to LHCII

G. Transcripts related to PSII biogenesis

H. Transcripts related to PSII subunits

I. Transcripts related to LHCI

Supplemental Fig. 4. continued

K. Transcripts related to Cytochrome b_6f subunits

L. Transcripts related to ATP synthase subunits

Supplemental Fig. 4. continued

M. Transcripts related to Calvin-Benson cycle

N. Transcripts related to carbon concentrating mechanism (CCM)

O. Transcripts related to photoprotection

P. Transcripts related to cyclic electron flow around PSI (CEF)

Q. Transcripts related to chlorophyll biosynthesis

R. Transcripts related to ROS scavenging: peroxidase (POX)

S. Transcripts related to ROS scavenging: ascorbate peroxidase (APX)

T. Transcripts related to ROS scavenging: catalase (CAT)

U. Transcripts related to ROS scavenging: glutathione reductase (GR)

V. Transcripts related to ROS scavenging: glutathione S-transferase (GST)

Supplemental Fig. 4. Continued

W. Transcripts related to cell cycle

X. Transcripts related to cell wall

Supplemental Fig. 4. Continued

Y. Transcripts related to protein synthesis, ribosomal protein, prokaryotic

Z. Transcripts related to protein synthesis, ribosomal protein, chloroplast

Supplemental Fig. 4. Continued

AA. Transcripts related to mitochondrial electron transport / ATP synthesis (F1-ATPase)

AB. Transcripts related to mitochondrial electron transport / ATP synthesis

Supplemental Fig. 4. Continued

AC. Transcripts related to RNA

AD. Transcripts related to RNA processing

Supplemental Fig. 4. Continued

AE. Transcripts related to transport. phosphate

AF. Transcripts related to signaling. mitogen-activated protein kinase (MAP kinase)

Supplemental Fig. 4. Continued

AG. Transcripts related to cell motility. eukaryotes. axonemal dyneins

AH. Transcripts related to cell motility

Supplemental Fig. 4. Log$_2$(fold change, FC) of mean normalized read counts of groups of transcripts based on MapMan function annotation or manual curation. Plots are Log$_2$FC of transcripts related to the indicated function groups under six different treatments, as compared to the pre-heat time points. RNA-seq data with normalized read counts were used for this analysis. Black vertical lines mark the start of heat treatments or the corresponding time point at constant 25°C. Each line represents one unique transcript in the indicated function group. ANOVA test was applied to the time course (five time points) of one transcript for each condition. If a transcript differed significantly during the time course (or underwent a global significant change) of one condition, it is colored purple (ANOVA significant); blue lines had no significant changes during the time course. Different intensities of purple and blue colors were used to distinguish different transcripts. The total number of transcripts plotted in each function group is marked on the top left panel. See interactive figures with gene IDs and annotations in Supplemental Dataset 2.

Supplemental Fig. 5A

Acetate depleting 25vs35

-log$_{10}$ (p value)

Supplemental Fig. 5B

Acetate depleting 25vs40

-log$_{10}$ (p value)

Supplemental Fig. 5C

Constant acetate 25vs35

Supplemental Fig. 5D

Constant acetate 25vs40

-log$_{10}$ (p value)

Supplemental Fig. 5E

Temperature

Category	Count
mitochondrial electron transport / ATP synthesis.NADH-DH (type I)	55/58 **
mitochondrial electron transport / ATP synthesis	108/131 **
mitochondrial electron transport / ATP synthesis.NADH-DH (type I).complex I	33/34 **
transport.p- and v-ATPases.H+-transporting two-sector ATPase	17/17 **
protein.synthesis.ribosomal protein.prokaryotic	52/67 **
mitochondrial electron transport / ATP synthesis.NADH-DH (type I).localisation not clear	22/24 **
protein.synthesis.ribosomal protein.prokaryotic.chloroplast	32/38 **
transport	347/565 **
protein.synthesis.initiation	44/59 **
TCA / organic transformation	38/50 **
PS.lightreaction.photosystem I	19/22 *
amino acid metabolism.synthesis.aspartate family.lysine	10/10 *
TCA / organic transformation.TCA	22/27 *
mitochondrial electron transport / ATP synthesis.F1-ATPase	15/17 *
protein.synthesis.ribosomal protein.prokaryotic.chloroplast.30S subunit	12/13 *
lipid metabolism.lipid degradation	44/63
amino acid metabolism.synthesis.aspartate family	31/42
misc.nitrilases, nitrile lyases, berberine bridge enzymes, reticuline oxidases, troponine reductases	11/12
protein.synthesis.ribosomal protein.prokaryotic.chloroplast.50S subunit	20/25
amino acid metabolism.degradation.branched chain group	16/19
gluconeogenesis / glyoxylate cycle	11/12
mitochondrial electron transport / ATP synthesis.cytochrome c oxidase	16/19
lipid metabolism	141/228
amino acid metabolism	138/223
PS.lightreaction.photosystem I.LHC-I	10/11
transport.metabolite transporters at the mitochondrial membrane	34/48
TCA / organic transformation.other organic acid transformations	12/14
transport.p- and v-ATPases	30/42
protein.synthesis.ribosome biogenesis.pre-rRNA processing and modifications.snoRNPs	9/10
protein.folding.immunophilins (IMM).FKBPs	17/22
RNA.regulation of transcription.HDA	15/19
lipid metabolism.lipid degradation.lipases	27/38
lipid metabolism.phospholipid synthesis	22/30
protein.targeting	120/197
protein.targeting.mitochondria	20/27
glycolysis	43/65
transport.metal	32/47
transport.metabolite transporters at the envelope membrane	19/26
glycolysis.unclear/dually targeted	10/12
nucleotide metabolism.degradation	17/23
amino acid metabolism.synthesis	91/149
secondary metabolism.isoprenoids.carotenoids	18/25

$-\log_{10}$ (p value)

Supplemental Fig. 5F

Medium

Supplemental Fig. 5G

Supplemental Fig. 5. MapMan ontology enrichment revealed differentially regulated biological processes under different conditions. Ontology enrichment was performed for transcripts with significantly differential expression in the indicated comparisons using the extended MapMan ontology. The p values were calculated using a hypergeometric test. All functional sets with a p value less than 0.05 and more than 10 annotated transcripts

are shown. The ratio at the end of each bar indicates the number of significant transcripts between the two compared conditions vs detected transcripts in our RNA-seq data contained in the respective function bin. Multiple testing correction was performed by Benjamini & Hochberg FDR, * FDR < 0.1, ** FDR < 0.05. The bars are sorted based on the FDR p values, from top to bottom, from the small to big FDR values. **(A)** acetate-depleting 25°C vs acetate-depleting 35°C; **(B)** acetate-depleting 25°C vs acetate-depleting 40°C; **(C)** constant-acetate 25°C vs constant-acetate 35°C; **(D)** constant-acetate 25°C vs constant-acetate 40°C; **(E)** temperature effect: (constant-acetate 35°C and acetate-depleting 35°C) vs (constant-acetate 40°C and acetate-depleting 40°C); **(F)** medium effect: (constant-acetate 35°C and constant-acetate 40°C) vs (acetate-depleting 35°C and acetate-depleting 40°C); **(G)** acetate-depleting 25°C vs constant-acetate 25°C. See interactive figures and detailed enrichment analysis in Supplemental Dataset 3.

Supplemental Fig. 6

Supplemental Fig. 6. Chlamydomonas cells had different morphologies at different temperatures under the acetate-depleting condition. Light microscopic images of Chlamydomonas cells under the acetate-depleting condition. Images shown are representative results from at least three biological replicates. The algal cultivation and heat treatments were the same as in Fig. 1B.