Moderate High Temperature is Beneficial or Detrimental

Depending on Carbon Availability in the Green Alga *Chlamydomonas reinhardtii*

Ningning Zhang[1,#], Benedikt Venn[2,#], Catherine E. Bailey[1], Ming Xia[1], Erin M. Mattoon[1, 3],

Timo Mühlhaus[2], Ru Zhang[1,*]

[1]Donald Danforth Plant Science Center, St. Louis, Missouri 63132, USA;

[2] RPTU Kaiserslautern, Kaiserslautern, 67663, Germany;

[3]Plant and Microbial Biosciences Program, Division of Biology and Biomedical

Sciences, Washington University in Saint Louis, St. Louis, MO 63130, USA.

Ningning Zhang: lucy.nnzhang@gmail.com, co-first-author[#]

Benedikt Venn: venn@rptu.de, co-first-author[#]

Ming Xia: celine.xia@wustl.edu, present address[3]

Catherine E. Bailey: cbailey@danforthcenter.org

Erin M. Mattoon: EMattoon@danforthcenter.org

Timo Mühlhaus: timo.muehlhaus@rptu.de

Ru Zhang: rzhang@danforthcenter.org, corresponding author*

Running title: Algal heat responses interact with carbon availability.

Highlight

We revealed the overlooked, dynamic effects of moderate high temperature in algae

depending on carbon availabilities and demonstrated the importance of carbon

metabolism in thermotolerance of photosynthetic cells.

Abstract

High temperatures impair plant/algal growth and reduce food/biofuel production, but the

underlying mechanisms remain elusive. The unicellular green alga *Chlamydomonas*

reinhardtii is a superior model to study heat responses in photosynthetic cells due to its

fast growth rate, many similarities in cellular processes to land plants, simple and

sequenced genome, and ample genetic/genomics resources. Chlamydomonas grows in
light by photosynthesis and/or with supplied carbon source, acetate. The majority
published research about Chlamydomonas heat responses used acetate-containing
medium. How organic carbon sources affect heat responses is important for algal
biofuel/bioproduct industry but understudied. We cultivated Chlamydomonas wild-type
cultures under highly controlled conditions in photobioreactors at control of 25°C,
moderate high temperature of 35°C, or acute high temperature of 40°C with and without
constant acetate supply for 1- or 4-days. Our results showed that 35°C increased algal
growth with constant acetate supply but reduced algal biomass and growth without
sufficient acetate. The overlooked and dynamic effects of 35°C can be explained by
induced carbon metabolisms, including acetate uptake/assimilation, glyoxylate cycle,
gluconeogenesis pathways, and glycolysis. Acute high temperature at 40°C for more than
2 days was lethal to algal cultures with and without constant acetate supply. Our research
provides insights to understand algal heat responses and help improve thermotolerance
in photosynthetic cells.

Key words: moderate high temperature, acute high temperature, heat responses,
Chlamydomonas reinhardtii, acetate, glyoxylate cycle, gluconeogenesis, glycolysis,
photobioreactors

Abbreviations:
Photobioreactors (PBRs), Tris-acetate-phosphate (TAP) medium, Tris-phosphate (TP)
medium

Introduction
Many biological processes in photosynthetic cells are sensitive to high temperatures
(Mittler *et al.*, 2012; Schroda *et al.*, 2015; Guihur *et al.*, 2022). Global warming increases
the intensity, duration, and frequency of high temperatures in the field, reducing plant
growth and food/biofuel production (Janni *et al.*, 2020). A recent model revealed high
temperature as the primary climatic driver for yield loss in crops from 1981-2017 in US
(Ortiz-Bobea *et al.*, 2019). Considering the increasing global temperature and human

population, it is imperative to understand how photosynthetic cells respond to high
temperatures and improve plant thermotolerance.

High temperatures in the field have different intensities and durations. For many land
plants or algae grown in temperate regions, moderate high temperatures refer to heat
slightly above the optimal temperature for growth (at or around 35°C), while acute high
temperatures refer to heat at or above 40°C, which significantly reduces or completely
halts growth (Zhang *et al.*, 2022*a*). Moderate high temperatures are often long-lasting and
frequent in nature with mild effects on photosynthetic cells, while acute high temperatures
are often short-term but severely damaging. Most previous heat-stress experiments in
plants and algae used acute high temperatures at or above 40°C (Hemme *et al.*, 2014;
Rütgers, 2017; Balfagón *et al.*, 2019; Kim *et al.*, 2020; Ji *et al.*, 2021), likely due to the
rapid onset and easily quantifiable phenotypes as compared to moderate high
temperatures. Although the impact of moderate high temperatures in photosynthetic cells
can be difficult to investigate due to comparatively mild phenotypes, moderate high
temperatures are physiologically relevant stresses in field conditions and their frequent
and long-lasting features could reduce agricultural yield significantly (Delorge *et al.*, 2014;
Anderson *et al.*, 2021). Arabidopsis seedlings treated with 5-day heat at 35°C had
reduced growth and viability (Song *et al.*, 2021). Additionally, global warming can further
increase the frequency and duration of moderate high temperatures in nature. However,
the effects of moderate high temperatures in photosynthetic cells are often overlooked
and underexplored.

The unicellular green alga *Chlamydomonas reinhardtii* (Chlamydomonas throughout) is a
superior model to study heat responses in photosynthetic cells (Schroda *et al.*, 2015).
Chlamydomonas can grow in light under photoautotrophic conditions using
photosynthesis (Minagawa and Tokutsu, 2015). It can also grow with supplied organic
carbon source (acetate) in light (mixotrophic) or dark (heterotrophic) conditions (Sasso *et*
al., 2018), providing a platform to study heat responses under different light/carbon
conditions. Additionally, it has a haploid, sequenced, annotated, small genome (111 Mb,
17,741 protein-encoding genes) with simpler gene families and fewer gene duplications

than land plants (Merchant *et al.*, 2007; Karpowicz *et al.*, 2011). Furthermore, several

well-established gene editing, cloning, high-throughput, and functional genomic tools are

available in Chlamydomonas (Shimogawara *et al.*, 1998; Li *et al.*, 2016, 2019; Greiner *et*

al., 2017; Crozet *et al.*, 2018; Wang *et al.*, 2019; Dhokane *et al.*, 2020; Emrich-Mills *et al.*,

2021).

Algae have great potential for production of biofuels and bioproducts (Scranton *et al.*,

2015; Mathimani and Pugazhendhi, 2019). However, how algal cells respond to high

temperatures is under-investigated as compared to land plants (Schroda *et al.*, 2015).

Outdoor algal ponds frequently experience moderate high temperatures around 35°C

during the summer time (Mata *et al.*, 2010), but the effects of moderate high temperatures

on algal growth have been overlooked. Previously published algal heat treatments were

often conducted in flasks in pre-warmed water baths at or above 42°C with sharp

temperature switches and without control of nutrients (Hemme *et al.*, 2014; Rütgers *et al.*,

2017). While previous research was highly valuable to understand algal heat responses,

high temperatures in the field or outdoor ponds often increase gradually. Rapid increases

to high temperatures largely reduce algal viability (Zhang *et al.*, 2022*a*); similar effects

have also been reported in land plants (Mittler *et al.*, 2012). Acute high temperatures at

or above 40°C inhibit algal cell division (Mühlhaus *et al.*, 2011*b*; Hemme *et al.*, 2014;

Zachleder *et al.*, 2019; Ivanov *et al.*, 2021), while moderate high temperature at 35°C only

transiently inhibits cell division during the first 4-8 hour (h) heat (Zhang *et al.*, 2022*a*).

Thus, conducting long-term experiments at moderate high temperatures in flasks can

result in overgrown cultures, nutrient depletion, and light limitation, complicating data

interpretation.

Consequently, it is advantageous to investigate algal heat responses under highly-

controlled conditions in photobioreactors (PBRs) (Zhang *et al.*, 2022*a*; Mattoon *et al.*,

2022), which have several evident strengths: (1) precise temperature regulation with a

sterile temperature probe inside each algal culture; (2) controlled heating to mimic the

heating speed in nature; (3) the availability of turbidostatic control based on defined

parameters (e.g., chlorophyll contents) to enable frequent culture dilutions with constant

medium supply; (4) precisely controlled cultivation condition, including temperature, light,

nutrient, and air agitation, allowing for reproducible experiments; (5) automatic recording

of culture status every minute, e.g. growth conditions and optical densities, enabling

quantitative growth rate measurements; (6) being able to simulate nutrient depletion by

turning off the turbidostatic control to investigate how nutrient availability affects algal heat

responses. Utilization of PBRs for algal cultivation and heat treatments can largely reduce

compounding effects during high temperature treatments and improve our understanding

of algal heat responses.

Recently, we conducted systems-wide analyses in wild-type (WT) Chlamydomonas

cultures during and after 24-h heat of 35°C and 40°C in PBRs with constant nutrient

supply and light in acetate-containing medium (mixotrophic condition) under the

turbidostatic control (Zhang *et al.*, 2022*a*). We chose moderate high temperature of 35°C

and acute high temperature of 40°C based on the algal growth rates under different

temperatures; and the 24-h heat duration allowed us to balance time-course sample

collection over a sufficient duration with maintaining culture viability (heat at 40°C for

longer time killed algal cells); our experimental conditions enabled us to investigate algal

heat responses during and after physiological-relevant high temperatures (Zhang *et al.*,

2022*a*). Our results showed that 40°C inhibited algal growth while 35°C increased algal

growth with constant acetate supply. The growth inhibition at 40°C could be explained by

reduced photosynthesis, impaired respiration, and arrested cell cycle, while these cell

parameters had minor changes in algal cultures treated by 35°C. Our proteomics data

indicated that several proteins involved in acetate uptake/assimilation, glyoxylate cycle,

and gluconeogenesis were up-regulated during 35°C (Zhang *et al.*, 2022*a*).

Chlamydomonas uptakes acetate and feeds it into the glyoxylate cycle and

gluconeogenesis for starch biosynthesis (Johnson and Alric, 2012, 2013). The main

function of glyoxylate cycle is to allow growth when glucose is unavailable and two-carbon

compounds, e.g., acetate, are the organic carbon source; glyoxylate cycle is a shunt of

the tricarboxylic acid (TCA) cycle in mitochondria but without CO_2 releases to allow for

the anabolism of simple carbon compounds in gluconeogenesis, a process to make

sugars, e.g., glucose (Johnson and Alric, 2012; Chew *et al.*, 2019; Walker *et al.*, 2021).

We hypothesized that the increased growth in Chlamydomonas at 35°C with constant

acetate supply was attributable to up-regulated acetate metabolisms. The majority of

published high temperature research in Chlamydomonas has been conducted in acetate-

containing medium (Voß *et al.*, 2010; Mühlhaus *et al.*, 2011*a*; Hemme *et al.*, 2014;

Rütgers *et al.*, 2017; Zhang *et al.*, 2022*a*), but the interface between acetate supply and

algal heat responses is understudied. To validate our hypothesis and address these

unknown questions, we cultivated the same Chlamydomonas WT strain in PBRs as

before but heated the cultures at 35°C or 40°C with and without constant acetate supply

for 1 or 4 days. Our previous work was conducted at 35°C and 40°C for 24 h with constant

acetate supply (Zhang *et al.*, 2022*a*). In this paper, to investigate the effects of carbon

availability on algal thermotolerance, we included 3 new different heat treatments at 35°C

and 40°C: acetate-depleting condition for 24 h, constant-acetate condition for 4 days, no-

acetate condition for 4 days (the last two conditions with turbidostatic control and constant

nutrient supply). Our new results confirmed our hypothesis about the role of acetate

metabolism in algal thermotolerance and revealed the overlooked effects of moderate

high temperature of 35°C on algal growth. Heat of 35°C can be beneficial or detrimental

based on acetate availability. Acute high temperature of 40°C is lethal to algal cultures,

with and without constant acetate supply. Our research provided insights to understand

algal heat responses and help improve thermotolerance in photosynthetic cells.

Materials and methods

Algal cultivation

Chlamydomonas reinhardtii wild-type strain CC-1690 (also called *21gr*, mating type plus)

(Pröschold *et al.*, 2005; Zhang *et al.*, 2022*b*; Sager, 1955-7) was purchased from the

Chlamydomonas resource center and used in all experiments. Algal cultures were grown

in Tris-acetate-phosphate (TAP, with acetate) or Tris-phosphate (TP, without acetate)

medium with revised trace elements (Kropat *et al.*, 2011) in 400 mL photobioreactors

(PBRs) (Photon System Instruments, FMT 150/400-RB) as described before (Zhang *et*

al., 2022*a*). Each medium type used individual PBRs in independent experiments.

Cultures were illuminated with constant 100 μmol photons $m^{-2} s^{-1}$ light (red: blue, 1:1 ratio),

and agitated by bubbling with filtered air at a flow rate of 1 L min^{-1}. Algal cultures with

targeted cell density around 1-2 $\times 10^6$ cells mL^{-1} (~4-6 μg mL^{-1} chlorophyll content) were

maintained at 25°C for 4 days with constant nutrient supply through a turbidostatic mode

before different temperature treatments. The turbidostatic mode was controlled by OD_{680}

(optical density at 680 nm), which is proportional to chlorophyll contents (μg mL^{-1}) and

was monitored once per min automatically. When OD_{680} increased to a maximum value

slightly above the target value (for target cell density) due to algal growth, OD_{680} signaled

the control computer to turn on a turbidostatic pump to add fresh medium to dilute the

culture until a minimum OD_{680} slightly below the target value was reached, then the

turbidostatic pump was turned off automatically. Because of the small OD range we used,

the PBR cultures had exponential growth between dilution events through the

turbidostatic mode with constant nutrient supply (Fig. 1A). The OD_{680} data during

exponential growth phases in between dilution events was log_2 transformed, and the

relative growth rate was calculated using the slope of $log_2(OD_{680})$. The relative growth

rate is an inverse of the doubling time of algal cultures. For the treatments without

constant nutrient supply (acetate-depleting condition), the turbidostatic pumps were

turned off after PBR cultures reached steady growth and at the start of different

temperature treatments.

High temperature treatments in PBRs

After algal cultures in PBRs reached steady growth with turbidostatic control at 25°C for

4 days, PBR temperatures were increased to moderate or acute high temperatures (35°C

or 40°C in different PBRs) for the indicated duration with indicated nutrient conditions.

PBR temperatures were changed from 25°C to 35°C or 40°C gradually over the course

of 30 min with controlled heating speeds. PBR cultures grown under constant 25°C with

the same nutrient status and treatment duration served as controls. Algal cultures were

harvested from PBRs at different time points during different treatments for various

measurements.

Algal biomass quantification

Algal cultures (90 mL) were harvested from PBRs, centrifuged to remove supernatants,

flash frozen in liquid nitrogen, then stored in a -80°C freezer until analysis. Algal cell

pellets were freeze dried for 24 h over liquid nitrogen. The remaining algal dry biomass

were weighted and quantified.

Acetate quantification assay

Algal cultures were harvested from PBRs (2 mL) and centrifuged to collect 500 µL top

clear supernatants to a new tube. The supernatant was stored in a -80°C freezer until

analysis. The acetate content in the supernatant was quantified using the Acetate

Colorimetric Assay Kit (Sigma, Cat No. MAK086) according to manufacture instructions.

Cell imaging using light microscopy

Chlamydomonas cultures harvested at select time points with different temperature

treatments were fixed with 0.2% glutaraldehyde (VWR, Cat No. 76177-346). Algal cells

were imaged with a Leica DMI6000 B microscope and a 63x (NA1.4) oil-immersion

objective.

Chlorophyll quantification

Chlorophyll contents in algal cells were quantified as before (Zhang *et al.*, 2022*a*). PBR

cultures of 1 mL were harvested in 1.5-mL tubes with 2.5 µL 2% Tween20 (Sigma, P9416-

100ML, to help cell pelleting), and centrifuged at 18,407 g at 4°C. After removing

supernatant, cell pellets were stored in a -80°C freezer until quantification. Later, cell

pellets were thawed, resuspended in 1 mL of HPLC grade methanol (100%, Sigma,

34860-4L-R), vortexed for 1 min, incubated in the dark at 4°C for 5 min, and centrifuged

at 15,000 g at 4°C for 5 min. Supernatant was pipetted out to a two-sided disposable

plastic cuvettes (VWR, 97000-586) for chlorophyll (Chl) quantification at 652 nm and 665

nm in a spectrophotometer (IMPLEN Nonophotometer P300) using the following

equations: Chl a + Chl b = $22.12*A_{652} + 2.71*A_{665}$ (in µg mL^{-1} algal cultures) (Zhang *et al.*,

2022*a*). Chlorophyll concentrations were also normalized to cell densities (Chl pg cell^{-1})

or algal cell volume (Chl pg μm^{-3}). Cell density and mean cell volume were measured

using a Coulter Counter (Multisizer 3, Beckman Counter, Brea, CA).

RNA extraction

PBR cultures of 2 mL were pelleted with Tween-20 (0.005%, v/v) by centrifugation at

11,363 x g and 4^{o}C for 2 min, followed by supernatant removal. The cell pellet was flash

frozen in liquid nitrogen and then stored in a -80^{o}C freezer until processing. RNA

extraction and RT-qPCR analysis were performed as before with minor modifications

(Zhang *et al.*, 2022*a*). Total RNA was extracted with TRIzol reagent (Thermo Fisher

Scientific, Cat No. 15596026), digested on-column with RNase-free DNase (Qiagen, Cat

No. 79256), purified by RNeasy mini-column (Qiagen, Cat No. 74106), and quantified with

Qubit™ RNA BR Assay Kit, (Life technology, Cat No. Q10210).

RT-qPCR analysis

Total 0.4 μg RNA was reverse transcribed with oligo dT primers using SuperScript® III

First-Strand Synthesis System (Life technology, Cat No. 18080-051) following

manufacturer instructions. Quantitative real-time PCR (RT-qPCR) analysis was

performed using a CFX384 Real-Time System (C 1000 Touch Thermal Cycler, Bio-Rad,

Hercules, California) using SensiFAST SYBR No-ROS kit (Bioline, BIO-98020) following

this set-up: (1) 95^{o}C (2 min); (2) 40 cycles of 95^{o}C (5 s), 60^{o}C (10 s), and 72^{o}C (15 s); (3)

final melt curve at 60°C (5 s), followed by continuous temperature ramping from 60°C to

99°C at a rate of 0.5°C s^{-1}. Melting curves and qPCR products were tested to ensure

there were no primer dimers or unspecific PCR products. All RT-qPCR products were

sequenced for confirmation. Primers and gene IDs for RT-qPCR were included in

Supplementary Table 1. *CBLP* (*β-subunit-like polypeptide*, Cre06.g278222), (Schloss,

1990; Xie *et al.*, 2013) and *EIF1A* (*Eukaryotic translation initiation factor 1A,*

Cre02.g103550) (Strenkert *et al.*, 2019) had stable expression among all time points, and

were used as reference genes for RT-qPCR normalization. The relative gene expressions

were calculated relative to the pre-heat time point by using the $2^{-\Delta\Delta CT}$ method (Livak and

Schmittgen, 2001; Hellemans *et al.*, 2007; Remans *et al.*, 2014). Three biological

replicates were included for each time point and treatment. RT-qPCR was used to verify
some selected transcripts and compared with the results of RNA-seq for samples
harvested under the acetate-depleting conditions (Supplemental Fig. 2). RNA-seq for
samples harvested under the constant-acetate conditions were verified by RT-qPCR
previously (Zhang *et al.*, 2022*a*).

RNA-seq analysis
Raw sequencing data for the constant-acetate experiments at 35°C and 40°C were
obtained from (Zhang *et al.*, 2022*a*). Raw sequencing data for the constant-acetate
experiment at 25°C and acetate-depleting experiments at 25°C, 35°C, and 40°C were
collected in this study. All raw sequencing data were processed using the same pipeline
as described below.

RNA libraries were prepared and sequenced by the Joint Genome Institute (JGI,
Community Science Program) using the NovaSeq platform generating 151-nt paired-end
reads. Plate-based RNA sample preparation was performed on the PerkinElmer Sciclone
NGS robotic liquid handling system using Illumina's TruSeq Stranded mRNA HT sample
prep kit utilizing poly-A selection of mRNA following the protocol outlined by Illumina in
their user guide: https://support.illumina.com/sequencing/sequencing_kits/truseq-
stranded-mrna.html, and with the following conditions: total RNA starting material was
1000 ng per sample and 8 cycles of PCR was used for library amplification. The prepared
libraries were quantified using KAPA Biosystems' next-generation sequencing library
qPCR kit and ran on a Roche LightCycler 480 real-time PCR instrument. Sequencing of
the flowcell was performed on the Illumina NovaSeq sequencer using NovaSeq XP V1.5
reagent kits, S4 flowcell, following a 2x151 indexed run recipe.

Samples were quality control filtered using the JGI BBDuk and BBMap pipelines (BBDuk
2022). Samples were quality assessed using FastQC (Andrews) and mapped to the
Chlamydomonas reinhardtii v5.6 genome (Merchant *et al.*, 2007) using HISAT2 version
2.2.0 (Kim, Langmead, and Salzberg 2015). Reads per feature were counted via
featureCounts (Liao, Smyth, and Shi 2014). The count matrix was combined with existing

data (Zhang *et al.*, 2022*a*), resulting in a 16,403x101 count matrix (Supplemental Dataset
4, 5). The normalization of the count matrix was conducted using the median of ratios
method (Love, Huber, and Anders 2014). Prior to imputation of two missing time points in
the constant-acetate-25°C condition (see details below), transcripts were filtered to have
nonzero counts in at least 90% of the samples.

Under constant 25°C, acetate, and light with turbidostatic control (constant-acetate-25°C),
the algal growth rates and cell physiologies stay constant (Fig. 5). We had RNA-seq data
for 8 time points (pre-heat, 0, 0.5, 1, 2, 24, 26, and 48 h) under constant-acetate-25°C
and these samples were highly correlated with each other (with Pearson correlation ≥
0.993) (Supplemental Fig. 1). RNA samples for the time points of 4 and 8 h under the
constant-acetate-25°C condition were unavailable; thus RNA-seq data for the 4 and 8 h
under the constant-acetate-25°C condition were imputed from other extra time points (0,
0.5, 1, 26, 48 h) of the same condition. For the time points of 4 and 8 h under the constant-
acetate-25°C condition, each transcript triplicates were sampled randomly using the 5
extra time points under the same condition mentioned above. It was ensured that both
missing time points were not imputed with the same count data. The final count matrix
consists of 14,893 transcripts in 90 samples (5 time points measured at 6 conditions as
triplicates). The natural logarithm was determined for all normalized counts before
averaging triplicates. Pearson correlation coefficients were calculated for all sample pairs.

The count data were tested for differential expression using DESeq2 v1.38.3 (Love,
Huber, and Anders 2014). DESeq2 applies a Benjamini Hochberg correction for multiple
testing. The design matrix was constructed to test for global kinetic differences between
the three different temperatures and both growth media over time. The following tests
were performed (Supplemental Dataset 1, 3, 5): (A) acetate-depleting 25°C vs acetate-
depleting 35°C; (B) acetate-depleting 25°C vs acetate-depleting 40°C; (C) constant-
acetate 25°C vs constant-acetate 35°C; (D) constant-acetate 25°C vs constant-acetate
40°C; (E) temperature effect: (constant-acetate 35°C and acetate-depleting 35°C) vs
(constant-acetate 40°C and acetate-depleting 40°C); (F) medium effect: (constant-

acetate 35°C and constant-acetate 40°C) vs (acetate-depleting 35°C and acetate-

depleting 40°C); (G) acetate-depleting 25°C vs constant-acetate 25°C.

Functional descriptions were determined for each transcript (Merchant *et al.*, 2007;

Usadel *et al.*, 2009; Venn and Mühlhaus, 2022). Transcripts were grouped according to

their functional description. Time courses of normalized transcript counts were visualized

as $\log_2$(fold change). An ANOVA was performed for each transcript at each treatment to

elucidate whether a transcript underwent a relevant change during its time course (Venn

et al., 2022a). Heatmaps (Supplemental Dataset 1) and functional sets figures

(Supplemental Dataset 2) were created using Plotly.NETv4.0.0 (Schneider *et al.*, 2022).

Ontology enrichment was performed using extended MapMan annotations (e.g.

"PS.lightreaction.LHC" became "PS.Lightreaction.LHC", "PS.Lightreaction" and "PS")

(Supplemental Dataset 3). The measured and filtered transcripts served as background.

If a transcript showed significant differential expression in the respective comparison

(FDR < 0.05), it was considered as significant for its annotations. Enrichment p values

were determined using hypergeometric tests. Multiple testing correction was performed

using the Benjamini-Hochberg method (Benjamini and Hochberg, 1995; Venn *et al.*,

2022).

Results

To investigate how the availability of organic carbon source affected algal heat responses,

we first cultivated WT Chlamydomonas cells (CC-1690, *21gr*) in PBRs in Tris-acetate-

phosphate (TAP, acetate as carbon source) medium at 25°C with constant nutrient supply

through turbidostatic control (providing frequent fresh medium and culture dilution) (Fig.

1A). After algal cultures reached steady growth rates in PBRs, the turbidostatic control

was turned off and cultures were switched to 35°C, or 40°C, or stayed at 25°C for 24 h

without constant acetate supply, defined as the acetate-depleting condition. This

experiment allowed us to quantify acetate consumption under different temperatures,

which could not be performed in cultures with turbidostatic control or frequent medium

dilution. It also enabled us to investigate algal heat responses under the acetate-depleting

condition, similar to shaker cultures, which were used by most previous algal heat

experiments in the published literatures (Hemme *et al.*, 2014; Rütgers *et al.*, 2017). Under

the acetate-depleting condition, algal cultures were harvested at different time points to

analyze cell physiologies, biomass, and transcriptome (Fig. 1B). Algal dry biomass

quantification showed that cultures treated with 35°C had increased biomass at 8-h heat

but decreased biomass at 24-h heat as compared to 25°C (Fig. 1C). Acetate quantification

in the supernatant of algal cultures showed that 35°C-treated cultures had increased

acetate consumption and depleted acetate faster than 25°C or 40°C (Fig. 1D).

We hypothesized the transiently increased and then decreased algal biomass in 35°C-

treated cultures was due to increased acetate uptake/usage followed by acetate

starvation in the end under the acetate-depleting condition (Fig. 1). Chlamydomonas

uptakes acetate and feeds it into the glyoxylate and gluconeogenesis cycles to produce

starch, which can be used in glycolysis to generate ATP (Johnson and Alric, 2012, 2013)

(Fig. 2A). To verify this hypothesis, we performed RNA-seq analysis and investigated key

transcripts involved in the acetate uptake/assimilation, glyoxylate, gluconeogenesis, and

glycolysis cycles under the acetate-depleting condition by comparing with the constant-

acetate condition (Fig. 2B-P). Under the acetate-depleting condition, almost all these

transcripts related to acetate uptake/metabolism were up-regulated under 35°C as

compared to 25°C. Similar 35°C effects on these transcripts were held under the constant-

acetate condition, although the up-regulation of these transcripts were stronger and lasted

longer at 35°C under the constant-acetate condition than the acetate-depleting condition.

In contrast, these transcripts related to acetate uptake/metabolism were down-regulated

under 40°C as compared to 25°C under both the acetate-depleting and constant-acetate

conditions. Our RT-qPCR results of selected transcripts showed similar trends as our

RNA-seq data (Supplemental Fig. 2). These results showed that the 35°C promoted

acetate metabolism, from acetate uptake/assimilation to glyoxylate, gluconeogenesis,

and glycolysis cycles; however, these acetate metabolism pathways were reduced under

40°C.

13-40

Besides the acetate metabolism pathways mentioned above, we also checked transcripts
related to other processes that are often affected by high temperatures (Supplemental
Fig. 3, 4). Chlamydomonas has 2 heat shock transcription factors, HSF1 and HSF2; HSF1
is the master heat regulator but the function of HSF2 is unclear (Schulz-Raffelt *et al.*,
2007). The transcripts of both *HSF*s were induced by heat, with higher induction at 40°C
than 35°C under the same acetate condition or higher induction under the acetate-
depleting than constant-acetate condition at the same high temperature (Supplemental
Fig. 3A, B). Heat shock proteins (HSPs) are the hallmarks of heat responses and HSPs
function as chaperones to help refold denatured proteins, fold nascent polypeptides, and
prevent denatured protein aggregates (Altschuler and Mascarenhas, 1982; Schroda *et*
al., 2015; Haq *et al.*, 2019). Based on molecular mass, HSPs are grouped to large HSPs
(HSP100/90/70/60, >50 kDa) and small HSPs (10-43 kDa) (Jee, 2016; Bourgine and
Guihur, 2021). Under the acetate-depleting condition, many transcripts related to large
HSPs, *HSP60/70/90s*, were down-regulated at 24 h of 25°C and 35°C; such down-
regulation of *HSP60/70/90s* was absent at 24 h of 25°C and 35°C under the constant-
acetate condition or 24 h of 40°C under both acetate conditions (Supplemental Fig. 4A-
C). Additionally, transcripts of *HSP100* and most of the *small HSPs* did not have such
down-regulation at 24 h of 25°C and 35°C under the acetate-depleting condition
(Supplemental Fig. 4D, E). Almost all transcripts related to *HSPs* we examined, including
HSP60/70/90/100s and *small HSPs*, were up-regulated at 2 h of 40°C, with similar
induction fold change under either the acetate-depleting or constant-acetate condition
(Supplemental Fig. 4A-E). It suggested the early induction of these *HSPs* under 40°C
were temperature-dependent (present at 40°C while much less so under 35°C) but
acetate-availability-independent. Alternatively, the acetate-availability may not differ that
much at 2 h of the acetate-depleting and constant-acetate condition (Fig. 1D).

The global transcriptome changes related to light reaction and Calvin-Benson cycle of
photosynthesis suggested more disturbance of these transcripts under the acetate-
depleting than the constant-acetate conditions at the same temperatures, and more
disturbance of these transcripts at 40°C than 35°C under the same acetate conditions
(Supplemental Fig. 4F-M). Chlamydomonas employs a cellular structure, pyrenoid, to

concentrate CO_2 for improved carbon fixation efficiency, called carbon concentrating
mechanisms (CCM) (Meyer *et al.*, 2017; Hennacy and Jonikas, 2020). Under the acetate-
depleting conditions, several transcripts related to CCM were up-regulated at 24 h under
25°C and 35°C, but not under the same temperature of the constant-acetate condition,
suggesting the induction of CCM when acetate was depleted, and cell densities increased
at 24 h of 25°C and 35°C under the acetate-depleting condition (Supplemental Fig. 4N).
Such induction of the CCM transcripts was absent at 40°C under either acetate condition,
which may correlate with the altered pyrenoid structures at 40°C (Zhang *et al.*, 2022*a*).
Transcripts related to photoprotection had larger induction at 24 h under all three
temperatures of the acetate-depleting than the constant-acetate conditions at the same
temperature, indicating compromised photosynthesis and the need for photoprotection
when acetate was depleted (Supplemental Fig. 4O).

The compromised photosynthesis under high temperatures lead to increased
accumulation of reactive oxygen species (ROS) (Pospíšil, 2016; Janni *et al.*, 2020;
Niemeyer *et al.*, 2021). Some transcripts related to ROS scavenging were down-regulated
at 24 h of 25°C and 35°C under the acetate-depleting condition but much less so under
the constant-acetate condition at the same temperatures (Supplemental Fig. 4R-V). This
may be related to the increased cell density and reduced light/oxidative stress at 24 h of
25°C and 35°C under the acetate-depleting condition. Such down-regulation of transcripts
related to ROS scavenging was not present at 24 h of 40°C under the acetate-depleting
condition. This may suggest relatively increased ROS accumulation and ROS scavenging
activity at 40°C than 25°C and 35°C under the acetate-depleting conditions. Additionally,
transcripts related to ROS scavenging had very similar transcriptional changes at 40°C
under either the acetate-depleting or constant-acetate condition, suggesting 40°C had
more dominant effects on the ROS accumulation and ROS scavenging activities than
acetate.

Transcripts related to cell cycle and cell wall had similar transcriptional patterns under
each of the six conditions (Supplemental Fig. 4W, X). Under diurnal cycles, cell wall genes
were up-regulated following the up-regulation of cell cycle genes and cell walls form

around daughter cells during cell division (Zones *et al.*, 2015). These transcripts were

down-regulated at 24 h of 25°C under the acetate-depleting condition, suggesting the

transition to cell cycle arrest due to acetate-depletion. The reduction of these transcripts

was larger and more prolonged under 40°C than 35°C, consistent with complete and

transient inhibition of cell cycle under 40°C and 35°C, respectively (Hemme *et al.*, 2014;

Zhang *et al.*, 2022*a*). At 24 h of 40°C and 35°C, these transcripts were more reduced

under the acetate-depletion condition than the constant-acetate condition.

Almost all transcripts mentioned above had little changes at 25°C during all time points

of the constant-acetate condition, validating the steady growth and constant cell status

under the turbidostatic control with constant nutrient/light but without heat treatments in

our algal cultures grown in PBRs (Supplemental Fig. 4).

We performed statistical analysis to investigate global transcript kinetic differences

between different conditions (Supplemental Dataset 1, 3). We then applied ontology

enrichments analysis for transcripts that were significantly expressed in a defined

comparison to identify differentially regulated biological processes (Supplemental Fig. 5,

Supplemental Dataset 3). The top enriched function group that differed at 25°C and 35°C

under both acetate-depleting and constant-acetate condition is the nucleus-encoded

ribosomal proteins that are involved in chloroplast protein synthesis (Supplemental Fig.

5A, C). More than half of the ribosomal related transcripts were down-regulated at 4 h

heat of 35°C as compared to 25°C under both acetate conditions (Supplemental Fig. 4Y,

Z), suggesting 35°C may compromise chloroplast function. The top enriched function

group that differs between 25°C and 40°C under the acetate-depleting condition is the

mitochondrial electron transport (Supplemental Fig. 5B). Many transcripts related to

mitochondrial electron transport were down-regulated at 2 h heat of 40°C as compared

to 25°C under the acetate-depleting condition (Supplemental Fig. 4AA, AB). When

comparing 35°C and 40°C regardless the acetate availability, the top enriched function

group is also the mitochondrial electron transport (Supplemental Fig. 5E). The results

suggested that mitochondrial electron transport was more sensitive to 40°C than 25°C

and 35°C. Under the constant-acetate condition, 40°C affected RNA processes the most

as compared to 25°C (Supplemental Fig. 5D, 4AC, AD). When comparing the acetate-
depleting and constant-acetate conditions regardless the high temperatures, the top two
enriched groups are phosphate transport and mitogen-activated protein kinase (MAP
kinase) cascades signaling (Supplemental Fig. 5F). MAP kinase signaling pathways has
important roles in plant growth and stress responses and involve protein
phosphorylation/dephosphorylation (Bigeard and Hirt, 2018; Jagodzik *et al.*, 2018; Zhang
and Zhang, 2022). The biggest differences in transcripts related to the MAP kinase
signaling pathways between the acetate-depleting and constant-acetate conditions were
at the 24 h time point at 25°C and 35°C when the acetate was depleted (Supplemental
Fig. 4AF), suggesting potential roles of the MAP kinase signaling pathways in nutrient
sensing and growth regulation. When comparing the acetate-depleting and constant-
acetate conditions at 25°C, the top enriched function groups are dominantly related to cell
motility (Supplemental Fig. 5G). More than half of the transcripts related to cell motility at
25°C were down-regulated at 24 h under the acetate-depleting condition but not the
constant-acetate condition (Supplemental Fig. 4AG, AH). Microscopic images showed
that cells had flagella at 24 h of 25°C under the acetate-depleting condition (Supplemental
Fig. 6). Thus, the down-regulation of transcripts related to cell motility may suggest the
transition from motile to non-motile phase when the acetate was depleted. Cell motility
needs energy and ciliary beating frequency corelates with the cytosolic ATP concentration
in Chlamydomonas (Takano *et al.*, 2021).

To further understand our transcriptome data and the effects of acetate, we compared
cell physiologies at different temperatures between the acetate-depleting and constant-
acetate conditions (Fig. 3). Under the acetate-depleting condition without medium dilution,
cell density increased over time at 25°C and 35°C but no changes at 40°C because of
cell cycle arrest at 40°C (Fig. 3A). Cell density was lower at 35°C than 25°C at 8 h and 24
h time points, suggesting reduced growth rates at 35°C at the end of the acetate-depleting
condition. Algal cells treated at 35°C had transiently increased then decreased cell size
while 40°C-treated algal cells had steadily increased cell size due to heat-inhibited cell
division (Fig. 3B). Under the acetate-depleting condition, chlorophyll content per culture
volume (mL) had no significant differences between 25°C and 40°C but was significantly

higher at 8-h of 35°C than 25°C (Fig. 3C), consistent with the increased biomass at the 8-h of 35°C (Fig. 1C). Chlorophyll content normalized to cell density revealed constant chlorophyll per cell during 25°C, but transiently and constantly increased cellular chlorophyll during 35°C and 40°C, respectively (Fig. 3D). Chlorophyll content normalized to cell volume showed the increased cellular chlorophyll during 40°C and 35°C could not be completely explained by increased cell volume (Fig. 3E).

Based on our previously published data, we summarized cell parameters in algal cultures grown in the same PBRs but under the constant-acetate condition through turbidostatic control for 24 h (Zhang *et al.*, 2022*a*) (Fig. 4F-J). Cell parameters at different time points during 25°C had little or no changes as compared to the pre-heat time points, demonstrating the steady algal growth and effectiveness of the turbidostatic control with constant acetate supply. Under the constant acetate condition, the smaller changes of cell density and chlorophyll per mL culture as compared to the acetate-depleting conditions were due to the turbidostatic control and frequent dilution of the culture (Fig. 3F and 3H). The changes of cell volume and chlorophyll content per cell during 35°C and 40°C had the similar trends under both the acetate-depleting and constant-acetate condition, but the increase of these parameters was larger under 40°C in cultures with the constant-acetate than the acetate-depleting condition (Fig. 3B, D, G, I). This was supported by the fold changes of cell parameters by comparing the data under the acetate-depleting and constant-acetate conditions (Fig. 3K-O).

In addition to the cellular parameters mentioned above using time-course harvesting, we next utilized non-disruptive methods to quantify algal growth under different temperatures. The acetate-depleting condition without turbidostatic control allowed us to have a large range of cell densities under different temperature treatments to investigate the optical density (OD) vs cell density relationship and validate automatic culture monitoring under different temperature treatments. OD_{680} (optical density at 680 nm) monitors chlorophyll content per mL culture (Chapman *et al.*, 2015; Xiao *et al.*, 2015; Young *et al.*, 2022). OD_{750} (optical density at 750 nm) monitors light scattering and is thought to be proportional to cell density or cell volume (Chioccioli *et al.*, 2014; Young *et al.*, 2022). Our

PBRs can measure OD_{680} and OD_{720}, but no OD_{750}. However, OD_{720} serves as a proximity

for OD_{750} for light scattering. Thus, we investigated the dynamic changes of OD_{680} and

OD_{720} relative to chlorophyll content, cell density, and cell volume before and during heat

treatments under the acetate-depleting condition, in medium starting with acetate but

without constant acetate supply (Fig. 4). The change of OD_{680} and OD_{720} (Fig. 4A, B)

mimicked the change of chlorophyll content per mL algal culture for all three temperature

treatments (Fig. 3C). But the change of OD_{720} (Fig. 4B) mimicked the change of cell

density for 25^oC and 35^oC but not 40^oC (Fig. 3A). Combining all data from different

temperature treatments and time points, both OD_{680} and OD_{720} were linearly proportional

to chlorophyll content per mL algal culture (Fig. 4C, D), but they were much less

proportional to cell volume or cell density (Fig. 4E-J). Our results showed that OD_{680} and

OD_{720} can be used to accurately estimate chlorophyll content in algal cultures with

different heat treatments, thus they can be used to estimate algal relative growth rates

based on chlorophyll accumulation during an exponential growth phase between dilution

events with turbidostatic control in PBRs (Zhang *et al.*, 2022*a*) (See methods for details).

The relative growth rates calculated from both OD_{680} and OD_{720} yielded similar results.

Because OD_{680} had larger values and higher signal/noise ratios than OD_{720}, we used

OD_{680} to estimate relative growth rates of PBR cultures over several days under the

turbidostatic controls with and without acetate supply to investigate how acetate affected

algal heat responses at 35^oC and 40^oC in long-term (Fig. 5-9). This research goal could

not be achieved without turbidostatic controls because our acetate-depleting experiment

revealed that acetate was fully depleted in 24 h during all three temperature treatments

of 25^oC, 35^oC and 40^oC without constant acetate supply (Fig. 1D).

To investigate the effects of acetate on long-term heating, after algal cultures reached

steady growth at 25^oC in PBRs with turbidostatic control, we conducted all three

temperature treatments with continuously turbidostatic control and constant acetate

supply for 4 days (Fig. 5). Without heat treatments, algal growth rates and cell parameters

stayed constant at 25^oC, demonstrating the effectiveness of our turbidostatic control for

steady algal growth (Fig. 5). At 35^oC with constant acetate supply, the relative growth

rates increased first, but the increase was reduced to a smaller degree after 2-day heating

at 35°C and stabilized thereafter (Fig. 5A). At 40°C, the relative growth rates decreased

steadily, and the cultures died after 2-day heating at 40°C, suggesting algal cells could

not acclimate to long-term constant heat at 40°C, even with constant acetate supply. Cells

treated with 40°C had more than 4-fold increase of cell volume as compared to the pre-

heat condition, followed by reduced cellular chlorophyll and cell death (Fig. 5C-F). The

turbidostatic mode by OD_{680} tightly controlled the chlorophyll contents in unit of ug per mL

culture in PBRs during the treatments of 25°C and 35°C (Fig. 5D), but not 40°C due to

the cell cycle arrest and overwhelmingly increased chlorophyll per cell, and eventually

cell death under 40°C (Fig. 5B and E).

To further test the effects of acetate on algal heat responses, we conducted a different

sets of experiments using algal cultures grown in PBRs, under the turbidostatic control

with constant nutrient supply but in photoautotrophic medium (no acetate), with different

temperature treatments for 4 days (Fig. 6). Without heat treatments, algal growth rates

and cell parameters stayed constant at 25°C, demonstrating the effectiveness of our

turbidostatic control for steady algal growth under photoautotrophic conditions. Without

acetate but with constant medium supply, algal growth had gradual decreased growth

rates during 35°C (Fig. 6A), unlike the increased growth rates during 35°C with constant

medium and acetate supply (Fig. 5A), strongly supporting the role of acetate in algal heat

tolerance. Without acetate, algal cells with 35°C treatment had increased cell volume and

chlorophyll contents (Fig. 6B-F). Our microscopic images showed that the dramatically

increased cell volume during 35°C was due to the formation of palmelloids (Fig. 7).

Palmelloids are clumps of non-motile cells, which often results from a failure of hatching

or release from the mother cell wall under stressful conditions (Harris, 2009). It is reported

that Chlamydomonas cells aggregated to form palmelloids after 3 min acute heat shock

at 50°C in the acetate-containing medium (de Carpentier *et al.*, 2022). The dramatically

increased cell volume due to palmelloids did not occur in the constant-acetate-35°C

condition (Fig. 5C), suggesting 35°C was more stressful without acetate than with acetate.

The reduction of growth rates during 40°C was comparable with and without acetate (Fig.

5A and 6A).

To better compare algal growth and cell physiologies under the turbidostatic control with

and without acetate, we normalized the data with heat treatments to the constant 25°C

with the same medium type (Fig. 8). Algal cells treated with 35°C without acetate had

similar pattern of cell density and cell volume as algal cells treated with 40°C with acetate

(Fig. 8B, C), supporting 35°C was more stressful without acetate than with acetate. The

normalized data also revealed the increased chlorophyll per mL culture, per cell, and per

cell volume during 35°C without acetate (Fig. 8D-F).

Our PBRs could also monitor photosystem II (PSII) operating efficiency using chlorophyll

fluorescence measurement automatically (Fig. 9). Heat at 40°C reduces PSII operating

efficiency dramatically, with more reduction without acetate than with acetate during the

first two days of heating (Fig. 9B, C, D). The acetate effects on PSII operating efficiency

disappeared in algal cells treated by 40°C for more than 2 days (Fig. 9D). The reduction

of PSII operating efficiency during 35°C was much smaller than during 40°C, with higher

PSII operating efficiency in 35°C-treated algal cells with acetate than without acetate (Fig.

9D).

Discussion

By performing algal cultivation under highly controlled conditions in PBRs with and without

a constant organic carbon source, acetate, we investigated how the availability of acetate

affected the growth of Chlamydomonas under moderate (35°C) and acuate high

temperatures (40°C) (Fig. 10).

Heat of 35°C was beneficial or detrimental depending on carbon availability

Under the turbidostatic control without nutrient limitation, moderate high temperature of

35°C increased algal growth rates with constant acetate supply (Fig. 5A), but decreased

agal growth rates without acetate in photoautotrophic medium (Fig. 6A), confirming the

important role of acetate in heat tolerance. The increased heat tolerance to 35°C with

acetate is not strain-dependent because the acetate effects were similar in two different

algal strains: CC-1690 (this study) and CC-5325 (Mattoon *et al.*, 2022). In medium starting

with acetate but without constant acetate supply (the acetate-depleting condition), algal

biomass increased first but then decreased at 35°C compared to 25°C, which can be

explained by increased acetate uptake/usage initially followed by acetate

depletion/starvation by the end of the 24-h 35°C treatments (Fig. 1C, D).

Under the acetate-depleting condition, besides acetate, other nutrients may also be

depleted gradually during 24 h. We checked known transcription factors and regulators

responding to starvation of other nutrients, including *nitrogen starvation responsive*

regulator 1 (NRR1) (Boyle *et al.*, 2012), *sulfur starvation responsive regulator 2.1*

(SNRK2.1) (González-Ballester *et al.*, 2010), *phosphorus starvation response 1 (PSR1)*

(Bajhaiya *et al.*, 2016), *copper deficiency response regulator 1 (CRR1)* (Sommer *et al.*,

2010), and *iron deficiency response regulator (MAPK3)* (Fei *et al.*, 2017). If these

nutrients were depleted, the corresponding transcription factors and regulators would be

up-regulated. Most of these nutrient-deficiency transcription factors and regulators had

minimal or no induction at 24 h of the acetate-depleting condition (Supplemental Fig. 3C-

G). One exception is the *NRR1*, which was up-regulated at 25°C or 35°C but was down-

regulated under 40°C at 24 h of the acetate-depleting condition (Supplemental Fig. 3C).

The results suggested minimal or no limitation of sulfur, phosphorus, copper, and iron

during the acetate-depleting condition, but some nitrogen deficiency at 24 h of the

acetate-depleting condition. One of the responses to nitrogen deficiency is reduction of

chlorophyll (Schmollinger *et al.*, 2014). During our acetate-depleting condition, most

transcripts involved in chlorophyll biosynthesis were stable (Supplemental Fig. 4Q).

Additionally, there was no change of the chlorophyll per cell or per cell volume at 24 h of

25°C as compared to the pre-heat time point which did not have acetate-limitation; these

parameters were higher under 35°C and 40°C than 25°C at 24 h of the acetate-depleting

condition (Fig. 3D, E). Collectively, the results indicated that the nitrogen limitation may

have just started at 24 h of the acetate-depleting condition, thus would not impact the cell

physiology too much. Finally, our results from the 4-day constant-acetate and no-acetate

conditions under the turbidostatic control strongly pointed to the effects of acetate on algal

heat tolerance but not other nutrients (Fig. 5, 6). Thus, the acetate effects are the focus

of our research.

Chlamydomonas uptakes acetate and feeds it into the glyoxylate cycle and
gluconeogenesis for starch biosynthesis and starch is broken down through glycolysis to
make cellular energy (Johnson and Alric, 2012, 2013) (Fig. 2A). The dynamic effects of
$35^{o}C$ on algal growth under different acetate conditions can be explained by the $35^{o}C$-
induced carbon metabolisms, including the up-regulation of transcripts related to acetate
uptake/assimilation, glyoxylate cycle and gluconeogenesis pathways, and glycolysis (Fig.
2). The up-regulation of these transcripts lasted longer with higher induction under the
constant-acetate than the acetate-depleting condition (Fig. 2). Energy produced from
glycolysis can be used for energy-requiring cellular activities to increase thermotolerance
(Olas *et al.*, 2021), e.g., production of heat shock proteins and repair pathways related to
photosynthesis (Murata and Nishiyama, 2018; Bourgine and Guihur, 2021). Pyruvate
kinase catalyzes the final step of glycolysis and it converts phosphoenolpyruvate and one
ADP to pyruvate and one ATP (Baud *et al.*, 2007; Wulfert *et al.*, 2020). Chlamydomonas
mutants deficient in pyruvate kinase were heat-sensitive under $35^{o}C$ in acetate-containing
medium (Mattoon *et al.*, 2022), supporting the important roles of glycolysis in
thermotolerance of $35^{o}C$. Although ATP production mainly comes from mitochondrial
respiration, the ATP production from glycolysis can be important under stressful
conditions where energy availability is limited (van Dongen *et al.*, 2011). Under low
oxygen conditions, plants increase activity of pyruvate kinases to produce more ATP (van
Dongen *et al.*, 2011). Heat-treated barley leaves utilized glycolysis as an alternative
energy source for thermotolerance based on proteomics analysis (Rollins *et al.*, 2013).
699

Additionally, acetate may protect photosynthesis from heat-induced photoinhibition.
Acetate is proposed to protect PSII against photoinhibition by replacing the bicarbonate
associated to the non-heme iron at the acceptor side of PSII, changing the environment
of plastoquinone, and affecting PSII charge recombination (Roach *et al.*, 2013). Our PSII
efficiency data supported this hypothesis because PSII efficiency was higher with acetate
than without acetate during $25^{o}C$, $35^{o}C$ or $40^{o}C$ treatments (Fig. 9D). Chlamydomonas
grown in acetate-containing medium produced less singlet O_2 (one kind of ROS) than
those grown in non-acetate-containing, photoautotrophic medium (Roach *et al.*, 2013).
Furthermore, experimental and modeling analysis suggested that acetate promoted cyclic

electron flow (CEF) around photosystem I (PSI); reducing equivalents produced during

the acetate metabolism reduce plastoquinone pools and increase CEF activity (Johnson

and Alric, 2012; Lucker and Kramer, 2013; Chapman *et al.*, 2015). CEF cycles

photosynthetic electrons around PSI, producing only ATP but no NADPH and providing

extra ATP needed for photosynthesis and other cellular activities (Munekage *et al.*, 2004;

Baker *et al.*, 2007). CEF balances the ATP/NADPH ratio, contributes to the generation of

transthylakoid proton motive force, and protects both PSI and PSII from photo-oxidative

damage (Johnson, 2011; Yamori and Shikanai, 2016). The increased acetate uptake

under $35^{\circ}C$ (Fig. 1D) was coupled with induced CEF activity measured by P700

oxidation/reduction in Chlamydomonas (Zhang *et al.*, 2022*a*). Our transcriptome analysis

also showed up-regulation of transcripts related to PSI and CEF at 2h of $35^{\circ}C$ treatments

under either the acetate-depleting or constant-acetate condition, consistent with heat-

induced CEF in the presence of acetate (Supplemental Fig. 4J, P).

The effects of acetate on the improved heat tolerance under the moderate high

temperature may also be related to the accumulation of thermoprotective metabolites

(osmolytes, including amino acids, sugars, and sugar alcohols) (Guihur *et al.*, 2022) or

compatible solutes (Hemme *et al.*, 2014; Schroda *et al.*, 2015). Compatible solutes

accumulate from the catabolism of large molecules upon the onset of heat to stabilize

membranes and prevent protein unfolding/aggregation (Schroda *et al.*, 2015). In

Chlamydomonas, these compatible solutes include glycerophos-phoglycerol (GPG),

trehalose, β_alanine, isoleucine, asparagine, glutamine, and guanosine (Hemme *et al.*,

2014). The elevated acetate metabolism at $35^{\circ}C$ with constant acetate supply may

increase the accumulation of these thermoprotective metabolites. This is an understudied

but exciting area. Future metabolite analysis may provide more information about this.

With constant acetate supply, $35^{\circ}C$ is beneficial, increasing carbon metabolisms and

energy production, thus improving thermotolerance and growth (Fig. 5). If algal cells can

be cultivated under sterile conditions with constant organic carbon supply, $35^{\circ}C$ could be

used to promote algal growth and increase biofuel/bioproduct generation under

mixotrophic conditions in light. Culture temperature of $35^{\circ}C$ in closed, outdoor algal ponds

may not be difficult to reach with natural sunlight heating in the summertime of the

moderate climate regions and year-round in the tropical regions.

Without constant acetate supply, 35°C is detrimental. Under the acetate-depleting

condition, the 35°C-elevated carbon metabolism without sufficient carbon input may

deplete cellular carbon reserves. This was not sustainable in the long term and eventually

reduced biomass accumulation (Fig. 1C). With constant acetate supply, heat at 35°C

transiently arrested the cell cycle, which recovered after 8 h at 35°C (Fig. 3G). In

photoautotrophic medium without acetate, 35°C reduced algal growth rates and inhibited

cell division (stressed cells aggregated to form palmelloids) (Fig. 6, 7). Such effects of

35°C can particularly compromise the yields of outdoor algal ponds because they usually

do not contain carbon sources but frequently experience moderate high temperatures

(Mata *et al.*, 2010; El-Sheekh *et al.*, 2019).

It is reported that the growth rate of another Chlamydomonas WT strain (CC-125) at 35°C was

about 50% of that at 28°C (0.083 vs 0.166) in photoautotrophic medium under constant light and

continuous dilution but with 2% CO_2 (Vítová *et al.*, 2011). The 50% growth reduction is

comparable to the reduced growth of our data at 35°C without acetate (Fig. 6A). We used air level

CO_2, which is 400 ppm or 0.04% CO_2 to mimic the nature condition (1/50th of 2% CO_2 as in (Vítová

et al., 2011)). CO_2 provides the inorganic carbon source for algal growth. But the equivalent 50%

growth reduction at 35°C as compared to the control temperature with (Vítová *et al.*, 2011) and

without (this study) high CO_2 suggests CO_2 (inorganic carbon source) is not as effective as acetate

(organic carbon source), to improve thermotolerance in algal cells. Furthermore, different algal

strains with changes in cultivation conditions may have varying moderate high temperature

ranges. For example, another Chlamydomonas WT strain (CC-137) had optimal growth at 35°C

in photoautrophic medium with 2% CO_2 in day/night cycle with semi-dilution (one dilution per day

at the end of dark phase) (Lien and Knutsen, 1979). Such semi-dilution may result in a large range

of cell densities, which may affect the light perceived by each cell. Additionally. the day/night cycle

may also alleviate photooxidative stress as photosynthesis only occurs in light. However, the

author mentioned that cytokinesis and sporulation became inhibited at temperature above 36°C

(Lien and Knutsen, 1979), indicating 35°C may be close to the moderate high temperature range

that would reduce algal growth in this strain under their cultivation condition without acetate.

The up-regulated carbon metabolism under moderate high temperatures may also occur

in land plants, especially in sink tissues (e.g., roots) or mixotrophic tissues (e.g.,

developing green seeds) (Koley *et al.*, 2022). Heat-induced glyoxylate cycle,

gluconeogenesis, and glycolysis have been reported in plants (Rollins *et al.*, 2013; Zhang

et al., 2013; Aprile *et al.*, 2013). Carbon metabolism and sugar availability were shown to

be essential for heat tolerance in Arabidopsis (Olas *et al.*, 2021).

Heat of 40°C was detrimental to algal cells with and without constant acetate supply

Acute high temperatures at or above 40°C inhibit the algal cell cycle (Fig. 3, 5-7)

(Mühlhaus *et al.*, 2011*b*; Hemme *et al.*, 2014; Zachleder *et al.*, 2019; Ivanov *et al.*, 2021),

alter thylakoid membranes, reduce photosynthesis, and damage respiration (Zhang *et al.*,

2022*a*). These cellular damages took place quickly, within 4 h of heat at 40°C with

constant acetate supply (Zhang *et al.*, 2022*a*). However, 40°C-treated cells grew bigger

and had more chlorophyll under the constant-acetate condition than the acetate-depleting

condition in the short-term (24-h heat) (Fig. 3B, G, D, I, L, N), although the constant

acetate supply could not prevent ultimate cell death with 40°C-heat longer than 2-days

(Fig. 5). Similarly, without acetate, Chlamydomonas cells also died after 2-day heat at

40°C (Fig 6). The death of Chlamydomonas cultures was also reported at 39°C for 33 h

in photoautotrophic medium without acetate (Zachleder *et al.*, 2019; Ivanov *et al.*, 2021).

Thus, constant heating at 40°C is lethal for Chlamydomonas cells independent of carbon

availability.

Transcripts of many genes involved in carbon metabolism, e.g., acetate

uptake/assimilation, glyoxylate cycle and gluconeogenesis pathways, and glycolysis were

significantly down-regulated during 40°C (Fig. 2). Most of the reducing power from acetate

assimilation is used in mitochondrial respiration to produce ATP (Johnson and Alric, 2012,

2013). Our previous results showed that 40°C reduced mitochondrial activities

significantly under the 24-h constant-acetate condition (Zhang *et al.*, 2022*a*). When

comparing 40°C vs 25°C under the acetate-depleting condition and 40°C vs 35°C

regardless the acetate condition, the top enriched function group was related to

mitochondrial electron transport (Supplemental Fig. 5B, E), suggesting mitochondria were

particularly sensitive to 40°C. The compromised mitochondrial activities may restrict
acetate uptake/assimilation during 40°C heat treatment. Under the acetate-depleting
conditions, the acetate uptake was slightly lower in 40°C-treated cultures than 25°C (Fig.
1D). The decreased acetate uptake/usage, together with reduced photosynthesis and
respiration (Zhang *et al.*, 2022*a*) and cell cycle arrest (Fig. 3, 5-7), may contribute to the
significantly reduced algal dry biomass in cultures treated with 8h-heat of 40°C than 25°C
(Fig. 1C). By the end of 24-h treatment under the acetate-depleting condition, there were
comparable amount of dry biomass in cultures treated with 40°C and 25°C (Fig. 1C),
which could be explained by the arrested cell cycle and almost eliminated energy/biomass
consumption for cell division during 40°C. Such an inefficient system with reduced carbon
input and little energy output was not sustainable, eventually killing all the algal cells by
the end of 2-day heat of 40°C, even with constant acetate supply (Fig. 5). Our results
suggested that the increased thermotolerance by acetate did not apply to long-term heat
at 40°C.
818
At 24-h of the acetate-depleting condition, the transcripts related to large *HSP*s,
HSP60/70/90s, were down-regulated at 25°C and 35°C, but not under 40°C
(Supplemental Fig. 4A-C). The down-regulation of large *HSP*s may be related to the
increased cell density and light shading, thus reduced light stress at 24 h of 25°C and
35°C under the acetate-depleting condition (Fig. 3A). However, the amount of chlorophyll
per mL culture was comparable at 24 h of 25°C, 35°C, and 40°C under the acetate-
depleting condition, suggesting comparable level of light shading and light stress under
all three temperatures (Fig. 3A), although photosynthesis may have more damages at
40°C than 25°C/35°C. The up-regulated transcripts related to photoprotection at 24 h of
all three temperatures under the acetate-depleting condition may deemphasize the
effects of light shading (Supplemental Fig. 4O). Alternatively, the different regulation of
large *HSP*s between 25°C/35°C and 40°C at 24 h of the acetate-depleting condition may
suggest the energy distribution between growth and heat tolerance. Most large HSPs are
ATP-dependent chaperons, especially HSP90s (Al-Whaibi, 2011; Xu *et al.*, 2012; Guihur
et al., 2022). Down-regulation of these large HSPs when acetate was depleted may be
helpful to reserve energy for cell growth when the temperature stress was none (25°C) or

moderate (35°C). However, under the acute heat of 40°C, thermotolerance has higher

priority than growth, thus the expression of these large *HSPs* were maintained while cell

cycle was arrested to preserve energy.

As compared to 40°C, the induction of *HSP*s at 35°C was minimal or none under either

acetate condition (Supplemental Fig. 4A-C), suggesting the 35°C-induced

thermotolerance by acetate may not be related to the accumulation of HSPs. This may

suggest some none-overlapping heat sensing/signaling pathways between 35°C and

40°C, as HSPs have important roles in the classical heat shock responses (Guihur *et al.*,

2022). Several heat sensors have been proposed (Vu *et al.*, 2019), including membrane

fluidity (Martinière *et al.*, 2011), Ca^{2+} channels in the plasma membrane (Gao *et al.*, 2012;

Finka *et al.*, 2012), unfolded proteins (Rütgers *et al.*, 2017; Li *et al.*, 2020), and

phytochromes (Jung *et al.*, 2016; Legris *et al.*, 2016), and some circadian clock

components (e.g. EARLY FLOWERING 3, ELF3) (Jung *et al.*, 2020). Among these, the

heat effects on phytochromes and ELF3 have been reported around the temperature

ranges of 30-35°C (Jung *et al.*, 2016, 2020; Legris *et al.*, 2016), providing potential heat

sensor candidates for moderate high temperatures. Future research to investigate the

shared and unique heat sensors for moderate and acuate high temperatures may provide

more information about how plants sense and adapt to different high temperatures.

Most previous heat treatments in Chlamydomonas used cultures grown with acetate-

containing medium in flasks without turbidostatic controls (Hemme *et al.*, 2014; Rütgers

et al., 2017). Our results showed that such heat treatment could not last more than 8 h at

35°C or 24 h of 40°C due to acetate-depletion (Fig. 1D). In fact, many previous heat

experiments were conducted for a short-term, e.g., 1 h (Légeret *et al.*, 2016; Rütgers *et*

al., 2017). Using our highly controlled PBR systems, we could not only conduct heat

experiments under the 24-h acetate-depleting condition but also during 4-day heating with

and without acetate under the turbidostatic control with constant nutrient supply to

uniquely reveal the effects of carbon availability on thermotolerance.

In summary, by using highly controlled cultivation systems in PBRs and the model green

alga Chlamydomonas, we revealed how the availability of an organic carbon source,

acetate, interacted with different intensities and duration of high temperatures in

photosynthetic cells. Our research revealed the overlooked effects of moderate high

temperature of 35°C, which can be beneficial with constant carbon supply or detrimental

with insufficient or no carbon supply. We also showed that the damaging effects of acute

high temperature of 40°C is dominant and independent of carbon availability. Our

research not only helps us understand heat responses in photosynthetic cells but also

provides insights for high temperature effects on the production of algal biofuels and

bioproducts.

Supplementary data

Supplemental Table 1: Gene IDs and primers for RT-qPCR analysis in Fig. 2 and

Supplemental Fig. 2 and 3.

Supplemental Dataset 1. RNA-seq data and plots of each transcript detected, including

mean normalized read count plots, mean normalized read count heat map, $\log_2$(fold

change) of mean normalized read counts relative to the pre-heat time point under the

same condition, and statistical analysis using False Discovery Rate (FDR).

Supplemental Dataset 2. $\log_2$(fold change, FC) of mean normalized read counts of

groups of transcripts based on MapMan function annotation (MapMan folder) or manual

curation (Manual_curation folder). Manual curation was based on the gene lists from this

paper (Zhang *et al.*, 2022*a*). Each line in the plots represent one unique transcript in that

function category under one of six conditions. RNA-seq data with normalized read counts

were used for this analysis. $\log_2$FC is relative to the pre-heat time point under the same

condition. Black vertical lines indicated the start of heat treatments under 35°C or 40°C

or the equivalent time point at 25°C. ANOVA test was applied to the time course (five time

points) of one transcript for each condition. If a transcript differed significantly during the

time course (or underwent a global significant change) of one condition, it is colored

purple (ANOVA significant); blue lines had no significant changes during the time course.

Different intensities of purple and blue colors were used to distinguish different transcripts.

Each file opens as a web page. Please use the interactive feature of the figures to see

the gene IDs for each panel.

Supplemental Dataset 3. MapMan ontology enrichment. Ontology enrichment was

performed for transcripts with significantly differential expression in the indicated

comparisons using the extended MapMan ontology. The p values were calculated using

a hypergeometric test and corrected for multiple testing using the Benjamini-Hochberg

method.

Supplemental Dataset 4. Normalized RNA-seq read counts.

Supplemental Dataset 5. RNA-seq data summary, including gene IDs, MapMan function

groups, annotation, adjusted False Discovery Rate (FDR) values for comparisons of

different conditions, mean and standard deviation (sd) of normalized read counts.

Supplemental figures -6 (see figure file)

Acknowledgements

We would like to thank Dr. Michael Schroda for discussing the initial experimental idea

under the acetate-depleting conditions and Dr. Trent Northen for recommending the

acetate quantification assay. We also want to thank the team of the Department of Energy

(DOE) Joint Genome Institute (JGI) Community Science Program (Anna Lipzen, Christa

Pennacchio, Kerrie Barry, Dr. Jeremy Schmutz, and others) for managing and

sequencing our RNA samples.

Author contributions

RZ designed the experiments. CB and MX operated and maintained the algal growth in

photobioreactors. NZ, CB, and MX harvested algal samples at different time points and

quantified biomass. NZ and MX performed acetate quantification assay. NZ performed

RT-qPCR analysis and extracted RNA for RNA-seq. CB and NZ performed light

microscope imaging. CB and MX quantified cell density, cell size, and chlorophyll
contents, and recorded OD_{608}/OD_{720} readings. CB and MX performed 4-day heating
experiments and quantified growth rates as well as cell parameters. CB analyzed PSII
data. BV, TM, RZ did RNA-seq analysis. EMM did initial analysis of the acetate pathways.
NZ, BV, CB, RZ, MX, and EMM prepared the figures. BV wrote the RNA-seq method and
RZ wrote the rest of the manuscript. RZ, EMM, BV, and CB revised the manuscript.
933

934 **Conflict of interest**
935 No conflict of interest declared.
936

Funding statement
This work was supported by the start-up funding from the Donald Danforth Plant Science
Center (DDPSC) and the DOE Biological & Environmental Research (BER) grant (DE-
SC0020400) to RZ. EMM was supported by the William H. Danforth Fellowship in Plant
Sciences and the DDPSC start-up funding (to RZ). BV and TM were supported by the
Germany funding TRR 175 (Project No: 270050988). The RNA-seq service support was
from the Department of Energy (DOE) Joint Genome Institute (JGI) Community Science
Program (CSP, 503414) to RZ. The NovaSeq for RNA-seq (proposal:
10.46936/10.25585/60001126) was conducted by the U.S. Department of Energy Joint
Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is
supported by the Office of Science of the U.S. Department of Energy operated under
Contract No. DE-AC02-05CH11231.

Data availability
All data supporting the findings of this study are available within the paper and within its
supplementary materials published online.
953

References

Altschuler M, Mascarenhas JP. 1982. Heat shock proteins and effects of heat shock in plants. Plant molecular biology **1**, 103–115.

Al-Whaibi MH. 2011. Plant heat-shock proteins: A mini review. Journal of King Saud University - Science **23**, 139–150.

Anderson CM, Mattoon EM, Zhang N, *et al.* 2021. High light and temperature reduce photosynthetic efficiency through different mechanisms in the C4 model Setaria viridis. Communications Biology **4**, 1092.

Aprile A, Havlickova L, Panna R, *et al.* 2013. Different stress responsive strategies to drought and heat in two durum wheat cultivars with contrasting water use efficiency. BMC genomics **14**, 821.

Bajhaiya AK, Dean AP, Zeef LAH, Webster RE, Pittman JK. 2016. PSR1 Is a Global Transcriptional Regulator of Phosphorus Deficiency Responses and Carbon Storage Metabolism in Chlamydomonas reinhardtii. Plant physiology **170**, 1216–1234.

Baker NR, Harbinson J, Kramer DM. 2007. Determining the limitations and regulation of photosynthetic energy transduction in leaves. Plant, cell & environment **30**, 1107–1125.

Balfagón D, Sengupta S, Gómez-Cadenas A, Fritschi FB, Azad RK, Mittler R, Zandalinas SI. 2019. Jasmonic acid Is required for plant acclimation to a combination of high light and heat stress. Plant physiology **181**, 1668–1682.

Baud S, Wuillème S, Dubreucq B, de Almeida A, Vuagnat C, Lepiniec L, Miquel M, Rochat C. 2007. Function of plastidial pyruvate kinases in seeds of Arabidopsis thaliana. The Plant journal: for cell and molecular biology **52**, 405–419.

BBDuk. 2022. BBMap short read aligner, and other bioinformatic tools.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society. Series B, Statistical methodology **57**, 289–300.

Bigeard J, Hirt H. 2018. Nuclear Signaling of Plant MAPKs. Frontiers in plant science **9**, 469.

Bourgine B, Guihur A. 2021. Heat Shock Signaling in Land Plants: From Plasma Membrane Sensing to the Transcription of Small Heat Shock Proteins. Frontiers in plant science **12**, 710801.

Boyle NR, Page MD, Liu B, *et al.* 2012. Three acyltransferases and nitrogen-responsive regulator are implicated in nitrogen starvation-induced triacylglycerol accumulation in Chlamydomonas. The Journal of biological chemistry **287**, 15811–15825.

de Carpentier F, Maes A, Marchand CH, Chung C, Durand C, Crozet P, Lemaire SD, Danon A. 2022. How abiotic stress-induced socialization leads to the formation of massive aggregates in Chlamydomonas. Plant physiology **190**, 1927–1940.

Chapman S, Paget C, Johnson G, Schwartz J-M. 2015. Flux balance analysis reveals acetate metabolism modulates cyclic electron flow and alternative glycolytic pathways in Chlamydomonas reinhardtii. Frontiers in plant science **6**.

Chew SY, Chee WJY, Than LTL. 2019. The glyoxylate cycle and alternative carbon metabolism as metabolic adaptation strategies of Candida glabrata: perspectives from Candida albicans and Saccharomyces cerevisiae. Journal of Biomedical Science **26**.

Chioccioli M, Hankamer B, Ross IL. 2014. Flow Cytometry Pulse Width Data Enables Rapid and Sensitive Estimation of Biomass Dry Weight in the Microalgae Chlamydomonas reinhardtii and Chlorella vulgaris. PLoS ONE **9**, e97269.

Crozet P, Navarro FJ, Willmund F, *et al.* 2018. Birth of a photosynthetic chassis: A MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas reinhardtii*. ACS synthetic biology **7**, 2074–2086.

Delorge I, Janiak M, Carpentier S, Van Dijck P. 2014. Fine tuning of trehalose biosynthesis and hydrolysis as novel tools for the generation of abiotic stress tolerant plants. Frontiers in plant science **5**.

Dhokane D, Bhadra B, Dasgupta S. 2020. CRISPR based targeted genome editing of *Chlamydomonas reinhardtii* using programmed Cas9-gRNA ribonucleoprotein. Molecular biology reports **47**, 8747–8755.

van Dongen JT, Gupta KJ, Ramírez-Aguilar SJ, Araújo WL, Nunes-Nesi A, Fernie AR. 2011. Regulation of respiration in plants: a role for alternative metabolic pathways. Journal of plant physiology **168**, 1434–1443.

El-Sheekh MM, Gheda SF, El-Sayed AE-KB, Abo Shady AM, El-Sheikh ME, Schagerl M. 2019. Outdoor cultivation of the green microalga Chlorella vulgaris under stress conditions as a feedstock for biofuel. Environmental science and pollution research international **26**, 18520–18532.

Emrich-Mills TZ, Yates G, Barrett J, *et al.* 2021. A recombineering pipeline to clone large and complex genes in Chlamydomonas. The Plant Cell **33**, 1161–1181.

Fei X, Yu J, Li Y, Deng X. 2017. CrMAPK3 regulates the expression of iron-deficiency-responsive genes in Chlamydomonas reinhardtii. BMC biochemistry **18**, 6.

Finka A, Cuendet AFH, Maathuis FJM, Saidi Y, Goloubinoff P. 2012. Plasma membrane cyclic nucleotide gated calcium channels control land plant thermal sensing and acquired thermotolerance. The Plant cell **24**, 3333–3348.

Gao F, Han X, Wu J, Zheng S, Shang Z, Sun D, Zhou R, Li B. 2012. A heat-activated calcium-permeable channel - Arabidopsis cyclic nucleotide-gated ion channel 6 - is involved in heat shock responses. The Plant Journal **70**, 1056–1069.

González-Ballester D, Casero D, Cokus S, Pellegrini M, Merchant SS, Grossman AR. 2010. RNA-seq analysis of sulfur-deprived Chlamydomonas cells reveals aspects of acclimation critical for cell survival. The Plant cell **22**, 2058–2084.

Greiner A, Kelterborn S, Evers H, Kreimer G, Sizova I, Hegemann P. 2017. Targeting of photoreceptor genes in *Chlamydomonas reinhardtii* via zinc-finger nucleases and CRISPR/Cas9. The Plant cell **29**, 2498–2518.

Guihur A, Rebeaud ME, Goloubinoff P. 2022. How do plants feel the heat and survive? Trends in biochemical sciences **47**, 824–838.

Haq S, Khan A, Ali M, Khattak AM, Gai W-X, Zhang H-X, Wei A-M, Gong Z-H. 2019. Heat Shock Proteins: Dynamic Biomolecules to Counter Plant Biotic and Abiotic Stresses. International journal of molecular sciences **20**, 5321.

Harris EH. 2009. *The Chlamydomonas Sourcebook: Introduction to Chlamydomonas and Its Laboratory Use: Volume 1*. Academic Press.

Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. 2007. QBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. Genome biology **8**, R19.

Hemme D, Veyel D, Mühlhaus T, *et al.* 2014. Systems-wide analysis of acclimation responses to long-term heat stress and recovery in the photosynthetic model organism *Chlamydomonas reinhardtii*. The Plant cell **26**, 4270–4297.

Hennacy JH, Jonikas MC. 2020. Prospects for engineering biophysical CO_2 concentrating mechanisms into land Plants to enhance yields. Annual review of plant biology **71**, 461–485.

Ivanov IN, Zachleder V, Vítová M, Barbosa MJ, Bišová K. 2021. Starch production in *Chlamydomonas reinhardtii* through supraoptimal temperature in a pilot-scale photobioreactor. Cells **10**, 1084.

Jagodzik P, Tajdel-Zielinska M, Ciesla A, Marczak M, Ludwikow A. 2018. Mitogen-Activated Protein Kinase Cascades in Plant Hormone Signaling. Frontiers in plant science **9**, 1387.

Janni M, Gullì M, Maestri E, Marmiroli M, Valliyodan B, Nguyen HT, Marmiroli N. 2020. Molecular and genetic bases of heat stress responses in crop plants and breeding for increased resilience and productivity. Journal of experimental botany **71**, 3780–3802.

Jee H. 2016. Size dependent classification of heat shock proteins: a mini-review. Journal of exercise rehabilitation **12**, 255–259.

Ji S, Siegel A, Shan S-O, Grimm B, Wang P. 2021. Chloroplast SRP43 autonomously protects chlorophyll biosynthesis proteins against heat shock. Nature plants **7**, 1420–1432.

Johnson GN. 2011. Physiology of PSI cyclic electron transport in higher plants. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1807**, 384–389.

Johnson X, Alric J. 2012. Interaction between starch breakdown, acetate assimilation, and photosynthetic cyclic electron flow in *Chlamydomonas reinhardtii*. The Journal of biological chemistry **287**, 26445–26452.

Johnson X, Alric J. 2013. Central carbon metabolism and electron transport in *Chlamydomonas reinhardtii*: metabolic constraints for carbon partitioning between oil and starch. Eukaryotic cell **12**, 776–793.

Jung J-H, Barbosa AD, Hutin S, *et al.* 2020. A prion-like domain in ELF3 functions as a thermosensor in Arabidopsis. Nature **585**, 256–260.

Jung J-H, Domijan M, Klose C, *et al.* 2016. Phytochromes function as thermosensors in Arabidopsis. Science **354**, 886–889.

Karpowicz SJ, Prochnik SE, Grossman AR, Merchant SS. 2011. The GreenCut2 resource, a phylogenomically derived inventory of proteins specific to the plant lineage. The Journal of biological chemistry **286**, 21427–21439.

Kim S-C, Guo L, Wang X. 2020. Nuclear moonlighting of cytosolic glyceraldehyde-3-phosphate dehydrogenase regulates Arabidopsis response to heat stress. Nature communications **11**, 3439.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory requirements. Nature methods **12**, 357–360.

Koley S, Chu KL, Mukherjee T, Morley SA, Klebanovych A, Czymmek KJ, Allen DK. 2022. Metabolic synergy in Camelina reproductive tissues for seed development. Science advances **8**, eabo7683.

Kropat J, Hong-Hermesdorf A, Casero D, Ent P, Castruita M, Pellegrini M, Merchant SS, Malasarn D. 2011. A revised mineral nutrient supplement increases biomass and growth rate in Chlamydomonas reinhardtii. The Plant journal: for cell and molecular biology **66**, 770–780.

Légeret B, Schulz‑Raffelt M, Nguyen HM, Auroy P, Beisson F, Peltier G, Blanc G, Li‑Beisson Y. 2016. Lipidomic and transcriptomic analyses of *Chlamydomonas reinhardtii* under heat stress unveil a direct route for the conversion of membrane lipids into storage lipids. Plant, cell & environment **39**, 834–847.

Legris M, Klose C, Burgie ES, Rojas CCR, Neme M, Hiltbrunner A, Wigge PA, Schäfer E, Vierstra RD, Casal JJ. 2016. Phytochrome B integrates light and temperature signals in Arabidopsis. Science **354**, 897–900.

Li X, Fauser F, Jinkerson RE, *et al.* 2019. A genome-wide algal mutant library and functional screen identifies genes required for eukaryotic photosynthesis. Nature genetics **51**, 627–635.

Li Z, Tang J, Srivastava R, Bassham DC, Howell SH. 2020. The transcription factor bZIP60 links the unfolded protein response to the heat stress response in maize. The Plant cell **32**, 3559–3575.

Li X, Zhang R, Patena W, *et al.* 2016. An indexed, mapped mutant library enables reverse genetics studies of biological processes in *Chlamydomonas reinhardtii*. The Plant cell **28**, 367–387.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics **30**, 923–930.

Lien T, Knutsen G. 1979. Synchronous growth of *Chlamydomonas reinhardtii*(chlorophyceae): a review of optimal conditions. Journal of phycology **15**, 191–200.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods **25**, 402–408.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology **15**, 550.

Lucker B, Kramer DM. 2013. Regulation of cyclic electron flow in *Chlamydomonas reinhardtii* under fluctuating carbon availability. Photosynthesis research **117**, 449–459.

Martinière A, Shvedunova M, Thomson AJW, Evans NH, Penfield S, Runions J, McWatters HG. 2011. Homeostasis of plasma membrane viscosity in fluctuating temperatures. The New phytologist **192**, 328–337.

Mata TM, Martins AA, Caetano NS. 2010. Microalgae for biodiesel production and other applications: A review. Renewable and Sustainable Energy Reviews **14**, 217–232.

Mathimani T, Pugazhendhi A. 2019. Utilization of algae for biofuel, bio-products and bio-remediation. Biocatalysis and agricultural biotechnology **17**, 326–330.

Mattoon EM, McHargue W, Bailey CE, *et al.* 2022. High-throughput Identification of Novel Heat Tolerance Genes via Genome-wide Pooled Mutant Screens in the Model Green Alga Chlamydomonas reinhardtii. bioRxiv, 2022.07.13.499508.

Merchant SS, Prochnik SE, Vallon O, *et al.* 2007. The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. Science **318**, 245–250.

Meyer MT, Whittaker C, Griffiths H. 2017. The algal pyrenoid: key unanswered questions. Journal of experimental botany **68**, 3739–3749.

Minagawa J, Tokutsu R. 2015. Dynamic regulation of photosynthesis in *Chlamydomonas reinhardtii*. The Plant journal: for cell and molecular biology **82**, 413–428.

Mittler R, Finka A, Goloubinoff P. 2012. How do plants feel the heat? Trends in biochemical sciences **37**, 118–125.

Mühlhaus T, Weiss J, Hemme D, Sommer F, Schroda M. 2011*a*. Quantitative Shotgun Proteomics Using a Uniform 15N-Labeled Standard to Monitor Proteome Dynamics in Time Course Experiments Reveals New Insights into the Heat Stress Response of Chlamydomonas reinhardtii. Molecular & Cellular Proteomics **10**, M110.004739.

Mühlhaus T, Weiss J, Hemme D, Sommer F, Schroda M. 2011*b*. Quantitative shotgun proteomics using a uniform ^{15}N-labeled standard to monitor proteome dynamics in time course experiments reveals new insights into the heat stress response of *Chlamydomonas reinhardtii*. Molecular & cellular proteomics: MCP **10**, M110.004739.

Munekage Y, Hashimoto M, Miyake C, Tomizawa K-I, Endo T, Tasaka M, Shikanai T. 2004. Cyclic electron flow around photosystem I is essential for photosynthesis. Nature **429**, 579–582.

Murata N, Nishiyama Y. 2018. ATP is a driving force in the repair of photosystem II during photoinhibition. Plant, cell & environment **41**, 285–299.

Niemeyer J, Scheuring D, Oestreicher J, Morgan B, Schroda M. 2021. Real-time monitoring of subcellular H2O2 distribution in *Chlamydomonas reinhardtii*. The Plant cell.

Olas JJ, Apelt F, Annunziata MG, John S, Richard SI, Gupta S, Kragler F, Balazadeh S, Mueller-Roeber B. 2021. Primary carbohydrate metabolism genes participate in heat-stress memory at the shoot apical meristem of Arabidopsis thaliana. Molecular plant **14**, 1508–1524.

Ortiz-Bobea A, Wang H, Carrillo CM, Ault TR. 2019. Unpacking the climatic drivers of US agricultural yields. Environmental research letters: ERL [Web site] **14**, 064003.

Porra RJ, Thompson WA, Kriedemann PE. 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. Biochimica et Biophysica Acta (BBA) - Bioenergetics **975**, 384–394.

Pospíšil P. 2016. Production of reactive oxygen species by photosystem II as a response to light and temperature stress. Frontiers in plant science **7**, 1950.

Pröschold T, Harris EH, Coleman AW. 2005. Portrait of a species: *Chlamydomonas reinhardtii*. Genetics **170**, 1601–1610.

Remans T, Keunen E, Bex GJ, Smeets K, Vangronsveld J, Cuypers A. 2014. Reliable gene expression analysis by reverse transcription-quantitative PCR: reporting and minimizing the uncertainty in data accuracy. The Plant cell **26**, 3829–3837.

Roach T, Sedoud A, Krieger-Liszkay A. 2013. Acetate in mixotrophic growth medium affects photosystem II in Chlamydomonas reinhardtii and protects against photoinhibition. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1827**, 1183–1190.

Rollins JA, Habte E, Templer SE, Colby T, Schmidt J, von Korff M. 2013. Leaf proteome alterations in the context of physiological and morphological responses to drought and heat stress in barley (Hordeum vulgare L.). Journal of experimental botany **64**, 3201–3212.

Rütgers M, Muranaka LS, Schulz-Raffelt M, Thoms S, Schurig J, Willmund F, Schroda M. 2017. Not changes in membrane fluidity but proteotoxic stress triggers heat shock protein expression in *Chlamydomonas reinhardtii*. Plant, cell & environment **40**, 2987–3001.

Sager R. 1955-7. Inheritance in the green alga *Chlamydomonas reinhardtii*. Genetics **40**, 476–489.

Sasso S, Stibor H, Mittag M, Grossman AR. 2018. From molecular manipulation of domesticated Chlamydomonas reinhardtii to survival in nature. eLife **7**, e39233.

Schloss JA. 1990. A *Chlamydomonas* gene encodes a G protein β subunit-like polypeptide. Molecular & general genetics: MGG **221**, 443–452.

Schmollinger S, Mühlhaus T, Boyle NR, *et al.* 2014. Nitrogen-sparing mechanisms in *Chlamydomonas* affect the transcriptome, the proteome, and photosynthetic metabolism. The Plant cell **26**, 1410–1435.

Schneider K, Venn B, Mühlhaus T. 2022. Plotly.net: A fully featured charting library for .NET programming languages. F1000Research **11**, 1094.

Schroda M, Hemme D, Mühlhaus T. 2015. The *Chlamydomonas* heat stress response. The Plant journal: for cell and molecular biology **82**, 466–480.

Schulz-Raffelt M, Lodha M, Schroda M. 2007. Heat shock factor 1 is a key regulator of the stress response in *Chlamydomonas*. The Plant journal: for cell and molecular biology **52**, 286–295.

Scranton MA, Ostrand JT, Fields FJ, Mayfield SP. 2015. *Chlamydomonas* as a model for biofuels and bio-products production. The Plant journal: for cell and molecular biology **82**, 523–531.

Shimogawara K, Fujiwara S, Grossman A, Usuda H. 1998. High-efficiency transformation of *Chlamydomonas reinhardtii* by electroporation. Genetics **148**, 1821–1828.

Sommer F, Kropat J, Malasarn D, Grossoehme NE, Chen X, Giedroc DP, Merchant SS. 2010. The CRR1 nutritional copper sensor in Chlamydomonas contains two distinct metal-responsive domains. The Plant cell **22**, 4098–4113.

Song Z-T, Zhang L-L, Han J-J, Zhou M, Liu J-X. 2021. Histone H3K4 methyltransferases SDG25 and ATX1 maintain heat-stress gene expression during recovery in Arabidopsis. The Plant journal: for cell and molecular biology **105**, 1326–1338.

Strenkert D, Schmollinger S, Gallaher SD, *et al.* 2019. Multiomics resolution of molecular events during a day in the life of *Chlamydomonas*. Proceedings of the National Academy of Sciences of the United States of America **116**, 2374–2383.

Takano W, Hisabori T, Wakabayashi K-I. 2021. Rapid estimation of cytosolic ATP concentration from the ciliary beating frequency in the green alga Chlamydomonas reinhardtii. The Journal of biological chemistry **296**, 100156.

Usadel B, Poree F, Nagel A, Lohse M, Czedik-Eysenberg A, Stitt M. 2009. A guide to using MapMan to visualize and compare Omics data in plants: a case study in the crop species, Maize. Plant, cell & environment **32**, 1211–1229.

Venn B, Mühlhaus T. 2022. CSBiology/OntologyEnrichment: Release 0.0.1 (0.0.1). Zenodo. https://doi.org/10.5281/zenodo.6340412.

Venn B, Weil L, Schneider K, Zimmer D, Mühlhaus T. 2022. fslaborg/FSharp.Stats. Zenodo. https://doi.org/10.5281/zenodo.6337056.

Vítová M, Bišová K, Hlavová M, Kawano S, Zachleder V, Cížková M. 2011. *Chlamydomonas reinhardtii*: duration of its cell cycle and phases at growth rates affected by temperature. Planta **234**, 599–608.

Voß B, Meinecke L, Kurz T, Al-Babili S, Beck CF, Hess WR. 2010. Hemin and Magnesium-Protoporphyrin IX Induce Global Changes in Gene Expression in Chlamydomonas reinhardtii. Plant physiology **155**, 892–905.

Vu LD, Gevaert K, De Smet I. 2019. Feeling the Heat: Searching for Plant Thermosensors. Trends in plant science **24**, 210–219.

Walker RP, Chen Z-H, Famiani F. 2021. Gluconeogenesis in Plants: A Key Interface between Organic Acid/Amino Acid/Lipid and Sugar Metabolism. Molecules **26**.

Wang L, Yang L, Wen X, Chen Z, Liang Q, Li J, Wang W. 2019. Rapid and high efficiency transformation of *Chlamydomonas reinhardtii* by square-wave electroporation. Bioscience reports **39**, BSR20181210.

Wulfert S, Schilasky S, Krueger S. 2020. Transcriptional and Biochemical Characterization of Cytosolic Pyruvate Kinases in Arabidopsis thaliana. Plants **9**.

Xiao L, Young EB, Grothjan JJ, Lyon S, Zhang H, He Z. 2015. Wastewater treatment and microbial communities in an integrated photo-bioelectrochemical system affected by different wastewater algal inocula. Algal Research **12**, 446–454.

Xie B, Bishop S, Stessman D, Wright D, Spalding MH, Halverson LJ. 2013. *Chlamydomonas reinhardtii* thermal tolerance enhancement mediated by a mutualistic interaction with vitamin B12-producing bacteria. The ISME journal **7**, 1544–1555.

Xu Z-S, Li Z-Y, Chen Y, Chen M, Li L-C, Ma Y-Z. 2012. Heat shock protein 90 in plants: molecular mechanisms and roles in stress responses. International journal of molecular sciences **13**, 15706–15723.

Yamori W, Shikanai T. 2016. Physiological functions of cyclic electron transport around Photosystem I in sustaining photosynthesis and plant growth. Annual review of plant biology **67**, 81–106.

Young EB, Reed L, Berges JA. 2022. Growth parameters and responses of green algae across a gradient of phototrophic, mixotrophic and heterotrophic conditions. PeerJ **10**, e13776.

Zachleder V, Ivanov I, Vítová M, Bišová K. 2019. Cell cycle arrest by supraoptimal temperature in the alga *Chlamydomonas reinhardtii*. Cells **8**, 1237.

Zhang N, Mattoon EM, McHargue W, *et al.* 2022*a*. Systems-wide analysis revealed shared and unique responses to moderate and acute high temperatures in the green alga Chlamydomonas reinhardtii. Communications biology **5**, 460.

Zhang N, Pazouki L, Nguyen H, *et al.* 2022*b*. Comparative phenotyping of two commonly used Chlamydomonas reinhardtii background strains: CC-1690 (21gr) and CC-5325 (The CLiP mutant library background). Plants **11**, 585.

Zhang X, Rerksiri W, Liu A, Zhou X, Xiong H, Xiang J, Chen X, Xiong X. 2013. Transcriptome profile reveals heat response mechanism at molecular and metabolic levels in rice flag leaf. Gene **530**, 185–192.

Zhang M, Zhang S. 2022. Mitogen-activated protein kinase cascades in plant signaling. Journal of integrative plant biology **64**, 301–341.

Zones JM, Blaby IK, Merchant SS, Umen JG. 2015. High-resolution profiling of a synchronized diurnal transcriptome from *Chlamydomonas reinhardtii* reveals continuous cell and metabolic differentiation. The Plant cell **27**, 2743–2769.

Figures

Fig. 1.

Fig. 1. **Under the acetate-depleting condition, moderate high temperature of 35°C transiently increased but then decreased Chlamydomonas biomass accumulation.**
(A) Before heat treatments, Chlamydomonas cells (CC-1690, also called *21gr*, wild-type) were grown in photobioreactors (PBRs) in Tris-acetate-phosphate (TAP) medium (acetate as an organic carbon source) with a light intensity of 100 μmol photons m^{-2} s^{-1} and constantly bubbling of air. Algal cultures were maintained turbidostatically within a small range of OD$_{680}$, which monitors chlorophyll content (μg chlorophyll mL^{-1} cultures). When PBR cultures grew to the set maximum OD$_{680}$, pumps were turned on to add fresh medium and dilute the cultures to the set minimal OD$_{680}$, then pumps were turned off to allow for exponential growth to the maximum set OD$_{680}$. Figure cited and modified from a supplemental figure of this paper (Zhang *et al.*, 2022*a*). **(B)** Experimental outline for the acetate-depleting condition. The PBR cultures were first maintained at 25°C with constant

nutrient supply using the turbidostatic mode as mentioned above. After the cultures reached steady growth (pre-heat time point), the turbidostatic pumps were turned off and the cultures were grown under the indicated temperature for 24 hours (h) without fresh medium supply (thus no constant acetate supply, defined as the acetate-depleting condition). Algal cultures were harvested at different time points for the indicated parameters. Each temperature treatment was conducted in an individual PBR with at least 3 replicates. **(C)** Under the acetate-depleting condition, algal dry biomass increased with 8-h heat but decreased with 24-h heat of 35°C as compared to the constant 25°C. Mean $\pm$ SE, n=3-8 biological replicates. **(D)** Heat at 35°C accelerated acetate consumption. Algal cultures were harvested at different time points during the acetate-depleting condition and acetate content in algal supernatant was quantified using the Acetate Colorimetric Assay Kit. The red dashed line marks the start of heating at 35°C or 40°C or equivalent time point at the control 25°C. **(C, D)** Statistical analyses were performed using two-tailed t-test assuming unequal variance; *, p<0.05, the colors of the asterisks match the heated condition as compared to 25°C at the same time points; #, p<0.05, for the comparisons between 35°C and 40°C at the same time points. ns, not significant.

Fig. 2.

Fig. 2. Moderate high temperature of 35ºC up-regulated transcripts related to acetate uptake/assimilation, glyoxylate cycle, gluconeogenesis, and glycolysis pathways. (A) Simplified pathways of acetate uptake/assimilation, glyoxylate cycle,

gluconeogenesis, and glycolysis based on these papers (Johnson and Alric 2013; Durante et al. 2019). Key enzymes are in boxes and were analyzed via RNA-seq in panels B-P. Acetate uptake/assimilation enzymes: GFY3, putative acetate transporter; ACS2/3, acetyl-CoA synthases; ACK2, acetate kinase. Glyoxylate cycle key enzymes: ICL1, isocitrate lyase; MAS1, malate synthase; CIS2, Citrate synthase; MDH1/2/4, NAD-dependent malate dehydrogenases. Gluconeogenesis key enzymes: PCK1, phosphoenolpyruvate carboxykinase; FBP1, fructose-1,6-bisphosphatase. Glycolysis key enzymes: PYK2/3/4, pyruvate kinase 2/3/4. See all gene IDs and annotations in Supplementary Table 1. PEP, phosphoenolpyruvate. Fru-6-P, fructose-6-phosphate. Fru-1,6-P$_2$, fructose-1,6-bisphosphate. **(B-P)** Transcriptional change of the indicated genes under six different conditions, presented as log$_2$(fold change) of mean normalized read counts relative to the pre-heat time point. In each panel, each line represents the transcriptional time course of the indicated gene under one condition as indicated by the legend. Solid lines for the acetate-depleting condition and dashed lines for the constant-acetate condition. Black, brown, red lines for 25^{o}C, 35^{o}C, and 40^{o}C temperature treatments, respectively. See Supplemental Dataset 1 for bigger pictures and more information.

Fig. 3.

Fig. 3. Algal cell physiologies with different temperature treatments under the acetate-depleting condition were compared with those under the constant-acetate condition. (A-E) Cell parameters from algal cultures under the acetate-depleting condition. The algal cultivation and heat treatments were the same as in Fig. 1B. **(F-J)** Cell parameters from algal cultures under the constant-acetate condition, with constant acetate supply via turbidostatic mode, data plotted based on the results from this paper

(Zhang *et al.*, 2022*a*). (**A-J**) Two panels on the same row share the same y axis. The red dashed line marks the start of heat. **(K-O)** Fold change of the indicated cell parameters from algal cultures under the acetate-depleting condition (filled bars) and under the constant-acetate condition (empty bars). (**A-O**) Mean ± SE, *n* = 3-7 biological replicates. Statistical analyses were performed with two-tailed t-test assuming unequal variance; *, $p<0.05$, for the comparisons between 35°C or 40°C with 25°C at the same time point under the same acetate condition, the colors of the asterisks match the heat condition; #, $p<0.05$, for the comparisons between 35°C and 40°C at the same time point under the same acetate condition; ^, $p<0.05$, for the comparisons between the acetate-depleting and constant-acetate conditions at the same time point and under the same temperature. **(D, E, I, J)** Cell parameters at different time points during constant 25°C had little change, not significantly different from the pre-heat time points ($p>0.05$).

Fig. 4.

Fig. 4. Both OD$_{680}$ and OD$_{720}$ were linearly proportional to chlorophyll contents under different temperatures and can be used to estimate algal growth rates. Black, brown, red lines/symbols represent 25°C, 35°C, and 40°C treatments, respectively. **(A, B)** OD$_{680}$ and OD$_{720}$ automatically monitored algal growth every one min in algal cultures grown in photobioreactors with different temperatures under the acetate-depleting conditions (medium starting with acetate but no turbidostatic control when heat started). The algal cultivation and heat treatments were the same as in Fig. 1B. Multiple lines with the same colors represent biological replicates (n= 4-8) under the same condition. **(C-J)** OD$_{680}$ and OD$_{720}$ were linearly proportional to Chlorophyll (Chl) contents per mL cultures but not cell volume or cell density. Chl contents (**C, D**), cell volume (**E, F**), and cell densities (**G-J**) were plotted against OD$_{680}$ or OD$_{720}$ readings in algal cultures with different treatments mentioned above. Dashed blue lines are linear trendlines of best fit and R-squared values are displayed. **(I, J)** The same data as in panel G-H but with smaller ranges for X values.

Fig. 5.

Fig. 5. Under the constant-acetate condition with turbidostatic control, moderate (35°C) and acute high temperatures (40°C) increased and reduced algal growth during 4-day heating, respectively. Black, brown, red lines/symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat at 35°C or 40°C. **(A)** Chlamydomonas cultures were grown in photobioreactors (PBRs) under turbidostatic conditions at different temperatures in acetate-containing medium. Algal cultures were first acclimated at 25°C for 4 days before the temperature was switched to 35°C or 40°C or stayed at 25°C for 4 days. Relative growth rates were calculated based on the cycling of OD_{680} caused by the turbidostatic control (see Fig. 1A and method for details). Each temperature treatment had 3 biological replicates in separate PBRs. **(B-F)** Cell parameters were quantified from algal cultures harvested at different time points with different treatments. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing 35°C or 40°C with 25°C at the same time point. *, $p < 0.05$, the colors of the asterisks match the heat condition; #, $p < 0.05$, for the comparisons between 35°C and 40°C at the same time point. Chl, Chlorophyll.

Fig. 6.

11-53

Fig. 6. Under the no-acetate condition (photoautotrophic medium) with turbidostatic control, both moderate (35°C) and acute high temperatures (40°C) reduced algal growth during 4-day heating. Black, brown, red lines/symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat at 35°C or 40°C. **(A)** Chlamydomonas cultures were grown in photobioreactors (PBRs) under turbidostatic conditions at different temperatures in photoautotrophic medium without acetate. Algal cultures were first acclimated at 25°C for 4 days before the temperature was switched to 35°C or 40°C or stayed at 25°C for 4 days. Relative growth rates were calculated based on the cycling of OD_{680} caused by the turbidostatic control (see Fig. 1A and method for details). Each temperature treatment had at least 3 biological replicates in separate PBRs. **(B-F)** Cell parameters were quantified from algal cultures harvested at different time points with different treatments. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing 35°C or 40°C with 25°C at the same time point. *, $p < 0.05$, the colors of the asterisks match the heat condition; #, $p < 0.05$, for the comparisons between 35°C and 40°C at the same time point. Chl, Chlorophyll.

Fig. 7

Fig. 7. Under the no-acetate condition (photoautotrophic medium) with turbidostatic control, 35°C-treated cells formed palmelloids and 40°C-treated cells were bleached after 4-day heat treatments. Light microscopic images of Chlamydomonas cells under different conditions. The algal cultivation and heat treatments were the same as in Fig. 6. Images shown are representative results from three biological replicates. Black scale bar on each panel represents 10 μm.

Fig. 8

14-53

Fig. 8. Under the turbidostatic control without nutrient limitation, moderate high temperatures (35°C) increased algal growth rates with constant acetate but reduced algal growth rates without acetate. Plotted data were from Fig. 5 and 6 but were normalized to the corresponding controls at the 25°C with the same medium type to compare the relative effects of high temperatures with (filled symbols and solid lines) and without (open symbols and dashed lines) acetate. Black, brown, red lines/symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat at 35°C or 40°C. **(A)** Chlamydomonas cultures were grown in photobioreactors (PBRs) under turbidostatic conditions at different temperatures with and without acetate. Algal cultures were first acclimated at 25°C for 4 days before the temperature was switched to 35°C or 40°C or stayed at 25°C for 4 days. Relative growth rates were calculated based on the cycling of OD_{680} caused by the turbidostatic control (see Fig. 1A and method for details), normalized to the corresponding controls at the 25°C with the same medium type. Each temperature treatment had at least 3 biological replicates in separate PBRs. **(B-F)** Cell parameters were quantified from algal cultures harvested at different time points with different treatments, normalized to the corresponding controls at the 25°C with the same medium type. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing the data with and without acetate under the same high temperature treatment. *, $p < 0.05$, the colors of the asterisks match the heat condition, brown asterisks for 35°C and red asterisks for 40°C comparisons. Chl, Chlorophyl.

15-53

Fig. 9

Fig. 9. Under the turbidostatic control and no nutrient limitation, algal cultures had higher photosystem II (PSII) operating efficiency with acetate than without acetate during treatment of 25°C, 35°C, or 40°C, although the acetate effects diminished after 2-day heat of 40°C due to cell death. PSII operating efficiency was measured automatically in photobioreactors under different conditions. Black, brown, red lines/symbols represent 25°C, 35°C, and 40°C treatments, respectively. The red dashed lines mark the start of heat at 35°C or 40°C. **(A)** Acetate-depleting condition at 25°C, 35°C, or 40°C for 24 h, in medium started with acetate but no turbidostatic control, thus no constant acetate supply. **(B)** Constant-acetate condition at 25°C, 35°C, or 40°C for 4 days, in medium with acetate and turbidostatic control, thus no nutrient limitation. **(C)** No-acetate condition at 25°C, 35°C, or 40°C for 4 days, in photoautotrophic medium without acetate but with turbidostatic control, thus no nutrient limitation. **(D)** The comparison of PSII operating efficiency in the constant-acetate (data from panel B, darker color bars) and no-acetate (data from panel C, light color bars) conditions during the indicated periods at 25°C, 35°C, or 40°C for 4 days. Statistical analyses were performed with two-tailed t-test assuming unequal variance by comparing the indicated groups. *, $p < 0.05$, comparing 35°C or 40°C with 25°C with the same medium type at the same time; #, $p < 0.05$, for the comparisons between 35°C and 40°C with the same medium type at the same time; ^, $p < 0.05$, for the comparisons between the same temperature treatment at the same time but with and without acetate. The colors of the significant marks match the heat condition, brown for 35°C and bright red color for 40°C related comparisons, and dark red for 35°C/40°C comparisons.

Fig. 10.

Fig. 10. Carbon (acetate) availability affected Chlamydomonas heat tolerance under the moderate high temperature (35°C), but acetate had much less effects under the acute high temperature (40°C). In medium containing the organic carbon source, acetate, heat of 35°C accelerated carbon metabolisms by upregulating transcripts related to acetate uptake/assimilation, glyoxylate cycle, gluconeogenesis, and glycolysis pathways. With constant acetate supply, 35°C increased algal growth. Under the acetate-depleting condition, 35°C increased algal growth transiently followed by decreased growth and biomass accumulation due to acetate starvation. Without acetate supply in photoautotrophic medium, 35°C reduced algal growth rates. Under either constant-acetate or depleting-acetate condition, heat of 40°C down-regulated transcripts related to acetate uptake, glyoxylate cycle, gluconeogenesis, and glycolysis pathways, but up-regulated transcripts related to heat shock proteins (HSPs) and reactive oxygen species

(ROS) scavenging. In medium with or without acetate, 40°C decreased algal growth rates, inhibited cell division, damaged PSII efficiency, reduced chlorophyll contents, and eventually led to cell death. Thus, Chlamydomonas cells could not survive continuous heat of 40°C for longer than 2-days, independent of acetate supply. Brown and red labels/lines represent the effects of 35°C and 40°C, respectively. Purple, blue, and green labels/lines represent the acetate-depleting, constant-acetate, and no-acetate conditions, respectively.

See supplemental figures below.

Supplemental Fig. 1.

Supplemental Fig. 1. RNA-seq data from algal samples harvested from photobioreactors (PBRs) under the constant-acetate-25°C with turbidostatic control were highly correlated with each other. The Pearson correlations were generated by comparing all RNA-seq samples to the first sample (marked by *) of the constant-acetate-25°C condition. Under constant acetate, 25°C, and light (constant-acetate-25°C) with turbidostatic control in PBRs, the algal growth rates and cell physiologies stay constant (Fig. 5). We had RNA-seq data for 8 time points under the constant-acetate-25°C condition, and these samples were highly correlated with each other (with Pearson correlation ≥ 0.993). The 0 h of the constant-acetate-25°C condition corresponds to the time when PBRs reach expected high temperatures during heat treatments, which takes about 30 min from 25°C to 35°C or 40°C. All pre-heat samples were harvested from PBRs under the same constant-acetate-25°C condition with turbidostatic control and these RNA-seq samples had Pearson correlation > 0.95.

Because the cell status and transcriptome stay steady under the constant-acetate-25°C, we used the five extra time points under the constant-acetate-25°C condition (with underlines) to impute the 2 h and 4 h time points under the constant-acetate-25°C condition. See method for details.

Supplemental Fig. 2. RT-qPCR analysis was consistent with RNA-seq results. Transcript fold-changes of four selected transcripts involved in the glyoxylate cycle (ICL1 and MAS1) and gluconeogenesis cycle (PCK1 and FBP1) as in Fig. 2A were calculated based on RT-qPCR (**A-D**) or RNA-seq (**E-H**) results. Red dashed lines represent the start of heat treatments. For RT-qPCR results, the relative fold changes were calculated by normalizing the transcript expression to the reference genes *CBLP, EIF1A* and the pre-heat level. For RNA-seq results, the relative fold changes were calculated based on normalized RNA-seq read counts relative to the pre-heat level. Values are mean ± SE, n = 3 biological replicates. Glyoxylate cycle key enzymes: ICL1, isocitrate lyase; MAS1, malate synthase. Gluconeogenesis key enzymes: PCK1, phosphoenolpyruvate carboxykinase; FBP1, fructose-1,6-bisphosphatase (See all gene IDs and RT-qPCR primers in Supplemental Table 1).

Supplemental Fig. 3

Supplemental Fig. 3 continued

Supplemental Fig. 3 continued

Supplemental Fig. 3. Log₂(fold change) of the indicated transcript relative to pre-heat under different conditions. Log₂(fold change) were determined on the mean normalized transcript abundance, relative to the pre-heat time point. Each line represents one condition as indicated by the legend. Solid lines for the acetate-depleting condition and dashed lines for the constant-acetate condition. Black, brown, red lines for 25°C, 35°C, and 40°C temperature treatments, respectively. See all gene IDs in Supplemental Table 1 and Supplemental Dataset 1 for bigger pictures and more information.

Supplemental Fig. 4.

A. Transcripts related to HSP60s (HSP, heat shock protein)

B. Transcripts related to HSP70s

Supplemental Fig. 4. Log$_2$(fold change, FC) of mean normalized read counts of groups of transcripts based on MapMan function annotation or manual curation. Plots are Log$_2$FC of transcripts related to the indicated function groups under six different treatments, as compared to the pre-heat time points. RNA-seq data with normalized read counts were used for this analysis. Black vertical lines indicate the start of heat treatments under 35^{o}C or 40^{o}C or the equivalent time point at 25^{o}C. Each line represents one unique transcript in the indicated function group. ANOVA test was applied to the time course (five time points) of one transcript for each condition. If a transcript differed significantly during the time course (or underwent a global significant change) of one condition, it is colored purple (ANOVA significant); blue lines had no significant changes during the time course. Different intensities of purple and blue colors were used to distinguish different transcripts. The total number of transcripts plotted in each function group is marked on the top left panel for each sub-figure. See the interactive figures with gene IDs and annotations in Supplemental Dataset 2.

See more panels of Supplemental Fig. 4 below.

Supplemental Fig. 4. continued

C. Transcripts related to HSP90s

D. Transcripts related to HSP100s

Supplemental Fig. 4. continued

E. Transcripts related to small HSPs

F. Transcripts related to LHCII

Supplemental Fig. 4. continued

G. Transcripts related to PSII biogenesis

H. Transcripts related to PSII subunits

Supplemental Fig. 4. continued

I. Transcripts related to LHCI

J. Transcripts related to PSI subunits

K. Transcripts related to Cytochrome *b₆f* subunits

L. Transcripts related to ATP synthase subunits

Supplemental Fig. 4. continued

M. Transcripts related to Calvin-Benson cycle

N. Transcripts related to carbon concentrating mechanism (CCM)

O. Transcripts related to photoprotection

P. Transcripts related to cyclic electron flow around PSI (CEF)

Supplemental Fig. 4. continued

Q. Transcripts related to chlorophyll biosynthesis

Supplemental Fig. 4. continued

S. Transcripts related to ROS scavenging: ascorbate peroxidase (APX)

T. Transcripts related to ROS scavenging: catalase (CAT)

Supplemental Fig. 4. continued

U. Transcripts related to ROS scavenging: glutathione reductase (GR)

V. Transcripts related to ROS scavenging: glutathione S-transferase (GST)

W. Transcripts related to cell cycle

X. Transcripts related to cell wall

Supplemental Fig. 4. Continued

Y. Transcripts related to protein synthesis, ribosomal protein, prokaryotic

Z. Transcripts related to protein synthesis, ribosomal protein, chloroplast

Supplemental Fig. 4. Continued

AA. Transcripts related to mitochondrial electron transport / ATP synthesis (NADH-DH type 1)

AB. Transcripts related to mitochondrial electron transport / ATP synthesis

Supplemental Fig. 4. Continued

AC. Transcripts related to RNA

AD. Transcripts related to RNA processing

Supplemental Fig. 4. Continued

AE. Transcripts related to transport. phosphate

Time, hour

AF. Transcripts related to signaling. mitogen-activated protein kinase (MAP kinase)

Time, hour

Supplemental Fig. 4. Continued

AG. Transcripts related to cell motility. eukaryotes. axonemal dyneins

AH. Transcripts related to cell motility

Supplemental Fig. 4. Log$_2$(fold change, FC) of mean normalized read counts of groups of transcripts based on MapMan function annotation or manual curation. Plots are Log$_2$FC of transcripts related to the indicated function groups under six different treatments, as compared to the pre-heat time points. RNA-seq data with normalized read counts were used for this analysis. Black vertical lines indicate the start of heat treatments under 35°C or 40°C or the equivalent time point at 25°C. Each line represents one unique transcript in the indicated function group. ANOVA test was applied to the time course (five time points) of one transcript for each condition. If a transcript differed significantly during the time course (or underwent a global significant change) of one condition, it is colored purple (ANOVA significant); blue lines had no significant changes during the time course. Different intensities of purple and blue colors were used to distinguish different transcripts. The total number of transcripts plotted in each function group is marked on the top left panel for each sub-figure. See the interactive figures with gene IDs and annotations in Supplemental Dataset 2.

Supplemental Fig. 5A

Acetate depleting 25vs35

-log$_{10}$ (p value)

Supplemental Fig. 5B

Acetate depleting 25vs40

Supplemental Fig. 5C

Constant acetate 25vs35

Supplemental Fig. 5D

Constant acetate 25vs40

Supplemental Fig. 5E

Temperature

-log$_{10}$ (p value)

Supplemental Fig. 5F

Supplemental Fig. 5G

Acetate depleting 25vsConstant acetate 25

Supplemental Fig. 5. MapMan ontology enrichment revealed differentially regulated biological processes under different conditions. Ontology enrichment was performed for transcripts with significantly differential expression in the indicated comparisons using the extended MapMan ontology. The p values were calculated using a hypergeometric test. All functional sets with a p value less than 0.05 and more than 10 annotated transcripts

are shown. The ratio at the end of each bar indicates the number of significant transcripts between the two compared conditions vs detected transcripts in our RNA-seq data contained in the respective function bin. Multiple testing correction was performed by Benjamini & Hochberg FDR, * FDR < 0.1, ** FDR < 0.05. The bars are sorted based on the FDR p values, from top to bottom, from the small to big FDR values. **(A)** acetate-depleting 25°C vs acetate-depleting 35°C; **(B)** acetate-depleting 25°C vs acetate-depleting 40°C; **(C)** constant-acetate 25°C vs constant-acetate 35°C; **(D)** constant-acetate 25°C vs constant-acetate 40°C; **(E)** temperature effect: (constant-acetate 35°C and acetate-depleting 35°C) vs (constant-acetate 40°C and acetate-depleting 40°C); **(F)** medium effect: (constant-acetate 35°C and constant-acetate 40°C) vs (acetate-depleting 35°C and acetate-depleting 40°C); **(G)** acetate-depleting 25°C vs constant-acetate 25°C. See interactive figures and detailed enrichment analysis in Supplemental Dataset 3.

Supplemental Fig. 6

Supplemental Fig. 6. **Chlamydomonas cells have different morphologies at different temperatures under the acetate-depleting condition.** Light microscopic images of Chlamydomonas cells under the acetate-depleting condition. Images shown are representative results from at least three biological replicates. The algal cultivation and heat treatments were the same as in Fig. 1B.