Cerebral oxygenation and cardiac output responses during short repeated-sprints exercise and

modulatory effect of glucose ingestion

Paulo A. S. Armada-da-Silva[1,2,*]; Mingzhu, Hu[3,*]; Zongze, Wu[3]; Linjian, Wen[3]; Ruisen, Feng[3]; Zeng, Xinglin[4];

Yuan, Zhen[4]; Kong, Zhaowei[3]

[1] Faculdade de Motricidade Humana, Universidade de Lisboa, Portugal

[2] CIPER, Faculdade de Motricidade Humana, Universidade de Lisboa, Portugal

[3] Faculty of Education, University of Macao, Macao SAR, China

[4] Centre for Cognitive and Brain Sciences, University of Macao, Macao SAR, China

Running head: Cerebral oximetry during short, repeated sprints

Keywords: High-intensity interval exercise, brain oximetry, haemodynamic function, near infrared

spectroscopy

Corresponding author: Armada-da-Silva, P. Faculdade de Motricidade Humana, Estrada da Costa,

Portugal, parmada@fmh.ulisboa.pt

* P.A.S. and H.M. have contributed equally to this work.

Abstract

In this study we investigated changes in concentrations of cerebral oxyhaemoglobin (O_2Hb),

deoxyhaemoglobin (HHb), and total haemoglobin (tHb) during repeated sprints and their relationship

with cardiac output. We also examined the effect of glucose ingestion and acute hyperglycaemia on

cerebral haemoglobin responses. Ten young male participants ingested either a glucose drink (70 g) or a

placebo before performing a set of 10 repeated 6-second sprints on a cycle ergometer. Heart rate, stroke

volume, and cardiac output were measured continuously using impedance cardiography, while changes

in O_2Hb, HHb, and tHb in the frontal region of the cerebral cortex were measured using near-infrared

spectroscopy (NIRS). The results showed that each sprint elicited a transient increase in O_2Hb and to

lesser extent HHb concentrations, which was enhanced with the number of sprint repetitions and

correlated with cardiac output. After each sprint, O_2Hb and HHb quickly returned to baseline, while

cardiac output remained elevated. At the end of the repeated sprints, O_2Hb was decreased to below pre-

exercise levels, while HHb and tHb were elevated. After a recovery of 10-15 min, O_2Hb returned to pre-

exercise values in the placebo trial, but increased to above pre-exercise values (reactive hyperaemia) in

the glucose trial. Our findings suggest that short sprint exercise increases O_2Hb, HHb, and tHb levels

during exertion in parallel with cardiac output. However, in addition to the transient increase in cerebral

haemoglobin, a progressive decline in cerebral oxygen saturation occurs during repeated sprints. Glucose

ingestion does not alter cerebral haemoglobin responses to sprint exercise but appears to be associated

with faster recovery of O_2Hb.

Introduction

Studies investigating cerebral blood circulation have reported an increase in regional cerebral blood flow

(rCBF) during dynamic exercise of moderate-to-severe exercise intensity [i.e., ~ 70% maximal oxygen

uptake ($\dot{V}O_{2max}$)]. However, when the work rate increases to maximal exhaustive levels, rCBF decreases

towards baseline [1,2]. In addition to augmented CBF, dynamic exercise also results in an increase in

cerebral metabolism, as evidenced by a lowered cerebral metabolic ratio [3,4], indicating enhanced neural

activity [5]. This response is typically observed during exhaustive exercise [6], exercise in hyperthermic and

hypoxic conditions, or exercise under increased mental effort [4].

In addition to changes in rCBF, dynamic exercise also affects cerebral oxygenation [7]. Changes in

concentration of oxyhaemoglobin (O_2Hb) and deoxyhaemoglobin (HHb) in the cerebral tissue can be non-

invasively measured with near-infrared spectroscopy (NIRS) with a reasonable spatial resolution (~1-2

cm) and high time resolution. Two independent meta-analyses of NIRS studies have shown that O_2Hb

concentration in the frontal region of the cerebral cortex increases during submaximal exercise but

decreases as exercise approaches maximal intensity, suggesting a mismatch between oxygen delivery and

demand at the end stages of incremental maximal exercise [8,9]. Such mismatch has been related with a

cerebral vasoconstriction in response to exercise-induced hyperventilation and hypocapnia [2,10]. The

lowered cerebral oxygenation at near-maximal exercise, combined with muscle deoxygenation, may

impair muscle recruitment and trigger fatigue [2].

Extremely short all-out sprint interval exercise (<10 s) is an attractive option for achieving

cardiorespiratory fitness [11], but the cerebrovascular response to this type of exercise is unclear. One

study showed that a 30-second all-out cycling exercise at normal ambient oxygen levels and in hypoxia

resulted in a decrease in middle cerebral artery blood flow mean velocity (MCAv) and cerebral

oxygenation, which were unrelated to changes in end-tidal carbon dioxide partial pressure ($P_{ET}CO_2$) [12]. In

addition, the MCAv decrease during the 30-second all-out cycling did not follow the changes in cardiac

output and mean blood pressure, which increased during the exercise [12]. In another study, participants

performed seven 30-second exercise bouts on a cycle ergometer at a work rate of 150% maximal oxygen

uptake, where O_2Hb concentration increased during the first bout but decreased in the last three bouts,

and HHb concentration increased from resting levels, without a clear correlation between

cerebrovascular response and $P_{ET}CO_2$ [13]. These two studies suggest that, unlike submaximal and maximal

exercise, cerebrovascular response during supramaximal exercise is not strictly dependent on arterial CO_2

levels, mean blood pressure, or cardiac output. However, the impact of repeated short sprint efforts on

cerebral blood flow and oxygenation, and how it affects performance, remain unknown. It is worth noting

that dynamic cerebral autoregulation, which is another major mechanism regulating cerebrovascular

function, has a time delay similar to the duration of short sprint bouts (~6 s) [14], which may impair its

protective role in preventing a steep rise in cerebral blood flow and the risk of hyperperfusion during all-

out exercise.

Blood glucose levels play a role in determining cerebral metabolism and CBF [15]. Although glucose is the

primary energy substrate utilized by the brain, acute hyperglycaemia has been shown to decrease CBF

and the cerebral metabolic rate of both oxygen and glucose [15,16], although opposite findings have been

reported [17]. On the other hand, glucose ingestion has been shown to improve cerebral glucose uptake

and reduce perceived exertion during prolonged exercise, suggesting a potential ergogenic effect of

glucose through central mechanisms [18]. Hence, we also tested the effect of acute glucose intake and

subsequent rise in glycaemia in cerebral haemoglobin concentration changes during repeated sprints and

how it relates with performance.

Therefore, this study aimed to investigate changes in cerebral O_2Hb, HHb, and tHb during the

performance of repeated cycling sprints and their relationship with both cardiac output and power

output. The effect of acute hyperglycaemia on cerebral (prefrontal) oxygenation during the repeated

sprints was also investigated. A NIRS probe scanning the entire prefrontal cortex was used to assess brain

oximetry, which allows compensating for interindividual cerebral circulatory regional differences and to

investigate differences in cerebrovascular responses to short intermittent cycling sprints across the

frontal region of the cerebral cortex.

Methods

Participants and experimental design

Eleven healthy male adults volunteered (Table 1) to participate in this study after providing written

informed consent. Participants were free of any metabolic, cardiovascular, or neurological disease and

did not regularly participate in exercise training. The study protocol was reviewed and approved by the

ethics review board of the University of Macao (Special Administrative Region of Macao, China) and

conformed to the ethical principles for research involving human participants (approved number:

BSERE22-APP011-FED).

Each participant visited the laboratory on three separate occasions at least one week apart. In the first

visit, each participant performed a maximal incremental cycling exercise to determine the individual

$\dot{V}O_{2max}$. In the following visits, each participant performed a repeated sprint exercise test on a cycle

ergometer after ingesting 70 g of glucose (GLU) or a taste-matched non-caloric placebo (PLA), in a

randomised counterbalanced and single-blind design.

Instrumentation and measures

Maximal incremental test

The maximal incremental test was carried out on a cycle ergometer (Monark 839E, Varberg, Sweden).

After a warm-up at 25 W for 3 min, participants started cycling at 50 W and a cadence of 60 ± 5

revolutions per minute (rpm), with the workload increasing by 25 W every 3 min. The test was completed

when the participant reached volitional exhaustion, with a respiratory exchange ratio (RER) ≥ 1.15,

ratings of perceived exertion (RPE) ≥18, and peak HR ≥ 90% of age-predicted maximum (220-age) [19].

Respiratory gases were collected by the Vmax Encore System (CareFusion Corp., San Diego, CA, USA)

throughout the test, and $\dot{V}O_{2max}$ was calculated as the highest oxygen consumption value averaged over

15 s before the end of the test [11].

Repeated sprints cycling exercise

During the two experimental sessions, participants performed a repeated sprint set of 10 repetitions of 6

s maximal sprints interspersed by 24 s resting, performed on a cycle ergometer (Monark Ergomedic 894E,

Monark Exercise AB, Vansbro, Sweden). Each participant sat upright on the cycle ergometer keeping the

head as steady as possible with the help of a neck cushion to minimise motion artefacts during NIRS

recordings. Before the repeated sprints, a 5-min warm-up, cycling at a power output of 60 W (resistance,

1 kg; cadence, 60 rpm) was completed. During the sprints, participants were instructed to use the same

ready position, sitting upright and with one leg ready to push powerfully against the pedal and maximally

accelerate the cycling crank at the "go" signal. Each participant was free to choose his leading leg in the

first session and was asked to use the same leg to initiate all subsequent sprint repetitions. Five seconds

into the next sprint, participants adopted the ready position and awaited the verbal go command at the

end of a 3-s countdown. A weight equal to 6.5% body mass was applied to the flywheel controlled by the

Monark Anaerobic Test Software at the instant the pedalling cadence reached 90 rpm. The controlling

software recorded the mean power output for each sprint.

NIRS

A portable 2 x 4-channel continuous-wave NIRS system (Octamon, Artinis Medical Systems, Einsteinweg,

The Netherlands) was used to measure changes (Δ) in O_2Hb, HHb, and tHb concentrations using two

wavelengths at 750 and 850 nm. Data were acquired with the OxySoft software (OxySoft, Artinis Medical

Systems, Einsteinweg, The Netherlands) at a frequency of 50 Hz with a differential path length factor of

six. The optodes were attached to a dark ambient light-protecting probe holder, tightly secured to the

head by adjustable headbands and placed over the forehead and the underlying prefrontal cortex (four

channels on each side). The two light detectors were aligned on Fp1 and Fp2 locations of the 10-20

system for the electroencephalography and surrounded by four light-emitting optodes with an inter-

optode distance of 35 mm, providing a total of eight NIRS channels.

Impedance cardiography

Cardiac output was measured continuously during the sessions by signal morphology impedance

cardiography (PhysioFlow PF-05, Manatec biomedical, Paris, France). After skin preparation to minimise

electrical resistance, six Ag/AgCl electrodes (Hewlett Packard 40493 E) were placed on the chest wall

along the xiphoid process and at the V1 and V6 precordial locations, and at the base of the neck to record

the thoracic impedance and electrocardiographic signals. Resting blood pressure was measured by

automatic oscillometry to calibrate the cardiac impedance data acquisition following the manufacturer

instructions. Stroke volume and cardiac output were recorded for each cardiac cycle and measured

continuously throughout the trial.

Blood glucose and lactate

Blood glucose measurements were collected with a glucometer (ACCU–CHEK Performa, USA) and the

proper test strips (F. Hoffmann-La Roche AG, Basel, Switzerland). Blood lactate was also measured with

the Lactate Pro 2 handheld device and respective test strips (AKRAY Europe B.V., Amstelveen, the

Netherlands). Both blood analytes were measured in duplicate from capillary blood drawn from a

lanceted fingertip at three time points: 1) at baseline, soon after arrival to the laboratory; 2) immediately

before starting the exercise protocol; and 3) immediately after completion of the repeated sprints

exercise. The duplicate blood sample measurements were averaged for analysis.

Procedures

Participants arrived at the laboratory between 8:00 and 9:00 am after a 12-hours fast. Soon after arrival,

participants ingested 300 ml of a drink containing 70 g of glucose (Cambridge Isotope Laboratories,

Cambridge, MA, USA) or a taste-matched placebo containing artificial sweeteners (Truvia Calorie-Free

Sweetener; Cargill, IL, USA). Next, they rested sitting for one hour. Fifteen minutes to the end of the one-

hour resting, participants were fitted with the cardiac output, and NIRS instrumentations and the two

devices were set up for acquisition. The trial started with 10 minutes of seated resting, while wearing a

cushion around the neck to reduce head movement.

After the 10-min rest, participants mounted the cycle ergometer and completed the 5-min warm-up

followed by the 10 x 6-s repeated sprints. After exercise, there were another 10 minutes resting. Ratings

of perceived exertion were obtained at the end of the fifth and tenth sprint repetitions using the Borg's

scale.

Data analysis

The NIRS data pre-processing was performed using the Homer2 toolbox [20] for Matlab (Matlab version

9.5.0, release R2018b, The Mathworks Inc., Natick, Massachusetts, USA). First, light absorption data were

converted into optical density and haemoglobin concentrations using the modified Beer-Lambert law

with a differential path length factor of six. Next, all NIRS signals were low pass filtered at a cut-off

frequency of 0.5 Hz to remove heartbeat-associated changes in O_2Hb, HHb and tHb, using a finite impulse

response fourth-order Butterworth filter, implemented in the forward and inverse directions to avoid

phase filtering artefacts [20]. After filtering, the whole NIRS signal was epoched into the individual sprints,

starting at minus 5 s and ending at 25 s after the sprint start. The median was used to quantify the

changes in haemoglobin concentrations in the time intervals 0-15 s and 15-25 s, to measure the response

during the sprints and during the rest interval between sprints, respectively. The median was chosen

because of its higher robustness against potential biasing effects caused by motion artefacts and outliers.

The time interval 0-15 s was selected following a visual inspection of the transient ΔO_2Hb response

accompanying the performance of the sprints (Fig. 3). Due to the non-normal distribution of some

ΔO_2Hb, ΔHHb, and ΔtHb values (Supporting information, Figs. S1-S3 and Table S1), a natural log-

transformation was applied to these variables for the purpose of statistical analysis.

Statistical analysis

The Shapiro-Wilk test was used to verify the normal distribution of the data. To assess the responses to

repeated sprint exercise, a repeated measures ANOVA with a complete factorial design (cerebral

oximetry data: 2 trials x 8 channels x 10 sprint repetitions; power output and cardiovascular data: 2 trials

x 10 sprint repetitions) was conducted. The Greenhouse-Geiser correction was applied to within-subject

factors violating the sphericity assumption (Mauchly's test p-value < 0.05). Post hoc tests were performed

employing one-way ANOVA with Bonferroni's correction for family-wise error rate. Planned contrasts

were performed using polynomial fitting. This approach treated the repeated sprints as a quantitative

variable with even spacing. While the polynomial contrast analysis does not directly compare two specific

sprint repetitions, it provides valuable information about the overall trend effect of the repeated sprints

on the response variables. In case of a significant linear contrast, backward differences contrasts were

computed (the contrast between two consecutive sprint repetitions, including resting). Further contrasts

between the first repeated sprint and the following sprints were also calculated. One sample *t*-test was

used for testing the significance of post-exercise mean ΔO_2Hb, ΔHHb and ΔtHb against pre-exercise

values (since continuous NIRS measurements are relative, O_2Hb and HHb baseline concentrations are set

to zero). Correlation values were computed using the Pearson's coefficient. All statistical tests were

performed in RStudio (v. 1.3.1093, 2009-2020 RStudio, PBC) using functions from Rstatix (v. 0.7.0) and R

base (v. 4.0.3) packages. Unless otherwise stated, all data are reported as mean $\pm$ standard deviation

(SD). Significance was accepted at $p < 0.05$.

Results

Due to large noise affecting the NIRS signals, one participant was excluded from the final analysis,

resulting in a final sample size of 10 participants. In another participant, the NIRS signals pertaining to

three of the sprints in the two trials were removed from analysis because of significant motion artefacts.

The missing data (48 out of 1600 data points) were imputed using the average values from the same

individual, channel, and the other sprints. One participant in the PLA trial had missing mean power

output data for the two final repeated sprints due to equipment failure. The last recorded mean power

output was used to replace the missing values.

Power output and RPE

Figure 1 displays the mean power output for each sprint and trial. The mean power output was found to

be higher during the GLU trial than the PLA trial [GLU: 648 ± 183 W; PLA: 558 ± 161 W; $F_{(1, 9)} = 6.312$, p =

0.033, η^2 = 0.098], particularly in the later sprint repetitions [trial x sprint repetition interaction effect:

$F_{(2.79, 25.12)}$ = 12.314, p < 0.000, η^2 = 0.203]. According to one-way ANOVA, mean power output was

significantly different between GLU and PLA in the first (PLA higher than GLU), seventh and tenth sprint

repetitions (Bonferroni's corrected p < 0.05, Fig. 1). At the end of the fifth repetition, perceived exertion

was moderate (GLU: 11.9 $\pm$ 2.1; PLA: 13.2 $\pm$ 1.9 units), but RPE reached near-maximal scores at the end of

the sprint set (GLU: 19.3 $\pm$ 2.1; PLA: 19.3 $\pm$ 0.5 units; main effect of sprint repetition, $F_{(1, 9)}$ = 232.66, p <

0.000, η^2 = 0.856), with no differences between trials.

Blood glucose and lactate

At baseline, blood glucose concentration was similar in the two trials (GLU: 5.7 $\pm$ 0.5 mmol·l^{-1}; PLA: 5.9 $\pm$

0.6 mmol·l^{-1}; p = 0.310). However, following glucose ingestion, the blood glucose concentration rose and

was significantly higher in the GLU trial compared to the PLA trial prior to exercise (GLU: 8.1 $\pm$ 1.7 mmol·l^{-}

^{1}; PLA: 6.1 $\pm$ 0.8 mmol·L^{-1}; p = 0.016). By the end of the repeated sprints, the blood glucose concentration

returned to similar levels in both trials (GLU: 6.5 $\pm$ 1.6 mmol·l^{-1}; PLA: 5.9 $\pm$ 0.9 mmol·l^{-1}; p = 0.254).

At baseline, the blood lactate concentration was at the same level in the GLU and PLA trials (GLU: 1.7 $\pm$

0.9 mmol·l^{-1}; PLA: 1.4 $\pm$ 0.4 mmol·l^{-1}; p = 0.335). However, after glucose ingestion, the blood lactate

concentration increased significantly (GLU: 2.4 $\pm$ 0.8 mmol·l^{-1}; PLA: 1.6 $\pm$ 0.5 mmol·l^{-1}; p = 0.023).

Immediately after the repeated sprints, the blood lactate concentration was markedly elevated and

significantly higher in the PLA trial than in the GLU trial (GLU: 10.8 $\pm$ 1.1 mmol·l^{-1}; PLA: 11.7 $\pm$ 1.4 mmol·l^{-}

^{1}; p = 0.047).

Heart rate, stroke volume, and cardiac output

Figure 2 shows the changes in heart rate, stroke volume and cardiac output during the repeated sprints.

The heart rate increased progressively throughout the sprints set, reaching its highest value of 163 $\pm$ 5

beats·min^{-1} in the final sprint [main effect for sprint repetition, excluding resting values: $F_{(3.45, 31.04)}$ =

347.947, p < 0.000, η^2 = 0.884], with no significant difference between GLU and PLA [$F_{(1, 9)}$ = 0.018, p =

0.897, η^2 < 0.000].

The stroke volume increased from 84.2 $\pm$ 5.8 ml at rest to 97.3 $\pm$ 8.4 ml during the first sprint, but

remained at a plateau after the fourth repetition [main effect for sprint repetition, excluding resting

values: $F_{(2.47, 22.25)}$ = 39.546, p < 0.000, η^2 = 0.380]. There was no significant difference in stroke volume

between GLU and PLA trials [$F_{(1, 9)}$ = 0.195, p = 0.669, η^2 = 0.005].

Cardiac output increased in a curvilinear pattern during the repeated sprints, rising from a resting value

of 7.2 $\pm$ 0.7 l·min^{-1} to a cardiac output of 10.7 $\pm$ 1.0 l·min^{-1} in the first sprint and 19.1 $\pm$ 1.1 l·min^{-1} at the

end of the repeated sprints [main effect for sprint repetition, excluding resting values: $F_{(2.59, 23.31)}$ =

214.561, $p < 0.000$, $\eta^2 = 0.792$]. There was no significant difference in cardiac output between GLU and

PLA [$F_{(1, 9)} = 0.280$, $p = 0.610$, $\eta^2 = 0.005$], as well as no interaction effect between trial and sprint

repetition [$F_{(1.93, 17.38)} = 1.122$, $p < 0.346$, $\eta^2 = 0.025$].

During the time interval between sprints, cardiac output remained high, but decreased slightly, although

significantly, from the level attained during the sprints [overall mean $\pm$ SD during sprinting, 16.2 ± 2.8

$l \cdot min^{-1}$; in between sprints, $16.1 \pm 2.8 \, l \cdot min^{-1}$; $F_{(1, 9)} = 6.084$, $p = 0.036$, $\eta^2 = 0.003$]. Polynomial contrasts

showed a significant linear and quadratic effect of sprint repetition on heart rate, stroke volume, and

cardiac output responses. Several backward differences contrasts were also significant (see Fig. 2 for

details).

Brain oximetry

Table 2 presents the results of ΔO_2Hb, ΔHHb, ΔtHb during the repeated sprints and the rest periods

between the sprint repetitions.

Oxyhaemoglobin

During the sprints, O_2Hb levels rapidly increased, reaching its peak at the end of each sprint (Fig. 3;

Supporting information, Fig. S4). The O_2Hb response to the sprints increased with the increasing number

of sprint repetitions [main effect of sprint repetition: $F_{(9, 81)} = 3.380$, $p = 0.001$, $\eta^2 = 0.074$]. However,

there was no effect of glucose ingestion [$F_{(1, 9)} = 0.930$, $p = 0.360$, $\eta^2 = 0.005$] or NIRS channel [$F_{(7, 63)} =$

10.222, $p = 0.979$, $\eta^2 = 0.002$] on this response. Polynomial fitting demonstrated a significant linear effect

of sprint repetition on ΔO_2Hb, which did not improve with higher-order components. Additionally,

contrast analysis showed significantly lower ΔO_2Hb in the first sprint compared to the fourth and

subsequent sprints ($p < 0.05$). After each sprint, O_2Hb concentrations returned to pre-sprint levels or

lower (Fig. 3, Table 2). There were no significant differences in ΔO_2Hb values between trials, channel, or

sprint repetitions for the rest periods between the sprints.

Deoxyhaemoglobin

In comparison to the changes in O_2Hb, the changes in HHb in response to sprinting were less pronounced

and not affected by the sprint repetitions [$F_{(9,81)} = 1.446$, $p = 0.182$, $\eta^2 = 0.01$] (Table 2, Fig. 3; Supporting

information Fig. S4). Also, there were no significant differences in the changes of HHb between the two

trials [$F_{(1, 9)} = 0.476$, $p = 0.508$, $\eta^2 = 0.000$] (Table 2) or between the NIRS channels [$F_{(7,63)} = 1.531$, $p = 0.173$,

$\eta^2 = 0.02$]. During the rest interval between the sprints, HHb concentration returned to pre-exercise

levels with no differences observed between the trials [$F_{(1, 9)} = 0.287$, $p = 0.605$, $\eta^2 = 0.000$] or NIRS

channels [$F_{(7,63)}$= 1.162, p = 0.337, η^2 = 0.01], but with a significant main effect of sprint repetition [$F_{(9, 81)}$= 2.152, p = 0.034, η^2 = 0.012]. Polynomial contrasts revealed only marginal linear (p = 0.052) and quadratic (p = 0.057) effects of the repeated sprints in ΔHHb in the rest interval between the sprints.

Total haemoglobin

The changes in tHb followed a similar pattern to that observed for O$_2$Hb, with a marked increase during each sprint, followed by a rapid decline towards the baseline immediately after the sprint. Glucose ingestion [$F_{(1, 9)}$= 0.970, p = 0.350, η^2 = 0.007] and NIRS channels [$F_{(7, 63)}$= 0.403, p = 0.897, η^2 = 0.006] had no effect on the changes in tHb. A significant difference was observed in ΔtHb across the repeated sprints [$F_{(9, 81)}$= 2.796, p = 0.007, η^2 = 0.054], with a significant linear effect (p = 0.008) and lower ΔtHb observed during the first sprint compared to most of the subsequent sprints (0.003 < p < 0.046). No differences in ΔtHb during the time interval between sprints were observed between trials [$F_{(9, 81)}$= 2.017, p = 0.189, η^2 = 0.002], NIRS channels [$F_{(7, 63)}$= 1.016, p = 0.429, η^2 = 0.017], or sprint repetitions [$F_{(9, 81)}$ = 0.866, p = 0.560, η^2 = 0.010].

The concentrations of O$_2$Hb, HHb, and tHb were compared before, at the end of exercise, and 10-15 minutes post-exercise (see Fig. 3, top and middle panels). The repeated sprint performance caused a significant reduction in cerebral O$_2$Hb concentration (one sample *t*-test, combining GLU and PLA: t(8) = -2.644; p = 0.029, n = 9) that was comparable between the two trials [GLU: -1.16 $\pm$ 4.13 μM; PLA: -3.11 $\pm$ 2.87 μM; $F_{(1, 7)}$ = 2.453, p = 0.161, η^2 = 0.096, n = 9]. On the other hand, there was a steady increase in cerebral HHb concentration during the repeated sprints (one sample *t*-test, combining GLU and PLA: t(8) = 3.248, p = 0.011, n = 9), with no effect of glucose ingestion on this response [GLU: 3.80 $\pm$ 5.95 μM; PLA: 6.44 $\pm$ 3.82 μM; $F_{(1, 7)}$ = 1.541, p = 0.254, η^2 = 0.042, n = 9]. Due to the increase in HHb concentration, tHb also increased after the sprint set (one sample *t*-test, combining GLU and PLA: t(8) = 3.119, p = 0.014, n = 9) regardless of glucose ingestion [GLU: 3.98 $\pm$ 5.60 μM; PLA: 5.75 $\pm$ 3.86 μM; $F_{(1, 7)}$ = 0.670, p = 0.440, η^2 = 0.014, n = 9].

Following the exercise, there was an increase in cerebral O$_2$Hb back to baseline levels, particularly in the GLU trial [GLU: 2.54 $\pm$ 1.80 μM; PLA: 0.95 $\pm$ 2.32 μM; $F_{(1, 7)}$ = 11.263, p = 0.012, η^2 = 0.140, n = 9]. As a result, post-exercise O$_2$Hb was higher than pre-exercise levels in the GLU trial (one sample *t*-test, t(8) = 8.099, p < 0.000, n = 9) but not in the PLA trial (one sample *t*-test, t(8) = 2.018, p = 0.08, n = 8). At 10-15 min post-exercise, HHb remained elevated in both trials [GLU: 3.42 $\pm$ 4.02 μM; PLA: 3.55 $\pm$ 3.15 μM; $F_{(1, 7)}$ = 0.023, p = 0.884, η^2 < 0.000, n = 9] compared to pre-exercise values (one sample *t*-test, combining GLU and PLA: t(8) = 4.557, p = 0.001, n = 9). Following changes in HHb and O$_2$Hb, tHb concentration also increased after the repeated sprint exercise in both trials [Glu: 3.29 $\pm$ 4.07 μM; Pla: 3.12 $\pm$ 3.31 μM; $F_{(1, 7)}$

= 0.050, p = 0.829, η^2 < 0.000, n = 9], and at 10-15 min post-exercise, it was higher than at pre-exercise

(one sample *t*-test, combining GLU and PLA: t(8) = 3.882, p = 0.005, n = 9).

Cardiac output and ΔO_2Hb, ΔHHb and ΔtHb relationship

Figure 4 illustrates the relationship between cardiac output, power output, and cerebral haemoglobin

concentrations (see also Supporting information, Fig. S5). During the sprints, positive correlations were

found between cardiac output and the changes in O_2Hb and ΔtHb in both GLU (cardiac output: ΔO_2Hb, R

= 0.85, p = 0.0019; cardiac output: ΔtHb, R = 0.86, p = 0.0013) and PLA trials (cardiac output: ΔO_2Hb, R =

0.87, p = 0.0012; cardiac output: ΔtHb, R = 0.85, p = 0.0016). No significant correlation was observed

between cardiac output and ΔHHb during the repeated sprints, nor between cardiac output and ΔO_2Hb,

ΔHHb and ΔtHb during the time interval between the sprints. Conversely, in the GLU trial, power output

was negatively correlated with ΔO_2Hb values during the time interval between the sprints.

Discussion

In this study, our aim was to outline the time profile of cerebral O_2Hb, HHb and tHb responses to a series

of brief, repeated sprints and investigate the correlation between this response and cardiac output.

Furthermore, we sought to investigate the impact of acute hyperglycaemia on the changes in O_2Hb, HHb,

and tHb following the repeated sprint exercise. Our findings indicated that each sprint produced a rapid

and significant surge in cerebral O_2Hb concentration, and to a lesser extent HHb, leading to a transient

increase in cerebral blood volume (i.e., increased tHb) and oxygenation. During the interval between the

repeated sprints, O_2Hb and HHb decreased and returned to, or below, pre-exercise levels rapidly. The

variations in ΔO_2Hb and ΔtHb followed a curvilinear trend similar to that observed for heart rate and

cardiac output in response to the repeated sprint exercise.

During each of the short sprints, there was an increase in O_2Hb, which was approximately ten times

higher than the pulsatile peak increase due to the heartbeat at rest (~0.5 μM). Regulation of cerebral

blood flow during rest and exercise is closely tied to changes in arterial blood gases (particularly carbon

dioxide), neural activity, blood pressure, and cardiac output [21–23]. An increased cardiac output and

cerebral perfusion pressure, caused by an elevation of arterial blood pressure, are two possible

explanations for the transient increase in O_2Hb and HHb that was observed during each short sprint

[12,24,25]. Although we did not record blood pressure, previous studies have shown that heavy-intensity

dynamic (e.g., rowing exercise) and resistance (e.g., leg-press exercise) exercise result in a rapid increase

in diastolic and systolic blood pressures, therefore it is reasonable to expect that mean blood pressure

would increase during all-out sprint exercise [12,25–27]. Although enhanced central drive and activation of the

metaboreflex may contribute to the sprint-induced elevation of blood pressure [28], the immediate and

transient increase in O_2Hb and HHb during the short sprint exercise suggests a role for mechanical and

haemodynamic factors associated with the powerful contraction of the leg and trunk muscles, resulting in

brief Valsalva-like manoeuvres and synchronous elevation of arterial blood pressure [24,27,29]. Indeed, the

production of Valsalva-like manoeuvres is commonly observed during heavy static and dynamic

resistance exercise dynamic exercise [27,30]. However, this manoeuvre can also occur during dynamic

exercise, such as during the catch phase of rowing [24,29], and is associated with the cyclical increase in

MCAv in synchrony with the elevations of intra-abdominal pressure, central venous pressure, and catch

force [24]. Similarly, the onset of bilateral static leg extension simultaneously raises mean arterial blood

pressure, MCAv, and cerebral O_2Hb concentration with or without a superimposed Valsalva strain [30].

Previous studies have shown a direct influence of cardiac output on cerebral blood flow and cerebral

oxygenation [23,31]. In this study, we found that cardiac output had a moderate and positive correlation

with changes in O_2Hb and tHb during the sprints. This suggests that cardiac output plays a role in the rise

of O_2Hb recorded during the sprints. However, the relationship between cardiac output and cerebral

blood flow is still unclear and in the present study O_2Hb concentration declined during rest between the

sprint bouts despite the increased cardiac output. While previous research has reported a direct

relationship between cardiac output and MCAv during rest and moderate intensity cycling exercise (Ogoh

et al., 2005), changes in MCAv during static leg exercise seem to develop independently of changes in

cardiac output [30]. Regarding our observations, we cannot refute the possibility that with increasing

number of repeated cycling sprints there was an enhanced blood pressure response resulting from

heightened exercise pressor reflex, central drive, and strongest Valsalva manoeuvre, thus increasing the

elevation in O_2Hb concentration across the sprints set, regardless of changes in cardiac output [28,32].

Despite its relationship with the changes in cerebral O_2Hb and tHb concentration during sprints, cardiac

output seems to have little influence on cerebral haemoglobin concentration levels between sprints.

Following each sprint, cerebral oxygenation levels rapidly returned to baseline, while cardiac output

remained elevated, indicating that a high cardiac output alone is insufficient to maintain elevated

cerebral oxygenation levels. The rapid decrease in cerebral oxygenation levels with the end of each sprint

could have been caused by a sudden decrease in mean arterial pressure upon cessation of the exercise [26].

Curtelin et al. [12] reported a drop in mean blood pressure immediately after the end of a Wingate test,

which was approximately three times larger than its increase during the effort. This decrease in mean

blood pressure was accompanied by a significant decline in MCAv, which lasted for approximately 40 s

before returning to pre-exercise levels [12]. However, a decrease in CBF at the end of exercise might occur

independently from changes in arterial blood pressure. For instance, Pott et al. [24] reported a continuous

decrease in MCAv after each rowing catch movement despite stable mean blood pressure, which the

authors suggest could have been caused by delayed response of dynamic cerebral autoregulation. A

similar event could have taken place during the repeated sprints, given the match between the sprints'

duration and the phase delay of cerebral autoregulation [14].

The findings of this study showed that the repeated sprints exercise led to a progressive decrease in

cerebral O_2Hb and an increase in HHb, with considerable interindividual variation (Fig 3). The rise in HHb

outweighed the decrease in O_2Hb, resulting in an increase in cerebral tHb at the end of the repeated

sprints exercise. These changes in cerebral oxygenation partially agree with MCAv data recorded during

the Wingate test, revealing decreased CBF and cerebral oxygenation levels caused by augmented

cerebral vascular resistance at later sprint stages [12]. A similar response is observed during heavy

resistance exercise, with cerebrovascular resistance increasing during exercise but lowering rapidly once

the muscular activity stops [26]. Thus, the data obtained from this study extends these observations,

showing similarly restrained cerebral blood flow and cerebral oxygen delivery during short cycling sprints,

even with moderate perceived exertion levels.

Dynamic cerebral autoregulation is a primary mechanism for regulating cerebrovascular resistance. While

the effect of exercise on this mechanism is controversial, with studies indicating decreased efficacy of

dynamic autoregulation during exhaustive aerobic exercise [33], other studies report unchanged or even

enhanced cerebral autoregulation during aerobic exercise [34]. The protracted CBF recovery from

hypotension after heavy aerobic exercise [35] and sprinting [12] suggests a diminished cerebral

autoregulation efficacy, possibly combined with a reduced gain of the arterial baroreflex [35], maybe to

compensate previous mismatch between oxygen delivery and demand. While the present study cannot

address the impact of short sprint performance on dynamic autoregulation, the rapid decrease of O_2Hb at

the end of the sprints when cardiac output was elevated suggests normal cerebral autoregulation during

this type of exercise. Future studies may test the effect of different sprint/recovery cycle durations on

cerebral haemoglobin responses. Importantly, a decline in O_2Hb during repeated sprints contrasts with

the increase in cerebral oxygenation during lower body resistance exercise [36]. Since decreased brain

oxygen saturation is observed in other sprint studies [12], it may be concluded that the extreme intensity of

sprint exercise and the recruitment of a large muscle mass reduce cerebral blood flow and lower cerebral

oxygenation. In addition, the cessation of the repeated sprints exercise was accompanied by a cerebral

reactive hyperaemic response, with a steep increase in O_2Hb within minutes following exercise (Fig. 3).

This strengthens the claim that cerebral oxygen delivery is reduced during repeated sprint exercise.

Finally, the absence of significant increases in both O_2Hb and tHb during the sprint set suggests that

repeated sprint performance does not cause a large increase in cerebral blood flow that could

endangered the brain.

In this experiment, we investigated the effect of glucose ingestion and acute hyperglycaemia on the

response of cerebral haemoglobin to sprint exercise. Although glucose is the primary energy source for

the brain, hyperglycaemia has been shown to decrease the cerebral metabolic rate of oxygen [16] and brain

connectivity [37] in healthy young participants at rest. Furthermore, acute and chronic hyperglycaemia

decreases cerebral blood flow [38]. In our study, we found that glucose ingestion did not affect O_2Hb and

HHb responses to the repeated sprints exercise, despite improving performance. However, glucose

ingestion significantly enhanced reactive hyperaemia, resulting in faster and higher levels of O_2Hb

concentration during the post-exercise period. The relationship between post-exercise hyperaemia and

any baseline changes in CBF or cerebral blood volume caused by glucose ingestion requires further

investigation.

Study limitations

There are several limitations in this study that should be acknowledged. One significant limitation is the

unknown effect of extracerebral haemodynamics (scalp and skull) on cerebral oximetry measurements.

Since the emitted light from NIRS travels through extracerebral tissues and the cerebral cortex, changes

in blood volume and flow in superficial tissues can affect the NIRS measurements. The design of the NIRS

probe used in this study, with rigid circular optodes holders firmly secured to apply pressure on the

forehead skin underneath the optodes, helped to minimise changes in skin blood flow and their impact

on NIRS recordings. Studies have shown that facial skin blood flow and vascular conductance increase

with dynamic exercise intensity without plateauing [39], while ocular circulation is similar to cerebral

circulation, decreasing from elevated values to baseline in response to exhaustive dynamic exercise [40].

Based on these findings, it can be inferred that the oximetry data obtained during the repeated sprints

exercise primarily reflect the changes in cerebral circulation. Future studies may assess the contribution

of forehead skin to changes in haemoglobin concentration in the frontal region of the cerebral cortex

during repeated sprints.

Another limitation of this study is the absence of measurements of $P_{ET}CO_2$ and blood pressure, which

could have provided a more comprehensive understanding of the physiological changes occurring during

the repeated sprints exercise.

Perspective

We provide evidence that the performance of short cycling sprints produces a significant impact on

cerebral oxygenation. We recorded a temporary increase in O_2Hb, HHb and tHb concentrations during

the sprints, with the magnitude of these changes growing with the number of sprint repetitions. In

addition, we found a positive correlation between changes in O_2Hb and cardiac output, which

progressively increased throughout the repeated sprint set and remained elevated during the rest

periods between sprint repetitions. However, O_2Hb rapidly returned to baseline or below during these

rest periods. Contrary to the sprint response, we observed a decrease in O_2Hb concentration throughout

the repeated sprint set. Conversely, HHb increased, suggesting inadequate brain oxygen delivery and

insufficient cerebral oxygenation oxygen delivery during the all-out repeated cycling. Although not

affecting haemoglobin concentration responses, glucose ingestion did improve performance and was

associated with a faster recovery of O_2Hb levels during the post-exercise period. In conclusion, our
findings suggest that repeated spring exercise elicits a transient and modest increase in cerebral O_2Hb
and tHb during the physical effort itself. However, there is an overall decrease in cerebral oxygenation,
leading to a robust post-exercise cerebral reactive hyperaemia in certain individuals. Importantly, our
data does not indicate a marked elevation in O_2Hb or cerebral blood flow during short all-out cycling
sprints in young, non-trained individuals, thereby reducing the likelihood of cerebral hyperperfusion
during this type of exercise.

References

1.   Ide K, Boushel, Sørensen, et al. Middle cerebral artery blood velocity during exercise with β-1
adrenergic and unilateral stellate ganglion blockade in humans. *Acta Physiol Scand*. 2000;170(1):33-38.
doi:https://doi.org/10.1046/j.1365-201x.2000.00757.x

2.   Subudhi AW, Lorenz MC, Fulco CS, Roach RC. Cerebrovascular responses to incremental exercise
during hypobaric hypoxia: Effect of oxygenation on maximal performance. *Am J Physiol Heart Circ*
Physiol. 2008;294(1):164-171. doi:10.1152/ajpheart.01104.2007

3.   Ide K, Horn A, Secher NH. Cerebral metabolic response to submaximal exercise. *J Appl Physiol*.
1999;87(5):1604-1608. doi:10.1152/jappl.1999.87.5.1604

4.   Dalsgaard MK, Ide K, Cai Y, Quistorff B, Secher NH. The intent to exercise influences the cerebral
O(2)/carbohydrate uptake ratio in humans. *J Physiol*. 2002;540(Pt 2):681-689.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=
11956354

5.   Dienel GA. The metabolic trinity, glucose–glycogen–lactate, links astrocytes and neurons in brain
energetics, signaling, memory, and gene expression. *Neurosci Lett*. 2017;637:18-25.
doi:10.1016/j.neulet.2015.02.052

6.   Rasmussen P, Nielsen J, Overgaard M, et al. Reduced muscle activation during exercise related to brain
oxygenation and metabolism in humans. *Journal of Physiology*. 2010;588(11):1985-1995.
doi:10.1113/jphysiol.2009.186767

7.   González-Alonso J, Dalsgaard MK, Osada T, et al. Brain and central haemodynamics and oxygenation
during maximal exercise in humans. *J Physiol*. 2004;557(1):331-342. doi:10.1113/jphysiol.2004.060574

8.   Rooks CR, Thom NJ, McCully KK, Dishman RK. Effects of incremental exercise on cerebral oxygenation
measured by near-infrared spectroscopy: A systematic review. *Prog Neurobiol*. 2010;92(2):134-150.
doi:https://doi.org/10.1016/j.pneurobio.2010.06.002

9.   De Wachter J, Proost M, Habay J, et al. Prefrontal Cortex Oxygenation During Endurance Performance:
484   A Systematic Review of Functional Near-Infrared Spectroscopy Studies. *Front Physiol*.
485   2021;12(October). doi:10.3389/fphys.2021.761232

486   10.   Ogoh S, Ainslie PN. Cerebral blood flow during exercise: Mechanisms of regulation. *J Appl Physiol*.
487   2009;107(5):1370-1380. doi:10.1152/japplphysiol.00573.2009

488   11.   Shi Q, Tong TK, Sun S, et al. Influence of recovery duration during 6-s sprint interval exercise on time
489   spent at high rates of oxygen uptake. *J Exerc Sci Fit*. 2018;16(1):16-20. doi:10.1016/j.jesf.2018.01.001

490   12.   Curtelin D, Morales-Alamo D, Torres-Peralta R, et al. Cerebral blood flow, frontal lobe oxygenation and
491   intra-arterial blood pressure during sprint exercise in normoxia and severe acute hypoxia in humans.
492   *Journal of Cerebral Blood Flow and Metabolism*. 2018;38(1):136-150. doi:10.1177/0271678X17691986

493   13.   Shibuya KI, Tanaka J, Kuboyama N, Ogaki T. Cerebral oxygenation during intermittent supramaximal
494   exercise. *Respir Physiol Neurobiol*. 2004;140(2):165-172. doi:10.1016/j.resp.2003.11.004

495   14.   Fantini S, Sassaroli A, Tgavalekos KT, Kornbluth J. Cerebral blood flow and autoregulation: current
496   measurement techniques and prospects for noninvasive optical methods. *Neurophotonics*.
497   2016;3(3):031411. doi:10.1117/1.nph.3.3.031411

498   15.   Xu F, Liu P, Pascual JM, Xiao G, Huang H, Lu H. Acute effect of glucose on cerebral blood flow, blood
499   oxygenation, and oxidative metabolism. *Hum Brain Mapp*. 2015;36(2):707-716.
500   doi:10.1002/hbm.22658

501   16.   Ishibashi K, Wagatsuma K, Ishiwata K, Ishii K. Alteration of the regional cerebral glucose metabolism in
502   healthy subjects by glucose loading. *Hum Brain Mapp*. 2016;37(8):2823-2832. doi:10.1002/hbm.23210

503   17.   Tsuchida T, Sadato N, Nishizawa S, Yonekura Y, Itoh H. Effect of postprandial hyperglycaemia in non-
504   invasive measurement of cerebral metabolic rate of glucose in non-diabetic subjects. *Eur J Nucl Med*.
505   2002;29(2):248-250. doi:10.1007/s00259-001-0701-5

506   18.   Nybo L, Moller K, Pedersen BK, Nielsen B, Secher NH. Association between fatigue and failure to
507   preserve cerebral energy turnover during prolonged exercise. *Acta Physiol Scand*. 2003;179(1):67-74.
508   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=
509   12940940

510   19.   Fox 3rd SM, Naughton JP, Haskell WL. Physical activity and the prevention of coronary heart disease.
511   *Ann Clin Res*. 1971;3(6):404-432.

512   20.   Huppert TJ, Diamond SG, Franceschini MA, Boas DA. HomER: a review of time-series analysis methods
513   for near-infrared spectroscopy of the brain. *Appl Opt*. 2009;48(10):D280-D298.
514   doi:10.1364/ao.48.00d280

515   21.   Ainslie PN, Barach A, Murrell C, Hamlin M, Hellemans J, Ogoh S. Alterations in cerebral autoregulation
516   and cerebral blood flow velocity during acute hypoxia: Rest and exercise. *Am J Physiol Heart Circ*
517   *Physiol*. 2007;292(2):976-983. doi:10.1152/ajpheart.00639.2006

518   22.   Ogawa Y, Iwasaki KI, Aoki K, Shibata S, Kato J, Ogawa S. Central hypervolemia with hemodilution
519   impairs dynamic cerebral autoregulation. *Anesth Analg*. 2007;105(5):1389-1396.
520   doi:10.1213/01.ane.0000281910.95740.e4

521   23.   Ogoh S, Brothers M, Barnes Q, et al. The effect of changes in cardiac output on middle cerebral artery
522   mean blood velocity at rest and during exercise. *Journal of Physiology*. 2005;569(2):697-704.
523   doi:10.1113/jphysiol.2005.095836

524   24.   Pott F, Knudsen L, Nowak M, Nielsen HB, Hanel B, Secher NH. Middle cerebral artery blood velocity
525   during rowing. *Acta Physiol Scand*. 1997;160(3):251-255. doi:https://doi.org/10.1046/j.1365-
526   201X.1997.00144.x

527   25.   Cifford PS, Hane B, Secher NH. Arterial blood pressure response to rowing. *Med Sci Sports Exerc*.
528   1994;26(6).

529   26.   Edwards MR, Martin DH, Hughson RL. Cerebral hemodynamics and resistance exercise. *Med Sci Sports*
530   *Exerc*. 2002;34(7).

531   27.   Perry BG, Lucas SJE. The Acute Cardiorespiratory and Cerebrovascular Response to Resistance Exercise.
532   *Sports Med Open*. 2021;7(1). doi:10.1186/s40798-021-00314-w

533   28.   Gandevia SC, Hobbs SF. Cardiovascular responses to static exercise in man: central and reflex
534   contributions. *J Physiol*. 1990;430(1):105-117. doi:https://doi.org/10.1113/jphysiol.1990.sp018284

535   29.   Sejersen C, Fischer M, Mattos JD, Volianitis S, Secher NH. Fluctuations in cardiac stroke volume during
536   rowing. *Scand J Med Sci Sports*. 2021;31(4):790-798. doi:10.1111/sms.13901

537   30.   Pott F, Van Lieshout JJ, Ide K, Madsen P, Secher NH. Middle cerebral artery blood velocity during
538   intense static exercise is dominated by a Valsalva maneuver. *J Appl Physiol*. 2003;94(4):1335-1344.
539   doi:10.1152/japplphysiol.00457.2002

540   31.   Lie SL, Hisdal J, Høiseth LØ. Cerebral blood flow velocity during simultaneous changes in mean arterial
541   pressure and cardiac output in healthy volunteers. *Eur J Appl Physiol*. 2021;121(8):2207-2217.
542   doi:10.1007/s00421-021-04693-6

543   32.   Stebbins CL, Walser B, Jafarzadeh M. Cardiovascular responses to static and dynamic contraction
544   during comparable workloads in humans. *Am J Physiol Regul Integr Comp Physiol*. 2002;283(3 52-
545   3):568-575. doi:10.1152/ajpregu.00160.2002

546   33.   Ogoh S, Dalsgaard MK, Yoshiga CC, et al. Dynamic cerebral autoregulation during exhaustive exercise in
547   humans. *Am J Physiol Heart Circ Physiol*. 2005;288(3 57-3):1461-1467.
548   doi:10.1152/ajpheart.00948.2004

549   34.   Brys M, Brown CM, Marthol H, Franta R, Hilz MJ. Dynamic cerebral autoregulation remains stable
550   during physical challenge in healthy persons. *Am J Physiol Heart Circ Physiol*. 2003;285(3 54-3):1048-
551   1054. doi:10.1152/ajpheart.00062.2003

552   35.   Ogoh S, Dalsgaard MK, Secher NH, Raven PB. Dynamic blood pressure control and middle cerebral
553   artery mean blood velocity variability at rest and during exercise in humans. *Acta Physiologica*.
554   2007;191(1):3-14. doi:10.1111/j.1748-1716.2007.01708.x

555   36.   Cherouveim ED, Margaritelis N V., Koulouvaris P, et al. Skeletal muscle and cerebral oxygenation levels
556   during and after submaximal concentric and eccentric isokinetic exercise. *J Sports Sci*. 2022;40(2):195-
557   202. doi:10.1080/02640414.2021.1983248

558   37.   Ishibashi K, Sakurai K, Shimoji K, Tokumaru AM, Ishii K. Altered functional connectivity of the default
559   mode network by glucose loading in young, healthy participants. *BMC Neurosci*. 2018;19(1):1-8.
560   doi:10.1186/s12868-018-0433-0

561   38.   Shemesh D, Bokobza N, Rozenberg K, Rosenzweig T, Abookasis D. Decreased cerebral blood flow and
562   hemodynamic parameters during acute hyperglycemia in mice model observed by dual-wavelength
563   speckle imaging. *J Biophotonics*. 2019;12(8):1-12. doi:10.1002/jbio.201900002

564   39.   Hayashi N, Kashima H, Ikemura T. Facial Blood Flow Responses to Dynamic Exercise. *Int J Sports Med*.
565   2020;42(03):241-245.

566   40.   Ikemura T, Hayashi N. Ocular circulatory responses to exhaustive exercise in humans. *Eur J Appl*
567   *Physiol*. 2012;112(9):3313-3318. doi:10.1007/s00421-012-2313-0

Conflicts of interests

The authors report no conflict of interests.

Author contributions

P.A.S. and H.M. conceptualised the study, conducted data collection, analysis, and interpretation, and

drafted the original manuscript. W.Z., W.L. and F.R. contributed to conducting the experiments, data

collection, and analysis. X.Z. assisted in the acquisition and analysis of NIRS data. Z.Y. and Z.K. contributed

to the study conception and provided guidance in manuscript writing and revision.

Funding

This study was partly supported by the multiannual grant UID/DTP/00447/2019 by Fundação para a

Ciência e a Tecnologia, Portugal.

Acknowledgements

The authors thank the participants involved in this study.

Table 1: Participants characteristics. Data are presented as mean $\pm$ standard deviation; n = 11.

Age (years)	Height (cm)	Weight (kg)	BMI (kg·m⁻²)	% Body fat (%)	$\dot{V}O_{2peak}$ (ml·kg⁻¹·min⁻¹)	HR_{peak} (beats·min⁻¹)	$\dot{W}_{peak}$ (W)
25 ± 3	174 ± 7	89 ± 11	29 ± 3	30 ± 5	32 ± 2	172 ± 4	132 ± 12

585   BMI, body mass index; $\dot{V}O_{2peak}$, peak value for the rate of oxygen consumption during the incremental maximal exercise test; HR_{peak}, peak value for heart rate

586   during the incremental maximal exercise test; $\dot{W}_{peak}$, peak value for power output during the incremental maximal exercise test.

588   Table 2: Mean ± SD values for oxyhaemoglobin, deoxyhaemoglobin, and total haemoglobin concentration changes during the repeated sprints; n = 10.

Sprint repetition	Trial	$\Delta O_2Hb_{0-15\ sec}$ µM	$\Delta O_2Hb_{15-25\ sec}$ µM	$\Delta HHb_{0-15\ sec}$ µM	$\Delta HHb_{15-25\ sec}$ µM	$\Delta tHb_{0-15\ sec}$ µM	$\Delta tHb_{15-25\ sec}$ µM
1st	Placebo	1.81 ± 1.69	-0.18 ± 1.82	0.02 ± 0.39	-0.25 ± 0.66	1.83 ± 1.84	-0.44 ± 2.06
	Glucose	1.68 ± 1.00	0.30 ± 2.49	-0.11 ± 0.39	-0.35 ± 0.52	1.51 ± 1.96	-0.03 ± 2.42
	Mean ± SD	1.75 ± 1.84	0.06 ± 2.19	-0.04 ± 0.40	-0.30 ± 0.59	1.67 ± 1.90	-0.24 ± 2.25
2nd	Placebo	2.75 ± 1.33	-0.61 ± 1.56	0.25 ± 0.54	0.01 ± 0.49	3.01 ± 1.51	-0.62 ± 1.36
	Glucose	1.99 ± 1.82	-1.04 ± 1.42	0.16 ± 0.33	0.14 ± 0.43	2.16 ± 1.20	-0.88 ± 1.47
	Mean ± SD	2.37 ± 1.64	-0.82 ± 1.31	0.21 ± 0.45	0.08 ± 0.46	**2.59 ± 1.81**[a, b]	-0.75 ± 1.42
3rd	Placebo	2.61 ± 1.52	-0.38 ± 1.40	-0.01 ± 1.18	-0.22 ± 1.25	2.67 ± 1.61	-0.59 ± 2.01
	Glucose	2.24 ± 1.71	-0.55 ± 0.99	0.02 ± 0.29	-0.07 ± 0.43	2.30 ± 1.76	-0.60 ± 1.08
	Mean ± SD	2.42 ± 1.62	-0.47 ± 1.21	0.00 ± 0.85	-0.15 ± 0.94	2.48 ± 1.69	-0.60 ± 1.61
4th	Placebo	3.05 ± 1.71	-0.62 ± 1.43	-0.04 ± 0.82	-0.41 ± 0.93	3.11 ± 2.03	-1.04 ± 2.14
	Glucose	2.67 ± 1.39	-0.51 ± 1.26	0.01 ± 0.48	-0.16 ± 0.46	2.77 ± 1.57	-0.68 ± 1.45

	Mean ± SD	**2.86 ± 1.56**[a]	-0.57 ± 1.34	-0.02 ± 0.67	-0.28 ± 0.74	**2.94 ± 1.82**[a]	-0.86 ± 1.83
5th	Placebo	3.08 ± 1.95	-0.59 ± 0.67	0.10 ± 0.66	-0.16 ± 0.43	3.20 ± 2.22	-0.72 ± 0.82
	Glucose	3.17 ± 1.70	-0.24 ± 1.26	-0.05 ± 0.52	-0.19 ± 0.81	3.16 ± 1.60	-0.41 ± 1.79
	Mean ± SD	**3.12 ± 1.83**[a]	-0.42 ± 1.02	0.02 ± 0.60	-0.18 ± 0.64	**3.18 ± 1.93**[a]	-0.57 ± 1.49
6th	Placebo	3.18 ± 1.66	-0.17 ± 1.08	0.25 ± 0.73	0.00 ± 0.83	3.44 ± 2.00	-0.17 ± 1.70
	Glucose	2.88 ± 1.97	-0.59 ± 1.11	-0.04 ± 0.51	-0.12 ± 0.52	2.93 ± 2.11	-0.73 ± 1.39
	Mean ± SD	**3.03 ± 1.83**[a]	-0.38 ± 1.11	0.10 ± 0.64	-0.06 ± 0.69	**3.19 ± 2.07**[a]	-0.45 ± 1.58
7th	Placebo	2.75 ± 1.52	-0.47 ± 1.25	0.15 ± 0.77	-0.28 ± 0.70	2.94 ± 1.87	-0.74 ± 1.45
	Glucose	2.37 ± 1.25	-0.68 ± 1.02	0.07 ± 0.41	-0.12 ± 0.36	2.49 ± 1.49	-0.80 ± 1.16
	Mean ± SD	**2.56 ± 1.34**[a]	-0.57 ± 1.14	0.11 ± 0.62	-0.20 ± 0.56	**2.71 ± 1.70**[a]	-0.77 ± 1.31
8th	Placebo	3.03 ± 2.78	-0.60 ± 1.49	0.40 ± 1.02	-0.07 ± 0.92	3.47 ± 3.27	-0.66 ± 1.24
	Glucose	3.15 ± 2.34	-0.07 ± 1.10	0.09 ± 0.44	-0.15 ± 0.52	3.26 ± 2.45	-0.21 ± 0.93
	Mean ± SD	**3.09 ± 2.56**[a]	-0.33 ± 1.33	0.25 ± 0.80	-0.11 ± 0.75	**3.36 ± 2.88**[a]	-0.44 ± 1.12

		$\Delta O_2Hb_{0-15\,sec}$	$\Delta O_2Hb_{15-25\,sec}$	$\Delta HHb_{0-15\,sec}$	$\Delta HHb_{15-25\,sec}$	$\Delta tHb_{0-15\,sec}$	$\Delta tHb_{15-25\,sec}$
9th	Placebo	3.23 ± 2.70	-0.81 ± 1.33	0.02 ± 1.08	-0.54 ± 1.14	3.28 ± 3.15	-1.37 ± 2.11
	Glucose	2.72 ± 1.42	-0.26 ± 1.01	-0.04 ± 0.45	-0.35 ± 0.57	2.69 ± 1.55	-0.59 ± 1.30
	Mean $\pm$ SD	$\mathbf{2.98 \pm 2.17}$[a]	-0.54 ± 1.21	-0.01 ± 0.82	-0.44 ± 0.90	$\mathbf{2.98 \pm 2.49}$[a]	-0.98 ± 1.79
10th	Placebo	3.40 ± 3.01	-0.36 ± 1.46	0.10 ± 0.82	-0.29 ± 0.68	3.57 ± 3.56	-0.60 ± 1.70
	Glucose	3.50 ± 1.80	-0.35 ± 1.44	-0.22 ± 0.92	-0.58 ± 1.77	3.35 ± 1.78	-0.92 ± 2.22
	Mean $\pm$ SD	$\mathbf{3.45 \pm 2.47}$[a]	-0.35 ± 1.44	-0.06 ± 0.88	-0.43 ± 1.35	$\mathbf{3.46 \pm 2.81}$[a]	-0.76 ± 1.98
Overall mean $\pm$ SD	Placebo	2.89 ± 2.10	-0.48 ± 1.35	0.12 ± 0.84	-0.22 ± 0.85	3.05 ± 2.44	-0.70 ± 1.73
	Glucose	2.64 ± 1.84	-0.40 ± 1.41	-0.01 ± 0.51	-0.19 ± 0.77	2.66 ± 1.92	-0.59 ± 1.60

$\Delta O_2Hb_{0-15\,sec}$, median change in oxyhaemoglobin concentration in the time interval $0 - 15$ s (sprint); $\Delta O_2Hb_{15-25\,sec}$, median change in oxyhaemoglobin concentration in the time interval $15 - 25$ s (recovery); $\Delta HHb_{0-15\,sec}$, median change in deoxyhaemoglobin concentration in the time interval $0 - 15$ s (sprint); $\Delta HHb_{15-25\,sec}$, median change in deoxyhaemoglobin concentration in the time interval $15 - 25$ s (recovery); $\Delta tHb_{0-15\,sec}$, median change in total haemoglobin concentration in the time interval $0 - 15$ s (sprint); $\Delta tHb_{15-25\,sec}$, median change in total haemoglobin concentration in the time interval $15 - 25$ s (recovery). Positive and negative values indicate increase and decrease in haemoglobin concentrations, respectively, from the time immediately before the onset of the respective sprint repetition (baseline value = 0). Mean $\pm$ SD is the average and standard deviation of values from GLU and PLA trials within each sprint. Overall mean $\pm$ SD is the average and

standard deviation for all values in the respective trial. [a] Significantly different from the first sprint, $p < 0.05$; [b] significantly different from the immediately

preceding sprint, $p < 0.05$.

Figure 1

Figure 1: Mean power output during each repeated sprint for GLU (red line) and PLA (blue line) trials (left
panel). Pooled mean power output for GLU (red bar) and PLA (blue bar) trials (right panel). Data
presented as mean ± SD, ns indicates non-significant difference, sample size: n = 10.

Figure 2

Figure 2: Heart rate, stroke volume and cardiac output (mean ± SD) at rest and during each repeated

sprint for GLU (red lines) and PLA (blue lines) trials. The insert is the output of the two-way repeated

measures ANOVA for the significant main effect of sprint repetition (excluding the resting values). P-

values are from backward difference contrasts, comparing each sprint repetition with the immediately

preceding one, including the first sprint and resting values. Sample size: n = 10.

Figure 3

Figure 3: Changes in O₂Hb, HHb and tHb during the repeated sprints exercise. Top and middle graphs display the concentration changes of O₂Hb (red line), HHb (blue line) and tHb (green line) recorded in two participants. The records start immediately before the warm-up. In both participants, a transient increase in O₂Hb, HHb, and tHb during each sprint (black dots) is clearly observed. In the top graph, there was a notable decrease in O₂Hb concentration and increase in HHb throughout the repeated sprints exercise, followed by a return of both O₂Hb and HHb towards pre-exercise values during the recovery period. The participant in the middle graph exhibited lesser changes in O₂Hb and HHb concentration during exercise but displayed substantial post-exercise cerebral hyperaemia. The bottom panel presents the overall average for ΔO₂Hb and ΔHHb during the repeated sprints, revealing the rapid increase in cerebral haemoglobin concentration during the sprint and its quick decrease soon after the end of the sprint. Each tracing is the mean of 10 participants, 10 sprints and 8 channels. The shadowed areas represent the standard error of the mean; sample size: n = 10.

Figure 4

Figure 4: Scatter plots illustrating the relationship between ΔO_2Hb (upper left graph), ΔHHb (upper right

graph), and ΔtHb (lower left graph) with cardiac output, as well as between ΔO_2Hb and power output

(lower right graph), during the sprints (solid lines) and the time interval between sprints (dotted lines) in

GLU (red lines and red circles) and PLA (blue lines and blue triangles) trials. The Pearson's R coefficients

and respective p-values are presented at the top of the graphs for values during the sprint (left column)

and the time interval between the sprints (right column) for GLU (red font) and PLA (blue font) trials.

During the sprint, a significant correlation was found between cardiac output and ΔO_2Hb, as well as ΔtHb.

No significant correlation was found between cardiac output and ΔHHb, and between cardiac output and

ΔO_2Hb or ΔtHb during the time interval between sprints. In the PLA trial, mean power output exhibited a

negative correlation with ΔO_2Hb during the sprints, while in the GLU trial, mean power output showed a

negative correlation with ΔO_2Hb during the time interval between sprints.