Vitamin A-treated NK cells reduce IFN-γ production and support regulatory T cell differentiation

Running Title:

Vitamin A-induced NK cell immune regulation

Authors:

Mingeum Jeong[1], Jia-Xiang See[1,2], Carolina De La Torre[3], Adelheid Cerwenka[1, 2] and Ana Stojanovic[1]

Affiliation:

[1]Department of Immunobiochemistry, Mannheim Institute of Innate Immunosciences (MI3), Medical Faculty Mannheim, Heidelberg University, Mannheim, 68167, Germany.

[2]European Center for Angioscience (ECAS), Medical Faculty Mannheim, Heidelberg University, Mannheim, 68167, Germany.

[3]NGS Core Facility, Medical Faculty Mannheim, Heidelberg University, Mannheim, 68167, Germany

Corresponding Author:

Ana Stojanovic

Contact information:

Department of Immunobiochemistry, Mannheim Institute for Innate Immunoscience (MI3), Medical Faculty Mannheim, Heidelberg University, Ludolf Krehl-Straße 13-17, D-68167 Mannheim, Germany

Telephone: +49 (0)621/383-71502

E-Mail: Ana.Stojanovic@medma.uni-heidelberg.de

The final character count:

Abstract

NK cells are innate cytotoxic lymphocytes that contribute to immune responses against stressed, transformed or infected cells. The effector functions of NK cells are regulated by microenvironmental factors, including cytokines, metabolites and nutrients. Vitamin A is an essential micronutrient that plays an indispensable role in embryogenesis and development, but was also reported to regulate immune responses. However, the role of vitamin A in regulating NK cell functions remains poorly understood. Here, we show that the most prevalent vitamin A metabolite, all-*trans* retinoic acid (atRA), induces transcriptional and functional changes in NK cells leading to altered metabolism and reduced IFN-γ production in response to a wide range of stimuli. atRA-exposed NK cells displayed a reduced ability to support dendritic cell (DC) maturation and were inefficient in eliminating immature DCs. Moreover, they supported the polarization and proliferation of regulatory T cells. These results imply that in vitamin A-enriched environments, NK cells can acquire regulatory-like functions and might support tolerogenic immunity and/or immunosuppression.

Keywords: all-*trans* retinoic acid, NK cells, NK-DC cross-talk, IFN-γ, regulatory T cells, Vitamin A

Introduction

Natural killer (NK) cells are innate lymphocytes that play important roles in anti-tumor and anti-viral immunity (Hamerman, Ogasawara, & Lanier, 2005; Vivier, Tomasello, Baratin, Walzer, & Ugolini, 2008). The balance of signals delivered by the activating and inhibitory receptors expressed on the cell surface determines NK cell activation (Pegram, Andrews, Smyth, Darcy, & Kershaw, 2011). NK cells can eliminate target cells via activating receptor-triggered secretion of cytotoxic granules or by inducing apoptosis through the engagement of death receptors expressed by target cells (Smyth et al., 2005). Furthermore, soluble factors, such as cytokines, can modulate NK cell functions (Marcais et al., 2013). For example, cytokines IL-12 and IL-18 synergistically trigger NK cell activation and production of pro-inflammatory cytokine IFN-γ (Okamura, Kashiwamura, Tsutsui, Yoshimoto, & Nakanishi, 1998).

NK cells contribute to regulation of adaptive immunity through direct interaction with T cells, as well as by regulating the activation and function of dendritic cells (DCs) (Walzer, Dalod, Robbins, Zitvogel, & Vivier, 2005). IL-12 and IL-18 released by DCs, or the engagement of the activating receptor NKp30 by its ligand(s) expressed by DCs, were shown to enhance NK cell cytotoxicity and their cytokine production, including IFN-γ and TNF-α (Borg et al., 2004; Ferlazzo et al., 2002; Piccioli, Sbrana, Melandri, & Valiante, 2002; Yu et al., 2001). Reciprocally, IFN-γ or TNF-α secreted by NK cells induced the maturation of DCs, characterized by elevated expression of MHC class I and co-stimulatory molecules (Piccioli et al., 2002; Vitale et al., 2005). At a high NK/DC ratio, NK cells were shown to kill immature DCs (Ferlazzo et al., 2002; Piccioli et al., 2002). During T cell priming, NK cells are recruited to lymph nodes and support the differentiation of T helper type 1 (Th1) cells by providing IFN-γ (Martin-Fontecha et al., 2004; Morandi, Bougras, Muller, Ferlazzo, & Munz, 2006). On the contrary, NK cells can also restrict the responses of activated T cells via NKG2D-mediated cytotoxicity (Cerboni et al., 2007; Rabinovich et al., 2003).

Microenvironmental factors, such as micronutrients or oxygen tension, can modulate activation and effector functions of NK cells (Littwitz-Salomon et al., 2021; Ni et al., 2020). Vitamin A is an essential fat-soluble micronutrient that is digested and absorbed in the small intestine, and stored mainly in the liver (Blomhoff, Green, Berg, & Norum, 1990). NK cells can be recruited to these tissues upon inflammation, to perform cytotoxicity against abnormal cells and to regulate immune responses (Colonna, 2009; Papaioannou et al., 2022; Tian, Chen, & Gao, 2013). Therefore, NK cells can be exposed to vitamin A-enriched microenvironment. Previous studies have demonstrated that animals fed with vitamin A-deficient diet showed a decreased number of NK cells in blood, while splenic NK cells displayed impaired effector functions, which could be restored by the oral administration of retinoic acid (Fraker, Halter, & Forbes, 1986; Zhao & Ross, 1995). Contrarily, the cytotoxicity of the human NK92 cell line was suppressed by *in vitro* treatment with atRA (A. Li et al., 2007). Thus, the exact roles of vitamin A and its metabolites in regulating NK cell phenotype and functions remains incomplete.

Here, we show that NK cells treated with the most prevalent vitamin A metabolite, all-*trans* retinoic acid (atRA), acquire regulatory-like functions. Compared to controls, atRA-treated NK cells produce less IFN-γ, favor an immature phenotype of DCs, and support the differentiation of regulatory T (Treg) and regulatory Th17 cells, as well as their proliferation. Our data provide a novel insight in NK cell regulation by vitamin A, which might affect not only tissue homeostasis and immune tolerance, but also support immunosuppression during an active immune response.

Results

atRA induces transcriptional and phenotypic changes in NK cells

To address the effect of vitamin A, mouse splenic NK cells exposed for 7 days to the active form of vitamin A metabolite, atRA, or the equivalent volume of solvent, DMSO, were subjected to transcriptome analysis. atRA-treated NK displayed an altered transcriptome, with over 400 differentially regulated transcripts (Fig 1A). Expression of genes involved in retinoic acid and lipid metabolism (Dhr3, Dgat1, and Pparg) was upregulated, while the expression of genes involved in NK cell activation and effector functions (Ifng, Il18r1, Tnf, Klrb1c, and Ncr1) was downregulated upon the exposure to atRA (Fig 1A-B). Accordingly, pathway analysis revealed enriched retinol metabolism in atRA-treated NK cells compared to control NK cells, while the nuclear factor kappa B (NFκB) pathway and NK cell-mediated cytotoxicity were downregulated (Fig 1C).

Next, we analyzed protein expression of selected molecules whose transcripts were differentially expressed in atRA-treated NK cells (Fig 1A-B), including the transcription factor Eomes, activating receptors NK1.1, NKp46, NKG2D, and DNAM-1, death-receptor ligand TRAIL, and nucleotide-metabolizing ectoenzymes CD73, CD39, and CD38. The expression of Eomes was decreased in atRA-treated NK cells compared to controls, as well as the surface expression of NK1.1 and NKG2D, while the expression of NKp46 remained unchanged (Fig 1D). atRA-treated NK cells comprised higher frequencies of DNAM-1- and TRAIL-expressing cells (Fig 1E). Similarly, the expression of CD73, CD39, and CD38 increased over time during culture of NK cells with atRA compared to control treatment (Fig 1F and Fig EV1). As retinoic acid was previously reported to induce the expression of gut-homing receptors, such as CCR9 and α4β7, in T cells and group 3 innate lymphoid cells (ILC3s) (Iwata et al., 2004; Kim, Taparowsky, & Kim, 2015), we examined the expression of adhesion molecules and homing-receptors on atRA-treated NK cells. Among atRA-treated NK cells, CXCR3-expressing cells were enriched, the percentages of CD62L- and CCR9-expressing cells were lower, while the frequencies of CXCR6-expressing cells remained comparable to control-treated NK cells (Fig EV 2). Together, these data demonstrate that the exposure to atRA induces comprehensive transcriptional and phenotypic alterations in NK cells.

atRA-treated NK cells display altered metabolism

Mitochondria are organelles that contribute to cellular metabolism by oxidizing pyruvate, fatty acids, and amino acids to generate energy (Spinelli & Haigis, 2018). As atRA-treated NK cells displayed changes in the expression of genes that are involved in several metabolic pathways (Fig 1C), we assessed the features of mitochondria in NK cells upon treatment with atRA. Compared to controls, atRA-treated NK cells displayed a reduced mitochondrial mass and membrane potential, as well as reduced production of reactive oxygen species (ROS) (Fig 2A). Accordingly, mitochondrial respiration and glycolysis, measured via the oxygen consumption rate (OCR) and the extracellular acidification rate (ECAR), respectively, were altered by exposure to atRA; atRA-treated NK cells showed reduced basal and maximal OCR, and reduced ECAR (Fig 2B-C). In parallel, atRA-treated NK cells were smaller (Fig 2D), while the cell expansion in culture (Fig EV 3A) and the uptake of fatty acid, measured by the ability to uptake fluorescently labelled palmitic acid (Fig EV 3B), remained unchanged compared to controls. These results indicate that the exposure to atRA induced reduction in mitochondrial mass and function, as well as altered metabolism of NK cells.

atRA attenuates IFN-γ production of NK cells

To explore the effect of atRA on NK cell effector function, we assessed the cytokine and chemokine expression by atRA-treated and control NK cells. atRA-treated NK cells displayed reduced transcript amounts of Ifng, while Csf2, Ccl5, and Ccl1 expression was elevated compared to controls (Fig 3A). Therefore, we first analyzed the ability of atRA-treated NK cells to produce IFN-γ in response to various stimuli. When stimulated with IL-12 and IL-18, which synergistically induce IFN-γ release by NK cells (Piccioli et al., 2002), the cytokine amount and percentage of IFN-γ-producing atRA-treated NK cells was lower compared to controls (Fig 3B). In addition to pro-inflammatory cytokines, the activation of NK cells can be triggered by the engagement of activating receptors (Lanier, 2001). When activated by cross-linking of the activating receptors NK1.1, NKG2D, NKp46, or DNAM-1, atRA-treated NK cells produced less IFN-γ compared to controls (Fig 3C). Similar results were obtained when NK cells were incubated with tumor cells, RMA-S and YAC-1 lymphoma (Fig 3D). These results imply that the exposure to atRA downregulates IFN-γ production by NK cells in response to various stimuli.

atRA-treated NK cells comprised less IL-18Rα-expressing cells, compared to controls (Fig 4A), which correlated with reduced IFN-γ production in response to IL-18, but not to IL-12 (Fig 4B). To further investigate the mechanism of reduced IL-18-induced IFN-γ production, we analyzed the phosphorylation of proteins involved in the NFκB pathway, which mediates the responses downstream of IL-18R (Rex et al., 2020), and was transcriptionally affected in NK cells treated by atRA (Fig 1C). IL-12 and IL-18 were reported to support the production of IFN-γ via the STAT4 signaling pathway (Nakahira et al., 2002), and NFκB target gene encoding inhibitor of kappa B-zeta (IκBζ) (Miyake et al., 2010), respectively. NK cells increased the expression of IκBζ and NFκB p50 upon stimulation with IL-18, but not with IL-12 (Fig 4C), and the treatment with atRA reduced the IL-18-induced expression of IκBζ and NFκB p50 in NK cells. On the contrary, the phosphorylation of STAT4, which was induced by IL-12, but not IL-18, was not altered by pre-exposure of NK cells to atRA (Fig EV 4A). Similarly, the phosphorylation of NFκB p65 and IκB kinase alpha/beta (IKK α/β) upon stimulation with IL-12 or IL-18 was unchanged compared to control NK cells (Fig EV 4B). Furthermore, we investigated the activation of mTORC1, which drives glycolysis and lipid synthesis by regulating the surface expression of nutrient transporters (Marcais et al., 2014), using the phosphorylation of ribosomal protein S6 as a read-out, and the expression of cMyc, which is required for cytokine-induced metabolism and effector functions in NK cells (Loftus et al., 2018). Phosphorylation of S6 was induced with either IL-12 or IL-18, while cMyc expression increased in NK cells upon exposure to IL-18 (Fig 4D). However, atRA-exposed NK cells failed to activate mTORC1 and to increase cMyc expression in response to IL-18 (Fig 4D). Altogether, atRA-treated NK cells produced less IFN-γ compared to control NK cells in response to IL-18, but not IL-12, and this correlated with reduced activation of NFκB-IκBζ and the mTORC1 pathway, and the reduced expression of cMyc.

atRA-treated NK cells favor immature phenotype of DCs

It was previously reported that NK cells release IFN-γ while interacting with DCs, and promote DC maturation and the expression of antigen presenting and co-stimulatory molecules on DCs (Piccioli et al., 2002; Vitale et al., 2005). Reciprocally, DCs were shown to produce IL-12 and IL-18 during the interaction with NK cells (Piccioli et al., 2002; Semino, Angelini, Poggi, & Rubartelli, 2005). To address if atRA-treated NK cells regulate DC phenotype, we co-cultured atRA-treated or control NK cells with bone marrow-derived dendritic cells (BM-DCs) (Fig 5A), and analyzed maturation and survival of BM-DCs. In accordance to previous literature (Piccioli et al., 2002; Vitale et al., 2005), DCs displayed increased expression of CD86 and MHC II upon co-culture with control NK cells, while atRA-treated

NK cells showed a reduced ability to upregulate the expression of these molecules on DCs (Fig 5B). Accordingly, we detected lower amounts of IFN-γ (Fig 5C) and lower percentages of apoptotic DCs (Fig 5D) in the co-culture of BM-DCs with atRA-treated NK cells. These results indicate that NK cells exposed to atRA favor immature DC phenotype, and this correlates with a lower ability of NK cells to secrete IFN-γ in response to DCs.

atRA-treated NK cells support regulatory T cell differentiation

NK cells and NK cell-derived IFN-γ were also reported to support the differentiation of Th1 cells (Laouar, Sutterwala, Gorelik, & Flavell, 2005; Martin-Fontecha et al., 2004). To determine whether atRA affects the ability of NK cells to regulate T cell differentiation, we co-cultured atRA-treated or control NK cells with naïve CD4[+] T cells in Th1-, Th17-, or Treg-polarizing conditions. In Th1-polarizing conditions, the expression of the transcription factor T-bet and the cytokine IFN-γ by CD4[+] T cells was not affected by the NK cells, irrespective of their pre-treatment (Fig EV 5A). However, control NK cells reduced the ability of naive T cells to differentiate into FoxP3-expressing Tregs, while the frequency of differentiated Tregs was not affected by atRA-treated NK cells (Fig 6A). In Th17-polarizing conditions, the majority of CD4[+] T cells acquired the expression of the transcription factor Rorγt, which was not affected by NK cells (Fig 6B). Among Rorγt-expressing CD4[+] T cells, a subpopulation of cells co-expressed the transcription factor FoxP3, and the percentage of FoxP3-expressing CD4[+] Th17 cells increased in the presence of atRA-treated NK cells, compared to both control NK cells and to T cells cultured alone (Fig 6B).

To address the mechanism by which atRA-treated NK cells supported the differentiation of FoxP3-expressing Th17 cells, we first prevented a direct NK cell-to-T cell contact using transwell-chamber system. In this setup, FoxP3 expression by CD4[+] T cells cultured with atRA-treated NK cells was not affected (Fig 6C), indicating that soluble molecules supported the differentiation of FoxP3-expressing CD4[+] cells. Among these, IL-2-receptor signaling via Signal transducer and activator of transcription 5 (STAT5) was shown to be required for the development of FoxP3-expressing Treg cells (Chinen et al., 2016; Vang et al., 2008), and adenosine, IL-10, CCL1, and CCL5 were previously reported to support Treg differentiation (Hoelzinger et al., 2010; Kuehnemuth et al., 2018; Morita et al., 2019; Ohta & Sitkovsky, 2014). Since our transcriptomic and flow-cytometric data showed that the expression of adenosine-metabolizing enzymes (CD73, CD39, and CD38; Fig 1F), and the expression of Ccl1 and Ccl5 transcripts (Fig 3A) were increased in NK cells exposed to atRA, we analyzed whether these molecules regulate regulatory T cell differentiation in our NK-Th17 co-culture system. Antagonist-mediated blockade of the adenosine A2A receptor (A2AR), or the mAb-mediated neutralization of IL-10, CCL1, or CCL5, did not affect FoxP3 expression by CD4[+] T cells upon co-culture with atRA-treated or control NK cells (Fig EV 5B-D). However, although the percentage of CD25 (IL-2Rα)-expressing T cells was comparable (Fig 6D), the frequency of phosphorylated-STAT5[+]-expressing T cells was higher in co-cultures with atRA-treated compared to control NK cells (Fig 6E). This correlated with a higher percentage of Ki67[+] FoxP3-expressing CD4[+] T cells in the co-cultures with atRA-treated NK cells (Fig 6F and Fig EV 5E), suggesting that NK cells exposed to atRA could support the proliferation of FoxP3-expressing CD4[+] T cells via the phosphorylation of STAT5 downstream of IL-2 receptor engagement.

Discussion

Vitamin A is an essential lipid-soluble nutrient, which promotes tolerogenic immunity in the gastrointestinal tract (Hall, Grainger, Spencer, & Belkaid, 2011). Primary role of vitamin A in gut is to

create microenvironment in which immune cell functions are skewed towards promoting immune regulation. For, example, retinoic acid (RA)-imprinted DCs were shown to support the Treg differentiation and their IL-10 production (Bakdash, Vogelpoel, van Capel, Kapsenberg, & de Jong, 2015; Sun et al., 2007), as well as the expression of gut-homing receptors on B cells and Tregs (Bakdash et al., 2015; Mora et al., 2006). Similarly, group 3 innate lymphoid cells (ILC3s) were reported to increase the expression of gut-homing receptors, CCR9 and α4β7, upon the exposure to RA (Eksteen et al., 2009; Iwata et al., 2004; Zeng et al., 2013). In mice fed with vitamin A-deficient diet, the ratio of IL-22-producing ILC3s to ILC2s declined, resulting in an increased host susceptibility to bacterial infection (Spencer et al., 2014). Our data show that atRA does not increase CCR9 and CD62L expression in NK cells, indicating that vitamin A might not imprint a "classical" lymph node- and/or gut-homing phenotype in NK cells, and might rather regulate NK cell phenotype in vitamin A-enriched microenvironments other than gut. Besides gut, the organ with highest amounts of stored vitamin A and its derivatives is the liver (Kane, Folias, & Napoli, 2008). In the steady-state, the liver tissue operates under tolerogenic immunoregulatory environment. It was reported that T cells primed in the liver predominantly differentiate towards a regulatory phenotype, in a process that is orchestrated by local antigen presentation (Bottcher et al., 2014), inhibitory checkpoints molecules (Diehl et al., 2008) and/or microenvironmental cytokines (Osei-Bordom, Bozward, & Oo, 2020). In the gut mucosa, Treg and Th17 cells are enriched to restrain inflammatory responses and protect the host from infection, respectively. Despite their different functions, they share similar differentiation modules, tightly regulated by microenvironmental factors, such as IL-6, TGF-β, IL-1β, IL-21, short-chain fatty acids and RA (Y. K. Lee, Mukasa, Hatton, & Weaver, 2009; Pandiyan et al., 2019; Ueno et al., 2018). It was reported that the outcome of naive T cell differentiation towards Treg or Th17 effector cell depends on relative concentrations of key cytokines regulating these cell fates (M. O. Li, Sanjabi, & Flavell, 2006; Veldhoen, Hocking, Atkins, Locksley, & Stockinger, 2006). For example, RA supports Treg differentiation by enhancing TGF-β over IL-6 signaling (Xiao et al., 2008). Th17 cells appears to be particularly plastic, showing high adaptability to local cytokine milieus. They can acquire both IFN-γ production and Th1-like phenotype, or to co-express FoxP3 favoring IL-10 secretion and immune regulation (Duhen et al., 2013; Zielinski et al., 2012). However, it is not clear how FoxP3$^+$ cells develop within the Th17 differentiation program. Gagliani et al. also reported the existence of intestinal "ex-Th17 cells" under homeostasis, that irreversibly lost the ability to produce IL-17, and acquired Tr1-like phenotype (Gagliani et al., 2015). Although key cytokine players that shape the Th17-Treg plasticity and trans-differentiation have been identified, the contribution of other environmental factors, including interactions with neighboring cells, especially in tissues other than gut, is still elusive. Here, we show that in both Th17- and Treg-polarizing conditions, atRA-treated NK cells supported the differentiation and/or proliferation of FoxP3-expresing CD4$^+$ T cells. This effect was mediated by soluble factors, and correlated with increased phosphorylation of STAT5 in T cells. STAT5 is a transcription factor activated upon cytokine-receptor engagement, including receptors for IL-3, IL-4, IL-5, GM-CSF and cytokines signaling via the common γc chain (Lischke et al., 1998; Mui, Wakao, O'Farrell, Harada, & Miyajima, 1995). As STAT5 signaling and IL-2 were reported to be essential for the development of Tregs (Chinen et al., 2016; Vang et al., 2008), and suppression of Th17 program (Laurence et al., 2007), IL-2-IL-2R might drive atRA-mediated effects in co-culture of NK cells with T cells. In addition, our findings also indicate the potential of atRA-treated NK cells to produce adenosine, due to co-expression of the adenosine-generating enzymes CD73, CD39 and CD38, at both transcriptional and protein level. Adenosine is immunosuppressive mediator derived from extracellular ATP that accumulates in pathological conditions, such as in hypoxic solid tumor tissue (Ohta, 2016). Stimulation of adenosine receptor A2AR in naive CD4$^+$ T cells was shown to promote Treg development and FoxP3 expression (Zarek et al., 2008). Although blocking the adenosine receptor did not indicate its role in our co-culture system, this axis might support pleiotropic immunoregulatory functions of NK cells in vitamin A-

enriched tissues. In the liver, where NK cells comprise the majority of lymphocytes (Gao, Jeong, & Tian, 2008), the adenosine pathway plays an important roles in regulating inflammation and immune responses in liver disease (W. Wang et al., 2021; Zimmerman et al., 2013). Liver tissue injury leads to concomitant release of intracellular ATP from damaged and necrotic cells and to the depletion of vitamin A storage from hepatic stellate cells (HSCs) (Blomhoff et al., 1990). Released retinol can be converted to its active form(s) by HSCs and by neighboring stromal and parenchymal cells (Friedman, 2008), thus becoming available for ongoing tissue responses, while ATP is being converted to adenosine. How these microenvironmental milieus regulate NK cell responses has still not been addressed.

Our results show that exposure to atRA primarily reduces NK cell pro-inflammatory responses, such as IFN-γ production. IFN-γ is an important effector cytokine in both innate and adaptive immunity, which can be produced by both NK cells and activated T cells (Handa, Suzuki, Matsui, Shimizu, & Kumagai, 1983; Kasahara, Hooks, Dougherty, & Oppenheim, 1983). atRA was shown to downregulate the production of IFN-γ by CD8$^+$ T cells (Van et al., 2009), as well as by NKT cells in mice with concanavalin A -induced hepatitis (K. A. Lee et al., 2012). An important role of IFN-γ during early immune response activation is to promote differentiation and maturation of monocytes and DCs (Ardavin et al., 2001; Young & Hardy, 1995), and NK cell-derived IFN-γ was shown to support the expression of co-stimulatory molecules and antigen-presenting machinery in maturing DCs (Piccioli et al., 2002; Vitale et al., 2005). At the same time, by eliminating immature DCs, NK cells can assure the generation of an activated mature DC pool to drive efficient T cell priming. We show that atRA-treated NK cells secreted less IFN-γ during co-culture with DCs, which correlated with lower expression of co-stimulatory molecules, and higher survival of immature DCs. Several receptor-ligand interaction, as well as soluble factors were reported to contribute to NK-DC cross-talk resulting in maturation and/or cytotoxicity (Ferlazzo, Semino, & Melioli, 2001; Ferlazzo et al., 2002; Hayakawa et al., 2004; Piccioli et al., 2002). atRA-mediated reduction of IFN-γ production by NK cells seems to comprise a rather general mechanism that operates downstream of various stimuli, including cytokines, activating receptor triggering or encounter of tumor target cells. The underlying mechanism targets two signaling axes that regulate NK cell effector functions and cellular fitness. First, our results show that NK cells exposed to atRA were unable to efficiently engage mTORC1 and increase cMyc expression in response to IL-18. mTORC1 and cMyc are metabolic regulators that increase glycolysis rate and support pro-inflammatory effector responses in T cells (Powell, Lerner, & Schwartz, 1999; R. Wang et al., 2011; Zheng et al., 2007). NK cells require mTORC1 for development, as well as for the IL-15-induced IFN-γ production by elevating glycolytic rate (Marcais et al., 2014). cMyc protein expression in NK cells was shown to be regulated by glutamine availability and SLC7A5-dependent amino acid transport (Loftus et al., 2018). cMyc-deficient NK cells expanded in presence of IL-15, failed to undergo blastogenesis, to increase mitochondrial mass and metabolic flux, and produced less IFN-γ in response to IL-2 and IL-12 (Loftus et al., 2018). Similarly, atRA-exposed NK cells displayed reduced rates of both glycolysis and oxidative phosphorylation compared to controls, reduced cell size, and reduced mitochondrial mass and membrane potential. Although previous studies reported the impact of atRA on mitochondria and cellular metabolism that appears to be cell- and organ-specific (Chidipi et al., 2021; Rigobello et al., 1999; Tourniaire et al., 2015), it remains to be investigated if atRA mediates these effects through regulation of cMyc.

In addition to mTORC1 and cMyc, the attenuated IFN-γ production by atRA-treated NK cells correlated with lower expression of genes involved in the NFκB pathway, as well as lower expression of NFκB p50 in response to stimulation of the IL-18 receptor. Stimulation with IL-12 was not affected by atRA, indicating that atRA modulated preferentially NFκB-dependent responses. The NFκB signaling cascade operates downstream of several other receptors expressed by NK cells, including NKG2D, DNAM-1,

CD16 or Ly49H (Kwon et al., 2016), which might explain the reduced IFN-γ production when atRA-exposed NK cells were triggered via these receptors, or with tumor cells previously reported to express their cognate ligands (Chan et al., 2010; Diefenbach, Jensen, Jamieson, & Raulet, 2001; Oberg et al., 2004). NFκB was reported to directly bind to the *Ifng* promoter (Sica et al., 1997). However, NFκB also induces the expression of transcription factor Iκbζ that also regulates *Ifng* transcription (Miyake et al., 2010). We have previously reported that in hypoxic conditions, IFN-γ production by NK cells was attenuated through HIF-1α-mediated inhibition of IL-18-NFκB-Iκbζ pathway, and IL-18 was necessary for tumor growth control by HIF-1α-deficient NK cells (Ni et al., 2020). In addition, Iκbζ-deficient mice demonstrate reduced IFN-γ production by NK cells, but not other cells (Miyake et al., 2010), indicating that along with metabolic reconfiguration, the NFκB-Iκbζ axis might be the predominant checkpoint for regulating IFN-γ release by NK cells.

In summary, we identified a novel role of vitamin A metabolite, atRA, in regulating NK cell phenotype, metabolism and function. atRA-exposed NK cells display a reduced size, mitochondrial mass and membrane potential, associated with reduced metabolic rates. They favor an immunoregulatory microenvironment through a lower ability to remove immature DCs and to produce IFN-γ in response to wide range of stimuli. In addition, atRA-treated NK cells support FoxP3[+] T cell differentiation and/or proliferation. Therefore, in vitamin A-enriched tissue environments, NK cells can contribute to the maintenance of homeostasis, or to regulation and/or resolution of inflammatory responses in pathological settings.

Materials and methods

Mice
Rag2-deficient mice were bred and maintained at the animal facility of Medical Faculty Mannheim, University of Heidelberg. C57BL6/N mice were purchased from Janvier. Both male and female mice were used for experiments (8-40 weeks of age). All animal experiments were approved by the local authorities.

Preparation of single-cell suspension from spleen and bone marrow
Single-cell suspensions were obtained from spleens after red blood cell lysis with buffered chloride potassium phosphate solution (ACK buffer) for 2-5 min at room temperature. Single-cell suspensions from bone marrow were obtained by flushing femurs and tibias with phosphate buffered saline (PBS). Subsequently, cells were collected by centrifugation, and red blood cells were lysed with ACK buffer.

NK cell isolation and culture
NK cells were enriched from splenic single-cell suspensions of Rag2-deficient mice after removal of adherent cell fraction by short culture at 37°C. Cells were then cultured for 5-7 days in Roswell Park Memorial Institute (RPMI)-1640 medium (Sigma-Aldrich) supplemented with 10% heat-inactivated fetal calf serum (FCS), 2 mM glutamine, 1% non-essential amino acids, 1 mM sodium pyruvate, 100 U/ml Penicillin/Streptomycin, 50 µM β-mercaptoethanol (all from GIBCO) and 1700 U/ml recombinant human IL-2 (National Institutes of Health). All-*trans* retinoic acid (atRA; Sigma-Aldrich) was dissolved in dimethyl sulfoxide (DMSO; Sigma-Aldrich) and added at the concentrations indicated in the Results. Equivalent volumes of DMSO were used as control.

Tumor cell lines
The mouse lymphoma cell lines RMA-S and YAC-1 were cultured in RPMI-1640 medium supplemented with 10% FCS and 100 U/ml Penicillin/Ampicillin. The mouse melanoma cell line B16 was cultured in Dulbecco's Modified Eagle Medium (DMEM; Sigma-Aldrich) supplemented with 10% FCS and 100 U/ml Penicillin/ Streptomycin.

In vitro stimulation of NK cells
After 7 days of culture with atRA, NK cells were cultured for 5 hours with RMA-S, YAC-1 or B16 cells (1:1), or stimulated with cytokines (1 ng/mL of IL-12 and 10 ng/mL of IL-18), plate-bound antibodies (Abs) anti-NK1.1 (clone PK136; Biolegend), anti-NKG2D (clone A10; Biolegend), or anti-NKp46 (polyclonal goat IgG; R&D Systems), or plate-bound CD155-Fc chimera protein (CD155-IEGRMDP-Mouse IgG2a; R&D Systems). For plate coating with Abs or Fc, flat-bottomed 96-well plates were incubated overnight at 4°C with 50 µL of 10 µg/ml Ab or 2 µg/ml chimera protein solution in PBS. Stimulation was performed in the presence of fluorophore-labeled anti-CD107a mAb (clone 1D4B; Biolegend) or the isotype-matched mAb control. GolgiStop™ (BD Biosciences) was added 4 hours before the end of stimulation.

Flow-cytometry and cell sorting
Cells were incubated with Fc-receptor-blocking reagent (10% supernatant of αCD16/CD32-producing hybridoma 2.4G2), followed by incubation with fluorochrome-conjugated monoclonal antibodies directed against CD45.1 (clone A20), CD45.2 (clone 104), CD4 (clone GK1.5), CD3ε (clone 145-2C11), NK1.1 (clone PK136), NKG2D (clone CX5), NKp46 (clone 29A1.4, CD226 (clone 10E5), CD200R (clone OX-2R), CD11b (clone M1/70), CD27 (clone LG.3A10), LAG3 (clone C9B7W), TRAIL (clone N2B2), CD62L (clone MEL-14), KLRG1 (clone 2F1), CD39 (clone Duha59), CD73 (clone TY/11.8),

CD38 (clone 90), IL-18Rα (clone A17071D), CD86 (clone GL-1), CD25 (clone PC61) (all from Biolegend), and I-A/I-E (clone M5/114.15.2; BD Biosciences). For the detection of cytokines and transcription factors, antibodies specific for IFN-γ (clone XMG1.2; Biolegend), Ki67 (clone 16A8; Biolegend), IκBζ (clone LK2NAP; Invitrogen), cMyc (clone D84C12; Cell Signaling Technology), NFκB-p50 (clone D4P4D; Cell Signaling Technology), Eomes (clone X4-83; BD Bioscience), FoxP3 (clone FJK-16s; eBiosciences), Rorγt (clone Q31-378; BD Biosciences) and T-bet (clone 4B10; Biolegend), and eBioscience™ Foxp3/Transcription Factor Staining Buffer Set (Thermo Fisher) were used according to the manufacturer's instructions. For the detection of phosphorylated proteins, including phosphorylated STAT4 (clone D2E4), phosphorylated NFκB-p65 (clone 93H1), phosphorylated S6 (clone D57.2.2E), and phosphorylated IKKα/β (clone 16A6) (all from Cell Signaling Technology), BD Cytofix/Cytoperm™ and BD Phosflow™ buffer (BD Biosciences) were used according to the manufacturer's instructions. 7-aminoactinomycin D (7-AAD; BD Bioscience) or Zombie Aqua™ (Biolegend) were used to exclude dead cells. For the analysis of lipid uptake and mitochondria, Bodipy™ FL C16 (Invitrogen) or CellROX™ Deep Red dye, MitoTracker™ Green FM dye, and MitoProbe™ tetramethylrhodamine methyl ester (TMRM) dye (all from Thermo Fischer) were used, according to the manufacturer's instructions. For the detection of apoptosis, Annexin V Apoptosis Detection Kit (Biolegend) was used. Flow-cytometry was performed with LSRFortessa™ (BD Biosciences), and data were analyzed using FlowJo™ software (BD Biosciences). For the gene expression analysis, NK cells (NK1.1[+]NKp46[+]TRAIL[neg]CD200R[neg]) were sorted from pre-gated live/single/CD45[+] splenocytes of Rag2-deficient mice using BD FACSAria™ Fusion (BD Biosciences).

Measurement of oxygen consumption rate and extracellular acidification rate
The analysis of oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) was performed using Seahorse™ XF HS Mini Analyzer (Agilent). After 7 days of culture with atRA or an equivalent volume of DMSO, NK cells were resuspended in XF RPMI medium (Agilent) containing 10 mM glucose, 2 mM L-glutamine, and 1 mM sodium pyruvate. The OCR and ECAR were analyzed in response to 1.5 µM oligomycin, 2.5 µM BAM15 ((2-fluorophenyl){6-[(2-fluorophenyl)amino](1,2,5-oxadiazolo[3,4–e]pyrazin-5-yl)}amine), and 0.5 µM Rotenone and Antimycin A (all from Agilent), according to the manufacturer's recommendations.

Co-culture of NK cells and DCs
To obtain dendritic cells (DCs), bone-marrow cells of C57BL6/N mice were cultured in DMEM medium (Sigma-Aldrich) supplemented with 10% FCS, 2 mM glutamine, 1% non-essential amino acids, 1 mM sodium pyruvate, 100 U/ml Penicillin/Streptomycin (all from GIBCO) and GM-CSF (10% supernatant of GM-CSF-producing X6310 cell line). NK cells were co-cultured with DCs at ratio 1:2 for the time indicated in the Result sections. Collected supernatants were assessed for cytokine content using ELISA MAX™ Standard Set Mouse IFN-γ (Biolegend). DCs were analyzed for the expression of CD86 and MHC class II using flow cytometry. DC apoptosis was assessed by measuring Annexin V binding to the cell surface.

Differentiation of CD4[+] T cells
Naïve CD4[+] T cells were isolated from spleens of C57BL6/N mice using MACS™ isolation kit (Miltenyi). The purity of naïve T cells (CD45[+]CD3ε[+]CD4[+]CD62L[+]CD44[neg]) was ≥ 90%. T cells were then activated with plate-bound anti-CD3ε (clone 145-2C11; Biolegend) mAbs (plates were prepared by incubating overnight at 4°C with 50 µL of 2 µg/ml Abs in PBS) and 0.5 µg/ml of soluble anti-CD28 (clone 37.51; Biolegend) mAbs. Following combinations of cytokines and antibodies were used for the differentiation of T cells: for Th1 cells - 1 µg/ml anti-IL-4 (clone 11B11; Biolegend), 30 U/ml rhIL-2 (National Institutes of Health), and 10 ng/ml IL-12 (Peprotech); for Th17 - 1 µg/ml anti-IFN-γ (clone

R4-6A2), 1 µg/ml anti-IL-2 (clone JES6-1A12), 1 µg/ml anti-IL-4 (clone 11B11) (all from Biolegend), 20 ng/ml IL-6, 20 ng/mL IL-1β (both from Peprotech), 1 ng/ml TGFß1, and 10 ng/ml IL-23 (both from Biolegend); for Treg - 1 µg/ml anti-IFN-γ (clone R4-6A2), 1 µg/ml anti-IL-4 (clone 11B11) (both from Biolegend), 30 U/ml rhIL-2 (National Institutes of Health), and 2 ng/ml TGF-ß1 (Biolegend). NK cells treated with atRA for 5 days were collected and cultured with naïve T cells at a ratio of 1:1 for 2 days. To measure T cell cytokine production, cells were stimulated with 50 ng/mL of Phorbol myristate acetate (PMA) and 1 µM of ionomycin. In some experiments, 5 µM of ZM 241385, 5 µM of SCH 58261 (both A2AR agonists, Tocris), 7.5 µg/mL of anti-IL-10 antibody (clone JES5-2A5, BioXCell), 4 µg/mL of anti-CCL5 Ab (polyclonal goat IgG; R&D Systems), or 4 µg/mL of anti-CCL1 Ab (clone 148113; R&D Systems), was added to the cultures.

RNA isolation and quantitative PCR

After 7 days of culture with atRA or equivalent volume of DMSO, NK cells were lysed with RLT buffer containing 1% of β-mercaptoethanol, and RNA extraction was performed using RNeasy® Mini Kit (Qiagen) according to manufacturer's instructions. Genomic DNA was removed by using TURBO DNA-free™ kit (ThermoFisher). RNA concentration was determined using plate reader Infinite 200 pro (TECAN). cDNA was synthesized from a total of 400 ng of RNA, using RT^2 First Strand Kit (Qiagen). For the analysis of relative mRNA expression, RT² Profiler™ PCR array mouse cytokines & chemokines (Qiagen) was used according to manufacturer's instructions. mRNA expression was calculated using $2^{-\Delta CP}$ method, relative to b2m transcript.

Gene expression analysis

NK cells isolated from spleens of Rag2$^{-/-}$ mice using flow-cytometric sort were cultured with atRA or equivalent volume of DMSO for 7 days, and then used for gene expression analysis. Gene expression profiling was performed using Clariom™ D Mouse Arrays (Thermo Fisher). Biotinylated antisense cDNA was prepared according to the standard labeling protocol with the GeneChip® WT Plus Reagent Kit and the GeneChip® Hybridization, Wash and Stain Kit (both from Thermo Fisher). Thereafter, chip hybridization was performed using GeneChip Hybridization oven 640, then dyed in the GeneChip Fluidics Station 450, and afterwards scanned with a GeneChip Scanner 3000 (all from Affymetrix). Raw fluorescence intensity values were normalized by applying quantile normalization and RMA background correction. A custom CDF version 22 with ENTREZ based gene definitions was used to annotate the arrays (Dai et al., 2005). Differentially expressed transcripts were obtained by One Way-analysis of variance (ANOVA) using a commercial software package SAS JMP15 Genomics, version 10, from SAS (SAS Institute). Parameters applied were a false positive rate of α = 0.05 with FDR correction as the level of significance. Data are visualized using EnhancedVolcano (https://github.com/kevinblighe/EnhancedVolcano) and Pheatmap package in R (https://github.com/raivokolde/pheatmap). For pathway enrichment analysis, following criteria were used to choose specifically expressed genes: fold change > 1.5 or < -1.5 and adjusted p-value < 0.05. Pathway enrichment analysis was based on the pathway collection in Kyoto Encyclopedia of Genes and Genomes (KEGG) (https://www.genome.jp/kegg/ pathway.html).

Quantification and statistical analysis

Experimental results are shown as mean ± SEM; n represents numbers of experiments, and is specified in Figure Legends. The statistical significance was assessed with the paired Student's t-test for normally distributed data, Welch's t-test for data with unequal variances, One Way-ANOVA followed by Tukey's post-hoc test for multiple testing, all using with GraphPad Prism software (GraphPad Software). The differences between experimental groups were considered significant when *p < 0.05, **p < 0.01, ***p < 0.001, ****p <0.0001.

Data Availability

The microarray gene expression data from this publication have been deposited to the GEO database (https://www.ncbi.nlm.nih.gov/geo) and assigned the identifier GSE218594 (available per acceptance for publication, access to reviewers is provided via secured token).

References

Ardavin, C., Martinez del Hoyo, G., Martin, P., Anjuere, F., Arias, C. F., Marin, A. R., . . . Hernandez, H. (2001). Origin and differentiation of dendritic cells. *Trends Immunol, 22*(12), 691-700. doi:10.1016/s1471-4906(01)02059-2

Bakdash, G., Vogelpoel, L. T., van Capel, T. M., Kapsenberg, M. L., & de Jong, E. C. (2015). Retinoic acid primes human dendritic cells to induce gut-homing, IL-10-producing regulatory T cells. *Mucosal Immunol, 8*(2), 265-278. doi:10.1038/mi.2014.64

Blomhoff, R., Green, M. H., Berg, T., & Norum, K. R. (1990). Transport and storage of vitamin A. *Science, 250*(4979), 399-404. doi:10.1126/science.2218545

Borg, C., Jalil, A., Laderach, D., Maruyama, K., Wakasugi, H., Charrier, S., . . . Zitvogel, L. (2004). NK cell activation by dendritic cells (DCs) requires the formation of a synapse leading to IL-12 polarization in DCs. *Blood, 104*(10), 3267-3275. doi:10.1182/blood-2004-01-0380

Bottcher, J. P., Schanz, O., Garbers, C., Zaremba, A., Hegenbarth, S., Kurts, C., . . . Knolle, P. A. (2014). IL-6 trans-signaling-dependent rapid development of cytotoxic CD8+ T cell function. *Cell Rep, 8*(5), 1318-1327. doi:10.1016/j.celrep.2014.07.008

Cerboni, C., Zingoni, A., Cippitelli, M., Piccoli, M., Frati, L., & Santoni, A. (2007). Antigen-activated human T lymphocytes express cell-surface NKG2D ligands via an ATM/ATR-dependent mechanism and become susceptible to autologous NK- cell lysis. *Blood, 110*(2), 606-615. doi:10.1182/blood-2006-10-052720

Chan, C. J., Andrews, D. M., McLaughlin, N. M., Yagita, H., Gilfillan, S., Colonna, M., & Smyth, M. J. (2010). DNAM-1/CD155 interactions promote cytokine and NK cell-mediated suppression of poorly immunogenic melanoma metastases. *J Immunol, 184*(2), 902-911. doi:10.4049/jimmunol.0903225

Chidipi, B., Shah, S. I., Reiser, M., Kanithi, M., Garces, A., Cha, B. J., . . . Noujaim, S. F. (2021). All-Trans Retinoic Acid Increases DRP1 Levels and Promotes Mitochondrial Fission. *Cells, 10*(5). doi:10.3390/cells10051202

Chinen, T., Kannan, A. K., Levine, A. G., Fan, X., Klein, U., Zheng, Y., . . . Rudensky, A. Y. (2016). An essential role for the IL-2 receptor in Treg cell function. *Nat Immunol, 17*(11), 1322-1333. doi:10.1038/ni.3540

Colonna, M. (2009). Interleukin-22-producing natural killer cells and lymphoid tissue inducer-like cells in mucosal immunity. *Immunity, 31*(1), 15-23. doi:10.1016/j.immuni.2009.06.008

Dai, M., Wang, P., Boyd, A. D., Kostov, G., Athey, B., Jones, E. G., . . . Meng, F. (2005). Evolving gene/transcript definitions significantly alter the interpretation of GeneChip data. *Nucleic Acids Res, 33*(20), e175. doi:10.1093/nar/gni179

Diefenbach, A., Jensen, E. R., Jamieson, A. M., & Raulet, D. H. (2001). Rae1 and H60 ligands of the NKG2D receptor stimulate tumour immunity. *Nature, 413*(6852), 165-171. doi:10.1038/35093109

Diehl, L., Schurich, A., Grochtmann, R., Hegenbarth, S., Chen, L., & Knolle, P. A. (2008). Tolerogenic maturation of liver sinusoidal endothelial cells promotes B7-homolog 1-dependent CD8+ T cell tolerance. *Hepatology, 47*(1), 296-305. doi:10.1002/hep.21965

Duhen, R., Glatigny, S., Arbelaez, C. A., Blair, T. C., Oukka, M., & Bettelli, E. (2013). Cutting edge: the pathogenicity of IFN-gamma-producing Th17 cells is independent of T-bet. *J Immunol, 190*(9), 4478-4482. doi:10.4049/jimmunol.1203172

Eksteen, B., Mora, J. R., Haughton, E. L., Henderson, N. C., Lee-Turner, L., Villablanca, E. J., . . . Adams, D. H. (2009). Gut homing receptors on CD8 T cells are retinoic acid dependent and not maintained by liver dendritic or stellate cells. *Gastroenterology, 137*(1), 320-329. doi:10.1053/j.gastro.2009.02.046

Ferlazzo, G., Semino, C., & Melioli, G. (2001). HLA class I molecule expression is up-regulated during maturation of dendritic cells, protecting them from natural killer cell-mediated lysis. *Immunol Lett, 76*(1), 37-41. doi:10.1016/s0165-2478(00)00323-0

Ferlazzo, G., Tsang, M. L., Moretta, L., Melioli, G., Steinman, R. M., & Munz, C. (2002). Human dendritic cells activate resting natural killer (NK) cells and are recognized via the NKp30 receptor by activated NK cells. *J Exp Med, 195*(3), 343-351. doi:10.1084/jem.20011149

Fraker, L. D., Halter, S. A., & Forbes, J. T. (1986). Effects of orally administered retinol on natural killer cell activity in wild type BALB/c and congenitally athymic BALB/c mice. *Cancer Immunol Immunother, 21*(2), 114-118. doi:10.1007/BF00199858

Friedman, S. L. (2008). Hepatic stellate cells: protean, multifunctional, and enigmatic cells of the liver. *Physiol Rev, 88*(1), 125-172. doi:10.1152/physrev.00013.2007

Gagliani, N., Amezcua Vesely, M. C., Iseppon, A., Brockmann, L., Xu, H., Palm, N. W., . . . Flavell, R. A. (2015). Th17 cells transdifferentiate into regulatory T cells during resolution of inflammation. *Nature, 523*(7559), 221-225. doi:10.1038/nature14452

Gao, B., Jeong, W. I., & Tian, Z. (2008). Liver: An organ with predominant innate immunity. *Hepatology, 47*(2), 729-736. doi:10.1002/hep.22034

Hall, J. A., Grainger, J. R., Spencer, S. P., & Belkaid, Y. (2011). The role of retinoic acid in tolerance and immunity. *Immunity, 35*(1), 13-22. doi:10.1016/j.immuni.2011.07.002

Hamerman, J. A., Ogasawara, K., & Lanier, L. L. (2005). NK cells in innate immunity. *Curr Opin Immunol, 17*(1), 29-35. doi:10.1016/j.coi.2004.11.001

Handa, K., Suzuki, R., Matsui, H., Shimizu, Y., & Kumagai, K. (1983). Natural killer (NK) cells as a responder to interleukin 2 (IL 2). II. IL 2-induced interferon gamma production. *J Immunol, 130*(2), 988-992. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/6294182

Hayakawa, Y., Screpanti, V., Yagita, H., Grandien, A., Ljunggren, H. G., Smyth, M. J., & Chambers, B. J. (2004). NK cell TRAIL eliminates immature dendritic cells in vivo and limits dendritic cell vaccination efficacy. *J Immunol, 172*(1), 123-129. doi:10.4049/jimmunol.172.1.123

Hoelzinger, D. B., Smith, S. E., Mirza, N., Dominguez, A. L., Manrique, S. Z., & Lustgarten, J. (2010). Blockade of CCL1 inhibits T regulatory cell suppressive function enhancing tumor immunity without affecting T effector responses. *J Immunol, 184*(12), 6833-6842. doi:10.4049/jimmunol.0904084

Iwata, M., Hirakiyama, A., Eshima, Y., Kagechika, H., Kato, C., & Song, S. Y. (2004). Retinoic acid imprints gut-homing specificity on T cells. *Immunity, 21*(4), 527-538. doi:10.1016/j.immuni.2004.08.011

Kane, M. A., Folias, A. E., & Napoli, J. L. (2008). HPLC/UV quantitation of retinal, retinol, and retinyl esters in serum and tissues. *Anal Biochem, 378*(1), 71-79. doi:10.1016/j.ab.2008.03.038

Kasahara, T., Hooks, J. J., Dougherty, S. F., & Oppenheim, J. J. (1983). Interleukin 2-mediated immune interferon (IFN-gamma) production by human T cells and T cell subsets. *J Immunol, 130*(4), 1784-1789. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/6403613

Kim, M. H., Taparowsky, E. J., & Kim, C. H. (2015). Retinoic Acid Differentially Regulates the Migration of Innate Lymphoid Cell Subsets to the Gut. *Immunity, 43*(1), 107-119. doi:10.1016/j.immuni.2015.06.009

Kuehnemuth, B., Piseddu, I., Wiedemann, G. M., Lauseker, M., Kuhn, C., Hofmann, S., . . . Anz, D. (2018). CCL1 is a major regulatory T cell attracting factor in human breast cancer. *BMC Cancer, 18*(1), 1278. doi:10.1186/s12885-018-5117-8

Kwon, H. J., Choi, G. E., Ryu, S., Kwon, S. J., Kim, S. C., Booth, C., . . . Kim, H. S. (2016). Stepwise phosphorylation of p65 promotes NF-kappaB activation and NK cell responses during target cell recognition. *Nat Commun, 7*, 11686. doi:10.1038/ncomms11686

Lanier, L. L. (2001). On guard--activating NK cell receptors. *Nat Immunol, 2*(1), 23-27. doi:10.1038/83130

Laouar, Y., Sutterwala, F. S., Gorelik, L., & Flavell, R. A. (2005). Transforming growth factor-beta controls T helper type 1 cell development through regulation of natural killer cell interferon-gamma. *Nat Immunol, 6*(6), 600-607. doi:10.1038/ni1197

Laurence, A., Tato, C. M., Davidson, T. S., Kanno, Y., Chen, Z., Yao, Z., . . . O'Shea J, J. (2007). Interleukin-2 signaling via STAT5 constrains T helper 17 cell generation. *Immunity, 26*(3), 371-381. doi:10.1016/j.immuni.2007.02.009

Lee, K. A., Song, Y. C., Kim, G. Y., Choi, G., Lee, Y. S., Lee, J. M., & Kang, C. Y. (2012). Retinoic acid alleviates Con A-induced hepatitis and differentially regulates effector production in NKT cells. *Eur J Immunol, 42*(7), 1685-1694. doi:10.1002/eji.201142322

Lee, Y. K., Mukasa, R., Hatton, R. D., & Weaver, C. T. (2009). Developmental plasticity of Th17 and Treg cells. *Curr Opin Immunol, 21*(3), 274-280. doi:10.1016/j.coi.2009.05.021

Li, A., He, M., Wang, H., Qiao, B., Chen, P., Gu, H., . . . He, S. (2007). All-trans retinoic acid negatively regulates cytotoxic activities of nature killer cell line 92. *Biochem Biophys Res Commun, 352*(1), 42-47. doi:10.1016/j.bbrc.2006.10.132

Li, M. O., Sanjabi, S., & Flavell, R. A. (2006). Transforming growth factor-beta controls development, homeostasis, and tolerance of T cells by regulatory T cell-dependent and -independent mechanisms. *Immunity, 25*(3), 455-471. doi:10.1016/j.immuni.2006.07.011

Lischke, A., Moriggl, R., Brandlein, S., Berchtold, S., Kammer, W., Sebald, W., . . . Friedrich, K. (1998). The interleukin-4 receptor activates STAT5 by a mechanism that relies upon common gamma-chain. *J Biol Chem, 273*(47), 31222-31229. doi:10.1074/jbc.273.47.31222

Littwitz-Salomon, E., Moreira, D., Frost, J. N., Choi, C., Liou, K. T., Ahern, D. K., . . . Finlay, D. K. (2021). Metabolic requirements of NK cells during the acute response against retroviral infection. *Nat Commun, 12*(1), 5376. doi:10.1038/s41467-021-25715-z

Loftus, R. M., Assmann, N., Kedia-Mehta, N., O'Brien, K. L., Garcia, A., Gillespie, C., . . . Finlay, D. K. (2018). Amino acid-dependent cMyc expression is essential for NK cell metabolic and functional responses in mice. *Nat Commun, 9*(1), 2341. doi:10.1038/s41467-018-04719-2

Marcais, A., Cherfils-Vicini, J., Viant, C., Degouve, S., Viel, S., Fenis, A., . . . Walzer, T. (2014). The metabolic checkpoint kinase mTOR is essential for IL-15 signaling during the development and activation of NK cells. *Nat Immunol, 15*(8), 749-757. doi:10.1038/ni.2936

Marcais, A., Viel, S., Grau, M., Henry, T., Marvel, J., & Walzer, T. (2013). Regulation of mouse NK cell development and function by cytokines. *Front Immunol, 4*, 450. doi:10.3389/fimmu.2013.00450

Martin-Fontecha, A., Thomsen, L. L., Brett, S., Gerard, C., Lipp, M., Lanzavecchia, A., & Sallusto, F. (2004). Induced recruitment of NK cells to lymph nodes provides IFN-gamma for T(H)1 priming. *Nat Immunol, 5*(12), 1260-1265. doi:10.1038/ni1138

Miyake, T., Satoh, T., Kato, H., Matsushita, K., Kumagai, Y., Vandenbon, A., . . . Takeuchi, O. (2010). IkappaBzeta is essential for natural killer cell activation in response to IL-12 and IL-18. *Proc Natl Acad Sci U S A, 107*(41), 17680-17685. doi:10.1073/pnas.1012977107

Mora, J. R., Iwata, M., Eksteen, B., Song, S. Y., Junt, T., Senman, B., . . . von Andrian, U. H. (2006). Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science, 314*(5802), 1157-1160. doi:10.1126/science.1132742

Morandi, B., Bougras, G., Muller, W. A., Ferlazzo, G., & Munz, C. (2006). NK cells of human secondary lymphoid tissues enhance T cell polarization via IFN-gamma secretion. *Eur J Immunol, 36*(9), 2394-2400. doi:10.1002/eji.200636290

Morita, H., Kubo, T., Ruckert, B., Ravindran, A., Soyka, M. B., Rinaldi, A. O., . . . Akdis, C. A. (2019). Induction of human regulatory innate lymphoid cells from group 2 innate lymphoid cells by retinoic acid. *J Allergy Clin Immunol, 143*(6), 2190-2201 e2199. doi:10.1016/j.jaci.2018.12.1018

Mui, A. L., Wakao, H., O'Farrell, A. M., Harada, N., & Miyajima, A. (1995). Interleukin-3, granulocyte-macrophage colony stimulating factor and interleukin-5 transduce signals through two STAT5 homologs. *EMBO J, 14*(6), 1166-1175. doi:10.1002/j.1460-2075.1995.tb07100.x

Nakahira, M., Ahn, H. J., Park, W. R., Gao, P., Tomura, M., Park, C. S., . . . Fujiwara, H. (2002). Synergy of IL-12 and IL-18 for IFN-gamma gene expression: IL-12-induced STAT4 contributes to IFN-gamma promoter activation by up-regulating the binding activity of IL-18-induced activator protein 1. *J Immunol, 168*(3), 1146-1153. doi:10.4049/jimmunol.168.3.1146

Ni, J., Wang, X., Stojanovic, A., Zhang, Q., Wincher, M., Buhler, L., . . . Cerwenka, A. (2020). Single-Cell RNA Sequencing of Tumor-Infiltrating NK Cells Reveals that Inhibition of Transcription Factor HIF-1alpha Unleashes NK Cell Activity. *Immunity, 52*(6), 1075-1087 e1078. doi:10.1016/j.immuni.2020.05.001

Oberg, L., Johansson, S., Michaelsson, J., Tomasello, E., Vivier, E., Karre, K., & Hoglund, P. (2004). Loss or mismatch of MHC class I is sufficient to trigger NK cell-mediated rejection of resting

lymphocytes in vivo - role of KARAP/DAP12-dependent and -independent pathways. *Eur J Immunol, 34*(6), 1646-1653. doi:10.1002/eji.200424913

Ohta, A. (2016). A Metabolic Immune Checkpoint: Adenosine in Tumor Microenvironment. *Front Immunol, 7*, 109. doi:10.3389/fimmu.2016.00109

Ohta, A., & Sitkovsky, M. (2014). Extracellular adenosine-mediated modulation of regulatory T cells. *Front Immunol, 5*, 304. doi:10.3389/fimmu.2014.00304

Okamura, H., Kashiwamura, S., Tsutsui, H., Yoshimoto, T., & Nakanishi, K. (1998). Regulation of interferon-gamma production by IL-12 and IL-18. *Curr Opin Immunol, 10*(3), 259-264. doi:10.1016/s0952-7915(98)80163-5

Osei-Bordom, D., Bozward, A. G., & Oo, Y. H. (2020). The hepatic microenvironment and regulatory T cells. *Cell Immunol, 357*, 104195. doi:10.1016/j.cellimm.2020.104195

Pandiyan, P., Bhaskaran, N., Zou, M., Schneider, E., Jayaraman, S., & Huehn, J. (2019). Microbiome Dependent Regulation of Tregs and Th17 Cells in Mucosa. *Front Immunol, 10*, 426. doi:10.3389/fimmu.2019.00426

Papaioannou, S., See, J.-X., Jeong, M., De La Torre, C., Reiners-Koch, P.-S., Sati, A., . . . Stojanovic, A. (2022). Liver sinusoidal endothelial cells orchestrate NK cell recruitment and activation in acute inflammatory liver injury. *bioRxiv*, 2022.2007.2015.500206. doi:10.1101/2022.07.15.500206

Pegram, H. J., Andrews, D. M., Smyth, M. J., Darcy, P. K., & Kershaw, M. H. (2011). Activating and inhibitory receptors of natural killer cells. *Immunol Cell Biol, 89*(2), 216-224. doi:10.1038/icb.2010.78

Piccioli, D., Sbrana, S., Melandri, E., & Valiante, N. M. (2002). Contact-dependent stimulation and inhibition of dendritic cells by natural killer cells. *J Exp Med, 195*(3), 335-341. doi:10.1084/jem.20010934

Powell, J. D., Lerner, C. G., & Schwartz, R. H. (1999). Inhibition of cell cycle progression by rapamycin induces T cell clonal anergy even in the presence of costimulation. *J Immunol, 162*(5), 2775-2784. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/10072524

Rabinovich, B. A., Li, J., Shannon, J., Hurren, R., Chalupny, J., Cosman, D., & Miller, R. G. (2003). Activated, but not resting, T cells can be recognized and killed by syngeneic NK cells. *J Immunol, 170*(7), 3572-3576. doi:10.4049/jimmunol.170.7.3572

Rex, D. A. B., Agarwal, N., Prasad, T. S. K., Kandasamy, R. K., Subbannayya, Y., & Pinto, S. M. (2020). A comprehensive pathway map of IL-18-mediated signalling. *J Cell Commun Signal, 14*(2), 257-266. doi:10.1007/s12079-019-00544-4

Rigobello, M. P., Scutari, G., Friso, A., Barzon, E., Artusi, S., & Bindoli, A. (1999). Mitochondrial permeability transition and release of cytochrome c induced by retinoic acids. *Biochem Pharmacol, 58*(4), 665-670. doi:10.1016/s0006-2952(99)00149-5

Semino, C., Angelini, G., Poggi, A., & Rubartelli, A. (2005). NK/iDC interaction results in IL-18 secretion by DCs at the synaptic cleft followed by NK cell activation and release of the DC maturation factor HMGB1. *Blood, 106*(2), 609-616. doi:10.1182/blood-2004-10-3906

Sica, A., Dorman, L., Viggiano, V., Cippitelli, M., Ghosh, P., Rice, N., & Young, H. A. (1997). Interaction of NF-kappaB and NFAT with the interferon-gamma promoter. *J Biol Chem, 272*(48), 30412-30420. doi:10.1074/jbc.272.48.30412

Smyth, M. J., Cretney, E., Kelly, J. M., Westwood, J. A., Street, S. E., Yagita, H., . . . Hayakawa, Y. (2005). Activation of NK cell cytotoxicity. *Mol Immunol, 42*(4), 501-510. doi:10.1016/j.molimm.2004.07.034

Spencer, S. P., Wilhelm, C., Yang, Q., Hall, J. A., Bouladoux, N., Boyd, A., . . . Belkaid, Y. (2014). Adaptation of innate lymphoid cells to a micronutrient deficiency promotes type 2 barrier immunity. *Science, 343*(6169), 432-437. doi:10.1126/science.1247606

Spinelli, J. B., & Haigis, M. C. (2018). The multifaceted contributions of mitochondria to cellular metabolism. *Nat Cell Biol, 20*(7), 745-754. doi:10.1038/s41556-018-0124-1

Sun, C. M., Hall, J. A., Blank, R. B., Bouladoux, N., Oukka, M., Mora, J. R., & Belkaid, Y. (2007). Small intestine lamina propria dendritic cells promote de novo generation of Foxp3 T reg cells via retinoic acid. *J Exp Med, 204*(8), 1775-1785. doi:10.1084/jem.20070602

Tian, Z., Chen, Y., & Gao, B. (2013). Natural killer cells in liver disease. *Hepatology, 57*(4), 1654-1662. doi:10.1002/hep.26115

Tourniaire, F., Musinovic, H., Gouranton, E., Astier, J., Marcotorchino, J., Arreguin, A., . . . Landrier, J. F. (2015). All-trans retinoic acid induces oxidative phosphorylation and mitochondria biogenesis in adipocytes. *J Lipid Res, 56*(6), 1100-1109. doi:10.1194/jlr.M053652

Ueno, A., Jeffery, L., Kobayashi, T., Hibi, T., Ghosh, S., & Jijon, H. (2018). Th17 plasticity and its relevance to inflammatory bowel disease. *J Autoimmun, 87*, 38-49. doi:10.1016/j.jaut.2017.12.004

Vang, K. B., Yang, J., Mahmud, S. A., Burchill, M. A., Vegoe, A. L., & Farrar, M. A. (2008). IL-2, -7, and -15, but not thymic stromal lymphopoeitin, redundantly govern CD4+Foxp3+ regulatory T cell development. *J Immunol, 181*(5), 3285-3290. doi:10.4049/jimmunol.181.5.3285

Veldhoen, M., Hocking, R. J., Atkins, C. J., Locksley, R. M., & Stockinger, B. (2006). TGFbeta in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity, 24*(2), 179-189. doi:10.1016/j.immuni.2006.01.001

Vitale, M., Della Chiesa, M., Carlomagno, S., Pende, D., Arico, M., Moretta, L., & Moretta, A. (2005). NK-dependent DC maturation is mediated by TNFalpha and IFNgamma released upon engagement of the NKp30 triggering receptor. *Blood, 106*(2), 566-571. doi:10.1182/blood-2004-10-4035

Vivier, E., Tomasello, E., Baratin, M., Walzer, T., & Ugolini, S. (2008). Functions of natural killer cells. *Nat Immunol, 9*(5), 503-510. doi:10.1038/ni1582

Walzer, T., Dalod, M., Robbins, S. H., Zitvogel, L., & Vivier, E. (2005). Natural-killer cells and dendritic cells: "l'union fait la force". *Blood, 106*(7), 2252-2258. doi:10.1182/blood-2005-03-1154

Wang, R., Dillon, C. P., Shi, L. Z., Milasta, S., Carter, R., Finkelstein, D., . . . Green, D. R. (2011). The transcription factor Myc controls metabolic reprogramming upon T lymphocyte activation. *Immunity, 35*(6), 871-882. doi:10.1016/j.immuni.2011.09.021

Wang, W., Chen, N. Y., Ren, D., Davies, J., Philip, K., Eltzschig, H. K., . . . Weng, T. (2021). Enhancing Extracellular Adenosine Levels Restores Barrier Function in Acute Lung Injury Through Expression of Focal Adhesion Proteins. *Front Mol Biosci, 8*, 636678. doi:10.3389/fmolb.2021.636678

Xiao, S., Jin, H., Korn, T., Liu, S. M., Oukka, M., Lim, B., & Kuchroo, V. K. (2008). Retinoic acid increases Foxp3+ regulatory T cells and inhibits development of Th17 cells by enhancing TGF-beta-driven Smad3 signaling and inhibiting IL-6 and IL-23 receptor expression. *J Immunol, 181*(4), 2277-2284. doi:10.4049/jimmunol.181.4.2277

Young, H. A., & Hardy, K. J. (1995). Role of interferon-gamma in immune cell regulation. *J Leukoc Biol, 58*(4), 373-381. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/7561512

Yu, Y., Hagihara, M., Ando, K., Gansuvd, B., Matsuzawa, H., Tsuchiya, T., . . . Kato, S. (2001). Enhancement of human cord blood CD34+ cell-derived NK cell cytotoxicity by dendritic cells. *J Immunol, 166*(3), 1590-1600. doi:10.4049/jimmunol.166.3.1590

Zarek, P. E., Huang, C. T., Lutz, E. R., Kowalski, J., Horton, M. R., Linden, J., . . . Powell, J. D. (2008). A2A receptor signaling promotes peripheral tolerance by inducing T-cell anergy and the generation of adaptive regulatory T cells. *Blood, 111*(1), 251-259. doi:10.1182/blood-2007-03-081646

Zeng, R., Oderup, C., Yuan, R., Lee, M., Habtezion, A., Hadeiba, H., & Butcher, E. C. (2013). Retinoic acid regulates the development of a gut-homing precursor for intestinal dendritic cells. *Mucosal Immunol, 6*(4), 847-856. doi:10.1038/mi.2012.123

Zhao, Z., & Ross, A. C. (1995). Retinoic acid repletion restores the number of leukocytes and their subsets and stimulates natural cytotoxicity in vitamin A-deficient rats. *J Nutr, 125*(8), 2064-2073. doi:10.1093/jn/125.8.2064

Zheng, Y., Collins, S. L., Lutz, M. A., Allen, A. N., Kole, T. P., Zarek, P. E., & Powell, J. D. (2007). A role for mammalian target of rapamycin in regulating T cell activation versus anergy. *J Immunol, 178*(4), 2163-2170. doi:10.4049/jimmunol.178.4.2163

Zielinski, C. E., Mele, F., Aschenbrenner, D., Jarrossay, D., Ronchi, F., Gattorno, M., . . . Sallusto, F. (2012). Pathogen-induced human TH17 cells produce IFN-gamma or IL-10 and are regulated by IL-1beta. *Nature, 484*(7395), 514-518. doi:10.1038/nature10957

Zimmerman, M. A., Grenz, A., Tak, E., Kaplan, M., Ridyard, D., Brodsky, K. S., . . . Eltzschig, H. K. (2013). Signaling through hepatocellular A2B adenosine receptors dampens ischemia and reperfusion injury of the liver. *Proc Natl Acad Sci U S A, 110*(29), 12012-12017. doi:10.1073/pnas.1221733110

Figure Legends

Figure 1. atRA induces transcriptomic and phenotypic alterations in NK cells.

A   NK cells were cultured with 1 µM of atRA (or DMSO as a control solvent) in the presence of IL-2 for 7 days. The differential transcript expression in atRA-treated compared to control NK cells. Cut-off (red): Fold-Change > 1.2 or < -1.2, and p-value<0,01.

B   Hierarchical clustering of selected differentially expressed candidate transcripts.

C   Pathway enrichment analysis based on Kyoto Encyclopedia of Genes and Genomes (KEGG) for atRA-treated compared to control NK cells (FC > 1.5 or < 1.5 and adjusted p-value < 0.05). NES, normalized enrichment score; MFI, mean florescence intensity.

D-F   Control and atRA-treated NK cells were analyzed by flow cytometry (n=5-22). Graphs indicate mean ± SEM. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$ (paired Student's t-test; Welch's t-test for analysis of TRAIL expression).

Figure 2. atRA-treated NK cells show differential mitochondrial phenotype and reduced metabolic activity.

A   NK cells were cultured with 1 µM of atRA (or DMSO as a control solvent) in the presence of IL-2 for 7 days. Cells were analyzed for mitochondrial mass (Mitotracker™), mitochondrial membrane potential (TMRM) and reactive oxygen species (ROS) production (CellRox™) by flow cytometry. (n=3-4; mean ± SEM; *$p<0.05$by paired Student's t-test)

B-C   Oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) were analyzed upon treatment with oligomycin (oligo), BAM15, and Rotenone and Antimycin A (Rot/AntiA). Graphs show OCR and ECAR over 100 min time-lapse, and quantification of basal and maximal respiration, and baseline ECAR (n=3, mean ± SEM; *$p<0.05$ by paired Student's t-test).

D   Representative histograms (left) and quantification (right) of relative cell size, based on forward scatter area (FSC-A) assessment, and cellular granularity, based on side scatter area (SSC-A) assessment (n=15; mean ± SEM; **$p<0.01$ by paired Student's t-test).

Figure 3. atRA-treated NK cells produce lower amounts of IFN-γ than control NK cells upon stimulation.

A   NK cells were cultured with 1 µM of atRA (or DMSO as a control solvent) in the presence of IL-2 for 7 days. Relative mRNA expression by atRA-treated NK cells normalized to expression of control NK cells.

B-D   Representative contour-plots (right) and quantification (left) of IFN-γ-expressing NK cells, and the amount of IFN-γ in NK cells (in B) in response to IL-12 and IL-18 (n=6) (B), activating receptor triggering (n=4) (C), or tumor cells (n=3) (D). The graphs indicate mean ± SEM; *$p<0.05$, **$p<0.01$, ***$p<0.001$ by paired Student's t-test.

Figure 4. atRA-treated NK cells produce less IFN-γ and display lower activation of mTORC1 and NFκB pathway upon stimulation with IL-18.

A   NK cells were cultured with 1 μM of atRA (or DMSO as a control solvent) in the presence of IL-2 for 7 days. Representative contour-plots (left) and frequencies of IL-18Rα-expressing NK cells (right, n=10; mean ± SEM; *p<0.05 by paired Student's t-test).

B   Upon culture with atRA or DMSO, cells were stimulated with IL-18 or IL-12 for 4 hours and analyzed by flow cytometry. The percentage of IFN-γ-expressing NK cells in response to IL-18 or IL-12 (n=5; mean ± SEM; **p<0.01 by paired Student's t-test).

C-D   Representative histograms (left) and quantification (right) of (C) NFκB p50 and IκBζ, and (D) pS6 and cMyc expression by stimulated NK cells (n=4; mean ± SEM; *p<0.05, **p<0.01 by Tukey One Way-ANOVA).

Figure 5. atRA-treated NK cells favor immature BM-DC phenotype.

A   Schematic illustration depicting the co-culture of BM-DCs and atRA-treated NK cells (created with BioRender.com).

B   Representative histograms (left) and quantification (right) of CD86 and MHC II expression by BM-DCs cultured with atRA-treated or control NK cells for 24 hours (n=4; mean + SEM; *p<0.05 by Tukey One Way-ANOVA).

C   Quantification of secreted IFN-γ in the DC-NK cell co-culture (n=3; mean + SEM; *p<0.05 by paired Student's t-test).

D   The percentage of apoptotic (Annexin V-labeled) BM-DCs after 6 hours of co-culture with NK cells (n=4; mean + SEM. *p<0.05, **p<0.01 by Tukey One Way-ANOVA).

Figure 6. atRA-treated NK cells promote the differentiation and proliferation of FoxP3-expressing T cells.

A-B   CD45.1[+] NK cells were treated with 1 μM of atRA (or DMSO as a solvent control) in the presence of IL-2 for 5 days, and then co-cultured with freshly-isolated naïve CD45.2[+] CD4[+] T cells in Treg or Th17-polarizing conditions. T cells were analyzed for the expression of Rorγt, FoxP3, Ki67, CD25, and phosphorylated (p)-STAT5 by flow cytometry. Representative contour-plots (left) and frequencies (right) of FoxP3- and/or Rorγt-expressing CD4[+] T cells in (A) Treg-polarizing conditions (n=5-8) and (B) Th17-polarizing conditions (n=5-8). The graphs indicate mean ± SEM; *p<0.05 by Tukey OneWay-ANOVA

C   Frequencies of FoxP3-expressing CD4[+] T cells cultured in Th17-polarizing conditions with NK cells separated using permeable transwell membrane (n=3, mean + SEM; n.s. not significant and p=0.05 by Tukey OneWay-ANOVA).

D   Percentage of CD25-expressing CD4[+] T cells upon co-culture with NK cells (n=2).

E   Representative histograms (left) and quantification (right) of p-STAT5 in CD45.2[+] T cells cultured alone, with control NK cells, or with atRA-treated NK cells (n=5; mean ± SEM. *p<0.05, ***p<0.0001 by Tukey OneWay-ANOVA).

F   Frequency of FoxP3[+] Ki67[+] and FoxP3[neg] Ki67[+] CD4[+] T cells upon co-culture with NK cells (n=3; mean ± SEM. *p<0.05 by paired Student's t-test).

Expanded View Figure Legends

Expanded View Figure 1. atRA-treated NK cells upregulate DNAM-1, TRAIL, CD73, and CD39 in a time-dependent manner.

NK cells were cultured with 1 µM of atRA (or the equivalent volume of DMSO as a solvent control) in the presence of IL-2 for 2, 5, or 7 days. The expression of indicated molecules was analyzed by flow cytometry. Shown are the histograms representative of four independent experiments.

Expanded View Figure 2. Expression of CCR9, CD62L, CXCR3, and CXCR6 by atRA-treated and control NK cells.

NK cells were cultured with 1 µM of atRA (or the equivalent volume of DMSO as a solvent control) in the presence of IL-2 for 7 days. The expression of indicated molecules was analyzed by flow cytometry (n=4-9; mean ± SEM. **p<0.01, ***p<0.001 by paired Student's t-test).

Expanded View Figure 3. atRA does not affect NK cell expansion and lipid uptake.

A   NK cells were cultured with 1 µM of atRA (or the equivalent volume of DMSO as a control solvent) in the presence of IL-2 for 7 days. Quantification of cell expansion rates at the end of culture (day 7) calculated as final cell numbers divided by cell numbers seeded at the beginning of the culture (day 0), (n=15; mean ± SEM; not significant by paired Student's t-test)

B   Geometric mean fluorescence intensity (MFI) of Bodipy™ FL C16 staining analyzed by flow cytometry (n=4; mean ± SEM; not significant by paired Student's t-test).

Expanded View Figure 4. Expression pSTAT4, NFκB p-p65, and pIκB kinase alpha/beta (IKK α/β) by atRA-treated and control NK cells upon stimulation with cytokines.

A-B   NK cells were cultured with 1 µM of atRA (or the equivalent volume of DMSO as a control solvent) in the presence of IL-2 for 7 days, and then stimulated with IL-18 or IL-12 for additional 4 hours. Expression of phosphorylated (p)-STAT4, NFκB p-p65, and p-IKK α/β was analyzed by flow cytometry. Representative histograms (left) and quantification (right) of (A) p-STAT4, and (B) NFκB p-p65 and p-IKK α/β expression by NK cells are shown (n=3-4; mean ± SEM; not significant by paired Student's t-test).

Expanded View Figure 5. The co-culture with atRA-treated NK cells does not alter the differentiation of Th1 cells.

A   CD45.1[+] NK cells were treated with 1 µM of atRA (or the equivalent volume of DMSO as a control solvent) in the presence of IL-2 for 5 days. NK cells were then co-cultured with naïve CD45.2[+] CD4[+] T cells in Th1-polarizing conditions. Upon culture, cells were stimulated with PMA and ionomycin for 4 hours and analyzed by flow-cytometry. Representative contour-plots (up) and quantification (down) of T-bet expression and IFN-γ-production by CD4[+] T cells (n=3, mean + SEM; not significant by paired Student's t-test).

B-E   CD45.1[+] NK cells were treated with 1 µM of atRA (or the equivalent volume of DMSO as a control solvent) in the presence of IL-2 for 5 days, and then co-cultured with naïve CD45.2[+] CD4[+] T cells in Treg-polarizing conditions.

B-D   A2AR agonists, ZM 241385 or SCH 58261 (B), anti-IL-10 (C), anti-CCL5 or anti-CCL1 antibodies (D), were added to the co-culture of NK cells and T cells. Contour-plots show Rorγt and FoxP3 expression by CD4[+] T cells upon co-culture.

E   Frequencies of Ki67-expressing FoxP3[+] or FoxP3[neg] T cells upon co-culture with NK cells (n=3, mean ± SEM; **, $p<0.01$ by paired Student's t-test).

Acknowledgments

We thank Marian Wincher, Petra Bugert, Dalyan Devran and Patrick Matei for technical support; Sophia Papaioannou for help with experiments; the animal facility of the Medical Faculty Mannheim for assistance with animal care and experiments; and Dr. Margareta Correia and Prof. Guoliang Cui for support and discussion of the data. The project was supported by the Research Training Group at Heidelberg University train4CIM (to AC); intramural funding program SEED by Medical Faculty Mannheim (to AS); grants from the German Research Foundation [SPP1937 (CE 140/2-2 to AS and AC), SFB1366 (Project number 319 394046768-SFB 1366; C02 to AC), SFB-TRR156 (B10N to AC), RTG2727 – 445549683 Innate Immune checkpoints in cancer and tissue damage (B1.2 to AC and AS), TRR179 (TP07 to AC)]; ExU 6.1.11 (to AC), German Cancer Aid grant "NK fit against AML" (73000336 to AC); a network grant of the European Commission (H2020-MSCA-MC322, ITN-765104-MATURE-NK); and by the Angioformatics platform of the European Center for Angioscience (ECAS).

Author contributions
MJ and AS designed experiments and wrote the manuscript; MJ performed experiments and analyzed the data; JXS performed experiments; CDLT performed gene expression analysis. AS and AC supervised the study.

Conflict of interests
The authors declare no competing interests.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6