Title: Bridging Theory and Experiments of Priority Effects

Authors: Heng-Xing Zou[1], Volker H. W. Rudolf[2]

Article Type: Review

Keywords: priority effects, community assembly, phenology, species interactions

List of Elements: Figure 1, Box 1-3; Supplemental Material Table S1, Table S2, Appendix I-III (.docx); Supplementary Data (.xlsx)

ORCIDs: [1] 0000-0002-1965-6910, [2] 0000-0002-9214-2000

Address: [1,2] Program in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, 6100 Main St, Houston, TX 77005

Twitter: [1] @hengxing_zou, [2] @volker_rudolf

Corresponding Author: Zou, H.-X. (hengxingzou@rice.edu)

Words in Abstract: 119

Words in Main Text: 3497

Number of References: 102

Abstract

Priority effects play a key role in structuring natural communities, but considerable confusion remains about how they affect different ecological systems. Synthesizing previous studies, we show that this confusion arises because the mechanisms driving priority and the temporal scale at which they operate differ among studies, leading to divergent outcomes in species interactions and biodiversity patterns. We suggest grouping priority effects into two functional categories based on their mechanisms: "frequency-dependent" priority effects that arise from positive frequency dependence, and "trait-dependent" priority effects that arise from time-dependent changes in interacting traits. Through easy quantification of these categories from experiments, we can construct community models representing diverse biological mechanisms and interactions with priority effects, therefore better predicting their consequences across ecosystems.

Glossary

15   **Community assembly:** Changes in species number, identity, and abundance over time in an ecological community.

   Historical contingency: Scenarios when the order and timing of any biotic or abiotic events

18   affect community assembly. Priority effects can be drivers of historical contingency, but not all historical contingencies are driven by priority effects.

   Generation time: The average time interval between an individual's birth and the birth of its

21   offspring in a population.

   Phenology: Seasonal timing of life history events, e.g., breeding, migration, germination, leaf out, and recovery from hibernation.

24   **Positive frequency dependence:** The species with a higher frequency in the community have a higher population growth rate.

   Priority effects: Scenarios where the outcomes of species interaction depend on their relative

27   arrival times or initial abundances.

   Season: A period marked by regular environmental changes that affect vital processes of species. Seasons often refer to spring, summer, fall, and winter, but they can also describe a

30   period with suitable habitats, e.g., when ephemeral ponds have water. Communities often reassemble at the beginning of the season.

Priority Effects: Definition, Importance, and Issues

33   The outcome of species interaction often depends on the temporal sequence of **community assembly** (see **Glossary**). This phenomenon, termed the "**priority effect**," is found across a wide range of taxa and ecosystems [1–4]. By altering species interactions, priority effects can

36   affect community composition [5–7], consequently driving biodiversity patterns [8,9] and ecosystem functioning [10–12]. Despite the long history of research into priority effects in community ecology, uncertainty remains about when and how they affect various ecological

39   systems [7]. Much of this uncertainty arises because priority effects are used broadly to describe any relationship between the temporal sequence of arrival orders and changes in the outcomes of interactions or community states, regardless of the underlying biology and

42   temporal scales at which the system operates. Resolving this uncertainty is not only essential to understanding the dynamics and composition of natural communities but also key to predicting how communities will change in the future as climate change reshuffles species **phenology** and

45   the timing of species interactions over **seasons** [13–15].

      Here, we review current progress in priority effects and provide a conceptual framework to understand, interpret, and predict their consequences across different systems. We first classify

48   the diverse mechanisms causing priority effects into two categories based on the temporal scale at which they occur. We then use this framework to explain current discrepancies across empirical and theoretical studies. Finally, we provide guidelines for evaluating priority effects

51   across different temporal and organizational scales.

Unpacking Priority Effects: Mechanisms and Scales

54   Ever since the introduction of the term [16], priority effects have received much attention

for their roles in shaping community assembly, such as the presence of alternative states [17,18]

and **historical contingency** [19]. While the latter is often associated with priority effects,

57   historical contingency can arise from multiple other abiotic mechanisms (e.g., dispersal,

species sorting, environmental heterogeneity, disturbance [9,20–23]), many of which can also

cause priority effects (see [7] for a thorough review). Here, instead of abiotic mechanisms of

60   priority effects, we focus on the multitude of biological causes identified by experiments across

many systems (Box 1; Table S1), which often differ in underlying mechanisms and temporal

scales at which priority effects are measured.

A Multitude of Mechanisms

The biological mechanisms of documented priority effects are often either unknown,

66   indirectly inferred, or appear specific to a given study system (Table S1). For instance, in many

microbial systems, priority effects are only measured by changes in community composition to

differences in arrival times. While some studies point to more resolved mechanisms such as

69   sugar concentration or pH [3,27], others are often attributed to general factors like resource

preemption [24] or phylogenetic relatedness [25,26], with the specific mechanisms often

unknown. In coral reef fish, priority effects often arise because early arrivers become more

72   aggressive toward late arrivers [28–30]. Intertidal invertebrates that colonize first prevent the

colonization of late-comers by physically blocking the limited substrate surface [31]. In

dragonfly and damselfly communities, priority effects can be driven by intraguild predation

75   because early arrivers gain a size advantage that allows them to prey on smaller late arrivers

[32,33]. In plants, priority effects can arise from a range of different mechanisms, including

competition for light [34] or pollinators [35], and plant-soil feedback (e.g., via shifts in

78   microbiome or allelopathy) [36]. In host-pathogen systems, priority effects between two

parasites co-infecting the same host can be mediated via an immune response of the host (e.g.,

cross-immunity) [37–39]. This multitude of mechanisms will likely lead to different

81   consequences of priority effects in specific systems.

A Dichotomy of Temporal Scales

84   Given the diversity of study systems, it is not surprising that priority effects have been

studied at different temporal scales. While roughly half of the experiments we reviewed lasted

more than five generations of their study organisms, the other half lasted for only one

87   generation or less (Box 1). The two distinctive groups reveal a clear dichotomy that largely

reflects differences in the life history of focal organisms. Studies that focused on organisms

with relatively long **generation times**, such as plants and amphibians [1,40–42], usually last

90   less than one generation. Therefore, the differences in arrival times are usually less than the

length of their full life cycle (Figure 1) and reflect species phenology (i.e., seasonal timing

within a year). Consequently, these studies largely focus on changes in traits (morphology,

93   development or survival rates, etc.) of individuals and investigate specific biological

mechanisms that cause priority effects [30,32,43]. But they do not directly measure the

long-term consequences of priority effects which can only be inferred by models of population

96   dynamics parameterized by empirical data [2,42,44–46].

On the other hand, studies on organisms with short generation times such as zooplankton

and microbes focus on population and community dynamics and thus often run for tens to

99   hundreds of generations [3,47,48]. In these studies, arrival times between species typically

differ by multiple generations, allowing them to study the long-term consequences of priority

effects on communities (e.g., alternative stable states) [9,49,50]. However, because of the fast

102   generation time, they typically do not investigate changes in traits of participating organisms

and instead focus on environmental mechanisms that promote priority effects, such as

disturbance, dispersal, resource levels, presence of predators, and pH [3,8,20,27,48].

105   Because of the dichotomy of temporal scales, both groups measure different response

variables that are recorded at different scales (individual- vs. population-level). Therefore,

patterns or mechanisms identified in studies that focus on the individual short-term (within one

108   generation) scale may not emerge or be relevant for dynamics and patterns observed in studies

on long-term community dynamics.

111   **The Gap Between Theoretical Framework and Empirical Studies**

The theory of priority effects dates back to competition models by Lotka and Volterra, who

showed that when two species in the community limit each other more than themselves, the

114   outcome of competition is determined by the relative abundance of competitors, and the system

exhibits alternative stable states [51]. Therefore, whichever species arrives earlier can gain a

numeric advantage and exclude the other, leading to different final community states. This type

117   of priority effect has also been described in the context of modern coexistence theory [52–54].

Although intuitive, this theory and its recent extensions assume that priority effects arise from a

shift between alternative stable states driven by a change in the relative numeric abundances

120   (frequencies) of species (Supplementary Material Appenidix II). This naturally raises the

question: to which empirical studies does this theory apply?

We can answer this question by focusing on the key requirements for applying this

mathematical framework to priority effects: (1) early arrival must provide the species with a numeric advantage when the interaction starts, (2) the combination of interaction coefficients needs to ensure **positive frequency dependence** to promote alternative stable states, and (3)

the inter- (and intra-) specific per-capita interactions and demographic rates of species are fixed and independent of arrival times and sequence. The last condition effectively decouples arrival time from species interactions. It highlights that timing only matters when early arrival also

ensures a numeric advantage, but the numeric advantage does not necessarily require early arrival. In fact, many studies simulate this type of priority effect without manipulating the arrival times and instead simply start experiments with an initial numeric advantage [9,55,56].

A closer inspection of empirical studies on priority effects quickly reveals that these conditions are met in some studies but not in most. Systems with short generation times such as zooplankton or microbial community are the most likely to meet these conditions because

arrival times are typically separated by multiple generations, giving the early arriver enough time to reproduce, eventually outnumbering late arrivers [3,50]. However, per-capita effects are often not quantified in these studies and still could change between arrival times, e.g., due to

habitat modification, phenotypic plasticity, or rapid evolution [27,57–59].

In contrast, most short-term studies documenting priority effects in empirical systems with longer generation times such as terrestrial plants, fish, and amphibians, do not satisfy these

141   conditions. First, differences in arrival times are often too short (less than a generation) for a numeric advantage through reproduction, and many experiments keep initial densities constant across arrival time treatments. Second, studies often report that interactions in these systems

144   are not constant but change with the timing of arrival (Table S1). Early arrival advantages caused by aggression [29,30], behavioral interference [40,60], and plant-soil feedback

[36,43,61,62] are all mediated by changes in specific traits that alter the interaction between the early and the late arriver. These changes have been further quantified in insects, vertebrates, and plants [2,42,46]. These common changes in demographic rates with relative arrival time represent a mechanism of priority effects that is fundamentally different from frequency-dependent priority effects. Traditional theory based on positive frequency dependence does not explicitly include these changes in traits, and therefore cannot be applied to interpret or predict the long-term effects of the full diversity of different priority effects in nature. This gap between theory and empirical work clearly highlights current limitations and the need for a conceptual framework that captures this diversity of biological mechanisms.

Bridging the Gap: A Unifying Framework of Priority Effects

Here, we propose a unified conceptual framework with new categorizations of priority effects to bridge the gap between theory and nature. Priority effects are defined as scenarios where the *outcomes of species interactions depend on their relative arrival times* [7]. To understand how this is possible, consider a generic two-species competition model (e.g., Beverton-Holt model; Supplementary Material Appendix II). In this system, one species can only affect the other in two ways: its density, and per-capita interaction coefficients. However, the possible equilibrium states depend only on the interaction coefficients, and only if they promote alternative stable states does the realized outcome depend on the relative frequencies of both species. Therefore, priority effects can only alter long-term outcomes when differences in arrival times also alter (1) interaction coefficients or (2) the relative frequencies of both species. Considering these facts, we suggest grouping mechanisms of priority effects into two general types: (1) "trait-dependent" priority effects which arise when differences in arrival time

result in changes in per-capita interaction coefficients (Box 2), and (2) "frequency-dependent"

priority effects (Table 1) which arise when differences in arrival time allow the early arriver to

171   increase in relative frequency, leading to a priority effect that is maintained by positive

frequency dependence. Note that the concept of trait-dependent priority effects includes

previous concepts such as size- or stage-mediated priority effects [4,32], but is intentionally

174   broader to include any possible trait that influences species interactions in the local community

and beyond [7]. Frequency-dependent priority effects represent those studied in most

traditional theories and have the same requirements (see [53] and Supplementary Material

177   Appendix II). Note that positive frequency dependence can also arise from processes other than

reproduction: In plants and slow-growing fungi, higher initial biomass often confers a higher

growth rate; empirical studies on these systems often measure biomass, not fecundity, as a

180   proxy for fitness [63,64].

Distinguishing between both types of priority effects is crucial because they have

fundamentally different effects on the dynamic of the system. Trait-dependent priority effects

183   lead to concurrent changes in key parameters that determine the possible outcome of

interactions, e.g., competing species' fitness ratio and stabilization potential (Box 2).

Depending on the original and shifted position of the interaction, this may simply mean a

186   change in the relative frequencies of coexisting species, or a dramatic shift, e.g., from stable

coexistence to competitive exclusion (Box 2). This sets them fundamentally apart from

frequency-dependent priority effects, which arise from a shift between two alternative stable

189   states (i.e., either species wins) that is only mediated by changes in their initial frequencies.

Unlike frequency-dependent priority effects, trait-dependent priority effects are possible for

any combination of per-capita interactions, and they can arise within a generation and therefore

192 help explain priority effects in empirical systems with longer generation times where arrival

times are separated by less than a generation.

 Note that mechanisms leading to frequency- and trait-dependent priority effects can

195 co-occur in the same system but operate at different time scales. For example, different

biological mechanisms of resource competition can lead to either frequency- or trait-dependent

priority effects (Box 3). In another example, the early arriving bacteria in flower nectar lowers

198 the pH of the environment, impeding successful colonization by yeast [27]. This can be

categorized as trait-dependent priority effect because arriving late changes the yeast's

demographic rates. However, on a larger temporal scale, the pH alteration can further promote

201 bacterial growth, resulting in positive frequency dependence.

 The concepts of niche preemption and niche modification (*sensu* [7]) are broad

generalizations of specific mechanisms that can be categorized and modeled as frequency- or

204 trait-dependent priority effects. In the previous example, because the relative arrival times

change the demographic rates of the yeast, the priority effect arise from niche modification and

is trait-dependent. Within the microbiome hosted by a single polyp of *Hydra vulgaris*, one

207 strain of bacteria gains dominance by a higher initial abundance [65]; this is an example of

niche preemption but the underlying mechanism is positive frequency-dependent. When fish

first occupy a nesting site in coral reef and defend late colonizers [28], the priority effect again

210 matches niche preemption but the driving mechanism is trait-dependent because the order of

arrival changes behaviors, and the outcomes also depend on traits like body conditions [30].

 Our framework could even be extended to evolutionary time scales. If one species arrives

213 sufficiently early and adapts to local conditions, it could dominate the habitat and prevent other

species or even ancestral conspecifics from colonizing, i.e., the monopolization effect

[57,66–69]. At extremely large temporal scales, evolution can lead to a "phylogenetic priority

216   effect": early-arriving taxa could diversify and dominate in the local community [70–72].

These evolutionary priority effects can again arise from a combination of frequency- and

trait-dependent mechanisms: during the adaptation process, the early colonizer can also modify

219   the environment to lower the fitness of late colonizers (trait-dependent) or grow in large

populations to achieve positive frequency dependence.

222   **Why It Matters**

Our framework bridges the gap between theory and empirical work by providing a

straightforward pathway to integrate a diversity of priority effects observed in natural systems

225   into theoretical models (Box 2). This differentiation also allows researchers to critically

evaluate the role of priority effects across a wider range of natural systems, reconciling the

dichotomy between experiments on systems operating on different time scales (long- vs. short

228   generation times; Box 1). Trait-dependent priority effects expand existing theory by including

less-discussed biological mechanisms, such as evolutionary changes in species traits and fitness

[57,59], and facilitative effects from early arrivers [73,74]. The latter is especially important

231   because frequency-dependent priority effects, by definition, are only present in competitive

interactions, and therefore cannot characterize the widespread positive interactions that could

change with relative arrival times among species.

234   Furthermore, differentiating between the mechanisms of priority effects is essential to

understanding the dynamics of natural systems and how they will respond to changes in the

temporal structure (e.g., phenological shifts) or other types of perturbations. We already

237   outlined how the type of mechanisms determines the consequences of shifts in arrival time for

local populations, but these differences in local dynamics also scale up to drive differences in regional patterns. For instance, trait-dependent priority effects can limit the homogenizing

240 effect of high dispersal in metacommunities because competitive hierarchies depend on species' arrival times and not relative frequencies [8,75,76]. The mechanism of priority effects could also determine management decisions. For example, restoring a system that exhibits alternative

243 stable states due to frequency-dependent priority effects might require manipulations of relative frequencies of species, while restoring a system driven by trait-dependent priority effects might require manipulations of actual timing [73,77]. Our categorization provides a more applicable

246 theoretical framework of priority effects to empirical scenarios.

Looking Forward

249 *Realistic Models of Priority Effects*

Our classification framework emphasizes the importance of biological realism in research on priority effects, especially in theoretical studies. Models that do not capture the key

252 mechanisms driving population and community dynamics may not accurately predict how they change over time or respond to perturbations. Thus, a realistic model of priority effects should include two key components: a base demographic model, and a function that describes how

255 interacting species traits change with relative arrival times [15]. This integration captures the diversity of priority effects in natural systems and helps to isolate individual and synergistic effects of arrival times within and across generations.

258 This integration comes with new opportunities and challenges. Different mechanisms typically operate at different time scales (e.g., within vs. between generations). How models capture these time scales will ultimately depend on the life history of species, which is directly

261   related to the temporal scales of arrival time differences (Box 1). Previous studies suggest an

easy implementation in annual systems by modeling interaction coefficients as a function of

arrival times (e.g., the interaction-phenology function [15]; Box 2), which allows researchers to

264   maintain much of the original model structure and analytical tools for analyses. However, this

approach does not easily extend to systems where interactions last over multiple generations

within a year. These systems require different model formulations, like size/stage-structured

267   models or agent-based models [78]. In addition, variations in arrival times are widespread in

nature (e.g., between years and patches [14,79]), and individual and combined effects of

different types of priority effects could lead to different community dynamics and biodiversity

270   patterns [21,75,76]. We hope that our framework will stimulate a range of new theoretical

studies that will shed new light on the different consequences of priority effects across

biological systems.

Better Quantification of Different Priority Effects

       Traditionally, studies focused on the detection of priority effects by comparing dynamics

276   that arise from different arrival orders (e.g., early vs. late treatment; Box 1). However, this

simple approach is unreliable to detect the potential for priority effects. For instance, with

frequency-dependent priority effects, whether the early species excludes the later arriver

279   depends on whether its initial population is large enough or has enough time to increase

relative frequency to gain a sufficient numeric advantage (Box 3). Therefore, the community

dynamics depend not only on the arrival *order* but also on the arrival *time*.

282   To overcome these issues, future experimental research should aim to characterize priority

effects across a wider range of arrival times to quantify how interactions or outcomes scale

with relative arrival time (e.g., interaction-phenology function; Box 2). This would allow the

285 experiment to uncover an accurate representation of priority effects in the focal system even

when knowing the biological mechanism is difficult. The range of arrival times should scale

with the life history of species to capture differences in arrival times that cover at least one

288 generation (or length of a life stage). Relative arrival times can either be directly manipulated

or simulated by competing different combinations of stages/size classes of species [2,32,42].

The effect of timing can then be measured by vital rates such as survival, growth rate, and

291 fecundity. For systems without apparent size- or stage-mediated competition (e.g., bacteria,

single-cell fungi, protists), interspecific competition coefficients can be directly evaluated from

invasion growth rates [3,80]. Differences in arrival times may nevertheless allow the early

294 arriver to establish a numeric advantage in these systems, potentially requiring crossing

different initial frequencies of each species to the gradient of arrival times. The quantified

interaction-phenology relationship can then be used to model long-term community dynamics

297 under other conditions.

Although the interaction-phenology function is a powerful tool to characterize priority

effects, it is phenomenological and cannot detect the potential underlying biological

300 mechanism, such as size-mediated predation, resource competition, or environmental factors.

Additional measurements on other parameters, such as functional traits, demographic

parameters, or behavioral assays are required to identify the exact mechanism of priority

303 effects in a system. Different arrival times are also likely to influence intraspecific competition

in some systems with distinct ontogeny [81–83]; therefore, characterizing the relationship

between intraspecific competition and arrival time might also be necessary.

Concluding Remarks

Priority effects represent a time-explicit understanding of community dynamics, which

309   remains one of the understudied frontiers in community ecology [15,84]. We show that by

accounting for the specific mechanisms that link the relative arrival times of species to their

interactions, we can extend our current theoretical framework of priority effects to cover a

312   wider range of taxa and ecological conditions. Our framework provides new paths for

modeling priority effects and highlights the need for new experimental designs that manipulate

arrival times at finer temporal scales. We hope that this framework provides new perspectives

315   on classic themes of community assembly (e.g., alternative assembly paths [85,86], repeated

colonization [86,87], regional species and trait compositions [88,89], and environmental

heterogeneity [90,91]), and marks the first step in considering timing in classic and burgeoning

318   concepts in ecology (e.g., phenotypic plasticity [58,92,93], metacommunity [21,75,76], and

higher-order interactions [94,95]; **Outstanding Questions**). Resolving and predicting species

interactions in a temporally explicit context will be a powerful tool to decipher how natural

321   communities respond to widespread phenological shifts under climate change.

Acknowledgments

324   We thank Joshua Fowler, Tadashi Fukami, and Hannah Yin for their comments and

discussions on the manuscript. We thank members of the sPriority working group funded by

the Synthesis Center of the German Research Centre for Integrative Biodiversity Research

327   (iDiv, Germany) for sharing their compiled literature of experiments on priority effects.

Funding was provided by NSF DEB-1655626.

Figure Legends

Figure 1. A framework of trait- and frequency-dependent priority effects. (A)

Trait-dependent priority effects often emerge from differences in arrival times within one

333   generation, while frequency-dependent priority effects often emerge from a numeric advantage

of the early species, established by arriving multiple generations earlier. (B) – (C) With

trait-dependent priority effects, the timing of species interactions affects their interaction

336   strengths, visualized by the relationship between per-capita interspecific interaction coefficients

and relative arrival times. The relationship between interactions and relative arrival times is

typically either continuously nonlinear or in the extreme case piecewise (see Box 2 for

339   examples). With frequency-dependent priority effects, per-capita interactions between species

are not affected by their relative arrival times. (D) Trait-dependent priority effects allow for

shifts between all possible outcomes of interactions, while frequency-dependent priority effects

342   only lead to shifts between to two alternative states (e.g., which species eliminates the other,

Box 2).

Box 1. Empirical studies of priority effects: a literature review

We compiled 268 empirical studies of priority effects (see Supplementary Material Appendix I for methods). Despite the diverse study systems in our list of experiments, most studies were conducted on terrestrial plants (79, or 29.5%). Among all experiments, 46 (17.1%) did not explicitly manipulate arrival times but rather used initial community composition or length of natural colonization as proxies. Even among the remaining 222 studies that considered arrival time as a treatment, only 58 used a temporal gradient rather than a simple early/late/simultaneous arrival treatment (Figure IA).

We further calculated the proportion of studies with durations of less or equal to one generation (defined by the emergence of adults or reproduction within the experimental period), 2-5 generations, or more than five generations. We omitted 11 studies using taxa that are significantly different in life histories (e.g., insects and viruses). 109 of the remaining 257 studies (42.4%) were conducted over a time span of less than or equal to one generation and 121 studies (47.1%) lasted more than five generations of focal taxa (56, 41.2%). These results represented the slow- and fast-generating systems often used in experiments on priority effects. Indeed, studies spanning less than five generations (126, 49.0%) were conducted on mycorrhizal/wood-decomposing fungi, terrestrial/aquatic plants, amphibians, fish, and slow-generating insects (e.g., odonates), all of which require weeks to years for development and reproduction. In contrast, most studies spanning more than five generations were conducted on fast-generating microbial communities (bacteria and yeast), viruses, nematodes, trematodes, protists, phytoplankton, and zooplankton, with a small fraction of long-term studies on terrestrial plant communities (11, 4.3%). These results clearly indicate that the current body of experimental work on priority effects is strongly biased towards two extremes:

short-term studies on systems with long generation times and long-term studies on systems

369 with very short generation times (Figure IB).

In addition to system selection and experimental design, priority effects are also measured in diverse ways. The most common response variables include community composition [49],

372 functional traits [25,26], demographic rates such as growth, survival, and fecundity [2,42,61], and ecosystem function [96,97], but studies can also measure highly specific responses such as aggression, site occupancy [28,30,98], and in coinfection experiments, pathogen load, and host

375 response [38,99]. These measurements in part highlight the importance of priority effects in nature but also increase the difficulty of generalizing, quantifying, and comparing the consequences of priority effects across systems.

Figure I. Manipulation of time in experiments and the dichotomy of time scales of experimental systems. (A) Results from the literature review show that 46 out of 268

381 experiments (17.2%) did not manipulate the arrival times of species. (B) Results from the literature review show the dichotomy of temporal scales among experiments: almost all studies spanning less than five generations were conducted on slow-generating systems (e.g., plants,

384 amphibians, fish), while most studies spanning more than five generations were conducted on fast-generating systems (e.g., microbial communities, protists, planktons), with a small fraction of long-term studies on terrestrial plant communities. (C) Taxonomic breakdown of studies in

387 (B). Crustaceans include zooplankton and insects.

Box 2. Mathematical representations of trait-dependent priority effects

Trait-dependent priority effects assume a change in species interaction with their timing. This is often modeled by a shift in interspecific competition coefficients α_{12} and α_{21}, although other demographic rates may also contribute to this change. For example, consider a Beverton-Holt model where competition between two species is defined as a function of the difference between their arrival times (phenology), Δp_{12}:

$$\frac{N_1(t+1)}{N_1(t)} = \frac{\lambda_1}{1 + \alpha_{11}N_1(t) + f_{12}(\Delta p_{12})N_2(t)}$$

$$\frac{N_2(t+1)}{N_2(t)} = \frac{\lambda_2}{1 + \alpha_{22}N_2(t) + f_{21}(\Delta p_{21})N_1(t)}$$

(Eqn. B2). Here, changes in the density (N_i) between years depend on the intrinsic growth rate (λ) of a species which in turn is reduced by intra- and interspecific competition.

The key to the difference between frequency- and trait-dependent priority effects is termed the interaction-phenology function, $\alpha_{12} = f_{12}(\Delta p_{12})$, because it links per capita interaction strengths directly to their relative (typically seasonal) timing of arrival within a generation [15]. The shape of $f(\Delta p_{ij})$ depends on the specific system; experiments often suggest a nonlinear relationship [2,42,46], but $f(\Delta p_{ij})$ can be piecewise if species interaction only depends on whether another species is previously present, not how long it has been present (e.g., patch memory [75]; Figure 1). When the function is constant ($f_{12}(\Delta p_{12}) = \alpha_{12}$ for all Δp_{12}), species interactions are unrelated to relative arrival times, and the model collapses to frequency-dependent priority effects (Figure 1). By accommodating a wide range of interaction-phenology relationships, this method of modeling priority effects is versatile and can be used to parameterize any empirical system.

Because both the fitness ratio and stabilization potential (*sensu* [53]) depend on

408   interspecific competition coefficients, they also become functions of Δp_{12}:

$$\frac{\kappa_2}{\kappa_1} = \frac{\lambda_2 - 1}{\lambda_1 - 1} \sqrt{\frac{\alpha_{11} f_{12}(\Delta p_{12})}{\alpha_{22} f_{21}(\Delta p_{21})}}$$

$$1 - \rho = 1 - \sqrt{\frac{f_{12}(\Delta p_{12}) f_{21}(\Delta p_{21})}{\alpha_{11}\alpha_{22}}}$$

(Eqn. B3)

Therefore, any change in the relative timing of species would shift the position of the

411   species pair in the coexistence space. For instance, when $\Delta p_{12} = 0$ species 1 and 2 could

coexist (point D in Figure II), but when species 1 arrives earlier ($\Delta p_{12} > 0$) the corresponding

fitness ratio and stabilization potential may shift to point C, or species 1 wins. Note that these

414   shifts may not necessarily lead to observable changes in competition outcomes, and the nature

of these shifts depends on the working definition of priority effects: the arrival order changes

either the outcome of species interaction (by which it would not be a trait-dependent priority

417   effect) or the species interaction itself (by which it would). Note that $f\left(\Delta p_{ij}\right)$ can easily be

applied to coefficients in models with other types of interactions, such as facilitation and

mutualism, to calculate the corresponding fitness ratio and stabilizing potential [100,101].

Figure I. Frequency- and trait-dependent priority effects in a coexistence space. The

coexistence space is often used to map species interactions with measurements of stabilizing

423   mechanism (stabilizing potential on the x-axis) and equalizing mechanisms (fitness ratio on the

y-axis). Positive frequency dependence (PFD) is represented by a region where $\rho > f_2/f_1 >$

$1/\rho$ (point A). Trait-dependent priority effects are defined by changes in interaction strengths

426   due to changes in arrival times, represented by arrows. Note that a trait-dependent priority

effect does not necessarily lead to exclusion; a shift from exclusion to coexistence (points C to D), or in relative abundance of species is also a trait-dependent priority effect.

Box 3. Frequency- and trait-dependent priority effects from resource competition

Here, we illustrate the utility of our framework by unpacking a common mechanism of priority effects, resource competition [7]. In this system, the early arrival advantage could result from the higher abundance and subsequent population growth by positive frequency dependence [65]. However, the early arrival advantage could also arise from changes in the traits of individuals. For instance, the early arriver might have a larger size compared to the late arriver. This size advantage could lead to higher feeding rates in amphibians [40,102] or an advantage in light competition in plants [34].

To quantitatively capture the above examples, we analyze a modified, discrete-time version of the consumer-resource model used by Ke and Letten [53], where two consumer species compete for a shared resource. We model consumers with an annual cycle, and allow arrival times to vary within a year between species. We assume a higher feeding rate of the earlier arriver within a year. Therefore, the early arriving advantage in this model could arise from: (1) resource preemption by the early arriver, which is a numeric mechanism that leads to positive frequency dependence; (2) change in resource uptake rates, which is a trait-dependent mechanism. See Appendix III and Table S2 for model and simulation details.

When resource uptake rates are held constant, the two species display positive frequency dependence: a species wins if it arrives early over several years (Figure IA), regardless of the within-year arrival times (phenology) of the two species (Figure IC). However, when the early arriver within the year has a higher resource uptake rate, both arrival times within and across years matter: while one species could gain an advantage by establishing itself several years earlier than the other, it could still be excluded if it emerges late within a year (Figure IB, ID). These results highlight that: (1) one common biological mechanism of priority effects, resource

453   competition, can be driven by either frequency- or trait-dependent mechanism, or a

combination of both; (2) the two types of priority are not mutually exclusive, but operate on

different time scales; (3) in nature, population dynamics could be affected by arrival times both

456   within a year and across years.

Figure I. A model with both frequency- and trait-dependent priority effects. (A) Population

459   dynamics when species 1 establishes several years earlier than species 2; species 1 has a higher

density, showing a frequency-dependent priority effect. Vertical dash line indicates when

species 2 arrives. (B) Population dynamics when species 1 establishes several years earlier than

462   species 2 but emerges later than species 2 within a year; despite species 1's

frequency-dependent priority effect, species 2 has a higher feeding rate by emerging earlier

within a year, displaying a trait-dependent priority effect over species 1. (C) Without

465   trait-dependent priority effects, the relative emergence time within a year (Δp) does not affect

competition outcomes. (D) With both frequency- and trait-dependent priority effects, both the

relative arrival time over years (years early on the y-axis) and within a year (Δp on the x-axis)

468   affect competition outcomes.

Reference

1. Raczyński, M. *et al.* (2021) Size-mediated priority effects are trait-dependent and consistent across latitudes in a damselfly. *Oikos* 130, 1535–1547
2. Blackford, C. *et al.* (2020) Species Differences in Phenology Shape Coexistence. *Am. Nat.* 195, E168–E180
3. Grainger, T.N. *et al.* (2019) Applying modern coexistence theory to priority effects. *Proc. Natl. Acad. Sci. U. S. A.* 116, 6205–6210
4. Carter, S.K. and Rudolf, V.H.W. (2019) Shifts in phenological mean and synchrony interact to shape competitive outcomes. *Ecology* 100, 1–11
5. Holditch, Z. and Smith, A.D. (2020) Priority determines Tribolium competitive outcome in a food-limited environment. *PLoS ONE* 15, 1–14
6. Fragata, I. *et al.* (2022) Specific sequence of arrival promotes coexistence via spatial niche pre-emption by the weak competitor. *Ecol. Lett.* 25, 1629–1639
7. Fukami, T. (2015) Historical contingency in community assembly: integrating niches, species pools, and priority effects*Annual Review of Ecology, Evolution, and Systematics*, 461–23
8. Pu, Z. and Jiang, L. (2015) Dispersal among local communities does not reduce historical contingencies during metacommunity assembly. *Oikos* 124, 1327–1336
9. Toju, H. *et al.* (2018) Priority effects can persist across floral generations in nectar microbial metacommunities. *Oikos* 127, 345–352
10. Weidlich, E.W.A. *et al.* (2018) Priority effects caused by plant order of arrival affect below-ground productivity. *J. Ecol.* DOI: 10.1111/1365-2745.12829
11. Bittleston, L.S. *et al.* (2020) Context-dependent dynamics lead to the assembly of functionally distinct microbial communities. *Nat. Commun.* 11, 1–10
12. Catano, C.P. *et al.* (2023) Community assembly history alters relationships between biodiversity and ecosystem functions during restoration. *Ecology* 104, e3910
13. Kharouba, H.M. *et al.* (2018) Global shifts in the phenological synchrony of species interactions over recent decades. *Proc. Natl. Acad. Sci. U. S. A.* 115, 5211–5216
14. Carter, S.K. *et al.* (2018) Shifts in phenological distributions reshape interaction potential in natural communities. *Ecol. Lett.* 21, 1143–1151
15. Rudolf, V.H.W. (2019) The role of seasonal timing and phenological shifts for species coexistence*Ecology Letters*, 221324–1338
16. Slatkin, M. (1974) Competition and Regional Coexistence. *Ecology* 55, 128–134
17. Palmgren, A. (1926) Chance as an element in plant geography. *Proc. Int. Congr. Plant Sci.*
18. Sutherland, J.P. (1974) Multiple stable points in natural communities. *Am. Nat.* DOI: 10.1086/282961
19. Gause, G.F. (1934) *The Struggle for Existence: A Classic of Mathematical Biology and Ecology*, Courier Dover Publications
20. Chase, J.M. (2007) Drought mediates the importance of stochastic community assembly. *Proc. Natl. Acad. Sci.* 104, 17430–17434
21. Lu, M. (2021) Complex relationships between beta diversity and dispersal in meta-community models. *Ecography* 44, 1769–1780
22. Vannette, R.L. and Fukami, T. (2017) Dispersal enhances beta diversity in nectar microbes. *Ecol. Lett.* 20, 901–910
23. Shurin, J.B. *et al.* (2004) Alternative stable states and regional community structure. *J.*

Theor. Biol. 227, 359–368

24. Vannette, R.L. and Fukami, T. (2014) Historical contingency in species interactions: Towards niche-based predictions. *Ecol. Lett.* 17, 115–124

25. Tan, J. *et al.* (2012) Species phylogenetic relatedness, priority effects, and ecosystem functioning. *Ecology* 93, 1164–1172

26. Peay, K.G. *et al.* (2012) Phylogenetic relatedness predicts priority effects in nectar yeast communities. *Proc. R. Soc. B Biol. Sci.* DOI: 10.1098/rspb.2011.1230

27. Chappell, C.R. *et al.* (2022) Wide-ranging consequences of priority effects governed by an overarching factor. *eLife* 11, e79647

28. Almany, G.R. (2003) Priority effects in coral reef fish communities. *Ecology* DOI: 10.1890/0012-9658(2003)084[1920:PEICRF]2.0.CO;2

29. Geange, S.W. and Stier, A.C. (2009) Order of arrival affects competition in two reef fishes. *Ecology* DOI: 10.1890/08-0630.1

30. Poulos, D.E. and McCormick, M.I. (2015) Asymmetries in body condition and order of arrival influence competitive ability and survival in a coral reef fish. *Oecologia* 179, 719–728

31. Vieira, E.A. *et al.* (2018) Persistence and space preemption explain species-specific founder effects on the organization of marine sessile communities. *Ecol. Evol.* 8, 3430–3442

32. Rasmussen, N.L. *et al.* (2014) Linking phenological shifts to species interactions through size-mediated priority effects*Journal of Animal Ecology*, 831206–1215

33. Sniegula, S. *et al.* (2019) Size-mediated priority and temperature effects on intra-cohort competition and cannibalism in a damselfly. *J. Anim. Ecol.* 88, 637–648

34. D'Antonio, C.M. *et al.* (2001) Factors Influencing Dynamics of Two Invasive C4 Grasses in Seasonally Dry Hawaiian Woodlands. *Ecology* 82, 89–104

35. Waters, S.M. *et al.* (2020) Experimental shifts in exotic flowering phenology produce strong indirect effects on native plant reproductive success. *J. Ecol.* DOI: 10.1111/1365-2745.13392

36. van der Putten, W.H. *et al.* (2016) Where, when and how plant–soil feedback matters in a changing world. *Funct. Ecol.* 30, 1109–1121

37. Hoverman, J.T. *et al.* (2013) Does timing matter? How priority effects influence the outcome of parasite interactions within hosts. *Oecologia* DOI: 10.1007/s00442-013-2692-x

38. Devevey, G. *et al.* (2015) First arrived takes all: Inhibitory priority effects dominate competition between co-infecting Borrelia burgdorferi strains Ecological and evolutionary microbiology. *BMC Microbiol.* DOI: 10.1186/s12866-015-0381-0

39. Clay, P.A. *et al.* (2019) Within-host priority effects systematically alter pathogen coexistence. *Am. Nat.* DOI: 10.1086/701126

40. Alford, R.A. and Wilbur, H.M. (1985) Priority effects in experimental pond communities: competition between Bufo and Rana. *Ecology* DOI: 10.2307/1939161

41. Kardol, P. *et al.* (2013) Resource availability mediates the importance of priority effects in plant community assembly and ecosystem function*Oikos*, 12284–94

42. Rudolf, V.H.W. (2018) Nonlinear effects of phenological shifts link interannual variation to species interactions*Journal of Animal Ecology*, 871395–1406

43. Kardol, P. *et al.* (2007) Microbe-mediated plant--soil feedback causes historical contingency effects in plant community assembly*Ecological monographs*, 77147–162

44. Uricchio, L.H. *et al.* (2019) Priority effects and nonhierarchical competition shape species

composition in a complex grassland community. *Am. Nat.* 193, 213–226

45. Song, C. *et al.* (2020) Disentangling the effects of external perturbations on coexistence and priority effects. *J. Ecol.* 108, 1677–1689

46. Shorrocks, B. and Bingley, M. (1994) Priority effects and species coexistence: Experiments with fungal-breeding Drosophila. *J. Anim. Ecol.* 63, 799–806

47. Louette, G. and De Meester, L. (2007) Predation and priority effects in experimental zooplankton communities. *Oikos* 116, 419–426

48. Olito, C. and Fukami, T. (2009) Long-term effects of predator arrival timing on prey community succession. *Am. Nat.* 173, 354–362

49. Drake, J.A. (1991) Community-assembly mechanics and the structure of an experimental species ensemble. *Am. Nat.* 137, 1–26

50. Fukami, T. (2004) Assembly history interacts with ecosystem size to influence species diversity. *Ecology* 85, 3234–3242

51. May, R.M. (1971) Stability in multispecies community models. *Math. Biosci.* DOI: 10.1016/0025-5564(71)90074-5

52. Fukami, T. *et al.* (2016) A framework for priority effects. *J. Veg. Sci.* 27, 655–657

53. Ke, P.J. and Letten, A.D. (2018) Coexistence theory and the frequency-dependence of priority effects. *Nat. Ecol. Evol.* 2, 1691–1695

54. Chesson, P. (2018) Updates on mechanisms on maintenance of species diversity*Journal of Ecology*, 1061773–1794

55. Ladd, B. and Facelli, J.M. (2008) Priority effects produced by plant litter result in non-additive competitive effects. *Oecologia* 157, 687–696

56. Livingston, G. *et al.* (2013) The dynamics of community assembly under sudden mixing in experimental microcosms. *Ecology* 94, 2898–2906

57. Nadeau, C.P. *et al.* (2021) Adaptation reduces competitive dominance and alters community assembly. *Proc. R. Soc. B Biol. Sci.* 288, 20203133

58. Hess, C. *et al.* (2022) Phenotypic plasticity promotes species coexistence. *Nat. Ecol. Evol.* 6, 1256–1261

59. Yamamichi, M. *et al.* (2022) Integrating eco-evolutionary dynamics and modern coexistence theory. *Ecol. Lett.* 25, 2091–2106

60. Resetarits Jr., W.J. and Wilbur, H.M. (1989) Choice of Oviposition Site by Hyla Chrysoscelis: Role of Predators and Competitors. *Ecology* 70, 220–228

61. Grman, E. and Suding, K.N. (2010) Within-Year Soil Legacies Contribute to Strong Priority Effects of Exotics on Native California Grassland Communities. *Restor. Ecol.* 18, 664–670

62. Zhang, Z. *et al.* (2020) Soil-microorganism-mediated invasional meltdown in plants. *Nat. Ecol. Evol.* DOI: 10.1038/s41559-020-01311-0

63. Tilman, D. (1982) Resource competition and community structure. *Monogr. Popul. Biol.* DOI: 10.2307/4549

64. Leopold, D.R. *et al.* (2017) Priority effects are interactively regulated by top-down and bottom-up forces: evidence from wood decomposer communities. *Ecol. Lett.* 20, 1054–1063

65. Wein, T. *et al.* (2018) Carrying Capacity and Colonization Dynamics of Curvibacter in the Hydra Host Habitat. *Front. Microbiol.* 9

66. Urban, M.C. and De Meester, L. (2009) Community monopolization: local adaptation enhances priority effects in an evolving metacommunity. *Proc. R. Soc. B Biol. Sci.* 276,

606   4129–4138

67. De Meester, L. *et al.* (2016) Evolving Perspectives on Monopolization and Priority Effects. *Trends Ecol. Evol.* 31, 136–146

68. Zee, P.C. and Fukami, T. (2018) Priority effects are weakened by a short, but not long, history of sympatric evolution. *Proc. R. Soc. B Biol. Sci.* 285

69. Leibold, M.A. *et al.* (2019) Regional neutrality evolves through local adaptive niche evolution. *Proc. Natl. Acad. Sci. U. S. A.* 116, 2612–2617

70. Dupuis, L.V. *et al.* (2015) Littoral cladoceran community reassembly following the cessation of disturbance. *J. Paleolimnol.* 54, 121–135

71. Leopold, D.R. *et al.* (2015) Evolutionary priority effects in New Zealand alpine plants across environmental gradients. *J. Biogeogr.* 42, 729–737

72. Brandt, A.J. *et al.* (2016) Precipitation alters the strength of evolutionary priority effects in forest community assembly of pteridophytes and angiosperms. *J. Ecol.* 104, 1673–1681

73. Weidlich, E.W.A. *et al.* (2021) Priority effects and ecological restoration. *Restor. Ecol.* 29, e13317

74. Halliday, F.W. *et al.* (2020) Facilitative priority effects drive parasite assembly under coinfection. *Nat. Ecol. Evol.* 4, 1510–1521

75. Miller, Z.R. and Allesina, S. (2021) Metapopulations with habitat modification. *Proc. Natl. Acad. Sci.* 118

76. Zou, H.-X. and Rudolf, V.H.W. (2023) Priority effects determine how dispersal affects biodiversity in seasonal metacommunities. *Am. Nat.* DOI: 10.1086/725039

77. Young, T.P. *et al.* (2017) Using priority effects to manipulate competitive relationships in restoration. *Restor. Ecol.* 25, S114–S123

78. Carter, S.K. and Rudolf, V.H.W. (2022) Exploring conditions that strengthen or weaken the ecological and evolutionary consequences of phenological synchrony. *Am. Nat.* DOI: 10.1086/720899

79. Theobald, E.J. *et al.* (2017) Climate drives phenological reassembly of a mountain wildflower meadow community. *Ecology* 98, 2799–2812

80. Godwin, C.M. *et al.* (2020) An empiricist's guide to modern coexistence theory for competitive communities. *Oikos* DOI: 10.1111/oik.06957

81. Miller, T.E.X. and Rudolf, V.H.W. (2011) Thinking inside the box: Community-level consequences of stage-structured populations. *Trends Ecol. Evol.* 26, 457–466

82. Rudolf, V.H.W. *et al.* (2014) Resolving the roles of body size and species identity in driving functional diversity. *Proc. R. Soc. B Biol. Sci.* 281

83. Murillo-Rincón, A.P. *et al.* (2017) Intraspecific priority effects modify compensatory responses to changes in hatching phenology in an amphibian. *J. Anim. Ecol.* 86, 128–135

84. White, E.R. and Hastings, A. (2020) Seasonality in ecology: Progress and prospects in theory. *Ecol. Complex.* 44, 100867

85. Fukami, T. and Nakajima, M. (2011) Community assembly: Alternative stable states or alternative transient states? *Ecol. Lett.* 14, 973–984

86. Song, C. *et al.* (2021) Untangling the complexity of priority effects in multispecies communities. *Ecol. Lett.* 24, 2301–2313

87. Robinson, J.F. and Dickerson, J.E. (1987) Does Invasion Sequence Affect Community Structure? *Ecology* 68, 587–595

88. Fukami, T. *et al.* (2005) Species divergence and trait convergence in experimental plant community assembly. *Ecol. Lett.* 8, 1283–1290

89. Spasojevic, M.J. *et al.* (2018) Integrating species traits into species pools. *Ecology* 99, 1265–1276

90. Daniel, J. *et al.* (2019) Stochastic and deterministic processes drive wetland community assembly across a gradient of environmental filtering. *Oikos* 128, 1158–1169

91. Little, C.J. and Altermatt, F. (2018) Do priority effects outweigh environmental filtering in a guild of dominant freshwater macroinvertebrates? *Proc. R. Soc. B Biol. Sci.* 285

92. Hausch, S. *et al.* (2018) Effects of intraspecific phenotypic variation on species coexistence. *Ecology* 99, 1453–1462

93. Pérez-Ramos, I.M. *et al.* (2019) Functional traits and phenotypic plasticity modulate species coexistence across contrasting climatic conditions. *Nat. Commun.* 10, 2555

94. Kleinhesselink, A.R. *et al.* (2022) Detecting and interpreting higher-order interactions in ecological communities. *Ecol. Lett.* 25, 1604–1617

95. Gibbs, T. *et al.* (2022) Coexistence in diverse communities with higher-order interactions. *Proc. Natl. Acad. Sci.* 119, e2205063119

96. Fukami, T. *et al.* (2010) Assembly history dictates ecosystem functioning: Evidence from wood decomposer communities. *Ecol. Lett.* DOI: 10.1111/j.1461-0248.2010.01465.x

97. Ploughe, L.W. *et al.* (2020) Priority effects: How the order of arrival of an invasive grass, Bromus tectorum, alters productivity and plant community structure when grown with native grass species. *Ecol. Evol.* 10, 13173–13181

98. Shulman, M.J. *et al.* (1983) Priority effects in the recruitment of juvenile coral reef fishes. *Ecology* DOI: 10.2307/1937505

99. Natsopoulou, M.E. *et al.* (2015) Interspecific competition in honeybee intracellular gut parasites is asymmetric and favours the spread of an emerging infectious disease. *Proc. R. Soc. B Biol. Sci.* 282, 20141896

100. Koffel, T. *et al.* (2021) From competition to facilitation and mutualism: a general theory of the niche. *Ecol. Monogr.* 91, e01458

101. Spaak, J.W. *et al.* (2021) Mapping species niche and fitness differences for communities with multiple interaction types. *Oikos* 130, 2065–2077

102. Lawler, S.P. and Morin, P.J. (1993) Temporal overlap, competition, and priority effects in larval anurans. *Ecology* 74, 174–182

	Trait-dependent	**Frequency-dependent**

A. Scales of relative arrival time vs. generation time

Relative arrival times usually within one generation

Relative arrival times usually longer than one generation

B. Timing affects...

Interaction strengths

Population size

C. How per-capita effects change with relative arrival times

D. Potential outcomes of interaction (see Box 2)

Exclusion, coexistence, or alternative stable states; may include shifts in relative abundances

Outcomes change continuously along the gradient of relative arrival times

Shifts between alternative stable states via positive frequency dependence

Outcomes change abruptly depending only on the arrival order