**Interactions between *Mycoplasma mycoides mycoides* and bovine macrophages under**
**physiological conditions**

Philippe totté*[1], Tiffany Bonnefois[1], Lucia Manso-Silvan[1]

[1] ASTRE, CIRAD, INRAE, Univ Montpellier – Montpellier, France

*Corresponding author
Correspondence: philippe.totte@cirad.fr

**ABSTRACT**
We investigated the interactions of unopsonized and opsonized *Mycoplasma mycoides* subsp.
*mycoides* (*Mmm*) with bovine macrophages *in vitro*. *Mmm* survived and proliferated extracellularly on
bovine macrophage cell layers in the absence of *Mmm*-specific antisera. Bovine complement used at
non-bactericidal concentrations did neither have opsonizing effect nor promoted intracellular
survival, whereas *Mmm*-specific antisera substantially increased phagocytosis and *Mmm* killing. A
phagocytosis-independent uptake of *Mmm* by macrophages occurred at high MOIs, also found to
induce the production of TNF, and both responses were unaffected by non-bactericidal doses of
bovine complement. Bovine complement used at higher doses killed *Mmm* in cell-free cultures and
completely abrogated TNF responses by macrophages. These results provide a framework to identify
*Mmm* antigens involved in interactions with macrophages and targeted by potentially protective
antibodies and point towards a pivotal role of complement in the control of inflammatory responses
in contagious bovine pleuropneumonia.

*Keywords:* Contagious bovine pleuropneumonia, *Mycoplasma mycoides*, macrophages, complement,
TNF

## Introduction

Mycoplasma mycoides subsp. mycoides (*Mmm*) is a wall-less bacterium and the causative agent of contagious bovine pleuropneumonia (CBPP), a highly contagious respiratory disease of cattle notifiable to the World Organization for Animal Health (Woha, 2015). CBPP lung lesions and chronic carrier state are suggestive of an uncontrolled inflammatory response by the host's immune system and immune evasion mechanisms developed by *Mmm* respectively (Provost et al., 1987). Although macrophages patrolling the surface of respiratory mucosae represent the first line of defense against pathogens invading the lung, little is known of the interactions of these cells with *Mmm*. Macrophages containing *Mmm* antigens are found in the lungs of infected animals (Scanziani et al., 1997) and they produce the archetypal pro-inflammatory cytokine TNF upon infection *in vitro* (Jungi et al., 1996). However, studies investigating the fate of *Mmm* in the presence of macrophages are lacking.

Another major contributor to the host defense against infection is the complement system. Complement proteins are present in extracellular fluids allowing direct killing and non-specific opsonization, followed by increased phagocytosis of pathogens (Volanakis, 1998). Non-specific opsonization can result in killing but also in intracellular survival if pathogens are able to prevent fusion of phagosomes with lysosomes (Cosma et al., 2003; Horwitz, 1983). Upon activation, the complement promotes inflammation through the induction of pro-inflammatory cytokines (Okusawa et al., 1988; Laudisi et al., 2013). These pleiotropic effects of complement are not exclusive and may be dose-dependent. For example, the complement membrane attack complex responsible for the pathogen killing activity of complement can also induce the production of pro-inflammatory cytokines by phagocytes when used at non-bactericidal doses (Laudisi at al., 2013). Thus, in CBPP, interactions between *Mmm*, macrophages and complement may either benefit the host, through direct or opsonization-mediated killing of *Mmm* or: paradoxically; harm the host, through opsonization-mediated intracellular survival of *Mmm* and amplification of inflammatory responses.

Although previous work suggests that mycoplasmas are generally phagocytosed by macrophages only in the presence of specific antiserum (Marshall et al., 1995), the question remains open for *Mmm*. Also, effects of complement other than opsonization, such as direct killing and cytokine production, may impact disease outcome in CBPP. Therefore, the following study was undertaken to investigate the interaction of bovine macrophages with *Mmm* under physiological conditions, *i.e.*, in the presence of complement prepared from the same cell donor. Firstly, the effect of bovine complement on the viability of *Mmm* was analyzed to determine non-bactericidal concentrations. Secondly, the uptake of *Mmm* by macrophages and subsequent extra- and intracellular survival in the presence of bovine complement at non-bactericidal doses was investigated. Finally, we addressed the possibility that complement acts as a potentiator of *Mmm*-induced TNF responses of macrophages.

## Methods

### Bacterial strains

Non-fluorescent *Mmm* strain 8740-Rita and *Mycoplasma bovis* strain Oger2, and fluorescent mNeonGreen-expressing *Mmm* mutant strain 8740-Rita clone 6 (RN6), were grown in PPLO-based medium as described previously (Bonnefois et al., 2016). Mycoplasmas obtained from log-phase cultures were washed once in PBS by centrifugation at 10000 x $g$ for 10 min before resuspension in macrophage culture medium (see below) at $1\text{-}2\text{x}10^9$ CFU/ml, as determined through dilution and plating on PPLO-based media supplemented with 1 % Noble Agar (Difco, USA).

### Bovine complement and antiserum

Non-decomplemented bovine sera obtained from 3 adult Jersey cattle housed at Cirad's animal facility served as a source of complement. Sera were aseptically collected from whole blood tubes after clotting for 30 min at room temperature, followed by centrifugation at 800 x g for 20 min. Single-use aliquots of non-decomplemented sera were kept in liquid nitrogen until use after quick thawing at 37°C.

82  Bactericidal capacities of bovine complement were assessed by incubation of $1x10^6$-$1x10^8$ viable *Mmm*
83  with different concentrations of non-decomplemented bovine sera at 100 μL/well in 96-well flat-
84  bottom-plates for 1 h at 37°C and 5% $CO_2$ before vortexing and CFU titration. The specific *Mmm*
85  antiserum was obtained by pooling sera from CBPP convalescent animals. Prior to use, *Mmm* antiserum
86  was decomplemented by heating at 56°C for 30min.
87
**Macrophages and bacterial infection**
Macrophages were derived from monocytes obtained from the same three animals used to
prepare non-decomplemented bovine sera and approved by the Languedoc-Roussillon regional ethics
committee (French CE-LR#36) in the authorized project using animals for scientific
purposesAPAFIS#27628. Briefly, monocytes were purified from PBMC by positive selection of
CD14+ cells using anti-human CD14 magnetic beads (Miltenyi, Germany) according to the
manufacturer's instructions. Monocytes were resuspended in Iscove's Modified Dulbecco's Medium
(IMDM) supplemented with 2 mM L-glutamine, 50 μM 2-mercaptoethanol, 0.4 mg/mL ampicillin and
10% heat-inactivated fetal calf serum (FCS; Eurobio AbCys, France), seeded at 3 x $10^5$cells/well of 96-
well flat-bottom-plates in 200 μl, and incubated for 6 days at 37°C and 5% $CO_2$ with a change of half
of the medium at day 3. Before infection, the medium was removed and replaced with IMDM lacking
FCS to which the following stimuli were added to reach a final volume of 100 μl/well: i) appropriate
amounts of mycoplasmas to obtain MOIs of 1, 10 and 100; and ii) non-decomplemented autologous
bovine sera or antiserum or FCS. Non-decomplemented bovine sera were used at non-bactericidal doses
ranging from 5 to 10% v/v depending on the donor, whereas antiserum was used at 20% v/v and FCS
at 10% v/v. After 1 h incubation, the medium was replaced and 100μl of complete IMDM medium was
added to the wells before harvesting or further incubation. For harvesting macrophages, monolayers
were scrapped off 200μl tips, pelleted by centrifugation at 500 x g for 5 min, resuspended in 200 μL
PBS, and passaged 5 times through a 27gauge needle, which lysed more than 80% of cells as shown by
Trypan blue viability counts. CFU *Mmm* titers were determined from 10-fold serial dilution by plating
20 μL on agar by counting single colonies after 4 days incubation. For gentamycin assays, an MOI of
10 was used, as it was shown in preliminary macrophage-free assays that addition of 400 μg/ml of
gentamicin to the culture for 3 hours reliably killed 100% of mycoplasmas. Cultures infected with *M.*
*bovis* strain Oger2 were used as positive controls of intracellular survival in the presence of gentamycin
(Suleman et al., 2016).
113
**Flow cytometry**
Macrophages were infected with fluorescent and non-fluorescent *Mmm* at MOIs of 500-1000
as described above. Due to the high MOI, non-decomplemented bovine sera could be used at up to 40%
v/v without affecting *Mmm* viability. Before harvesting, the medium was replaced by 100 μl/well of
fresh medium and cells were scraped off, homogenized by pipetting up and down, and transferred to 5
ml cytometry tubes. After addition of 10 μL/tube of propidium iodide (PI, BD Biosciences, USA) and
10min incubation at 4°C in the dark, 100 μL of sheath fluid was added to each tube.  Preliminary
macrophage-free tests showed that 100% of *Mmm* took up PI in these conditions. Since PI and
mNeonGreen fluoresce in different wavelength, it was possible to gate out PI positive macrophages to
exclude dead cells (i.e., with loss of membrane integrity) and extracellular *Mmm*. This allowed analysis
of strictly intracellular mNeonGreen-dependent fluorescence. CytochalasinD (Sigma, France), an
inhibitor of actin-dependent phagocytosis, was used at 10 μg/ml. At least 5000 events were analyzed
within the PI negative gate with a FacsCanto flow cytometer (BD Biosciences) equipped with the
FacsDiva software (BD Biosciences). Gates and quadrants were used to delineate populations of interest
and calculate statistics respectively. Results were expressed in percentages of cells fluorescing above
background (*i.e.*, fluorescence produced by macrophages infected with non-fluorescent *Mmm*) and
geometric mean fluorescence intensity (MFI), which is proportionally related to numbers of
mNeonGreen+ *Mmm* per cell.
132
**ELISA**
Supernatants were collected from macrophage cultures infected as above for 24 h with *Mmm* at
MOIs ranging from 100 to 1000 and stored at -20°C until use. Non-decomplemented bovine sera were

used at both bactericidal and non-bactericidal concentrations and the highest dose was also used for
FCS. Lipopolysaccharide (LPS) from *Escherichia coli* 0111:B4 (Invivogen, France), used at 2 µg/ml,
served as a positive control. To quantify bovine TNF, a sandwich ELISA was performed using a
matched antibody pair (BioRad, France) as previously described (Totté et al., 2015). Supernatants were
tested in duplicates and results obtained from an ELISA reader were expressed in mean optical densities.
141
142 **Statistics**
143     A non-parametric Mann–Whitney U-test (https://www.socscistatistics.com/tests/mannwhitney/)
144 was used to analyze differences between responses obtained with various stimuli. Single and double
145 asterisks represent *p* values of <0.05 and <0.01 respectively. A difference was considered to be
146 significant at a *p* value of <0.05.
147
148 <center>**Results**</center>
149
**Bactericidal effect of bovine complement on *Mmm* viability in the absence of macrophages**
A dose-dependent effect of non-decomplemented bovine sera, containing bovine complement,
on *Mmm* CFU titers was observed *in vitro* (Fig1). Depending on the dose and animals used the decrease
in CFU titers varied between 0 and 3 logs, indicating a potent effect of bovine sera and a high variability
among animals. There was no sign of *Mmm* aggregation induced by non-decomplemented serum when
cultures were checked under a light microscope, and passage through a 27G needle had no impact on
CFU titers, indicating that reduced viability occurred through direct cytotoxicity. The killing effect was
due to complement since it was neutralized by heating sera at 56°C for 30 min (results not shown). The
mycoplasmacidal activity and inter-animal variability of bovine sera was stable over the 3 years study
period (Suppl.Fig1).
160

161
162     **Figure1:** Cytotoxic effect of bovine complement on *Mmm* in vitro. *Mmm* was incubated for 1
163 h with decomplemented sera (FCS) or different concentrations (10%, 20%) of non-decomplemented
164 bovine sera from 3 different animals (s+animal number). Results from 3 different experiments are
165 shown as mean+/-SD of CFU expressed in log10. Asterisks and bars indicate statistically significant
166 differences with FCS.
167
**Effect of bovine complement on *Mmm* opsonization and survival in the presence of macrophages**
A protocol was used to focus on *Mmm* associated to macrophages and to exclude non-adherent
and plastic-adherent mycoplasmas from the analysis (see methods). Furthermore, no cytopathological
effects of *Mmm* on macrophage monolayers were detected at any MOI used. In the absence of bovine
complement and antiserum, complete survival of *Mmm* occurred at 24 h and growth of up to 1 log was
observed at 48h, indicating no killing activity by macrophages (Fig2). Similar trends were observed at
MOIs of 1 and 100 whereas only survival of *Mmm* without growth was observed in the absence of
macrophages (data not shown). The addition of non-decomplemented bovine sera at non-bactericidal
concentrations had no effect on *Mmm* viability at any time-points (Fig2). As expected, addition of
*Mmm*-specific antiserum induced significant killing activity after 24 h and up to 2.5 logs decrease in
CFU titers at 48h (Fig2).
179

180
**Figure 2**: Macrophages do not kill *Mmm* even in the presence of bovine complement.
Macrophages were infected with *Mmm* at an MOI of 10 in the absence (FCS) or presence (serum) of
non-decomplemented bovine sera at non-bactericidal concentrations and incubated for 1, 24, and 48h
before assessment of macrophages-associated *Mmm* titers. Anti-Mmm antiserum (antiserum) was
added to additional wells as positive controls of bactericidal activity. Results from 3 different
experiments using 3 animals are shown as mean+/-SD of CFU expressed in log10. Asterisks indicate
statistically significant differences with FCS.

188
In order to analyze intracellular survival of *Mmm* experiments were repeated in the presence of
gentamicin to kill all extracellular mycoplasmas. This was performed only at an MOI of 10, which
represented a concentration of $3\times10^6$ CFU/ml of *Mmm*, since gentamicin did not kill 100% extracellular
mycoplasmas at higher *Mmm* concentrations. As seen in Fig.3, gentamicin assays indicated that the
overwhelming majority of *Mmm* survival at 24 h was due to extracellular mycoplasmas and that bovine
complement had no effect on *Mmm* survival inside macrophages. This was not due to a bias in
gentamicin assay since significant survival was observed with *M. bovis* Oger2 (Fig3).

197
**Figure 3**: Bovine complement has no impact on *Mmm* survival inside macrophages.
Macrophages were infected with *Mmm* at an MOI of 10 in the absence (FCS) or presence (serum) of
bovine complement at non-bactericidal concentrations and incubated for 24 h. Cells were then treated
for 4-hours with gentamycin before harvesting and assessment of macrophage-associated *Mmm* titers.
Wells containing macrophages infected with *M. bovis* (Oger2) were used as positive controls of
intracellular survival in the presence of gentamycin. Results from 3 different experiments using 3
animals are shown as mean+/-SD of CFU expressed in log10.

205
**Effect of bovine complement on the uptake of fluorescent *Mmm* by macrophages**
We were able to analyze *Mmm*-dependent intracellular fluorescence by gating out extracellular
fluorescence as explained in "Methods". As seen in Fig4, in the absence of bovine complement or
antiserum, more than 90% of macrophages had engulfed fluorescent *Mmm* but at low levels as indicated
by low FITC fluorescence intensity ("Li" in Fig4a). Indeed, FITC signals did not exceed one log above
background level and could not be detected when MOIs of 100 or less were used (data not shown). No
differences were detected between 1 and 24 h PI in both percentages of positive cells and MFI (Fig4a
and Fig4b). Non-decomplemented bovine sera used at non-bactericidal concentrations had no effect on

FITC fluorescent levels at any time-points (Fig4a and Fig4b). Data at 48h were not exploitable due to
high background fluorescence. The low-level uptake was not affected by cytochalasinD ("Li" in Fig4c)
indicating that it was not dependent on phagocytosis. A substantially stronger FITC fluorescence
intensity was observed in the presence of antiserum indicating a strong increase in *Mmm* uptake by
macrophages ("Hi" in Fig4a). At 24 h PI, the % of cells within the "Hi" gate increased while the mean
fluorescence intensity decreased (gate P5 and MIF in Fig4b) suggesting partial destruction of
fluorescent *Mmm* within macrophages. Moreover, the vast majority of highly fluorescent cells
disappeared in the presence of cytochalasinD (FIg4c) indicating that the uptake of *Mmm* by
macrophages in the presence of antiserum was dependent on phagocytosis.
The lack of effect of bovine complement on the uptake of *Mmm* by macrophages was confirmed
for all three animals tested (Table1). Indeed, the addition of non-decomplemented bovine sera at non-
bactericidal concentrations had no impact on numbers of positive cells and MFI when compared to FCS
at 1 h and 24 h PI. In addition, the positive effect of antiserum on *Mmm* uptake was confirmed as well
as the increase in percentage of positive cells at 24 h PI accompanied by a decrease in MIF, indicating
progressive destruction of phagocytized fluorescent *Mmm*.
229

230
**Figure 4**: Bovine complement does not impact *Mmm* uptake by macrophages. Macrophages were
infected with either non-fluorescent or fluorescent *Mmm* at an MOI of 500-1000, and either in the
absence (FCS) or in the presence (serum) of non-decomplemented bovine sera at non-bactericidal
concentrations. They were incubated for 1 h (a) and 24 h (b) before analysis of intracellular FITC
fluorescence by flow cytometry (see Methods). Anti-*Mmm* antiserum (antiserum) was added to
additional wells as positive control of specific phagocytosis. Typical histogram plots are shown for
one representative animal as an example. Histograms and arrows display background (bgd, hatched
line), and low (Li, green line) and high (Hi, green line) fluorescence intensities produced by
macrophages infected with non-fluorescent and fluorescent *Mmm* respectively. The effect of
cytochalasinD, an inhibitor of phagocytosis, on *Mmm* uptake by macrophages is shown in c).
Percentages of highly fluorescent cells and MFI are given by the P5 interval.

**Table 1:** Effect of bovine complement (serum) from three different animals (mean+/-SD) on the uptake of fluorescent *Mmm* by
autologous macrophages as measured by flow cytometry 1 h and 24 h post infection

| Stimuli | Low fluorescence intensity[a] | | | | High fluorescence intensity[a] | | | |
|---|---|---|---|---|---|---|---|---|
| | % pos [b] | | MFI[c] | | % pos [b] | | MFI[c] | |
| | 1 h | 24 h | 1 h | 24 h | 1 h | 24 h | 1 h | 24 h |
| FCS | 93+/-2 | 96+/-5 | 1013+/-33 | 995+/-33 | 7+/-3 | 10+/-1 | NA | NA |
| Serum | 92+/-3 | 99+/-3 | 1008+/-52 | 1050+/-42 | 8+/-4 | 9+/-2 | NA | NA |
| Antiserum[d] | NA | NA | NA | NA | 63+/-10 | 94+/-5 | 14518 +/-1313 | 6905 +/-948 |

(a): low and high fluorescence intensities above background fluorescence (see also Figure 4)
(b): Percentages of positive cells
(c): Mean fluorescence intensity
(d): sera from CBPP convalescent animals
NA: not applicable. The effect of antiserum is measurable only within the high fluorescence intensity whereas, in the presence of
complement, the vast majority of positive cells were found in the low fluorescence intensity range.
261

**Effect of bovine complement on *Mmm*-induced TNF production by macrophages**

As shown in Fig5, in the absence of bovine sera and antiserum, *Mmm* induced the production of TNF by macrophages when MOIs of 500-1000 were used. At an MOI of 100, TNF was not detected (not shown). There was no effect of bovine complement at non-bactericidal doses (Fig.5, serum1). In order to analyze the effect of non-decomplemented bovine sera at bactericidal doses, concentrations of both non-decomplemented bovine sera and FCS had to be increased. For one animal, a substantial increase in background (*i.e.,* TNF induced by FCS alone in the absence of *Mmm*) prevented its use. Nevertheless, non-decomplemented bovine sera from the remaining two animals, used at bactericidal concentrations inducing at least 1log decrease in *Mmm* viability, almost completely abrogated the production of TNF induced by *Mmm* (Fig5, serum2).

**Figure 5** : Effect of bovine complement on *Mmm*-induced release of TNF by autologous macrophages. Macrophages were either left uninfected (*Mmm-*) or infected (*Mmm+*) with *Mmm* at an MOI of 500-1000 in either the absence (FCS) or presence of non-decomplemented bovine sera at non-bactericidal (serum1) and bactericidal (serum2) concentrations and incubated for 24 h before TNF titration of supernatants. Lipopolysaccharide (LPS) was added to additional wells as positive controls. Results from 3 different experiments using complement from two animals are shown as mean+/-SD of optical densities (OD). Asterisks indicate statistically significant differences.

## Discussion

Given their pivotal role in first-line immune defenses against mucosal pathogens, we have
investigated the interactions between macrophages and *Mmm*, the causal agent of CBPP. Non-
decomplemented bovine sera were used as a source of complement to mimic physiological conditions
present at infected sites and cryopreservation in liquid nitrogen was instrumental in overcoming the
instability of complement. The strong bactericidal effect of some non-decomplemented bovine sera on
*Mmm* viability that we observed in this study is in conflict with previous work (Gaurivaud et al., 2014).
One likely explanation is that we used bovine sera instead of guinea pig sera. Similarly, non-
decomplemented caprine sera was found to reduce the viability of *Mycoplasma mycoides* subsp. *capri*
(*Mmc*), *Mmm*'s closest relative (Schieck et al., 2016). Given its stability over time *in vivo* and variable
potency between animals, it would be interesting to assess whether a correlation exists between the anti-
*Mmm* effect of bovine sera and the resistance of animals to CBPP. Considering the pleiotropic effects
of complement on immune responses to pathogens, we have explored the possible impact of bovine
complement, used at non-bactericidal concentrations, on important functions of macrophages such as
non-specific opsonization-mediated phagocytosis and production of pro-inflammatory TNF.
In the first place, we analyzed the kinetics of *Mmm* survival in the presence of bovine
macrophages and in the absence of complement and antiserum. Lysis of macrophages prior to *Mmm*
titration allowed taking into account both intracellular and membrane-associated mycoplasmas. *Mmm*
CFU titers remained stable after 24 h and increased by up to 1log after 48h, indicating the absence of
measurable killing by macrophages. Little or negligible numbers of macrophage-associated *Mmm*
survived intracellularly in the presence of gentamicin, suggesting that viable *Mmm* were mainly located
on cell membranes. Absence of killing by macrophages or even growth on macrophages in the absence
of specific antiserum and complement has been reported previously for other mycoplasmas (Marshall
et al., 1995; Howard et al., 1976). We found no measurable effect of bovine complement on *Mmm* CFU
titers, suggesting that neither non-specific opsonization-mediated killing nor intracellular survival
occurred under our experimental conditions. It should be noted that pre-incubation of *Mmm* with bovine
complement for 30 min at 37°C before infecting macrophages yielded similar results (not shown).
However, we cannot exclude the possibility that the presence of *Mmm* growing on macrophages masks
a low-level killing activity. On the other hand, *Mmm* titers substantially decreased over time when anti-
*Mmm* antiserum was added to the medium indicating that intrinsic killing capacities of macrophages
were not altered. Again, absence of killing by macrophages, even in the presence of complement, has
been shown previously for several mycoplasmas (Marshall et al., 1995). It is possible that the presence
of a polysaccharide capsule (Buttery, 1970) protects non-opsonized and non-specifically opsonized
*Mmm* from phagocytosis, which may be further investigated using non capsulated variants (Gaurivaud
et al., 2014).
In order to gain a better insight on the interactions of *Mmm* with bovine macrophages, we used
a more direct approach based on flow cytometry-assisted monitoring of intracellular fluorescence of
phagocytes after infection with fluorescent *Mmm*. Our results show that, in the absence of complement
and antiserum, more than 90% of macrophages take up fluorescent *Mmm* at low levels, as indicated by
low mean fluorescence intensities. Unfortunately, gentamicin was not 100% bactericidal at MOIs used
for flow cytometry, thus preventing assessment of the viability of intracellular *Mmm*. The lack of
decrease in fluorescence over time suggests that either no killing occurred or that it was not sufficient
to cope with ongoing growth of *Mmm* on macrophages. CytochalasinD had no effect on the number of
cells harboring low intracellular fluorescence, suggesting that the uptake of *Mmm* by macrophages was
independent of phagocytosis. Interestingly, it has recently been shown that *Mmc* interacts with C-type
lectin receptors (CTLRs) (Lindenwald et al., 2020) some of which mediate endocytosis. Several CTLRs
are present on macrophages, which warrants further studies on their potential role in CBPP. Moreover,
pathogen-associated molecular patterns (PAMPs) of *Mmm* recognized by these receptors have potential
as virulent factors. Bovine complement used at non-bactericidal concentrations had no effect on the
uptake of *Mmm* by macrophages at any time points, confirming viability experiments using CFU
titration. Also, it should be noted that bovine complement did not improve *Mmm* capacity to bind to the
surface of macrophages, as measured by PI-dependent extracellular fluorescence (not shown). On the

other hand, anti-*Mmm* antiserum induced a strong and phagocytosis-dependent increase in *Mmm* uptake leading to more than 90% of cells located in the high fluorescence intensity gate at 24 hpi. Moreover, the mean fluorescence intensity within that gate decreased at 24 hpi confirming the progressive destruction of *Mmm* as seen with CFU titration. These last results are intriguing since *Mmm* and related mycoplasmas possess a two-protein system capable of cleaving immunoglobulins G (IgG) (Arfi et al., 2016). Decomplementation of anti-*Mmm* antisera did not affect their opsonizing potency indicating that IgGs are most likely involved.

Our results confirm previous work demonstrating the capacity of *Mmm* to induce TNF production by macrophages in the absence of either complement or antiserum (Jungi at al., 1996). Interestingly, TNF was induced only at high MOIs (i.e., >500) also shown by flow cytometry to be associated with phagocytosis-independent engulfment of *Mmm*. This suggests a possible involvement of autophagy in the pro-inflammatory response of macrophages to *Mmm*. There was no beneficial effect of bovine complement on the TNF response of infected macrophages when used at non-bactericidal concentrations. The effect of bovine complement at bactericidal doses was also analyzed, considering the possible release of pro-inflammatory molecules from *Mmm* undergoing lysis. We found that, when used at bactericidal concentrations, bovine complement efficiently abrogated the production of TNF by macrophages in response to *Mmm* infection.

In summary, using a combination of CFU titration, gentamicin assays, fluorescent *Mmm*, and TNF ELISA, we were able to show that: i) macrophages were not capable of killing *Mmm* efficiently in the absence of antiserum; ii) bovine complement used at non-bactericidal doses neither had opsonizing effect nor did it promote intracellular survival; iii) bovine complement did not impact the phagocytosis-independent *Mmm* uptake by macrophages that occurred in the absence of antiserum; iv) bovine complement used at non bactericidal concentrations did not amplify *Mmm*-induced TNF responses of macrophages, and even abrogated that response when used at concentrations inducing bactericidal activity. These results provide a framework to identify PAMPs as well as *Mmm* antigens recognized by CTLRs and protective antibodies. In addition, they point towards a role for complement in the control of TNF-dependent inflammatory responses in CBPP.

## Data, scripts, code, and supplementary information availability

Supplementary File1 is available online: XXXXDOI of the webpage hosting the data https://doi.org/10.5802/fake.doi

## Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

## Funding

This work was financially supported by the French Agricultural Research Centre for International Development (CIRAD) in France.

## References

Arfi Y, Minder L, Di Primo C, Le Roy A, Ebel C, Coquet L, Claverol S, Vashee S, Jores J, Blanchard A, Sirand-Pugnet P. (2016) MIB-MIP is a mycoplasma system that captures and cleaves immunoglobulin G. Proc Natl Acad Sci U S A. May 10;113(19):5406-11. https://doi.org/10.1073/pnas.1600546113

Bonnefois T, Vernerey MS, Rodrigues V, Totté P, Puech C, Ripoll C, Thiaucourt F, Manso-Silván L. (2016) Development of fluorescence expression tools to study host-mycoplasma interactions and validation in two distant mycoplasma clades. J Biotechnol. Oct 20;236:35-44. https://doi.org/10.1016/j.jbiotec.2016.08.006

Buttery, S.H. (1970) Hapten inhibition of the reaction between Mycoplasma mycoides polysaccharide and bovine antisera. Immunochemistry, 7, 305-310. https://doi.org/10.1016/0019-2791(70)90168-0

Cosma CL, Sherman DR, Ramakrishnan L. (2003) The secret lives of the pathogenic mycobacteria. Annu Rev Microbiol 57:641-76. https://doi.org/10.1146/annurev.micro.57.030502.091033

Gaurivaud P, Lakhdar L, Le Grand D, Poumarat F, Tardy F. (2014) Comparison of in vivo and in vitro properties of capsulated and noncapsulated variants of Mycoplasma mycoides subsp. mycoides strain Afadé: a potential new insight into the biology of contagious bovine pleuropneumonia. FEMS Microbiol Lett. Oct;359(1):42-9. https://doi.org/10.1111/1574-6968

Horwitz, M. A. (1983) The Legionnaires' disease bacterium (Legionella pneumophila) inhibits phagosome-lysosome fusion in human monocytes. J. Exp. Med. 158:2108- 2126. https://doi.org/10.1084/jem.158.6.2108.

Howard CJ, Taylor G, Collins J, Gourlay RN. (1976) Interaction of Mycoplasma dispar and Mycoplasma agalactiae subsp. bovis with bovine alveolar macrophages and bovine lacteal polymorphonuclear leukocytes. Infect Immun. Jul;14(1):11-7. https://doi.org/10.1128/iai.14.1.11-17.1976

Jungi TW, Krampe M, Sileghem M, Griot C, Nicolet J (1996) Differential and strain-specific triggering of bovine alveolar macrophage effector functions by mycoplasmas. Microb Pathog. 2:487–498. https://doi.org/10.1006/mpat.1996.0078

Laudisi F, Spreafico R, Evrard M, Hughes TR, Mandriani B, Kandasamy M, Morgan BP, Sivasankar B, Mortellaro A. (2013) Cutting edge: the NLRP3 inflammasome links complement-mediated inflammation and IL-1β release. J.Immunol. Aug 1;191(3):1006-10. https://doi.org/10.4049/jimmunol.1300489

Lindenwald DL, Monteiro JT, Rautenschlein S, Meens J, Jung K, Becker SC, Lepenies B. (2020) Ovine C-type lectin receptor hFc-fusion protein library - A novel platform to screen for host-pathogen interactions. Vet Immunol Immunopathol. Apr 18;224-230. https://doi.org/10.1016/j.vetimm.2020.110047

Marshall AJ, Miles RJ, Richards L. (1995) The phagocytosis of mycoplasmas. J Med Microbiol. Oct;43(4):239-50. https://doi.org/10.1099/00222615-43-4-239

Okusawa S, Yancey KB, van der Meer JW, Endres S, Lonnemann G, Hefter K, Frank MM, Burke JF, Dinarello CA, Gelfand JA. (1988) C5a stimulates secretion of tumor necrosis factor from human mononuclear cells in vitro. Comparison with secretion of interleukin 1 beta and interleukin 1 alpha.  J Exp Med. 1;168(1):443-8. https://doi.org/10.1084/jem.168.1.443

Provost A, Perreau P, Bréard A. Contagious bovine pleuropneumonia (1987) Rev Sci Tech Off. int. Epiz., 6 (3), 625-679. https://doi.org/10.20506/rst.6.3.306

Scanziani E, Paltrinieri S, Boldini M, Grieco V, Monaci C, Giusti AM, Mandelli G (1997) Histological and immunohistochemical findings in thoracic lymph nodes of cattle with contagious bovine pleuropneumonia. J Comp Pathol. Aug;117(2):127-36.https://doi.org/10.1016/s0021-9975(97)80029-1

Schieck E, Lartigue C, Frey J, Vozza N, Hegermann J, Miller RA, Valguarnera E, Muriuki C, Meens J, Nene V, Naessens J, Weber J, Lowary TL, Vashee S, Feldman MF, Jores J. (2016) Galactofuranose in Mycoplasma mycoides is important for membrane integrity and conceals adhesins but does not contribute to serum resistance. Mol Microbiol. Jan;99(1):55-70. https://doi.org/10.1111/mmi.13213.

Suleman M, Prysliak T, Clarke K, Burrage P, Windeyer C, Perez-Casal J. (2016) Mycoplasma bovis isolates recovered from cattle and bison (Bison bison) show differential in vitro effects on PBMC proliferation, alveolar macrophage apoptosis and invasion of epithelial and immune cells. Vet Microbiol. Apr 15;186:28-36. https://doi.org/10.1016/j.vetmic.2016.02.016

Totté P, Puech C, Rodrigues V, Bertin C, Manso-Silvan L, Thiaucourt F. (2015) Free exopolysaccharide from Mycoplasma mycoides subsp. mycoides possesses anti-inflammatory properties. Vet Res. Oct 21;46:122. https://doi.org/10.1051/vetres:2007046

Volanakis J.E. (1998) Overview of the complement system, in: Volanakis J.E., Frank M.M., (Eds.), The human complement system in health and disease, Marcel Dekker Inc., New York, 1998, pp. 9–32. https://doi.org/10.1201/9780367800772

World Organization for Animal Health (2015). http://oie.int/en/animalhealth-in-the-world/oie-listed-diseases-2015/. Accessed 20 Sept 2022