Is niche divergence more likely in parapatry? A test in *Sclerurus mexicanus* sensu lato

(Aves: Furnariidae)

Jacob C. Cooper[1,2,3,5] and Diego Barragán[1,4]

1. University of Kansas Biodiversity Institute, 1345 Jayhawk Boulevard, Lawrence, Kansas

66045

2. Committee on Evolutionary Biology, Culver Hall 402, University of Chicago, Chicago,

Illinois 60637

3. Field Museum of Natural History, Life Sciences Division, 1400 S. Lake Shore Drive,

Chicago, Illinois 60605

4. Futuro Azul, Jaime Roldos Aguilera E6-14 & Velasco Ibarra, Quito, Ecuador

5. Contact author email: jccooper@uchicago.edu

15**Note**: This manuscript was written by the authors in 2015-2017, with this version approved for

16upload to Biorxiv in 2019, with minimal edits made in Dec 2022. This article was submitted and

17rejected pending revision; it is being posted here as a citable manuscript until the work can be

18revised appropriately and resubmitted. - JCC, 6 Dec 2022. Revision with proper GBIF DOI

19uploaded 12 Dec 2022.

20**ABSTRACT**

21**Aim**

22While species' niches are understood to diverge through time, the mode of these divergences is
23not fully understood. The null hypothesis of niche divergence in the present is that of
24conservatism; that species are likely to occur in the same environment as their ancestors and,
25therefore, as allopatric sister species. Alternatively, species are expected to diverge when
26selective pressure, such as parapatry with a congener, occurs. Here, we analyse niche divergence
27in a mosaic of allo- and parapatric *Sclerurus mexicanus* sensu lato populations to determine if
28niche divergence is more likely in parapatry.

29**Location**

30Neotropics.

31**Methods**

32We created a dataset of 1,100 vetted localities for *S. mexicanus* sensu lato, and assigned each
33point to each of the seven described populations. We created individual dispersal areas for each
34species in Maxent following: 1) a broad approach in which training areas where allowed to
35overlap with parapatric congeners; and 2) a narrow approach in which species dispersal areas
36adhered to strict parapatry and there was no overlap in training areas. We complemented these
37ENMs with 'random' models drawn from points within each training area, thus creating 'null'
38ENMs against which we could test niche divergence using the metric Schoener's D.

39**Results**

40There was no significant difference in the performance of broad and narrow training areas. Few
41significant divergences were found, and all of those that were found consisted of allopatric
42populations. Partial divergences were frequently recovered when comparing parapatric taxa.

43Main conclusions

44In contrast to our expectation, we found no significant evidence for increased niche divergence

45in parapatric versus allopatric species. Possible explanations for these findings include true

46conservatism even among parapatric lineages (*e.g.*, additional environmental or biological

47factors may delimit distributions) or artefactual errors inherent to model creation. We discuss the

48implications of these findings, and discuss ways to improve upon tests of niche divergence in the

49future, especially when dealing with cases of parapatry.

50Keywords

51Allopatry, ecological niche modelling, Maxent, niche divergence, parapatry, Schoener's D,

52*Sclerurus mexicanus*.

54INTRODUCTION

55   Since the conception of ecological niche theory, attempts have been made to quantify the

56differences between the occupied niche space of different taxa (Grinnell, 1917; Elton, 1927;

57Hutchinson, 1957; Colwell & Futuyma, 1971; Carnes & Slade, 1982; Austin et al., 1990; Glor &

58Warren, 2011). These differences have been hypothesized to be key or causal to species'

59diversification (Orr & Smith, 1998), especially in cases of related species' groups that inhabit

60spatially parapatric niches. In recent years, new methods have been put forth to test how species'

61niches differ in ecological space, all utilizing different approaches to Ecological Niche Modeling

62(ENM), and all with the overarching goal of testing whether or not species truly inhabit different

63multivariate environmental (*i.e.*, Hutchinsonian) niches (Hutchinson, 1957; Warren et al., 2008;

64Blonder et al., 2014; Nunes & Pearson, 2016).

65   As our knowledge of species' diversification has increased, it has become apparent that

66most species diverge in allopatry by means of genetic drift, differential selection, and a lack of

67genetic mixing (Price, 2008). Likewise, our expectation of niche divergence has approached the

68null; that is, when species become allopatric, we hypothesize that they are more likely retain

69their ancestral niche and therefore appear 'conserved' via ENM tests (Peterson et al., 1999;

70McCormack et al., 2009; Peterson, 2011; Ryan Shipley et al., 2013; Khaliq et al., 2015; Skeels &

71Cardillo, 2017). Furthermore, research has shown that competition and the re-establishment of

72contact can drive niche (and character) divergence, and thereby result in the para- and sympatric

73shifts we observe in the present (Jang & Gerhardt, 2006; Kirschel et al., 2009; Wooten & Gibbs,

742012; Freeman, 2015; Miller et al., 2016).

75   While ecological niche divergence has been examined in allopatric complexes (e.g.,

76McCormack et al., 2009), it has been less studied in parapatric species complexes. Given the

77expectation in allopatric speciation that niches remain conserved until secondary contact with a

78competitor occurs, it is therefore logical to hypothesize that the magnitude of niche divergence

79will be greater when species have a contact zone during their speciation cycle (Nunes & Pearson,

802016). This hypothesis is reinforced by our understanding of niche shifts that accompany

81speciation events involving species moving into parapatric habitats or geographic realms

82(Benkman et al., 2009; Fjeldså et al., 2011; Knowles & Massatto, 2016), and has provided the

83basis of a novel methodology for testing ecological niche divergence (Nunes & Pearson, 2016).

84There are few tests of the assumption of niche divergence within monophyletic parapatric groups

85(Reino et al., 2016), and it is difficult to fully assess the other factors that may be influencing

86species' realized niches in ENMs.

87   Soberón & Peterson (2005)elucidate ways to address additional factors in their **BAM**

88framework, highlighting three factors that are responsible for limiting species' distributions: (**B**)

89biotic interactions; (**A**) abiotic environmental conditions (*i.e.*, scenopoetic variables *sensu*

90Soberón*, 2007);* and (**M**) motility restrictions, such as geographic barriers (*i.e.*, an invasible

91area). Biotic interactions, collectively known as a species' Eltonian niche (Soberón, 2007), are

92often assumed to be implicitly accounted for in ENMs by a study species' absence from areas

93with a biotic amensal or without a biotic commensal. Linking these factors directly into an ENM

94is often done by restricting a species' explorable area in the initial steps of ENM creation,

95thereby removing regions of potential sympatry from model training (Saupe et al., 2012; de

96Araújo et al., 2014; Cooper & Soberón, 2018). ENMs can also opt to ignore these biotic factors,

97as many of the localized competitors and commensals a species encounters may not be as

98important as abiotic factors in large-scale spatial analyses (*i.e.*, the Eltonian Noise Hypothesis;

99Soberón, 2007; Soberón & Nakamura, 2009).

100   In order to better understand niche divergence in parapatry, we performed a multi-

101directional niche-difference test of a widespread species group, *Sclerurus mexicanus* sensu lato

102(Cooper & Barragán, 2016). Now known to comprise of at least five major groups of taxa

103(d'Horta et al., 2013; Cooper & Cuervo, 2017), populations of *S. mexicanus* occur in allo- and

104parapatric populations from central Mexico to southern Brazil, and exhibit remarkable levels of

105ecomorphic and phenotypic conservatism, with field reports of species' responding to unrelated,

106allopatric population's songs (Remsen, 2003; Cooper & Barragán, 2016). Given their lack of

107known sympatry, it is likely that parapatric populations are competitors with other members of

108the super species, and therefore provide an excellent mosaic of parapatric and allopatric

109populations in which to test the magnitude of niche divergence in the presence and absence of

110amensals.

111   Here, we perform two sets of niche divergence analyses on this species complex under

112two alternative hypotheses: 1) that species can freely disperse into regions occupied by

113parapatric congeners, and 2) that species are prohibited from dispersal into regions where

114congeners are known occur. While both of these methods will model a species' absence in areas

115with direct competitors, their mechanisms in reaching these conclusions differ. The first

116hypothesis excludes environments that are more optimal for competitive congeners by

117accounting for these as "absent" environments for the species of interest, while the second

118hypothesis manually thresholds environments and excludes these areas from analyses altogether.

119The first hypothesis also assumes that species have differential competitive abilities in different

120parts of their shared niche space or differ to at least some extent in their fundamental niches,

121while the second hypothesis assumes that competitive exclusion is masking the limits of a

122species' ecological niche. While there is no way to get around these errors for attempting to

123assess a species' complete fundamental niche, it is still possible to analyze a species' realized

124niche and determine whether or not species are significantly differing in their occupied

125ecological space. Furthermore, a test of both methods will help reveal whether biotic exclusion is

126significantly affecting our interpretations of ecological niche divergence.

128**METHODS**

129   We downloaded occurrence data for *Sclerurus mexicanus* sensu lato from the Global

130Biodiversity Informatics Facility (GBIF.org, 2015) and from eBird (EBD_relMay-2015; Sullivan

131et al., 2009; eBird, 2012) . These data were combined with a set of georeferenced recordings

132from Cooper & Cuervo (2017), and were subsequently separated into species groups

133corroborated by the recordings. We recognized five species within the complex sensu Cooper &

134Cuervo (2017), but separated subspecies of *S. macconnelli* for niche analyses in light of the

135genetic monophyly of *peruvianus* with respect to *macconnelli* and the uncertain relationships of

136*bahiae* (d'Horta et al., 2013). We thus modeled seven recognized populations from *S. mexicanus*

137sensu lato, ordered in this manuscript from north to south and listed here with the number of

138vetted occurrences used for each: *Sclerurus mexicanus* (*n* = 532); *S. pullus* (*n* = 358); *S.*

139*obscurior* (*n* = 19); *S. andinus* (*n* = 24); *S. macconnelli peruvianus* (*n* = 77); *S. macconnelli*

140*macconnelli* (*n* = 41); and *S. macconnelli bahiae* (*n* = 49; Fig. 1). Localities were cross-

141referenced with the literature, and we omitted records that were unidentifiable by range or lacked

142physical documentation, most notably: intermediate elevations (*c.* 1200-1800 m) in the western

143Andes of Colombia and Ecuador where *S. obscurior* and *S. andinus* potentially overlap; an

144unidentified population from intermediate elevations from southern Ecuador; and an unidentified

145population from Ceará, Brazil (Cooper & Barragán, 2016).

146   Based on these occurrence data, we created accessible areas in which to train the models

147(hereafter, **M**s, sensu Soberón & Peterson, 2005 and Cooper & Soberón, 2018) using QGIS 2.8.2

148(QGIS Development Team, 2016). **M**s were bounded by major biogeographic barriers (*e.g.*,

149mountain ridgelines, major rivers, ecoregion boundaries [World Wildlife Fund, 2010], etc.) and

150incorporated what is hypothesized to be the entire accessible area for the species (Soberón &

151Peterson, 2005; Guisan et al., 2006). We created two types of **M**: broad **M**s that encompassed the

152geographic distribution of a species and included regions of potential dispersal, including areas

153in which a parapatric species occurs (*i.e.*, hypothesis 1); and narrow **M**s that excluded the

154entirety of the distribution of a parapatric species from the training area (and are thus a subset of

155the broad **M**s), with divisions approximated halfway between known populations (*i.e.*,

156hypothesis 2; Fig. 2).

157   Analyses were performed in R 3.2.5 (R Core Team, 2016), with our study following the

158methodology of Warren et al. (2008). Bioclimatic data were downloaded at a resolution of 2.5

159arcminutes (*c.* 5 km at the equator) for four bioclimatic variables: mean temperature of the

160warmest quarter, mean temperature of the coldest quarter, mean precipitation of the wettest

161quarter, and mean precipitation of the driest quarter (Hijmans et al., 2005). Suitability models

162were created using the average of five raw, unclamped outputs of MaxEnt 3.3.3k (Phillips et al.,

1632004), using the packages 'dismo' (Hijmans et al., 2015), 'raster' (Hijmans, 2015), and 'rJava'

164(Urbanek, 2013). Suitability models were projected into a larger 'clade' area (an **M**

165encompassing the entire super species) so that all populations could be modeled within each

166others' distributions and suitability comparisons could be made in identical spatial and

167environmental extents.

168   Sets of random points were drawn from each population's appropriate **M** to create

169environmentally 'null' suitability predictions in MaxEnt for contrast against actual population

170comparisons using the R package 'maptools' (Bivand & Lewin-Koh, 2015). These random point

171sets are equal in number to the known occurrences and exclude exact coordinates of known

172occurrences. We used these points to create null ENMs within the **M** of each species for each

173hypothesis using the same methodology outlined for individual species' models. Following

174Warren et al. (2008), we used the metric of Schoener's D to assess niche similarity, where the

175metric *D* is an indicator of the similarity between MaxEnt outputs, with total dissimilarity being

1760 and identical outputs being 1. We replicated these analyses 100 times bi-directionally for each

177of the 21 pairwise comparisons for each hypothesis to create two independent random

distributions of Schoener's D using the R package 'dismo' (Warren et al., 2008; Glor & Warren,

2011; Peterson et al., 2011; Hijmans et al., 2015). If the Schoener's D value of the direct

comparison was less than the critical value (i.e. within the lower 5%) of both pseudoreplicate

distributions, the comparison was scored as a "2" for being significantly divergent and the null

hypothesis of niche conservatism was rejected; if the value was significant with regard to only

one distribution, the comparison was ranked as "1" for being potentially divergent, but the results

were not considered a rejection of overall niche conservatism; no differences between niches was

ranked as "0", and indicated a failure to reject the null of niche conservatism. Plots for all

comparisons were visually inspected using the R package 'ggplot2' (Wickham, 2009). We did

not assess extreme conservatism, as this is indistinguishable from the null hypothesis for

ecological niches between allopatric populations. Differences in performance between the two

methods was assessed via a Fisher's Exact Test on the number of successes in allopatry and

parapatry for broad and narrow **M**s and on the sum of the net amount of divergence for these

categories.

RESULTS

A total of 21 pair-wise comparisons of *Sclerurus mexicanus* sensu lato populations for

both the broad and the narrow **M**s were completed (Table 1). Using broad training areas, three of

these comparisons showed significant divergence (i.e. a "2") among the populations compared,

with all three of these involving comparisons with *S. macconnelli bahiae.* Twelve comparisons

did not reject the null hypothesis of niche conservatism, but indicated potential unidirectional

(slight) divergence (i.e. a "1") while the remaining six comparisons showed no evidence for

ecological niche divergence of any kind. In almost every case of potential divergence, the critical

201value fell within the range of the values derived from a montane population's model compared to

202pseudoreplicates of its lowland congener, and not vice versa (Fig 3).

203   When species were modeled in narrow regions, the number of highly significant tests

204dropped, with only one "2" result obtained (*S. mexicanus* vs. *S. macconnelli bahiae*). In total,

205five comparisons lost value (e.g., decreased from 1 to 0), eleven remained the same, and three

206gained value (always from 0 to 1), with no significant difference in **M** performance. There was

207no observable trend as to whether allo- or parapatry increased the likelihood of obtaining a

208significant result with regards to niche divergence, nor was there any evidence that particular

209combination of habitat increased the probability of significance changing.

211**DISCUSSION**

212   Evidence for niche divergence was strongest (i.e., a 2) only in comparisons of the

213Atlantic forest *S. macconnelli bahiae* to the montane and semi-montane *S. mexicanus*, *S. pullus*,

214and *S. andinus*. Both **M** scenarios appeared to perform equally well, with the greatest

215divergences restricted to broad models between aforementioned allopatric populations. These

216results confirm the null hypothesis that niches will remain largely conserved through time, and

217that niche divergence in allopatric populations only occurs when coupled with differential

218selective pressures. In our tests, we failed to find strong support for increased niche divergence

219in parapatric versus allopatric taxa, despite the expectation that secondary contact drives species

220into different environments.

221   Despite a lack of evidence of increased divergence in parapatric taxa, it is worth noting

222that almost every instance of potentially diverging niches was the result of a direct comparison

223falling within the Schoener's D distribution derived from comparing the montane population to

224random points from a lowland population. Given that *Sclerurus* as a whole is predominately a

225lowland genus, these results may indicate competitive exclusion of highland taxa from lowland

226areas. Alternatively, these results may indicate elevational stacking of species upon secondary

227contact where species segregate based on the elevations at which they outcompete one another.

228Such a scenario would result in different realized niches, but we would be unable to recover

229significant differences via ENM tests as the environmental niches would be abutting and

230potentially overlapping. Field tests of populations' physiological tolerances and habitat

231preferences will be needed to determine whether there are uni- or bidirectional amensal

232relationships dictating elevational turnover within *Sclerurus mexicanus* sensu lato.

233   Conversely, these skewed divergences may be an artefact of model design, wherein

234montane habitats are usually proportionately smaller than adjacent lowland habitats and the

235probability of selecting montane points for a null model is thus lower. Thus, lowland taxa whose

236**M**s abut mountain ranges are more likely to have random points drawn from lowlands, whereas

237montane taxa are also more likely to have lower elevational limits drawn in a random process.

238Narrow models, while limiting the amount of habitat at different elevations that can be drawn,

239fail to account for potential biotic interactions between related taxa and assume *a priori* that taxa

240that are not explicitly sympatric are amensal. Further studies into the best ways to mitigate

241inherently unequal areas of habitat between species with different elevational ranges are needed

242to understand the best way to compensate for these inequalities, and the usage of presence-only

243techniques may be warranted (*e.g.*, minimum volume ellipsoids; Van Aelst & Rousseeuw, 2009).

244   We are aware of one other study that assessed multiple related taxa in a mosaic of allo-

245and parapatric distributions: McCormack et al. (2009). They likewise failed to find strong

246support for niche divergence in allopatric populations, and noted the apparent conservatism of

247niches within their study genus, *Aphelocoma* (Aves: Corvidae). McCormack et al. (2009) used

248defined distributions to determine areas for drawing random points and filtering occurrence data,

249but apparently did not use **M**s to constrain taxa-specific ENMs. Similar to this study, many of the

250taxa in question are restricted to montane habitats and thus have a predilection for random points

251being drawn from lowland habitats, especially when dispersal limitations are not enforced for

252individual ENMs. Coincidentally, McCormack et al.'s (2009) sympatric species pair possessed

253Schoener's D distributions closely resembling our parapatric comparisons, wherein the test

254indicates unidirectional niche divergence.

256**Implications**

257   Given the results of this and other studies using Warren et al.'s (2008) methodology for

258determining niche divergence, it is possible that parapatric species may not be recoverable as

259possessing different ecological niches using this framework. We know from specimen and vocal

260data that members of *Sclerurus mexicanus* sensu lato occupy elevational gradients in the Andes,

261and thus infer that these species must possess some differences in their ecological niches.

262However, the topography of these regions and the nature of the modeling programs used may

263make quantifying full niche divergence impossible. Furthermore, the potential biotic element of

264these scenarios – amensal relationships with congeners – implies that species' distributions may

265be extremely proximate in nature and thus difficult to fully distinguish using broad-scale ENM

266procedures. The absence of a competitor in part of a species' distribution may also influence the

267recovered environmental niche breadth; *i.e.*, species may overlap environmentally with

268competitors in models due to records from areas where competitors are not present and thus not

269limiting a species' niche or distribution (Connell, 1961).

270   Investigations into presence-only methods for determining niche divergence are needed,

271as well as continued explorations of lesser-used variables such as elevation and tree cover (Cord

272et al., 2014; Title & Bemmels, 2017). While differences in fundamental niche will necessitate

273field research into species' physiological tolerances, presence-only methods will serve to

274circumvent topographical and areal biases that are introduced from pseudoabsence data. Further

275examinations of niche difference tests are also needed to determine if potential unidirectional

276divergences are typical for proximate species that differ in their realized ecological niches. While

277the results we present are indicative of widespread niche conservatism, it is entirely plausible

278that these taxa still exhibit some realized niche divergence with respect to some other ecological

279factor, and further investigation is warranted.

280**ACKNOWLEDGMENTS**

281   We are grateful to the following individuals and groups for their input on this project:
282John M. Bates, The Field Museum; Andrés Cuervo, Universidad de Los Andes; Chris Hensz, A.
283Townsend Peterson, & Jorge Soberón, University of Kansas Biodiversity Institute; the ENM
284group of the University of Kansas; and the ornithology group of The Field Museum. This work
285was part of JCC's Masters research funded by NSF 1208472 and by a University of Kansas
286summer fellowship, and was part of DB's undergraduate research at the same institution.

287**LITERATURE CITED**

288Van Aelst S. & Rousseeuw P. (2009) Minimum volume ellipsoid. *Wiley Interdisciplinary*

289   *Reviews: Computational Statistics*, **1**, 71-82.

290de Araújo C.B., Marcondes-Machado L.O., & Costa G.C. (2014) The importance of biotic

291   interactions in species distribution models: a test of the Eltonian noise hypothesis using

292   parrots. *Journal of Biogeography*, **41**, 513–523.

293Austin M.P., Nicholls A.O., & Margules C.R. (1990) Measurement of the Realized Qualitative

Niche: Environmental Niches of Five Eucalyptus Species. *Ecological Monographs*, **60**,

161–177.

Benkman C.W., Smith J.W., Keenan P.C., Parchman T.L., & Santisteban L. (2009) A New

Species of the Red Crossbill (Fringillidae: *Loxia*) from Idaho. *The Condor*, **111**, 169–176.

Bivand R. & Lewin-Koh N. (2015) maptools: Tools for Reading and Handling Spatial Objects. R

package version 0.8-36.

Blonder B., Lamanna C., Violle C., & Enquist B.J. (2014) The n-dimensional hypervolume.

Global Ecology and Biogeography, **23**, 595–609.

Carnes B.A. & Slade N.A. (1982) Some Comments on Niche Analysis in Canonical Space.

Ecology, **63**, 888–893.

Colwell R.K. & Futuyma D.J. (1971) On the Measurement of Niche Breadth and Overlap.

Ecology, **52**, 567–576.

Connell J.H. (1961) The influence of interspecific competition and other factors on the

distribution of the barnacle *Chthamalus stellatus*. *Ecology*, **42**, 710–723.

Cooper J.C. & Barragán D. (2016) Tawny-throated Leaftosser (*Sclerurus mexicanus*).

Neotropical Birds Online (T. S. Schulenberg, Editor).

Cooper J.C. & Cuervo A.M. (2017) Vocal variation and species limits in the *Sclerurus*

mexicanus complex. *The Wilson Journal of Ornithology*, **129**, 13–24.

Cooper J.C. & Soberón J. (2018) Creating individual accessible area hypotheses improves

stacked species distribution model performance. *Global Ecology and Biogeography*, **27**,

156-165.

Cord A.F., Klein D., Gernandt D.S., Pérez de la Rosa J.A., & Dech S. (2014) Remote sensing

data can improve predictions of species richness by stacked species distribution models: a

case study for Mexican pines. *Journal of Biogeography*, **41**, 736–748.

d'Horta F.M., Cuervo A.M., Ribas C.C., Brumfield R.T., & Miyaki C.Y. (2013) Phylogeny and

comparative phylogeography of *Sclerurus* (Aves: Furnariidae) reveal constant and cryptic

diversification in an old radiation of rain forest understorey specialists. *Journal of*

Biogeography, **40**, 37–49.

eBird (2012) eBird: An online database of bird distribution and abundance [web application].

Elton C.S. (1927) *Animal Ecology.* Sidgwick and Jackson, London, United Kingdom,

Fjeldså J., Bowie R.C.K., & Rahbek C. (2011) The Role of Mountain Ranges in the

Diversification of Birds. *Annual Review of Ecology, Evolution, and Systematics*, **43**, 249–

265.

Freeman B.G. (2015) Competitive interactions upon secondary contact drive elevational

divergence in tropical birds. *The American Naturalist*, **186**, 470–479.

GBIF.org. (2015) GBIF Occurrence Download https://doi.org/10.15468/dl.39f5dl

Glor R.E. & Warren D. (2011) Testing ecological explanations for biogeographical boundaries.

Evolution, **65**, 673–683.

Grinnell J. (1917) The niche-relationships of the California Thrasher. *The Auk*, **34**, 427–433.

Guisan A., Lehmann A., Ferrier S., Austin M., Overton J.M., Aspinall R., & Hastie T. (2006)

Making better biogeographical predictions of species' distributions. *Journal of Applied*

Ecology, **43**, 386–392.

Hijmans R.J. (2015) raster: Geographic Data Analysis and Modeling. R package version 2.3-40. .

Hijmans R.J., Cameron S.E., Parra J.L., Jones P.G., & Jarvis A. (2005) Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology*,

25, 1965–1978.

Hijmans R.J., Phillips S.J., Leathwick J., & Elith J. (2015) dismo: Species Distribution

Modeling. .

Hutchinson G.E. (1957) Concluding Remarks. *Cold Springs Harbor Symposia on Quantitative*

Biology, **22**, 415–427.

Jang Y. & Gerhardt H.C. (2006) Divergence in the calling song between sympatric and allopatric

populations of the southern wood cricket *Gryllus fultoni* (Orthoptera: Gryllidae). *Journal of*

Evolutionary Biology, **19**, 459–472.

Khaliq I., Fritz S.A., Prinzinger R., Pfenninger M., Böhning-Gaese K., & Hof C. (2015) Global

variation in thermal physiology of birds and mammals: evidence for phylogenetic niche

conservatism only in the tropics. *Journal of Biogeography*, **42**, 2187–2196.

Kirschel A.N.G., Blumstein D.T., & Smith T.B. (2009) Character displacement of song and

morphology in African tinkerbirds. *Proceedings of the National Academy of Sciences*, **106**,

8256–8261.

Knowles L.L. & Massatto R. (2017) Distributional shifts - not geographic isolation - as a

probable driver of montane species divergence. *Ecography*, **40**, 1475-1485.

McCormack J.E., Zellmer A.J., & Knowles L.L. (2009) Does Niche Divergence Accompany

Allopatric Divergence in Aphelocoma Jays As Predicted Under Ecological Speciation?:

Insights From Tests With Niche Models. *Evolution*, **65**, 1–14.

Miller E.T., Wagner S.K., Harmon L.J., & Ricklefs R.E. (2016) Radiating despite a Lack of

Character: Ecological Divergence among Closely Related, Morphologically Similar

Honeyeaters (Aves: Meliphagidae) Co-occurring in Arid Australian Environments. *The*

American Naturalist, **189**, E000–E000.

Nunes L.A. & Pearson R.G. (2016) A null biogeographic test for assessing ecological niche

evolution. *Journal of Biogeography*, 1–7.

Orr M.R. & Smith T.B. (1998) Ecology and speciation. *Trends in Ecology & Evolution*, **13**, 502–

6.

Peterson A.T. (2011) Ecological niche conservatism: a time-structured review of evidence.

Journal of Biogeography, **38**, 817–827.

Peterson A.T., Soberón J., Pearson R.G., Anderson R.P., Martínez-Meyer E., Nakamura M., &

Araújo M.B. (2011) *Ecological niches and geographic distributions.* Princeton University

Press, Princeton, New Jersey.

Peterson A.T., Soberón J., & Sanchez-Cordero V. (1999) Conservatism of ecological niches in

evolutionary time. *Science*, **285**, 1265–1267.

Phillips S.J., Dudík M., & Schapire R.E. (2004) A maximum entropy approach to species

distribution modeling. *Proceedings of the Twenty-First International Conference on*

Machine Learning, 655–662.

Price T.D. (2008) *Speciation in Birds.* Roberts & Company Publishers, Greenwood Village,

Colorado.

QGIS Development Team (2016) QGIS Geographic Information System. *Open Source*

Geospatial Foundation Project, .

R Core Team (2016) R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Au.

Reino L., Ferreira M., Martínez-Solano Í., Seguardo P., Xu C., & Barbosa A.M. (2017)

Favourable areas for co-occurrence of parapatric species: niche conservatism and niche

divergence in Iberian tree frogs and midwife toads. *Journal of Biogeography*, **44**, 88-98.

Remsen J.V.J. (2003) Family Furnariidae (ovenbirds). *Handbook of the Birds of the World: Vol.*

8. Broadbills to Tapaculos (ed. by J. del Hoyo, A. Elliott, and D.A. Christie), pp. 162–357.

Lynx Ediciones, Barcelona.

Ryan Shipley J., Contina A., Batbayar N., Bridge E.S., Peterson A.T., & Kelly J.F. (2013) Niche

conservatism and disjunct populations. *The Auk*, **130**, 476–486.

Saupe E.E., Barve V., Myers C.E., Soberón J., Barve N., Hensz C.M., Peterson A.T., Owens

H.L., & Lira-Noriega A. (2012) Variation in niche and distribution model performance: The

need for a priori assessment of key causal factors. *Ecological Modelling*, **237–238**, 11–22.

Skeels A. & Cardillo M. (2017) Environmental niche conservatism explains the accumulation of

species richness in Mediterranean-hotspot plant genera. *Evolution*, **71**, 582-594.

Soberón J. (2007) Grinnellian and Eltonian niches and geographic distributions of species.

Ecology Letters, **10**, 1115–1123.

Soberón J. & Nakamura M. (2009) Niches and distributional areas: Concepts, methods, and

assumptions. *Proceedings of the National Academy of Sciences*, **106**, 19644–19650.

Soberón J. & Peterson A.T. (2005) Interpretation of Models of Fundamental Ecological Niches

and Species' Distributional Areas. *Biodiversity Informatics*, **2**, 1–10.

Sullivan B.L., Wood C.L., Iliff M.J., Bonney R.E., Fink D., & Kelling S. (2009) eBird: An

online database of bird distribution and abundance [web application]. *Biological*

Conservation, **142**, 2282–2292.

Title P.O. & Bemmels J.B. (2018) ENVIREM: An expanded set of bioclimatic and topographic

variables increases flexibility and improves performance of ecological niche modeling.

Ecography, **41**, 291-307.

Urbanek S. (2013) rJava: Low-level R to Java interface. R package version 0.9-6.

Warren D.L., Glor R.E., & Turelli M. (2008) Environmental niche equivalency versus

conservatism: Quantitative approaches to niche evolution. *Evolution*, **62**, 2868–2883.

Wickham H. (2009) *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag, New York.

Wooten J.A. & Gibbs H.L. (2012) Niche divergence and lineage diversification among closely

related Sistrurus rattlesnakes. *Journal of Evolutionary Biology*, **25**, 317–328.

World Wildlife Fund (2010) Terrestrial ecoregions. *List of ecoregions.*

Figures

Figure 1. The estimated geographic distribution of different *Sclerurus mexicanus* sensu lato populations from and basemap from Cooper & Cuervo (2017) with an overlay of training points from this study. Question marks indicate areas in which the population's identity has not been corroborated with vocal data or genetic data.

Figure 2. Narrow (left) and broad (right) training areas for three study taxa, listed from north to south: *S. pullus, S. obscurior,* and *S. andinus*. In the narrow training areas, strict parapatry (*i.e.*, no sympatry) is enforced.

Figure 3. A comparison of broad models for *Sclerurus andinus* and *Sclerurus m. macconnelli*. The red (left) distribution is the random iterations of *S. andinus* compared to the true *S. m. macconnelli*, while the blue-green (right) distribution is the random iterations of *S. m. macconnelli* compared to *S. andinus*. Vertical dashed lines represent the critical values for this test (i.e. the point at which any test to the left has a *P* value of 0.05 or lower), while the solid vertical is the test statistic of the true ENM of *S. andinus* compared to the true ENM of *S. m. macconnelli*. As with many tests of montane vs. lowland taxa, the test was significant for the distribution of random lowland derived models, but not for random montane models.

432Figure 1.

435Figure 2.

438Figure 3.

439**Tables**

440Table 1. Pairwise comparisons of Schoener's D for all combinations of *S. mexicanus*

441populations. Significance is ranked thusly: 2 = comparison was significant for the tails of both

442bidirectional comparisons; 1 = test was significant for one of the bidirectional comparisons; 0 =

443test was not significant.

Populations	Broad *M* Results	Narrow *M* Results	Allopatric?	Parapatric?
andinus-bahiae	2	1	Yes	
andinus-macconnelli	1	1	Yes	
andinus-mexicanus	0	0	Yes	
andinus-obscurior	1	1		Yes
andinus-peruvianus	0	0		Yes
andinus-pullus	1	0	Yes	
bahiae-macconnelli	0	1	Yes	
bahiae-mexicanus	2	2	Yes	
bahiae-obscurior	1	1	Yes	
bahiae-peruvianus	1	0	Yes	
bahiae-pullus	2	0	Yes	
macconnelli-mexicanus	1	1	Yes	
macconnelli-obscurior	1	0	Yes	
macconnelli-peruvianus	1	1		Yes
macconnelli-pullus	1	1	Yes	
mexicanus-obscurior	1	1	Yes	
mexicanus-peruvianus	0	0	Yes	
mexicanus-pullus	0	1	Yes	
obscurior-peruvianus	1	1	Yes	
obscurior-pullus	1	1		Yes
peruvianus-pullus	0	1	Yes	

444
445Table 1.