A widespread oscillatory network encodes an aggressive internal state

Yael S. Grossman[2,*], Austin Talbot[2,7]*, Neil M. Gallagher[2,3], Gwenaëlle E. Thomas[3], Alexandra J. Fink[6], Kathryn K. Walder-Christensen[2] , Scott J. Russo[6], David E. Carlson[5,8,†], Kafui Dzirasa[1,2,3,4,9,†].

[1]Howard Hughes Medical Institute, Chevy Chase, Maryland 20815, USA; [2]Dept. of Psychiatry and Behavioral Sciences, [3]Dept. of Neurobiology, [4] Dept. of Neurosurgery, [5]Dept. of Biostatistics and Bioinformatics, Duke University Medical Center, Durham, North Carolina 27710, USA; [6]Fishberg Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, New York, *10029* USA; [7]Department of Statistical Science, [8]Dept. of Civil and Environmental Engineering, [9]Dept. of Biomedical Engineering, Duke University, Durham North Carolina 27708, USA;

*These authors contributed equally

†Senior authors; Contributed equally

Correspondence should be sent to:

Kafui Dzirasa, M.D. Ph.D.
Dept. of Psychiatry and Behavioral Sciences
Duke University Medical Center
Durham, NC 27710, USA
Email: kafui.dzirasa@duke.edu
Twitter: @KafuiDzirasa

or

David E. Carlson, PhD
Department of Biostatistics and Bioinformatics
Department of Civil and Environmental Engineering
Duke University
Durham, NC 27708, USA
Email: david.carlson@duke.edu

Keywords

aggression; social behavior; electome network; closed loop stimulation; machine learning; prefrontal cortex

Abstract

Social aggression is an innate behavior that can aid an organism in securing access to resources[1], or it can impair group function and survival in behavioral pathology[2-4]. Since many brain regions contribute to multiple social behaviors[5-7], expanded knowledge of how the brain distinguishes between social states would enable the development of interventions that suppress aggression, while leaving other social behaviors intact. Here we show that a murine aggressive internal state is encoded by a widespread network. This network is organized by prominent and synchronized theta (4-11Hz) and beta (14-30Hz) oscillations that relay through the prefrontal cortex, and couples to widespread cellular firing. Strikingly, network activity during social isolation encodes the trait aggressiveness of mice, and causal cellular manipulations known to impact aggression can bidirectionally regulate the network's activity. Finally, we use closed-loop stimulation of prefrontal cortex and causal mediation analysis to establish that the network is a mediator of aggressive behavior. Thus, we define a widespread network that encodes an aggressive internal state within and across mice.

Social behavior reflects the integration of sensory information with internal affective states. Many subcortical brain regions contribute to aggressive behavior in mammals including lateral septum (LSN) [8, 9], nucleus accumbens [2, 10], lateral habenula [11, 12], the ventrolateral portion of ventromedial hypothalamus [5, 13-17], and medial amygdala [3, 18]. Prefrontal cortex stimulation has been shown to mitigate aggressive behavior in both humans [19, 20] and rodents [21], implicating cortical regions in regulating aggression. Finally, sensory regions, such as those underlying olfaction, also contribute to aggression [22].

To appropriately regulate aggressive behavior, the brain must integrate information across these and other cortical and subcortical regions. Moreover, since many of these regions also regulate non-aggressive social behaviors [5, 23], the brain must ultimately utilize information from overlapping regions to segregate aggression from other social behavioral states. Though efforts have revealed several cellular-level processes within one of these regions that contribute to this mechanism [13, 16, 24], the complementary network level process that

integrates information across regions to distinguish aggressive states from prosocial states

remains unknown. Addressing this knowledge gap is of major importance as 1) mammals

regularly select from a repertoire of social behaviors based on external sensory cues to ensure

their survival [6, 25], and 2) a range of psychiatric disorders are broadly marked by a failure to

appropriately match behavior with evolving social contexts [26].

We implanted mice with microwire electrodes across eleven brain regions implicated in

regulating complex social behavior. We then recorded electrical activity from these brain

regions, concurrently, as mice engaged in social encounters that induced aggressive attack

behavior and non-attack social behavior. After confirming that statistical models based on

single brain regions could independently differentiate attack behavior from non-attack social

behaviors, we asked whether this code was also reflected at a network level, where

millisecond-timescale information was integrated across all the 11 brain areas. For this analysis,

we used a machine learning approach that models variations in natural patterns of activity

within and between implanted brain regions across seconds of time (a timescale that we

reasoned would allow us to capture socially-relevant internal states)[27]. Our approach also

tuned the model to optimally encode attack vs. non-attack social behavior.

We then applied multiple levels of validation of increasing generalization to our single-region

and network-level models [28]. 1) We tested whether the models generalized to data collected

from time periods that had not been used to train our model (i.e., hold out sessions). 2) We

tested whether our models generalized to new animals, and 3) we tested an orthogonal

behavioral context associated with aggression. Only the model that was based solely on activity

from ventral hippocampus and the network model survived this third level of validation. 4)

Next, we tested whether these two models encoded an aggressive state, more broadly, and not

simply attack behavior. Only the network model encoded the social context of mice during

periods immediately prior to or following aggressive and prosocial interactions. Moreover,

network activity prior to and following attacks correlated with trait aggressiveness on a mouse-

by-mouse basis. 5) Finally, we subjected the network to two high levels of experimental

validation, each based on causal cellular manipulations, that bidirectionally impact attack

behavior. Specifically, we developed a close-loop optogenetic stimulation approach that

detected when the brain transitioned into an aggressive state, on a second-by-second basis,

and stimulated the prefrontal cortex in a manner previously known to suppress attack behavior.

We also employed a designer receptor exclusively activated by designer drug (DREADD)-based

approach to selectively activate cells in ventral medial hypothalamus in a manner that has been

shown to induce attack behavior. These manipulations bidirectionally impacted network

activity. 6) As this latter manipulation was also performed in mice on a different genetic

background, we further confirmed the generalizability of our network-level findings. Thus, we

firmly establish a network-level architecture whereby the brain encodes an aggressive state in a

manner that generalizes across context and individuals.

Direct stimulation of prefrontal cortex broadly suppresses social behaviors

We initially focused our efforts to selectively suppress aggressive behavior on modulating the

medial prefrontal cortex (i.e., prelimbic and infralimbic cortex in mice), since this brain region

had been implicated in social behaviors [29-31]. Prior work had shown that optogenetic

stimulation of prefrontal cortex was sufficient to suppress attack behavior and increase non-

aggressive social behaviors in CD1 strain mice [21]. Further highlighting the translational

potential of such an approach, clinical studies had shown that direct transcranial stimulation of

prefrontal cortex decreased aggressive feelings in violent offenders [32], and in individuals with

methamphetamine use disorder [33].

We performed our optogenetic stimulation experiments during these social encounters using a

protocol modeled after prior work, where blue light stimulation is used to activate

channelrhodopsin 2 in the medial prefrontal cortex for the entirety of a social encounter

(473nm, 5mW, 20Hz, 3ms pulse width, Fig. 1a) [21]. Control experiments were performed using

yellow light stimulation (593.5nm, 5mW, 20Hz, 3ms pulse width), which does not activate

channelrhodopsin 2[34, 35]. CD1 mice show periods of attack behavior, defined by biting,

boxing (kicking/clawing), or tussling behavior, when a male C57BL6/J (C57) mouse is introduced

into their home cage [36]. On the other hand, they exhibit periods of non-attack social

interactions such as sniffing, grooming, or resting (placing nose or forepaws against the subject

mouse, but not moving) when exposed to a female intruder [16]. Consistent with the prior

report, medial prefrontal cortex stimulation suppressed attack behavior towards C57 male mice

and tended to increase non-attack social behavior [N=8; t_7=3.43; P=0.0055, and t_7=-2.35;

P=0.051 using a two-tailed paired t-test, for attack and non-attack behavior, respectively,

significance determined by a Benjamini-Hochberg false discovery rate (FDR) correction, Fig. 1a].

To ensure that this suppression of aggression was selective, we also tested mice in a second

social paradigm in which they interacted with a female. Importantly, the CD1 mice do not show-

attack behavior during this social context. Here, we found that cortical stimulation decreased

non-attack social behavior toward female intruders (t_7=3.647, P=0.0041, using paired t-test,

significance determined by FDR correction). Thus, medial prefrontal cortex stimulation was

unable to selectively suppress attack behavior. Rather, stimulation suppressed multiple types of

social behavior.

<u>Multiple brain regions fail to independently encode attack behavior across mice and contexts</u>

After failing to selectively suppress CD1 aggressive social behavioral by targeting the prefrontal

cortex, we set out to probe for brain regions that may exhibit such selectivity. We implanted

CD1 mice across multiple cortical and subcortical brain regions known to contribute to social

behavior, including infralimbic cortex[21], orbitofrontal cortex[37], prelimbic cortex[21], lateral

septum [8, 9], nucleus accumbens [2, 10], lateral habenula [11, 12], mediodorsal thalamus[38],

ventromedial hypothalamus [5, 13-17], medial amygdala [3, 18], ventral hippocampus [39], and

primary visual cortex.

Following surgical recovery, we recorded neural activity while the CD1 mice freely interacted

with an intact male C57 mouse and a female C57 mouse for 300 seconds each. We repeated

these encounters over six sessions, yielding a total of 1800 seconds of neural data and behavior

for each exposure (Fig. 1b, see also Supplemental Fig. S1). We recorded a subset of these CD1

mice (N=9 of the 20 mice) as they interacted with a castrated male mouse intruder. Since CD1

mice do not generally exhibit attack behavior towards castrated males [40], this encounter

provided neurophysiology data during additional non-attack social behaviors that were

unpaired with female sensory cues. We also acquired neural activity while the CD1 mice were

isolated in their home cage.

We first verified that each of the implanted brain regions encoded social behavior using

discriminative cross spectral factor analysis non-negative matrix factorization (dCSFA-NMF, see

Fig. 1e) [41]. dCSFA-NMF utilizes supervised machine learning to generate a statistical model

that is both descriptive [integrates brain local field potential (LFP) activity features across time]

and predictive (discovers networks that distinguish between types of external behavior). LFPs

reflect the activity of populations of neurons, and these signals can be consistently sampled

across mice. The electrical functional connectome networks (*Electome Networks*) generated

from dCSFA-NMF integrate LFP power (oscillatory amplitude across 1-56 Hz; a correlate of

cellular and synaptic activity within brain regions), LFP synchrony (how two regions' LFP

frequencies synchronize across time; a correlate of brain circuit function), and LFP Granger

synchrony (Granger causality testing; a correlate of directional transfer of information across a

brain circuit). Furthermore, dCSFA-NMF generates electome network activity scores (an

indicator of the strength of each network) at a temporal resolution sufficient to capture brain

states underlying the external behavior under observation (in this instance, a resolution of one

second). The electome networks are designed to learn patterns that explain interpretable

correlates of neural activity whose expression relate to measured behavior, facilitating an

overall interpretable model [28]. Any given brain region can belong to multiple electome

networks, and each electome network may incorporate any number of brain regions. dCSFA-

NMF thus integrates spatially distinct brain regions and circuits into electome networks that

encode behavior.

To explore whether there was a generalized activity pattern within individual regions that

encoded social behavior, we designed a series of dCSFA-NMF models based on LFP oscillatory

power in frequencies from 1-56Hz. Each single-region model was trained using observations

pooled from 20 CD1 mice to separate periods when mice were socially isolated from periods

when they were engaged in social behavior (e.g., attack behavior towards the intact males and

non-attack social behavior towards male and female mice). We trained our models with one

supervised network to discover the patterns of LFP activity that encoded social behavior. We

also trained three unsupervised networks to account for the remaining variance in neural

activity that was not directly related to social behavior (see methods for justification of

hyperparameter selection). We tested the accuracy of the models in new CD1 mice recorded

under all three conditions (N=9 mice).

We observed high accuracy in decoding periods of isolation from social behavior, using the

supervised network, for each implanted brain area (p<0.05 using Wilcoxon rank-sum,

significance determined by FDR correction for 44 comparisons, Fig. 1c). Next, we designed a

new series of models to separate one of the three social behavioral states from the other three

(e.g., attack behavior toward intact males vs. non-attack interactions with female, intact male,

and castrated male mice). These models were built using observations pooled from the same

20 CD1 mice, and again based on LFP power. We then tested the accuracy of the supervised

networks in the models for the same nine hold-out mice. Thus, we trained and tested the

model's generalizability to decode each class of social behavior from the other three for each of

the 11 implanted brain areas (i.e., 33 additional models). Using this approach, we found that

five of the brain region-based statistical models decoded attack behavior versus non-attack

social behavior: infralimbic cortex, lateral habenula, ventral hippocampus, medial amygdala,

and medial dorsal thalamus. V1 successfully decoded the non-attack male interaction from the

other social conditions as well (P<0.05 using one-tailed Wilcoxon rank-sum test, significance

determined by FDR correction for 44 comparisons, see Fig. 1c). None of the other implanted

brain regions showed this selectivity. Thus, five regions independently encoded attack behavior

vs. non-attack social behavior in a manner that generalized across mice.

Our broad goal was to identify a neural signature that could be used to suppress aggression

while leaving other social behaviors intact. Thus, we tested whether the putative aggression

codes we discovered for the five regions generalized to another context associated with

aggression. Specifically, urine from other male mice has been found to elicit aggressive and

dominance behavior in CD1 males [22, 42, 43]. As such, the most aggressive mice from the

training and testing groups (N=8) were allowed to freely explore a clean inanimate object or an

object covered in urine from another intact CD1 male mouse (seven sessions). We then tested

whether each of the five regions' putative aggression codes could distinguish periods where

mice explored the clean objects from those where mice explored objects covered in urine.

Though only ventral hippocampus model tended to decode behavior in this new context, none

of the brain regions showed statistically significant encoding at the mesoscale-level (LFP)

following multiplicity correction (AUC=0.56±0.06, P=0.04 for ventral hippocampus, using one-

tailed Wilcoxon rank-sum test, significance determined by FDR correction for 5 comparisons,

Fig. 1d).

<u>An aggressive state is encoded at the network-level</u>

After failing to robustly decode attack behavior using LFPs independently from any of the 11

brain regions, we established that a brain network integrated information across all the

implanted brain regions to encode an aggressive state. This network-level encoding mechanism

generalized to multiple new contexts associated with aggression. Critically, the network also

encoded attack behavior with a predictive efficacy that exceeded independent ventral

hippocampus activity.

For this analysis, we trained a new dCFSA-NMF model using data from all the implanted brain

regions. This model utilized LFP power for each region, and the coherence and Granger

directionality between them. The model utilized one supervised network that was trained to

encode periods of attack-behavior (positive class) vs. social behavior in castrated male and

female social context (negative class). We also included non-attack social behavior towards

intact males in the negative class to discourage the network from simply learning non-

aggressive sensory cues specific to the intact male (i.e., supervision, electome network #1; see

Fig. 1e). Based on our hyperparameter selection approach using the Bayesian Information

Criteria (see methods and Supplemental Fig. S2), seven additional unsupervised networks were

trained to account for the variance in neural activity that was not related to attack vs. non-

attack social behavior. We then validated our supervised network using the set of nine holdout

CD1 mice from our single area coding test analysis. Again, none of these mice were used to

train the electome networks. We found that the supervised network (network #1) successfully

discriminated between attack behavior and non-attack social behavior in the test mice (N=9,

Wilcoxon signed rank, p=0.0020, Fig. 1f). Incidentally, we also observed that one of the

unsupervised networks (Network #6) showed strong encoding in the nine hold-out test mice

(AUC=0.65±0.03, $P<5\times10^{-5}$ using Wilcoxon rank sum).

Attack behavior is indicative of an aggressive brain state. We also reasoned that it was possible

for a mouse to be in an aggressive internal state, even though it was not actively exhibiting

attack behavior. Since such a context was likely to be present immediately prior to or following

attack behavior, we tested whether network activity pooled from the 3 seconds preceding and

3 seconds following social behaviors encoded the distinct social conditions (Fig. 2a). Critically,

these data windows were not used to train the network model since they did not contain attack

or non-attack social behavior. Activity of the supervised network (network #1) was lower in the

intervals surrounding attack behavior compared to non-attack social interactions with males or

females ($F_{3,67}$=28.6, P<0.0001 using Friedman's test, followed by P<0.05 using Wilcoxon signed-

rank test, significance determined by FDR correction, Fig. 2b, top left). Strikingly, network #1

activity during periods of isolation also negatively correlated with the time mice spent

exhibiting attack behavior towards other males (R=-0.58; P=0.016 using Spearman's rank

correlation, Fig. 2b, top right), encoding aggression on a mouse-by-mouse basis. Thus, electome

network #1 [hereafter referred to as *EN-Aggression Inhibition (EN-AggINH)*] represented a

network that putatively inhibited aggression when its activity was highest. In contrast to the

models developed for each of the brain regions independently, *EN-AggINH* activity also

encoded the exposure to male urine (N=8, Network Activity = 9.1±1.0 and 8.3±1.1 for clean and

urine covered objects, respectively; P=0.012 using one tailed Wilcoxon signed-rank test;

AUC=0.57±0.02, data not shown). Network #6 activity failed to generalize to this urine context

(AUC=0.48±0.01, data not shown). Thus, only *EN-AggINH* generalized to this second aggression

context.

Next, we tested the model for ventral hippocampus since we observed a trend towards

decoding attack behavior in the urine context. This model failed to encode the aggressive state.

Specifically, activity from ventral hippocampus was statistically indistinguishable between

periods surrounding attack behavior and non-attack social interactions with females and

activity during social isolation ($F_{3,67}$=7.9, P=0.047 using Friedman's test; P=0.36 and 0.72,

respectively, using Wilcoxon signed-rank test, significance determined by FDR correction, Fig.

2b, bottom left). Moreover, there was no relationship between ventral hippocampal activity

during isolation and the innate aggressiveness of mice (P=0.60 using Spearman's rank, Fig. 2b,

bottom right). Thus, only network activity encoded the aggressive internal state.

EN-Aggression Inhibition activity couples to cellular firing

EN-AggINH was composed of prominent theta frequency activity (4-11 Hz) in medial amygdala

and beta frequency activity (14-30 Hz) in medial amygdala and prelimbic cortex (Fig. 3a-b).

Prominent synchrony was also observed in the theta and beta frequency bands. Indeed, when

we quantified directionality across these synchronized bands, we saw that activity flowed from

orbital frontal cortex and primary visual cortex, relayed through medial dorsal thalamus, and

infralimbic cortex, and flowed into medial amygdala and ventral hippocampus (Fig. 3c-d, and

Supplemental Fig. S3).

We verified that the activity of *EN-AggINH* truly reflects biological activity, by relating the

electome network to neural firing, as in previous work [44]. To achieve this, we analyzed the

correlation between cellular activity across the implanted brain regions and the activity of *EN-*

AggINH, as cell activity is an undisputed measure of biological function. We then used a

permutation test to rigorously test our findings (Fig. 3e). Specifically, we shuffled cellular firing

within social behavioral conditions, maintaining the relationship between cell firing and

behavior. We then repeated this procedure 1000 times to generate a null distribution for which

only 5% of cells would be expected to exhibit firing coupled to network activity. We found that

~18% of cells showed firing that was coupled to the activity of *EN-AggINH*, far more than could

be explained by chance (χ^2=16.4, p=0.00005). Specifically, of the 186 cells recorded, nine

(4.8%) showed firing activity that was positively correlated with *EN-AggINH* and 25 (13.4%)

showed activity that was negatively correlated (Fig. 3f). Thus, most cells that showed coupling

to *EN-AggINH* were inhibited when network activity increased. These analyses confirmed that

EN-AggINH activity reflects the dynamics of cellular activity across the brain.

EN-Aggression Inhibition generalizes to new biological contexts related to aggression

To further validate *EN-AggINH*, we established that activity in this network was modulated by

orthogonal biological conditions that have been shown to induce or suppress aggressive

behavior in mice. In most cases, we performed this analysis in new animals, which is considered the gold standard of model validation in machine learning [45]. We transformed LFP data recorded from these new sessions into our original network model.

For our first gold-standard validation experiment, we tested whether our network generalized to new mice on a different genetic background engaging in a new aggression context (Fig. 4a-b). Specifically, this approach also used a validated cellular manipulation that causally induces aggression under a behavioral condition that would otherwise not yield aggressive attack behavior (i.e., we used female social partners). We expressed an excitatory DREADD (AAV-hSyn-DIO-hM3Dq) in the ESR1+ cells of ventromedial hypothalamus, since it has been shown that direct excitation of these cells induces aggressive behavior towards female mice [17, 46, 47]. Experiments were performed in the male F1 offspring of female CD1 strain mice crossed with ESR1-Cre male mice on a C57 strain background. Subsequently, we implanted the mice with recording electrodes to target the same brain regions as our initial experiment used to train the network model. Following recovery, we performed behavioral and neural recording when mice were exposed to a female mouse. The experimental mice were either treated with saline or CNO (Clozapine N-oxide, which activates the excitatory DREADD), in a pseudorandomized order, prior to the repeated testing sessions.

As anticipated, treatment with CNO induced attack behavior towards the female mice (N=8; P= 0.0039 for both attack latency and attack number using one-tailed Wilcoxon sign rank; Fig. 4c). When we probed neural activity across the entire exposure to the female intruder, we found that treatment with CNO also suppressed *EN-AggINH* activity (N=8, P=0.0039 using one-tailed Wilcoxon sign-rank; Fig. 4d). Thus, the network model generalized to a second aggression context induced by a cellular manipulation, and was robust to different genetic backgrounds. Critically, network activity was also lower during the time intervals surrounding attack/non-attack social behavior for the CNO vs. saline treatment sessions (P=0.02 using one-tailed Wilcoxon sign-rank; Fig. 4d), again demonstrating that *EN-AggINH* encoded an aggressive state. Our observations also established that *EN-AggINH* does not simply encode sensory cues associated with male intruders, since the network responses observed in the CNO treated mice were induced by a female intruder.

EN-Aggression Inhibition mediates attack behavior

We used mediation analysis to determine whether *EN-AggINH* activity putatively played a

mechanistic role in suppressing attack behavior. Mediation analysis is a framework to

determine whether the impact of a "treatment" (manipulation) on an outcome (attack

behavior) is mediated by a change in an intermediate variable (*EN-AggINH* activity). If so, the

intermediate variable is viewed, at least in part, as a mechanistic route (a mediator) for how the

treatment impacts the outcome. Three components were necessary to optimally implement

test our mediation analysis models: a manipulation that causally modulated 1) attack behavior

and 2) *EN-AggINH* activity, and 3) an approach to deliver the manipulation during levels of *EN-*

AggINH activity that would predict the emergence of attack behavior. We chose to build such

an approach based on prefrontal cortex optogenetic stimulation, since we had previously found

that such a manipulation causally suppressed attack behavior [21].

Specifically, we set out to preferentially stimulate medial prefrontal cortex when *EN-AggINH*

was naturally suppressed in the brain (signaling the onset of attack behavior). First, we built a

closed-loop system that estimated the activity of *EN-AggINH* in real time (i.e., within 200ms,

Fig. 5a). This approach employed a new network encoded solely based on power and coherence

measures (i.e., a reduced network, Fig. 5b), because the processing time to calculate Granger

directionality was prohibitive for real-time implementations. While this new network lacked the

interpretive power of dCSFA-NMF, it enabled us to predict attack behavior in real time (Fig. 5c).

In principle, when the activity of *EN-AggINH* fell below an established threshold (signaling the

onset of attack behavior), our closed-loop approach would deliver a one-second light

stimulation (5mW, 20Hz, 3ms pulse width) to prefrontal cortex. To verify that this real-time

estimation system worked as designed, we tested whether light stimulation was triggered by a

decrease in *EN-AggINH* activity. Indeed, network activity was significantly lower one second

prior to stimulation than it was two seconds prior to stimulation, demonstrating that our

approach successfully identified when the *EN-AggINH* activity decreased below the threshold

that signaled the onset of aggression (N=9; P<0.005 for within-subject comparison of *EN-*

AggINH activity 1 vs. 2 seconds prior to yellow light stimulation using one-tailed signed-rank

test, Fig. 5d). Importantly, we found that prefrontal cortex stimulation acutely increased *EN-*

AggINH activity (N=9, P<0.01 for comparison of *EN-AggINH* activity one second after blue vs.

yellow stimulation, using one-tailed signed-rank test, see Fig. 5d). Thus, our closed-loop

stimulation approach satisfied two of the components needed to implement our mediation

approach. Next, we tested whether increasing *EN-AggINH* activity via prefrontal cortex

stimulation as the brain transitioned into a putative attack state would suppress aggressive

behavior. We found our closed-loop stimulation approach significantly suppressed attack

behavior (see Fig. 6a; N=9 mice that were not used to train the initial model; t_8=6.1, P=0.0003,

comparing blue vs. yellow light stimulation using two-tailed paired t-test for attack behavior,

significance determined by FDR correction). Thus, our closed-loop manipulation suppressed

attack behavior, satisfying the remaining component needed to implement our mediation

analysis approach.

We first used the classic Baron and Kenny approach [48] to determine whether *EN-AggINH*

activity mediates the effect of neurostimulation on aggressive behavior. According to this

statistical approach, there is a mediated effect of network activity on behavior if three

conditions are met: 1) stimulation modulates network activity, 2) network activity correlates

with behavior, and 3) modeling the behavior from network activity and stimulation together is

better than modeling behavior from stimulation alone. Indeed, we had identified a significant

direct effect of stimulation on attack behavior (P<0.005, see Fig. 6a) and network activity

(P<0.0005, Fig. 5d). To optimally match the conditions between the treatment and control

cases, we used windows during the closed-loop stimulation procedure where the laser was

triggered, and then compared blue laser stimulation (treatment) to yellow laser stimulation

(control). Thus, the data points used for our mediation analysis predicted imminent or ongoing

attack behavior, and network activity prior to the stimulation in both the control (yellow light)

and treatment (blue light) case were similar. A statistical model of behavior using network

activity and stimulation (see Fig. 5e, model 2) significantly outperformed the model using only

stimulation (see Fig. 5e, model 1; nested logistic regression models, P<0.01, likelihood ratio

test), satisfying the necessary conditions to show that *EN-AggINH* is a mediator.

After establishing that *EN-AggINH* activity mediated the impact of PFC stimulation on behavior,

we set out to evaluate the significance of the average causal mediation effect (ACME) and the

average direct effect (ADE) during the same stimulated closed-loop windows using causal

mediation analysis [49]. ACME is the causal effect of stimulation on behavior due to the change

in *EN-AggINH* activity (see. Fig 5e, model 3), and ADE is the causal effect on behavior from

prefrontal cortex stimulation not explained by the change in *EN-AggINH* activity. We found that

there was a significant ACME (P<0.01), but not a significant ADE (P=0.48). This analysis

suggested that *EN-AggINH* activation is the primary mechanism whereby prefrontal cortex

stimulation suppresses aggression.

Next, we tested models where *EN-AggINH* activity functioned as a biomarker, rather than a

mediator of attack behavior. In these models, the manipulation modifies another neural

process, which in turn, simultaneously impacts attack behavior and *EN-AggINH* activity (Fig. 5g,

model 5). First, we evaluated whether theta power in 11 different brain regions could serve as a

mediator in lieu of *EN-AggINH* activity (Fig. 5f, model 4). We chose this frequency band since it

was prominently featured in *EN-AggINH* and within the network we previously found to encode

social appetitive behavior[7]. Across these 11 models, only orbitofrontal cortex had a significant

average causal mediation effect (*uncorrected* p-value of 0.038, see Fig. 5f). Critically, this model

did not survive a correction for multiple comparisons, and its ACME estimate was dwarfed by

the size of the ACME estimate for *EN-AggINH* (the estimate for the EN-AggINH model was

49.7% larger). This evidence suggests that EN-AggINH is a much better mediator than any of

these other potential 'biomarkers' by themselves.

After failing to identify any significant mediation effect of theta activity within each of the

eleven brain regions, we tested whether including theta activity as an intermediary in our

causal graph would disrupt *EN-AggINH's* role as a mediator in attack behavior (Fig. 5g, model 5).

Here, we corrected for the role of theta power in the model of how EN-AggINH changes as a

function of stimulation, as well as correcting for theta power in forecasting attack behavior. As

such, this framework dictates that *EN-AggINH* cannot mediate behavior that is already

explained by changes in theta power in a specified region. When we ran eleven models, one

model for each brain region, we found that *EN-AggINH* still significantly mediated attack

behavior in all of them (P<0.05 for all models; see Fig. 5g, bottom). Thus, even after accounting

for these potential intermediate variables, our findings still supported *EN-AggINH* as a mediator

of attack behavior.

<u>Validation of temporal activity and spatial spectral features of *EN-AggINH*</u>

We validated the temporal activity and spatial spectral features of *EN-AggINH* by establishing

that they could be utilized to selectively suppress aggression. Specifically, after determining

that our closed-loop manipulation suppressed aggression, we also quantified the impact of this

stimulation protocol on non-aggressive interactions with other male and female mice. We

found that closed-loop PFC stimulation increased non-attack behavior towards the intact C57

males [N=9; t_8=-2.3, P=0.049 comparing blue vs. yellow light stimulation using two-tailed paired

t-test for attack behavior and non-attack behavior, significance determined by FDR correction,

Fig. 6a). No differences in non-attack social behavior were observed during exposure to female

mice (t_8=0.74, P=0.48 using two-tailed paired t-test, significance determined by FDR correction,

Fig. 6a). Thus, closed-loop PFC stimulation selectively reduced aggression.

To verify that this selective modulation of aggression was due to synchronization of the light

stimulation with endogenous *EN-AggINH* activity, and not simply due to the dynamic pattern of

stimulation delivered using this method, we performed an additional control experiment where

we used the stimulation patterns from our closed-loop experiments to drive stimulation in a

new group of animals (e.g., randomly copying patterns from another mouses brain, analogous

to a "sham" in neurofeedback experiments). Thus, for these sessions, prefrontal cortex

stimulation occurred in a manner that mirrored our closed-loop stimulation experiments,

except that stimulation was not fixed to endogenous *EN-AggINH* activity (i.e., open loop –

nonsynchronous; Fig. 6b). Nonsynchronous stimulation failed to suppress aggressive behavior

($F_{1,21}$=4.87, P =0.039 for light type × stimulation pattern effect for post-hoc analysis using a

mixed effects model two-way ANOVA; t_{13}=0.09, P=0.93 for nonsynchronous stimulation using

paired t-test; see Fig. 6b), verifying that the suppression of attack behavior driven by closed-

loop stimulation was indeed due to delivery of stimulation timed to endogenous *EN-AggINH*

activity. Incidentally, nonsynchronous stimulation had no impact on non-attack social behavior

towards intact males or females (N=14; t_{13}=1.79, P=0.097; and t_{13}=0.54, P=0.60, for interaction

with males and females, respectively, comparing blue vs. yellow light stimulation using two-

tailed paired t-test). Thus, we validated the temporal activity component of *EN-AggINH.*

After establishing that we could selectively reduce aggression by temporally targeting PFC

based on the activity state of *EN-AggINH*, we tested whether we could reduce aggression by

spatially targeting stimulation based on the sub-components of PFC output circuity that

composed the network. We identified potential spatially specific targets by looking at the

relative LFP spectral Granger directionality from prefrontal cortex that occurred in the

aggressive internal state. Our initial visualization of *EN-AggINH* was constrained to the absolute

information flow at the strongest synchronies (Fig 3c-d). On the other hand, the relative

measures provide a measure of which circuits decrease their information flow prior to and

during attack behavior since *EN-AggINH* activity decreases during aggression (see Fig. 6c). In

other words, the relative Granger directionality measures quantified information flow pathways

that decreased the most during aggression. We focused our analysis on the Granger

directionality between PFC [prelimbic (PL) and infralimbic cortex (IL)] to nucleus accumbens

(PFC→NAc), medial amygdala (PFC→MeA) and lateral habenula (PFC→LHb), since *EN-AggINH's*

relative LFP spectral energy was highest for PFC→NAc and PFC→MeA circuitry and lowest in

the PFC→LHb circuit. Thus, a prominent decrease in information flow in PFC→NAc and

PFC→MeA circuitry was associated with aggression, while no such change was observed in PFC-

LH activity. Critically, all three circuits consisted of monosynaptic projections, enabling direct

targeting using optogenetics. We next quantified the relative spectral energy of these circuits at

20Hz since stimulating PFC at this frequency was sufficient to suppress the aggressive internal

state (Fig. 5d) and attack behavior (Fig. 1a). Given their representation in *EN-AggINH*, we

reasoned that driving PFC→NAc or PFC→MeA activity at 20Hz should selectively suppress

aggression, while driving PFC→LHb activity should not.

We causally activated these three circuits at 20Hz and measured their impact on social

behaviors. To selectively stimulate the terminals of PFC neurons in each target region (NAc,

MeA, or LHb), we injected mice with a retrograde AAV-Cre (rAAV-Cre) virus in one target region

and an AAV-DIO-channel rhodopsin-2 virus in PFC (N=8-9 per group). A stimulating fiber was

placed above the target region injected with rAAV-Cre. Social behavior was quantified during

20Hz stimulation with yellow vs. blue light (5mW, 20Hz, 3ms pulse width).

Blue light stimulation of PFC→NAc or PFC→MeA decreased aggression (t_8=2.4, P=0.04; t_7=5.9,

P=0.001 for NAc and MeA stimulation, respectively for blue vs. yellow light using two tailed-

paired t-test; N=8-9 mice per group, see Fig. 6d-e). This stimulation also increased non-attack

social behavior towards the male C57 mice (t_8=3.1, P=0.015; t_7=3.8, P=0.007 for NAc and MeA

stimulation, respectively). Neither of these stimulation protocols impacted social behavior

towards female C57 mice (t_8=1.2, P=0.27; t_7=0.8, P=0.46 for NAc and MeA stimulation,

respectively). On the other hand, PFC→LHb stimulation had no impact on aggression (t_7=0.38,

P=0.71; using two-tailed paired t-test, N=7 mice, see Fig. 6f), or non-attack social behavior

towards C57 males (t_7=0.24, P=0.82 using two-tailed paired t-test). Though this stimulation

protocol tended to increase social interaction with C57 females, these results did not reach

statistical significance (t_7=2.2, P=0.06 using two-tailed paired t-test). These results

demonstrated that directly stimulating the PFC subcircuits that normally showed the greatest

decreases in aggression-related activity causally and selectively suppressed aggression. On the

other hand, stimulating a PFC subcircuit with minimal activity changes during aggression had no

impact on social behavior towards male mice. Thus, these findings validated the spatial spectral

features of *EN-AggINH.*

Discussion

Here, we set out to discover the internal state that regulates whether an animal will exhibit

aggressive or pro-social behavior. We reasoned that attack behaviors emerge from an

aggressive internal brain state. Thus, we used machine learning to discover the mesoscale

neural architecture of the brain when an animal exhibited attack vs. non attack social

behaviors. Like other well-defined internal brain states, such as sleep, we found that the

network distinguishing attack behavior incorporated state-dependent patterns of neural

activity across every brain region we measured. For multiple regions, differences were

observed in local oscillatory power, while others exhibited differences in oscillatory synchrony

with a broader collection of regions. Each brain region showed selectivity in the frequencies of

oscillations that contributed to the network. For example, prelimbic cortex showed strong

activity in the beta frequency range, while medial amygdala showed strong activity in the beta

and theta frequency range. No brain region showed prominent activity contributions across all

frequencies. We also observed differences in the activity profile of a primary sensory region,

V1, which may reflect a change in encoding, or differences in visual sensory input observed

during attack behavior. Critically, the brain state identified during attack behavior was better

captured by the activity across all recorded brain regions as an integrated network, rather than

the independent activity within each brain region.

Though behavioral output has been classically utilized to infer the internal state of a brain, we

reasoned that an internal brain state was also likely present during intervals immediately

preceding and following behavioral output. Thus, we tested whether the aggression network

showed distinct activity profiles in the time intervals surrounding attack and non-attack social

behaviors. Indeed, network activity was lower during interval surrounding attack behavior.

Strikingly, we also found that network activity when animals were isolated in their home cage

encoded their trait aggression. Thus, the network did not simply encode behavioral output

since it was observed separately from attack behavior. Rather, the network encoded an

aggressive internal brain state.

Interestingly, this aggressive brain state was encoded by decreased activity in the network.

Given that we identified more cells that increased their firing rates as network activity

decreased, the discovery of a network that decreases its activity during aggression does not

indicate that overall brain activity is suppressed during aggressive states. Rather, these findings

argue that the aggressive state is encoded by a network that decreases its activity relative to

when mice are socially isolated or engaged in pro-social behavior. Indeed, our data suggested

that several common regions/circuits were activated during aggressive and pro-social behavior.

These common circuits need not be reflected in our network since our model was trained to

differentiate attack vs. non-attack social behavior. Nevertheless, our discovery of a network

that decreased its activity during aggression raises the intriguing hypothesis that the brain

actively inhibits aggression during pro-social engagement. When activity in this inhibition

network is suppressed, aggressive behavior emerges.

This interpretation is supported by our validation experiments where we directly activated

ESR1+-Cre neurons in ventromedial hypothalamus. Our findings showed that direct activation

of these cells induced the aggressive brain state (suppressed *EN-AggINH* activity). When mice

treated with CNO were exposed to a stimulus that would generally produce non-attack social

behavior (i.e., a female mouse), attack behavior emerged. Thus, the presence of the aggressive

brain state changed the mapping between sensory input and behavior output. Similarly, direct

stimulation of medial prefrontal cortex biased mice towards exhibiting non-attack social

behavior when they were exposed to a stimulus that would generally induce attack behavior

(i.e., a male intruder). Our findings showed that medial prefrontal cortex stimulation decreased

the aggressive internal state (increased *EN-AggINH* network activity). Critically, our findings

using mediation analysis argue that the brain state represented by *EN-AggINH* contributes to

the mediation of medial prefrontal cortex stimulation to a suppression of attack behavior.

Supporting this finding, our mediation analysis performed using data from the ESR1-Cre

experiment showed that *EN-AggINH* also mediated the impact of CNO treatment (see

Supplemental Fig. S5). Thus, *EN-AggINH* reflects the internal brain state that suppresses basal

aggression.

Here, we framed internal brain states as a collection of functions that transform sensory input

into behavior. Indeed, we found that when *EN-AggINH* activity is suppressed, the brain

transforms both male and female social sensory cues into attack behavior. It is also widely

appreciated that sensory input can also cause the brain to transition from one internal state to

another. For example, a loud sound can cause an animal to transition from sleeping to a hyper

aroused internal state. Along this line, we found that exposure to male mice could promote an

aggressive internal state in CD1 mice even prior to attack behavior, while exposure to a female

mouse did not (under normal conditions). In this framework, one would also anticipate that

many modulatory strategies that regulate attack behavior could mediate their effect by driving

the brain out of the state represented by low *EN-AggINH* activity. Indeed, we predict that

delivering a bright visual cue or a strong sensory cue (i.e., air puff) timed to decreases in *EN-*

AggINH activity could also potentially be used suppress attack behavior, since many circuits and

sensory inputs likely converge onto the internal state represented by *EN-AggINH*.

Our closed-loop stimulation approach was developed using a neural-network based

approximation technique for which the features were substantially constrained relative to

dCSFA-NMF. Nevertheless, we found that the reduced encoder was sufficient to identify the

precise time windows when the brain transitioned into aggression, as marked by a decrease in

EN-AggINH activity. In the future, novel approaches may allow for further improvement in the

precision of our real-time stimulation approach. For example, future work could exploit

convolutional neural networks to bypass the feature extraction step. These neural network

encoders could be altered to predict both aggressive and pro-social states, such as the

generalized social appetitive network that we recently discovered [7]. By using both networks

concurrently to actuate a closed-loop system, it may be possible to further suppress aggressive

behavior relative to pro-social behavior. Indeed, our current findings also pointed to a network

that exhibits increased activity during aggressive behavior (Electome Network #6, see Fig. 1f,

and Supplemental Fig. S6). Though the network failed to encode the urine paradigm, it is

possible that it contains activity that synergizes with *EN-AggINH* to encode aggressive social

states more optimally. If future studies demonstrate this potential, imitation encoders for both

Electome Network 6 and *EN-AggINH* could be integrated to further optimize closed-loop

approaches to selectively suppress aggression.

Multiple neuropsychiatric disorders including mood disorders, psychotic disorders,

neurodevelopmental disorders, and neurodegenerative disorders are associated with deficits in

regulating social behavior, including aggression. While multiple pharmacological approaches

have been instituted to suppress aggressive behavior towards self and others, many of these

strategies act by sedating the individual and can disrupt aspects of pro-social function. Our

discovery of a brain network that encoded an aggressive state raises the potential for novel

approaches to suppress aggressive behavior that spare pro-social behavior. Indeed, compared

to a standard open-loop stimulation protocol (20Hz stimulation) which suppressed both attack

and non-attack pro-social behavior, our closed-loop stimulation approach spared non-attack

social behavior towards males or females. Intriguingly, like other open-loop PFC stimulation

studies [50, 51], our 20Hz stimulation protocol induced behavioral hyperactivity in experimental

mice (see Supplemental Fig. S7). On the other hand, our closed-loop stimulation protocol did

not (see Supplemental Fig. S7). Thus, our findings also show that closed-loop stimulation may

limit off-target behavioral effects that are induced by classic stimulation approaches.

Overall, our findings establish a generalized network-level signature whereby the brain

suppresses aggression via active inhibition. Moreover, they highlight the exciting potential for

state-specific neuromodulation to regulate internal states.

Acknowledgements

We would like to thank Stephen Mague, Karim Abdelaal, and Ashleigh Rawls and Jean M. Zarate

for comments on this work; and Timothy Nyangacha for technical support. This work was

supported by WM Keck Foundation and Hope for Depression Research Foundation grants to

KD; NIH grants R01MH120158 to KD, 1R01EB026937 to DEC, and 1R01MH125430 to S.R., DEC,

and KD. A special thanks to Freeman Hrabowski, Robert and Jane Meyerhoff, and the

Meyerhoff Scholarship Program.

Figure Legends

Figure 1. A widespread network encodes attack behavior. a) Direct stimulation of prefrontal

cortex suppresses social behavior. Schematic of optogenetic stimulation (left) and social

encounters utilized for testing (middle). Prefrontal cortex stimulation suppressed attack

behavior, increased non-attack social behavior towards male mice, and suppressed non-attack

social behavior towards females (*$P<0.05$ for each comparison). **b)** Schematic for electrical

recordings, showing targeted brain regions (left), and representative local field potentials

(middle) recorded during repeated exposure to social contexts that produce attack and non-

attack social behavior (right). **c)** Framework to test individual brain regions' encoding of social

states (left). All implanted regions encoded social engagement; however, only five selectively

encoded the attack behavior vs. non-attack behavior (right). Pink shading indicates $P<0.05$ with

FDR correction. **d)** Attack codes discovered from the five brain regions failed to encode

aggressive behavior induced by male urine (gray shading indicates P<0.05 prior to but not

following FDR correction). **e)** Schematic of machine-learning model used to discover network

encoding attack behavior (left). The inputs to the model included LFP activity from the 11 brain

regions, the aggression class (+/-), and the social condition (IM-Intact male, CM-Castrated Male,

F-Female) for each 1-second data window. **f)** Encoding across eight learned networks. The

supervised network (purple, *EN-AggINH*) showed the strongest encoding. Data shown as mean

± SEM.

Figure 2. *EN-Aggression Inhibition* encodes an aggressive internal state. a) Neural activity was

sampled while mice were socially isolated (blue) and during intervals preceding and following

social behavior. **b)** Network activity during these intervals encoded attack behavior vs. male and

female non-attack social behaviors, while ventral hippocampal activity did not (P<0.05 using

Friedman's test followed by Wilcoxon sign-rank test). During isolation (blue) Network activity,

but not ventral hippocampus activity, encoded the subsequent total attack time of individual

mice (P<0.05 using Spearman's Rank Correlation).

Figure 3. Dynamics and biological components of *EN-Aggression Inhibition*. a) Prominent

oscillatory frequency bands composing *EN-AggINH* are highlighted for each brain region around

the rim of the circle plot. Prominent synchrony measures are depicted by lines connecting brain

regions through the center of the circle. The plot is shown at relative spectral energy of 0.4.

Theta (4-11 Hz) and beta (14-30 Hz) frequency components are highlighted in blue and green,

respectively. **b)** Example relative LFP spectral energy plots for three brain regions corresponding

to the circular plot in A (See Supplemental Fig. S3-4 for full description of network features). **c)**

Granger offset measures were used to quantify directionality within *EN-AggINH*. Prominent

directionality was observed across the theta and beta frequency bands (shown at spectral

density threshold of 0.4 and a directionality offset of 0.3). Histograms quantify the number of

leading and lagging interactions between brain regions. **d)** Schematic depicting directionality

within *EN-AggINH*. **e-f)** *EN-AggINH* maps to cellular activity. e) Three cells from LHb, VMH, and

MeA showing firing activity that is negatively correlated with *EN-AggINH* activity (red) and a

VHip cell showing positively correlated firing (blue). f) *EN-AggINH* activity correlated with

cellular firing across the brain across the brain. Single- and multi-units were used for analyses.

Figure 4. *EN-Aggression Inhibition* encodes distinct aggression contexts. a) Experimental approach for causally inducing aggression via direct activation of ESR1+ cells in ventromedial hypothalamus. b) Cellular activation induced attack behavior towards female mice (P<0.001 using sign-rank test), c) decreased *EN-AggINH* activity during social interactions with female mice (P<0.01 using one-tailed Wilcoxon sign-rank test) and d) intervals surrounding these interactions (P<0.05 using Wilcoxon sign-rank test).

Figure 5. *EN-Aggression Inhibition* activity is causally related to aggression. a) Schematic for closed-loop manipulation of *EN-AggINH* activity. **b)** Real-time estimation of aggression. Receiver operating characteristic depicting detection of aggressive behavior in a mouse using *EN-AggINH* activity vs. real-time reduced encoder is shown to the right. Dashed blue line corresponds to the established detection threshold. **c)** Detection of aggression using reduced encoder vs. *EN-AggINH* across mice (N=9; P=0.43 using two-tailed paired Wilcoxon sign-rank). **d)** *EN-AggINH* activity relative to light stimulation during closed-loop manipulation. Network activity significantly decreased one second prior to yellow light stimulation (N=9, [##]P<0.005 using one-tailed sign rank test; note that activity was normalized for each mouse to the average network activity during isolation). Activity was also higher one second after stimulation with blue light vs. yellow light ([**]P<0.01 using one-tailed signed-rank test). **e)** Directed graph with the inferred modes of action derived from mediation analysis. There is a causal relationship from stimulation to behavior and from stimulation to *EN-AggINH* expression (model 1; P<0.01 using signed rank and paired t-tests). *EN-AggINH* is a mediator from stimulation to behavior (P<0.01 using nested logistic regression models, likelihood ratio test; model 2), *EN-AggINH* activation is the primary mechanism whereby prefrontal cortex stimulation suppresses aggression (P<0.01 using average causal mediation effect, model 3). **f-g)** Directed graph testing f) local theta power as the primary mechanism whereby prefrontal cortex stimulation suppresses aggression (model 4) and g) *EN-AggINH* activation as the primary mechanism whereby prefrontal cortex stimulation suppresses aggression when local power is included as an intermediary (model 5). The uncorrected P values for each brain area in both models are shown below as -log(P).

Figure 6. Validation of spatiotemporal features of *EN-Aggression Inhibition*. a) Portion of windows stimulated during social behaviors using a closed-loop approach (P=0.002 using one-

tailed sign-rank test, left). Behavioral effects of closed-loop stimulation, right). **b)** Schematic for

nonsynchronous control stimulation (left). Nonsynchronous stimulation does not impact

aggressive or non-attack social behavior towards males or females. **c)** Granger Coherence for

PFC-dependent subcircuits within *EN-AggINH* (shown as relative spectral energy, see also

Supplemental Figure S4) **d)** Viral targeting strategy (left) and behavioral impact of PrL→NAc

circuit stimulation (right). **e)** Viral targeting strategy (left) and behavioral impact of PrL→MeA

circuit stimulation (right). **f)** Viral targeting strategy (left) and behavioral impact of PrL→LH

circuit stimulation (right). **P<0.005, *P<0.05 using two tailed paired t-test. Order of blue and

yellow light stimulation trials is show next to social condition diagrams.

Author contributions

Conceptualization and Methodology – YSG, AT, NMG, SR, DEC, and KD; Formal Analysis – YSG,

AT, NMG, AJF, KKW, DEC, and KD; Investigation – YSG, AT, NMG, GET, AJF, KKW, SR, DEC, and

KD; Resources – YSG, AT, NMG, GET, AJF, KKW, SR, DEC, KD; Writing – Original Draft, YSG, AT,

NMG, SR, DEC, and KD; Writing – Review & Editing, YSG, AT, NMG, KKW, SR, DEC, and KD;

Visualization – YSG, AT, DEC, and KD; Supervision –KKW, DEC, and KD; Project Administration

and Funding Acquisition –SR, DEC, and KD; See Supplemental materials for detailed author

contributions.

Declaration of Interests

The authors declare no competing interests

Materials and Methods

<u>Animal care and use</u>

All procedures were approved by the Duke University Institutional Animal Care and Use

Committee in compliance with National Institute of Health (NIH) Guidelines for the Care and

Use of Laboratory Animals. Mice were maintained on a reverse 12-hr light cycle with *ad libitum*

access to food and water.

Twenty-nine six-month retired breeder male CD1 strain mice (Charles River Laboratories,

Wilmington, Massachusetts) were used to discover a network that encoded aggression,

hereafter called *EN-AggINH*. Another fifty-five CD1 mice were used to probe the behavioral and

network responses to optogenetic stimulation. Mice were singly housed with enrichment.

ESR1-Cre mice on a C57/Bl6J background were provided by Scott Russo. These mice were

crossed with CD1 females in the Bryan Vivarium at Duke University to obtain F1 offspring. Eight

fourteen-week-old virgin ESR1-cre F1 male offspring were used to validate *EN-AggINH*. All F1

offspring were group-housed 2-5 mice per cage until they received viral injections in the

ventromedial hypothalamus at 7-8 weeks. After surgery, these mice were singly housed with

enrichment. All partner mice (C57BL/6J: two to seven intact males, two to seven females, and

two to seven castrated males per experimental mouse) were 7-14 weeks old. These mice were

purchased from Jackson Laboratories (Bar Harbor, Maine). All stimulus mice were housed 5 per

cage with enrichment. All behavioral testing and neurophysiological recordings occurred during

the dark cycle.

<u>Castration of C57 male mice</u>

Eighteen male mice were anesthetized with 1% isoflurane. The scrotal sac was sanitized with

betadine and 70% ethanol. The testes were then moved into the sac by gently palpating the

lower abdomen. Next, an incision was made in the sac and the testes were extracted. After

blood flow was cut off to the testes using a thread tourniquet, the testes were removed. The

remaining fatty tissue was placed back into the scrotum, which was then sutured. Mice were

allowed 10 days for recovery prior to experimental use.

<u>Electrode implantation surgery</u>

Mice were anesthetized with 1% isoflurane and placed in a stereotaxic device. Anchor screws

were placed above the cerebellum, right parietal hemisphere, and anterior cranium. The

recording bundles designed to target prelimbic cortex, infralimbic cortex, medial amygdala,

ventral hippocampus, primary visual cortex, mediodorsal thalamus, lateral habenula, lateral

septum nucleus, nucleus accumbens, ventrolateral portion of the ventromedial hypothalamus,

and orbitofrontal cortex were centered based on stereotaxic coordinates measured from

bregma. [Orbitofrontal cortex: anterior/posterior (AP) 2.35mm, medial/lateral (ML) 1.0mm,

dorsal/ventral (DV) from dura -2.75mm; infralimbic cortex and prelimbic cortex: AP 1.8mm, ML

0mm, DV -2.7mm from dura; medial amygdala: AP -1.25, ML 2.7mm, DV -4.3 from dura; lateral

septum and nucleus accumbens: AP 1.0mm, ML 0mm, DV -4.0mm from dura; ventromedial

hypothalamus, lateral habenula, and medial dorsal thalamus: AP -1.47mm, ML 0mm, DV -

5.4mm from dura; central hippocampus and primary motor cortex: AP -3.0mm, ML 2.6mm, DV -

3.0mm from dura]. We targeted infralimbic cortex and prelimbic cortex by building a 0.6mm DV

stagger into the bundle. We targeted lateral septum and nucleus accumbens by building a

0.3mm ML and 1.5mm DV stagger into the bundle. We targeted lateral habenula, medial dorsal

thalamus, and ventral medial hypothalamus by building a 0.3mm ML, and 1.9mm and 2.5mm

DV stagger into our electrode bundle microwires. We targeted primary motor cortex and

ventral hippocampus using a 0.3mm ML and 2.5mm DV stagger in our electrode bundle

microwires. For optogenetic stimulation experiments, the addition of a Mono Fiberoptic

Cannula coupled to a 2.5mm metal ferrule (NA: 0.22, 100mm [inner diameter], 125mm buffer

[outer diameter], MFC_100/125-0.22, Doric Lenses, Quebec) was built into the prefrontal

cortex bundle. The tip of the fiber was secured 300mm above the tip of the IL microwire. Mice

were allowed 10-15 days for recovery from surgery before behavioral testing.

<u>Viral surgery</u>

For optogenetic experiments targeting PFC soma [21], we used CD1 mice that showed an attack

latency < 60s when exposed to an intact C57 male. Thirty-five CD1 mice were anesthetized with

1% isoflurane and placed in a stereotaxic device. The mice were unilaterally injected with AAV2-

CamKII-ChR2-EYFP (purchased from the Duke Viral Vector Core, Durham, NC; courtesy of K.

Deisseroth), based on stereotaxic coordinates from bregma (left Infralimbic cortex: AP 1.8mm,

ML 0.3mm, DV -2.0mm from the brain). A total of 0.5mL of virus was infused at the injection

site at a rate of 0.1mL/min over five minutes, and the needle was left in place for ten minutes

after injection. For the open-loop stimulation experiment, CD1 mice were implanted with an
optic fiber (Mono Fiberoptic Cannula coupled to a 2.5mm metal ferrule (NA: 0.22, 100mm
[inner diameter], 125mm buffer [outer diameter], MFC_100/125-0.22, Doric Lenses, Quebec))
0.3mm above the injection site immediately after viral syringe was removed. These mice were
allowed 3 weeks for recovery prior to behavioral testing. For the closed-loop experiments, CD1
mice were allowed 3 weeks for viral expression prior to implantation with an optrode.

For the ESR1-Cre validation experiment, thirteen F1 offspring were bilaterally injected with
AAV2-hSyn-DIO-GqDREADD (obtained from Addgene) based on stereotaxic coordinates
measured from bregma (AP -1.5mm, ML ±0.7mm, DV -5.7mm from the dura). A total of 0.3mL
of virus was infused bilaterally at a rate of 0.1mL/min, and the needle was left in place for five
minutes after injection. Two weeks after viral infusion, F1 males were screened for aggressive
behavior towards females. The F1 males received i.p. injections of CNO (1mg/kg) at the start of
the screening session. Thirty-five minutes after injection, a novel C57 female was placed in the
home cage for 5 minutes. Screening was repeated one week and two weeks later. Only F1
males who attacked females for at least two of the three screening sessions (9/13 mice) were
implanted with electrodes [17]. The eight mice that showed good surgical recovery were
subjected to further experiments.

For PFC projection-targeting experiments, we used forty-four male CD1 mice that showed an
attack latency <60s and initiated attacks at least three times within three minutes when
exposed to an C57 male mouse. These mice were unilaterally injected with AAV2-EF1a-DIO-
ChR2-eYFP (obtained from Addgene) in the left prefrontal cortex based on stereotaxic
coordinates measured from bregma (AP 1.8mm, ML 0.3mm, DV -2.0mm from the dura), and
AAVrg-EF1a-Cre-mCherry (obtained from Addgene) was injected in the downstream target
region based on stereotaxic coordinates measured from bregma (NAc: AP 1.0mm, ML 0.9mm,
DV -3.8mm from the dura; MeA: AP -1.25mm, ML 2.75mm, DV -4.3mm from the dura; or LHb:
AP -1.6mm, ML 0.4mm, DV -2.2mm from the dura). A total of 0.3mL of virus was infused in the
prefrontal cortex and 0.3mL of virus was infused in the downstream target region at a rate of
0.1mL/min. The needle was left in place for five minutes after injection. Immediately after the
viral syringe was removed from the downstream target region, mice were implanted with an

optic fiber (Mono Fiberoptic Cannula coupled to a 2.5mm metal ferrule (NA: 0.22, 100mm

[inner diameter], 125mm buffer [outer diameter], MFC_100/125-0.22, Doric Lenses, Quebec))

0.3mm above the downstream target region injection site. Three weeks after viral

infusion/optic fiber implantation, CD1 mice were screened for aggression. Thirty-three

implanted mice that continued to show an attack latency <60s and initiated attacks at least

three times within three minutes were used for testing effects of projection targeted

stimulation on aggressive behavior.

<u>Histological analysis</u>

Histological analysis of implantation sites was performed at the conclusion of experiments to

confirm recording sites and viral expression. Animals were perfused with 4% paraformaldehyde

(PFA), and brains were harvested and stored for at least 96 hrs in PFA. Brains were

cryoprotected with sucrose and frozen in OCT compound and stored at -80C. Brains were later

sliced at 40μm. Brains from mice used to train and validate the network were stained using

NeuroTrace fluorescent Nissl Stain (N21480, ThermoFisher Scientific, Waltham, MA) using

standard protocol. Specifically, Nissl staining for brain tissue occurred on a shaker table at room

temperature. Tissue was washed in PBST (0.1% Triton in phosphate-buffered saline solution) for

10 minutes. It was then washed for five minutes in PBS twice. The tissue was then protected

from light for the remainder of the protocol. The tissue was incubated in 1:300 Nissl diluted in 2

mL PBS for 10 minutes. After the Nissl incubation, tissue was washed once in 0.1% PBST for 10

minutes, then twice in PBS for 5 minutes. Brains from ESR1 mice and mice used for 20 Hz or

closed-loop stimulation were mounted in Vectashield mounting medium containing DAPI (H-

1200-10, Vector Laboratories, Newark, CA). Images were obtained at 10x using an Olympus

fluorescent microscope. Of the 297 total implantation sites in the training and testing set of

mice, 17 were mistargeted (~5.7% error rate). Of these mistargeted implants, 13 were within

200μm of the targeted structure. Given our prior work demonstrating high LFP spectral

coherence (in the 1-55Hz frequency range) across microwires separated by 250um, in both

cortical and subcortical brain regions [44], we chose to retain these animals in our analysis. The

other four mistargeted implants were within 350μm of the targeted structure. The most

reliably mistargeted site was ventral medial hypothalamus for which 4 animals were implanted

within 200um of the target, and 2 animals were implanted within 350 μm of the target.

Machine learning analysis typically benefits from larger data sets. Thus, we concluded that

maintaining a higher number of data points likely outweighed the effect of a small number of

mistargeted brain regions, particularly since our LFP measures were robust to the targeting

inaccuracies we observed histologically. As such, we pooled data from all 20 implanted animals

to learn our initial model. We employed a similar strategy for our validation analysis, where an

animal was only removed from the validation set if there was clear histological confirmation of

mistargeting >200μm for any of the recorded regions. Specifically, presuming accurate targeting

with 94.3% certainty and targeting within 200um at a higher certainty, we included animals

with missing or damaged histological slices in our analysis. However, if there was clear

histological confirmation of mistargeting for any of the recorded regions (as was the case for 1

mouse), the animal was removed from the validation testing. Critically, our validation

procedure implies that the machine learning models were robust regardless of any slight

imprecision in the animals we utilized for training.

<u>Neurophysiological data acquisition</u>

Mice were connected to a data acquisition system (Blackrock Microsystems, UT, USA) while

anesthetized with 1% isoflurane. Mice were allowed 60 minutes in their home cage prior to

behavioral and electrophysiological recordings. Local field potentials (LFPs) were bandpass

filtered at 0.5-250Hz and stored at 1000Hz. An online noise cancellation algorithm was applied

to reduce 60Hz artifact (Blackrock Microsystems, UT, USA). Neural spiking data was referenced

online against a channel recording from the same brain area that did not exhibit a SNR>3:1.

After recording, cells were sorted using an offline sorting algorithm (Plexon Inc., TX) to confirm

the quality of the recorded cells. Only cell clusters well-isolated compared to background noise,

defined as a Mahanalobis distance greater than 3 compared to the origin, were used for the

unit-electome network correlation analysis. We used both single (well isolated clusters with

ISI<1.5) and multi-units (well isolated clusters with ISI<1.5; N=186 total neurons) for our

analysis as our objective was to determine whether electome network activity was reflective of

cellular activity. Neurophysiological recordings were referenced to a ground wire connected to
anchor screws above the cerebellum and anterior cranium.

<u>Behavioral recordings and analysis for training/testing models</u>

The CD1 mice used for training and testing the electome model were first subjected to
screening to assess their basal level of aggressiveness. Screening occurred once a day for three
consecutive days prior to surgical implantation. Animals were screened in cohorts. For each
screening session, an intact male C57 was placed in the CD1's home cage for 5 minutes and the
latency to first attack was recorded. To ensure that our network generalized broadly across CD1
mice, we used a training and testing set for which ~50% of the mice showed high aggression
during screening (i.e., latency to attack < 60s), and ~50% of the mice showed low to moderate
aggression (i.e., latency to attack > 60s). Animals that showed no aggression during screening
(16/45 mice) were excluded from further experiments.

All screening/testing occurred within the home cage of mice except for the quantification of
cortical stimulation-induced gross locomotor activity. These latter experiments were performed
in a 44cm × 44cm × 35cm (L×W×H) open field arena. Subject mice (CD1 and ESR1 males) were
acclimated to the recording tether for three days prior to the first recording session. Each
acclimation session involved anesthetizing the mouse with 1% isoflurane, tethering the subject
mouse, allowing 60 minutes to recover from isoflurane, then placing a male C57 in the home
cage for 5 minutes. Mice were then anesthetized with isoflurane again and detached from the
tether. The aggression level of experimental mice was determined based on average latency to
attack partner mice during the second and third acclimation sessions.

After screening, twenty-nine mice were implanted, and data was acquired across 1-6 behavioral
testing/recording sessions following recovery. Sessions were separated by 5-7 days. Recordings
for all social encounters were performed in the home cages of the CD1 mice. Each behavioral
testing session began with 5 minutes of recording prior to introduction of the social stimulus.
All mice were subjected to encounters with an intact C57 male mouse and a female C57 mouse.
A subset of eighteen CD1 mice were also subjected to an encounter with a castrated male
mouse, and another subset of eighteen mice were subjected to exposure to objects covered in

CD1 mouse urine. Object pairs included yellow duplex blocks, curved red duplex blocks,

weighted 5 mL conicals, glassware tops, and objects assembled from black legos®. The CD1

mice were exposed to a different pair of objects during each session. Order of exposure to

stimulus mice and objects was shuffled for every session. Six of the CD1 mice were recorded

under all four conditions. Data observations (1 second each) were pooled across eleven CD1

mice for training the network model. Object pairs included yellow duplex blocks, curved red

duplex blocks, weighted 5 mL conicals, glassware tops, and objects assembled from black

legos®. The CD1 mice were exposed to a different pair of objects during each session. Order of

exposure to stimulus mice and objects was shuffled for every session.

For ESR1 male behavioral testing, eight mice were injected with either saline or CNO (1mg/kg,

i.p.) after the five-minute baseline recording. Thirty-five minutes after this injection, mice were

exposed to an intact male C57, a castrated male C57, and a female C57, presented in

pseudorandom order. Mice were subjected to six total recording sessions (three in which they

were treated with saline and three in which they were treated with CNO), again in

pseudorandom order. Sessions were separated by 5 days to allow an adequate washout of

CNO[52].

Behavior was scored for each second as an "attack", "non-attack social interaction", or "non-

interaction". One-second windows were identified as "aggressive" if the mouse was engaged in

biting, boxing (kicking/clawing), or tussling behavior [36]. Windows were labeled as "non-attack

social interaction" if the mouse had his nose or forepaws touching the stimulus mouse (intact

male/female/castrated male) or object, but was not biting, boxing, or tussling. Examples of

behaviors labeled "non-attack social behavior" included sniffing, grooming, or resting (placing

nose or forepaws against the subject mouse, but not moving). If the stimulus mouse had

his/her forepaws or nose on the CD1, but it was not reciprocated, this was labeled "non-

interaction". CD1 straight approach, sideways approach, and chasing of the stimulus mouse

could result in attack (biting/kicking/tousling), non-attack social behavior (nose or paw touch),

or withdrawal without any touch. Thus, while sideways approach and chasing are regularly

labeled as "aggressive" in the literature [36, 53, 54], and straight approach is regularly labeled

as "pro-social", these behaviors lacked consistent resolutions. Moreover, mice also

demonstrated these behaviors towards female and castrated mice (non-attack social context).

One-second windows containing these behaviors were labeled "non-interaction". All other

timepoints not labeled "attack" or "non-attack social" were also labeled "non-interaction".

These behavioral criteria were selected to include ethologically aggression-related behaviors

and maximize the likelihood that the CD1 was aware of the presence of the stimulus mouse or

object during the behavioral window, while remaining confident in the classification of "attack"

and "non-attack social" window labels.

While tail rattling is not an attack behavior like the other behaviors that were labeled as

"attack", it was consistently only demonstrated by aggressive mice towards intact male mice.

Moreover, tail rattling is well-recognized in the literature as an aggressive behavior. Thus, we

included this behavior in the "attack" behavior category. In our subset of 20 mice used for

training the network, tail rattling was observed 8 ± 4s out of the 135 ± 26s "attack" windows

per mouse.

The videos used to generate the labels for training and testing our machine learning model

were hand-scored by a trained researcher. Videos from ESR1 mice and optogenetic stimulation

were automatically tracked using DeepLabCut [55, 56]. This information was then used for

creating behavioral classifiers in SimBA [57].

<u>LFP preprocessing and signal artifact removal</u>

Each LFP signal was segmented into 1s non-overlapping windows. If there were multiple intact

channels implanted in a region, they were averaged to produce a single signal. Windows with

non-physiological noise were removed using an established automated heuristic [7]. Briefly, the

envelope of the signal in each channel was estimated using the magnitude of the Hilbert

transform. The Median Absolute Deviation (MAD) of the magnitude was then calculated on

each channel of each recording. Signal was marked as non-physiological if the envelope

exceeded a high threshold (5x MAD, which is roughly 4x the standard deviation for a normally

distributed signal). Any data adjacent to non-physiological data that had an envelope value

above a smaller threshold (0.167 MAD) was also considered non-physiological. All data marked

in this way was ignored when averaging channels for each region. Any channels with standard

deviation less than 0.01 were removed as well. If no channels were usable for a given region
within a window, that whole window was removed from the data. This set of heuristics
resulted in 34.7±5.1% of the data being excluded from analysis. After this, 60Hz line artifact was
further removed using a series of Butterworth bandpass filters at 60Hz and harmonics up to
240Hz with a stopband width of 5Hz and stopband attenuation of -60dB. Finally, the signal was
downsampled to 500Hz.

<u>Estimation of LFP oscillatory power, cross-spectral coherence, and Granger directionality.</u>

Signal processing was performed using Matlab (The MathWorks, Inc. Natick MA). For LFP
power, spectral power was estimated using Welch's method using a 250-millisecond window
and 125-millisecond steps. Windows were zero-padded to give a 1Hz resolution. Cross-spectral
coherence was estimated pairwise between all regions using Welch's method and magnitude-
squared coherence defined as

$$C_{AB}(\omega) = \frac{|Psd_{AB}(\omega)|^2}{Psd_{AA}(\omega)Psd_{BB}(\omega)},$$

where A and B are two regions and $Psd_{AA}(\omega)$ and $Psd_{AB}(\omega)$ are the power and cross spectra
at a given frequency ω, respectively.

Spectral Granger Causality features were estimated using the Multivariate Granger Causality
(MVGC) MATLAB toolbox [58]. To get stable Granger Causality estimates, a 6th order highpass
Butterworth filter – with a stopband at 1Hz and a passband starting at 4 Hz – was applied to the
data using the *filtfilt* function (MATLAB, The MathWorks, Inc. Natick MA). Granger Causality
values for each window were estimated with a 20-order AR model at 1 Hz intervals to align with
the power and coherence features. Granger features were processed identically to a previously
reported approach [7]. Briefly, Granger features were exponentiated to approximately maintain
the additivity assumption made implicitly by NMF models [7, 59] as, $\exp(f_{A\rightarrow B}(\omega))$, where
$f_{A\rightarrow B}(\omega)$ is the Granger Causality at frequency ω from region A to region B. The exponentiated
feature is a ratio of total power to unexplained power. Exponentiated Granger feature values
were truncated at 10 to prevent implausible values.

<u>Data for single-region and network-level machine learning analyses</u>

We used 21460 seconds of data, pooled across the twenty mice, to train/validate our single

region and network models. This included a total of 4680 seconds while mice were socially

isolated in their home cage, 14890 seconds where CD1 mice exhibited non-attack social

behavior (3542 seconds towards intact males, 9067 seconds towards females, and 2281

seconds towards castrated), and 1890 seconds where mice exhibited attack behavior towards

the intact males.

<u>Discriminative Cross-Spectral Factor Analysis – Nonnegative Matrix Factorization</u>

The network was trained to distinguish between behavioral windows when the CD1 mice

showed aggressive behavior towards intact C57 males, and windows where they exhibited pro-

social behavior. These latter windows comprised pro-social interactions towards intact C57

males, castrated C57 males, or C57 females. Here, we used data from twenty-nine mice to learn

the final model, with a split of 20 and 7 for model training and internal validation.

We used Discriminative Cross-Spectral Factor Analysis – Nonnegative Matrix Factorization

(dCSFA-NMF) model [41]. This approach assumes each window of is an independent stationary

observation and examines dynamics in brain activity only at the scale of windows. A one-second

window was chosen to balance capturing fine-grained transient behavior with sufficient length

to properly estimate spectral features [7]. Each window has associated spectral power,

coherence, and Granger Causality features (in total $p = 9,586$ features), which is represented

as $x_i \in \mathbb{R}^p$ for the i^{th} window. Each window was associated with a behavioral label that

identified which condition the CD1 mouse was subjected to during that window (intact male,

castrated male, or female) and whether the CD1 mouse was engaged in aggressive or non-

aggressive behavior during that window, and the aggressive behavior was coded as $y_i \in \{0,1\}$.

As a short description of the dCSFA-NMF model, the features are described as an additive

positive sum of K non-nonnegative electrical functional connectome (electome) networks. This

model is learned using a supervised autoencoder. The objective we use to learn the parameters

is

$$\min_{W,d,\phi} \sum_{i=1}^{N} Loss_X\big(x_i, Wf(x_i; \phi)\big) + \lambda \, Loss_Y\big(y_i, df(x_i; \phi)\big) + \mu \, Loss_{EN}(A).$$

Here, $Loss_X$ is the reconstruction loss of the features derived from electrophysiology, which for

our work was an L_2 loss. Our predictive loss $Loss_Y$ is the cross-entropy loss commonly used for

binary classification. Each of the K networks is represented in vector form and combined to

make a matrix $W \in R^{p \times K}$. The electome network scores are given by the multi-output function

$f(x; \phi): R^p \to R^K$, where ϕ represent the parameters of the function. In our model, the multi-

output function was an affine transformation of $Ax + b$ followed by a softplus rectification,

defined as $softplus(x) = \log(1 + \exp x)$, thus $\phi = \{A, b\}$. Parameters $d \in R^k$ represent the

relationship between the electome networks and the behavior. A sparsity constraint is enforced

so that $d = [d_1, 0, ..., 0]$, meaning that only a single electome is used to predict behavior,

simplifying interpretation. λ is a weighting parameter used to control the relative importance

of prediction. We chose a value that kept the two losses approximately equal during training,

which corresponded to 1.

Previous work has also found that the reconstruction loss can reduce overfitting and make the

learned predictions more robust [60]. To further reduce overfitting of the predictive aspect of

the encoder, we applied an elastic net loss [61] on the encoder $Loss_{EN}$ with a weighting μ and

the ratio between the L_1 and L_2 losses set to .5. The value for μ was set to a small value that

had worked well previously. In this work, power features were also upweighted by a factor of

10 to accommodate that there were many more Granger features and truncated at 6 to

prevent outliers from dominating the predictions.

These models and statistical analyses were implemented with Python 3.7 and Tensorflow

version 2.4. Parameters were learned with stochastic gradient descent using the Nesterov

accelerated ADAM optimizer [62]. Learning was performed for 30000 iterations, which was

observed to be ample for parameter convergence. The learning rate and batch size were set to

1e-3 and 100 respectively, values that have empirically performed well in similar applications.

Predictive performance was evaluated in new mice not involved in learning the network. Given

processed data from the new mice, network scores were estimated as an evaluation of the

encoder learned during training of the dCSFA-NMF model.

<u>Hyper-parameter selection</u>

The dCSFA-NMF procedure requires selection of several settings in the algorithm. Specifically,

we must choose the number of electome networks K, the importance of the supervised task λ,

the relative importance of the power features, coherence features, and Granger features, and

the parameterization of the mapping function $f(x_i; \phi)$. Besides K, these settings were chosen

to match previously used values or follow heuristics. Specifically, in our prior work, we

demonstrated that the inferred model is highly insensitive to λ [27]. Thus, we chose a λ value to

give roughly equal weight to the predictive and generative tasks. Similarly, since the former

task grows linearly with brain regions and the latter task grows quadratically, we chose to

weight the power features to rough match the coherence features. Since the decoder is also

linear, we chose a linear mapping function followed by a softplus to ensure non-negativity. This

approach served to limit complexity.

To choose the value of K, we evaluated the reconstruction error (Mean Squared Error) on the

seven internal validation mice, which evaluates how well the electome networks describe the

neural measurements. As the goal for our analysis was to maintain high reconstruction and

effectively predict the behavior, an elbow analysis was used to choose the number of electome

networks K after which we observed minimal gains in explaining the data.

Specifically, our previous work has demonstrated that latent dimensionality is not an important

parameter in terms of predictive performance [27]. Thus, we trained one supervised network

for all the models tested in this study. We also trained multiple unsupervised networks for each

model to explain variance in brain activity that was not directly related to predictive

performance. Since our previous work had found that the supervised network has relatively low

variance, we used the Bayesian Information Criterion (BIC) to select the number of

unsupervised networks (latent factors d) to use in the final network model. The BIC is defined

as:

$$BIC(d) = k \log N - 2 \log(\hat{L}),$$

where $k = p \cdot d$ is the number of model parameters (p is the number of spectral features), N is

the number of samples, and $\hat{L} = p(X|\hat{\theta})$ is the likelihood of the observed data using the

estimated model parameters. This criterion balances the model fit quantified by $\hat{L}$ with the

complexity quantified by $k \log N$. In this work, $-\log(\hat{L})$ is an L_2 loss, corresponding to a

Gaussian observational likelihood. The model parameter $\hat{\theta}$ was estimated on 80% of the data

while model parameter $\hat{L}$ was evaluated on a 20% holdout set to avoid overfitting. The BIC was

evaluated for all dimensionalities from 1-20 networks, and the lowest value was selected as the

best model. Since 7 unsupervised networks provided the best fit (a BIC of 5457701, see also

Supplemental Fig. S2), our final network model utilized a total of 8 networks, 1 supervised and 7

unsupervised, across all 11 regions.

For each single-region model, we trained 3 unsupervised networks and a single supervised

network. Here, we reduced the number of networks as compared to the full network model,

given the dramatic reduction in the number of covariates considered by the model. Critically,

our objective was to compare the predictive performance of the single-region models against

each other and the full network model. Since the predictive performance is driven by the

supervised network[27], the smaller latent dimensionality of the single region models had no

impact on our final conclusions.

Single-region decoding

To test the efficacy of any single brain region as a biomarker for aggression, we extracted

power at 1 Hz frequency bins over 1-56 Hz from each region. One-second windows were pooled

from the twenty CD1 mice and used to generate a series of dCSFA-NMF models for each of the

11 brain regions. The models were trained to distinguish behavioral windows of one social state

exhibited by CD1 mice, from windows of two other social states. These three social states

included 1) male-directed attack, 2) female non-attack social interactions, and 3) castrated male

non-attack social interactions. We also developed a model to distinguish 4) periods where CD1

mice were isolated in their home cage from any of the three social states. Each model was then

tested on data from a holdout set of nine mice. The Area under the receiver operating curve

(AUC) was calculated for each holdout mouse to determine the performance of the model.

False discovery rate was used to correct for multiple hypothesis testing.

Validating Model Dimensionality

A frequent concern of latent variable models (including dCSFA-NMF) is the dependence of the networks and encoder on the choice of latent dimensionality. To address this concern, we performed a sensitivity analysis on the supervised network to determine the extent to which the choice of this dimensionality influenced the learned aggression electome network and encoder. In this sensitivity analysis, we estimated a dCSFA-NMF model allowing the number of total networks to range from two to twenty. We then compared the similarity between each learned encoder and decoder to our model with eight networks (the final model used in this work). This was quantified using the cosine similarity, which measures the angle between two networks (or encoders), ranging from -1 to 1. A value of 1 indicates perfect alignment (pointing in the same direction), 0 is orthogonal, and -1 indicates that the vectors point in opposite directions (Supplemental Figure S2).

We found that the supervised network maintained a strong consistency across most dimensions, particularly between 5-10 networks, as shown by the cosine similarities being greater than 0.95. The supervised encoders were virtually identical across all the models except the one that utilized three networks. This model learned a network that was positively associated with aggression.

To evaluate the robustness of the similarities across most of the supervised networks, we created a null distribution of the similarities across randomly chosen generative networks. These later similarities were substantially lower for both the network composition and the encoder. This indicates that as far as the supervised electome and encoder are concerned, latent dimensionality is not particularly influential on the resulting network, and by extension the biological interpretation.

Decoder Information Content

The amount of information contained in the predictive model was quantified by the reduction in uncertainty. The associated formula for this reduction in uncertainty, known as the Bernoulli entropy, is $1-1*(p \log_2(p) +(1-p)\log_2(1-p))$, where p is the accuracy of the model. At the extremes, an accuracy of 0.5 (random guessing) removes no uncertainty, whereas an accuracy of 1 or 0 completely eliminates uncertainty.

<u>Single-cell correlation to Electome Network activity</u>

Data acquired during the third behavioral testing session was from the twenty implanted mice

were used for cellular analysis. We used Spearman correlation to quantify the relationship

between cellular firing windows and the activity of the electome network used to classify attack

behavior. We performed 1000 permutations of randomly shuffling 1 second windows within

each class for attack and non-attack social interactions with male, female and castrated C57

mice. This approach maintained the relationship between network activity and behavior and

the relationship between cell firing and behavior. We then calculated the Spearman correlation

between network activity and cell firing for each permutation. A cell was deemed positively

correlated if its unshuffled Spearman Rho was above 97.5% of the permutated distribution and

negatively correlated if it was below 2.5%.

<u>Real-Time Encoder Approximation</u>

Because Granger Causality features were too computationally demanding for real-time

calculation, we developed a 'fast' dCSFA-NMF model that relied only on power and coherence

features for estimation of aggressive state to use in the closed-loop stimulation experiments.

This 'fast' model was trained on the same data. The model was trained using regularized

regression to best predict the output of the full encoder. As such, this reduced encoder is also

an affine transformation followed by a SoftPlus activation with a smaller parameter set,

$\phi_r = \{A_r, b_r\}$. This approximation explained a large component of the variance of the

supervised network score on the hold-out validation mice ($R = 0.47$, p-value $<10^{-16}$).

<u>Optogenetic stimulation</u>

Mice were anesthetized with 1% isoflurane, then tethered to an optic patch cable placed over

the optic fiber cannula. For closed-loop experiments, nine mice were also connected to the

recording system. The mice were then allowed 60 minutes for recovery prior to session

recording. For the fiber-only optogenetic stimulation experiments, CD1 mice experienced two

stimulation sessions. For closed-loop optogenetic stimulation, CD1 mice experienced three

sessions of behavioral screening followed by two sessions of closed-loop stimulation.

Stimulation sessions were separated by 5-7 days between sessions. For behavioral screening,

CD1 mice were exposed to intact C57 males, females, and castrated male mice for 5 minutes
each. Screening sessions two and three were used to determine a reduced network threshold
at which 40% of aggressive behavioral windows could be detected. For each session, mice were
recorded for 3 minutes of baseline in their home cage, then during the three social encounters.
Mice were recorded in an open field for 5 minutes after each session. The order of the three
social encounters were shuffled for each session. During each condition, the CD1 mouse
received segments of alternating blue (473nm, Crystal Laser LC, Reno, NV. Model No. DL473-
025-O) and yellow (593.5nm, OEM Laser Systems, Model No. MGL-F-593.5/80mW) light
stimulation, for two minutes each.

For open-loop stimulation targeting PFC soma, CD1 mice received light stimulation for the
entirety of the two-minute segment. For closed-loop, mice received stimulation for one second
when the reduced network score dropped below threshold.

For nonsynchronous stimulation, each of the fourteen CD1 mouse was psuedorandomly
matched to a different mouse that had been used for closed-loop stimulation. Each non-
synchronous mouse was then subjected to the identical order of conditions and yellow and blue
light stimulation blocks as their individually matched closed-loop mouse. Light stimulation was
delivered using the pattern implemented for the closed-loop partner mouse.

For projection targeting stimulation testing, CD1 males were exposed to one testing session
composed of three six-minute blocks of light stimulation. The sequence of light stimulation was
a yellow light stimulation segment, a blue light stimulation segment, then a final segment of
yellow light stimulation. Within each six-minute stimulation block, an intact male C57 and a
female C57 were sequentially placed in the CD1 cage for three-minutes each.

Immediately prior to experiments, light levels were calibrated using a power meter (ThorLabs,
Model No. P0025297 and 11070530), and delivered using a Waveform generator (Agilent
Technologies, Model No. 33210A) for the open-loop experiment. For closed-loop and
nonsynchronous stimulation experiments, the laser was activated using analog output from the
Cerebus recording system.

1128

<u>Mediation Analysis</u>

For the Baron and Kenny approach [48] to establish that *EN-AggINH* expression mediated the

behavioral effect of the neurostimulation, we first used two results previously described in the

methods to establish that there was an effect from the neurostimulation on network

expression and on behavior. We next constrained the data used to the most relevant case,

which is on the closed-loop stimulation. Specifically, we focused on windows of LFP data during

the closed-loop experiment when either the blue or yellow laser was activated to match the

cases between the treatment and control as closely as possible. As only blue light should

significantly manipulate neural activity, this is viewed as the treatment, and the yellow light is

set as the control. We followed the procedures outlined in "LFP preprocessing and signal

artifact removal" to preprocess the data and remove data with significant artifacts. *EN-AggINH*

expression was calculated by projecting the data into the learned model. The remaining data

was then fit into two logistic regression models to predict behavior using the statsmodel

package in python [63]. The first model used only the stimulation to predict behavior (behavior

~ const + stimulation), and the second model used stimulation and network expression to

predict behavior (behavior ~ const + network_expression + stimulation). These two models

were compared by using a likelihood ratio test to evaluate whether the second model was

significantly better.

For the causal mediation analysis, we again need to roughly balance treatment and control

groups. We used the same data as described above in the classic mediation analysis. We define

the treatment as blue versus yellow light stimulation, the mediator as *EN-AggINH* expression,

and the outcome as aggressive versus non-aggressive behavior. These data were then used in

the causal mediation analysis approach proposed by Kosuke, Keele, and Tingley [49] by using

the statsmodels package in python [63].

1153

<u>Statistics</u>

GraphPad Prism and Matlab were used for statistical analyses of behavior and network activity.

Paired T-tests were used for comparing within-subject behavioral response to optogenetic

stimulation or CNO application and corrected for false discovery rate for multiple hypothesis
testing through the Benjamini-Hochberg procedure. One-tailed Wilcoxon signed-rank tests
were used to compare within-subject mean network score responses to optogenetic
stimulation, stimulus mouse exposure and interaction, and CNO injection. Data is presented as
mean ± standard error of measurement, throughout the paper, unless otherwise specified.

Code and data availability

This learning algorithm is publicly available code at https://github.com/carlson-
lab/encodedSupervision. Data will be made available for replication purposes and pre-agreed
upon scientific extensions with a material transfer agreement.

1166

References

1. Lindenfors, P. and B.S. Tullberg, *Evolutionary aspects of aggression the importance of*
sexual selection. Adv Genet, 2011. **75**: p. 7-22.

2. Lee, S.C., T. Yamamoto, and S. Ueki, *Characteristics of aggressive behavior induced by*
nucleus accumbens septi lesions in rats. Behav Neural Biol, 1983. **37**(2): p. 237-45.

3. Haller, J., *The role of central and medial amygdala in normal and abnormal aggression:*
A review of classical approaches. Neurosci Biobehav Rev, 2018. **85**: p. 34-43.

4. Bak, M., et al., *The pharmacological management of agitated and aggressive behaviour:*
A systematic review and meta-analysis. Eur Psychiatry, 2019. **57**: p. 78-100.

5. Hashikawa, K., et al., *The neural circuits of mating and fighting in male mice.* Curr Opin
Neurobiol, 2016. **38**: p. 27-37.

6. Chen, P. and W. Hong, *Neural Circuit Mechanisms of Social Behavior.* Neuron, 2018.
98(1): p. 16-30.

7. Mague, S.D., et al., *Brain-wide electrical dynamics encode individual appetitive social*
behavior. Neuron, 2022. **110**(10): p. 1728-1741 e7.

8. Wong, L.C., et al., *Effective Modulation of Male Aggression through Lateral Septum to*
Medial Hypothalamus Projection. Curr Biol, 2016. **26**(5): p. 593-604.

9. Brayley, K.N. and D.J. Albert, *Suppression of VMH-lesion-induced reactivity and*
aggressiveness in the rat by stimulation of lateral septum, but not medial septum or
cingulate cortex. J Comp Physiol Psychol, 1977. **91**(2): p. 290-9.

10.   Couppis, M.H. and C.H. Kennedy, *The rewarding effect of aggression is reduced by*
nucleus accumbens dopamine receptor antagonism in mice. Psychopharmacology (Berl),
2008. **197**(3): p. 449-56.

11.   Flanigan, M., et al., *An emerging role for the lateral habenula in aggressive behavior.*
Pharmacol Biochem Behav, 2017. **162**: p. 79-86.

12.   Golden, S.A., et al., *Basal forebrain projections to the lateral habenula modulate*
aggression reward. Nature, 2016. **534**(7609): p. 688-92.

13.   Falkner, A.L., et al., *Decoding ventromedial hypothalamic neural activity during male*
mouse aggression. J Neurosci, 2014. **34**(17): p. 5971-84.

14.   Falkner, A.L., et al., *Hypothalamic control of male aggression-seeking behavior.* Nat
Neurosci, 2016. **19**(4): p. 596-604.

15.   Hashikawa, Y., et al., *Ventromedial Hypothalamus and the Generation of Aggression.*
Front Syst Neurosci, 2017. **11**: p. 94.

16.   Lin, D., et al., *Functional identification of an aggression locus in the mouse*
hypothalamus. Nature, 2011. **470**(7333): p. 221-6.

17.   Takahashi, A., et al., *Establishment of a repeated social defeat stress model in female*
mice. Sci Rep, 2017. **7**(1): p. 12838.

18.   Shibata, S., T.Y. Yamamoto, and S. Ueki, *Differential effects of medial, central and*
basolateral amygdaloid lesions on four models of experimentally-induced aggression in
rats. Physiol Behav, 1982. **28**(2): p. 289-94.

19.   Dambacher, F., et al., *Reducing proactive aggression through non-invasive brain*
stimulation. Soc Cogn Affect Neurosci, 2015. **10**(10): p. 1303-9.

20.   Choy, O., A. Raine, and R.H. Hamilton, *Stimulation of the Prefrontal Cortex Reduces*
Intentions to Commit Aggression: A Randomized, Double-Blind, Placebo-Controlled,
Stratified, Parallel-Group Trial. J Neurosci, 2018. **38**(29): p. 6505-6512.

21.   Takahashi, A., et al., *Control of intermale aggression by medial prefrontal cortex*
activation in the mouse. PLoS One, 2014. **9**(4): p. e94657.

22.   Jones, R.B. and N.W. Nowell, *The effect of urine on the investigatory behaviour of male*
albino mice. Physiol Behav, 1973. **11**(1): p. 35-8.

23.   Lischinsky, J.E. and D. Lin, *Neural mechanisms of aggression across species.* Nat
Neurosci, 2020. **23**(11): p. 1317-1328.

24. Nair, A., et al., *An approximate line attractor in the hypothalamus encodes an aggressive*
state. Cell, 2023. **186**(1): p. 178-193 e15.

25. de la Zerda, S.H., et al., *Social recognition in laboratory mice requires integration of*
behaviorally-induced somatosensory, auditory and olfactory cues.
Psychoneuroendocrinology, 2022. **143**: p. 105859.

26. APA, *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR*, ed. APA. 2000,
Washington, D.C.: American Psychiatric Association.

27. Talbot, A., et al., *Estimating a brain network predictive of stress and genotype with*
supervised autoencoders. J R Stat Soc Ser C Appl Stat, 2023. **72**(4): p. 912-936.

28. Vu, M.T., et al., *A Shared Vision for Machine Learning in Neuroscience.* J Neurosci, 2018.

29. Levy, D.R., et al., *Dynamics of social representation in the mouse prefrontal cortex.* Nat
Neurosci, 2019. **22**(12): p. 2013-2022.

30. Murugan, M., et al., *Combined Social and Spatial Coding in a Descending Projection from*
the Prefrontal Cortex. Cell, 2017. **171**(7): p. 1663-1677 e16.

31. Padilla-Coreano, N., et al., *Cortical ensembles orchestrate social competition through*
hypothalamic outputs. Nature, 2022. **603**(7902): p. 667-671.

32. Molero-Chamizo, A., et al., *Bilateral Prefrontal Cortex Anodal tDCS Effects on Self-*
reported Aggressiveness in Imprisoned Violent Offenders. Neuroscience, 2019. **397**: p.
31-40.

33. He, J., et al., *Effect of transcranial direct current stimulation over the left dorsolateral*
prefrontal cortex on the aggressive behavior in methamphetamine addicts. J Psychiatr
Res, 2023. **164**: p. 364-371.

34. Carlson, D., et al., *Dynamically Timed Stimulation of Corticolimbic Circuitry Activates a*
Stress-Compensatory Pathway. Biol Psychiatry, 2017. **82**(12): p. 904-913.

35. Mattis, J., et al., *Principles for applying optogenetic tools derived from direct*
comparative analysis of microbial opsins. Nat Methods, 2011. **9**(2): p. 159-72.

36. Miczek, K.A., et al., *Aggressive behavioral phenotypes in mice.* Behav Brain Res, 2001.
125(1-2): p. 167-81.

37. Coccaro, E.F., et al., *Amygdala and orbitofrontal reactivity to social threat in individuals*
with impulsive aggression. Biol Psychiatry, 2007. **62**(2): p. 168-78.

38.   Siegel, A., H. Edinger, and A. Koo, *Suppression of attack behavior in the cat by the*
1249   *prefrontal cortex: role of the mediodorsal thalamic nucleus.* Brain Res, 1977. **127**(1): p.
1250   185-90.

1251   39.   Chang, C.H. and P.W. Gean, *The Ventral Hippocampus Controls Stress-Provoked*
1252   *Impulsive Aggression through the Ventromedial Hypothalamus in Post-Weaning Social*
1253   *Isolation Mice.* Cell Rep, 2019. **28**(5): p. 1195-1205 e3.

1254   40.   Novotny, M., et al., *Synthetic pheromones that promote inter-male aggression in mice.*
1255   Proc Natl Acad Sci U S A, 1985. **82**(7): p. 2059-61.

1256   41.   Talbot, A., et al., *Supervised Autoencoders Learn Robust Joint Factor Models of Neural*
1257   *Activity.* arxiv, 2020.

1258   42.   Chen, A.X., et al., *Specific Hypothalamic Neurons Required for Sensing Conspecific Male*
1259   *Cues Relevant to Inter-male Aggression.* Neuron, 2020. **108**(4): p. 763-774 e6.

1260   43.   Lee, D.L. and J.L. Wilson, *Urine from Sexually Mature Intact Male Mice Contributes to*
1261   *Increased Cardiovascular Responses during Free-Roaming and Restrained Conditions.*
1262   ISRN Vet Sci, 2012. **2012**.

1263   44.   Hultman, R., et al., *Brain-wide Electrical Spatiotemporal Dynamics Encode Depression*
1264   *Vulnerability.* Cell, 2018. **173**(1): p. 166-180 e14.

1265   45.   Roelofs, R., et al., *A Meta-Analysis of Overfitting in Machine Learning.* Advances in
1266   Neural Information Processing Systems 32 (NeurIPS 2019) 2019.

1267   46.   Deonaraine, K.K., et al., *Sex-specific peripheral and central responses to stress-induced*
1268   *depression and treatment in a mouse model.* J Neurosci Res, 2020. **98**(12): p. 2541-2553.

1269   47.   Li, L., et al., *Social trauma engages lateral septum circuitry to occlude social reward.*
1270   Nature, 2022.

1271   48.   Baron, R.M. and D.A. Kenny, *The moderator-mediator variable distinction in social*
1272   *psychological research: conceptual, strategic, and statistical considerations.* J Pers Soc
1273   Psychol, 1986. **51**(6): p. 1173-82.

1274   49.   Imai, K., L. Keele, and D. Tingley, *A general approach to causal mediation analysis.*
1275   Psychol Methods, 2010. **15**(4): p. 309-34.

1276   50.   Anikeeva, P., et al., *Optetrode: a multichannel readout for optogenetic control in freely*
1277   *moving mice.* Nat Neurosci, 2011.

1278   51.   Kumar, S., et al., *Cortical Control of Affective Networks.* J Neurosci, 2013. **33**(3): p. 1116
1279   −1129.

52. Jendryka, M., et al., *Pharmacokinetic and pharmacodynamic actions of clozapine-N-oxide, clozapine, and compound 21 in DREADD-based chemogenetics in mice.* Sci Rep, 2019. **9**(1): p. 4522.

53. Miczek, K.A. and J.M. O'Donnell, *Intruder-evoked aggression in isolated and nonisolated mice: effects of psychomotor stimulants and L-dopa.* Psychopharmacology (Berl), 1978. **57**(1): p. 47-55.

54. Kwiatkowski, C.C., et al., *Quantitative standardization of resident mouse behavior for studies of aggression and social defeat.* Neuropsychopharmacology, 2021. **46**(9): p. 1584-1593.

55. Mathis, A., et al., *DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.* Nat Neurosci, 2018. **21**(9): p. 1281-1289.

56. Nath, T., et al., *Using DeepLabCut for 3D markerless pose estimation across species and behaviors.* Nat Protoc, 2019. **14**(7): p. 2152-2176.

57. Nilsson, S.R., et al., *Simple Behavioral Analysis (SimBA): an open source toolkit for computer classification of complex social behaviors in experimental animals.* bioRxiv, 2020.

58. Barnett, L. and A.K. Seth, *The MVGC multivariate Granger causality toolbox: a new approach to Granger-causal inference.* J Neurosci Methods, 2014. **223**: p. 50-68.

59. Gallagher, N.M., et al., *Cross-Spectral Factor Analysis* Advances in Neural Information Processing Systems, 2017. **30**.

60. Le, L., A. Patterson, and M. White, *Supervised autoencoders: Improving generalization performance with unsupervised regularizers.* NeuroIPS, 2018.

61. Zou, H. and T. Hastie, *Regularization and variable selection via the elastic net.* Journal of the Royal Statistical Society, 2005(67): p. 301-320.

62. Tato, A. and R. Nkambou, *Infusing Expert Knowledge Into a Deep Neural Network Using Attention Mechanism for Personalized Learning Environments.* Front Artif Intell, 2022. **5**: p. 921476.

63. Seabold, S., Perktold, J., *Statsmodels: Econometric and Statistical Modeling*
with Python. PROC. OF THE 9th PYTHON IN SCIENCE CONF., 2010.

1309

Figure 1

a

Isolation   Male   Female

Aggressive Internal State

3s

Attack Behavior   Attack Behavior

3s

Non-Aggressive Internal State

3s

Non-Attack Social Behavior   Non-Attack Social Behavior

3s

Figure 2

Figure 3

Figure 4

Figure 5