1   **Title: Parasite transmission in size-structured populations**

2   **Authors: Kelsey E Shaw[1], Rebecca E Cloud[3], Raeyan Syed[2], David J Civitello[2]**

3   1. Department of Biological Sciences, University of Notre Dame, South Bend, Indiana,
4   46556, USA
5   2. Department of Biology, Emory University, Atlanta, GA, 30322, USA
6   3. School of Integrative Biology, University of Illinois, Urbana-Champaign, IL, 61801,
7   USA
8
9   **Author emails:**
10   1. Kelsey E Shaw, kshaw7@nd.edu
11   2. Rebecca E Cloud, rcloud3@illinois.edu
12   3. Raeyan Syed, raeyan.syed@emory.edu
13   4. David J Civitello, david.james.civitello@emory.edu
14

15   **Contact/Corresponding author:** Kelsey Shaw

16   1510 Clifton Road NE Atlanta, GA 30322

17   Phone: (404) 727-7208

18   Fax: 404-727-2880

19   Email: kshaw7@nd.edu

Open Research Statement: All experimental data and novel code used for data analysis are
currently available on Figshare via this private link:
https://figshare.com/s/7d70b88220b912e7eec5. Data will be made public on Figshare upon
acceptance. Data set utilized for Figure 4 are from the following source:
Rumi, A., D. Gutiérrez Gregoric, and A. Roche. 2009. Tendencias Del Crecimiento Individual en
Poblaciones Naturales de Biomphalaria spp. (Gastropoda, Planorbidae) en la Cuenca del Plata,
Argentina. *Comunicaciones de la Sociedad Malacológica del Uruguay*. URL:
https://www.redalyc.org/articulo.oa?id=52414008002
29
30

31   **Keywords:** Parasite transmission; heterogeneity; body size; host traits; schistosomiasis;

32   transmission interference

Abstract

Host heterogeneity can impact parasite transmission, but determining underlying traits and incorporating them into transmission models remains challenging. Body size is easily measured and influences numerous ecological interactions, including transmission. In the snail-schistosome system, larger snails have a higher exposure to parasites but lower susceptibility to infection per parasite. We quantified the impact of size-based heterogeneity on population-level transmission by conducting transmission trials in differently size-structured snail populations and competing size-dependent transmission models. Populations with greater proportions of large snails had lower prevalence, and small snails were shielded from infection by co-occurring large conspecifics. Using the winning size-explicit model, we then estimated that schistosome transmission varies dramatically across time due to seasonal changes in snail population size structure. Thus, incorporating traits such as body size, which are impacted by and directly affect host ecology, into transmission models could yield insights for natural dynamics and disease mitigation in many systems.

I. Introduction:

Parasite transmission drives the ecological and evolutionary dynamics of host-parasite interactions. Therefore, demystifying the host and parasite factors that influence transmission is critical for characterizing disease dynamics and successfully managing any host-parasite system. In particular, transmission theory that excludes host variation and instead treats hosts as uniform individuals with identical infection risk matches data poorly and cannot address critical themes in disease ecology, such as superspreading, parasite aggregation, and host contact networks (Lloyd-Smith et al. 2005, Van der Waal and Ezenwa 2016). For example, early in the Sars-CoV-2 pandemic, models that did not include variability in traits like infectiousness or contacts often overestimated values for R_0, with implications for public health recommendations (Gomes et al. 2022). In order to more accurately predict disease dynamics in natural populations, it is important to consider the sources of host variation and their consequences for transmission because hosts can differ in many ways (Dwyer et al. 1997, Hall et al. 2007, Strauss et al. 2019).

Body size is a one host trait that can be highly heterogenous and may impact parasite transmission in many systems. For example, larger individuals may provide a greater target area for vectors (Poulin 2004, Yan et al. 2021), may provide more resources and thus tolerate a larger burden of macroparasites (Hechinger et al. 2019), may be more apparent to parasites that actively seek out their host using chemical cues (Théron et al. 1998, Mayer et al. 2021), or may encounter a greater abundance and diversity of parasites if they are exploring larger territories(Han et al. 2015). Furthermore, body size has the potential to vary drastically among individuals in a population due to age, genetics, feeding history, and/or environmental stressors (Kooijman,

S.A.L.M. 2010). The snail-schistosome system is an excellent study system for investigating the

impact of host variation in body size on parasite transmission.

Schistosomes are the causative agent of schistosomiasis, a Neglected Tropical Disease that is

second only to malaria in its worldwide impact on human health and disproportionately affects

individuals in poor, rural environments (Gryseels et al. 2006). Over 200 million people

worldwide are currently infected, and schistosomiasis accounts for approximately 200,000 deaths

per year (World Health Organization 2022). Schistosomes have a complex life cycle and

obligately cycle between mammalian and snail hosts. One source of host heterogeneity that

impacts schistosome transmission is snail body size: larger snails experience higher exposure

rates but are less likely to become successfully infected on a per parasite basis (Niemann and

Lewis 1990, Théron et al. 1998). Snail body size can be influenced by many factors such as age,

genetics, and resource availability, but this relationship between body size, exposure, and

susceptibility has been demonstrated to occur regardless of the underlying size-determining

mechanism (Richards et al. 1992, Larson et al. 2014). The importance of body size in

transmission to the snail host becomes even more apparent when considering the vast range in

host snail body size. For example, a neonate *Biomphalaria glabrata*, an intermediate host species

for *Schistosoma mansoni*, is born at ~0.75 mm and can grow up to 40 mm under natural

conditions over its lifetime (Jarne et al. 2011). This growth trajectory corresponds to a ~150,000-

fold increase in biomass, akin to the difference between a mouse and an elephant (White 2010).

Despite the recognition that body size influences these mechanistic components of schistosome

transmission decades ago in isolated, individual snails, the consequences of this relationship

when scaled to size-structured populations have remained uninvestigated. Schistosome exposure

in a host is irreversible, meaning a schistosome cannot leave a snail once it has entered, even if it

fails to successfully infect the host. Therefore, snails that are invaded by parasites indirectly

reduce the infection risk of others in the population (King et al. 2011). If size-dependent

transmission effects scale up to the population level, then we predict that the presence of larger

hosts will contribute to a within-species dilution effect by shielding their smaller conspecifics

from parasite exposure while also being less likely to become infected themselves.

We investigated this hypothesis by conducting laboratory transmission trials of *Schistosoma*

mansoni in *Biomphalaria glabrata* populations of varying size structures. We then used these

data to compete candidate models of size-dependent transmission, and additionally used existing

field data of *B. glabrata* population size structures to estimate how parasite success may change

under natural conditions over time solely due to the dynamics of population size structure. In

many host-parasite systems researchers are still in the nascent phase of breaking down the

complex interaction of host traits that result in heterogeneous parasite transmission (Sweeny and

Albery 2022). In the snail-schistosome system, body size is a measurable trait with the potential

to influence parasite transmission via multiple mechanisms, but it remains relatively unexplored

how this individual-level trait scales up to influence transmission within and between

populations. Incorporating this fundamental host trait into our understanding of schistosome

transmission could yield important insights into transmission risk and disease control of this

widespread and debilitating parasite, and this approach can be applied broadly to other disease

systems and critical traits.

II. Methods

Snail Maintenance. *B. glabrata* snails of the NMRI strain were maintained under favorable

conditions. Snails were kept in HHCOMBO artificial lake water (Baer and Goulden 1998) at

26°C with a 12:12 light:dark cycle. Snails were fed a diet of fish flakes (Omega One) and

chicken feed (Nutrena Meatbird Crumbles) suspended in 1% agar *ad libitum.*

Experimental Design. We conducted the experiment using a fully factorial design: five

population size-structures, each with a total of 18 snails, were exposed to three densities of

parasites in 24-hour transmission trials in 15-liter tanks, creating 15 total treatment combinations.

B. glabrata were first assigned into three size classes by shell diameter: "small" (2-3 mm),

"medium" (6-8 mm), and "large" (12-15 mm). Different size-structures were determined by the

ratio of snails from each of the three size classes. The uniform size structure consisted of 18

snails from the same size class: "uniform small" (1:0:0), "uniform medium" (0:1:0), and

"uniform large" (0:0:1). "Equal" size-structured mesocosms had 6 snails from each size class

(1:1:1). "Small skewed" size structures contained 12 snails from the smallest size class and 3

snails from both the medium and large size classes (4:1:1). Snail size classes and population size

structures were chosen to encompass the range of size variability observed in nature (see Fig 4)

and also to span a sufficient gradient to fit and evaluate size-dependent transmission models.

Parasite densities also varied between the tanks: 36 (2 miracidia/snail), 144 (8 miracidia/snail),

and 256 parasites per tank (14 miracidia/snail). We obtained *S. mansoni* eggs from

experimentally infected mice livers and purified and hatched eggs to obtain miracidia

(Dinguirard et al. 2018). We conducted four randomized blocks of these trials in which each

block contained one replicate of each treatment. To control for potential batch effects on parasite

infectivity across the temporal blocks, a control group consisting of all medium snails

individually placed in 24 well plates was simultaneously exposed to each of the parasite

densities.

To quantify size-based susceptibility we also conducted transmission experiments in which 120

snails ranging from 0.5mm to 15mm were individually placed in well plates with 5 mL COMBO.

These snails were exposed to either 2 or 6 miracidia over 24 hours. Due to the small volume of

water used in the individual well plate exposures, we assume that snails were exposed to all

parasites present in these trials, an assumption justified by prior experiments (Niemann and

Lewis 1990).

Infection Diagnosis. 24 hours post-exposure, all snails were collected, sorted by size and

treatment group, and maintained for 5 weeks (longer than the minimal prepatent period for *S.*

mansoni in favorable conditions). Snails were diagnosed visually by parasite shedding in

individual well plates at 4- and 5-weeks post-exposure to obtain prevalence data (Asch 1972).

Snails were diagnosed as "infected" if we observed *S. mansoni* cercariae in their well plate, or

"uninfected" if no cercariae were detected. Snails infected at week 4 were sacrificed, while

uninfected snails were returned to their tanks to repeat this process at week 5. Snails that died

while prepatent or between weeks 4 and 5 of the shedding process were not included in our

results (6.25% of all snails).

To compare prevalence data amongst the various size structures and size classes, we used the

glmmTMB package in R (Brooks et al. 2017) to create a generalized linear mixed model

(GLMM) accounting for the random effects between individual tanks and exposure dates

(bionomial probability distribution with a logit-link function). We then used the emmeans

package to conduct an estimated marginal means (least-squares means) *post hoc* test (Lenth

2022).

Model Creation and Maximum Likelihood Estimation of Parameters. We then built

deterministic size-dependent transmission models for parameterization and competition using the

experimental data (Box 1). The models used coupled ordinary differential equations to track the

densities of susceptible snails (S_i) and infected snails (I_i) in each size class (i) as well as free

living parasites in the water (P). In our experiments, snails and parasites were all added at t_0 and

hosts were removed 24 hours later at t_1, therefore we only focus on a single timestep and do not

include processes on a longer time scale such as host birth and death. Each model separates the

transmission process into two underlying processes: exposure to parasites and susceptibility to

infection given exposure(Civitello and Rohr 2014). The models varied based on whether these

processes were functions of snail body size (represented as the longest shell diameter, hereafter

"length"). In the null model, all functions of length are constants, therefore no variation in

prevalence by size-structure can be predicted. We also considered two models in which one

parameter varied with body size: (1) the size-dependent exposure-only, in which only exposure is

a function of host body size, and (2) the size-dependent susceptibility only, in which only

susceptibility is a function of body size. Finally, we considered a fully size-dependent model,

which varies both exposure (ε) and susceptibility (σ) with host body size. All models yield an

analytical solution for snail infection prevalence at time t_1 (Box 1). The three models that contain

some element of size-dependence all predict that different size classes within a single size-

structured population can experience different levels of infection prevalence and that populations

of differing size-structures can vary in their overall population infection prevalence. For all

models with a size-dependent component, we investigated two alternative parsimonious

representations: linear and exponential functions of host length (Appendix S1: Figure S1).

For each model, we solved for the expected prevalence in each size class present analytically,

given the size structure and parasite density of that treatment (Box 1). We then used the bbmle

package in R to conduct maximum likelihood estimation to estimate parameters for ε and σ for

each model using the binomial error distribution with the probability of "success", i.e., infection,

equal to the expected prevalence for that size class given the experimental conditions and

parameter values (Allen, Linda 2010, Strauss et al. 2019, Bolker and R Development Core Team

2021). This probability distribution provides the likelihood for our observations of infection

status for snail hosts in each size class, which we combined to generate an overall likelihood for

all observations. Lastly, to control for potential differences in infectivity among parasites across

the different temporal blocks, we incorporated a block effect for the baseline susceptibility

parameter, σ_0. Using corrected Akaike's information criteria (AICc) in R (R Core Team 2021),

we competed the fully size-dependent model against the size-dependent exposure only, the size-

dependent susceptibility only, and the size-independent null model.

Application to field data

We then estimated the influence of population size structure on schistosome transmission for a

snail population in an endemic setting experiencing natural variation in size structure throughout

the year. We used WebPlotDigitizer (Rohatgi 2020) to gather snail size data over time from a

population of *Biomphalaria tenagophilia* in del Plata Basin, Argentina from 2000-2001 (Rumi et

al. 2009). WebPlotDigitizer is a free, open source, web-based platform that allows for

measurement and quantification of images. We took the histogram plots of snail sizes across

time in Figure 1 of Rumi et al. 2009 and imported them into WebPlotDigitizer. Then, we labeled

the corresponding value of the X axis and the Y axis in the original figure and marked the data

points. After that, WebPlotDigitizer can automatically extract the specific values of the marked

data points. The code of WebPlotDigitizer is available at

https://github.com/ankitrohatgi/WebPlotDigitizer. Then, using these size structures, we used the

fully size-dependent transmission model to estimate the probability of successful infection by a

schistosome given that schistosome makes irreversible contact with a snail in the population for

each time point. We took this approach due to the lack of data on waterbody size in this dataset,

and therefore we were unable to estimate snail infection prevalence since snail density was

unknown. However, the availability of snail abundance and body size data allowed us to take the

point of view of a parasite encountering the waterbody and confronted with different snail size

structures at different points in time. Specifically, to estimate probability of a parasite succeeding

in infecting a snail, for each size class (1 mm increment) of snail we estimated the probability

that the schistosome miracidium would contact and subsequently infect a snail in that single size

class based on the relative abundance of snails in that size class and the size-dependent exposure

and susceptibility traits of all snails in the population at that time. We then summed infection

probabilities across all snail size classes for each time point to get an overall probability of

infection (Eq. 12):

$$P(Infection|Snail\ contact) = \sum_{i=1}^{n} 1 - e^{-\sigma_0 * \left(\frac{\varepsilon_0}{\varepsilon_0 * S_i}\right) * P_0 * (1 - e^{-t(\varepsilon_0 * S_i)})} \quad \text{(Eq. 12)}$$

We estimated 95% confidence intervals for this quantity by drawing estimated transmission

parameters from the joint 95% confidence interval set inferred from the experimental results

(Bolker 2008). We contrasted these estimates to the 95% confidence interval for infection

success given exposure for the size-independent (null) model, which makes a constant prediction

of parasite success, independent of size structure.

III. Results

Experimental Transmission Experiments

Population prevalence for snails varied by size structure and parasite density (Fig. 1). The
uniform small size structure had the greatest prevalence at each parasite density (36 parasites:
0.37 ± 0.026 [mean $\pm$ SE]; 144 parasites: 0.69 ± 0.078; 256 parasites: 0.83 ± 0.034) and this
difference was significantly greater than the uniform large (z ratio = -5.425, p<.0001), uniform
medium (z ratio = -3.868，p=0.0010) and equal size structures (z ratio = -4.203, p= 0.0003). The
uniform large size structure had the smallest prevalence of the tanks at the lowest and highest
parasite densities (36 parasites: 0.13 ± 0.011; 256 parasites: 0.32 ± 0.13), and prevalence for the
uniform large size structure was significantly less than the small skewed (z ratio = 3.480, p=
0.0046) and the uniform small (z ratio = -5.425, p <.0001).

Within size classes, the average prevalence for small snails varied by population size structure
and parasite density (Fig 2). Regardless of population size structure, large snails were less likely
to be infected than small snails (z ratio = -3.67, p=0.0022). Small snails in the uniform small size
structure had the highest prevalence across parasite densities (Fig 2.A), and were significantly
more likely to be infected than small snails in the equal size structure treatment (z ratio = -5.102,
p<.0001). Small snails in the small skewed size structure were also significantly more likely to
be infected than small snails in the equal size structure (z ratio = -3.552, p = 0.0011). Medium
and large snails did not show significant differences in infection between size structures (Fig
2.B,C).

Transmission model competition

The size-independent null model performed poorly (Table 1). It either over or underestimated
transmission depending on the population size-structure (Fig 1D). Exposure-only size

dependence and susceptibility-only size dependence performed better (Table 1), but still tended

to underestimate schistosome prevalence especially at lower parasite densities (Fig 1B, 1C). A

fully size-dependent model performed far better than other candidate models (Table 1), despite

similar issues in underestimation at the lowest parasite density. An exponential relationship

between body size and exposure and susceptibility outperformed a linear relationship (Appendix

S1:Figure S1, Appendix S1:Table S1), therefore we focus in the main text on comparing size-

based models with exponential function(s) to the null model. Maximum likelihood estimates for

size-dependent exposure and susceptibility from the fully size-dependent model allow for

predictions of exposure(ε), susceptibility(σ), and transmission ($\beta = \varepsilon \times \sigma$) based on body size

(Fig 3). Experimental data on size-based susceptibility correspond well to these predictions (Fig

3B) and the transmission parameter is more strongly influenced by susceptibility, resulting in a

prediction of overall lower transmission with increasing body size (Fig 3C).

Implications for *S. mansoni* dynamics

The size structure of a natural *Biomphalaria* population varied substantially during the year (Fig

4A). Using these size structure data and our winning fully size-dependent model, we estimated

that the probability of establishing a successful infection for a single schistosome miracidium

encountering a snail in this population would also vary over the course of the year due to the

influence of snail body size on transmission. We took this approach of generating predictions

from our model due to the lack of available data from field settings that includes both snail body

size and infection prevalence. When small snails are overrepresented in the population, e.g. day

150, the predicted success of a schistosome is much higher than when the population is skewed

towards larger individuals, e.g. day 208 (Fig 4B). Importantly, the null and size-dependent

models yielded similar estimates in only 3 of 14 months. Overall, the size-dependent model

predicted lower parasite success than the null model, suggesting that ignoring size structure

could lead to overestimation of transmission potential.

Discussion

Natural populations are heterogenous, and in order to create ecological theory that can reflect

observed dynamics or predict future outcomes we must find tenable ways to address this

heterogeneity (Bolnick et al. 2011). It is particularly important to include relevant heterogeneity

in transmission, because this process shapes the ecological and evolutionary trajectory of disease

dynamics (McCallum et al. 2001). Transmission models that do not incorporate heterogenous

host traits often fit data poorly (Dwyer et al. 2010), compromising their applicability to natural

populations. Therefore, identifying the traits that drive transmission and the directionality of their

effects can improve theoretical predictions for natural dynamics and better assess interventions.

However, we do not suggest that disease ecologists embark on an endless hunt to characterize all

heterogenous host traits. Instead, focusing on traits that either integrate numerous aspects of a

host's ecology, have the potential to regulate multiple downstream processes impacting

transmission, or both, can greatly reduce unexplained variability while still being tenable to

include mathematically and empirically. Body size is trait that satisfies both of these criteria for

many disease systems, as it varies amongst individuals due to many processes such as age and

nutrition (Kooijman, S.A.L.M. 2010), and it profoundly impacts many ecological processes such

as predation, resource competition, and reproductive success(Wellborn 1994, Hopcraft et al.

2010). This leads to its utility as a tractable trait to include in transmission models because often

an individual's history resulting in their current body size is usually impossible to reconstruct for

wild organisms. Further exploring the role of body size in other host-parasite systems may reveal

correlations and trade-offs between host physiology, behavior, and transmission.

For decades, it has been known that larger *Biomphalaria* snails experience a higher exposure to

schistosomes but a lower susceptibility to infection (Niemann and Lewis 1990). However, the

relevance of this phenomenon to epidemiological dynamics has not been determined. Here we

demonstrate the epidemiological relevance of parasite transmission in size-structured

populations. We showed that this individual-level phenomenon can have population-level

consequences for schistosome prevalence: differently size-structured populations experienced

very different schistosome transmission dynamics in our experimental trials. Populations with a

greater proportion of large sized snails had lower schistosome prevalences, while populations of

uniformly small snails experienced the greatest prevalence. From the perspective of an individual

snail, the body size of immediate neighbors is also important for transmission: small snails had

lower prevalence of schistosomes when in populations with a greater proportion of large

conspecifics. This observation supports the hypothesis that individuals with relatively high

exposure rates to parasites can shield their conspecifics from infection, a form of transmission

interference akin to the much-discussed dilution effect that is typically considered in multi-

species host communities (Keesing et al. 2006, Shaw and Civitello 2021). While our

experimental design only captured short-term transmission dynamics, explicitly incorporating

host body size into schistosome transmission models will allow for future extensions of this work

to explore longer term outcomes and include ecological interactions that can impact body size

such as resource competition (Malishev and Civitello 2020). Furthermore, this size-explicit

transmission model can now be incorporated into models that evaluate the entire schistosome

lifecycle, and better predict transmission risk to humans based both on the impact of snail size at

the time of transmission to the snail and also snail production of cercariae, the life stage which is

infectious to people, because cercarial production can be size and resource dependent (Civitello

et al. 2022).

Incorporating size structure into schistosome transmission models could also improve our

understanding of seasonal variation in schistosome transmission to intermediate snail hosts. We

found that for a well-characterized natural *B. glabrata* population, temporal variation in size

structure was large enough to cause substantial variation in predicted transmission success in

snails for schistosomes. It's important to note that this is not an empirical result: it is using our

mechanistic model to generate a hypothesis on the impact of size-structure on schistosome

dynamics in endemic settings. To our knowledge, no available public datasets contain the

necessary data on snail density, body size, and infection prevalence to directly compare to our

model predictions. Our hope is that the stark predicted difference in parasite infection success

between the null and the fully size-dependent model will motivate future work to collect the

necessary data to directly test this hypothesis. Numerous biotic and abiotic factors, such as

predation, resource competition, temperature, and nutrient availability, can impact host snail

growth and therefore population size structure across space and time. Furthermore, schistosome

intervention efforts can be costly and labor-intensive, but characterizing the seasonal patterns in

size structure may allow for more targeted approaches at times when the chance of transmission

is highest, such as following a large reproductive burst when the population has a large

proportion of neonates.

Hosts are not identical, despite the plethora of transmission models that treat them as such.

Scaling up the effects of heterogenous host traits on transmission to investigate population-level

effects can allow for better inference and prediction, especially when using traits such as body

size that are measurable and can reflect correlations between numerous ecological processes. Our

work shows the importance of this approach in the snail-schistosome system. Tackling the

influence of host heterogeneity in transmission may be key for resolving many unanswered

questions in disease ecology.

Acknowledgements: We thank members of the Civitello Lab for their comments on this

manuscript. KS was supported by the National Institute of Allergy and Infectious Diseases of the

National Institutes of Health under Award Number T32AI138952

and Award Number F31AI147611. DJC was supported by NIH 1R01 AI150774-01. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health.

Author Contributions: KES and DJC conceived and designed the study. KES, REC, RS

performed experiments. KES, REC, DJC analyzed data. KES and REC wrote the manuscript

with input from DJC.

Conflict of Interest Statement: The authors have no conflicts of interest to report.

References

Allen, Linda. 2010. An Introduction to Stochastic Processes with Applications to Biology. 2nd edition. Chapman and Hall/CRC.

Baer, K. N., and C. E. Goulden. 1998. Evaluation of a High-Hardness COMBO Medium and Frozen Algae forDaphnia magna. Ecotoxicology and Environmental Safety 39:201–206.

Bolker, B. M. 2008. Likelihood and All That. Pages 169–221 Ecological Models and Data in R. Princeton University Press.

Bolker, B., and R Development Core Team. 2021. bbmle: Tools for General Maximum Likelihood Estimation.

Bolnick, D. I., P. Amarasekare, M. S. Araújo, R. Bürger, J. M. Levine, M. Novak, V. H. W. Rudolf, S. J. Schreiber, M. C. Urban, and D. A. Vasseur. 2011. Why intraspecific trait variation matters in community ecology. Trends in Ecology & Evolution 26:183–192.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J. Skaug, and B. M. Maechler, Martin Bolker. 2017. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.

Civitello, D. J., T. Angelo, K. H. Nguyen, R. B. Hartman, N. C. Starkloff, M. P. Mahalila, J. Charles, A. Manrique, B. K. Delius, L. M. Bradley, R. M. Nisbet, S. Kinung'hi, and J. R. Rohr. 2022. Transmission potential of human schistosomes can be driven by resource competition among snail intermediate hosts. Proceedings of the National Academy of Sciences 119:e2116512119.

Civitello, D. J., and J. R. Rohr. 2014. Disentangling the effects of exposure and susceptibility on transmission of the zoonotic parasite Schistosoma mansoni. Journal of Animal Ecology 83:1379–1386.

Dinguirard, N., C. Heinemann, and T. P. Yoshino. 2018. Mass Isolation and In Vitro Cultivation

of Intramolluscan Stages of the Human Blood Fluke Schistosoma Mansoni.

JoVE:e56345.

Dwyer, G., J. S. Elkinton, and J. P. Buonaccorsi. 1997. Host heterogeneity in susceptibility and

disease dynamics. The American Naturalist 150:685–707.

Dwyer, G., J. S. Elkinton, and J. P. Buonaccorsi. 2010. The American Society of Naturalists

Host Heterogeneity in Susceptibility and Disease Dynamics : Tests of a Mathematical

Model Published by : The University of Chicago Press for The American Society of

Naturalists Stable URL : http://www.jstor.org/stable/2. East 150:685–707.

Gomes, M. G. M., M. U. Ferreira, R. M. Corder, J. G. King, C. Souto-Maior, C. Penha-

Gonçalves, G. Gonçalves, M. Chikina, W. Pegden, and R. Aguas. 2022. Individual

variation in susceptibility or exposure to SARS-CoV-2 lowers the herd immunity

threshold. Journal of Theoretical Biology 540:111063.

Gryseels, B., K. Polman, J. Clerinx, and L. Kestens. 2006. Human schistosomiasis. Lancet

368:1106–1118.

Hall, S. R., L. Sivars-Becker, C. Becker, M. A. Duffy, A. J. Tessier, and C. E. Cáceres. 2007.

Eating yourself sick: Transmission of disease as a function of foraging ecology. Ecology

Letters 10:207–218.

Han, B. A., A. W. Park, A. E. Jolles, and S. Altizer. 2015. Infectious disease transmission and

behavioural allometry in wild mammals. Journal of Animal Ecology 84:637–646.

Hechinger, R. F., K. L. Sheehan, and A. V. Turner. 2019. Metabolic theory of ecology

successfully predicts distinct scaling of ectoparasite load on hosts. Proceedings of the

Royal Society B: Biological Sciences 286:20191777.

Hopcraft, J. G. C., H. Olff, and A. R. E. Sinclair. 2010. Herbivores, resources and risks:

alternating regulation along primary environmental gradients in savannas. Trends in

Ecology & Evolution 25:119–128.

Jarne, P., J.-P. Pointier, and P. David. 2011. Biosystematics of Biomphalaria spp. with an

Emphasis on Biomphalaria glabrata. Pages 1–32 *in* R. Toledo and B. Fried, editors.

Biomphalaria Snails and Larval Trematodes. Springer New York, New York, NY.

Keesing, F., R. D. Holt, and R. S. Ostfeld. 2006. Effects of species diversity on disease risk.

Ecology Letters 9:485–498.

King, K. C., J. Jokela, and C. M. Lively. 2011. Trematode parasites infect or die in snail hosts.

Biology Letters 7:265–268.

Kooijman, S.A.L.M. 2010. Dynamic Energy Budget Theory For Metabolic Organisation.

Cambridge University Press.

Larson, M. K., R. C. Bender, and C. J. Bayne. 2014. Resistance of biomphalaria glabrata 13-16-

R1 snails to schistosoma mansoni PR1 is a function of haemocyte abundance and

constitutive levels of specific transcripts in haemocytes. International Journal for

Parasitology 44:343–353.

Lenth, R. V. 2022. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Lloyd-Smith, J. O., S. J. Schreiber, P. E. Kopp, and W. M. Getz. 2005. Superspreading and the

effect of individual variation on disease emergence. Nature 438:355.

Malishev, M., and D. J. Civitello. 2020. Modelling how resource competition among snail hosts

affects the mollusciciding frequency and intensity needed to control human schistosomes.

Functional Ecology 34:1678–1689.

Mayer, M., L. Schlippe Justicia, R. Shine, and G. P. Brown. 2021. Host defense or parasite cue:

Skin secretions mediate interactions between amphibians and their parasites. Ecology

Letters 24:1955–1965.

McCallum, H., N. Barlow, and J. Hone. 2001. How should pathogen transmission be modelled?

Trends in Ecology & Evolution 16:295–300.

Niemann, G. M., and F. A. Lewis. 1990. Schistosoma mansoni: Influence of Biomphalaria

glabrata size on susceptibility to infection and resultant cercarial production.

Experimental Parasitology 70:286–292.

Poulin, R. 2004. Macroecological patterns of species richness in parasite assemblages. Basic and

Applied Ecology 5:423–434.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Richards, C. S., M. Knight, and F. A. Lewis. 1992. Genetics of Biomphalaria glabrata and its

effect on the outcome of Schistosoma mansoni infection. Parasitology Today 8:171–174.

Rohatgi, A. 2020, July. WebPlotDigitizer.

Rumi, A., D. Gutiérrez Gregoric, and A. Roche. 2009. TENDENCIAS DEL CRECIMIENTO

INDIVIDUAL EN POBLACIONES NATURALES DE Biomphalaria spp.

(GASTROPODA, PLANORBIDAE) EN LA CUENCA DEL PLATA, ARGENTINA.

Comunicaciones de la Sociedad Malacológica del Uruguay.

Shaw, K. E., and D. J. Civitello. 2021. Re-emphasizing mechanism in the community ecology of

disease. Functional Ecology 35:2376–2386.

Strauss, A. T., J. L. Hite, D. J. Civitello, M. S. Shocket, C. E. Cáceres, and S. R. Hall. 2019.

Genotypic variation in parasite avoidance behaviour and other mechanistic, nonlinear

components of transmission. Proceedings. Biological sciences 286:20192164.

Sweeny, A. R., and G. F. Albery. 2022. Exposure and susceptibility: The Twin Pillars of

infection. Functional Ecology 36:1713–1726.

Théron, A., A. Rognon, and J.-R. Pagès. 1998. Host choice by larval parasites: a study of

Biomphalaria glabrata snails and Schistosoma mansoni miracidia related to host size.

Parasitology Research 84:727–732.

Van der Waal, K. L., and V. O. Ezenwa. 2016. Heterogeneity in pathogen transmission:

mechanisms and methodology. Functional Ecology 30:1606–1622.

Wellborn, G. A. 1994. Size-Biased Predation and Prey Life Histories: A Comparative Study of

Freshwater Amphipod Populations. Ecology 75:2104–2117.

White, C. R. 2010. There is no single p. Nature 464:691–693.

World Health Organization. 2022. Schistosomiasis Fact Sheet.

Yan, J., L. Gangoso, S. Ruiz, R. Soriguer, J. Figuerola, and J. Martínez-de la Puente. 2021.

Understanding host utilization by mosquitoes: determinants, challenges and future

directions. Biological Reviews 96:1367–1385.

Box 1. Model construction.

We constructed deterministic transmission models that tracked changes in the densities of

susceptible hosts, S_i and infected hosts, I_i, in an arbitrary number of host size classes, i, as well as

free-living parasites, P, with differential equations:

$$\frac{dS_i}{dt} = -\beta_i * S_i * P \qquad\qquad\qquad \text{Eq. 1}$$

$$\frac{dI_i}{dt} = \beta_i * S_i * P \qquad\qquad\qquad \text{Eq. 2}$$

$$\frac{dP}{dt} = -\sum_{i=1}^{n}(\varepsilon_i * (S_i + I_i)) * P \qquad\qquad \text{Eq. 3}$$

We parameterized these models based on a short-term infection experiment, therefore we

ignored processes that occur at longer time scales such as birth and deaths of hosts and parasites

(Civitello and Rohr 2014). For each model we broke down the transmission rate (β_i) into two

main components, exposure (ε_i) and susceptibility (σ_i), that we allowed to vary with size (L_i):

Size-independent (null): $\qquad \varepsilon_0 * \sigma_0 \qquad\qquad\qquad$ Eq. 4

Size-dependent exposure: $\qquad (\varepsilon_0 * e^{(\varepsilon_{sz}*L)}) * \sigma_0 \qquad\qquad$ Eq. 5

Size-dependent susceptibility: $\quad \varepsilon_0 * (\sigma_0 * e^{(\sigma_{sz}*L)}) \qquad\qquad$ Eq. 6

Fully size-dependent: $\qquad (\varepsilon_0 * e^{(\varepsilon_{sz}*L)}) * (\sigma_0 * e^{(\sigma_{sz}*L)}) \qquad$ Eq. 7

Prevalence can be directly calculated within each size class (i), based on the initial density of

susceptible individuals in each size class, $S(0)_i$, :

Size-independent (null): $\qquad 1 - e^{(-\sigma_0\left(\frac{\varepsilon_0}{\sum \varepsilon_0 S(0)_i}\right)P_0\left(1-e^{\left(-t(\sum \varepsilon_0 S(0)_i)\right)}\right)}$ Eq. 8

Size-dependent exposure: $\qquad 1 - e^{(-\sigma_0\left(\frac{\varepsilon_i}{\sum \varepsilon_i S(0)_i}\right)P_0\left(1-e^{\left(-t(\sum \varepsilon_i S(0)_i)\right)}\right)}$ Eq. 9

Size-dependent susceptibility: $\quad 1 - e^{(-\sigma_i\left(\frac{\varepsilon_0}{\sum \varepsilon_0 S(0)_i}\right)P_0\left(1-e^{\left(-t(\sum \varepsilon_0 S(0)_i)\right)}\right)}$ Eq. 10

Fully size-dependent: $\qquad 1 - e^{(-\sigma_i\left(\frac{\varepsilon_i}{\sum \varepsilon_i S(0)_i}\right)P_0\left(1-e^{\left(-t(\sum \varepsilon_i S(0)_i)\right)}\right)}$ Eq. 11

Table 1. Parameter estimates and model selection statistics
516

Model	Maximum Likelihood Parameter Estimates				AIC	ΔAIC	Akaike weight
	ε_0	ε_L	σ_0	σ_L			
Fully size-dependent	0.64	--	0.151	--	995.1	0	>.999
Size-dependent susceptibility	1.04	-0.085	0.162	--	1029.8	34.7	<0.001
Size-dependent exposure	0.669	--	0.305	-0.105	1083.6	88.5	<0.001
Size-independent (null)	0.25	0.184	0.609	-0.214	1116.8	121.6	<0.001

Figure Captions

Figure 1. Schistosome prevalence in snail populations with differing size structures. Points represent mean ± SE prevalence in each population structure and lines represent best model fits. Each panel depicts one of the candidate transmission models; experimental data is identical between panels. In our transmission trials, uniform small populations had a significantly greater prevalence than the uniform large uniform medium and equal size structures, and prevalence in the uniform large size structure was significantly less than the small skewed and the uniform small. A fully size-dependent transmission model (**A**) in which exposure and susceptibility vary with body size outperformed all other candidate models. Allowing only exposure to vary with body size (**B**) tended to underestimate prevalence in populations with a greater proportion of small individuals, while a model in which only susceptibility varies with body size (**C**) performed better than exposure-dependence, it did especially poorly at predicting prevalence of uniformly large populations. A null model with no size-dependence (**D**) is unable to make different predictions in differently size-structured populations, and thus has a very poor fit to the observed data.

Figure 2. Schistosome prevalence within size classes in snail populations with differing size structures. Points represent average prevalence in each population structure, n=4 replicates per treatment; lines represent fits from the winning candidate model with full size dependence. A) Small snails B) Medium snails C) Large snails. Size structures with a greater proportion of large conspecifics resulted in lower infection prevalence in the small snails within those populations. Medium and large snails did not show significant differences in infection between size structures.

Figure 3. Estimates for exposure, ε(L), (A), susceptibility,σ(L), (B), and overall transmission rate, β(L) = ε(L)*σ(L) (C) as functions of snail length. Lines represent the maximum likelihood of estimated parameters across body size of snails. Shading represents the minimum and maximum from the 95% confidence interval of parameters estimated. Points in (B) are from experimentally observed schistosome susceptibility across snail sizes from individual exposures, while the other two parameters are inferred from the full dataset.

Figure 4. Predicted infection success of a single schistosome parasite given exposure of a snail in a natural population with varying body size structure over time. A) Distribution of body sizes from a population of *Biomphalaria tenagophilia* in del Plata Basin, Argentina from 2000-2001(Rumi et al. 2009). Number of snails sampled at each timepoint represented as n=. B) Points are the mean ± 95% confidence interval of the predicted infection success of a single schistosome introduced into this snail population from simulated transmission trials using the fully size-dependent model. Grey line is the predicted successful infection from a transmission model that ignores size-based exposure and susceptibility. Substantial changes in snail population size structure cause month-to-month estimates of schistosome success to diverge substantially from predictions that ignore size dependence. In particular, the size-null prediction tends to overestimate schistosome success except at the times in which the population is heavily dominated by small snails.

580   **Figure 1**

602 **Figure 2**
603

625   **Figure 3**

A.

B.

Days since June 18, 2000