1 **Parasite Transmission in Size-Structured Populations**

2 **Kelsey E Shaw*[1], Rebecca E Cloud[2], Raeyan Syed[1], David J Civitello[1]**

3   1. Department of Biology, Emory University, Atlanta, GA
4   2. School of Integrative Biology, University of Illinois, Urbana-Champaign
5
6
7 *Corresponding Author: kshaw@emory.edu
8

Abstract:

Host heterogeneity can impact parasite transmission, but determining underlying traits and

incorporating them into transmission models remains challenging. Body size is easily measured

and influences numerous ecological interactions, including transmission. In the snail-

schistosome system, larger snails have a higher exposure to parasites but lower susceptibility to

infection per parasite. We quantified the impact of size-based heterogeneity on population-level

transmission by conducting transmission trials in differently size-structured snail populations and

competing size-dependent transmission models. Populations with greater proportions of large

snails had lower prevalence, and small snails were shielded from infection by co-occurring large

conspecifics. Using the winning size-explicit model, we then estimated that schistosome

transmission potential varies dramatically across time due to seasonal changes in snail population

size structure. Thus, incorporating traits such as body size, which are impacted by and directly

affect host ecology, into transmission models could yield insights for natural dynamics and

disease mitigation in many systems.

I. Introduction:

Parasite transmission drives the ecological and evolutionary dynamics of host-parasite interactions. Therefore, demystifying the host and parasite factors that influence transmission is critical for characterizing disease dynamics and successfully managing any host-parasite system (Lloyd-Smith *et al.* 2009). In particular, transmission theory that ignores host variation and instead treats hosts as uniform individuals with identical infection risk matches data poorly and cannot address critical themes in disease ecology, such as superspreading and parasite aggregation (Lloyd-Smith *et al.* 2005). In order to more accurately predict disease dynamics in natural populations, it is important to consider the sources of host variation and their consequences for transmission because hosts can differ in many ways (Dwyer *et al.* 1997; Hall *et al.* 2007; Strauss *et al.* 2019).

Schistosomes are the causative agent of schistosomiasis, a Neglected Tropical Disease that is second only to malaria in its worldwide impact on human health and disproportionately affects individuals in poor, rural environments (Gryseels *et al.* 2006). Over 200 million people worldwide are currently infected, and schistosomiasis accounts for approximately 200,000 deaths per year (World Health Organization 2022). Schistosomes have a complex life cycle and obligately cycle between mammalian and snail hosts. One source of host heterogeneity that impacts schistosome transmission is snail body size: larger snails experience higher exposure rates but are less likely to become successfully infected on a per parasite basis (Niemann & Lewis 1990; Théron *et al.* 1998). The importance of body size in transmission to the snail host becomes even more apparent when considering the vast range in host snail body size. For example, a neonate *Biomphalaria glabrata*, an intermediate host species for *Schistosoma*

mansoni, is born at ~0.75 mm and can grow up to 40 mm under natural conditions over its

lifetime (Jarne *et al.* 2011). This growth trajectory corresponds to a ~150,000-fold increase in

biomass, akin to the difference between a mouse and an elephant (White 2010).

Despite the recognition that body size influences these mechanistic components of schistosome

transmission decades ago in isolated, individual snails, the consequences of this relationship

when scaled to size-structured populations have remained uninvestigated. Schistosome exposure

in a host is irreversible, meaning a schistosome cannot leave a snail once it has entered, even if it

fails to successfully infect the host. Therefore, snails that are invaded by parasites indirectly

reduce the infection risk of others in the population (King *et al.* 2011). If size-dependent

transmission effects scale up to the population level, then we predict that the presence of larger

hosts will contribute to a within-species dilution effect by shielding their smaller conspecifics

from parasite exposure while also being less likely to become infected themselves.

We investigated this hypothesis by conducting laboratory transmission trials of *Schistosoma*

mansoni in *Biomphalaria glabrata* populations of varying size structures. We then used these

data to compete candidate models of size-dependent transmission, and additionally used existing

field data of *B. glabrata* population size structures to estimate how parasite success may change

under natural conditions over time solely due to the dynamics of population size structure. In

many host-parasite systems researchers are still in the nascent phase of breaking down the

complex interaction of host traits that result in heterogeneous parasite transmission (Sweeny &

Albery 2022). In the snail-schistosome system, body size is a measurable trait with the potential

to influence parasite transmission via multiple mechanisms, but it remains relatively unexplored

how this individual-level trait scales up to influence transmission within and between

populations. Incorporating this fundamental host trait into our understanding of schistosome

transmission could yield important insights into transmission risk and disease control of this

widespread and debilitating parasite, and this approach can be applied broadly to other disease

systems and critical traits.

II. Methods

Snail Maintenance. *B. glabrata* snails of the NMRI strain were maintained under favorable

conditions. Snails were kept in HHCOMBO artificial lake water (Baer & Goulden 1998) at 26°C

with a 12:12 light:dark cycle. Snails were fed a diet of fish flakes (Omega One) and chicken feed

(Nutrena Meatbird Crumbles) suspended in 1% agar *ad libitum.*

Experimental Design. We conducted the experiment using a fully factorial design: five

population size-structures, each with a total of 18 snails, were exposed to three densities of

parasites in 24-hour transmission trials in 15-liter tanks, creating 15 total treatment combinations.

B. glabrata were first assigned into three size classes by shell diameter: "small" (2-3 mm),

"medium" (6-8 mm), and "large" (12-15 mm). Different size-structures were determined by the

ratio of snails from each of the three size classes. The uniform size structure consisted of 18

snails from the same size class: "uniform small" (1:0:0), "uniform medium" (0:1:0), and

"uniform large" (0:0:1). "Equal" size-structured mesocosms had 6 snails from each size class

(1:1:1). "Small skewed" size structures contained 12 snails from the smallest size class and 3

snails from both the medium and large size classes (4:1:1). Parasite densities also varied between

the tanks: 36 (2 miracidia/snail), 144 (8 miracidia/snail), and 256 parasites per tank (14

miracidia/snail). We obtained *S. mansoni* eggs from experimentally infected mice livers and

purified and hatched eggs to obtain miracidia (Dinguirard *et al.* 2018). We conducted four

randomized blocks of these trials in which each block contained one replicate of each treatment.

To control for potential batch effects on parasite infectivity across the temporal blocks, a control

group consisting of all medium snails individually placed in 24 well plates was simultaneously

exposed to each of the parasite densities.

To quantify size-based susceptibility we also conducted transmission experiments in which 120

snails ranging from 0.5mm to 15mm were individually placed in well plates with 5 mL COMBO.

These snails were exposed to either 2 or 6 miracidia over 24 hours. Due to the small volume of

water used in the individual well plate exposures, we assume that snails were exposed to all

parasites present in these trials, an assumption justified by prior experiments (Niemann & Lewis

1990).

Infection Diagnosis. 24 hours post-exposure, all snails were collected, sorted by size and

treatment group, and maintained for 5 weeks (longer than the minimal prepatent period for *S.*

mansoni in favorable conditions). Snails were diagnosed visually by parasite shedding in

individual well plates at 4- and 5-weeks post-exposure to obtain prevalence data (Asch 1972).

Snails were diagnosed as "infected" if we observed *S. mansoni* cercariae in their well plate, or

"uninfected" if no cercariae were detected. Snails infected at week 4 were sacrificed, while

uninfected snails were returned to their tanks to repeat this process at week 5. Snails that died

while prepatent or between weeks 4 and 5 of the shedding process were not included in our

results (6.25% of all snails).

Model Creation and Maximum Likelihood Estimation of Parameters. We then built

deterministic size-dependent transmission models for parameterization and competition using the

experimental data (Box 1). Each model separates the transmission process into two underlying

processes: exposure to parasites and susceptibility to infection given exposure. The models

varied based on whether these processes were functions of snail body size (represented as the

longest shell diameter, hereafter "length"). In the null model, all functions of length are

constants, therefore no variation in prevalence by size-structure can be predicted. We also

considered two models in which one parameter varied with body size: (1) the size-dependent

exposure-only, in which only exposure is a function of host body size, and (2) the size-dependent

susceptibility only, in which only susceptibility is a function of body size. Finally, we considered

a fully size-dependent model, which varies both exposure (ε) and susceptibility (σ) with host

body size. The three models that contain some element of size-dependence all predict that

different size classes within a single size-structured population can experience different levels of

infection prevalence and that populations of differing size-structures can vary in their overall

population infection prevalence. For all models with a size-dependent component, we

investigated two alternative parsimonious representations: linear and exponential functions of

host length. For each model, we solved for the expected prevalence in each size class present

analytically, given the size structure and parasite density of that treatment. We then used the

bbmle package in R to conduct maximum likelihood estimation to estimate parameters for ε and

σ for each model using the binomial error distribution with the probability of "success", i.e.,

infection, equal to the expected prevalence for that size class given the experimental conditions

and parameter values (Strauss *et al.* 2019; Bolker & R Development Core Team 2021). Lastly, to

control for potential differences in infectivity among parasites across the different temporal

blocks, we incorporated a block effect for the baseline susceptibility parameter, σ_0. Using

corrected Akaike's information criteria (AICc) in R (R Core Team 2021), we competed the fully

size-dependent model against the size-dependent exposure only, the size-dependent susceptibility

only, and the size-independent null model.

Box 1. Model construction.
We constructed deterministic transmission models that tracked changes in the densities of
susceptible hosts, S_i and infected hosts, I_i, in an arbitrary number of host size classes, i, as well as
free-living parasites, P, with differential equations:

$$\frac{dS_i}{dt} = -\beta_i * S_i * P \qquad\qquad \text{Eq. 1}$$

$$\frac{dI_i}{dt} = \beta_i * S_i * P \qquad\qquad \text{Eq. 2}$$

$$\frac{dP}{dt} = -\sum_{i=1}^{n}(\varepsilon_i * (S_i + I_i)) * P \qquad\qquad \text{Eq. 3}$$

We parameterized these models based on a short-term infection experiment, therefore we
ignored processes that occur at longer time scales such as birth and deaths of hosts and parasites
(Civitello & Rohr 2014). For each model we broke down the transmission rate (β_i) into two main
components, exposure (ε_i) and susceptibility (σ_i), that we allowed to vary with size (L_i).
Prevalence can be directly calculated within each size class (i), based on the initial density of
susceptible individuals in each size class, $S(0)_i$, :

Model	Transmission rate (β)	Prevalence Equation
Size-independent (Null)	$\varepsilon_0 * \sigma_0$	$1 - e^{\left(-\sigma_0\left(\frac{\varepsilon_0}{\sum \varepsilon_0 S(0)_i}\right)P_0\left(1 - e^{\left(-t(\sum \varepsilon_0 S(0)_i)\right)}\right)\right)}$
Size-dependent exposure	$(\varepsilon_0 * e^{(\varepsilon_{sz}*L)}) * \sigma_0$	$1 - e^{\left(-\sigma_0\left(\frac{\varepsilon_i}{\sum \varepsilon_i S(0)_i}\right)P_0\left(1 - e^{\left(-t(\sum \varepsilon_i S(0)_i)\right)}\right)\right)}$
Size-dependent susceptibility	$\varepsilon_0 * (\sigma_0 * e^{(\sigma_{sz}*L)})$	$1 - e^{\left(-\sigma_i\left(\frac{\varepsilon_0}{\sum \varepsilon_0 S(0)_i}\right)P_0\left(1 - e^{\left(-t(\sum \varepsilon_0 S(0)_i)\right)}\right)\right)}$
Fully size-dependent	$(\varepsilon_0 * e^{(\varepsilon_{sz}*L)}) * (\sigma_0 * e^{(\sigma_{sz}*L)})$	$1 - e^{\left(-\sigma_i\left(\frac{\varepsilon_i}{\sum \varepsilon_i S(0)_i}\right)P_0\left(1 - e^{\left(-t(\sum \varepsilon_i S(0)_i)\right)}\right)\right)}$

Application to field data

We then estimated the influence of population size structure on schistosome transmission for a

snail population in an endemic setting experiencing natural variation in size structure throughout

the year. We used WebPlotDigitizer (Rohatgi 2020) to gather snail size data over time from a

population of *Biomphalaria tenagophilia* in del Plata Basin, Argentina from 2000-2001 (Rumi *et*

al. 2009). Then, using these size structures, we used the fully size-dependent transmission model

to estimate the probability of successful infection by a schistosome given that schistosome makes

irreversible contact with a snail in the population for each time point. Specifically, for each size

class (1 mm increment), we estimated the probability that the miracidium would contact and

subsequently infect a snail in that size class based on the relative abundance of snails in that size

class and the size-dependent exposure and susceptibility traits of all snails in the population at

that time. We then summed infection probabilities across all snail size classes for each time point

to get an overall probability of infection (Eq. 4):

$$P(Infection|Snail\ contact) = \sum_{i=1}^{n} 1 - e^{-\sigma_0 * \left(\frac{\varepsilon_0}{\varepsilon_0 * S_i}\right) * P_0 * (1 - e^{-t(\varepsilon_0 * S_i)})} \quad \text{(Eq. 4)}$$

We estimated 95% confidence intervals for this quantity by drawing estimated transmission

parameters from the joint 95% confidence interval set inferred from the experimental results

(Bolker 2008). We contrasted these estimates to the 95% confidence interval for infection

success given exposure for the size-independent (null) model, which makes a constant prediction

of parasite success, independent of size structure.

III. Results

Experimental Transmission Experiments

Population prevalence for snails varied by size structure and parasite density (Fig. 1). The

uniform small size structure had the greatest prevalence at each parasite density (36 parasites:

0.37 ± 0.026 [mean $\pm$ SE]; 144 parasites: 0.69 ± 0.078; 256 parasites: 0.83 ± 0.034). The uniform

large size structure had the smallest prevalence of the tanks at the lowest and highest parasite

densities (36 parasites: 0.13 ± 0.011; 256 parasites: 0.32 ± 0.13).

Figure 1. Schistosome prevalence in snail populations with differing size structures. Points represent mean ± SE

prevalence in each population structure and lines represent best model fits. Each panel depicts one of the candidate

transmission models; experimental data is identical between panels. A fully size-dependent transmission model (**A**)

in which exposure and susceptibility vary with body size outperformed all other candidate models. Allowing only

exposure to vary with body size (**B**) tended to underestimate prevalence in populations with a greater proportion of

small individuals, while a model in which only susceptibility varies with body size (**C**) performed better than

exposure-dependence, it did especially poorly at predicting prevalence of uniformly large populations. A null model

with no size-dependence (**D**) is unable to make different predictions in differently size-structured populations, and

thus has a very poor fit to the observed data.

201 Within size classes, the average prevalence for small snails varied by population size structure and

202 parasite density (Fig 2). Small snails in the uniform small size structure had the highest prevalence

203 across parasite densities (Fig 2.A, while medium and large snails did not show large differences

204 between size structures (Fig 2.B,C).

Figure 2. Schistosome prevalence within size classes in snail populations with differing size structures. Points

represent average prevalence in each population structure, n=4 replicates per treatment; lines represent fits from the

winning candidate model with full size dependence. A) Small snails B) Medium snails C) Large snails.

Transmission model competition

The size-independent null model performed poorly (Table 1). It either over or underestimated

transmission depending on the population size-structure (Fig 1D). Exposure-only size dependence

and susceptibility-only size dependence performed better (Table 1), but still tended to

underestimate schistosome prevalence especially at lower parasite densities (Fig 1B, 1C). A fully

size-dependent model performed far better than other candidate models (Table 1). An exponential

relationship between body size and exposure and susceptibility outperformed a linear relationship

217 (Table S1, Fig S1), therefore we focus in the main text on comparing size-based models with

218 exponential function(s) to the null model.

219 **Table 1.** Parameter estimates and model selection statistics
220

Model	Maximum Likelihood Parameter Estimates				AIC	ΔAIC	Akaike weight
	ε_0	ε_L	σ_0	σ_L			
Fully size-dependent	0.64	--	0.151	--	995.1	0	>.999
Size-dependent susceptibility	1.04	-0.085	0.162	--	1029.8	34.7	<0.001
Size-dependent exposure	0.669	--	0.305	-0.105	1083.6	88.5	<0.001
Size-independent (null)	0.25	0.184	0.609	-0.214	1116.8	121.6	<0.001

222 Maximum likelihood estimates for size-dependent exposure and susceptibility from the fully size-

223 dependent model allow for predictions of exposure(ε), susceptibility(σ), and transmission ($\beta = \varepsilon \times$

224 σ) based on body size (Fig 3). Experimental data on size-based susceptibility correspond well to

225 these predictions (Fig 3B) and the transmission parameter is more strongly influenced by

226 susceptibility, resulting in a prediction of overall lower transmission with increasing body size (Fig

227 3C).

Figure 3. Estimates for exposure, ε(L), (A), susceptibility,σ(L), (B), and overall transmission rate, β(L) =

ε(L)*σ(L) (C) as functions of snail length. Lines represent the maximum likelihood of estimated parameters across

body size of snails. Shading represents the minimum and maximum from the 95% confidence interval of parameters

estimated. Points in (B) are from experimentally observed schistosome susceptibility across snail sizes from

individual exposures, while the other two parameters are inferred from the full dataset.

Implications for *S. mansoni* dynamics

The size structure of a natural *Biomphalaria* population varied substantially during the year (Fig

4A). Using these size structure data, we estimated that the probability of establishing a successful

infection for a single schistosome miracidium encountering a snail in this population would also

vary over the course of the year due to the influence of snail body size on transmission. When

small snails are overrepresented in the population, e.g. day 150, the predicted success of a

schistosome is much higher than when the population is skewed towards larger individuals, e.g.

day 208 (Fig 4B). Importantly, the null and size-dependent models yielded similar estimates in

only 3 of 14 months. Overall, the size-dependent model predicted lower parasite success than the

null model, suggesting that ignoring size structure could lead to overestimation of transmission

potential.

Figure 4. Predicted infection success of a single schistosome parasite given exposure of a snail in a natural

population with varying body size structure over time. A) Distribution of body sizes from a population of

Biomphalaria tenagophilia in del Plata Basin, Argentina from 2000-2001(Rumi *et al.* 2009). Points are the mean ±

95% confidence interval of predicted schistosome success in this snail population from simulated transmission trials

using the fully size-dependent model. Grey line is the predicted success from a transmission model that ignores size-

based exposure and susceptibility. Substantial changes in snail population size structure cause month-to-month

estimates of schistosome success to diverge substantially from predictions that ignore size dependence. In particular,

the size-null prediction tends to overestimate schistosome success except at the times in which the population is

heavily dominated by small snails.

Discussion

Natural populations are heterogenous, and in order to create ecological theory that can reflect

observed dynamics or predict future outcomes we must find tenable ways to address this

heterogeneity (Bolnick *et al.* 2011). It is particularly important to include relevant heterogeneity

in transmission, because this process shapes the ecological and evolutionary trajectory of disease

dynamics (McCallum *et al.* 2001). Transmission models that do not incorporate heterogenous

host traits often fit data poorly (Dwyer *et al.* 2010), compromising their applicability to natural

populations. Therefore, identifying the traits that drive transmission and the directionality of their

effects can improve theoretical predictions for natural dynamics and better assess interventions.

However, we do not suggest that disease ecologists embark on an endless hunt to characterize all

heterogenous host traits. Instead, focusing on traits that either integrate numerous aspects of a

host's ecology, have the potential to regulate multiple downstream processes impacting

transmission, or both, can greatly reduce unexplained variability while still being tenable to

include mathematically and empirically. Body size is trait that satisfies both of these criteria for

many disease systems, as it profoundly impacts many ecological processes such as predation,

resource competition, and reproductive success (Wellborn 1994; Hopcraft *et al.* 2010). Body size

has the potential to vary drastically among individuals in a population and can reflect the

integration of many processes such as age, nutrition, and environmental stressors (Kooijman,

S.A.L.M. 2010).This leads to its utility as a tractable trait to include in transmission models

because often an individual's history resulting in their current body size is usually impossible to

reconstruct for wild organisms. Body size may interact with transmission via multiple

mechanisms: for example, larger individuals may provide a greater target area for vectors (Poulin

2004), may be more apparent to parasites that actively seek out their host using chemical cues

(Théron *et al.* 1998), or may encounter a greater abundance and diversity of microparasites if

they are exploring larger territories (Han *et al.* 2015). Further exploring the role of body size in

other host-parasite systems may reveal correlations and trade-offs between host physiology,

behavior, and transmission.

For decades, it has been known that larger *Biomphalaria* snails experience a higher exposure to

schistosomes but a lower susceptibility to infection (Niemann & Lewis 1990). However, the

relevance of this phenomenon to epidemiological dynamics has not been determined. Here we

demonstrate the epidemiological relevance of parasite transmission in size-structured

populations. We showed that this individual-level phenomenon can have population-level

consequences for schistosome prevalence: differently size-structured populations experienced

very different schistosome transmission dynamics in our experimental trials. Populations with a

greater proportion of large sized snails had lower schistosome prevalence, while populations of

uniformly small snails experienced the greatest prevalence. From the perspective of the snail, the

body size of immediate neighbors is also critically important for transmission: small snails had

lower prevalence of schistosomes when in populations with a greater proportion of large

conspecifics. This observation supports the hypothesis that individuals with relatively high

exposure rates to parasites can shield their conspecifics from infection, a form of transmission

interference akin to the much-discussed dilution effect that is typically considered in multi-

species host communities (Keesing *et al.* 2006; Shaw & Civitello 2021).

Incorporating size structure into schistosome transmission models could also improve our

understanding of seasonal variation in schistosome transmission. We found that for a well-

characterized natural *B. glabrata* population, temporal variation in size structure was large

enough to cause substantial variation in predicted transmission success for schistosomes.

Numerous biotic and abiotic factors, such as predation, resource competition, temperature, and

nutrient availability, can impact host snail growth and therefore population size structure across

space and time. Furthermore, schistosome intervention efforts can be costly and labor-intensive,

but characterizing the seasonal patterns in size structure may allow for more targeted approaches

at times when the chance of transmission is highest, such as following a large reproductive burst

when the population has a large proportion of neonates.

Hosts are not identical, despite the plethora of transmission models that treat them as such.

Scaling up the effects of heterogenous host traits on transmission to investigate population-level

effects can allow for better inference and prediction, especially when using traits such as body

size that are measurable and can reflect correlations between numerous ecological processes. Our

work shows the importance of this approach in the snail-schistosome system. Tackling the

influence of host heterogeneity in transmission may be key for resolving many unanswered

questions in disease ecology.

Acknowledgements: We thank members of the Civitello Lab for their comments on this

manuscript. KS was supported by the National Institute of Allergy and Infectious Diseases of the

National Institutes of Health under Award Number T32AI138952

and Award Number F31AI147611. DJC was supported by NIH 1R01 AI150774-01. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health.

References

Baer, K.N. & Goulden, C.E. (1998). Evaluation of a High-Hardness COMBO Medium and
Frozen Algae forDaphnia magna. *Ecotoxicology and Environmental Safety*, 39, 201–206.
Bolker, B. & R Development Core Team. (2021). bbmle: Tools for General Maximum
Likelihood Estimation.
Bolker, B.M. (2008). Likelihood and All That. In: *Ecological Models and Data in R*. Princeton
University Press, pp. 169–221.
Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., *et al.* (2011).
Why intraspecific trait variation matters in community ecology. *Trends in Ecology &*
Evolution, 26, 183–192.
Dinguirard, N., Heinemann, C. & Yoshino, T.P. (2018). Mass Isolation and In Vitro Cultivation
of Intramolluscan Stages of the Human Blood Fluke Schistosoma Mansoni. *JoVE*,
e56345.
Dwyer, G., Elkinton, J.S. & Buonaccorsi, J.P. (1997). Host heterogeneity in susceptibility and
disease dynamics. *The American Naturalist*, 150, 685–707.
Dwyer, G., Elkinton, J.S. & Buonaccorsi, J.P. (2010). The American Society of Naturalists Host
Heterogeneity in Susceptibility and Disease Dynamics : Tests of a Mathematical Model
Published by : The University of Chicago Press for The American Society of Naturalists
Stable URL : http://www.jstor.org/stable/2. *East*, 150, 685–707.
Gryseels, B., Polman, K., Clerinx, J. & Kestens, L. (2006). Human schistosomiasis. *Lancet*, 368,
1106–1118.
Hall, S.R., Sivars-Becker, L., Becker, C., Duffy, M.A., Tessier, A.J. & Cáceres, C.E. (2007).
Eating yourself sick: Transmission of disease as a function of foraging ecology. *Ecology*
Letters, 10, 207–218.
Han, B.A., Park, A.W., Jolles, A.E. & Altizer, S. (2015). Infectious disease transmission and
behavioural allometry in wild mammals. *Journal of Animal Ecology*, 84, 637–646.
Hopcraft, J.G.C., Olff, H. & Sinclair, A.R.E. (2010). Herbivores, resources and risks: alternating
regulation along primary environmental gradients in savannas. *Trends in Ecology &*
Evolution, 25, 119–128.
Jarne, P., Pointier, J.-P. & David, P. (2011). Biosystematics of Biomphalaria spp. with an
Emphasis on Biomphalaria glabrata. In: *Biomphalaria Snails and Larval Trematodes*
(eds. Toledo, R. & Fried, B.). Springer New York, New York, NY, pp. 1–32.
Keesing, F., Holt, R.D. & Ostfeld, R.S. (2006). Effects of species diversity on disease risk.
Ecology Letters, 9, 485–498.
King, K.C., Jokela, J. & Lively, C.M. (2011). Trematode parasites infect or die in snail hosts.
Biology Letters, 7, 265–268.

Kooijman, S.A.L.M. (2010). *Dynamic Energy Budget Theory For Metabolic Organisation*.
Cambridge University Press.
Lloyd-Smith, J.O., George, D., Pepin, K.M., Pitzer, V.E., Pulliam, J.R.C., Dobson, A.P., *et al.*
(2009). Epidemic Dynamics at the Human-Animal Interface. *Science*, 326, 1362–1367.
Lloyd-Smith, J.O., Schreiber, S.J., Kopp, P.E. & Getz, W.M. (2005). Superspreading and the
effect of individual variation on disease emergence. *Nature*, 438, 355.
McCallum, H., Barlow, N. & Hone, J. (2001). How should pathogen transmission be modelled?
Trends in Ecology & Evolution, 16, 295–300.
Niemann, G.M. & Lewis, F.A. (1990). Schistosoma mansoni: Influence of Biomphalaria glabrata
size on susceptibility to infection and resultant cercarial production. *Experimental*
Parasitology, 70, 286–292.
Poulin, R. (2004). Macroecological patterns of species richness in parasite assemblages. *Basic*
and Applied Ecology, 5, 423–434.
R Core Team. (2021). R: A language and environment for statistical computing.
Rohatgi, A. (2020). WebPlotDigitizer.
Rumi, A., Gutiérrez Gregoric, D. & Roche, A. (2009). TENDENCIAS DEL CRECIMIENTO
INDIVIDUAL EN POBLACIONES NATURALES DE Biomphalaria spp.
(GASTROPODA, PLANORBIDAE) EN LA CUENCA DEL PLATA, ARGENTINA.
Comunicaciones de la Sociedad Malacológica del Uruguay.
Shaw, K.E. & Civitello, D.J. (2021). Re-emphasizing mechanism in the community ecology of
disease. *Functional Ecology*, 35, 2376–2386.
Strauss, A.T., Hite, J.L., Civitello, D.J., Shocket, M.S., Cáceres, C.E. & Hall, S.R. (2019).
Genotypic variation in parasite avoidance behaviour and other mechanistic, nonlinear
components of transmission. *Proceedings. Biological sciences*, 286, 20192164.
Sweeny, A.R. & Albery, G.F. (2022). Exposure and susceptibility: The Twin Pillars of infection.
Functional Ecology, 36, 1713–1726.
Théron, A., Rognon, A. & Pagès, J.-R. (1998). Host choice by larval parasites: a study of
Biomphalaria glabrata snails and Schistosoma mansoni miracidia related to host size.
Parasitology Research, 84, 727–732.
Wellborn, G.A. (1994). Size-Biased Predation and Prey Life Histories: A Comparative Study of
Freshwater Amphipod Populations. *Ecology*, 75, 2104–2117.
White, C.R. (2010). There is no single p. *Nature*, 464, 691–693.
World Health Organization. (2022). Schistosomiasis Fact Sheet.
396