1

Female ticks (*Ixodes scapularis*) infected with *Borrelia burgdorferi* have increased

overwintering survival, with implications for tick population growth.

Short title: Infected ticks have increased overwintering survival

Amal El Nabbout[1], Laura V. Ferguson[2], Atsushi Miyashita[3], Shelley A. Adamo*[4]

1. Dept. Biology, Dalhousie University, Halifax, NS Canada

2. Dept. Biology, Acadia University, Wolfville, NS Canada

3. Institute of Medical Mycology, Teikyo University, Tokyo, Japan

4. Dept. Psychology and Neuroscience, Dalhousie University, NS Canada

*Corresponding author:

Shelley Adamo

Orcid number: 0000-0002-5973-571X

Dept. Psychology and Neuroscience, Dalhousie University, Halifax, NS Canada.   B3H 4R2

902 494 8853

sadamo@dal.ca

The tick, *Ixodes scapularis*, vectors pathogens such as *Borrelia burgdorferi*, the bacterium that

causes Lyme Disease. Over the last few decades *I. scapularis* has expanded its range,

introducing a novel health threat into these areas. Warming temperatures appear to be one cause

of its range expansion to the north. However, other factors are also involved. We show that

unfed adult female ticks infected with *B. burgdorferi* have greater overwintering survival than

uninfected female ticks. Locally collected adult female ticks were placed in individual

microcosms and allowed to overwinter in both forest and dune grass environments. In the spring

we collected the ticks and tested both dead and living ticks for *B. burgdorferi* DNA. Infected

ticks had greater overwintering survival compared with uninfected ticks every winter for three

consecutive winters in both forest and dune grass environments. We discuss the most plausible

explanations for this result. The increased winter survival of adult female ticks could enhance

tick population growth. Our results suggest that, in addition to climate change, *B. burgdorferi*

infection itself may be promoting the northern range expansion of *I. scapularis*. Our study

highlights how pathogens could work synergistically with climate change to promote host range

expansion.

37
38 Keywords: cold tolerance, ecological immunology, Lyme Disease, thermobiology, winter

Introduction

Winter is a stressful time for arthropods. Low temperatures can cause tissue injury and death, limiting the northern ranges of many insects and their relatives, such as ticks (Williams et al. 2015). The medically-important tick, *Ixodes scapularis*, dies at temperatures below -10° to -12°C (Burks et al. 1996; Vandyk et al. 1996), and even relatively mild winter temperatures (e.g. -3°C) may be lethal when humidity is high (i.e. due to inoculative freezing (Burks et al. 1996)). This cold intolerance may have historically excluded *I. scapularis* from northern regions of North America (Estrada-Peña 2002). Nevertheless, factors other than temperature play a role in determining overwintering survival (Lindsay et al. 1995; Lindsay et al. 1998; Brunner et al. 2014; Burtis et al. 2016; Burtis & Pflueger 2017; Clow et al. 2017; Kilpatrick et al. 2017; Hammond-Collins et al. 2022; Volk et al. 2022). For example, *I. scapularis* can successfully overwinter in regions that are below their thermal limits based on winter air temperatures (e.g. Northwestern Ontario (Lindsay et al. 1995), Upstate New York (Burtis et al. 2019), Maine (Volk et al. 2022)). Ticks can survive in these areas because they have adaptations, such as burrowing into leaf litter, that protect them from lethal temperatures (Brunner et al. 2012; Linske et al. 2019). However, overwintering survival is highly variable across years (Lindsay et al. 1998) (Volk et al. 2022), and our understanding of the various factors that determine overwintering success remains incomplete. We examine the possibility that infection with the bacterium *Borrelia burgdorferi* (the causative agent of Lyme Disease (Burgdorfer et al. 1982)) is also a factor in determining overwintering survival. We focus on the impact of infection on the overwintering survival of adult female ticks, because the overwintering survival of adult females has an impact on the size of the upcoming year's tick population. Despite their importance, few studies (Lindsay et al. 1995; Lindsay et al. 1998) examine overwintering in adult females. We

test whether infected adult female ticks have greater overwintering survival than non-infected

females, and what this effect may mean for its northward expansion.

After hatching from eggs, *I. scapularis* larvae climb on vegetation and position

themselves to find a host (i.e. quest) (for life cycle review see Eisen et al. 2012; Kilpatrick et al.

2017; Gray & Kahl 2022). Larvae attach themselves to a passing small mammal (e.g. mice).

Mice such as the white footed mouse (*Peromyscus leucopus*) can tolerate high numbers of *B.*

burgdorferi spirochetes in their blood when infected, allowing them to transmit *B. burgdorferi* to

I. scapularis ticks that feed on them (Eisen et al. 2012). After a single feeding, the larvae molt

into nymphs. *B. burgdorferi* persists in *I. scapularis* even after a molt (Helble et al. 2021; Pal et

al. 2021). Questing nymphs feed on a range of mammals. Nymphs that were infected as larvae

are able to transmit *B. burgdorferi* to its host (Gray & Kahl 2022). Uninfected nymphs can

acquire *B. burgdorferi* if they feed on an infected mammal that is capable of transmitting it.

After the second feeding, nymphs molt into adults. Adult females typically feed on larger

mammals, such as a deer, on which they typically find a mate (Gray & Kahl 2022). After

feeding, the female tick produces her eggs and then dies. Infected adult females are unlikely to

find a mammalian host capable of transmitting *B. burgdorferi* to another tick (Halsey et al.

2018), preventing the bacteria from continuing their life cycle. Large mammals such as deer do

not transmit *B. burgdorferi* to other ticks (Gray & Kahl 2022). However, although the *B.*

burgdorferi bacteria found in adult females will have little or no reproductive success, the

bacteria still interact physiologically with their host (Kurokawa et al. 2022).

Adults, nymphs and larvae can overwinter (Brunner et al. 2012). However, they have

limited capacity for movement below 4°C (Duffy & Campbell 1994), making them unable to

find a host or escape predators. Although the lack of questing (i.e. searching for a host) in winter

means that they are without food for the season, it does allow them to remain within their

insulated spaces and avoid the relatively lower ambient air temperatures (Burtis et al. 2019).

Leaf litter provides insulation, resulting in warmer temperatures and higher relative humidity

relative to the ambient air (Linske et al. 2019). However, overwintering survival can be low,

even when ticks are provided with leaf litter (Lindsay et al. 1998). Overwintering survival of

unfed adult females can vary from 27% to 78% depending on the year (Lindsay et al. 1998). If

infection enhances overwintering survival, some of this variability may be related to the

prevalence of infection in a population, that also varies from year to year (e.g. Scott et al. 2016).

Typically pathogens produce damage, reducing a host's condition (Boucias & Penland,

1994). Such pathology would be expected to reduce overwintering survival. However, *B.*

burgdorferi appears to have little or no negative impact on its host, although it does activate tick

immune responses (Kurokawa et al. 2020). When microbes have little negative impact, they can

enhance cold tolerance in arthropods (Ferguson et al. 2018). In the related tick *Ixodes ricinus,*

nymphs infected with *B. burgdorferi* were more likely to survive stressful environmental

conditions (e.g. low relative humidity (Herrmann & Gern 2010) and cold temperatures

(Herrmann & Gern 2013) than uninfected ticks. How pathogens alter host cold tolerance in

arthropods remains poorly understood (Overgaard & MacMillan 2017), but pathogens could

improve the winter survival of their hosts in a few ways. Pathogens could actively manipulate

hosts (e.g. ticks) to produce molecules that enhance host overwintering survival. This

mechanism has been suggested as an explanation for the effect of *Borrelia* on *I. ricinis*

(Herrmann & Gern, 2015). In *I. scapularis*, infection with the pathogen *Anaplasma*

phagocytophilum, leads to the production of the protein IAFGP (*Ixodes* anti-freeze glycoprotein)

(Neelakanta et al. 2010) (Heisig et al. 2014). This protein has both anti-microbial and

cryoprotectant properties (Heisig et al. 2014). It enhances cold tolerance in the short-term under

laboratory conditions (Neelakanta et al. 2010). Similarly, other molecules, such as heat shock

proteins, are produced by organisms in response to stressors such as pathogens (Adamo 2022),

and these molecules can protect organisms from a range of stressors (King & MacRae 2015),

including low temperatures (Overgaard & MacMillan 2017), a phenomenon known as cross-

tolerance (Sinclair et al. 2013). In this case, the pathogen is not manipulating the host, the host is

activating an integrated defense response that includes broad protection from a number of

stressors (Adamo 2022). *I. scapularis* infected with *B. burgdorferi* show increased gene

expression for heat shock proteins (Hart et al. 2022), and other molecules (Narashimhan et al.

2017) known to increase cold tolerance in arthropods (Overgaard & MacMillan 2017).

However, despite the existence of plausible mechanisms by which *B. burgdorferi* might enhance

I. scapularis overwintering survival, it has not been demonstrated that infection with *B.*

burgdorferi (or any other pathogen) increases overwintering success in *I. scapularis* under

natural winter conditions. We examine whether there is an interaction between infection and

winter survival in adult females.

If infected adult female ticks are more cold-tolerant, then this could increase their ability

to survive the winter. More adult females in the population could translate into a rise in the

number of larval ticks in the next generation. Enhanced growth of tick populations could also

contribute to the northward expansion of *I. scapularis*.

Materials and Methods

Field Study

Field work was conducted at the Harrison Lewis Coastal Discovery Centre (HLC), Port

L'Hebert, NS (HLC43°49'04.2" N 64°53'40.4" W), on the south shore of Nova Scotia. This area

is experiencing increasing tick populations (Hatchette et al. 2015). The work was conducted

over three winters (2018-2019, 2019-2020, and 2020-2021). Studies were approved by the

University Committee on Laboratory Animals at Dalhousie University (protocol number: I18-

06).

Ixodes scapularis were collected by dragging (Clow et al. 2017) from Thomas Raddall

Provincial Park, located adjacent to the HLC. Following collection, ticks were placed in

individual microcosms using a design modified from Burtis (Burtis 2017) as described in Adamo

et al. (2022) and repeated here. Adult female ticks were placed in individual microcosms topped

with maple/oak leaf litter (approximately 2 g). The microcosms were placed in the ground to the

depth of the leaf litter (about 5 cm). Microcosms were placed in either: a) a coastal forest

containing a mix of oak (*Quercus rubra*), maple (*Acer rubrum*) and spruce (*Picea rubens* and

Picea glauca) or, b) within the dune grass next to a sand beach. Both areas harbour *I. scapularis*

ticks in the spring, summer and fall. Microcosms were collected in the spring, and living and

dead ticks were tested for the presence of *B. burgdorferi* DNA using qPCR.

The microcosm was an open plastic tube (15.24 cm in length x 1.27 cm in width), with

mesh glued to each end, secured by a plastic cap. The plastic caps had a central hole that allowed

rain and snow to enter in the top and connected the soil with the open bottom of the tube. The

tubes contained sand and vermiculite (75% w/w sand and 25% w/w vermiculite) to a depth of 5

cm in the tube. The tubes were placed 5 cm into the ground. On top of the sand/vermiculite was

1 cm (approx. 2 g) of cut oak (*Quercus rubra* L.) and maple (*Acer rubrum* L.) leaves. A single

tick was placed into each microcosm and then the microcosm was placed under a coarse netting

(NuVue® (78.7cm L x 78.7 cm W x 83.8 cm H). A data logger (HOBO MX2301A) was placed

in each netting enclosure with a probe 5 cm underground between the microcosms; the logger

was covered with a solar radiation shield (Onset RS3-B) and the logger automatically recorded

temperature and relative humidity every hour. In 2018 (N=36), tick vials and data loggers were

placed in in either dune grass or in a mixed maple/oak/spruce stand of trees (Fig. 1). In years 2

and 3 (Fig. 1), we placed six microcosms under a single netting rather than placing all

microcosms for one study site under the same netting enclosure. Additionally, an extra study

location was created for each site, both of which were also located at the HLC; as a result, there

were two forest sites (forest 1 and forest 2) and two dune grass sites (dune grass 1 and dune grass

2) in years 2 and 3. In year 2, there were 145 ticks placed into individual microcosms. In year 3,

there were 120 ticks placed into individual microcosms with 6 microcosms per netting enclosure.

In Year 1 only, ticks were added to the microcosms as they were collected (from October

25, 2018 to Nov. 15, 2018). In years 2 and 3, ticks were held at 4°C and then all ticks were

placed in microcosms in the field on the same day (Year 2, Nov 7, 2019; Year 3 Nov. 28, 2020).

Ticks were retrieved on February 8, 2019 in Year 1. In year 2, ticks were retrieved on March 21,

2020, while in year 3 they were retrieved on March 23, 2021.

At the end of the field season, ticks were recovered by carefully sieving the tubes. Ticks

that were immobile after collection were exposed to both a gentle exhalation by the experimenter

(AEN) and probing with a paint brush. Ticks that showed no movement were tested again after

24 h at room temperature (approximately 22°C). If they still showed no movement they were

deemed to be dead. Both dead and living ticks were tested for the presence of *B. burgdorferi*

DNA using qPCR, following the procedure of Courtney et al. (2004), as described in Adamo et

al. (2022) (see Supplementary Information). Each DNA sample was tested to ensure that the

DNA was intact and not degraded. DNA integrity was assessed by gel electrophoresis (Aranda

et al. 2012) using a Gel Doc™ XR+ Digital Gel Imaging System® from Bio-Rad®. All of the

DNA we extracted from dead ticks was intact showing distinct bands on the gel.

Because of the importance of demonstrating that we could accurately assess *Borrelia*

DNA collected from dead ticks, we performed a lab study in which we simulated extreme winter

conditions (i.e. daily freeze/thaw). We assessed the presence of *Borrelia* in one half of a

sectioned adult female tick (n=30) immediately after cutting the tick in two, and the other half

was assessed 10 weeks later. The second half of the tick was incubated in a fluctuating thermal

regime incubator (-5°C to 10°C) in 12-hour increments for 10 weeks. We determined that we

could accurately measure infection in ticks that had been dead for 10 weeks under these

conditions (Supplementary Information, Table S2). There was 100% concordance between

results at time 0 and 10 weeks later (Table S2).

Statistical Analysis

Because of methodological differences across years, we analyzed the data using three 3-way

Log-Linear analyses. Data from all 3 years fit the assumptions for a log-linear model.   Post hoc

tests were controlled for a false discovery rate using the Benjamini-Hochberg procedure

(Benjamini & Hochberg 1995); www.biostathandbook.com/benjaminihochberg.xls). Tests were

performed using SPSS (ver. 27) and Prism (GraphPad, ver. 9.4.1).

Results

Infected ticks were more likely to survive the winter than uninfected ticks.

Ticks infected with *B. burgdorferi* had higher overwintering success than uninfected ticks

in each of the 3 years (Three-way Loglinear analysis for each year) (Fig. 1). In year 1 (N=36

ticks) the final model demonstrated significant interactions between infection and survival (X^2

(1)=4.73, p=0.03) as well as site and survival (X^2 (1)=6.1, p=0.01). In year 2 (N=145), the final

model demonstrated significant interactions between infection and survival (X^2 (1)= 19.6,

p<0.001) but not between site and survival (X^2 (1)=0.41, p=0.5). In year 3 (N=120) the final

model demonstrated a significant interaction between infection and survival (X^2 (1)= 5.75.

p=0.02), but marginal effects for site and survival (X^2 (1)=2.47, p=0.12).

Combining the results from all 3 years, uninfected ticks were more likely to die during

the winter than infected ticks (Factorial X^2 (1)=24.25, p<0001, N=301 ticks). The overall

relative risk of death during overwintering for uninfected ticks (49.25% dead, N=134) vs

infected ticks (22.16% dead, N=167) was 1.83 (95% confidence interval 1.4 – 2.4). In other

words, uninfected ticks were almost twice as likely to die during the winter as infected ticks.

Tick overwintering mortality varied significantly across years (Year 1: 58.3% (N=21/36,

Year 2: 22.1%% (N=32/145), Year 3: 41.7% (N=50/120), post hoc X^2 (2)= 21.77, *P*<0.001).

The infection rate also varied across years (Year 1: 30.6% (11/36), Year 2: (62.1% (90/145),

Year 3: 55.0% (66/120), post hoc X^2 (2)=14.8, p=0.0006).

Temperature and Relative Humidity conditions

The forest sites had higher minimum temperatures than the dune sites (Table S3). Dune

grass sites had a greater number of days with minimum temperatures below -5°C, a temperature

at which ticks are unlikely to show movement in the field (Clark 1995; Carroll & Kramer 2003).

29.7% days were below -5°C in the dune grass (106 days/357 days over 3 winters) compared

with 14.6% of days in the forest sites (52 days/357 days over 3 winters)(X^2 (1)=23.7, p<0.0001).

Year 1 had the lowest temperatures at the forest and dune grass sites (Table S3), and this year

also had the lowest rate of winter survival.

Implications for tick population growth

Although *B. burgdorferi* does not appear to be vertically transmitted (Rollend et al.

2013), an increase in the survival of infected adult females could lead to an increase in the

number of tick larvae in the next generation. Each additional infected female that survives the

winter could supply up to 3,000 extra larvae (Mount et al. 1997) to the next generation. After

five winters, an area in which 100% of the ticks are infected could have more than 6 times as

many ticks as an area with no infected ticks (see Supplementary Information for details).

Discussion

Overwintering mortality varied across the three years, as has been observed in other field

studies (Lindsay et al. 1998). Microclimate had an impact on mortality, but only in the first year

in which a higher proportion of ticks survived in the forested site compared to the dune grass

site. Other studies have also found that whether ticks have greater overwintering survival in a

particular microclimate varies across the years (e.g. Maine (Volk et al. 2022)). The first year

was also the coldest year in our study, and it may be that the effect of microclimate is most

pronounced during severe winters. Despite this variability in overwintering mortality across

years and sites, infected adult female ticks were more likely to survive the winter than uninfected

female ticks in each of the three winters at both sites (Fig. 1).   Therefore, this effect is robust

enough to be detectable across both years and microclimates.

Whether the correlation between infection and winter survival will impact tick population

growth depends on whether there is a causal link between infection and overwintering survival.

We did not infect our ticks, but collected them from the field. It is possible that the differential

survival was caused, not by infection, but by some pre-existing difference(s) between infected

and uninfected ticks. For example, it is possible that infected ticks must be in good condition to

survive the infection, while in the uninfected ticks, there may be a mix of ticks in good and poor

condition. This difference in initial condition between the two groups could explain the

difference in host mortality. We argue that this is unlikely. First, *B. burgdorferi* infection

produces no measurable damage to ticks and is not pathogenic (Couret et al. 2022). In a related

system, *Borrelia afzelii* infection had no effect on the survival of *I. ricinis* nymphs (Hurry et al.

2021). On the contrary, *B. burgdorferi* appears to benefit *I. scapularis* (Couret et al. 2022).

Changes in gene expression in infected ticks suggest that infected ticks have an enhanced feeding

ability, as well as other advantages (Couret et al. 2022).

Another possibility is that the number of *B. burgdorferi* spirochetes may decrease with

time. Therefore, some of our 'uninfected' ticks may have been infected ticks that had 'lost' their

Borrelia bacteria due to age. Older ticks are more likely to die, and this would inflate the death

rate of uninfected ticks. There is evidence of such a decline in the related tick, *I. ricinis* infected

with *Borellia afzelii* (Jacquet et al. 2017). However, even with the decline in bacterial number,

there was no change in the detection rate in *I. ricinis* (i.e. qPCR continued to identify *Borrelia*

DNA even at the lower bacterial load, Jacquet et al. 2017). In fact, the older the tick, the greater

the prevalence of *B. afzelii* in *I. ricinis* (Springer et al. 2022). Perhaps most importantly, the

bacterial load of *B. burgdorferi* does not decline in *I. scapularis* nymphs over a 12 month period

(see Fig. 2 in Samanta et al. 2022), unlike *I. ricinis* (Jacquet et al. 2017). There may be a

difference in the longevity of *Borrelia* in *I. ricinis* compared with its longevity in *I. scapularis*.

Finally, *B. burgdorferi* spirochetes have been found to persist in the gut for the entire lifespan of

I. scapularis (Pal et al. 2021), making it unlikely that they would be 'lost' over time. Therefore,

there is no evidence that bacterial loss with time would produce a significant false negative rate.

An additional concern is whether it is possible to reliably identify *Borrelia* DNA from

dead ticks. We demonstrated that we were able to measure *Borrelia* DNA in ticks that were

subjected to daily freeze/thaw for 10 weeks, using a greater temperature fluctuation than they

faced in the field (See Supplementary Information). DNA is an extremely stable molecule, with

an approximate half-life of 521 years (Allentoft et al. 2012). Moreover, *Borrelia* is found largely

in the midgut in the tick (Kurokawa et al. 2020). In humans, the intestinal microbiota of people

buried in cold climates from 100 to 3000 years ago can still be identified (Rollo et al. 2007).

And although nucleases in the environment do degrade DNA post mortem, low temperatures

inactivate these enzymes (Alaeddini et al. 2010) and DNA in dead cells remains intact when

frozen (Burkert et al. 2019). For organisms kept at 15°C, it would take 5,000 to 10,000 years for

an 800 bp fragment to degrade (Alaeddini et al. 2010). Moreover, the ticks in our study were

kept in microcosms, and these tubes isolated them from soil invertebrates and most other

organisms, decreasing the risk of degradation. Also, the *Borrelia* DNA was protected by the

ticks' hard exoskeleton during the winter, which was still intact in all of our dead ticks. Perhaps

most importantly, we tested the DNA integrity of every sample, and did not find evidence of

degradation in the DNA from dead ticks. For these reasons, we do not believe that our results

are due to an inability to measure *Borrelia* DNA in dead ticks.

B. burgdorferi infection could result in increased winter survival in *I. scapularis* by a

number of non-mutually exclusive mechanisms. *B. burgdorferi* infection is known to cause

physiological and behavioural changes in *I. scapularis* (Benelli 2020). The evidence suggests

that these changes could increase tick condition. For example, *B. burgdorferi* infection enhances

feeding in the tick (Courtet et al. 2022), allowing infected ticks to gain more resources during

their feeding period. This increase in resources could also result in increased overwintering

survival. *I. scapularis* also responds to a *B. burgdorferi* infection with an upregulation of a

number of immune genes (Narasimhan et al. 2017; Tang et al. 2021; Hart et al. 2022). In insects,

activating an immune response can enhance cold tolerance (Krams et al. 2011). Concomitant

with these changes, infection with *B. burgdorferi* induces the expression of a suite of protective

genes in *I. scapularis*. For example, *B. burgdorferi* infection results in the increased expression

of heat shock proteins (Hart et al. 2022), Glutathione peroxidase and Glutathione-S-transferase

(Narasimhan et al. 2017). These type of molecules are known to be cryoprotectant in insects

(King & MacRae, 2015; Overgaard & MacMillan, 2017). Therefore, *B. burgdorferi* could

enhance winter survival by providing an immune stimulus, leading to the co-activation of a

variety of protective mechanisms (e.g. Adamo 2022). We favour this hypothesis, especially as

B. burgdorferi does not appear to damage ticks (Courtet et al. 2022). In other words, one of the

results of the complex interaction between tick and spirochete (Kurokawa et al. 2020) may be

hormesis (Berry and Lopez-Martinez 2020), i.e. a mild stressor (i.e. *B. burgdorferi*) leads to

improved *I. scapularis* hardiness due to an upregulation of protective mechanisms.

Although *B. burgdorferi* and *I. scapularis* have co-existed for at least 20,000 year

(Walter et al. 2017), whether *B. burgdorferi* actively manipulates its host to increase its

overwintering survival remains unclear. The ability of an antimicrobial protein to also function

as a cryoprotective molecule (Neelakanta et al. 2010) makes the issue difficult to entangle. An

additional problem for the manipulation hypothesis, however, is that *B. burgdorferi* in the adult

tick has probably reached a dead end in terms of transmission. However, *B. burgdorferi* could

benefit by improving host winter survival in larvae or nymphs. Future studies should examine

whether infection in these life stages also increases winter survival.

A recent survey of ticks on the south shore of Nova Scotia found that about 4% of ticks

were infected with *A. phagocytophilum* (Guillot et al. 2020). Co-infections of *A.*

phagocytophilum and *B. burgdorferi* are typically rare, and occur at the rate expected if the

pathogens were assorting randomly (Steiner et al. 2008). Therefore, it is unlikely that the

enhanced overwintering survival we observed was due to *A. phagocytophilum* infection, not *B.*

burgdorferi. Moreover, although our focus was on *B. burgdorferi*, we did run primers for *A.*

phagocytophilum, and we did not find it in our samples (see Supplementary Information).

Finally, ticks infected with this pathogen could have been in either the *B. burgdorferi* infected or

uninfected group. Such a random assortment of *A. phagocytophilum* infected ticks would not

produce the large increase in winter survival that we found in ticks infected with *B. burgdorferi*.

Initial tick populations of *I. scapularis* are often *B. burgdorferi* free (Ogden et al. 2010;

Ogden et al. 2013).   Tick population size needs to reach a certain threshold for *B. burgdorferi* to

become endemic (Norman et al. 1999). Our study suggests that infection itself may increase the

likelihood of this event by boosting tick population size, especially during winters with low

overwintering survival for uninfected ticks.

The potential real world impact of *B. burgdorferi* infection on tick populations urges

further study of this phenomenon. If increased overwintering survival of adult females leads to

greater population growth, then tick population numbers should grow more rapidly once *B.*

burgdorferi infects the population. There is some evidence for this pattern (Elias et al. 2022).

But this pattern will only exist where other factors (such as the size of the mouse population, also

needed for tick survival) are conducive to a rapid increase in tick numbers (see Ogden et al.

2005; Eisen et al. 2016; Wallace et al. 2019; Ogden et al. 2021; Winter et al. 2021) for a

discussion of these factors). Therefore, the true impact of infection on tick numbers will vary

from region to region, and, in some cases infection may not provide ticks with an advantage (see

discussion in Lemoine et al. 2022).

Regardless of exact nature of the link between *B. burgdorferi* and the increased

overwintering survival of infected adult ticks, our results suggest that adult ticks found in the late

winter/early spring are more likely to be infected than adult ticks found in the late fall.

Unfortunately, we were unable to locate any data to show whether this occurs in areas with

endemic *I. scapularis* populations. Whether adult ticks are more likely to be infected with *B.*

burgdorferi by mid-spring appears to vary by year (Roome et al. 2018).

Once *B. burgdorferi* enters a population, infection could have a synergistic interaction

with other environmental changes (e.g. such as increasing temperatures due to climate change),

leading to boosted population numbers in previously marginal areas for *I. scapularis*. In some

regions, it may make the difference between the local extinction of a population and its growth.

Acknowledgements

This research was supported by a Canadian Lyme Foundation Venture Grant (SAA), a Discovery

Grant from the Natural Sciences and Engineering Council of Canada (NSERC) (SAA), MITACS

Accelerator Award (AEN) and Killam Post Doctoral Fellowship Award (LVF). We thank Dr. L.

McMillan and Dr. S. Mackinnon for helpful discussions and thank D. Adamo, A. Barkhouse, L.

Mace, and J. Zbarsky for help with DNA extraction and PCR. We thank the Harrison Lewis

Discovery Centre for hosting our field study and Thomas Raddall Provincial Park for permission

to drag for ticks. We thank the Public Health Agency of Canada (PHAC) for the gift of the

OspA for our positive control.

Disclosure

The authors declare that they have no conflicts of interest to disclose.

References

Adamo, S.A. (2022) The integrated defense system: optimizing defense against predators, pathogens,
and poisons. *Integrative and Comparative Biology, * **62**, 1536-1546.
Adamo, S.A., El Nabbout, A., Ferguson, L.V., Zbarsky, J.S. and Faraone, N. (2022) Balsam fir (*Abies*
balsamea) needles and their essential oil kill overwintering ticks (*Ixodes scapularis*) at cold
temperatures. *Scientific reports, * **12**, 12999.
Alaeddini, R., Walsh, S.J. and Abbas, A. (2010) Forensic implications of genetic analyses from
degraded DNA—a review. *Forensic science international: genetics, * **4**, 148-157.
Allentoft, M.E., Collins, M., Harker, D., Haile, J., Oskam, C.L., Hale, M.L., Campos, P.F., Samaniego,
J.A., Gilbert, M.T.P. and Willerslev, E. (2012) The half-life of DNA in bone: measuring decay
kinetics in 158 dated fossils. *Proceedings of the Royal Society B: Biological Sciences, * **279**, 4724-
4733.
Aranda, P.S., LaJoie, D.M., and Jorcyk, C.L. (2012) Bleach gel: A simple agarose gel for analyzing
RNA quality. *Electrophoresis*, 33: 366-369.
Benelli, G. (2020) Pathogens manipulating tick behavior—through a glass, darkly. *Pathogens, * **9**, 664.
Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: a practical and powerful
approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological), *
57, 289-300.
Berry Iii, R. and López-Martínez, G. (2020) A dose of experimental hormesis: When mild stress protects
and improves animal performance. *Comparative Biochemistry and Physiology Part A:*
*Molecular & Integrative Physiology, * **242**, 110658.
Boucias, D.G. and Penland, J.C. (1998) *Principles of insect pathology, * Kluwer, Boston.

Brunner, J.L., Killilea, M. and Ostfeld, R.S. (2014) Overwintering survival of nymphal Ixodes scapularis
(Acari: Ixodidae) under natural conditions. *Journal of Medical Entomology,* **49**, 981-987.

Burgdorfer, W., Barbour, A.G., Hayes, S.F., Benach, J.L., Grunwaldt, E. and Davis, J.P. (1982) Lyme
Disease: a tick-borne spirochetosis. *Science,* **216**, 1317-1319.

Burkert, A., Douglas, T.A., Waldrop, M.P. and Mackelprang, R. (2019) Changes in the active, dead, and
dormant microbial community structure across a Pleistocene permafrost chronosequence.
Applied and Environmental Microbiology, **85**, e02646-18.

Burks, C.S., Stewart Jr, R.L., Needham, G.R. and Lee Jr, R.E. (1996) The role of direct chilling injury
and inoculative freezing in cold tolerance of *Amblyomma americanum*, *Dermacentor variabilis*
and *Ixodes scapularis*. *Physiological Entomology,* **21**, 44-50.

Burtis, J.C. (2017) Method for the Efficient Deployment and Recovery of *Ixodes scapularis* (Acari:
Ixodidae) Nymphs and Engorged Larvae from Field Microcosms. *Journal of Medical*
Entomology, **54**, 1778-1782.

Burtis, J.C., Fahey, T.J. and Yavitt, J.B. (2019) Survival and energy use of Ixodes scapularis nymphs
throughout their overwintering period. *Parasitology,* **146**, 781-790.

Burtis, J.C., Ostfeld, R.S., Yavitt, J.B. and Fahey, T.J. (2016) The Relationship Between Soil
Arthropods and the Overwinter Survival of Ixodes scapularis (Acari: Ixodidae) Under
Manipulated Snow Cover. *Journal of Medical Entomology,* **53**, 225-229.

Burtis, J.C. and Pflueger, C. (2017) Interactions between soil-dwelling arthropod predators and Ixodes
scapularis under laboratory and field conditions. *Ecosphere,* **8**, 12.

Carroll, J.F. and Kramer, M. (2003) Winter activity of Ixodes scapularis (Acari : Ixodidae) and the
operation of deer-targeted tick control devices in Maryland. *Journal of Medical Entomology,* **40**,
238-244.

Clark, D.D. (1995) Lower temperature limits for activity of several Ixodid ticks: Effects of body size
and rate of temperature change. *Journal of Medical Entomology,* **32**, 449-452.

Clow, K.M., Ogden, N.H., Lindsay, L.R., Michel, P., Pearl, D.L. and Jardine, C.M. (2017) The influence
of abiotic and biotic factors on the invasion of *Ixodes scapularis* in Ontario, Canada. *Ticks and*
Tick-Borne Diseases, **8**, 554-563.

Couret, J., Schofield, S. and Narasimhan, S. (2022) The environment, the tick, and the pathogen–It is an
ensemble. *Frontiers in Cellular and Infection Microbiology*, 1644.

Courtney, J.W., Kostelnik, L.M., Zeidner, N.S. and Massung, R.F. (2004) Multiplex real-time PCR for

detection of *Anaplasma phagocytophilum* and *Borrelia burgdorferi. Journal of Clinical*

Microbiology, **42**, 3164-3168.

Difrisco, J. and Jaeger, J. (2019) Beyond networks: mechanism and process in evo-devo. *Biology &*

Philosophy, **34**, 24.

Difrisco, J. and Jaeger, J. (2020) Genetic Causation in Complex Regulatory Systems: An Integrative

Dynamic Perspective. *Bioessays,* **42**, 10.

Dolezal, T., Krejcova, G., Bajgar, A., Nedbalova, P. and Strasser, P. (2019) Molecular regulations of

metabolism during immune response in insects. *Insect Biochemistry and Molecular Biology,* **109**,

31-42.

Duffy, D.C. and Campbell, S.R. (1994) Ambient air temperature as a predictor of activity of adult *Ixodes*

scapularis (Acari: Ixodidae). *Journal of Medical Entomology,* **31**, 178-180.

Dunphy, G.B. and Downer, R.G.H. (1994) OCTOPAMINE, A MODULATOR OF THE HEMOCYTIC

NODULATION RESPONSE OF NONIMMUNE GALLERIA-MELLONELLA LARVAE.

Journal of Insect Physiology, **40**, 267-272.

Eisen, R.J., Eisen, L., Ogden, N.H. and Beard, C.B. (2016) Linkages of Weather and Climate With

I*xodes scapularis* and *Ixodes pacificus* (Acari: Ixodidae), Enzootic Transmission of *Borrelia*

burgdorferi, and Lyme Disease in North America. *Journal of Medical Entomology,* **53**, 250-261.

Eisen, R.J., Piesman, J., Zielinski-Gutierrez, E. and Eisen, L. (2012) What do we need to know about

disease ecology to prevent Lyme disease in the northeastern United States? *Journal of medical*

entomology, **49**, 11-22.

Elias, S.P., Witham, J.W., Schneider, E.F., Rand, P.W., Hunter Jr, M.L., Lubelczyk, C. and Smith Jr,

R.P. (2022) Emergence of *Ixodes scapularis* (Acari: Ixodidae) in a Small Mammal Population in

a Coastal Oak-Pine Forest, Maine, USA. *Journal of medical entomology,* **59**, 725-740.

Estrada-Peña, A. (2002) Increasing habitat suitability in the United States for the tick that transmits

Lyme disease: a remote sensing approach. *Environmental health perspectives,* **110**, 635-640.

Ferguson, L.V., Dhakal, P., Lebenzon, J.E., Heinrichs, D.E., Bucking, C. and Sinclair, B.J. (2018)

Seasonal shifts in the insect gut microbiome are concurrent with changes in cold tolerance and

immunity. *Functional ecology,* **32**, 2357-2368.

Galenza, A. and Foley, E. (2019) Immunometabolism: Insights from the Drosophila model.

Developmental and Comparative Immunology, **94**, 22-34.

Gray, J. and Kahl, O. (2022) Tick Ecology and the Eco-Epidemiology of *Borrelia burgdorferi* sensu
lato. *Lyme Borreliosis.* eds. K.P. Hunfeld & O. Kahls), pp. 31-45. Springer, Switzerland.

Guillot, C., Badcock, J., Clow, K., Cram, J., Dergousoff, S., Dibernardo, A., Evason, M., Fraser, E.,
Galanis, E. and Gasmi, S. (2020) Laboratory Biosafety: Sentinel surveillance of Lyme disease
risk in Canada, 2019: Results from the first year of the Canadian Lyme Sentinel Network
(CaLSeN). *Canada Communicable Disease Report,* **46**, 354.

Halsey, S.J., Allan, B.F. and Miller, J.R. (2018) The role of Ixodes scapularis, Borrelia burgdorferi and
wildlife hosts in Lyme disease prevalence: a quantitative review. *Ticks and tick-borne diseases,*
9, 1103-1114.

Hammond-Collins, K., Tremblay, M., Milord, F., Baron, G., Bouchard, C., Kotchi, S.O., Lambert, L.,
Leighton, P., Ogden, N.H. and Rees, E.E. (2022) An ecological approach to predict areas with
established populations of *Ixodes scapularis* in Quebec, Canada. *Ticks and Tick-borne Diseases,*
13, 102040.

Hart, C.E., Middleton, F.A. and Thangamani, S. (2022) Infection with Borrelia burgdorferi Increases the
Replication and Dissemination of Coinfecting Powassan Virus in Ixodes scapularis Ticks.
Viruses, **14**, 1584.

Hatchette, T.F., Johnston, B.L., Schleihauf, E., Mask, A., Haldane, D., Drebot, M., Baikie, M., Cole,
T.J., Fleming, S. and Gould, R. (2015) Epidemiology of Lyme Disease, Nova Scotia, Canada,
2002–2013. *Emerging infectious diseases,* **21**, 1751.

Hector, T.E., Sgro, C.M. and Hall, M.D. The influence of immune activation on thermal tolerance along
a latitudinal cline. *Journal of Evolutionary Biology*, 11.

Heisig, M., Abraham, N.M., Liu, L., Neelakanta, G., Mattessich, S., Sultana, H., Shang, Z., Ansari, J.M.,
Killiam, C. and Walker, W. (2014) Antivirulence properties of an antifreeze protein. *Cell*
Reports, **9**, 417-424.

Helble, J.D., Mccarthy, J.E. and Hu, L.T. (2021) Interactions between Borrelia burgdorferi and its hosts
across the enzootic cycle. *Parasite immunology,* **43**, e12816.

Herrmann, C. and Gern, L. (2010) Survival of *Ixodes ricinus* (Acari: Ixodidae) Under Challenging
Conditions of Temperature and Humidity Is Influenced by *Borrelia burgdorferi* sensu lato
Infection. *Journal of Medical Entomology,* **47**, 1196-1204.

Herrmann, C. and Gern, L. (2015) Search for blood or water is influenced by *Borrelia burgdorferi* in
Ixodes ricinus. *Parasites & vectors,* **8**, 1-8.

Hurry, G., Maluenda, E., Sarr, A., Belli, A., Hamilton, P.T., Duron, O., Plantard, O. and Voordouw,

M.J. (2021) Infection with Borrelia afzelii and manipulation of the egg surface microbiota have

no effect on the fitness of immature Ixodes ricinus ticks. *Scientific Reports,* **11**, 10686.

Jacquet, M., Genné, D., Belli, A., Maluenda, E., Sarr, A. and Voordouw, M.J. (2017) The abundance of

the Lyme disease pathogen Borrelia afzelii declines over time in the tick vector Ixodes ricinus.

Parasites & vectors, **10**, 1-10.

Kilpatrick, A.M., Dobson, A.D.M., Levi, T., Salkeld, D.J., Swei, A., Ginsberg, H.S., Kjemtrup, A.,

Padgett, K.A., Jensen, P.M., Fish, D., Ogden, N.H. and Diuk-Wasser, M.A. (2017) Lyme disease

ecology in a changing world: consensus, uncertainty and critical gaps for improving control.

Philosophical Transactions of the Royal Society B-Biological Sciences, **372**, 15.

King, A.M. and Macrae, T.H. (2015) Insect heat shock proteins during stress and diapause. *Annual*

review of entomology, **60**, 59-75.

Krams, I., Daukšte, J., Kivleniece, I., Krama, T. and Rantala, M.J. (2011) Overwinter survival depends

on immune defence and body length in male *Aquarius najas* water striders. *Entomologia*

Experimentalis et Applicata, **140**, 45-51.

Kurokawa, C., Lynn, G.E., Pedra, J.H., Pal, U., Narasimhan, S. and Fikrig, E. (2020) Interactions

between *Borrelia burgdorferi* and ticks. *Nature Reviews Microbiology,* **18**, 587-600.

Lemoine, M., Cornetti, L. and Tschirren, B. (2018) Does Borrelia burgdorferi sensu lato facilitate the

colonisation of marginal habitats by Ixodes ricinus? A correlative study in the Swiss Alps.

BioRxiv, 273490.

Lindsay, L.R., Barker, I.K., Surgeoner, G.A., Mcewen, S.A., Gillespie, T.J. and Addison, E.M. (1998)

Survival and development of the different life stages of *Ixodes scapularis* (Acari : Ixodidae) held

within four habitats on Long Point, Ontario, Canada. *Journal of Medical Entomology,* **35**, 189-

199.

Lindsay, L.R., Barker, I.K., Surgeoner, G.A., Mcewen, S.A., Gillespie, T.J. and Robinson, J.T. (1995)

Survival and development of *Ixodes scapularis* (Acari, Ixodidae) under various climatic

conditions in Ontario, Canada. *Journal of Medical Entomology,* **32**, 143-152.

Linske, M.A., Stafford, K.C., Williams, S.C., Lubelczyk, C.B., Welch, M. and Henderson, E.F. (2019)

Impacts of Deciduous Leaf Litter and Snow Presence on Nymphal *Ixodes scapularis* (Acari:

Ixodidae) Overwintering Survival in Coastal New England, USA. *Insects,* **10**.

Miyashita, A. and Adamo, S.A. (2020) Stayin' Alive: Endocrinological Stress Responses in Insects.

Advances in Invertebrate (Neuro)Endocrinology. eds. S. Saleuddin, A.B. Lange & I. Orchards),

pp. 283-324. Apple Academic Press/CRC Press, Burlington, ON.

Mount, G., Haile, D. and Daniels, E. (1997) Simulation of blacklegged tick (Acari: Ixodidae) population

dynamics and transmission of *Borrelia burgdorferi*. *Journal of Medical Entomology,* **34**, 461-

484.

Narasimhan, S., Schuijt, T.J., Abraham, N.M., Rajeevan, N., Coumou, J., Graham, M., Robson, A., Wu,

M.-J., Daffre, S. and Hovius, J.W. (2017) Modulation of the tick gut milieu by a secreted tick

protein favors Borrelia burgdorferi colonization. *Nature communications,* **8**, 184.

Neelakanta, G., Sultana, H., Fish, D., Anderson, J.F. and Fikrig, E. (2010) *Anaplasma phagocytophilum*

induces I*xodes scapularis* ticks to express an antifreeze glycoprotein gene that enhances their

survival in the cold. *Journal of Clinical Investigation,* **120**, 3179-3190.

Norman, R., Bowers, R., Begon, M. and Hudson, P.J. (1999) Persistence of tick-borne virus in the

presence of multiple host species: tick reservoirs and parasite mediated competition. *Journal of*

Theoretical Biology, **200**, 111-118.

Ogden, N., Bigras-Poulin, M., O'callaghan, C., Barker, I., Lindsay, L., Maarouf, A., Smoyer-Tomic, K.,

Waltner-Toews, D. and Charron, D. (2005) A dynamic population model to investigate effects of

climate on geographic range and seasonality of the tick *Ixodes scapularis*. *International journal*

for parasitology, **35**, 375-389.

Ogden, N.H., Beard, C.B., Ginsberg, H.S. and Tsao, J.I. (2021) Possible effects of climate change on

Ixodid ticks and the pathogens they transmit: Predictions and observations. *Journal of Medical*

Entomology, **58**, 1536-1545.

Ogden, N.H., Bouchard, C., Kurtenbach, K., Margos, G., Lindsay, L.R., Trudel, L., Nguon, S. and

Milord, F. (2010) Active and passive surveillance and phylogenetic analysis of *Borrelia*

burgdorferi elucidate the process of Lyme disease risk emergence in Canada. *Environmental*

Health Perspectives, **118**, 909-914.

Ogden, N.H., Lindsay, L.R. and Leighton, P.A. (2013) Predicting the rate of invasion of the agent of

Lyme disease Borrelia burgdorferi. *Journal of Applied Ecology,* **50**, 510-518.

Overgaard, J. and Macmillan, H.A. (2017) The integrative physiology of insect chill tolerance. *Annual*

review of physiology, **79**, 187-208.

Pal, U., Kitsou, C., Drecktrah, D., Yas, O.B. and Fikrig, E. (2021) Interactions between ticks and lyme
540   disease spirochetes. *Current issues in molecular biology,* **42**, 113-144.

541   Rollend, L., Fish, D. and Childs, J.E. (2013) Transovarial transmission of Borrelia spirochetes by Ixodes
542   scapularis: a summary of the literature and recent observations. *Ticks and tick-borne diseases,* **4**,
543   46-51.

544   Rollo, F., Luciani, S., Marota, I., Olivieri, C. and Ermini, L. (2007) Persistence and decay of the
545   intestinal microbiota's DNA in glacier mummies from the Alps. *Journal of archaeological*
546   *science,* **34**, 1294-1305.

547   Roome, A., Spathis, R., Hill, L., Darcy, J.M. and Garruto, R.M. (2018) Lyme disease transmission risk:
548   seasonal variation in the built environment. *Healthcare.* p. 84. MDPI.

549   Scott, J.D., Foley, J.E., Clark, K.L., Anderson, J.F., Durden, L.A., Manord, J.M. and Smith, M.L. (2016)
550   Established population of blacklegged ticks with high infection prevalence for the Lyme disease
551   bacterium, Borrelia burgdorferi sensu lato, on Corkscrew Island, Kenora District, Ontario.
552   *International journal of medical sciences,* **13**, 881.

553   Sinclair, B.J., Ferguson, L.V., Salehipour-Shirazi, G. and Macmillan, H.A. (2013) Cross-tolerance and
554   Cross-talk in the Cold: Relating Low Temperatures to Desiccation and Immune Stress in Insects.
555   *Integrative and Comparative Biology,* **53**, 545-556.

556   Springer, A., Jordan, D., Glass, A., Kahl, O., Fingerle, V., Girl, P., Chitimia-Dobler, L. and Strube, C.
557   (2022) Borrelia infections in ageing ticks: relationship with morphometric age ratio in field-
558   collected Ixodes ricinus nymphs. *Microorganisms,* **10**, 166.

559   Steiner, F.E., Pinger, R.R., Vann, C.N., Grindle, N., Civitello, D., Clay, K. and Fuqua, C. (2008)
560   Infection and co-infection rates of *Anaplasma phagocytophilum* variants, *Babesia* spp., *Borrelia*
561   *burgdorferi,* and the rickettsial endosymbiont in *Ixodes scapularis* (Acari: Ixodidae) from sites in
562   Indiana, Maine, Pennsylvania, and Wisconsin. *Journal of Medical Entomology,* **45**, 289-297.

563   Sterling, P. and Laughlin, S. (2015) *Principles of Neural Design,* MIT Press, Cambridge, MA.

564   Stoney, R., Robertson, D.L., Nenadic, G. and Schwartz, J.M. (2018) Mapping biological process
565   relationships and disease perturbations within a pathway network. *Npj Systems Biology and*
566   *Applications,* **4**, 11.

567   Tang, X., Cao, Y., Arora, G., Hwang, J., Sajid, A., Brown, C.L., Mehta, S., Marín-López, A., Chuang,
568   Y.-M. and Wu, M.-J. (2021) The Lyme disease agent co-opts adiponectin receptor-mediated
569   signaling in its arthropod vector. *Elife,* **10**, e72568.

Vandyk, J.K., Bartholomew, D.M., Rowley, W.A. and Platt, K.B. (1996) Survival of *Ixodes scapularis*

(Acari: Ixodidae) exposed to cold. *Journal of Medical Entomology,* **33**, 6-10.

Volk, M.R., Lubelczyk, C.B., Johnston, J.C., Levesque, D.L. and Gardner, A.M. (2022) Microclimate

conditions alter *Ixodes scapularis* (Acari: Ixodidae) overwinter survival across climate gradients

in Maine, United States. *Ticks and tick-borne diseases,* **13**, 101872.

Wallace, D., Ratti, V., Kodali, A., Winter, J.M., Ayres, M.P., Chipman, J.W., Aoki, C.F., Osterberg,

E.C., Silvanic, C. and Partridge, T.F. (2019) Effect of rising temperature on Lyme disease:

Ixodes scapularis population dynamics and *Borrelia burgdorferi* transmission and prevalence.

Canadian Journal of Infectious Diseases and Medical Microbiology, **2019**.

Walter, K.S., Carpi, G., Caccone, A. and Diuk-Wasser, M.A. (2017) Genomic insights into the ancient

spread of Lyme disease across North America. *Nature Ecology & Evolution,* **1**, 1569-1576.

Williams, C.M., Henry, H.A. and Sinclair, B.J. (2015) Cold truths: how winter drives responses of

terrestrial organisms to climate change. *Biological Reviews,* **90**, 214-235.

Winter, J.M., Partridge, T.F., Wallace, D., Chipman, J.W., Ayres, M.P., Osterberg, E.C. and Dekker,

E.R. (2021) Modeling the Sensitivity of Blacklegged Ticks (*Ixodes scapularis*) to Temperature

and Land Cover in the Northeastern United States. *Journal of Medical Entomology,* **58**, 416-427.

Author Contributions

588   SAA formulated the hypothesis, assisted with experimental design and data analysis, and led

589   the writing of the paper. AEN assisted with experimental design and data analysis, wrote a first

590   draft and performed the study. AM validated the qPCR for our study. LVF helped formulate the

591   question, assisted with experimental design and helped write the paper.

Data Availability

594   Data are available in the Mendeley repository at: El Nabbout, Amal (2023), "Data 2018-2021

595   HLCF", Mendeley Data, V1, doi: 10.17632/tzjdxmyctv.1

Figure Legends

598   **Figure 1.** Percentage of *I. scapularis* infected with *B. burgdorferi* that survive the winter

599   compared with uninfected ticks at each study site over the three winters. Numbers above each

600   bar indicate the sample size. Dark bars denote uninfected ticks, while light bars denote infected

601   ticks. Un= uninfected, In=Infected.

 Supplementary Information

606   Adult ticks (*Ixodes scapularis*) infected with *Borrelia burgdorferi* have increased overwintering

607   survival, with implications for tick population growth.

609   Amal El Nabbout[1], Laura V. Ferguson[2], Atsushi Miyashita[3], Shelley A. Adamo[4]

611   1. Dept. Biology, Dalhousie University, Halifax, NS Canada

612   2. Dept. Biology, Acadia University, Wolfville, NS Canada

613   3. Institute of Medical Mycology, Teikyo University, Tokyo, Japan

614   4. Dept. Psychology and Neuroscience, Dalhousie University, NS Canada

615
616

Determination of Tick Infection Status

The procedure used to determine tick infection status was as described in Courtney et al.,

(2004), and is repeated here. Ticks were identified using standard keys (Keirans and Likwack,

1989). DNA was extracted from *I. scapularis* following the Public Health Agency of Canada

(PHAC) protocol, adopted from Courtney et al. (2004). We used a Qiagen QIAamp© DNA Mini

Kit with an extra bead tube step (MN Bead Tubes Type D (Macherey-Nagel)) to improve the

DNA extraction process. MN Bead Tubes Type D (Macherey-Nagel) were filled with 200 μl of

Qiagen ATL buffer. Individual ticks were crushed thoroughly using a Bel-Art® Disposable

Pestle. Bead tubes were vortexed for 30 minutes. 20 μl of Qiagen Proteinase K was then pipetted

into each bead tube. Bead tubes were incubated at 56°C for 60-90 minutes. 200 μl of Qiagen AL

buffer was added to each bead tube and tubes were vortexed for 15 seconds and then incubated at

70°C for 10 minutes. Finally, 200 μl of 100% ethanol was pipetted into each bead tube and then

tubes were vortexed for 15 seconds. DNA extraction followed QIAamp© DNA mini kit

instructions.

DNA was then screened for evidence of *B. burgdorferi* infection using a multiplex real-

time PCR targeting the 23S rRNA of *B. burgdorferi*. *B. burgdorferi* infection was then

confirmed in positive samples using primers targeting the ospA gene. Each sample was tested in

triplicates and all plates included: a no template control (5 Microliters of milliQ water), a

25 Cq value positive control, and a 30 Cq value positive control. The thermocycling conditions

occurred as follows: activation of enzyme AmpErase at 50°C for 2 minutes, denaturation of

AmpErase and activation of AmpliTaq Gold® Polymerase at 95°C for 10 minutes, followed by

40 cycles of amplification at 95°C for 15 seconds and annealing at 60°C for 1 minute.

The integrity of DNA samples was assessed by gel electrophoresis® (Aranda et al., 2012)

using a Gel Doc™ XR+ Digital Gel Imaging System® from Bio-Rad. DNA quantity was

determined using a Qubit fluorometer.

Although we ran primers for *Anaplasma phagocytophilum*, we found no evidence for it in

our ticks, i.e. there was no increase in fluorescence even after 40 cycles.

Table S1. Primer sequences used in Real-time PCR [3].

	Primer or Probe	Sequence 5' to 3'
Borrelia	Bb 23Sf	cgagtcttaaaagggcgatttagt
burgdorferi	Bb 23Sr	gcttcagcctggccataaatag
	Bb 23S Probe FAM-TAMRA	agatgtggtagacccgaagccgagtg
	*ospA*f	ctggggaagtttcagttgaac
	*ospA*r	ttggtgccatttgagtcgta
	ospA Probe FAM-BBQ	ctgcagcttggaattcaggcactt
Anaplasma	Ap *msp2*f	atggaaggtagtgttggttatggtatt
phagocytophilum	Ap *msp2*r	ttggtcttgaagcgctcgta
	Ap *msp2* Probe VIC-TAMRA	tggtgccagggttgagcttgagattg

The efficiency of the Bb23S primer was 92.8%. For the ospA primer, the efficiency was

100.9%.

Detecting B. burgdorferi in Dead Ticks

Our experimental design depended on our ability to detect *B. burgdorferi* in ticks that

may die at various times throughout the field experiment. As a result, we performed experiments

to determine whether *B. burgdorferi* would be detectable in dead ticks subjected to freeze/thaw

for up to 10 weeks. Ticks were exposed to a fluctuating thermal regime (-5°C to 10°C) in 12-

hour increments for 10 weeks This temperature regime exposed these samples to more

freeze/thaw events than they would have experienced in the field (Table S3). We focused on the

number of freeze/thaw events because DNA in dead cells shows little or no degradation when

kept frozen (Burkert et al., 2019), but freeze/thaw cycles accelerate DNA degradation (Rollo et

al., 2007). We cut field collected *I. scapularis* ticks in half longitudinally using a sterile razor.

The first half was used immediately for DNA extraction. The second half of the tick had a small

aliquot of Vaseline added to seal the cut end. This method provided protection from dehydration

through the open cut, in the same way that the intact cuticle of a dead tick provides protection

from desiccation. These half ticks were stored in a fluctuating thermal regime incubator (-5°C to

10°C) in 12-hour increments for 10 weeks. These temperatures produce more extreme freeze and

thaw cycles than was normally observed in the field (Table S3), which assisted us in determining

the length of time the DNA is still measurable in *Ixodes scapularis* under Nova Scotian winter

conditions. After the 10 week incubation period was completed, we extracted the DNA as

previously described and tested for *B. burgdorferi* using qPCR.

Table S2 shows the high concordance (100%) of the results for the two tick halves.

Table S2. Concordance between the two halves of each tick tested 10 weeks apart.

	Infected 2nd 1/2	Uninfected 2nd 1/2
Infected 1st 1/2	10	0
Uninfected 1st 1/2	0	20

We also tested the integrity of the DNA from both the lab and field experiments. All

DNA was found to be intact, showing distinct bands on the gel (Aranda et al., 2012). These

results show that it is possible to reliably determine *B. burgdorferi* infection in the dead ticks we

collected.

Potential Effect of Infection on Tick Population Growth

For our estimate of the potential effect of infection on population growth, we assumed that

infection will only change overwintering survival and will not impact other aspects of tick

survival (e.g. the likelihood of finding a mammalian host). Using our data, we assumed that

infected females have an 80% chance of overwintering survival and uninfected females have a

50% chance of survival. We calculated the relative growth of the two populations, one with

100% *B. burgdorferi* prevalence, another with 0%. We assumed all adult females produced 1,000

larvae and died after laying eggs (Gray and Kahl, 2022). We also assumed that ½ of larvae will

be female, and we further assumed that all larvae will survive to overwinter as adult females.

We also assumed that infected adult female ticks pay a 10% fecundity cost, based on data from

Drosophila (McKean et al. 2008).

References

Adamo, S.A., El Nabbout, A., Ferguson, L.V., Zbarsky, J.S. & Faraone, N. (2022) Balsam fir (*Abies*

balsamea) needles and their essential oil kill overwintering ticks (*Ixodes scapularis*) at cold

temperatures. *Scientific Reports,* **12,** 12999.

Aranda, P.S., LaJoie, D.M., and Jorcyk, C.L. (2012) Bleach gel: A simple agarose gel for analyzing

RNA quality. *Electrophoresis*, 33: 366-369.

Burkert, A., Douglas, T.A., Waldrop, M.P. and Mackelprang, R. (2019) Changes in the active, dead, and
dormant microbial community structure across a Pleistocene permafrost chronosequence.
Applied and Environmental Microbiology, **85**, e02646-18.
697

Courtney, J. W., Kostelnik, L. M., Zeidner, N. S. and Massung, R. F. (2004). Multiplex real-time PCR
for detection of *Anaplasma phagocytophilum* and *Borrelia burgdorferi. Journal of Clinical*
Microbiology **42**, 3164-3168.

Gray, J. and Kahl, O. (2022) Tick Ecology and the Eco-Epidemiology of *Borrelia burgdorferi* sensu
lato. *Lyme Borreliosis.* eds. K.P. Hunfeld & O. Kahls), pp. 31-45. Springer, Switzerland.

Keirans, J. E. and Litwack, T. R. (1989). Pictorial key to the adults of hard ticks, family Ixodidae
(Ixodida: Ixodoidea), east of the Mississippi River. *Journal of Medical Entomology* **26**, 435-448.

McKean, K. A., Yourth, C. P., Lazzaro, B.P., and Clark, A.G. (2008). The evolutionary costs of
immunological maintenance and deployment. *BMC evolutionary biology* 8 (1): 1-19.

Rollo, F., Luciani, S., Marota, I., Olivieri, C. and Ermini, L. (2007) Persistence and decay of the
intestinal microbiota's DNA in glacier mummies from the Alps. *Journal of archaeological*
science, **34**, 1294-1305.
710

713
714