Title: MAPK/ERK Activation in Macrophages Promotes *Leishmania*

Internalization and Pathogenesis

Running title: ERK1/2 facilitates *Leishmania* internalization

Authors: Umaru Barrie[1, 2], Katherine Floyd[1, #], Arani Datta[1], and Dawn M.

Wetzel[1, 3*]

Affiliations: [1]Department of Pediatrics, University of Texas Southwestern

Medical Center, 5323 Harry Hines Blvd, Dallas, Texas, 75390, United States;

[2]Medical Scientist Training Program, University of Texas Southwestern Medical

Center, 5323 Harry Hines Blvd, Dallas, Texas, 75390, United States;

[3]Department of Biochemistry, University of Texas Southwestern Medical Center,

5323 Harry Hines Blvd, Dallas, Texas, 75390, United States.

*Corresponding author: Dawn M. Wetzel, M.D., Ph.D.

Departments of Pediatrics (Infectious Diseases) and Biochemistry

The University of Texas Southwestern Medical Center, Dallas, TX 75019

Phone: 214-645-9510; Email: Dawn.Wetzel@UTSouthwestern.edu

[#]**Current address:** Department of Internal Medicine and Molecular Microbiology,

Washington University School of Medicine, 660 S Euclid Ave, St. Louis, MO

63110, United States

Author contributions: U.B. and D.M.W. designed and performed experiments, analyzed data, and wrote the manuscript. K.F. conducted experiments, assisted with data analysis, and edited the manuscript. A.D. helped to design and perform experiments, analyzed data, and edited the manuscript.

This file includes:

Abstract word count: 225

Lay summary word count: 141

Manuscript word count: 6236

Number of references: 63

Number of figures: 6

Number of tables: 2

Supplemental figures: 4

Supplemental tables: 3

ABSTRACT: The obligate intracellular protozoan parasite *Leishmania* binds

several host cell receptors to trigger its uptake by phagocytic cells, ultimately

resulting in visceral or cutaneous leishmaniasis. After *Leishmania* engages

receptors on macrophages and other phagocytes, a series of signaling pathways

in the host cell are activated during its internalization, which are critical for

establishment and persistence of *Leishmania* infection. Thus, preventing

Leishmania internalization by phagocytes could be a novel therapeutic strategy

for leishmaniasis. However, the host cellular machinery that mediates

promastigote and amastigote uptake is not well understood. Here, using small

molecule inhibitors of Mitogen-activated protein kinases/Extracellular signal

regulated kinases (MAPK/ERK), we demonstrate that ERK1/2 mediates

Leishmania amazonensis uptake and (to a lesser extent) phagocytosis of beads

by macrophages. We find that inhibition of MEK1/2 or ERK1/2 leads to inefficient

amastigote uptake by macrophages. Moreover, using inhibitors and primary

macrophages lacking spleen tyrosine kinase (SYK) or Abl family kinases, we

show that SYK and Abl family kinases mediate Raf, MEK, and ERK1/2 activity

and are necessary for efficient uptake. Finally, we demonstrate that trametinib, a

MEK1/2 inhibitor used clinically to treat certain cancers, significantly reduces

disease severity and parasite burden in *Leishmania*-infected mice, even if it is

started significantly after lesions develop. Our results show that maximal

Leishmania infection requires MAPK/ERK and highlight the potential for

MAPK/ERK-mediated signaling pathways to be novel therapeutic targets for

leishmaniasis.

Keywords: *Leishmania*, kinase, MAPK/ERK, parasite, macrophage

Abbreviations: BCA, Bicinchoninic acid; BMDM, bone marrow-derived macrophages; CC, cytotoxic concentration; CR3, complement receptor 3; FcγR, Fc gamma Receptor; h, hour; IC, inhibitory concentration; IL, Interleukin; INFγ, Interferon gamma; Mϕ, macrophage; m, minutes; MAPK/ERK, Mitogen-activated protein kinases/Extracellular signal regulated kinases; PKC, Protein Kinase C; PVDF, polyvinylidene difluoride; RAF, Rapidly Accelerated Fibrosarcoma; RIPA, radioimmunoprecipitation assay; p-, phosphorylated; PMA, Phorbol 12-myristate 13-acetate; SD, standard deviation; SFK, Src family kinases; SYK, spleen tyrosine kinase; TNF, Tumor Necrosis Factor

Lay summary: *Leishmania* is a single-celled parasite that causes skin ulcers or a disseminated disease in humans. Our goal is to identify new drugs to treat *Leishmania* infection. *Leishmania* must live inside human immune cells to cause disease. If *Leishmania* is not able to enter human immune cells, it dies. Our studies demonstrate how *Leishmania* infection permits a set of proteins called MAP kinases to pass signals from one protein to the next within mammalian immune cells. The resulting signals allow *Leishmania* to enter into these immune cells and survive within its host. Importantly, trametinib, a drug that prevents these signals from MAP kinases, decreases the development of skin ulcers when it is given to mice that are infected with *Leishmania*. Our findings suggest that

84   leishmaniasis could be treated with drugs that act on kinases found in humans

85   rather than the parasites themselves.

87   **One sentence summary**: An Abl2-SYK-Raf-MEK-ERK pathway facilitates

88   *Leishmania* uptake by phagocytic cells and promotes disease severity in

89   *Leishmania*-infected mice.

1. INTRODUCTION

Leishmania spp. are obligate intracellular, protozoan parasites spread by the sand fly vector [1], causing leishmaniasis, a group of diseases ranging from self-healing cutaneous ulcers [2] to deadly disseminated visceral disease [3]. Leishmaniasis represents a major global health problem, since over 12 million people currently suffer from this disease. Approximately 2 million people are newly infected by *Leishmania* spp. annually [4]. Moreover, over 350 million people in 98 countries worldwide are at risk of developing leishmaniasis [4, 5], and the disease is now spreading into previously non-endemic areas, including the United States [6, 7]. *Leishmania*, particularly the species causing cutaneous disease, are understudied, and current treatments are few, expensive, poorly effective, and toxic. Resistance is also emerging [8-11]. Thus, identifying new drugs with novel mechanisms of action that circumvent resistance mechanisms is an urgent priority for effective drug development.

When an infected sand fly encounters its host, *Leishmania* promastigotes must be internalized by phagocytic cells such as macrophages (Mϕ) to persist in the host [12, 13]. This process is then repeated as the mammalian life cycle stage, the amastigote, replicates and spreads to other phagocytic cells to cause pathogenesis [14]. Amastigotes that do not enter phagocytic cells are thought to be cleared by the host immune system and/or die from lack of essential nutrients [14]. Major advancements have been made in understanding the immunomodulatory events involved in the pathogenesis of leishmaniasis.

However, little is known about the basic molecular events underlying the

pathogenesis, especially the host cellular signaling pathways, which facilitate the

uptake of *Leishmania* by Mϕs and other phagocytes.

Based on prior studies, the Fcγ classes of receptors (FcγR) are the primary

receptors on the host cell surface that permit amastigote uptake [14], and the

complement receptor CR3 is the primary mediator of promastigote uptake [15].

Our previous studies have identified the Src [16] family kinases (SFK, and

specifically Hck, Fgr, and Lyn), Abl1 (Abl) and Abl2 (Arg) [17] kinases, and

Spleen Tyrosine Kinase (SYK) [18] as major protein families involved in

phagocytosis and *Leishmania* uptake. In particular, we have demonstrated that

Abl facilitates complement-mediated phagocytosis, and Arg enhances

immunoglobulin/ FcγR-mediated phagocytosis [16, 17]. Modulation of these

kinases via genetic manipulation or small molecule inhibitors abrogates *L.*

amazonensis internalization by Mϕs, and thus limits disease progression in mice

[16, 17]. However, our studies also indicate that these kinases have a limited role

in the host's immune response to *Leishmania* or parasite clearance [16, 17]. A

drug that simultaneously decreases parasite uptake and promotes host-

advantageous immunological responses could provide reductions in infection and

disease burden. Host-cell active agents also could be used to combat other

intracellular pathogens that employ similar uptake mechanisms.

Exactly how SFK, Abl family kinases, and SYK would regulate the events that

occur during *Leishmania* uptake and phagocytosis has not been fully established.

Others and we have demonstrated that CrkII is activated after signals have been

relayed from SFK to Arg to SYK, leading to actin-rich phagocytic cup formation

and pathogen uptake [16-19]. Interestingly, studies have been published

primarily in cancer model systems indicating that SFK [20], Abl family kinases

[20], and SYK [21, 22] also can activate MAPK/ERK signaling. MAPK/ERK are

known to directly phosphorylate cytoplasmic or nuclear proteins, including

transcription factors, that control several fundamental cellular processes from

proliferation and differentiation to innate and adaptive immune responses [23-25].

In addition, Phorbol 12-myristate 13-acetate (PMA) is known to activate

MAPK/ERK through protein kinase C (PKC) [26].

It is not clear whether MAPK/ERK signaling would mediate the uptake of

Leishmania parasites by phagocytes. Research on *Plasmodium* indicated that

inhibiting MEK1/2, which is upstream of ERK1/2, affects both adaptive and innate

immune responses, which limits parasitemia and prevents pathogenesis [27].

Previous studies have also shown that MAPK/ERK signaling modulates the

immune response to *L. amazonensis* infection to favor chronic infection, where

subversion of MAPK/ERK signaling leads to parasite survival inside phagocytic

cells, although these results vary depending on species [28-32].

Correspondingly, limiting MAPK activity using the older-generation MAPK

inhibitors PD98059 or U0126 has been shown to enhance *L. amazonensis* killing

by activating reactive oxygen species (ROS) and repressing IL-10 production

[33-36]. These results implicate ERK1/2 phosphorylation as a mechanism

through which *L. amazonensis* promotes persistence and subverts the immune

response [33]. However, experiments characterizing the role of MAPK/ERK

signaling during parasite uptake by Mϕ have not been conducted.

Here, we uncover an additional role for MAPK/ERK during *Leishmania*

internalization by phagocytes. We show that Raf-1, MEK1/2, and ERK1/2 activity are

necessary for efficient uptake of amastigotes. Host MAPK/ERK is activated by Arg

and SYK to facilitate this process. In combination with prior results [16, 17, 33], our

molecular studies elucidate an FcγR-SFK-Arg-SYK-Raf-MEK-ERK signaling relay as

a major pathway involved in establishment of leishmaniasis by facilitating parasite

uptake. In addition, using a mouse model, we establish that trametinib, a selective

MEK inhibitor that is clinically approved to treat cancer, decreases footpad swelling

and parasite burden in cutaneous leishmaniasis, even if it is started significantly after

lesions are apparent. Our experiments highlight MAPK/ERK inhibitors as potential

treatment modalities for this devastating disease.

2. <u>MATERIALS AND METHODS</u>

2.1 Mice. C57BL/6 mice were procured from Jackson Laboratory (Bar Harbor, ME). *Syk^flox/flox Cre^+* and *Arg^{−/−}Abl^flox/flox LysM Cre^+* mice are either on a pure C57BL/6 background (SYK) [18] or have been backcrossed to C57BL/6 mice at least 5 times (Abl/Arg) [16], [17]. All the experiments involving knockout mice were conducted with WT littermates to ensure control for their genetic background. The Institutional Animal Care and Use Committee at University of Texas Southwestern Medical Center provided approval for all experimental protocols.

2.2 Mammalian cell culture. RAW 264.7 cells (ATCC, Manassas, VA) were cultured in Dulbecco's modified Eagle's medium (DMEM) plus 10% endotoxin-free, heat-inactivated, fetal bovine serum (FBS) (GeminiBio, Sacramento, CA) and 1% penicillin–streptomycin as described previously [16]. FBS was checked for LPS contamination using a highly sensitive Pierce™ Chromogenic Endotoxin Quant Kit (Thermo Scientific, Cat No. A39552), and was found to be endotoxin negative (< 0.1 EU/ml). For experiments with bone marrow-derived Mφs (BMDM), cells were obtained from the humeri, tibias, and femurs of *WT, Syk^flox/flox Cre^+*, or *Arg^{−/−}Abl^flox/flox LysM-Cre^+* mice. They were differentiated to *WT, Syk^flox/flox Cre^+*, or *Arg^{−/−}Abl^flox/flox LysM-Cre^+* BMDM by culturing them in DMEM, 10% FBS, and 20% L929 cell supernatant over 7 days. Differentiation to Mφ was confirmed [17] and samples were screened for *Mycoplasma* contamination [37].

2.3 Parasite culture. *L. amazonensis* promastigotes (strain IFLA/BR/67/PH8,

provided by Norma W. Andrews, University of Maryland, College Park, MD) were

maintained at 26°C in Schneider's Drosophila medium (Sigma, S9895, St. Louis,

MO) plus 15% FBS and 1% penicillin–streptomycin [16, 17]. *L. amazonensis*

amastigotes were grown axenically at 32°C in Schneider's Drosophila medium at

pH 5.5, plus 20% FBS, 1% penicillin–streptomycin, 0.1% hemin (25 mg/mL in

50% triethanolamine), 10 mM adenine, 5 mM L-glutamine, 0.25% glucose, 0.5%

trypticase, and 40 mM sodium succinate (all from Sigma) [16, 17]. For uptake

(internalization of amastigotes and promastigotes) assays, promastigotes were

incubated for 7 days to maximize infective metacyclics (*i.e*., those isolated

through a step Percoll gradient; Sigma). Amastigotes typically were grown

axenically for 5 days prior to use [17]. To assess for amastigote growth and

replication defects during incubation in MEK/ERK activators or inhibitors, medium

containing 10 µM trametinib (Selleck Chemicals, Houston, TX), PMA (Selleck

Chemicals), or 0.1% dimethyl sulfoxide (DMSO, Sigma) was used. To ensure

infectivity, parasites were periodically passed in C57/BL6 mice prior to isolation

of amastigotes from footpad lesions.

For some experiments, a transgenic stable mNeonGreen expressing line of *L.*

amazonensis parasites (*L. amazonensis*-mNeon), containing a marker for

hygromycin resistance, was generated [18]. *L. amazonensis*-mNeon

promastigotes and amastigotes were maintained as above, but the media was

supplemented with 100 µg/mL hygromycin (Sigma). *L. amazonensis* virulence

was maintained by passage in C57BL/6 mice [17]. *L. amazonensis*-mNeonGreen

parasites were detected with an excitation wavelength of 490–494 nm and

emission of 515 nm.

2.4 *Leishmania* uptake assays. Experiments were adapted and modified as

described previously [16, 17]. BMDM (*WT, Syk^{flox/flox} Cre^+*, *Arg^{−/−}Abl^{flox/flox} LysM-Cre^+*)

or RAW 264.7 cells were plated and treated with inhibitors (imatinib [Arg/Abl

inhibitor], PD0325901 [MEK1/2 inhibitor], SCH772984 [ERK1/2 inhibitor], trametinib

[MEK1/2 inhibitor]) (Selleck Chemicals, Houston, TX), PMA [ERK1/2 activator] or

DMSO, where appropriate. Metacyclic promastigotes were incubated in fresh mouse

serum for C3bi opsonization, and amastigotes were coated with anti-P8-

proteoglycolipid complex (monoclonal antibody IgG1) [18]. Next, samples with RAW

264.7 cells or BMDMs were incubated with C3bi-opsonized promastigotes at 10

parasites to 1 Mϕ or with IgG-opsonized amastigotes at 10 parasites to 1 Mϕ for 2

hours (h), unless otherwise indicated, and fixed with 3% formaldehyde for 20

minutes (m). Mouse anti-gp46 antibody was allowed to bind to external

promastigotes, whereas mouse anti-P8 antibody was incubated with external

amastigotes, as described previously [16, 17]. Samples were then incubated with

Alexa-Fluor-568-conjugated donkey anti-mouse-IgG secondary antibody. Any

internalized parasite did not receive any antibody tagging and fluoresces green.

Conversely, external parasites are visualized in both the red and green channels

(orange in a merged image). All antibodies are listed in **Supplemental Table 3**.

Thus, this three-color immunofluorescence assay [18] allowed distinguishing of

internal (green) and external *L. amazonensis* promastigotes or amastigotes (green +

red).

For quantification, samples were imaged with a BioTek Cytation 5 (Agilent, Santa

Clara, CA) confocal imaging reader using a 40x objective lens with automated

analysis performed [16]. At least 16 randomly selected fields were automatically

visualized per well with each field assessing the number of Mϕs and amastigotes or

promastigotes (internalized versus external). The absolute phagocytic index (the

number of internalized amastigotes or promastigotes per 100 Mϕs) and adhesion

index (the total number of amastigotes or promastigotes per 100 Mϕs) was

calculated for each condition. The absolute phagocytic index or adhesion index for

DMSO-treated conditions (control) was taken as the maximum value (100%) for

each experiment, and the phagocytic index for imatinib, SCH772984, trametinib, or

PMA-treated samples was determined and normalized relative to that of the control.

Means ± SD were calculated. Each experiment was performed with at least 3

biological repeats and contained 3 technical replicates. A one-sample Student's *t*-

test or two-way ANOVA was used to determine statistical significance, where

appropriate. The representative images shown were visualized using a Biotek

Cytation 5 imaging multimode reader. Parasites in representative images were

selected from files and linearly processed in Adobe Photoshop CS6 (version 13.0.6).

Where applicable, permeabilized samples were labeled with far-red phalloidin

(A22287, Thermo Fisher Scientific) at 1:100. For phagocytic cup analysis and

representative images, data were collected on a Zeiss LSM 880 inverted confocal

Airyscan microscope at 40X (analysis) or 63X (representative images). Images were

created via Image J (1.52a, http://imagej.nih.gov/ij) and processed in Adobe

Photoshop 13.0.6. [38].

2.5 Phagocytosis of beads. Experiments were adapted and modified as described

previously [16, 17]. RAW 264.7 cells were plated at 70% confluence and incubated

overnight in serum-free medium. Except where indicated, samples were pre-

incubated in medium containing 3.3 µM imatinib, 1 µM SCH772984, 1 µM trametinib,

250 nM PMA or 0.1% DMSO for 1-2 h. Next, latex yellow–green 2 µm beads

(Sigma) were coated with human IgM (Sigma, Cat. #: I-8260) and incubated in rabbit

anti-IgM (Sigma, Cat. #: 270A) for IgG opsonization (confirmed as described

previously [17]). Mφs were incubated with 10–15 beads/ Mφ for 2 h, unless otherwise

stated, at 37°C. They were fixed with 3% formaldehyde for 20 m, blocked using 2%

BSA in PBS (no permeabilization), and then incubated with rabbit anti-human IgM

and Hoechst 33258 dye (Sigma) to visualize nuclei. Next, samples were incubated

with an Alexa-Fluor-568-conjugated goat anti-rabbit-IgG secondary antibody

(Invitrogen, Cat.#: A11034). All antibodies are listed in **Supplemental Table 3**. This

multi-color immunofluorescence assay [39] allowed distinguishing of internalized

(green) and external beads (green + red), similar to described above for internal vs

external *L. amazonensis*. Quantification was done as above with a BioTek Cytation 5

confocal imaging reader using a 40x objective lens with automated analysis

performed by the Cytation 5.

2.6 Western Blot. To analyze protein/phosphoprotein expression, western blotting was performed. Briefly, RAW 264.7 cells or BMDMs from *WT, Syk$^{flox/flox}$ Cre$^+$*, or *Arg$^{-/-}$Abl$^{flox/flox}$ LysM-Cre$^+$* mice were incubated overnight in serum-free media (RAW 264.7 cells) or L929-free media (BMDM). Experiments were performed at ~70% confluence. Where appropriate, Mϕ were pre-incubated in imatinib, entospletinib, PD0325901, selumetinib, SCH772984, trametinib, PMA or DMSO. IgG-opsonized amastigotes or IgG-opsonized beads were added to adherent starved Mϕs for 30 m, unless otherwise indicated [40], and cells lysed with RIPA extraction buffer and halt protease and phosphatase inhibitor (Thermo Scientific, Cat. #: 78440, Richardson, TX). Protein purity and concentration were addressed using Coomassie Blue staining and BCA assay, respectively, following the manufacturers' protocols. Total proteins were resolved by polyacrylamide gel electrophoresis (12% gels) and transferred to PVDF membranes (BioRad). The membranes were blocked in 5% non-fat dry milk in Tris-buffered saline (TBS) (20 mM Tris at pH 7.6, 150 mM NaCl). The membrane was incubated overnight at 4°C with the primary antibody in 5% BSA and TBS-T. Next, horseradish peroxidase (HRP)-conjugated anti-mouse secondary antibody (#7076, Cell Signaling Technology) was added at 1:5000 in 5% non-fat dry milk in TBS-T for 1 h. All antibodies are provided in **Supplemental Table 3**. Membranes were incubated in SuperSignal West Pico Chemiluminescent Substrate (Thermo Scientific) for 5 m and visualized by using a phosphorimager (ImageQuant LAS 4000, GE Healthcare). For analysis, relative amounts of phosphorylated proteins were compared to total proteins with ImageJ analysis software (https://imagej.nih.gov/ij); membranes were stripped and reprobed). Each

experiment had at least 3 biological replicates. Two-way ANOVA was used for

statistical tests.

2.7 *Leishmania* animal model. For the mouse footpad model of cutaneous

leishmaniasis, 10 female C57BL/6 mice per group were infected between 6 and 8

weeks (w) of age as previously described [16, 17]. Briefly, 1×10^6 metacyclic

promastigotes were suspended in PBS and injected subcutaneously in the dorsal

side of the right hind foot. For the experiments in Fig 6A-C, mice were provided 2

mg/kg body weight of trametinib or drug vehicle (4% DMSO and 1% corn oil) by oral

gavage daily starting a w before infection and continuing until the mice were

euthanized. Two independent experiments were performed. For the experiments in

Fig 6D-F, mice were provided 2 mg/kg body weight of trametinib or drug vehicle by

oral gavage daily starting 6 w after infection and continuing until the mice were

euthanized. Two independent experiments were performed. Lesion growth was

documented with calipers every other w by an investigator who was blinded to

condition. The ratio of the infected: uninfected foot size was obtained [41]. ANOVA

was used to compare diluent and trametinib-treated mouse groups to swelling at

time 0. The number of parasites in lesions was determined by limiting dilution upon

termination of the experiment between w 14 to 16 [42].

Lymph nodes from infected diluent versus trametinib-treated mice were obtained for

cytokine and chemokine profiling [42, 43]. Briefly, lymph node cells were incubated

in RPMI 1640 (Invitrogen) plus 10% heat-inactivated FBS, 5×10^{-5} M 2-

mercaptoethanol (Sigma), and 1% penicillin-streptomycin. Cells were stimulated with

promastigote lysates (equivalent to 1×10^6 or 1×10^5 parasites per sample, as

indicated) or concanavalin A (ConA) (5 µg/ml; Sigma). Supernatants were harvested

at 72 h and the levels of the cytokines and chemokines IL-4, IL-10, IL-12, IL-13, IL-

17, IFN-γ, IL-1β, IL-1α, IP-10, MIP-1α (CCL3), MIP-2 (CXCL2), CXCL9, RANTES

(CCL5), and MIP-1β (CCL4) were profiled using the custom multiplex Luminex

Platform (R&D Systems, Inc, Minneapolis, MN) through the UT Southwestern

Microarray Core as described previously [44, 45]. Background cytokine levels were

calculated using unstimulated supernatants.

3. RESULTS

3.1 MAPK/ERK signaling in Mϕ facilitates *Leishmania* uptake

To directly test whether the MAPK/ERK pathway mediates *Leishmania* uptake,

we first explored the effect of the MEK1/2 inhibitor trametinib on the uptake of *L.*

amazonensis. We can grow both promastigotes and amastigotes of this species

in tissue culture without host cells (*i.e*., axenic culture), which facilitates our

studies of *Leishmania* uptake by Mϕs. Trametinib is a highly selective and potent

MEK1/2 inhibitor, with an IC_{50} for the purified kinase ranging from 0.92 nM to 3.4

nM in cell-free assays [46, 47]. Trametinib has no effect on the kinase activity of

c-Raf, B-Raf, or ERK1/2. We pretreated RAW 264.7 cells, a Mϕ–like cell line, for

2 hours (h) with trametinib or imatinib, an Abl/Arg inhibitor with an IC_{50} of 100-500

nM in kinase assays [35], as a positive control. We then added *Leishmania* to

samples to allow uptake and processed for immunofluorescence microscopy.

Visually, trametinib decreased the uptake of IgG-opsonized amastigotes in RAW

264.7 cells, comparable to the reductions seen after treatment with imatinib (**Fig.**

1A). Defects in internalization were quantified by calculating the phagocytic index

(PI), or number of internalized parasites per 100 Mϕ, which was lower in

trametinib-treated samples (**Fig. 1B**). No differences were seen in the total

(external + internal) number of *Leishmania* per 100 RAW 264.7 cells, which is

termed an adhesion index, among these conditions (**Fig. 1C**). We performed a

titration of trametinib's effects on amastigote uptake, and the maximal reduction

in PI was shown at 1 μM trametinib (**Fig. 1D, E).** We also determined that over

72 h, the CC_{50} (cytotoxic concentration required to kill 50% of viable cells) for

RAW 264.7 cells was greater than our maximum concentration of 2.5 μM.

Internalization defects remained even after prolonged incubation in trametinib (up

to 240 minutes, m) and were not due to reduced adhesion, as amastigotes

continued to be bound to trametinib-treated RAW 264.7 cells at levels

indistinguishable from controls (**Fig. 1F, G**). Treating Mϕs with other MAPK/ERK

pathway inhibitors [48] such as selumetinib and PD0325901 (MEK1/2 inhibitors),

as well as SCH772984 (an ERK1/2 inhibitor), also significantly reduced

amastigote uptake without affecting the total number of bound parasites

(**Supplemental Fig. 1A, B**). In addition, we found that treating RAW 264.7 cells

with the MEK1/2 inhibitor PD0325901 [49] decreased the internalization of

promastigotes to a degree that was comparable to imatinib (**Supplemental Fig.**

1C, D). Next, we characterized growth curves and found that trametinib showed

no impact on cell viability and motility of parasites in culture (**Fig. 1H**). Together,

these similar results using multiple MAPK/ERK inhibitors establish that targeting

MEK1/2 and ERK1/2 significantly attenuates internalization of both *L.*

amazonensis life cycle stages. In addition, as previously published for

MAPK/ERK inhibitors, amastigotes within Mϕs had a ~50% reduction in survival

when cultures were treated with trametinib for 72 h (**Supplemental Fig. 1E**).

3.2 cRaf, MEK1/2, and ERK1/2 are activated during amastigote uptake

Having established that MAPK/ERK are important for *Leishmania* uptake, we

assessed whether kinases in this pathway were activated in host cells during

amastigote internalization. Raf-1 (cRaf), MEK1/2, and ERK1/2 are key members of

the MAPK/ERK pathway [25, 50]. We thus measured cRaf, MEK1/2, and ERK1/2

phosphorylation in RAW 264.7 cells to examine the relative phosphorylation level of

each kinase during internalization of amastigotes. Essentially no phosphorylated

ERK or ERK was detected in amastigote samples (**Supplemental Fig. 2A**).

However, amastigotes potently induced ERK1/2 phosphorylation in Mϕs, with

phosphorylation levels increasing by 3 and 4.5-fold at 10 and 30 m post-infection,

respectively (**Fig. 2A**). Similarly, MEK1/2 (**Fig. 2B**) and cRaf (**Fig. 2C**) were

activated by amastigotes that were entering host cells**.** These data indicate that

amastigotes rapidly activate MAPK/ERK in a sustained fashion during their uptake

by Mϕs.

3.3 PMA rescues MAPK/ERK related uptake defects

PMA is known to activate ERK1/2 through PKC, with an IC_{50} of 11.7 nM against

purified kinase [26]. Consistent with this, although PKC activation was seen during

the uptake of amastigotes, PMA treatment increased PKC phosphorylation in RAW

264.7 cells to 8-fold higher levels (**Fig. 3A, B**). Importantly, 10 µM PMA did not affect

growth of amastigotes in culture (**Fig. 3C**), indicating that it does not act directly on

Leishmania. We next investigated whether PMA affected amastigote internalization.

To perform these experiments, and to confirm our prior results from a cell line using

primary cells, we isolated bone marrow derived-Mϕs (BMDM) from C57BL/6 mice.

PMA-treated BMDM were able to internalize IgG-coated amastigotes at higher levels

than DMSO-treated BMDM in a dose-dependent manner, reaching maximal

internalization at 2.5 µM PMA (**Fig. 3D-F**). Furthermore, PMA treatment rescued

trametinib inhibition of *Leishmania* uptake in BMDM to PI that were indistinguishable

from those of DMSO-treated controls (**Fig. 3G, H**). Collectively, our data indicate that

ERK1/2 kinase activity is necessary for maximal *Leishmania* uptake.

3.4 ERK1/2 is activated by Arg/Abl and SYK during amastigote uptake

Next, we investigated the mechanism by which *L. amazonensis* amastigotes induced

ERK1/2 phosphorylation during their uptake. Previous research has shown that Arg

facilitates *Leishmania* amastigote uptake downstream of SFK [16]. In addition,

inhibition of SYK, a tyrosine kinase implicated in FcγR immunoglobulin-mediated

signaling [18, 33, 51, 52] reduces ERK1/2 activation in a dose-dependent manner in

other systems [18, 33]. To test for roles for these kinases, small molecule inhibitors

(imatinib, entospletinib [IC_{50} for activity against purified SYK = 7.7 nM] [18],

SCH772984, trametinib), and PMA were added to RAW 264.7 cells before infection

with amastigotes. Inhibition of Arg and SYK with imatinib and entospletinib,

respectively, significantly reduced ERK1/2 phosphorylation, to a similar degree to

the MAPK/ERK inhibitors SCH772984 and trametinib (**Fig. 4A, B**). As expected,

PMA, the positive control, significantly increased ERK1/2 phosphorylation (**Fig. 4A,**

B). The inhibitors and activators followed similar trends for promastigote-induced

ERK activity (**Supplemental Fig. 2B**). Thus, we could modulate *Leishmania*-induced

ERK activity by both inhibitors and activators *in vitro*.

To further characterize the role of SYK and Arg in *Leishmania*-induced ERK

phosphorylation, we next isolated BMDM from mice lacking Abl family kinases

(*Arg*$^{-/-}$*Abl*$^{flox/flox}$*LysM Cre*$^+$ [hereafter termed *Arg*$^{-/-}$*Abl*$^{-/-}$ when discussing isolated

BMDM]), SYK (*SYK*$^{flox/flox}$ *Cre*$^+$, abbreviated *SYK*$^{-/-}$ when referring to isolated BMDM)

or wild type (*WT*) littermates. We have previously shown that amastigote uptake is

dependent upon Arg but not Abl [15-16]. Our immunoblots showed *Arg*$^{-/-}$*Abl*$^{-/-}$ and

SYK$^{-/-}$ BMDM nearly completely lack Arg or SYK, respectively, while ERK1/2

expression was maintained (**Supplemental Fig. 2C**). Consistent with our chemical

inhibition experiments, *SYK*$^{-/-}$ and *Arg*$^{-/-}$*Abl*$^{-/-}$ BMDM limited amastigote-mediated

ERK1/2 phosphorylation, demonstrating that SYK and Arg/Abl are necessary for

stimulating ERK1/2 phosphorylation following engagement of the FcγR (**Fig. 4C, D**).

At low doses (250 nM), PMA rescued ERK1/2 activity in *Arg*$^{-/-}$*Abl*$^{-/-}$ and *SYK*$^{-/-}$

BMDM. This suggested 1) amastigotes operate through a distinct pathway to

activate ERK1/2 compared to PMA, which activates MAPK/ERK through PKC [26]

and 2) during FcγR-mediated amastigote infection, Arg and SYK family kinases are

upstream of MAPK/ERK and are necessary to activate the MAPK/ERK pathway.

To directly test whether Abl family kinases are critical for ERK1/2-mediated

amastigote uptake, we treated *WT* and *Arg*$^{-/-}$*Abl*$^{-/-}$ BMDM with DMSO or PMA prior

to infection with IgG-opsonized amastigotes. We found that *Arg*$^{-/-}$ *Abl*$^{-/-}$ BMDM

displayed defects in IgG-mediated amastigote uptake, while PMA treatment partially

rescued these defects (**Fig. 4E-G**), returning uptake to 66.4 ± 16.6% (mean ± SD)

compared to 29.3 ± 0.9% with DMSO. Similarly, SYK was necessary for efficient

amastigote uptake, and the decreased PI of *SYK*$^{-/-}$ BMDM was partially rescued by

PMA treatment (**Supplemental Fig. 2D, E)**. Taken together, these results, in

combination with prior data, are consistent with a signaling pathway in which Arg/Abl

and SYK act upstream of and activate the ERK1/2 pathway (which is at least partly

independent from PKC) to facilitate amastigote uptake [26].

3.5 MAPK/ERK inhibition minimally impacts phagocytosis of opsonized beads

$Arg^{-/-}$ BMDM have been shown to have impaired FcR-mediated phagocytosis

overall, as demonstrated using IgG-coated (opsonized) beads [17]. Thus, we

hypothesized that the Arg-SYK-ERK pathway may be necessary for the general

process of phagocytosis, in addition to amastigote uptake. To delineate the specific

contributions of MAPK/ERK to phagocytosis, we examined bead uptake compared

to amastigote uptake by RAW 264.7 cells. Surprisingly, phagocytosis of IgG-

opsonized beads by RAW 264.7 cells was minimally affected by trametinib (**Fig. 5A,**

B) or SCH772984 (**Supplemental Fig. 3),** both of which resulted in only a ~30%

reduction in uptake. Conversely, the uptake of IgG-coated amastigotes (**Fig. 1)** was

reduced by ~70% with trametinib treatment. However, when RAW 264.7 cells were

treated with imatinib, similar decreases in phagocytosis of IgG-opsonized beads

versus amastigotes were observed. These results suggest that Arg/Abl play a more

generalized role in phagocytosis, as we have demonstrated previously [17].

Collectively, our data suggest that ERK1/2 activation is important for *Leishmania*

uptake specifically but may play a minor role in standard phagocytic mechanisms.

We also tested whether MAPK/ERK activity was induced following addition of beads

compared to amastigotes. Beads induced ERK1/2 phosphorylation, but to levels that

were lower than in seen during amastigote uptake (**Fig. 5C, D**). This phosphorylation

occurred in a time-dependent fashion, such that activation reached maximal levels

between 5 and 10 m, and ERK1/2 phosphorylation was not detectable after 30 m

(**Fig. 5C, D**). Similarly, beads activated upstream kinases MEK1/2 (**Fig. 5E, F)** and

cRaf (**Fig. 5G, H**); however, their phosphorylation was not sustained. Thus, in

contrast to the sustained activation of MAPK/ERK that we saw during amastigote

uptake, beads had only a transient effect on ERK1/2 activation.

3.6 Trametinib increases the intensity of actin labeled in phagocytic cups.

Next, we explored the mechanisms by which trametinib affected *Leishmania*

amastigote internalization. We first characterized trametinib's effects on phagocytic

cup formation by performing an internalization assay with amastigotes as described

above, with the addition of fluorescent phalloidin to label phagocytic cups. We found

that phalloidin labeling around internalizing parasites was significantly brighter in

trametinib-treated Mϕs on average than DMSO-treated Mϕs, suggesting direct

effects on actin polymerization during the formation of actin-rich phagocytic cups

(**Supplemental Fig. 4A-C**). We next calculated the percentages of Mϕs infected by

amastigotes and the average numbers of amastigotes per infected Mϕ. We found

that both were affected by trametinib at approximately the values expected from the

decrease in phagocytic indexes (**Supplemental Fig. 4D-E**).

3.7 Trametinib decreases lesion size and parasite burden in a murine model of

cutaneous leishmaniasis. To test whether the MAPK/ERK pathway is necessary

for *Leishmania* pathogenesis *in vivo*, we employed a C57BL/6 mouse model of

cutaneous leishmaniasis using *L. amazonensis.* Footpad swelling (*i.e.*, lesion size)

was measured in infected mice treated with 2 mg/kg of body weight/day of oral

trametinib or drug vehicle (control) starting 7 days (d) before inoculation and

continuing for the experimental duration of 14-16 weeks (w) [16, 17] (**Fig. 6A**).

Compared to controls, trametinib-treated mice developed smaller lesions (**Fig. 6B**),

which contained fewer parasites in lesions (**Fig. 6C**). The decreased number of

parasites found within lesions may help to explain these changes in lesion size.

Overall, in leishmaniasis, as a simplified paradigm, a Th1 response is thought to

be beneficial to the host, while a Th2 response is typically considered to be

deleterious. To determine whether the immunological effects of MEK1/2 inhibition

were associated with the healing seen in trametinib-treated mice, we isolated

draining lymph nodes from infected DMSO and trametinib-treated mice. We then

profiled cytokine and chemokine secretion after stimulation with high (1×10^6) or

low (1×10^5) doses of *L. amazonensis* lysates compared to unstimulated. A mild

shift in the overall ratio of Th1 to Th2 cytokines (Th1/Th2) towards a Th2

phenotype in trametinib-treated mice was seen (**Table 1**). Notably, lymph nodes

extracted from trametinib-treated mice released lower quantities of nearly all

cytokines in response to *L. amazonensis* than those isolated from diluent-treated

animals (**Table 2**). Trametinib-treated mice also demonstrated decreased

cytokine production in response to concanavalin A (Con A) stimulation. In

addition, we assessed chemokine secretion following MEK1/2 inhibition. The

production of several chemokines implicated in *Leishmania* pathogenesis,

including MIP-1α (CCL3) and MCP-1 (CCL2), were significantly decreased with

trametinib (**Table 2**), consistent with the decrease in cytokines/chemokines

overall.

We further tested whether MEK1/2 inhibition with trametinib reduces pathogenesis

after the development of symptoms by infecting mice as above. However, we started

diluent vs trametinib at 6 w after inoculation (*i.e.*, once lesions were clearly visible)

and continued therapy for the remaining experimental duration (**Fig. 6D**). Even with

this delayed treatment, trametinib-treated mice reproducibly had smaller lesions

(**Fig. 6E**). They also had significantly lower parasite burdens than those of controls

(**Fig. 6F**), which at least in part may explain these results. We also saw similar

immunological responses in mice treated with trametinib initiated partway through

the experiment as we did for the mice treated prior to *Leishmania* infection

(**Supplemental Table 1, Supplemental Table 2**) [53, 54]. Overall, our findings

demonstrate that trametinib administration limits the manifestations of leishmaniasis

in a mouse model, even if it is started after lesions have developed.

4. DISCUSSION

Here, we have shown that efficient *Leishmania* uptake by Mφs and infection in mice requires MAPK/ERK. While previous studies have shown that ERK1/2 activation post-infection by *Leishmania* alters various host signaling pathways, leading to increased intracellular survival and replication of these parasites [36, 55], our results establish an earlier role for host ERK1/2 activity during the *Leishmania* life cycle, and show that ERK1/2 is necessary for efficient *Leishmania* uptake into Mφs. Furthermore, our results delineate new activators of ERK signaling, and suggest that MAPK/ERK affects actin polymerization. In combination with prior data, we propose a model whereby, following engagement of the FcγR, host Arg/Abl and SYK stimulate Raf, MEK, and ERK1/2 phosphorylation, leading to amastigote uptake (**Fig. 6G**). Moreover, *in vivo* inhibition of MEK1/2 by trametinib in a mouse model of cutaneous leishmaniasis significantly limits lesion development and parasite burden, even when trametinib is started after these lesions have developed. Our results highlight a critical role for MAPK/ERK signaling in *Leishmania* uptake and further previously published studies of its role in the pathogenesis of leishmaniasis.

Our data indicate that Arg/Abl and SYK are required for amastigote-mediated ERK1/2 phosphorylation following FcγR activation, which has not been demonstrated previously. Our prior studies of host signaling required for *Leishmania* uptake revealed that amastigotes bind directly to FcγR [15] and activate SFK-Arg pathways [16] to induce uptake. We conclude from our combined results that an FcγR-SFK-Arg-SYK signaling pathway functions upstream of and activates

MAPK/ERK family kinases, which in turn affects the amount of polymerized actin in

phagocytic cups. We previously published similar effects on actin polymerization

using the SYK inhibitor entospletinib during *Leishmania* uptake [18]. Interestingly,

pharmacological inhibition of MEK1/2 and ERK1/2 in RAW 264.7 cells significantly

reduce, but does not completely abolish, *Leishmania* uptake. One explanation for

these results is that kinase inhibition via these compounds may be incomplete.

Another is that *Leishmania* uptake may be mediated via additional host signaling

pathways. Supporting the latter hypothesis, we find that low or sub-EC$_{50}$ doses of

PMA, which activates ERK1/2 through an SFK-Arg-SYK-independent manner,

rescues trametinib inhibition of *Leishmania* amastigote uptake in RAW 264.7 cells,

as well as in *SYK*$^{-/-}$ and *Arg*$^{-/-}$*Abl*$^{-/-}$ BMDM. PMA is a known activator of PKC [26,

56-60]. However, we find that PKC activity is only slightly increased during

amastigote infection of WT/DMSO-treated Mφs, suggesting that PKC may play only

a minor role during typical amastigote uptake. Ongoing work in the laboratory is

directed at uncovering additional novel uptake mechanisms for *Leishmania*. In

addition, although drug combinations typically are not used to treat leishmaniasis,

they are the standard of care for many other infections, such as HIV or tuberculosis.

Combination therapy with multiple kinase inhibitors to inhibit several uptake

modulators at once may provide additional benefit for the treatment of leishmaniasis

over single host cell kinase inhibitors alone.

Our data using MEK/ERK inhibitors indicate that there are differences in MAPK/ERK

signaling within Mφs depending on whether *Leishmania* amastigotes or IgG-coated

beads are undergoing internalization. These results differ from most available

literature that concludes that *Leishmania* primarily hijacks ongoing phagocytic

pathways to facilitate its uptake. We instead suggest that there may be alternative,

perhaps partly redundant, pathways that may support the uptake of *Leishmania* over

the phagocytosis of beads, and vice versa. Our future experiments will continue to

compare and contrast the signaling pathways required for these different uptake

mechanisms between opsonized beads and *Leishmania* parasites.

Our experiments using a mouse model for *Leishmania* infection, in combination with

prior studies [33], show that MAPK/ERK are required for maximal *Leishmania*

pathogenesis during cutaneous leishmaniasis. One highly plausible explanation for

the decreased lesion size and parasite burden that we see in trametinib-treated mice

is our previously documented inhibition of *Leishmania* uptake, in combination with

previous studies suggesting MAPK/ERK signaling affects parasite survival in Mϕs

[26-33]. Consistent with this hypothesis, fewer parasites are found in lesions from

trametinib-treated mice than in control mice. Alternately, lesion size variations after

L. amazonensis infection might result from known trametinib limitation of host

inflammatory and immunological responses. Many pathogens hinder immune

responses by targeting host intracellular signaling pathways, including MAPK

pathways [59, 61]; and MAPKs have been shown to play roles in the immune

response to leishmaniasis [26-33]. The murine immune response to leishmaniasis is

complicated and varies depending on the species of *Leishmania* and the mouse

strain used. As a simplified paradigm, Th1 responses are generally protective, but

Th2 responses are harmful to the host [28, 31, 32]. We find that all cytokines

assessed following *Leishmania* antigen stimulation in lymph nodes isolated from

infected trametinib-treated mice are reduced from the levels seen in diluent-treated

mice, similar to our prior results using bosutinib [16]. One possible explanation is that

fewer cytokines are generated because the lesions contain fewer parasites. Another

is that trametinib could instead dampen pathways important for maintaining an active

inflammatory response to the ongoing host response to infection. Supporting this

second possibility, cytokines measured after Con A stimulation (an antigen-

independent mitogen used as an alternative T cell stimulus) are significantly reduced

by MEK1/2 inhibition. Overall, the moderate shift in the total ratio of Th1 to Th2

cytokines (Th1/Th2) toward a Th2 phenotype that we see theoretically should

exacerbate disease rather than alleviate it. Reduced levels of proinflammatory

chemokines like MIP-1 (CCL3) and MCP-1 (CCL2), on the other hand, may be

responsible for reducing lesion size by restricting the number of immune cells,

especially phagocytic cells, which reach the site of infection. We do note a

surprisingly strong IL-10 response in the trametinib-treated mice, considering that

ERK activation has been shown to be required for IL-10 production by *Leishmania*-

infected Mɸs [62]. However, others have found that trametinib treatment increases

IL-10 production in mouse tissues [63]. Further work is needed to delineate the

regulation of immune responses to *Leishmania* infection and the resulting effects on

lesion worsening versus healing.

Finally, and of particular interest, we have demonstrated that there are reductions

in lesion size and parasite burden if trametinib is used in a *L. amazonensis*

mouse model of cutaneous leishmaniasis, even if it is started after foot swelling

has developed. Compared to our previous results, the data shown here strongly

suggest that trametinib is more effective for reducing lesion size than imatinib

[17] or bosutinib [16], which target signaling molecules in the same pathway.

Trametinib is used clinically to treat several cancers, making it feasible that this

drug could be repurposed to treat an infectious disease, and facilitating potential

clinical trials for its use against leishmaniasis. Our data imply that inhibiting a

signaling pathway downstream of SFK-Arg-SYK might improve results over

inhibiting kinases upstream in the pathway. Our results also give credence to our

overarching hypothesis that targeting host cell pathways, particularly ones that

are required for multiple aspects of the parasite life cycle and pathogenesis, may

permit treatment of an infectious disease such as leishmaniasis.

5. CONCLUSIONS

In summary, we have demonstrated that Arg-SYK signaling through MAPK/ERK

facilitates *Leishmania* uptake by Mϕ. Furthermore, ERK activity helps govern the

subsequent pathogenesis and disease progression of leishmaniasis in the mouse

model. Due to the relative specificity of newer-generation MAPK/ERK small

molecule inhibitors like trametinib, it is possible to selectively target MEK1/2 and

ERK1/2 for potential therapeutic intervention. Our findings demonstrate the

significance of MAPK/ERK signaling during *Leishmania* entry into phagocytic cells.

Our future studies will investigate the use of host-cell-active drugs, potentially in

conjunction with other kinase inhibitors or existing antiparasitics, to improve efficacy,

decrease toxicity, and limit the likelihood of developing resistance. Although

trametinib is not completely without side effects, this combination strategy could be

used to mitigate any risk of toxicity arising due to targeting essential signaling

pathways. Furthermore, our results support the novel possibility that medications

that target conserved host cell processes rather than infectious pathogens might be

employed to treat a wide spectrum of intracellular pathogens.

6. ACKNOWLEDGEMENTS

6.1 Overview. We thank Drs. Neal M. Alto, Melanie H. Cobb, Margaret A. Phillips, James J. Collins, and Michael L. Reese (all at UT Southwestern) for helpful discussions and feedback. Members of the UT Southwestern Departments of Pediatrics, Biochemistry, Microbiology, and Pharmacology provided valuable feedback and permitted common equipment use. Francis Khuong and Catherine Trice provided expert technical assistance and helped to edit this manuscript. Dr. Diane McMahon-Pratt (Yale University) generously donated antileishmanial antibodies. Dr. Anthony J. Koleske (Yale University) provided the *Arg$^{-/-}$Abl$^{flox/flox}$* mice. The UT Southwestern Microarray Core facilitated cytokine/chemokine profiling, the High Throughput Screening Core assisted with data visualization and quantification, and the Live Cell Imaging Core helped us with collecting confocal microscopy images.

6.2 Funding. This work was supported by a Children's Clinical Research Advisory Committee (CCRAC) Junior Investigator Award, a CCRAC Early Investigator Award, the National Institutes of Health (K08 AI103036 and R01 AI146349), Welch Foundation Grant I-2048, and the UT Southwestern Department of Pediatrics (to D.M.W.). U.B. was supported by a National Institutes of Health Supplement to Promote Diversity in Health-Related Research (R01 AI146349- S1) and Medical Scientist Training (MD/PhD) Grant NIH T32GM008014. K.F. was supported by a UT Southwestern-Amgen Summer Research Fellowship.

694   **6.3 Competing interests.** The authors have no competing financial interests.

696   **6.4 Data and materials availability.** All data are available within the main text of the

697   manuscript or the supplemental materials. Any materials generated for this

698   manuscript are freely available upon request.

7. REFERENCES AND NOTES

1.   Gossage SM, Rogers ME, Bates PA. Two separate growth phases during the

development of Leishmania in sand flies: implications for understanding the life cycle. Int

J Parasitol. 2003;33(10):1027-34. doi: 10.1016/s0020-7519(03)00142-5. PubMed PMID:

13129524; PubMed Central PMCID: PMCPMC2839921.

2.   Markle WH, Makhoul K. Cutaneous leishmaniasis: recognition and treatment. Am

Fam Physician. 2004;69(6):1455-60. PubMed PMID: 15053410.

3.   McCall LI, Zhang WW, Matlashewski G. Determinants for the development of

visceral   leishmaniasis   disease.   PLoS   Pathog.   2013;9(1):e1003053.   doi:

10.1371/journal.ppat.1003053. PubMed PMID: 23300451; PubMed Central PMCID:

PMCPMC3536654.

4.   Georgiadou SP, Makaritsis KP, Dalekos GN. Leishmaniasis revisited: Current

aspects on epidemiology, diagnosis and treatment. J Transl Int Med. 2015;3(2):43-50.

doi: 10.1515/jtim-2015-0002. PubMed PMID: 27847886; PubMed Central PMCID:

PMCPMC4936444.

5.   Steverding D. The history of leishmaniasis. Parasit Vectors. 2017;10(1):82. Epub

2017/02/17. doi: 10.1186/s13071-017-2028-5. PubMed PMID: 28202044; PubMed

Central PMCID: PMCPMC5312593.

6.   Gonzalez C, Wang O, Strutz SE, Gonzalez-Salazar C, Sanchez-Cordero V, Sarkar

S. Climate change and risk of leishmaniasis in North America: predictions from ecological

niche models of vector and reservoir species. PLoS Negl Trop Dis. 2010;4(1):e585. Epub

2010/01/26. doi: 10.1371/journal.pntd.0000585. PubMed PMID: 20098495; PubMed

Central PMCID: PMCPMC2799657.

7.   McIlwee BE, Weis SE, Hosler GA. Incidence of endemic human cutaneous leishmaniasis in the United States. JAMA Dermatol. 2018;154(9):1032-9. doi: 10.1001/jamadermatol.2018.2133. PubMed PMID: 30046836; PubMed Central PMCID: PMCPMC6143046.

8.   Croft SL, Sundar S, Fairlamb AH. Drug resistance in leishmaniasis. Clin Microbiol Rev. 2006;19(1):111-26. Epub 2006/01/19. doi: 10.1128/CMR.19.1.111-126.2006. PubMed PMID: 16418526; PubMed Central PMCID: PMCPMC1360270.

9.   Elmahallawy EK, Agil A. Treatment of leishmaniasis: a review and assessment of recent research. Curr Pharm Des. 2015;21(17):2259-75. Epub 2014/12/30. doi: 10.2174/1381612821666141231163053. PubMed PMID: 25543123.

10.   Mann S, Frasca K, Scherrer S, Henao-Martinez AF, Newman S, Ramanan P, et al. A review of leishmaniasis: current knowledge and future directions. Curr Trop Med Rep. 2021;8(2):121-32. Epub 2021/03/23. doi: 10.1007/s40475-021-00232-7. PubMed PMID: 33747716; PubMed Central PMCID: PMCPMC7966913.

11.   Ponte-Sucre A, Gamarro F, Dujardin JC, Barrett MP, Lopez-Velez R, Garcia-Hernandez R, et al. Drug resistance and treatment failure in leishmaniasis: A 21st century challenge. PLoS Negl Trop Dis. 2017;11(12):e0006052. Epub 2017/12/15. doi: 10.1371/journal.pntd.0006052. PubMed PMID: 29240765; PubMed Central PMCID: PMCPMC5730103.

12.   Handman E, Bullen DV. Interaction of *Leishmania* with the host macrophage. Trends Parasitol. 2002;18(8):332-4. doi: 10.1016/s1471-4922(02)02352-8. PubMed PMID: 12377273.

13.   Alexander J, Russell DG. The interaction of Leishmania species with
macrophages. Adv Parasitol. 1992;31:175-254. Epub 1992/01/01. doi: 10.1016/s0065-
308x(08)60022-6. PubMed PMID: 1496927.

14.   Kima PE. The amastigote forms of Leishmania are experts at exploiting host cell
processes to establish infection and persist. Int J Parasitol. 2007;37(10):1087-96. doi:
10.1016/j.ijpara.2007.04.007. PubMed PMID: 17543969; PubMed Central PMCID:
PMCPMC2043126.

15.   Ueno N, Wilson ME. Receptor-mediated phagocytosis of Leishmania: implications
for   intracellular   survival.   Trends   Parasitol.   2012;28(8):335-44.   doi:
10.1016/j.pt.2012.05.002.   PubMed   PMID: 22726697;   PubMed   Central   PMCID:
PMCPMC3399048.

16.   Wetzel DM, Rhodes EL, Li S, McMahon-Pratt D, Koleske AJ. The Src kinases Hck,
Fgr and Lyn activate Arg to facilitate IgG-mediated phagocytosis and Leishmania
infection. J Cell Sci. 2016;129(16):3130-43. Epub 2016/07/01. doi: 10.1242/jcs.185595.
PubMed PMID: 27358479; PubMed Central PMCID: PMCPMC5004897.

17.   Wetzel DM, McMahon-Pratt D, Koleske AJ. The Abl and Arg kinases mediate
distinct modes of phagocytosis and are required for maximal *Leishmania* infection. Mol
Cell Biol. 2012;32(15):3176-86. Epub 2012/06/06. doi: 10.1128/MCB.00086-12. PubMed
PMID: 22665498; PubMed Central PMCID: PMCPMC3434515.

18.   Ullah I, Barrie U, Kernen RM, Mamula ET, Khuong FTH, Booshehri LM, et al. Src-
and Abl-family kinases activate spleen tyrosine kinase to maximize phagocytosis and
Leishmania   infection.   J   Cell   Sci.   2023;136(14).   Epub   2023/06/26.   doi:

10.1242/jcs.260809.   PubMed   PMID: 37357611;   PubMed   Central   PMCID:

PMCPMC10399977.

19.   Greuber EK, Pendergast AM. Abl family kinases regulate FcgammaR-mediated

phagocytosis in murine macrophages. J Immunol. 2012;189(11):5382-92. Epub

2012/10/27. doi: 10.4049/jimmunol.1200974. PubMed PMID: 23100514; PubMed Central

PMCID: PMCPMC3504141.

20.   Ortiz MA, Mikhailova T, Li X, Porter BA, Bah A, Kotula L. Src family kinases,

adaptor proteins and the actin cytoskeleton in epithelial-to-mesenchymal transition. Cell

Commun Signal. 2021;19(1):67. Epub 2021/07/02. doi: 10.1186/s12964-021-00750-x.

PubMed PMID: 34193161; PubMed Central PMCID: PMCPMC8247114.

21.   Cortez D, Reuther G, Pendergast AM. The Bcr-Abl tyrosine kinase activates

mitogenic signaling pathways and stimulates G1-to-S phase transition in hematopoietic

cells. Oncogene. 1997;15(19):2333-42. Epub 1997/12/11. doi: 10.1038/sj.onc.1201400.

PubMed PMID: 9393877.

22.   Jin A, Kurosu T, Tsuji K, Mizuchi D, Arai A, Fujita H, et al. BCR/ABL and IL-3

activate Rap1 to stimulate the B-Raf/MEK/Erk and Akt signaling pathways and to regulate

proliferation,   apoptosis,   and   adhesion.   Oncogene.   2006;25(31):4332-40.   Epub

2006/03/07. doi: 10.1038/sj.onc.1209459. PubMed PMID: 16518411.

23.   Yang S, Liu G. Targeting the Ras/Raf/MEK/ERK pathway in hepatocellular

carcinoma. Oncol Lett. 2017;13(3):1041-7. Epub 2017/04/30. doi: 10.3892/ol.2017.5557.

PubMed PMID: 28454211; PubMed Central PMCID: PMCPMC5403244.

24.   Guo YJ, Pan WW, Liu SB, Shen ZF, Xu Y, Hu LL. ERK/MAPK signalling pathway

and tumorigenesis. Exp Ther Med. 2020;19(3):1997-2007. Epub 2020/02/28. doi:

10.3892/etm.2020.8454. PubMed PMID: 32104259; PubMed Central PMCID:

PMCPMC7027163.

25.   Eblen ST. Extracellular-Regulated Kinases: signaling from Ras to ERK substrates

to control biological outcomes. Adv Cancer Res. 2018;138:99-142. Epub 2018/03/20. doi:

10.1016/bs.acr.2018.02.004. PubMed PMID: 29551131; PubMed Central PMCID:

PMCPMC6007982.

26.   Verin AD, Liu F, Bogatcheva N, Borbiev T, Hershenson MB, Wang P, et al. Role

of ras-dependent ERK activation in phorbol ester-induced endothelial cell barrier

dysfunction. Am J Physiol Lung Cell Mol Physiol. 2000;279(2):L360-70. Epub 2000/08/05.

doi: 10.1152/ajplung.2000.279.2.L360. PubMed PMID: 10926560.

27.   Wu X, Dayanand KK, Thylur RP, Norbury CC, Gowda DC. Small molecule-based

inhibition of MEK1/2 proteins dampens inflammatory responses to malaria, reduces

parasite load, and mitigates pathogenic outcomes. J Biol Chem. 2017;292(33):13615-34.

Epub 2017/07/07. doi: 10.1074/jbc.M116.770313. PubMed PMID: 28679535; PubMed

Central PMCID: PMCPMC5566520.

28.   Martinez PA, Petersen CA. Chronic infection by *Leishmania amazonensis*

mediated through MAPK ERK mechanisms. Immunol Res. 2014;59(1-3):153-65. Epub

2014/05/20. doi: 10.1007/s12026-014-8535-y. PubMed PMID: 24838145; PubMed

Central PMCID: PMCPMC4686340.

29.   Shadab M, Ali N. Evasion of host defence by *Leishmania donovani*: subversion of

signaling pathways. Mol Biol Int. 2011;2011:343961. Epub 2011/11/18. doi:

10.4061/2011/343961. PubMed PMID: 22091401; PubMed Central PMCID:

PMCPMC3199940.

30.   Olivier M, Gregory DJ, Forget G. Subversion mechanisms by which Leishmania

parasites can escape the host immune response: a signaling point of view. Clin Microbiol

Rev. 2005;18(2):293-305. Epub 2005/04/16. doi: 10.1128/CMR.18.2.293-305.2005.

PubMed PMID: 15831826; PubMed Central PMCID: PMCPMC1082797.

31.   Ghalib HW, Whittle JA, Kubin M, Hashim FA, el-Hassan AM, Grabstein KH, et al.

IL-12 enhances Th1-type responses in human *Leishmania donovani* infections. J

Immunol. 1995;154(9):4623-9. Epub 1995/05/01. PubMed PMID: 7722314.

32.   Bhardwaj S, Srivastava N, Sudan R, Saha B. Leishmania interferes with host cell

signaling to devise a survival strategy. J Biomed Biotechnol. 2010;2010:109189. Epub

2010/04/17. doi: 10.1155/2010/109189. PubMed PMID: 20396387; PubMed Central

PMCID: PMCPMC2852600.

33.   Yang Z, Mosser DM, Zhang X. Activation of the MAPK, ERK, following *Leishmania*

amazonensis infection of macrophages. J Immunol. 2007;178(2):1077-85. doi:

10.4049/jimmunol.178.2.1077. PubMed PMID: 17202371; PubMed Central PMCID:

PMCPMC2643020.

34.   Mukbel RM, Patten C, Jr., Gibson K, Ghosh M, Petersen C, Jones DE.

Macrophage killing of *Leishmania amazonensis* amastigotes requires both nitric oxide

and superoxide. Am J Trop Med Hyg. 2007;76(4):669-75. Epub 2007/04/12. PubMed

PMID: 17426168.

35.   Calegari-Silva TC, Pereira RM, De-Melo LD, Saraiva EM, Soares DC, Bellio M, et

al. NF-kappaB-mediated repression of iNOS expression in *Leishmania amazonensis*

macrophage infection. Immunol Lett. 2009;127(1):19-26. Epub 2009/08/29. doi:

10.1016/j.imlet.2009.08.009. PubMed PMID: 19712696.

36.   Boggiatto PM, Martinez PA, Pullikuth A, Jones DE, Bellaire B, Catling A, et al.

Targeted extracellular signal-regulated kinase activation mediated by *Leishmania*

amazonensis requires MP1 scaffold. Microbes Infect. 2014;16(4):328-36. doi:

10.1016/j.micinf.2013.12.006. PubMed PMID: 24463270; PubMed Central PMCID:

PMCPMC4023638.

37.   Uphoff CC, Drexler HG. Eradication of mycoplasma contaminations. Methods Mol

Biol. 2013;946:15-26. Epub 2012/11/28. doi: 10.1007/978-1-62703-128-8_2. PubMed

PMID: 23179823.

38.   Ullah I, Gahalawat S, Booshehri LM, Niederstrasser H, Majumdar S, Leija C, et al.

An antiparasitic compound from the Medicines for Malaria Venture Pathogen Box

promotes *Leishmania* tubulin polymerization. ACS Infect Dis. 2020;6(8):2057-72. Epub

2020/07/21. doi: 10.1021/acsinfecdis.0c00122. PubMed PMID: 32686409.

39.   Wetzel DM, Hakansson S, Hu K, Roos D, Sibley LD. Actin filament polymerization

regulates gliding motility by apicomplexan parasites. Mol Biol Cell. 2003;14(2):396-406.

doi: 10.1091/mbc.E02-08-0458. PubMed PMID: 12589042; PubMed Central PMCID:

PMC149980.

40.   Pan AA, McMahon-Pratt D. Monoclonal antibodies specific for the amastigote

stage of *Leishmania pifanoi*. I. Characterization of antigens associated with stage- and

species-specific determinants. J Immunol. 1988;140(7):2406-14. PubMed PMID:

2450920.

41.   Champsi J, McMahon-Pratt D. Membrane glycoprotein M-2 protects against

Leishmania amazonensis infection. Infect Immun. 1988;56(12):3272-9. PubMed PMID:

3182080; PubMed Central PMCID: PMCPMC259734.

42.   Soong L, Chang CH, Sun J, Longley BJ, Jr., Ruddle NH, Flavell RA, et al. Role of

CD4+ T cells in pathogenesis associated with *Leishmania amazonensis* infection. J

Immunol. 1997;158(11):5374-83. PubMed PMID: 9164958.

43.   Soong L, Henard CA, Melby PC. Immunopathogenesis of non-healing American

cutaneous leishmaniasis and progressive visceral leishmaniasis. Semin Immunopathol.

2012;34(6):735-51. Epub 2012/10/12. doi: 10.1007/s00281-012-0350-8. PubMed PMID:

23053396; PubMed Central PMCID: PMCPMC4111229.

44.   Navas A, Vargas DA, Freudzon M, McMahon-Pratt D, Saravia NG, Gomez MA.

Chronicity of dermal leishmaniasis caused by *Leishmania panamensis* is associated with

parasite-mediated induction of chemokine gene expression. Infect Immun.

2014;82(7):2872-80. doi: 10.1128/IAI.01133-13. PubMed PMID: 24752514; PubMed

Central PMCID: PMCPMC4097649.

45.   Gonzalez-Fajardo L, Fernandez OL, McMahon-Pratt D, Saravia NG. *Ex vivo* host

and parasite response to antileishmanial drugs and immunomodulators. PLoS Negl Trop

Dis. 2015;9(5):e0003820. Epub 2015/05/30. doi: 10.1371/journal.pntd.0003820. PubMed

PMID: 26024228; PubMed Central PMCID: PMC4449175.

46.   Falchook GS, Lewis KD, Infante JR, Gordon MS, Vogelzang NJ, DeMarini DJ, et

al. Activity of the oral MEK inhibitor trametinib in patients with advanced melanoma: a

phase 1 dose-escalation trial. Lancet Oncol. 2012;13(8):782-9. Epub 2012/07/19. doi:

10.1016/S1470-2045(12)70269-3. PubMed PMID: 22805292; PubMed Central PMCID:

PMCPMC4109286.

47.   Cheng Y, Tian H. Current development status of MEK inhibitors. Molecules.

2017;22(10). Epub 2017/09/29. doi: 10.3390/molecules22101551. PubMed PMID:

28954413; PubMed Central PMCID: PMCPMC6151813.

48.   Wong DJ, Robert L, Atefi MS, Lassen A, Avarappatt G, Cerniglia M, et al. Antitumor

activity of the ERK inhibitor SCH772984 [corrected] against BRAF mutant, NRAS mutant

and wild-type melanoma. Mol Cancer. 2014;13:194. Epub 2014/08/22. doi:

10.1186/1476-4598-13-194. PubMed PMID: 25142146; PubMed Central PMCID:

PMCPMC4155088.

49.   Ciuffreda L, Del Bufalo D, Desideri M, Di Sanza C, Stoppacciaro A, Ricciardi MR,

et al. Growth-inhibitory and antiangiogenic activity of the MEK inhibitor PD0325901 in

malignant melanoma with or without BRAF mutations. Neoplasia. 2009;11(8):720-31.

Epub 2009/08/04. doi: 10.1593/neo.09398. PubMed PMID: 19649202; PubMed Central

PMCID: PMCPMC2713590.

50.   Sun Y, Liu WZ, Liu T, Feng X, Yang N, Zhou HF. Signaling pathway of MAPK/ERK

in cell proliferation, differentiation, migration, senescence and apoptosis. J Recept Signal

Transduct Res. 2015;35(6):600-4. Epub 2015/06/23. doi:

10.3109/10799893.2015.1030412. PubMed PMID: 26096166.

51.   Crowley MT, Costello PS, Fitzer-Attas CJ, Turner M, Meng F, Lowell C, et al. A

critical role for Syk in signal transduction and phagocytosis mediated by Fcgamma

receptors on macrophages. J Exp Med. 1997;186(7):1027-39. Epub 1997/10/07. doi:

10.1084/jem.186.7.1027. PubMed PMID: 9314552; PubMed Central PMCID:

PMCPMC2199061.

52.   Kiefer F, Brumell J, Al-Alawi N, Latour S, Cheng A, Veillette A, et al. The Syk

protein tyrosine kinase is essential for Fcgamma receptor signaling in macrophages and

neutrophils.   Mol   Cell   Biol.   1998;18(7):4209-20.   Epub   1998/06/25.   doi:

10.1128/MCB.18.7.4209.   PubMed   PMID: 9632805;   PubMed   Central   PMCID:

PMCPMC109005.

53.   Liu L, Mayes PA, Eastman S, Shi H, Yadavilli S, Zhang T, et al. The BRAF and

MEK inhibitors dabrafenib and trametinib: effects on immune function and in combination

with immunomodulatory antibodies Targeting PD-1, PD-L1, and CTLA-4. Clin Cancer

Res.   2015;21(7):1639-51.   Epub   2015/01/16.   doi: 10.1158/1078-0432.CCR-14-2339.

PubMed PMID: 25589619.

54.   Shi-Lin D, Yuan X, Zhan S, Luo-Jia T, Chao-Yang T. Trametinib, a novel MEK

kinase inhibitor, suppresses lipopolysaccharide-induced tumor necrosis factor (TNF)-

alpha   production   and   endotoxin   shock.   Biochem   Biophys   Res   Commun.

2015;458(3):667-73. Epub 2015/02/17. doi: 10.1016/j.bbrc.2015.01.160. PubMed PMID:

25684183.

55.   Boggiatto PM, Jie F, Ghosh M, Gibson-Corley KN, Ramer-Tait AE, Jones DE, et

al. Altered dendritic cell phenotype in response to *Leishmania amazonensis* amastigote

infection is mediated by MAP kinase, ERK. Am J Pathol. 2009;174(5):1818-26. doi:

10.2353/ajpath.2009.080905.   PubMed   PMID: 19349356;   PubMed   Central   PMCID:

PMCPMC2671270.

56.   Cheeseman KL, Ueyama T, Michaud TM, Kashiwagi K, Wang D, Flax LA, et al.

Targeting of protein kinase C-epsilon during Fcgamma receptor-dependent phagocytosis

requires   the   epsilonC1B   domain   and   phospholipase   C-gamma1.   Mol   Biol   Cell.

2006;17(2):799-813. Epub 2005/12/02. doi: 10.1091/mbc.e04-12-1100. PubMed PMID:

16319178; PubMed Central PMCID: PMCPMC1356590.

57.   Larsen EC, Ueyama T, Brannock PM, Shirai Y, Saito N, Larsson C, et al. A role

for PKC-epsilon in Fc gammaR-mediated phagocytosis by RAW 264.7 cells. J Cell Biol.

2002;159(6):939-44. Epub 2002/12/25. doi: 10.1083/jcb.200205140. PubMed PMID:

12499353; PubMed Central PMCID: PMCPMC2173999.

58.   Panetti TS, Wilcox SA, Horzempa C, McKeown-Longo PJ. Alpha v beta 5 integrin

receptor-mediated endocytosis of vitronectin is protein kinase C-dependent. J Biol Chem.

1995;270(31):18593-7. Epub 1995/08/04. doi: 10.1074/jbc.270.31.18593. PubMed

PMID: 7543105.

59.   Roy N, Chakraborty S, Paul Chowdhury B, Banerjee S, Halder K, Majumder S, et

al. Regulation of PKC mediated signaling by calcium during visceral leishmaniasis. PLoS

One. 2014;9(10):e110843. Epub 2014/10/21. doi: 10.1371/journal.pone.0110843.

PubMed PMID: 25329062; PubMed Central PMCID: PMCPMC4201563.

60.   Zheleznyak A, Brown EJ. Immunoglobulin-mediated phagocytosis by human

monocytes requires protein kinase C activation. Evidence for protein kinase C

translocation to phagosomes. J Biol Chem. 1992;267(17):12042-8. Epub 1992/06/15.

PubMed PMID: 1376316.

61.   Arthur JS, Ley SC. Mitogen-activated protein kinases in innate immunity. Nat Rev

Immunol. 2013;13(9):679-92. Epub 2013/08/21. doi: 10.1038/nri3495. PubMed PMID:

23954936.

62.   Buxbaum LU, Scott P. Interleukin 10- and Fcgamma receptor-deficient mice

resolve *Leishmania mexicana* lesions. Infect Immun. 2005;73(4):2101-8. Epub

2005/03/24. doi: 10.1128/IAI.73.4.2101-2108.2005. PubMed PMID: 15784551; PubMed

Central PMCID: PMCPMC1087424.

63.   Tada S, Anazawa T, Shindo T, Yamane K, Inoguchi K, Fujimoto N, et al. The MEK

inhibitor trametinib suppresses Major Histocompatibility Antigen-mismatched rejection

following pancreatic islet transplantation. Transplant Direct. 2020;6(9):e591. Epub

2020/08/28. doi: 10.1097/TXD.0000000000001045. PubMed PMID: 32851124; PubMed

Central PMCID: PMCPMC7423917.

8. LIST OF FIGURES

Figure 1. The MAPK/ERK pathway is required for optimal amastigote uptake.

Figure 2. MAPK/ERK are activated during *Leishmania* uptake by Mϕ.

Figure 3. PMA activates ERK1/2 to modulate amastigote uptake.

Figure 4. SYK and Arg/Abl stimulate ERK1/2 phosphorylation during amastigote

uptake.

Figure 5. ERK 1/2's role in *Leishmania* uptake is greater than for phagocytosis of

beads.

Figure 6. MAPK/ERK permit efficient infection in a mouse model of cutaneous

leishmaniasis.

Table 1. Ongoing trametinib treatment during *L. amazonensis* infection has

limited effects on Th1 vs Th2 responses.

Table 2. Cytokine and chemokine profiles in diluent versus trametinib-treated,

Leishmania-infected mice.

9. LIST OF SUPPLEMENTARY MATERIALS

Supplemental Figure 1. MAPK/ERK inhibitors decrease uptake and survival of

L. amazonensis promastigotes and amastigotes.

Supplemental Figure 2. Inhibiting kinases upstream of ERK during *L.*

amazonensis uptake decreases ERK phosphorylation.

Supplemental Figure 3. Inhibiting ERK1/2 has minimal effects on the uptake of

opsonized beads.

Supplemental Figure 4. Actin-based phagocytic cups are brighter in trametinib-

treated Mϕs taking up *Leishmania* amastigotes compared to DMSO-treated Mϕs.

Supplemental Table 1. Trametinib initiated during *L. amazonensis* infection has

limited effects on Th1 vs Th2 responses.

Supplemental Table 2. Cytokine and chemokine profiles from *Leishmania*-

infected mice treated once lesions are visible.

Supplemental Table 3. Antibodies.

<u>**10. FIGURES AND LEGENDS**</u>

Figure 1

Figure 1. The MAPK/ERK pathway is required for optimal amastigote

uptake. (A) Trametinib decreases IgG-opsonized amastigote uptake. RAW 264.7

cells were treated with 3.3 μM imatinib, 1 μM trametinib or DMSO for 2 h, and 10

IgG-opsonized amastigotes were incubated per RAW 264.7 cell for 2 h at 37°C.

An immunofluorescence assay distinguished between intracellular (green) and

extracellular (red + green) amastigotes. Nuclei are labeled with Hoescht 33342

(blue). Scale bar, 100 μm. **(B)** Graph showing the mean phagocytic index (PI) ±

standard deviation (SD) for RAW 264.7 cells normalized to DMSO for each

experiment (100%), quantified from n = 6 biological experiments. Comparisons

between imatinib and trametinib by ANOVA were nonsignificant. **(C)** Imatinib and

trametinib do not affect the total (external + internal) number of amastigotes per

100 RAW cells (termed an adhesion index). Shown are percentages of total

amastigotes per 100 imatinib-treated and trametinib-treated RAW 264.7 cells

relative to DMSO-treated cells for the experiment shown in B. **(D)** Titration of

trametinib demonstrates that the reduction of amastigote internalization is

maximal at ~100x the purified MEK1/2 IC_{50}. RAW 264.7 cells were treated with

increasing doses of trametinib (0.25, 0.5, 1, or 2.5 µM) or DMSO. Bar graph

shows the mean PI ± SD for trametinib-treated samples relative to DMSO-treated

samples for each experiment (100%) for 4 biological experiments. **(E)** The

adhesion index for amastigotes from experiment in D is not affected by

increasing doses of trametinib. Experiment performed as in C. **(F)** The relative

decrease in PI for trametinib-treated samples persists even after prolonged

incubation. RAW 264.7 cells were treated with DMSO (D) or 1 µM trametinib (T)

prior to addition of IgG-opsonized amastigotes for 1 h, 2 h, or 4 h, as indicated.

Shown is the mean PI ± SD for each time point for the trametinib-treated

condition, relative to the DMSO-treated condition, calculated from 4 biological

experiments. **(G)** Treating samples with trametinib for prolonged periods does not

affect the adhesion index for amastigotes. Shown are the total number of

amastigotes (internal + external) per 100 RAW 264.7 cells for the experiments

performed in F. For all experiments, ** *P* < 0.01, **** *P* < 0.0001, ns =

nonsignificant compared to the relevant DMSO-treated category by ANOVA. **(H)**

Trametinib is not toxic to amastigotes. Shown is one representative experiment

of five experiments with two technical replicates following the number of

amastigotes growing in untreated, DMSO-treated, or trametinib-treated media

over 5 days relative to on day 0. ns, $P > 0.05$ compared to DMSO category by

ANOVA.

1024

Figure 2

Figure 2. **MAPK/ERK are activated during *Leishmania* uptake by Mϕ.**

Amastigotes were added to $\geq$ 70% confluent RAW 264.7 cells. Equal amounts of

protein from whole lysates were employed for each category. **Top:**

Representative blots displaying relative activation of MAPK/ERK. Phosphorylated

(A) ERK1/2, **(B)** MEK1/2, **(C)** cRaf (Raf-1) and the corresponding total proteins

were detected by Western Blot. An unrelated protein, GAPDH, is shown as an

additional loading control. **Bottom:** Quantification of phosphorylated MAPK/ERK

levels. Graphs shown in **(A)** for ERK1/2, **(B)** for MEK1/2, and **(C)** for cRaf

represent mean $\pm$ SD of relative phosphorylated kinase levels, normalized to total

kinase levels, for at least 3 biological replicates per condition. * $P < 0.05$, ** $P <$

0.01, *** $P < 0.001$, **** $P < 0.0001$ by ANOVA compared to the 0 m time point

for each set of kinases.

1040 **Figure 3**

1041

Figure 3. PMA activates ERK1/2 to modulate amastigote uptake. **(A, B)** PKC

typically is only slightly activated during amastigote uptake. PMA serves as a

positive control. Amastigotes were added to monolayers of RAW 264.7 cells and

processed as above. Shown are **(A)** representative blots demonstrating and **(B)**

bar graphs quantifying relative activation of pPKC/PKC. *$P < 0.05$, *** $P < 0.001$,

**** $P < 0.0001$ compared with samples indicated by brackets (ANOVA). n = 3

experiments. **(C)** PMA is not toxic to amastigotes. Shown is one representative

experiment of five experiments with two technical repeats following the number of

amastigotes growing in DMSO-treated or PMA-treated media over 5 days relative

to on day 0. ns, $P > 0.05$ compared to DMSO category by ANOVA. **(D)**

Representative image demonstrating that PMA rescues trametinib inhibition of

Leishmania amastigote uptake. BMDM were treated with DMSO, 1 µM PMA, 1

µM trametinib or a combination of 1 µM trametinib and 1 µM PMA prior to

incubation with IgG-opsonized amastigotes for 2 hours. **(E)** PMA increases IgG-

opsonized amastigote uptake. BMDM were treated with DMSO or increasing

doses of PMA (0.25, 0.5, 1, 2.5 µM) prior to incubation with IgG-opsonized

amastigotes for the times indicated. n = 4 experiments. ** $P < 0.01$, *** $P < 0.001$,

**** $P < 0.0001$ compared to DMSO category by ANOVA. **(F)** PMA does not

affect the adhesion index for amastigotes. ns, not significant by ANOVA. **(G)**

Mean PI ± SD for BMDM treated with trametinib/PMA relative to DMSO-treated

samples for each experiment (100%), obtained from 3 biological experiments.

**** $P < 0.0001$, ns = nonsignificant by ANOVA compared to DMSO-treated

sample. **(H)** Trametinib/PMA does not affect amastigote adhesion for the

experiment shown in G.

1066

1067 **Figure 4**

1068

Figure 4. SYK and Arg/Abl stimulate ERK1/2 phosphorylation during

amastigote uptake. (A) Phosphorylation of ERK1/2 in RAW 264.7 cells is

induced upon amastigote uptake and is decreased by imatinib (Arg/Abl inhibitor),

entospletinib (SYK inhibitor), SCH772984 (ERK1/2 inhibitor), and trametinib

(MEK1/2 inhibitor). PMA serves as a positive control for pERK stimulation. RAW

264.7 cells were allowed to adhere to plates, and then treated with compounds or

DMSO prior to IgG-opsonized amastigote addition, lysis, and immunoblotting.

Representative immunoblots of pERK1/2 (top) and total ERK1/2 (bottom) [with

(+) or without (–) amastigotes] are shown. **(B)** Relative pERK levels, normalized

to ERK1/2 levels, among categories shown in A. Levels of pERK are normalized

to amastigote-exposed DMSO-treated RAW 264.7 cells (100%). 5 biological

replicates are quantified. *** $P < 0.001$, **** $P < 0.0001$ compared with

amastigote-stimulated, DMSO-treated RAW 264.7 cells (ANOVA). **(C)**

Phosphorylation of ERK1/2 induced upon FcγR engagement is decreased

in $SYK^{-/-}$ and $Arg^{-/-}Abl^{-/-}$ BMDM compared to WT BMDM. Shown is a

representative immunoblot of pERK1/2 (top) and total ERK1/2 (bottom) in WT,

$SYK^{-/-}$ and $Arg^{-/-}Abl^{-/-}$ BMDM treated with DMSO (D) or PMA (P), $\pm$ incubation

with amastigotes. **(D)** Graph showing relative levels of pERK1/2, normalized to

ERK1/2 levels, among WT, $SYK^{-/-}$, and $Arg^{-/-}Abl^{-/-}$ BMDM treated with DMSO

(D) or PMA (P), $\pm$ amastigotes (A). ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$, ns:

not significant by ANOVA (n = 6 experiments), compared to DMSO + amastigote

category for each condition. **(E-G)** Primary cells lacking Arg and Abl exhibit

decreases in amastigote uptake, but PMA can partially rescue this amastigote

internalization defect. **(E)** Representative image performed as in Fig 1. **(F)** Mean

phagocytic index ± SD for PMA-treated and/or $Arg^{-/-}Abl^{-/-}$ BMDM relative to WT

DMSO-treated BMDM. **(G)** DMSO and PMA do not affect the adhesion indexes

for the experiments shown in F. n for F-G = 3 experiments. ** $P < 0.01$; *** $P <$

0.001; **** $P < 0.0001$; ns, not significant compared with DMSO-treated WT

BMDM (ANOVA).

1098

 Figure 5

1100

Figure 5. ERK 1/2's role in *Leishmania* uptake is greater than for

phagocytosis of beads. **(A)** Trametinib mildly decreases bead internalization.

RAW 264.7 cells were treated with 3.3 µM imatinib, 1 µM trametinib or DMSO for

2 hr. prior to addition of 10 IgG-opsonized beads per RAW 264.7 cell for 2 h at

37°C. Treating RAW 264.7 cells with trametinib decreases uptake of IgG-

opsonized beads to a lesser degree than imatinib. Graph shows the mean PI ±

SD for each category, normalized to DMSO (100%). **** *P* < 0.0001, ns: not

significant compared to indicated categories by ANOVA (n = 6 biological

experiments). **(B)** Imatinib and trametinib does not affect the adhesion index for

beads in the experiments shown in A. **(C-H)** IgG-coated beads activate the

MAPK/ERK pathway, but phosphorylation is not sustained. Beads were added to

> 70% confluent RAW 264.7 cells and experiments were processed as in Fig 2.

Top: Phosphorylated forms of **(C)** ERK, **(E)** MEK1/2, **(G)** cRaf and the

1114   corresponding total kinase were detected by Western blotting. Bottom: Relative

1115   activation was quantified for **(D)** ERK, **(F)** MEK1/2, and **(H)** cRaf, using the

1116   methods described in Fig 2. * *P* < 0.05, ** P < 0.01, *** *P* < 0.001, **** *P* < 0.0001

1117   and ns: not significant compared to 0 m timepoint by ANOVA (n ≥ 3

1118   experiments).

1119

Figure 6

Figure 6. MAPK/ERK permit efficient infection in a mouse model of

cutaneous leishmaniasis. (A) Mouse footpad model for cutaneous

leishmaniasis. C57BL/6 mice diluent and trametinib-treated groups were injected

with promastigotes in the hind footpad. Mice were given 2 mg/kg body weight of

trametinib or an equal volume of diluent by oral gavage daily, starting 7 days

before infection and continuing until the mice were euthanized. Two experiments

were performed. Shown in the following panels is one representative experiment

containing 10 female mice per group. **(B)** Trametinib given prior to infection

reduces lesion size in mice. Depth of foot swelling was monitored with calipers by

a blinded investigator. Results represent the mean ± SD foot size increase over

the uninfected foot (normalized to 1). The control (diluent) and drug-treated

groups were compared to swelling at time 0 by ANOVA; *$P < 0.05$, ****$P <$

0.0001, ns, not significant. **(C)** Trametinib given prior to infection significantly

reduces parasite burden. Parasite burdens in infected footpads were determined

by limiting dilution as described in *Materials and Methods*. Plotted is the mean

number of parasites (*i.e.*, parasite burden) in millions at the end of the

experiment shown in B ± SD. In this example, there were 1.41×10^6 parasites in

the control mice versus 2.03×10^5 parasites in the trametinib-treated mice. *$P <$

0.05 (two-tailed *t*-test). **(D)** C57BL/6 mice diluent and trametinib-treated groups

were injected with *L. amazonensis* promastigotes as above. Starting 6 weeks

after inoculation (arrow), mice were given 2 mg/kg body weight of trametinib or

vehicle by oral gavage daily and continuing until the experiment was terminated.

(E) Trametinib given once lesions have begun reduces lesion size in trametinib-

treated mice compared to diluent-treated mice. Arrow shows start time of drug

treatment. Shown is one representative experiment containing 10 female mice

per group. Statistics calculated as in B. **(F)** Lesions in mice treated with

trametinib 6 weeks after infection contain fewer *Leishmania* parasites than

control mice. Plotted is the mean ± SD lesion parasite burden in millions at the

end of the experiment shown in E; in this example, there were 1.01×10^6

parasites in the control mice versus 8.29×10^4 parasites in the trametinib-treated

mice (>10-fold reduction). *P <0.05 (two-tailed *t*-test). (**G**) MAPK/ERK signal

transduction in Mɸ facilitates *Leishmania* uptake and pathogenesis. Upon FcγR

ligation by amastigotes, SFK, Arg, and SYK are activated and relay signals to the

MAPK/ERK pathway, which allows the internalization of *Leishmania*. Diagrams

created with Biorender.

1157

1158 **Table 1**

Experiment	Ratio	DMSO - High	Trametinib - High	DMSO - Low	Trametinib - Low
Trametinib prior to infection	**IFN Gamma/IL-4**	1.86	1.35	1.75	1.05
	IFN Gamma/IL-13	0.78	0.11	0.75	0.09
	IFN Gamma/IL-4+ IL-10+ IL-13	0.55	0.10	0.52	0.08

1159

Table 1. Ongoing trametinib treatment during *L. amazonensis* infection has

limited effects on Th1 vs Th2 responses. Effects of trametinib during *L.*

amazonensis infection on cytokine secretion. Shown are profiles of draining

lymph nodes isolated from diluent (DMSO) versus trametinib-treated mice treated

prior to inoculation as described in methods. High refers to high stimulation with a

lysate of 1 X 10^6 parasites and low refers to low stimulation with 1 X 10^5

parasites. Ratios of IFNγ: IL-4, IFNγ: IL-13, and IFNγ: IL-4+10+13 were

calculated from the data shown in Table 2. Trametinib may skew towards a Th2

response, which should not promote lesion healing.

1169

1170 **Table 2**

| Experiment | Cytokine | Treatment | Cytokine secretion (pg) after | | | | | |
| | | | Con A stimulation | | High stimulation | | Low treatment | |
			AVG	STD	AVG	STD	AVG G	STD
Trametinib prior to infection	**IFN Gamma**	DMSO	6.70	6.83	6.06	5.82	5.83	5.16
		Trametinib	0.48	0.64	0.73	1.05	0.58	0.75
		p-value	**0.01**		**0.01**		**0.005**	
	IL-1 BETA	DMSO	0.88	0.51	0.97	0.48	0.73	0.46
		Trametinib	0.45	0.12	0.42	0.11	0.39	0.09
		p-value	**0.02**		**0.002**		**0.03**	
	IL-12/IL23p40	DMSO	5.56	2.97	5.24	2.49	6.04	3.06
		Trametinib	3.28	1.18	3.39	0.79	3.53	1.31
		p-value	**0.04**		**0.04**		**0.03**	
	IL-13	DMSO	5.08	3.04	7.72	2.47	7.78	3.07
		Trametinib	7.01	2.27	6.83	1.10	6.57	1.15
		p-value	NS		NS		NS	
	IL-17A	DMSO	0.48	0.24	0.65	0.14	0.53	0.27
		Trametinib	0.59	0.09	0.74	0.25	0.53	0.25
		p-value	NS		NS		NS	
	IL-4	DMSO	2.36	1.26	3.26	1.89	3.32	2.57
		Trametinib	0.49	0.64	0.54	0.68	0.55	0.70

	p-value	**0.0006**		**0.0004**		**0.004**	
TNF ALPHA	DMSO	2.98	1.6	2.68	2.10	2.45	1.22
	Trametinib	2.85	2.30	2.23	0.96	1.75	1.45
	p-value	NS		NS		NS	
IL-10	DMSO	.008	.007	.02	.009	.009	.01
	Trametinib	8.94	8.78	8.36	10.44	5.16	1.06
	p-value	**0.005**		**0.02**		**0.0001**	
MCP-1	DMSO	25.56	14.87	25.07	12.31	27.90	10.54
	Trametinib	7.23	6.43	7.23	5.87	7.23	6.56
	p-value	**0.002**		**0.0006**		**0.0001**	
MIP-1 ALPHA	DMSO	4.69	3.95	3.93	2.77	4.62	2.96
	Trametinib	1.36	0.41	1.44	0.86	1.44	0.88
	p-value	**0.02**		**0.01**		**0.004**	

1171

Table 2. Cytokine and chemokine profiles in diluent versus trametinib-

treated, *Leishmania*-infected mice. Shown are profiles of draining lymph nodes

isolated from DMSO versus trametinib mice treated prior to infection as

described in methods. For high stimulation categories, a lysate containing 1 X

10^6 *L. amazonensis* was added, and for low stimulation categories, a lysate

containing 1 X 10^5 parasites was added. Post 14-16 weeks, mice were sacrificed

and multiplex chemokine ELISAs were performed on harvested lymph node

supernatants. *P* values were determined by two-tailed *t*-tests and are listed if $P <$

0.1. Bold: $P \leq 0.05$.

1181