Deep divergence and genomic diversification of gut symbionts of neotropical stingless bees

[#]Garance Sarton-Lohéac[a], [#]Carlos Gustavo Nunes da Silva[e], Florent Mazel[b], Gilles Baud[a],

Vincent de Bakker[a], Sudip Das[a], Yassine El Chazli[a], Kirsten Ellegaard[a], Marc Garcia-

Garcera[d], Natasha Glover[c], Joanito Liberti[a,b], Lorena Nacif Marçal[e], Aiswarya Prasad[a],

Vincent Somerville[a], SAGE class 2019-2020 and 2020-2021, Germán Bonilla-Rosso[a],

Philipp Engel[a]

[a]Department of Fundamental Microbiology, University of Lausanne, 1015 Lausanne,

Switzerland

[b]Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

[c]Swiss Institute of Bioinformatics, Lausanne, Switzerland

[d]Société des produits Nestlé, Lausanne, Switzerland

[e] Federal University of Amazonas, Instituto de Ciências Biológicas, Dept of Morphology,

Manaus, Brazil

[#]equally contributed

Correspondence:

Prof. Philipp Engel

Department of Fundamental Microbiology

University of Lausanne, CH-1015 Lausanne, Switzerland

Tel.: +41 (0)21 692 56 12

e-mail: philipp.engel@unil.ch

Abstract

Social bees harbor conserved gut microbiota that may have been acquired in a common ancestor of social bees and subsequently co-diversified with their hosts. However, most of this knowledge is based on studies on the gut microbiota of honey bees and bumble bees. Much less is known about the gut microbiota of the third and most diverse group of social bees, the stingless bees. Specifically, the absence of genomic data from their microbiota presents an important knowledge gap in understanding the evolution and functional diversity of the social bee microbiota. Here we combined community profiling with culturing and genome sequencing of gut bacteria from six neotropical stingless bee species from Brazil. Phylogenomic analyses show that most stingless bee gut isolates form deep-branching sister clades of core members of the honey bee and bumble bee gut microbiota with conserved functional capabilities, confirming the common ancestry and ecology of their microbiota. However, our bacterial phylogenies were not congruent with those of the host indicating that the evolution of the social bee gut microbiota was not driven by strict co-diversification, but included host switches and independent symbiont gain and losses. Finally, as reported for the honey bee and bumble bee microbiota, we find substantial genomic divergence among strains of stingless bee gut bacteria suggesting adaptation to different host species and glycan niches. Our study offers first insights into the genomic diversity of the stingless bee microbiota, and highlights the need for broader samplings to understand the evolution of the social bee gut microbiota.

Importance

Stingless bees are the most diverse group of the corbiculate bees and represent important pollinator species throughout the tropics and subtropics. They harbor specialized microbial communities in their gut that are related to those found in honey bees and bumble bees and that are likely important for bee health. Few bacteria have been cultured from the gut of stingless bees which has prevented characterization of their genomic diversity and functional potential. Here, we established cultures of major community members of the gut microbiota of six stingless bee species and sequenced their genomes. We find that most stingless bee isolates belong to novel bacterial species distantly related to those found in honey bees and bumble bees and encoding similar functional capabilities. Our study offers a new perspective on the evolution of the social bee gut microbiota and presents the basis to characterize the symbiotic relationships between gut bacteria and stingless bees.

Introduction

The eusocial corbiculate bees (hereafter social bees) comprise more than 700 species distributed in three distinct tribes: honey bees (Apini), bumble bees (Bombini), and stingless bees (Meliponini). Stingless bees and bumble bees form a monophyletic clade, which is sister to the honey bees, with whom they shared a common ancestor 80-100 million years ago [1-3]. Like mammals, social bees harbor dense and specialized bacterial communities in their gut that affect bee health and behavior [4-12]. The composition of the gut microbiota of social bees is relatively simple typically consisting of <10 bacterial phylotypes, i.e. sequence clusters sharing >97% identity in the 16S rRNA gene [13-20]. Five of these phylotypes (*Snodgrassella*, *Gilliamella*, Bombilactobacillus Firm-4, *Lactobacillus* Firm-5, and *Bifidobacterium*) have been referred to as the core gut microbiota of the social bees [20], because they are prevalent and abundant across honey bee, bumble bee, and stingless bees. Most members of the bee gut microbiota are culturable and gnotobiotic bees can be generated for several species [5, 21-23]. Together, these distinctive characteristics make the social bee microbiota a versatile model system for studying the evolution and ecology of host-associated microbial communities. Moreover, social bees are important pollinators that suffer from severe population declines [24, 25] which makes studies of their microbiota relevant in its own right.

Most of what is currently known about the gut microbiota of social bees stems from studies on honey bees and bumble bees. Genomic and experimental approaches have revealed that their gut bacteria are usually saccharolytic fermenters that utilize plant glycans derived from the pollen and nectar/honey diet of the host [22, 23, 26-32]. Further, it has been shown that the core members of the honey bee and bumble bee gut microbiota have substantially diversified [27-30, 33-36]. They consist of divergent sub-lineages (or species) and contain large extents of strain-level diversity and gene content variation. Most sub-lineages are host-specific [23, 27, 33, 36], and their phylogenetic relationships are to some extent congruent with the phylogeny

of the host (16, 33). Therefore, it has been suggested that the core members of the microbiota

were acquired in a common ancestor of the social bees (20) and possibly co-diversified with

the host (16, 33). In addition, studies in the Western honey bee (*Apis mellifera*) have shown

that the diversification of the bee gut microbiota was also driven by adaptation to different

spatial and metabolic niches within the gut (27, 30, 33-35). For example, strains of closely

related sub-lineages of *Lactobacillus* Firm-5 and *Bifidobacterium* can coexist in individual

bees. They carry distinct gene sets for the breakdown and utilization of pollen-derived

carbohydrates which allows them to partition the available dietary glycan niches in the gut (27,

30, 34).

In contrast to the microbiota of honey bees and bumble bees, much less is known about the gut

microbiota of the third group of social bees, the stingless bees (Meliponini). Previous studies

have focused on determining the taxonomic composition of the gut microbiota of these bees

using 16S rRNA gene sequencing (15, 17, 20, 37-39). However, only a few bacteria have been

cultured from stingless bees (40-42) and, except for two strains of *Bombilactobacillus* Firm-4

recently isolated from bees from Australia (43), no genomic data is currently available for core

members of the gut microbiota of stingless bees.

With >500 described species, stingless bees present the largest and most diverse group of the

social bees (44, 45). They are naturally distributed throughout the tropical and subtropical

regions of Africa, Asia, Australia, and the Americas and exhibit great variation in morphology,

diet, foraging range, social structure, and nesting habits (44, 45). As host phylogeny and

ecology are both key determinants of gut microbiota composition (46-51), we hypothesize that

genomic studies on bacterial isolates will help us to understand the functional diversity of gut

bacteria of stingless bees and provide novel insights into the evolution of these bacteria across

social bees, specifically in respect of the possible co-diversification with the host.

To address these questions, we looked at the gut microbiota of six neotropical species of

stingless bees from Brazil: *Frieseomelitta varia* (Fv), *Scaptotrigona polysticta* (Sp), *Melipona*

fuliginosa (Mf), *Melipona interrupta* (Mi), *Melipona seminigra* (Ms), and *Melipona lateralis*

(Ml). We determined the composition of the gut microbiota of these bees using 16S rRNA gene

sequencing, established a comprehensive culture collection of bacterial isolates, and conducted

genome sequencing and comparative genomics to determine the phylogenetic placement,

genomic diversity, and functional capabilities of these bacteria relative to those previously

isolated from honey bees and bumble bees.

Results

Six neotropical stingless bee species from Brazil harbor distinct gut microbiota dominated by nine bacterial families.

We sampled three colonies of six stingless bee species (Fv, Sp, Mf, Mi, Ms, Ml) from a meliponary located in the Amazonian rainforest near Manaus (**Supplementary Table 1**). For each colony, we pooled the guts of 15-60 worker bees (depending on the size of the bee species, see Methods) before DNA isolation. The V4 region of the 16S rRNA gene was amplified and sequenced with Illumina MiSeq 2x250bp resulting in a median depth of 64,713 (52,479 to 95,774) reads per sample. In total, we identified 277 amplicon sequence variants (ASVs, 29-63 ASVs per sample), which belonged to 36 different bacterial families (**Supplementary Table 2**). Despite this diversity, only nine families dominated the samples, representing together 97% of all quality-filtered reads (93-99% of the reads per sample): Acetobacteraceae, Bifidobacteriaceae, Enterobacteriaceae, Lactobacillaceae, Neisseriaceae, Orbaceae, Prevotellaceae, Streptococcaceae and Veillonellaceae (**Figure 1A**).

While Lactobacillaceae and Bifidobacteriaceae were abundant across all samples, there were clear differences in the distribution of some of the other bacterial families (**Figure 1A).** Neisseriaceae were abundant in the samples of Sp and Fv, but were only detected in three out of 12 samples from the genus *Melipona* (**Figure 1A, Supplementary Table 2**). In contrast, Acetobacteraceae and Streptococcaceae were present in most *Melipona* samples but rare across samples of Fv and Sp. Orbaceae and Enterobacteriaceae were mostly detected in the three Fv samples. Intriguingly, a single Enterobacteriaceae ASV constituted the most abundant community member in this bee species (21 – 36% of the reads per sample). According to these compositional differences, non-metric multidimensional scaling (NMDS) separated the samples into three distinct clusters: two clusters comprised all samples of Fv and Sp,

respectively, and the third cluster comprised all samples of the four *Melipona* species (Mf, Mi,

Ms, Ml) (**Supplementary Figure 1A**).

Related stingless bee species have overlapping community profiles.

We compared our results to a previously published amplicon sequencing dataset from stingless

bees (20) to assess the similarity of the communities to those of other stingless bee species.

After discarding samples with <5,000 reads to control for variation in sequencing depths, our

dataset comprised 135 samples from 19 different host species and three different countries. We

detected 688 ASVs in total with a median of 18 ASVs per sample (3 to 63 ASVs)

(**Supplementary Table 3**), spanning 53 bacterial families. Overall, the taxonomic patterns

were similar across the analyzed bee species. Apart from one sample of *Tetragonula*

fuscobalteata, for which 99% of the reads belonged to a single Weeksellaceae ASV, the nine

families dominating in the six bee species from our study were also abundant in the microbiota

of the samples from the previous study and represented between 34% to 100% of the total

number of reads (**Supplementary Figure 1B**, **Supplementary Table 3**).

NMDS based on ASV relative abundances separated the samples by location (i.e. samples from

Brazil were different from those from Australia and Malaysia**,** Permutational multivariate

analysis of variance (PERMANOVA)**: '**Location' pseudo-F: 20.17, p-value=0.001**)**, and by

bee genus (PERMANOVA: 'Genus' pseudo-F=10.49, p-value=0.001), although taxonomy and

geography are not independent (**Figure 1B**). In contrast, there was only weak clustering at the

species level ('Species' pseudo-F=1.96, p-value=0.03). Notably, only 30% of all ASVs (206

of 688 ASVs) were shared across host species (i.e. 70% of all ASVs are only found in one

species), and most of them (83.4%) only between 2-5 species (**Supplementary Figure 1C**).

However, the shared ASVs belonged to the nine predominant bacterial families and represented

a large fraction of the total number of reads per sample (53.3-99.7 %; except for the samples

of Fv and *P. helleri*: 10.6-15 % of the reads) (**Supplementary Figure 1D**). In particular, bees

sampled in the same country or belonging to the same bee genus shared the same ASVs,

explaining the clustering of these samples in the NMDS analysis. Together, these results show

that despite the large variability observed, the gut microbiota of most stingless bee species is

dominated by a few bacterial families and that bee species of the same genus, or with

overlapping geographic distribution, have similar community profiles at the 16S rRNA gene

level.

Establishment of a strain collection of gut bacteria isolated from stingless bees.

To enable genomic and experimental analyses of stingless bee gut bacteria, we established a

culture collection of bacteria isolated from Fv, Sp, Mf, Mi, Ms, and Ml. We plated

homogenized gut samples of the six bee species on eight different semi-solid media and under

three different atmospheres (microaerobic and anaerobic). This resulted in the cultivation of 98

distinct bacterial isolates (i.e. different 16S rRNA genotype or isolated from a different bee

species or colony) from 11 bacterial families (**Figure 2A, Supplementary Table 4**). Most

bacteria grew under both microaerobic or anaerobic conditions on generic growth media and

formed colonies after 2-4 days of growth. The 16S rRNA genotypes of the isolated strains

matched to 32 ASVs, accounting for 16 – 87 % of the overall community of the six stingless

bee species and including many shared ASVs (**Figure 2A** and **B**). BLASTN searches of the

16S rRNA gene sequences revealed that many of the isolates (55/98) were related to bacterial

strains obtained from the gut of honey bees and bumble bees (such as *Lactobacillus apis*,

Bifidobacterium commune, *Gilliamella sp.,* or *Snodgrassella alvi*) suggesting that they

represent stingless bee isolates of core members of the social bee microbiota. Other isolates

had best BLASTN hits to bacteria from other environments, such as *Floricoccus* sp. isolated

from flowers, various Enterobacteriaceae (e.g. *Pantoea* sp., *Klebsiella* sp., *Rosenbergiella*

epipactidis) isolated from humans and water, or *Fructobacillus* isolated from flowers and

fruits. The percent identity of many of the blast hits was relatively low (<98%) suggesting that

the isolated strains potentially correspond to new bacterial species (**Figure 2A** and **C,**

Supplementary Table 4).

Stingless bee isolates form deep-branching phylogenetic lineages related to bacteria

isolated from honey bees and bumble bees.

To assess the phylogenetic placement of the isolated stingless bee gut bacteria relative to gut

bacteria from honey bees and bumble bees, we selected 46 strains from 10 different bacterial

families for genome sequencing (**Figure 2A, Supplementary Table 4**). Using a combination

of Illumina and Oxford Nanopore sequencing, we obtained 23 complete and 23 draft genomes

(2-66 contigs). Genome size of the cultured isolates ranged from 1.2-6.3 Mb. *Fructobacillus*

ESL0730 (1.2Mb) and the two Streptococcaceae strains ESL0687 and ESL0729 (1.4 Mb)

harbored the smallest, and Leuconostocaceae ESL0723 the largest (6.3 Mb) genomes of the

sequenced strains (**Supplementary Table 4**). Genome comparisons with other bacteria,

including strains isolated from honey bees and bumble bees, showed that most stingless bee

gut bacteria had 80% average nucleotide identity with previously sequenced strains indicating

that we have isolated strains of novel bacterial species or genera (**Figure 2D**).

Accordingly, genome-wide phylogenies based on single-copy orthologs showed that most

isolates formed deep-branching, stingless bee-specific lineages, exclusive of any previously

sequenced strain. However, consistent with the results of the 16s rRNA gene analysis, several

of these lineages were related to major phylotypes of the honey bees and bumble bee gut

microbiota (**Figure 3A-D, Supplementary Figure 2-6**) such as *Snodgrassella*, *Gilliamella*,

Lactobacillus Firm-5, *Bifidobacterium*, and *Bombella*. In the case of *Snodgrassella*,

Gilliamella, and *Lactobacillus* Firm-5, the stingless bee-specific lineages formed a

monophyletic clade with lineages of honey bee and bumble bee isolates (**Figure 3A-C** and **3F**).

Notably, in all three cases, the bacteria from stingless bees presented the earliest branching

lineages, i.e. the honey bee and bumble bee gut bacteria diverged after the split from the

stingless bee gut bacteria. While these results suggest that these bacteria have derived from a

common ancestor that was already adapted to social bees, the bacterial phylogenies were

incongruent with current phylogenies of the host, which show that the honey bees (Apini)

diverged before the split of stingless bees (Meliponini) and bumble bees (Bombini) (**Figure

3E**). A different pattern was observed for *Bifidobacterium*. In this case, strains isolated from

stingless bees, honey bees, and bumble bees were not monophyletic. In fact, the stingless bee

isolates belonged to a different clade than the honey bee isolates, while the strains isolated from

bumble bees belonged to either of them (**Figure 3D** and **3G**). Similarly, the two

Acetobacteraceae strains (ESL0695 and ESL0709) were not monophyletic with the honey bee

isolates of the genus *Bombella*, although they belonged to the same Hymenoptera-associated

clade within this family **(Supplementary Figure 3).** This suggests that in both cases,

Bifidobacterium and Acetobacteriaceae, bacteria of distinct lineages have independently

adapted to the gut environment of social bees.

Of the 46 strains selected for sequencing, 13 were not directly related to bacteria previously

isolated from social bees. Most of these isolates matched to minor ASVs in our community

profiling analysis, with the exception of three strains (ESL0689, ESL0687, and ESL0729,

Figure 2A). ESL0689 corresponded to the Enterobacteriaceae ASV19 which dominated the

communities of all three Fv samples (20-35% of the reads per sample). This isolate was situated

on a long branch diverging between the genera *Klebsiella* and *Raoultella* (**Supplementary

Figure 4**). ESL0687 and ESL0729 corresponded to ASVs of the family Streptococcaceae

which were detected in several stingless bee species in our studies as well as in previous studies

(20, 37, 38). They formed a deep-branching sister clade of the flower-associated genus

Floricoccus (**Supplementary Figure 5A**). All three strains seem to be novel species based on

their divergence to previously sequenced bacteria and may represent specialized gut symbionts

of stingless bees.

Stingless bee gut bacteria have diversified into distinct species and reveal a high extent of

genomic diversity.

Stingless bee isolates belonging to the same lineage were often separated by long branches in

our phylogenies indicating substantial genomic divergence (**Figure 3F** and **G**, **Supplementary**

Figures 2-6). This was confirmed by comparing pairwise 16S rRNA gene identity to genome-

wide average nucleotide identity (ANI) between isolates of the same bacterial family. Despite

high similarity in 16S rRNA gene identity, ANI was often <95% suggesting that most lineages

of stingless bee gut bacteria contain several divergent species (**Figure 4A**). For example, the

12 sequenced strains of *Lactobacillus* Firm-5 fell into 8 distinct species-level clusters (i.e. ANI

<95%, **Figure 4B**). A similar pattern was observed for the 17 *Bifidobacterium* strains, which

fell into 14 distinct species-level clusters (**Figure 4C**), as well as the two *Gilliamella* and the

two Streptococcaceae strains, which both also fell below the species-level ANI cut-off.

Notably, some *Bifidobacterium* and *Lactobacillus* Firm-5 strains that were isolated from the

same sample belonged to different ANI clusters, indicating that divergent bacterial species can

co-occur in the same host species and colony. Inversely, strains belonging to the same ANI

cluster were sometimes isolated from different bee species, suggesting that these bacterial

species clusters are not necessarily host-specific (**Figure 4B** and **C**).

Core microbiota members in stingless bees have similar functional capabilities as in honey bees and bumble bees.

To assess the functional potential of stingless bee gut bacteria, we determined the genomic completeness of different metabolic pathways and functions in the genomes of the sequenced strains and compared it to related bacteria isolated from honey bees, bumble bees, or from elsewhere, and which were included in our phylogenomic analysis. We specifically looked at energy and carbon metabolism, amino acid, co-factor, and nucleoside biosynthesis, as well as secretion, motility, and adhesion (**Figure 5**).

Energy and carbon metabolism: Many of the sequenced strains (39/46) - including all Lactobacillaceae, the Bifidobacteria, and the two Streptococcaceae and *Gilliamella* strains – were missing key genes of the TCA cycle and for oxidative phosphorylation, but encoded functions for the breakdown (mostly glycoside hydrolases (GH)), and oxidation of sugars (Embden-Meyerhof-Parnas (EMP), pentose phosphate (PPP), and/or Entner-Doudorrof (ED) pathways). A detailed analyses of the enzymes used for carbohydrate breakdown by the core members *Lactobacillus* Firm-5, *Bifidobacterium*, and *Gilliamella* showed that the stingless bee gut bacteria encoded similar glycoside hydrolase family genes as honey bee and bumble bee isolates (**Figure 6**), including enzyme families for cleaving plant-derived glycans. For example, glycoside families GH5, GH30, GH31, GH42, GH43, and GH51 are involved in the degradation of hemicellulose (27, 30, 52). GH78 can be responsible for the cleavage of rhamnose residues from rutin, a major pollen-derived flavonoid that was demonstrated to be deglycosylated by honey bee isolates of *Lactobacillus* Firm-5 which encode GH78 genes (52). Another example is GH13 which includes neopullulanases and α-amylases for the breakdown of plant-derived starch. Together, these results suggest that stingless bee gut isolates of *Lactobacillus* Firm-5, *Bifidobacterium*, and *Gilliamella* are saccharolytic fermenters that

breakdown pollen or nectar-derived glycans, similar as previously reported for the

corresponding bacteria in the gut of honey bees or bumble bees. Notably, there was substantial

variation in the number and type of glycoside hydrolase family genes between divergent

strains, which goes in line with the extensive genomic diversity detected between stingless bee

gut isolates of these three phylotypes. A complete TCA cycle was only found in the genomes

of Neisseriaceae sp. ESL0693, *Acinetobacter* ESL0695, the Enterobacteriaceae, and the

Erwiniaceae. The same strains also harbored the most complete gene sets for oxidative

phosphorylation. Notably, Neisseriaceae sp. ESL0693 also lacked key genes in the EMP, PPP,

and the ED pathways and contained very few GH family genes (**Figure 5**). This suggests that

this bacterium cannot utilize sugars and obtains energy via aerobic respiration, as previously

found for *Snodgrassella* isolates of honey bees and bumble bees (23).

Amino acid, nucleoside, and co-factor biosynthesis: Differences between stingless bee gut

isolates of different taxonomic groups were also found in terms of their biosynthetic potential.

Strains belonging to the *Lactobacillus* Firm-5 clade were auxotrophic for the production of

most amino acids (i.e. all except for Lys, Gln, and Asn) as well as purine and several co-factors

(e.g. heme, vitamin B6 and B12) (**Figure 5**). Isolates of the Bifidobacteria, Streptococcaceae,

and Leuconostocaceae were also auxotrophic for many co-factors, but for much fewer amino

acids than *Lactobacillus* Firm-5. Interestingly, there was variation in auxotrophies among the

Bifidobacterial strains, especially for the production of purine, NAD+, Thr, Lys, Arg, Gly, and

chorismate. Other strains (such as those of *Gilliamella*, *Snodgrassella*, Acetobacteraceae,

Enterobacteriaceae, and Erwiniaceae) had fewer auxotrophies. Similar biosynthetic capability

profiles were found in related strains included in our phylogenies, which suggests that these

functional profiles are not specific to stingless bee gut bacteria but rather conserved across the

entire phylotype (**Supplementary Figures 2-6**).

Secretion, adhesion, motility: Secretion systems, pili, and flagella were mostly restricted to

the gram-negative bacteria of the isolated strains. Type 1, Type 5, and Type 6 secretion systems

were prevalent across these bacteria, whereas Type 2 and Type 4 secretion systems were only

present in a few strains **(Figure 5)**. Flagella were detected in the two Acetobacteraceae, all

Erwiniaceae, and Enterobacteriaceae strain ESL0689. Tad pili were not detected in any of the

analyzed bacteria, while Type IV pili components were mostly found in Neisseriaceae sp.

ESL0693 and *Acinetobacter* ESL0695, and to some extent also in Orbaceae, Erwiniaceae, and

Enterobacteriaceae ESL0689. Similar gene sets for secretion, adhesion, and motility were also

found in related gut bacteria from honey bees or bumble bees, as shown by the functional

profiles of all strains included in our phylogenies (**Supplementary Figures 2-6**).

Altogether, this first assessment of the gene content of the stingless bee gut bacteria show that

they have similar functional potential as isolates from honey bees and bumble bees suggesting

that they occupy similar ecological niches in the gut across social bees.

Discussion

Previous findings suggested that the core members of the bee gut microbiota have been acquired in a common ancestor of the social bees (20) and possibly co-diversified with their hosts over millions of years (16, 33). However, the lack of genomic data from gut bacteria of stingless bees has limited our view on the evolution of these specialized microbial communities. With the establishment of isolate genomes of diverse stingless bee gut bacteria our study fills an important knowledge gap and provides new lines of evidence that rule out strict co-diversification between the core microbiota members and social bees.

Our genome-wide phylogenies of *Snodgrassella*, *Lactobacillus* Firm-5, and *Gilliamella* revealed that the identified lineages of stingless bee gut bacteria branched off before the divergence of the lineages of honey bee and bumble bee gut bacteria. This pattern is not congruent with the topology of the host phylogeny, in which honey bees diverged before the split of stingless bees and bumble bees (53, 54). The basal split of the stingless bee gut bacterial lineages depends on the rooting of our trees being correct. However, all phylogenies were supported by high bootstrap values at the critical nodes suggesting robust phylogenetic signals in our datasets. Our phylogenies also showed that for both *Gilliamella* and *Snodgrassella*, honey bee isolates were not monophyletic, i.e. some lineages branched before and others after the divergence of the bumble bee clades, which is inconsistent with co-diversification. This was already noted in a previous study (33), and similar results were also obtained for Lactobacillus Firm-5 based on the phylogenetic analysis of single protein-coding genes cloned from different social bee species (20). Finally, our phylogenomic analysis showed that honey bee and stingless bee isolates of *Bifidobacterium* belonged to two separate clades of the Bifidobacteriaceae suggesting that these bacteria have independently adapted to the gut environment of social bees.

Co-diversification can only occur when symbionts and hosts exhibit a high degree of partner fidelity and are transmitted vertically from one generation to the next over many generations (55). While most core microbiota members of social bees indeed seem to have a host-restricted distribution (20), examples of lineages with a broader host range exist as well (56). Moreover, some strains have been experimentally shown to be able to colonize non-native hosts demonstrating that host jumps are possible (20, 27). Our study showed that closely related stingless bee species (i.e. from the same or related genera) can have overlapping community profiles with predominant ASVs being shared across hosts. Similar observations have been made in other 16S rRNA gene profiling studies of stingless bees (20, 37). While such analyses often provide insufficient resolution to discriminate between closely related strains or species, our genomic analyses confirmed that stingless bees isolates of *Bifidobacterium* and *Lactobacillus* do not necessarily cluster by host species. It is possible that the strength of host specificity varies across social bees, or bacterial lineages, depending on the symbiotic function of the gut bacteria, or the hosts' divergence, ecology, or geographic distribution. This would influence the extent to which gut bacteria can co-diversify within certain host lineages. To test this hypothesis, future studies would have to compare the strength of host specificity across multiple bees using representative sets of host species in each of the tree main bee clades.

Another piece of evidence indicating that the microbiota across social bees may be more variable than previously assumed comes from the observation that some of the designated core members were not always detected in the sampled bees. For example, while *Lactobacillus* and *Bifidobacterium* were prevalent across all six stingless bee species sampled in our study, they were absent from the gut microbial communities of some of the previously sampled bee species (15, 17, 37, 57). Likewise, stingless bees of the genus *Melipona* were shown to systematically

lack the two core members *Snodgrassella* and *Gilliamella* (37). Both taxa were also rare across

the four *Melipona* species analyzed in our study. However, three out of 12 analyzed colonies

had high abundances of *Snodgrassella*. This suggests that this bacterium is not completely

absent from this bee genus, but may be occasionally acquired from other bee species, varies in

prevalence depending on season, bee age, or development, or is restricted to only the *Melipona*

species analyzed in our study.

Finally, representatives of core members of the social bee gut microbiota were recently also

found in bees of the distant genus *Xylocopa* (Carpenter bees) (58, 59), suggesting that these

bacteria may have been associated with bees before the emergence of sociality, or that they

have a broader and less-specific distribution across bees than previously suspected.

In summary, our results together with previous findings indicate a rather dynamic evolutionary

background of the core members of the social bee gut microbiota. Rather than having strictly

co-diversified with their hosts extended periods of host-restricted evolution (and likely co-

diversification in some lineages) seem to have been interrupted by host switches, and

independent symbiont gains and losses. Our observation that the stingless bee isolates

repeatedly form the most a sister group to bumble bee and honey bee isolates (for *Gilliamella*,

Snodgrassella, and *Lactobacillus* Firm-5) is intriguing given the host phylogeny. It may

suggest that these core members have an origin in stingless bees and then spread to the other

two groups, especially for *Lactobacillus* where two stingless bee isolates clades split before

the split between bumble bee and honey bee isolates. However, given the large diversity of

social and solitary bees, it is clear that the currently available datasets are insufficient to explain

the distribution and phylogenetic relationships of these gut symbionts across hosts. Broader

samplings of stingless bees, honey bees, and bumble bees, combined with genome-resolved

approaches, are needed to fully understand the diversity, distribution, and evolutionary

trajectories of social bee gut bacteria and to accurately reconstruct the ancestral bee

microbiome composition. Formal analysis should be applied to test for co-diversification.

The importance of sampling biases when assessing patterns of co-diversification between hosts

and their gut bacteria is highlighted by the analyses of Bacteroidaceae gut symbionts of

hominids. While originally reported to have co-diversified with their hosts (60), re-examination

with increased sampling disrupted the co-diversification pattern observed earlier (61). In

contrast, a recent study identified strong signals of parallel evolutionary history between seven

(out of 56 tested) gut bacterial taxa and human populations (62), and phylogenetic congruency

has also been found for certain stinkbug insects and their primary gut symbiont (63). This

demonstrates that co-diversification has occurred between certain gut bacteria and their hosts.

Besides offering new insights into the evolution of the social bee gut microbiota, our genomic

analysis also revealed the functional potential of major gut symbionts of the analyzed stingless

bee species. All isolates of *Lactobacillus* Firm-5, *Bifidobacterium*, and *Gilliamella* carried

genes for the saccharolytic fermentation of diet-derived carbohydrates. In contrast,

Snodgrassella ESL0689 lacked such functions in its genome, but instead harbored genes for

aerobic respiration. These results are consistent with findings from honey bees and bumble

bees (22, 23, 26-28, 30) and hence suggest that the core microbiota members occupy similar

ecological niches across the three groups of social bees.

Another parallel to findings from honey bees and bumble bees was the extensive genomic

divergence present among strains of the core members *Lactobacillus* Firm-5, *Gilliamella*, and

Bifidobacterium, even when isolated from the same host species. Moreover, we found genomic

variation in carbohydrate breakdown and amino acid and nucleoside biosynthesis functions

among these strains. This suggests that the diversification of these bacteria has not only been

driven by isolation into different host species but also by the adaptation to different ecological

niches in the gut, similar as shown for bumble bees and honey bees (28, 33, 35, 64). These

parallels may not be surprising as the dietary preferences of the analyzed stingless bee species

are similar to those of honey bees and bumble bees. In the future, it will be interesting to look

at the functional potential of the core microbiota members in bees with different dietary habits,

such as the vulture bees (38), i.e. stingless bees that feed on raw meat instead of pollen yet

share a subset of the core members with other social bees.

Some of the isolate genomes we sequenced in our study did not come from any of the core

members of the social bee gut microbiota. They may present transient community members,

opportunistic pathogens, or host-specific gut symbionts with complementary functions to the

core microbiota. Therefore, the established genomes present an important resource for future

research. A particular strain that drew our attention was Enterobacteriaceae ESL0689, as it

belonged to an ASV that was present at high relative abundance in all three colonies of *Fv.*

ESL0689 harbored a complete TCA cycle and a respiratory chain and could also synthesize

most amino-acids and co-factors indicating a similar metabolic niche as *Snodgrassella* (23).

In conclusion, our study provides new insights into the evolution of the social bee gut

microbiota and presents a first step in characterizing the functional potential of major gut

bacteria present in stingless bees. However, given the large diversity of stingless bees with

hundreds of different species distributed throughout the tropical and subtropical regions of the

world, it is clear that our study only presents the starting point in characterizing their genomic

diversity and functional potential. More detailed studies and larger genomic survey combined

with experimental analyses will be needed to understand their evolution and assess their impact

on the host.

Figures

A

B

Figure 1. Community analysis of the gut microbiota of stingless bees. (A) 16S rRNA

gene-based community profiles of the gut microbiota of three colonies of six different

stingless bee species collected in Brazil. Fv – *Frieseomelitta varia,* Ms – *Melipona*

seminigra, Ml – *Melipona lateralis*, Mf – *Melipona fuliginosa*, Mi – *Melipona interrupta*, Sp

– *Scaptotrigona polysticta*. Relative abundance of ASVs is shown. ASVs are ordered and

colored at the family level according to the legend, families of core members are displayed in

bold. ASVs with <1% relative abundances are summed up as 'others' and shown in grey. **(B)**

NMDS based on ASV relative abundance (Bray-Curtis dissimilarity) across 136 16S rRNA

gene amplicon sequence samples including data from a previous study (20) and our study.

Figure 2. Bacterial strains isolated from the gut of six stingless bee species from Brazil.

(A) Maximum likelihood tree was inferred from nearly complete sequences of the 16S rRNA

gene of each isolate. Strain names of isolates for which we sequenced the genome are indicated

in bold. The genome size of each sequenced strain is noted next to it. The isolate host is

indicated as follows: Fv – *Frieseomelitta varia,* Ms – *Melipona seminigra*, Ml – *Melipona*

lateralis, Mf – *Melipona fuliginosa*, Mi – *Melipona interrupta*, Sp – *Scaptotrigona polysticta*.

Closest hit indicates best BlastN hit of the 16S rRNA gene sequence against the 16S ribosomal

RNA (NCBI: Bacteria and Archaea type strains) database. Colored circles indicate if the strain

of the best hit was isolated from the gut of a bumble bee, a honey bee, or from elsewhere. Bar

plots indicate % identity of the best BlastN hit. The matching ASV and its average relative

abundance across the nineteen analyzed stingless bee species are indicated. The names of the

stingless bee species from our study are given in bold. Note that many isolates matched the

same ASV. **(B)** Relative abundances of matching ASVs in each of the 18 samples of the six

stingless bees species sampled in our study, i.e. representativeness of the isolates in the

amplicon data. **(C)** Distribution of the isolated strains based on best 16S rRNA gene identity

to closest reference species. These are the same values as shown in Figure 2A. The dashed lines

indicate 95% and 97% identity corresponding to the genus and species threshold, respectively.

(D) Distribution of the isolated strains based on pairwise ANI with the closest reference

genome publicly available. The dashed line indicates the species threshold at 95% ANI.

Figure 3. Isolates of stingless bee gut bacteria present novel species and belong to deep-branching phylogenetic lineages. (A-D) Simplified genome-wide maximum likelihood phylogenies of *Snodgrassella*, *Lactobacillus* Firm-5, *Gilliamella*, and *Bifidobacterium* based on single copy gene orthologs. All branches shown are supported by >95/100 bootstrap replicates. Most of the branches have been collapsed. **(E)** Dendrogram depicting the topology of the social bee phylogeny (adapted from (53, 54))**. (F)** Detailed genome-wide phylogeny of the genus *Lactobacillus* with bacteria belonging to the social bee-specific phylotype *Lactobacillus* Firm-5 highlighted in different colors according to the host species/group. The

maximum likelihood tree was computed on the concatenated amino acid sequences of 355

single copy core genes using the substitution model LG+F+I+G4. **(G)** Detailed genome-wide

phylogeny of the genus *Bifidobacterium* with bacteria belonging to social bee-specific clades

highlighted in different colors according to the host species/group. The maximum likelihood

tree was computed on the concatenated amino-acid sequences of 151 single copy core genes

using the substitution model LG+F+I+G4.

Figure 4. Genomic divergence among isolates with high 16S rRNA gene similarity. (A) The average nucleotide identity (ANI) versus 16S rRNA gene identity was plotted for each pair of isolate genomes belonging to the same bacterial family. The two vertical dashed bars indicate the ANI thresholds of 95% and 83% demarcating the intra-species and inter-species ANI values, respectively (65). The horizontal dashed lines delineate the commonly used species (>97%) and genus (>95%) thresholds for the 16S rRNA gene similarity. Average nucleotide identity heatmaps of **(B)** *Lactobacillus* Firm-5 isolates and **(C)** *Bifidobacterium* isolates. The 95% ANI clusters are delimited in white. The isolate host is indicated, Fv – *Frieseomelitta varia,* Ms – *Melipona seminigra*, Ml – *Melipona lateralis*, Mf – *Melipona fuliginosa*, Mi – *Melipona interrupta*, Sp – *Scaptotrigona polysticta*.

Figure 5. Metabolic capabilities of the sequenced stingless bee gut isolates. (A) The heatmap indicates the genomic completeness of major metabolic pathways and functions related to energy and carbon metabolism (EMP: Embden–Meyerhof–Parnas), the biosynthesis of amino-acid, co-factors, and nucleoside, as well as secretion, adhesion, and motility across the sequenced bacterial isolates. The isolates are ordered into core and accessory members based on whether they are related to bacteria isolated from other social bees. **(B)** Total number of glycoside hydrolase (GH) and polysaccharide lyase (PL) genes in each isolate.

Figure 6. Carbohydrate-active enzyme profiles for *Bifidobacterium, Lactobacillus* Firm-5 and

***Gilliamella* isolates.** Distribution of genes in each GH and PL families respectively for *Bifidobacterium,*

Lactobacillus Firm-5 and *Gilliamella* isolates. The isolates were sorted according to their position in

the phylogenies, the hosts are indicated by the background colors. Glycoside hydrolases families

mentioned in the text are outlined with a red square. The color scale indicates the number of genes in

each family.

Methods

Bee sampling

Bees were collected from three different nests of each of the following six Meliponini species in February and March 2019: *Frieseomelitta varia*, *Scaptotrigona polysticta*, *Melipona fuliginosa*, *Melipona interrupta*, *Melipona lateralis* and *Melipona seminigra*. All nests were located in a rural meliponary in the vicinities of Iranduba municipality (Iranduba-AM, Brazil, 3°10'52.7"S 60°07'08.5"W), and the sampling was carried out under the SISGEN collection permission A256E82. Bees were sampled at the entrance of each nest and immediately immobilized by cooling at 4 °C. Then the entire gastrointestinal tract was dissected, and the hindgut separated from the anterior gut parts. Hindguts of bees from the same nest were pooled in two separate tubes. One pool per nest was mixed with 1x PBS and glycerol, homogenized using bead-beating, and subsequently cryopreserved at -80°C. This pool was used for bacterial culturing as described below. The other pool was cryopreserved at -80°C without homogenization and was subsequently used for DNA extraction and 16S rRNA gene analysis. For the sample used for sequencing, 20 bees per nest were pooled, while for the sample for culturing, only 3 bees per nest were pooled. For *Frieseomelitta varia* and *Scaptotrigona polysticta*, we pooled 40 and 10 bees, respectively, due to the small size of these bee species.

Bacterial culturing

For establishing a culture collection of primary isolates from the gut of the six sampled bee species, serial dilutions of the cryopreserved homogenates were plated on eight different media: CBA (Columbia Blood Agar supplemented with 5% defibrinated sheep blood (Thermofisher)), MRSA (De Man, Rogosa and Sharpe agar) supplemented with Fructose (2%) and L-Cysteine (0.1%), MRSA supplemented with Mannitol (2%), Chocolate Agar, TSA (tryptone soya agar), TYG (tryptone glucose yeast extract Agar), GC, LBA (Luria Bertani agar)

without NaCl, BHIA (brain heart infusion agar) and SDA (Sabouraud dextrose agar). Plates

were incubated in two different conditions: in a microaerobic 5% CO_2-enriched atmosphere

and in an anaerobic chamber (72% N_2, 8% H_2, 20% CO_2), both at 34°C. After 2-7 days of

incubation, colonies of different size and appearance were picked and re-grown on the same

media and culturing conditions. Cryo-stocks of bacterial strains of interest were prepared by

harvesting bacterial biomass in liquid media corresponding to the solid growth media and

supplemented with 20% glycerol. For DNA isolation, bacteria were grown from the stocks,

and a single colony was picked and re-grown on fresh media before harvesting bacterial

biomass.

Genotyping

All colonies that were selected for culturing were genotyped by PCR and Sanger sequencing

of a 16S rRNA gene fragment. To this end, a small amount of bacterial material was transferred

to a lysis buffer (Tris-HCl 1M pH=7.5, EDTA 0.5M, SDS 10%) containing 2.5 µl lysozyme

(20mg/mL) and 2.5 µl proteinase K (20mg/mL), and incubated for 10 min at 37°C, 20 min at

55°C, and 10 min at 95°C. PCR was performed with universal bacterial primers that amplify

the V1-V5 region of the 16S rRNA gene (27F - AGRGTTYGATYMTGGCTCAG and 907R-

CCGTCAATTCMTTTRAGTTT) using the following reagents and thermocycler program:

initial denaturing at 94°C for 5 min, followed by 32 cycles of denaturing at 94°C for 30 sec,

annealing at 56°C for 30 sec, and extension 72°C for 1 min, and a final extension at 72°C for

7 min. PCR results were checked on a 1% agarose gel. PCR reactions selected for Sanger

sequencing were purified using ExoSAP-IT™ (1µl ExoSAP 5x, 4µl ddH_2O) with the

thermocycler program: 30 min at 37°C followed by 15 min at 80°C. Purified samples were

then sent to Eurofins® for sequencing. Sanger sequences were analyzed with Geneious suite

(Geneious®) and compared to GenBank at NCBI using BLAST tools (66).

DNA isolation and genome sequencing

DNA isolation for Illumina sequencing was carried out using a customized SPRI bead-based

extraction method or the FastPure Bacteria DNA Isolation Mini Kit (Vazyme). For the SPRI-

bead method, bacteria were harvested and resuspended in tubes containing 200 mg of 0.1 mm-

sized acid washed glass beads and 200 µl of TER buffer (10 mM Tris-HCl, 1 mM EDTA, 100

µg /ml RNAse A, pH8.0). Samples were homogenized using a FastPrep-25 5G instrument (2

rounds of 30 sec with the power set to 6) and subsequently centrifuged at max speed for 10

min at room temperature. 40 µl of SPRI beads were added to 100 µl of supernatant,

immediately mixed thoroughly by repeated pipetting (>20x) and incubated for 5 min at room

temperature. After placing the tubes on a magnet stand the liquid was removed and discarded

and the beads washed two times with 200 µl 80% ethanol. After air drying the tubes on the

magnetic stand 22 µl of 5 mM Tris-HCl pH8 was added. For isolating bacteria with the

FastPure Bacteria DNA Isolation Mini Kit the manufacturer's protocol for gram-positive

bacteria was followed.

DNA isolation for Oxford Nanopore Technologies (ONT) sequencing was carried out using a

custom DNA extraction protocol for Gram-positive bacteria. Tubes were prepared with glass

beads and 160 µl of buffer P1 (Qiagen). Then bacteria were harvested and resuspended in these

tubes by intensive vortexing. Lysozyme (20 µl, 100 mg/ml) was added and after gentle mixing,

tubes were incubated at 56°C with shaking at 600 rpm for 30 min. Then 4 µl RNase A (100

mg/ml) were added to the tubes followed by 150 µl of buffer AL (lysis buffer, Qiagen). After

mixing by vortexing, tubes were incubated in a thermomixer (37°C, 900 rpm) for 20 min. Tubes

were centrifuged 10 min at 14'000 rpm to pellet the beads, the supernatant was transferred to

new tubes with 35 µl Na-acetate and 270 µl isopropanol and mixed by inverting. Following an

incubation of 1 h at 4°C, DNA was pelleted by centrifugation at 14'000 rpm for 10 min at

25°C. The supernatant was discarded, and the pellets were washed with 1 ml EtOH 80%. After

a second centrifugation (14'000 rpm, 10 min at 25°C) the ethanol was removed, and the pellet

left to dry at room temperature. DNA pellets were solubilized with 50 µl TER (10 mM Tris-

HCl, 1 M EDTA, pH 8.0, 2 mg/ml RNase A) and tubes incubated at 37°C for 15 min. The

solution was then transferred to PCR tubes. 40 µl NGClean beads were added, and the solution

mixed by repeated pipetting. After a 5 min incubation, PCR tubes were placed on magnetic

stands. When the solutions were clear, the liquid was removed, and the beads washed two times

with 200 µl of 80% EtOH. Upon complete drying, beads were resuspended with 22 µl of 5mM

Tris-HCl (pH 8.0). Finally, tubes were placed again on the magnetic stand and when the

solution were clear, the supernatant was transferred to new 1.5 ml Eppendorf tubes.

Illumina sequencing libraries were prepared using the Nextera DNA Flex Library Preparation

kit following instructions of the Illumina Reference Guide. This was followed by indexing,

dilution, and denaturation according to the 'Index Adapters Pooling Guide' and the 'MiniSeq

System Denature and dilute Libraries guide'. Libraries were checked and quantified using a

dsDNA fluorescent dye, before loading them on a MiniSeq High-output flowcell (150PE).

ONT libraries were prepared using the ligation-based approach (LSK109). The sequencing was

conducted on a ONT MinION for a duration of 72 hours with the high-accuracy base-calling

using Guppy (*v5.0.11*).

Genome assembly

Forty-six isolates from the stingless bees were sequenced with Illumina MiniSeq (150PE). Raw

reads were checked using FastQC *v0.11.9* (67) and trimmed by Trimmomatic v0.39 (68) using

parameters: PE -phred33 AllIllumina-PEadapters.fa:3:25:7 LEADING:9 TRAILING:9

SLIDINGWINDOW:4:15 MINLEN:60. For *de novo* assembly we used SPAdes (–careful

option, *v3.15.2*) (69). For assembly quality control, reads were mapped back against the

assembly with BWA *v0.7.17 (70)*, samtools *v1.12* (71) and plotted with R *v4.1.1*. Genomes

completeness was evaluated with checkM *v1.0.13* (72).

Additionally, 26 isolates were also sequenced with Oxford Nanopore to produce long reads.

Nanopore long reads were filtered with Filtlong *v0.2.0* (https://github.com/rrwick/Filtlong) for

a minimum length of 7'000 and minimum mean q-score of 10 (min_length 7000, min_mean_q

10, length_weight 10). ONT-based assemblies were computed with Flye *v2.7.1* (73) over 5

iterations. Graphmap *v0.5.2* (74) and Racon *v1.0.1* (75) were used to perform two rounds of

polishing. Finally, the racon-corrected assembly and Illumina reads were fed to Pilon *v1.24*

(76) for single base and indels corrections.

Genome annotation and analysis

Genomes were annotated with Prokka *v 1.13* (77). Phylogenies were computed for the bacterial

families for which we had an isolate. For each family, we identified a set of closely strains and

outgroup taxa and retrieved their genomes from NCIB and IMG/Mer (78). All genomes were

re-annotated with Prokka to ensure annotations consistency. Gene orthology was inferred with

OrthoFinder *v2.3.8 (79)*. Single copy ortholog genes were selected and their amino acid

sequences were aligned (mafft *v7.453) (80)*. An in-house script was used to trim the alignments

by removing positions with more than 50% gaps and sequences belonging to the same genome

were concatenated to produce a core gene alignment. This alignment was used to infer the

maximum likelihood phylogeny using IQTree (*v1.7.beta17* , -st AA -bb 1000 -seed 12345 -m

TEST) (81). For each phylogeny, the best evolutionary model was chosen according to BIC:

LG+I+G4 – Enterobacteriaceae ; *LG+F+I+G4* – Acetobacteraceae, *Bifidobacterium*,

Lactobacillus, Leuconostocaceae, Moraxellaceae, Neisseriaceae and Streptococcaceae ; and

JTTDMut+F+I+G4 – Orbaceae. Branch support of the trees were inferred using 1000 ultrafast

bootstrap (UFBoot) repetitions. Clades can be trusted when UFBoot values are >95%.

16S rRNA gene based community profiling

Region V4 of the 16S rRNA gene was amplified using the primers 515F-Nex and 806R-Nex

(TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGTGCCAGCMGCCGCGGTAA,

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGGACTACHVGGGTWTCTAAT

). The primers include Nextera XT index adapter sequences and the primers for the 16S rRNA

V4 region (82) described by Kešnerová et al. (2020). The two step PCR was performed as

following: the first PCR was performed using 12.5 µL of 2× Phanta Max Master Mix (Vazyme,

Nanjing, China), 5 µl of MilliQ water, 2.5 µl of each primer (5 µM), and 2.5 µl of template

DNA for a total volume of 25 µl. The PCR program started with a denaturation step at 98°C

for 30sec, followed by 25 cycles of amplification (10 sec at 98°C, 20 sec at 55°C, and 20 sec

at 72°C) and a 5 min final extension step at 72°C. The PCR products were verified by 2%

agarose gel electrophoresis, then purified with clean NGS purification-beads in a 1:0.8 ratio of

PCR product to beads and eluted in 27 µl Tris (10mM, pH 8.5). A second PCR step was

performed to append the unique dual indexes to each sample in a total volume of 25 µl using

12.5 µL of 2× Phanta Max Master Mix (Vazyme, Nanjing, China), 5 µl of MilliQ water, 2.5 µl

of Nextera XT index primers 1 and 2 (Nextera XT Index kit, Illumina) and 2.5 µl of templated

DNA. The PCR program started with a denaturation step at 95°C for 3 min, followed by 8

cycles of amplification (30 sec at 95°C, 30 sec at 55°C, and 30 sec at 72°C), and a 5 min final

extension step at 72°C. The libraries were then cleaned usingClean NGS purification beads

(1:1.1 ratio of PCR product to beads) and were eluted in 27.5 µl Tris (10mM, pH 8.5). Prior to

sequencing, the PCR product concentrations were quantified by PicoGreen and pooled in

equimolar concentrations; the negative controls and blank extractions were pooled in equal

volume. Sequencing was performed on an Illumina MiSeq sequencer (2 x 250bp) by the

Genomic Technology Facility of the University of Lausanne. We followed DADA2 (83)

pipeline to analyze the sequencing data. For the first part of the analysis, we executed the

pipeline only on the 18 samples from our study. To control for possible contaminants, we used

blank extractions and water as negative controls during the PCR. For the second part of the

analysis, where we combined our dataset with the one from (20), we executed the pipeline

independently a second time. Sequence quality control was performed with DADA2 integrated

function *plotQualityProfile*. Data from the three sequencing runs were processed

independently for the filtering, dereplication, the error rates calculation and the sample

inference. After merging the denoised forward and reverse pairs, we merged the three sequence

tables (*mergeSequenceTables*) and we applied the *collapseNoMismatch* function to unite

similar ASVs with shifts or length variation. We selected sequences in range 250-256bp and

removed chimeric sequences. Silva non-redundant SSU database v138.1 (83) was used for the

taxonomic assignment of the ASVs. We filtered out 62 ASVs for which the taxonomic

assignment was matching 'Eukaryota', 'Chloroplast' or 'Mitochondria'. Finally, we removed

samples with less than 5,000 reads. Non-metric MultiDimensional Scaling plots were

computed in R using Bray-Curtis distances (Phyloseq: *ordinate*). The adonis package was used

to compute permanovas.

Functional Profiler

We developed a genomic profiler using several software to annotate the genomes and compute

the completeness of key pathways and functions. The genomes were re-annotated with

GhostKOALA (84) to get KEGG annotations of genes. We created a set of rules to define the

steps of selected energy and metabolism pathways, and co-factors/nucleoside biosynthesis

pathways and computed their completeness. In brief, an in-house script parsed the KEGG

annotations of the genomes and for each pathways evaluated the completeness of each step.

The pathways completeness was then summarized by counting the number of steps present

relative to total number of steps needed. To obtain the completeness of amino acid biosynthesis

pathways, we used GapMind (85). Secretion systems and related appendages were detected in

the genomes by MacSyFinder's TXSScan module (86). Finally, we ran dbCan (87) for the

annotation of carbohydrate-active enzymes.

Acknowledgements

We would like to thank Julien Marquis and the team from the Lausanne Genomics Technology Facility and Alban Ramette and his team from the University of Berne for carrying out the Illumina sequencing and Oxford Nanopore sequencing, respectively. We also would like to thank the 'Ecole de Biologie' of the University of Lausanne for their financial support to sequence the presented genomes in the context of the course "Sequence-a-genome" (SAGE) as part of the Master of Science in Molecular Life Science. Finally, we would like to thank all SAGE students of the years 2019-2020 and 2020-2021 for their contribution to this course: Yami Ommar Arizmendi Cárdenas, Samuel Aubert, Alec Auston, Fabrice Battiston, Etienne Bellani, Valentina Benigno, Valentin Borgeat, Alessandro Brandulas Cammarata, Marion Brechet, Marine Bugnon, Jessica Burnier, Théo Cavinato, Giacomo Ceracchini, Jérémy Cherbuin, Lucas Culebras, Audrey Daina, Hugues de Villiers de la Noue, Joe Dickinson, Alexandre Dudt, Elise Eray, Sara Ezzat, Christopher Forbes-Jaeger, Auriane Form, Léo Franchi, Pablo Guridi Fernández, Manon Henna, Aya Iizuka, Nicolas Jacquemin, Richie Kalusivikako, Maroussia Liechti, Simon Maréchal, Achille Mariotti, Aoife Mc Nally, Mam Malick Sy Ndiaye, Jade Nicolet, Astrid Oliva, Claire Paltenghi, Priyanka Parmar, Nicolas Pellaton, Katharina Pfaller, Brenda Ríos-Ochoa, Eric Risse, Daniel Rodriguez, Artemiy Saukin, Matthieu Simeoni, Miloš Stojanov, Marina Sudário, François Sutter, James Tan, Emanuele Tettamanti, Jamille Viray, Grazia Vizzarro and Chaymae Ziyani. This work was funded by the ERC Starting Grant (MicroBeeOme), the NCCR Microbiomes, a National Centre of Competence in Research, funded by the Swiss National Science Foundation (grant no. 180575) and a Swiss National Science Foundation Spirit grant (grant no. IZSTZ0_189496) to P.E.. F.M. was funded by the Swiss National Science Foundation (grant nr 315230_184908: SOMETALP).

738 **Data and materials availability:**

739 The bacterial genomes and sequencing data are available in the NCBI's BioProjects

740 PRJNA906295 and PRJNA897721. Scripts used for the amplicon sequencing analysis, the

741 phylogenies and the metabolic profiler can be found on GitHub

742 https://github.com/gsartonl/Publication_Sarton-Loheac_2022.

Supplementary Figure 1. (A) NMDS based on ASV relative abundance in the 18 samples

collected from six different stingless bee species in Brazil. The samples clustered by bee

genus (PERMANOVA**:** 'Genus' pseudo F=11.05, p-value< 0.001). **(B)** Community profiles

of the gut microbiota of 136 samples from 19 different stingless bee species collected in

Brazil, Malaysia, and Australia. Relative abundance of ASVs is shown. ASVs are ordered

and colored at the genus level according to the legend. ASVs with <1% relative abundances

are summed up as 'others' and shown in grey. This plot includes the datasets from stingless

bees from (20) and our study. The samples are ordered by bee species : Aa – *Austroplebeia*

australis, Fv – *Frieseomelitta varia*, Gt – *Geniotrigona thoracica* , Hi – *Heterotrigona itama*,

Mf – *Melipona fuliginosa*, Mi – *Melipona interrupta*, Ml – *Melipona lateralis*, Ms –

Melipona seminigra, Ps – *Paratrigona subnuda*, Ph – *Partamona helleri*, Pd – *Plebeia*

droryana, Sp – *Scaptotrigona polysticta*, Tc – *Tetragonula carbonaria*, Td – *Tetragonula*

davenporti, Tf – *Tetragonula fuscobalteata*, Tl – *Tetragonula laeviceps*, Th – *Trigona*

hyalinata, Ts – *Trigona spinipes*. **(C)** Number of shared ASVs. For each family, we reported

how many ASVs were observed (indicated by the dot size) in how many bee species.

Families with an average abundance <1% are grouped as 'Other'. The sum of species-

specific or shared (by two or more bee species) ASVs is reported on the right side. Core

members families are highlighted in bold. **(D)** Relative abundances of the shared ASVs

across the 19 stingless bee species (samples from the same species were pooled).

Supplementary Figure 2. (A) Core genome phylogeny of the Neisseriaceae including

phylotype *Snodgrassella*. The maximum likelihood tree was computed with the amino-acid

sequences of 428 concatenated single copy core genes using LG+F+I+G4 substitution model.

Bee isolates form a monophyletic clade. The heatmap indicates the genomic completeness of

major metabolic pathways and functions related to energy and carbon metabolism (EMP:

Embden–Meyerhof–Parnas), the biosynthesis of amino-acid, co-factors, and nucleoside, as

well as secretion, adhesion, and motility across the sequenced bacterial isolates. **(B)** Core

genome phylogeny of the Orbaceae including *Gilliamella* phylotype. The maximum

likelihood tree was computed on the concatenated amino-acid sequences of 782 single copy

core genes using JTTDCMut+F+I+G4 substitution model. The heatmap indicates the

778   genomic completeness of major metabolic pathways and functions related to energy and

779   carbon metabolism (EMP: Embden–Meyerhof–Parnas), the biosynthesis of amino-acid, co-

780   factors, and nucleoside, as well as secretion, adhesion, and motility across the sequenced

781   bacterial isolates.

Supplementary Figure 3. Core genome phylogeny of the Acetobacteraceae. The maximum

likelihood tree was computed on the concatenated amino-acid sequences of 303 single copy

core genes using LG+F+I+G4 substitution model. Bee isolates are distributed in two distant

monophyletic clades. The heatmap indicates the genomic completeness of major metabolic

pathways and functions related to energy and carbon metabolism (EMP: Embden–Meyerhof–

Parnas), the biosynthesis of amino-acid, co-factors, and nucleoside, as well as secretion,

adhesion, and motility across the sequenced bacterial isolates.

Supplementary Figure 4. Core genome phylogeny of the Enterobacteriaceae. The maximum

likelihood tree was computed on the concatenated amino-acid sequences of 303 single copy

core genes using LG+F+I+G4 substitution model. Metabolic heatmap of the

Enterobacteriaceae isolates, the genomes are ordered according to the phylogeny. The

heatmap indicates the genomic completeness of major metabolic pathways and functions

related to energy and carbon metabolism (EMP: Embden–Meyerhof–Parnas), the

biosynthesis of amino-acid, co-factors, and nucleoside, as well as secretion, adhesion, and

motility across the sequenced bacterial isolates.

Supplementary Figure 5. (A) Core genome phylogeny of the Streptococcaceae including

Floricoccus isolates. The maximum likelihood tree was computed on the concatenated

amino-acid sequences of 303 single copy core genes using LG+F+I+G4 substitution model.

Metabolic heatmap of the Streptococcaceae isolates, the genomes are ordered according to

the phylogeny. The heatmap indicates the genomic completeness of major metabolic

pathways and functions related to energy and carbon metabolism (EMP: Embden–Meyerhof–

Parnas), the biosynthesis of amino-acid, co-factors, and nucleoside, as well as secretion,

adhesion, and motility across the sequenced bacterial isolates. **(B)** Core genome phylogeny of

the Leuconostocaceae. The maximum likelihood tree was computed on the concatenated

amino-acid sequences of 303 single copy core genes using LG+F+I+G4 substitution model.

Bee isolates are distributed in two distant monophyletic clades. Metabolic heatmap of the

Leuconostocaceae isolates, the genomes are ordered according to the phylogeny. The

heatmap indicates the genomic completeness of major metabolic pathways and functions

related to energy and carbon metabolism (EMP: Embden–Meyerhof–Parnas), the

biosynthesis of amino-acid, co-factors, and nucleoside, as well as secretion, adhesion, and

motility across the sequenced bacterial isolates. **(C)** Core genome phylogeny of the

Moraxellaceae. The maximum likelihood tree was computed on the concatenated amino-acid

sequences of 303 single copy core genes using LG+F+I+G4 substitution model. Metabolic

heatmap of the Moraxellaceae isolates, the genomes are ordered according to the phylogeny.

The heatmap indicates the genomic completeness of major metabolic pathways and functions

related to energy and carbon metabolism (EMP: Embden–Meyerhof–Parnas), the

biosynthesis of amino-acid, co-factors, and nucleoside, as well as secretion, adhesion, and

motility across the sequenced bacterial isolates**.**

Supplementary Figure 6. (A) Genome-wide phylogeny of the genus *Lactobacillus* including

bacteria belonging to the social bee-specific phylotype *Lactobacillus* Firm-5 highlighted in

different colors according to the host species/group. The maximum likelihood tree was

computed on the concatenated amino acid sequences of 355 single copy core genes using the

substitution model LG+F+I+G4. Metabolic heatmap of the *Lactobacillus* isolates, the

genomes are ordered according to the phylogeny. The heatmap indicates the genomic

completeness of major metabolic pathways and functions related to energy and carbon

metabolism (EMP: Embden–Meyerhof–Parnas), the biosynthesis of amino-acid, co-factors,

and nucleoside, as well as secretion, adhesion, and motility across the sequenced bacterial

isolates. **(B)** Genome-wide phylogeny of the genus *Bifidobacterium* with bacteria belonging

to social bee-specific clades. The maximum likelihood tree was computed on the

concatenated amino-acid sequences of 151 single copy core genes using the substitution

model LG+F+I+G4. Metabolic heatmap of the *Bifidobacterium* isolates, the genomes are

ordered according to the phylogeny. The heatmap indicates the genomic completeness of

major metabolic pathways and functions related to energy and carbon metabolism (EMP:

Embden–Meyerhof–Parnas), the biosynthesis of amino-acid, co-factors, and nucleoside, as

well as secretion, adhesion, and motility across the sequenced bacterial isolates.

Supplementary Tables legends

Supplementary Table 1. Gut Homogenates metadata of the 18 stingless bee hives. For each

bee species, individuals were collected from three hives in a rural meliponary

Supplementary Table 2.

ASV table of our 18 samples. For each ASV, the assigned taxonomy and the number of reads

in each sample are displayed. In the sheet RelAbundance, read counts were transformed to

relative abundance for each sample. In the Absence_Presence, 1 indicates that at least one

read matched the ASV, 0 indicates that the ASV was not found in the sample. Finally, the

Family_relAbundaces tab is a summary of the ASVs abundance from each taxonomic family

per sample.

Supplementary Table 3. ASV table. For each ASV, the assigned taxonomy and the number of

reads in each sample are displayed. This analysis groups our 18 samples and samples from

W. K. Kwong et al., Dynamic microbiome evolution in social bees. Sci Adv 3, 905 e1600513

(2017). In the sheet RelAbundance, read counts were transformed to relative abundance for each

sample. In the Absence_Presence, 1 indicates that at least one read matched the ASV, 0 indicates

that the ASV was not found in the sample. Finally, the Family_relAbundaces tab is a summary of the

ASVs abundance from each taxonomic family per sample.

Supplementary Table 4: Stingless bee strain collection. For each isolate, the bacterial

species, bee host are indicated. We blasted the 16S rRNA gene sequences against the NCBI

nr database and extracted for each isolate the reference type strain ID and the % of identity

869   between the type strain and our isolate 16S rRHA gene sequences. The Sequenced_Isolates

870   tab, indicates which strains were sequenced.   53

871
872
873

References

1.   P. J. Ascher J, Discover Life bee species guide and world checklist (Hymenoptera:
Apoidea: Anthophila). http://www.discoverlife.org/mp/20q?guide=Apoidea_species (2018).

2.   C. D. Michner, *The bees of the world* (The Johns Hopkins University Press, Baltimore,
2007).

3.   B. N. Danforth, S. Cardinal, C. Praz, E. A. Almeida, D. Michez, The impact of molecular
data on our understanding of bee phylogeny and evolution. *Annu Rev Entomol* **58**,
57-78 (2013).

4.   G. Bonilla-Rosso, P. Engel, Functional roles and metabolic niches in the honey bee
gut microbiota. *Curr Opin Microbiol* **43**, 69-76 (2018).

5.   W. K. Kwong, N. A. Moran, Gut microbial communities of social bees. *Nature reviews.*
Microbiology **14**, 374-384 (2016).

6.   K. Raymann, N. A. Moran, The role of the gut microbiome in health and disease of
adult honey bee workers. *Curr Opin Insect Sci* **26**, 97-104 (2018).

7.   Z. Zhang *et al.*, Honeybee gut Lactobacillus modulates host learning and memory
behaviors via regulating tryptophan metabolism. *Nat Commun* **13**, 2037 (2022).

8.   J. Liberti *et al.*, The gut microbiota affects the social network of honeybees. *Nat Ecol*
Evol **6**, 1471-1479 (2022).

9.   L. Li *et al.*, Gut microbiome drives individual memory variation in bumblebees. *Nat*
Commun **12**, 6588 (2021).

10.   K. Raymann, Z. Shaffer, N. A. Moran, Antibiotic exposure perturbs the gut microbiota
and elevates mortality in honeybees. *PLoS Biol* **15**, e2001861 (2017).

11.   E. V. S. Motta *et al.*, Oral or Topical Exposure to Glyphosate in Herbicide Formulation
Impacts the Gut Microbiota and Survival Rates of Honey Bees. *Appl Environ Microbiol*
86 (2020).

12.   E. V. S. Motta, K. Raymann, N. A. Moran, Glyphosate perturbs the gut microbiota of
honey bees. *Proc Natl Acad Sci U S A* **115**, 10305-10310 (2018).

13.   D. P. Cariveau, J. Elijah Powell, H. Koch, R. Winfree, N. A. Moran, Variation in gut
microbial communities and its association with pathogen infection in wild bumble
bees (Bombus). *ISME J* **8**, 2369-2379 (2014).

14.   D. L. Cox-Foster *et al.*, A metagenomic survey of microbes in honey bee colony
collapse disorder. *Science* **318**, 283-287 (2007).

15.   M. A. Hall *et al.*, Temporal changes in the microbiome of stingless bee foragers
following colony relocation. *FEMS Microbiol Ecol* **97** (2020).

16.   H. Koch, D. P. Abrol, J. Li, P. Schmid-Hempel, Diversity and evolutionary patterns of
bacterial gut associates of corbiculate bees. *Mol Ecol* **22**, 2028-2044 (2013).

17.   S. D. Leonhardt, M. Kaltenpoth, Microbial communities of three sympatric Australian
stingless bee species. *PLoS One* **9**, e105718 (2014).

18.   J. Li *et al.*, Two gut community enterotypes recur in diverse bumblebee species. *Curr*
Biol **25**, R652-653 (2015).

19.   N. A. Moran, A. K. Hansen, J. E. Powell, Z. L. Sabree, Distinctive gut microbiota of
honey bees assessed using deep sampling from individual worker bees. *PLoS One* **7**,
e36393 (2012).

20.   W. K. Kwong *et al.*, Dynamic microbiome evolution in social bees. *Sci Adv* **3**,
e1600513 (2017).

21. P. Engel *et al.*, Standard methods for research on Apis mellifera gut symbionts.
Journal of Apicultural Research **52** (2013).
22. L. Kesnerova *et al.*, Disentangling metabolic functions of bacteria in the honey bee
gut. *PLoS Biol* **15**, e2003467 (2017).
23. W. K. Kwong, P. Engel, H. Koch, N. A. Moran, Genomics and host specialization of
honey bee and bumble bee gut symbionts. *Proc Natl Acad Sci U S A* **111**, 11509-
11514 (2014).
24. M. A. Aizen, L. A. Garibaldi, S. A. Cunningham, A. M. Klein, Long-term global trends in
crop yield and production reveal no current pollination shortage but increasing
pollinator dependency. *Curr Biol* **18**, 1572-1575 (2008).
25. N. Gallai, J. M. Salles, J. Settele, B. E. Vaissiere, Economic valuation of the
vulnerability of world agriculture confronted with pollinator decline. *Ecol Econ* **68**,
810-821 (2009).
26. F. Bottacini *et al.*, Bifidobacterium asteroides PRL2011 genome analysis reveals clues
for colonization of the insect gut. *PLoS One* **7**, e44229 (2012).
27. K. M. Ellegaard *et al.*, Genomic changes underlying host specialization in the bee gut
symbiont Lactobacillus Firm5. *Mol Ecol* **28**, 2224-2237 (2019).
28. K. M. Ellegaard *et al.*, Extensive intra-phylotype diversity in lactobacilli and
bifidobacteria from the honeybee gut. *BMC Genomics* **16**, 284 (2015).
29. H. Zheng *et al.*, Metabolism of Toxic Sugars by Strains of the Bee Gut Symbiont
Gilliamella apicola. *MBio* **7** (2016).
30. H. Zheng *et al.*, Division of labor in honey bee gut microbiota for plant
polysaccharide digestion. *Proc Natl Acad Sci U S A* **116**, 25909-25916 (2019).
31. H. Zheng, J. E. Powell, M. I. Steele, C. Dietrich, N. A. Moran, Honeybee gut microbiota
promotes host weight gain via bacterial metabolism and hormonal signaling. *Proc*
Natl Acad Sci U S A **114**, 4775-4780 (2017).
32. P. Engel, V. G. Martinson, N. A. Moran, Functional diversity within the simple gut
microbiota of the honey bee. *Proc Natl Acad Sci U S A* **109**, 11002-11007 (2012).
33. Y. Li, S. P. Leonard, J. E. Powell, N. A. Moran, Species divergence in gut-restricted
bacteria of social bees. *Proc Natl Acad Sci U S A* **119**, e2115013119 (2022).
34. S. Brochet *et al.*, Niche partitioning facilitates coexistence of closely related honey
bee gut bacteria. *Elife* **10** (2021).
35. K. M. Ellegaard, P. Engel, Genomic diversity landscape of the honey bee gut
microbiota. *Nat Commun* **10**, 446 (2019).
36. K. M. Ellegaard, S. Suenami, R. Miyazaki, P. Engel, Vast Differences in Strain-Level
Diversity in the Gut Microbiota of Two Closely Related Honey Bee Species. *Curr Biol*
30, 2520-2531 e2527 (2020).
37. A. E. S. Cerqueira *et al.*, Extinction of anciently associated gut bacterial symbionts in
a clade of stingless bees. *ISME J* **15**, 2813-2816 (2021).
38. L. L. Figueroa, J. J. Maccaro, E. Krichilsky, D. Yanega, Q. S. McFrederick, Why Did the
Bee Eat the Chicken? Symbiont Gain, Loss, and Retention in the Vulture Bee
Microbiome. *Mbio* **12** (2021).
39. K. L. Haag *et al.*, Temporal Changes in Gut Microbiota Composition and Pollen Diet
Associated with Colony Weakness of a Stingless Bee. *Microb Ecol* 10.1007/s00248-
022-02027-3 (2022).
40. M. S. NGALIMAT *et al.*, A Review on the Association of Bacteria with Stingless Bees.
Sains Malaysiana **49**, 1853-1863 (2020).

41.   D. Tamarit *et al.*, Functionally Structured Genomes in Lactobacillus kunkeei
Colonizing the Honey Crop and Food Products of Honeybees and Stingless Bees.
Genome Biol Evol **7**, 1455-1473 (2015).
42.   N. A. I. Jaafar, S. A. S. Mohamad, W. R. W. A. Razak, Antimicrobial activity and
antibiotic resistance of lactic acid bacteria isolated from Malaysian stingless bee's
gut. *Malays J Microbiol* **15**, 333-341 (2019).
43.   S. A. Oliphant *et al.*, Apilactobacillus apisilvae sp. nov., Nicolia spurrieriana gen. nov.
sp. nov., Bombilactobacillus folatiphilus sp. nov. and Bombilactobacillus
thymidiniphilus sp. nov., four new lactic acid bacterial isolates from stingless bees
Tetragonula carbonaria and Austroplebeia australis. *Int J Syst Evol Microbiol* **72**
(2022).
44.   D. W. Roubik, Stingless Bee (Apidae: Apinae: Meliponini) Ecology. *Annu Rev Entomol*
10.1146/annurev-ento-120120-103938 (2022).
45.   C. Grüter, *Stingless Bees* (Springer Nature, 2020), 10.1007/978-3-030-60090-7.
46.   C. A. Gaulke *et al.*, Ecophylogenetics Clarifies the Evolutionary Association between
Mammals and Their Gut Microbiota. *mBio* **9** (2018).
47.   M. Groussin *et al.*, Unraveling the processes shaping mammalian gut microbiomes
over evolutionary time. *Nat Commun* **8**, 14319 (2017).
48.   N. D. Youngblut *et al.*, Host diet and evolutionary history explain different aspects of
gut microbiome diversity among vertebrate clades. *Nat Commun* **10**, 2200 (2019).
49.   A. T. Reese, R. R. Dunn, Drivers of Microbiome Biodiversity: A Review of General
Rules, Feces, and Ignorance. *MBio* **9** (2018).
50.   B. D. Muegge *et al.*, Diet drives convergence in gut microbiome functions across
mammalian phylogeny and within humans. *Science* **332**, 970-974 (2011).
51.   D. Levin *et al.*, Diversity and functional landscapes in the microbiota of animals in the
wild. *Science* 10.1126/science.abb5352 (2021).
52.   S. Brochet *et al.*, Niche partitioning facilitates coexistence of closely related honey
bee gut bacteria. *Elife* **10** (2021).
53.   S. Bossert *et al.*, Combining transcriptomes and ultraconserved elements to
illuminate the phylogeny of Apidae. *Mol Phylogenet Evol* **130**, 121-131 (2019).
54.   J. Romiguier *et al.*, Phylogenomics Controlling for Base Compositional Bias Reveals a
Single Origin of Eusociality in Corbiculate Bees. *Mol Biol Evol* **33**, 670-678 (2016).
55.   M. Groussin *et al.*, Elevated rates of horizontal gene transfer in the industrialized
human microbiome. *Cell* **184**, 2053-2067 e2018 (2021).
56.   E. Powell, N. Ratnayeke, N. A. Moran, Strain diversity and host specificity in a
specialized gut symbiont of honeybees and bumblebees. *Mol Ecol* **25**, 4461-4471
(2016).
57.   Y. H. Tola *et al.*, Loss and Gain of Gut Bacterial Phylotype Symbionts in Afrotropical
Stingless Bee Species (Apidae: Meliponinae). *Microorganisms* **9** (2021).
58.   J. C. Holley, M. N. Jackson, A. T. Pham, S. C. Hatcher, N. A. Moran, Carpenter Bees
(Xylocopa) Harbor a Distinctive Gut Microbiome Related to That of Honey Bees and
Bumble Bees. *Appl Environ Microbiol* **88**, e0020322 (2022).
59.   M. Y. Handy *et al.*, Incipiently social carpenter bees (Xylocopa) host distinctive gut
bacterial communities and display geographical structure as revealed by full-length
PacBio 16S rRNA sequencing. *Mol Ecol* 10.1111/mec.16736 (2022).
60.   A. H. Moeller *et al.*, Cospeciation of gut microbiota with hominids. *Science* **353**, 380-
382 (2016).

61. A. H. Nishida, H. Ochman, Captivity and the co-diversification of great ape
microbiomes. *Nat Commun* **12**, 5632 (2021).
62. T. A. Suzuki *et al.*, Codiversification of gut microbiota with humans. *Science* **377**,
1328-1332 (2022).
63. Y. Kikuchi *et al.*, Host-symbiont co-speciation and reductive genome evolution in gut
symbiotic bacteria of acanthosomatid stinkbugs. *BMC Biol* **7**, 2 (2009).
64. P. Engel, R. Stepanauskas, N. A. Moran, Hidden diversity in honey bee gut symbionts
detected by single-cell genomics. *PLoS Genet* **10**, e1004596 (2014).
65. M. Kim, H. S. Oh, S. C. Park, J. Chun, Towards a taxonomic coherence between
average nucleotide identity and 16S rRNA gene sequence similarity for species
demarcation of prokaryotes. *Int J Syst Evol Microbiol* **64**, 346-351 (2014).
66. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
search tool. *J Mol Biol* **215**, 403-410 (1990).
67. S. Andrews (2010) FastQC: a quality control tool for high throughput sequence data.
(Babraham Bioinformatics).
68. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
69. A. Bankevich *et al.*, SPAdes: a new genome assembly algorithm and its applications
to single-cell sequencing. *J Comput Biol* **19**, 455-477 (2012).
70. H. Li, R. Durbin, Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics* **26**, 589-595 (2010).
71. G. J. Etherington, R. H. Ramirez-Gonzalez, D. MacLean, bio-samtools 2: a package for
analysis and visualization of sequence and alignment data with SAMtools in Ruby.
Bioinformatics **31**, 2565-2567 (2015).
72. D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, G. W. Tyson, CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043-1055 (2015).
73. Y. Lin *et al.*, Assembly of long error-prone reads using de Bruijn graphs. *Proc Natl
Acad Sci U S A* **113**, E8396-E8405 (2016).
74. I. Sovic *et al.*, Fast and sensitive mapping of nanopore sequencing reads with
GraphMap. *Nat Commun* **7**, 11307 (2016).
75. R. Vaser, I. Sovic, N. Nagarajan, M. Sikic, Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Res* **27**, 737-746 (2017).
76. B. J. Walker *et al.*, Pilon: an integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).
77. T. Seemann, Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-
2069 (2014).
78. V. M. Markowitz *et al.*, IMG/M 4 version of the integrated metagenome comparative
analysis system. *Nucleic acids research* **42**, D568-573 (2014).
79. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol* **16**,
157 (2015).
80. K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic acids research* **30**,
3059-3066 (2002).
81. D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**, 518-522 (2018).

82.   J. G. Caporaso *et al.*, Global patterns of 16S rRNA diversity at a depth of millions of
1062   sequences per sample. *Proc Natl Acad Sci U S A* **108 Suppl 1**, 4516-4522 (2011).
1063   83.   M. R. C. McLaren, B. J. (2021) Silva 138.1 prokaryotic SSU taxonomic training data
1064   formatted for DADA2. (Zenodo).
1065   84.   M. Kanehisa, Y. Sato, K. Morishima, BlastKOALA and GhostKOALA: KEGG Tools for
1066   Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol* **428**,
1067   726-731 (2016).
1068   85.   M. N. Price, A. M. Deutschbauer, A. P. Arkin, GapMind: Automated Annotation of
1069   Amino Acid Biosynthesis. *mSystems* **5** (2020).
1070   86.   S. S. Abby, B. Neron, H. Menager, M. Touchon, E. P. Rocha, MacSyFinder: a program
1071   to mine genomes for molecular systems with an application to CRISPR-Cas systems.
1072   *PLoS One* **9**, e110726 (2014).
1073   87.   Y. Yin *et al.*, dbCAN: a web resource for automated carbohydrate-active enzyme
1074   annotation. *Nucleic acids research* **40**, W445-451 (2012).
1075