Species-specific LUBAC-mediated M1 ubiquitination

counteracts necroptosis by segregating the

cellular distribution and fate of activated MLKL

Nadine Weinelt [1], Kaja Nicole Wächtershäuser [2], Sonja Smith [1], Geoffroy Andrieux [3],

Tonmoy Das [3], Birte Jeiler [1], Jens Roedig [1], Leonard Feist [4], Björn Rotter [4], Melanie

Boerries [3,5], Francesco Pampaloni [2], Sjoerd J. L. van Wijk [1,6,]*

[1] Institute for Experimental Cancer Research in Pediatrics, Goethe University Frankfurt, Komturstrasse 3a, 60528 Frankfurt am Main, Germany

[2] Physical Biology Group, Buchmann Institute for Molecular Life Sciences (BMLS), Goethe University Frankfurt, Max-von-Laue-Strasse 15, 60438, Frankfurt am Main, Germany

[3] Institute of Medical Bioinformatics and Systems Medicine, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, 79110 Freiburg, Germany

[4] GenXPro GmbH, Altenhoeferallee 3, 60438, Frankfurt am Main, Germany

[5] German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ), partner site Freiburg, 79110 Freiburg, Germany

[6] German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ) partner site Frankfurt/Mainz, Frankfurt am Main, Germany

[*]Corresponding author: Sjoerd J. L. van Wijk (vanWijk@med.uni-frankfurt.de, [s.wijk@kinderkrebsstiftung-frankfurt.de)](mailto:s.wijk@kinderkrebsstiftung-frankfurt.de)

26   Further information and requests for resources and reagents should be directed to

27   and will be fulfilled by the Lead Contact, Sjoerd J. L. van Wijk (vanWijk@med.uni-

28   frankfurt.de, s.wijk@kinderkrebsstiftung-frankfurt.de)

Summary

Plasma membrane accumulation of phosphorylated mixed lineage kinase domain-like (MLKL) is a hallmark of necroptosis, leading to membrane rupture and inflammatory cell death. Pro-death functions of MLKL are tightly controlled by several checkpoints, including phosphorylation. Endocytosis and exocytosis limit MLKL membrane accumulation and counteract necroptosis, but the exact mechanisms remain poorly understood. Here, we identify linear ubiquitin chain assembly complex (LUBAC)-mediated M1 poly-ubiquitination (poly-Ub) as novel checkpoint for necroptosis regulation downstream of activated MLKL in human cells. Loss of LUBAC activity inhibits necroptosis, without affecting necroptotic signaling, but by preventing membrane accumulation of activated MLKL. Flotillin-1/2 act as putative necroptotic M1 poly-Ub targets that inhibit necroptosis suppression induced by LUBAC inhibition. Finally, we confirm LUBAC-dependent suppression of necroptosis in primary human pancreatic organoids. Our findings identify LUBAC as species-specific regulator of necroptosis which prevents MLKL membrane accumulation and pioneer primary human organoids to model necroptosis in near-physiological settings.

Keywords: LUBAC, MLKL, necroptosis, flotillin, linear ubiquitination, programmed cell death, organoids

Introduction

Caspase-independent necroptosis, characterized by plasma membrane rupture, release of cytokines, chemokines and damage-associated molecular patterns (DAMPs), elicit potent immunogenic responses [1,2]. Accordingly, necroptotic cell death underlies a wide variety of inflammatory and infectious diseases and tumor progression [3,4]. Induction of necroptosis upon tumor necrosis factor (TNF) receptor 1 (TNFR1) activation relies on recruitment of the pro-survival TNFR1 complex I consisting of receptor-interacting protein kinase 1 (RIPK1), TNFR1-associated death domain protein (TRADD), TNFR-associated factor 2 (TRAF2), cellular inhibitor of apoptosis 1/2 (cIAP1/2) and the linear ubiquitin chain assembly complex (LUBAC) [3]. cIAP1/2- and LUBAC-mediated K63 and M1 poly-Ub of complex I components, including RIPK1, initiate transforming growth factor β (TGFβ)-activated kinase 1 (TAK1) complex- and inhibitor of κB kinase (IKK) complex-dependent activation of pro-inflammatory mitogen-activated protein kinase (MAPK) and nuclear factor-κB (NF-κB) signaling [5-9]. Following complex I destabilization through cIAP1/2 depletion and upon caspase-8 inhibition or depletion, the formation of a cytosolic necrosome complex is triggered [3]. The necrosome is an amyloid-like structure composed of RIPK1 and RIPK3, interacting via their RIP homotypic interaction motif (RHIM) domains, leading to their activation by cross- and auto-phosphorylation [10-13]. Eventually, phosphorylated RIPK3 recruits and phosphorylates MLKL at T357 and S358, resulting in the exposure of the MLKL 4-helical bundle (4HB) domain, which induces MLKL oligomerization and subsequent translocation to the plasma membrane [14-20]. Activated MLKL accumulates in hotspots to control plasma membrane disruption and ion channel fluxes, ultimately leading to necroptotic cell death [17,18,21-23].

Necroptosis is controlled by checkpoints that include compartimentalization and
species-specific post-translational modifications, like phosphorylation and
ubiquitination of RIPK1, RIPK3 and MLKL [24-26]. Ubiquitination regulates protein
function, stability and localization by covalent modification of substrates with poly-Ub
chains, linked via K6, K11, K27, K29, K33, K48, K63, or M1-linked (linear) Ub [27-29].
M1 poly-Ub is mediated by the LUBAC E3 ligase, consisting of the catalytic subunit
HOIP (RNF31), HOIL-1 (RBCK1) and Sharpin [30-33]. HOIP belongs to the really
interesting new gene (RING)/in-between RING (IBR)/RING (RBR)-type family of E3
ligases and specifically generates M1 poly-Ub chains through a RING-HECT hybrid
reaction by transferring the activated Ub to its active site C885 and subsequently
conjugating it to the acceptor Ub located in the HOIP linear ubiquitin chain
determining domain (LDD) [34-36]. The catalytic activity of HOIP can be blocked by
HOIPIN-8, a small-molecule inhibitor that covalently binds to the HOIP catalytic C885
and interferes with several residues in the LDD of HOIP, thereby inhibiting the RING-
HECT hybrid reaction [37,38].

LUBAC generates pro-inflammatory and -survival signaling checkpoints upon
activation of immune receptors, such as TNFR1, CD40, IL-1R, nucleotide
oligomerization domain (NOD)-like receptors (NLRs) and toll-like receptors (TLRs)
via M1 poly-Ub of receptors, receptor-associated proteins or existing K63 poly-Ub
chains [5,6,39]. M1 poly-Ub chains act as a scaffold to recruit signaling complexes, such
as the TAK1 complex and the IKK complex to activate downstream MAPK and NF-κB
signaling [6,8,9,39,40]. Accordingly, LUBAC deficiency in mice, or in patients with genetic
mutations in *HOIP*, *HOIL-1* or *Sharpin* results in inflammatory disorders linked to
deregulated NF-κB activity leading to decreased gene-activatory signaling [30-32,41-45].
Similarly, genetic mouse models lacking functional LUBAC are sensitized towards

TNFR1- and RIPK1-dependent cell death [30,31,46-50] and HOIPINs mimic LUBAC loss-

of-function mutations by inhibiting cytokine-induced canonical NF-κB signaling and by

regulating TNFα-induced cell viability [37,51,52].

Apart from anti-necroptotic ubiquitination, also the removal of oligomerized MLKL

from the plasma membrane via endosomal sorting complex required for transport

(ESCRT)-mediated plasma membrane shedding and exocytic release of MLKL or via

endocytosis can counteract necroptotic cell death [22,53]. The homologous Flotillin-1

and -2 (FLOT1/2) proteins reside in lipid rafts, highly ordered membrane structures

which are enriched in sphingolipids, cholesterol and gangliosides in the plasma

membrane, the endoplasmic reticulum (ER), the Golgi apparatus and endocytic

compartments [54-57]. FLOT1/2 associate with cholesterol-rich membrane domains in

the inner leaflet of cell membranes via the N-terminal stomatin, prohibitin, flotillin,

HflK/C (SPFH) domain and oligomerize to heterotetramers through the C-terminal

flotillin domain to create FLOT platforms [58,59]. FLOTs mediate the regulation of

cytoskeleton-membrane interactions and cell-cell contacts, clathrin-independent

endocytosis and the establishment of protein complexes at the plasma membrane [60-

68]. Interestingly, FLOTs have recently been implicated in the protection from

necroptotic cell death through endocytic removal of activated MLKL from the plasma

membrane [53].

Despite recently acquired insights into the checkpoints that control the pro-

necroptotic functions of activated MLKL, it is still vastly unexplored how necroptosis is

regulated downstream of MLKL activation and oligomerization. In this study, we

identify LUBAC-mediated M1 poly-Ub as novel major and species-specific checkpoint

in necroptotic cell death in human cells. Suppression of LUBAC and M1 poly-Ub

blocks TNFα-induced necroptosis without affecting necroptotic phosphorylation of

RIPK1, RIPK3 or MLKL, necrosome formation and MLKL oligomerization. Loss of M1

poly-Ub suppresses MLKL membrane hotspot accumulation, as well as MLKL-

dependent release of pro-inflammatory signaling molecules, as determined by global

Massive Analyses of cDNA Ends sequencing (MACE-seq)-based transcriptome

analysis. We further identify FLOT1/2 as putative M1 poly-Ub targets during

necroptosis and demonstrate that loss of FLOT1/2 additionally decreases necroptotic

cell death associated with LUBAC inhibition. Finally, we confirm that loss of LUBAC

activity prevents necroptosis in primary human pancreatic organoids (hPOs). Taken

together, we identify a novel, species-specific role for LUBAC and M1 poly-Ub in

limiting membrane accumulation of activated MLKL and preventing necroptosis. In

addition, by modelling necroptotic cell death and LUBAC function in primary human

organoids, we provide a novel experimental platform to study programmed cell death

in intact human multicellular systems.

Results

LUBAC-mediated M1 poly-Ub is required for TNFα-induced necroptosis in human cells

Necroptotic cell death is tightly regulated and controlled by several molecular checkpoints, including receptor stimulation, phosphorylation of RIPK1, RIPK3 and MLKL as well as compartimentalization of phosphorylated MLKL [22,24,53]. To investigate how LUBAC and M1 poly-Ub regulate these necroptotic checkpoints, cell death was induced by TNFα (T), the Smac mimetic BV6 (B) and the pan-caspase inhibitor zVAD.fmk (Z). As anticipated, prominent necroptosis could be triggered by TBZ in human colon carcinoma HT-29 cells, as determined by high-content imaging-based quantification of propidium iodide (PI)-positive cells **(Figure 1A)**. HT-29 cells are highly sensitive towards TNFα-induced programmed cell death and are commonly used model cell lines to study necroptosis. Surprisingly, chemical inhibition of HOIP, the catalytic subunit of LUBAC, with the small molecular weight inhibitor HOIPIN-8 [38], rescued TBZ-induced necroptosis in a time-dependent manner **(Figure 1A)**. Since LUBAC subunit levels are differentially regulated in myeloid cells as compared to other cell types [69,70], the effect of HOIP inhibition was investigated in the human monocytic cell line THP-1. TBZ-induced necroptotic cell death in THP-1 cells could also be potently reversed by HOIPIN-8 **(Figure 1B)**, confirming a broader relevance of HOIPIN-8-mediated inhibition in necroptosis. In agreement with HOIPIN-8-mediated necroptosis inhibition, chemical inhibition of LUBAC activity with the mycotoxin gliotoxin [71] also effectively blocked necroptotic cell death, without inducing additional cytotoxicity **(Figures S1A and S1B)**. Importantly, the effects of HOIPIN-8- and gliotoxin-based HOIP inhibition on necroptosis could be confirmed by siRNA-mediated knock-down of HOIP expression that rescued necroptosis to a similar extent as HOIPIN-8- and gliotoxin-mediated LUBAC inhibition **(Figures 1C and**

S1C), making off-target effects unlikely. Importantly, total cellular levels of M1 poly-

Ub increased upon induction of necroptosis, which could be efficiently reduced by

HOIPIN-8 and siRNA-mediated knock-down of HOIP in HT-29 cells **(Figures 1D and**

1E) and THP-1 cells **(Figure S1D)**. Together, these observations reveal that LUBAC-

mediated M1 poly-Ub serves as a novel central checkpoint and conserved mediator

of TNFα-induced necroptotic cell death.

Up till now, the relevance of M1 poly-Ub generated by LUBAC in TNFα-mediated cell

fate signaling has been prominently demonstrated in genetic mouse models deficient

in *HOIP*, *HOIL-1* or *Sharpin*. Loss of either HOIP, HOIL-1 or Sharpin in mice

interferes with TNFR1-mediated signaling, leading to impaired NF-κB activation and

increased sensitivity towards TNFR1-mediated programmed cell death [30,46-50,72,73].

Therefore, the consequences of LUBAC inhibition on TNFα-induced necroptosis

were evaluated in mouse embryonic fibroblasts (MEFs) and L-929 mouse lung

fibroblasts. HOIPIN-8 acts by interfering with the active site residue C885 as well as

residues in the HOIP LDD [37], which are conserved in humans and mice **(Figure**

S1E). LUBAC inhibition efficiently abrogated TNFα-induced NF-κB activation and

stabilized phosphorylated IκBα in both HT-29 and MEFs **(Figures S1F and S1G)**,

consistent with the described pro-inflammatory roles of LUBAC and M1 poly-Ub and

in agreement with the abovementioned findings. However, to our surprise, HOIPIN-8-

mediated LUBAC inhibition did not rescue TBZ-induced cell death in MEFs **(Figure**

S1H) and even enhanced TBZ-induced cell death in L-929 cells, without inducing

cytotoxicity, **(Figure S1I)**, which could be effectively blocked by inhibition of RIPK3

with GSK'872 **(Figure S1I)**. This is in accordance with previous findings

demonstrating that RNAi-mediated silencing of HOIP or HOIL-1 sensitizes L-929 cells

to necroptotic cell death [73]. Of note, the cellular levels of M1 poly-Ub remained

relatively stable upon necroptosis induction in MEFs **(Figure S1J)**. These

observations, confirmed by a detailed analysis of cell death regulation in genetic

mouse models of LUBAC deficiency [30,46,47,72], suggest that LUBAC acts as species-

specific switch in the control of necroptosis.

Loss of LUBAC-catalyzed M1 poly-Ub does not affect necroptotic signaling and

necrosome formation

To mechanistically understand how LUBAC controls necroptosis, the role of M1 poly-

Ub on necroptotic checkpoints was analyzed in further detail. Importantly, necroptosis

induction did not affect the expression levels of the LUBAC subunits HOIP, HOIL-1

and Sharpin **(Figures 2A and S2A)**, suggesting that LUBAC activity is responsible

for the increased levels of M1 poly-Ub during necroptosis, instead of alterations in

LUBAC abundance. In addition, HOIPIN-8-mediated inhibition of LUBAC did not have

any influence on the expression patterns of individual LUBAC subunits, excluding the

possibility that selective loss of subunits might cause the observed HOIPIN-8-induced

effects on M1 poly-Ub during necroptosis **(Figures 2A and S2A)**.

Surprisingly, and in contrast to the prominent HOIPIN-8-mediated rescue of

necroptosis, HOIPIN-8 did not attenuate TBZ-induced phosphorylation of RIPK1 at

S166, RIPK3 at S227 and MLKL at S358, major checkpoints in necroptotic cell death

signaling **(Figure 2B and Figures S2B and S2C)**. In contrast, LUBAC inhibition

induced a slight accumulation of phosphorylated RIPK1 S166, RIPK3 S227 and

MLKL S358 **(Figures 2B and Figures S2B and S2C)**, suggesting alternative effects

of LUBAC on the core necroptotic cell death machinery.

A central hallmark of necroptosis consists in the formation of the necrosome, a specialized amyloid-like signaling platform, composed of activated RIPK1 and RIPK3, that eventually phosphorylates MLKL to drive necroptotic cell death [10,11,14,17]. Importantly, inhibition of LUBAC did not interfere with necrosome formation, as determined by co-immunoprecipitation of phosphorylated RIPK1 S166 and MLKL S358 with RIPK3 from untreated and HOIPIN-8-treated necroptotic cells **(Figure 2C)**.

Prior to membrane accumulation, activated MLKL oligomerizes into larger structures that are required for efficient necroptosis progression [17-20]. Upon induction of necroptosis, prominent MLKL oligomerization could be observed that, in contrast to GSK'872-mediated RIPK3 inhibition, could not be blocked by HOIPIN-8 **(Figure 2D)**. Furthermore, inhibiting MLKL with necrosulfonamide (NSA) did neither affect MLKL oligomerization nor phosphorylation **(Figure 2D)**, as NSA only prevents membrane accumulation of activated MLKL [17]. Similarly, siRNA-mediated loss of HOIP also did not have any effect on necroptosis-induced RIPK1 S166, RIPK3 S227 and MLKL S358 phosphorylation and did not attenuate MLKL oligomerization **(Figure 2E)**. In contrast to HOIPIN-8-mediated inactivation of LUBAC, inhibition of RIPK1 with necrostatin-1 (Nec-1s) or RIPK3 with GSK'872 influences necroptosis-induced phosphorylation of RIPK1 S166, RIPK3 S227 and MLKL S358, respectively, while blocking MLKL with NSA did not affect phosphorylation levels **(Figure 2F)**. Of note, in contrast to HOIPIN-8, Nec-1s, GSK'872 or NSA, as well as deletion of RIPK3 or MLKL, did not influence total M1 poly-Ub levels upon induction of necroptosis, suggesting that regulation of M1 poly-Ub occurs in parallel with necroptosis initiation and progression **(Figures 2F and S2C)**.

LUBAC-mediated M1 poly-Ub limits necroptosis-induced and MLKL-dependent production of inflammatory signaling molecules

Upon the induction of necroptosis, activated MLKL triggers the transcriptional upregulation and secretion of a variety of inflammatory signaling molecules, including cytokines and chemokines, like CXCL1 and CXCL10 [1,22]. Consistently with these findings, TBZ-triggered necroptosis strongly induced expression of CXCL1 and CXCL10 mRNA **(Figure 3A)** in an MLKL-dependent manner, since inhibition of MLKL with NSA and CRISPR/Cas9-mediated knock-out (KO) of MLKL expression largely prevented necroptosis-induced upregulation of CXCL1 and CXCL10 mRNA (**Figures 3A and S3A)**. Interestingly, LUBAC inhibition in necroptotic cells reduced CXCL1 and CXCL10 mRNA expression **(Figure 3A)** and similar findings were observed upon loss of HOIP expression **(Figure 3B)**. In agreement with the HOIP-dependent alterations in CXCL1 and CXCL10 mRNA expression, secretion of CXCL1 during necroptosis was also diminished in the presence of HOIPIN-8 or upon loss of HOIP expression **(Figures 3C and 3D)**. To further characterize the LUBAC-dependent regulation of the transcriptional changes associated with necroptosis, transcriptome profiling using unbiased MACE-seq was perfomed in necroptotic cells upon inhibition of LUBAC activity. Necroptosis prominently triggered the transcriptional upregulation of a highly pro-inflammatory gene expression profile **(Figure 3E)**, in agreement with previous observations [1]. Importantly, loss of LUBAC activity potently reversed the necroptotic inflammatory profile and strongly attenuated the expression of necroptosis-upregulated cytokines and chemokines, such as TNFα, CSF1, CXCL2 and CCL20, as well as the adhesion molecule ICAM1 **(Figure 3E, Figures S3B and S3C)**. Of note, necroptosis-induced expression of TNFα and ICAM1 could be largely prevented by inhibiton of MLKL with NSA **(Figures S3B and S3C)**, confirming that

MLKL is required for necroptosis-induced upregulation of regulatory and inflammatory

gene expression signatures.

Together, these data suggest that M1 poly-Ub counteracts the activated MLKL-

dependent secretion of signaling molecules, cytokines and chemokines, such as

CXCL1 and CXCL10, upon induction of necroptosis.

LUBAC and M1 poly-Ub segregate the subcellular distribution of activated MLKL

During the effector phase of necroptosis, activated MLKL oligomers grow to form

cytoplasmic clusters which translocate to the cell membrane to accumulate into

hotspots, preferentially at intercellular junctions [21]. To investigate whether LUBAC-

mediated M1 poly-Ub might control membrane translocation of activated MLKL, a

non-ionic detergent- and temperature-dependent fractionation of micelle-poor

(aqueous) and micelle-rich (detergent-enriched) fractions was applied to isolate

hydrophobic, membrane-associated proteins from aqueous solutions. In necroptotic

cells, phosphorylated MLKL accumulated in detergent-enriched fractions, which could

be efficiently blocked by NSA **(Figure 4A)**, suggesting extensive membrane

accumulation that is consistent with previous findings [17]. Surprisingly, LUBAC

inhibition prevented accumulation of phosphorylated MLKL in detergent-enriched

fractions, leading instead to an enrichment of phosphorylated MLKL in the aqueous

fraction **(Figure 4A)**. These data suggest that LUBAC-mediated M1 poly-Ub is a

prerequisite for the translocation of activated MLKL from the cytosol to membrane

compartments during necroptosis induction. Of note, necroptosis also triggered the

accumulation of M1 poly-Ub chains in both aqueous and detergent-enriched

fractions, that could be counteracted by LUBAC inhibition **(Figure S4A)**.

To further understand how LUBAC and M1 poly-Ub regulate the membrane

localization of phosphorylated MLKL during necroptosis, MLKL distribution was

assessed by immunofluorescence. Upon necroptosis induction, S358-phosphorylated

MLKL localized to membranes in large hotspots at intercellular junctions **(Figures 4B**

and S4B), corresponding with previous findings [21]. As anticipated, MLKL inhibition

with NSA blocked membrane accumulation of activated MLKL in necroptotic cells

(Figure S4B). In accordance with the altered cellular distribution of MLKL in

fractionation experiments, HOIPIN-8 completely abrogated the formation of

phosphorylated MLKL hotspots at the plasma membrane in necroptotic cells, leading

to the appearance of diffuse cytoplasmic punctate clusters of S358 phosphorylated

MLKL **(Figure 4B)**.

The cellular fate of phosphorylated MLKL determines the functional outcome of

necroptotic signaling and the regulation of the availability of necroptosis-prone

phosphorylated MLKL is tightly controlled. At one end, activated MLKL is internalized

through endocytosis and degraded through the lysosomal pathway [53]. At the other

end, the levels of activated MLKL are controlled by exocytosis into extracellular

vesicles [22,53,74,75]. TBZ-induced necroptosis strongly triggered exocytosis leading to

the formation of ALIX-, CD9- and vinculin-positive extracellular exosomes that

contained high levels of MLKL **(Figure 4C)**. HOIPIN-8 strongly reduced the

necroptosis-associated formation of extracellular vesicles and decreased the

accumulation of MLKL in exosomes **(Figure 4C)**. As expected, MLKL inhibition with

NSA completely blocked TBZ-induced exosome formation and release of MLKL

(Figure 4C). In line with these observations, HOIPIN-8-mediated inhibition of LUBAC

prevented lysosomal degradation of phosphorylated MLKL upon necroptosis

induction, as determined by washout of the necroptotic stimulus in the absence or

presence of the lysosomal inhibitor bafilomycin A1 (BafA1) **(Figure 4D)**. Intriguingly, LUBAC inhibition in necroptotic cells increased levels of phosphorylated MLKL S358 **(Figure 4D),** corresponding with previous data **(Figure S2B)**. Taken together, these data demonstrate that LUBAC-mediated M1 poly-Ub regulates necroptosis downstream of MLKL phosphorylation and oligomerization by controlling the availability and translocation of activated MLKL to membranes and MLKL hotspot formation at the plasma membrane.

FLOT1/2 license necroptosis in a M1 poly-Ub-dependent manner

Recent studies revealed that FLOT1/2 colocalize with MLKL during necroptosis and act as important mediators of MLKL endocytosis [53,75]. To understand the potential regulatory link between M1 poly-Ub, the cellular distribution of activated MLKL and FLOT1/2, enrichment of M1 poly-Ub chains, substrates and associated proteins was performed from necroptotic cells. Interestingly, high levels of FLOT1/2 could be enriched by capture of M1 poly-Ub via ubiquitin binding in ABIN and NEMO (UBAN)-based pulldowns from necroptotic cells **(Figures 5A and S5A)**. Of note, HOIPIN-8-mediated LUBAC inhibition completely prevented M1 poly-Ub-associated enrichment of FLOT1/2, without affecting total FLOT1/2 cellular expression levels **(Figures 5A and S5A)**. Furthermore, loss of FLOT1 or -2 individually, or combined loss of FLOT1/2, decreased necroptotic cell death and further potentiated the repressive effects of LUBAC inhibition on necroptosis **(Figures 5B and S5B)**. Loss of FLOT1/2 did not affect necroptosis-induced increases in M1 poly-Ub or HOIPIN-8-mediated decreases in necroptosis-triggered M1 poly-Ub **(Figure S5C)**. In addition, combined loss of FLOT1/2 did not interfere with necrosome formation **(Figure 5C)** and did not affect necroptosis-induced MLKL oligomerization in the absence or presence of HOIPIN-8 **(Figure 5D)**. Intruigingly, enrichment of FLOT1 in UBAN-based pulldowns

could be prevented by inhibition of RIPK1 with Nec-1s, RIPK3 with GSK'872 and

MLKL with NSA, indicating that activation and membrane translocation of MLKL is a

prerequisite for association of FLOT1 with M1 poly-Ub **(Figure 5E)**.

LUBAC-mediated M1 poly-Ub regulates necroptotic cell death in primary human

pancreatic organoids

Our data reveal that LUBAC and M1 poly-Ub act as a novel, species-specific

checkpoint that limit the membrane hotspot accumulation of activated MLKL and

thereby counteracts necroptotic cell death in human, but not in murine cell lines. To

further confirm this concept and to exclude artefacts in necroptotic signaling related

to the use of human cancer cell lines, LUBAC- and M1 poly-Ub-mediated regulation

of necroptosis was studied in primary human organoids. Organoids are stem cell-

derived three-dimensional cellular clusters consisting of organ-specific cell types that

self-organize and exhibit similar functionality and complexity as their tissue of origin

[76,77]. Organoids represent highly physiological models as compared to two-

dimensional monolayer cell culture models [78]. To model necroptosis, adult stem cell-

derived primary human pancreatic organoids (hPOs) recapitulate the phenotype of

the original pancreatic ductal epithelia and retain the pancreatic morphology and the

distribution of ductal, acinar, mesenchymal and endothelial cells in a hollow spherical

polarized cell monolayer enclosing a central lumen [79-82]. hPOs were maintained in

matrix for 3 to 5 days to allow the formation of organoid spheres composed of a

cellular monolayer surrounding a liquid-filled lumen, that expanded over time

(Figures 6A and S6A), in line with previous findings [83], after which necroptosis was

induced. For this, matrix-embedded primary hPOs were incubated with BV6 (B) and

the clinically-applicable pan-caspase inhibitor IDN-6556 (Emricasan; E), followed by

time-lapse imaging and subsequent fluorescein diacetate (FDA)- and PI-based

live/dead imaging. BV6 combined with Emricasan induced a prominent loss of

luminal hPO morphology, characterized by organoid collapse and compaction, with

individual cells migrating away from the collapsed organoids **(Figure 6A, Movie 1)**.

Live/dead staining of BE-incubated organoids revealed high levels of PI-positive cells

and low levels of FDA-positive cells, confirming prominent hPO cell death **(Figure**

6B). Similar organoid collapse and loss of luminal morphology were observed in

hPOs incubated with the Smac mimetic Birinapant (Bi) and Emricasan **(Figure S6A,**

Movie 2) and upon incubation with BV6 and zVAD.fmk (BZ) **(Figure S7)**. Importantly,

HOIPIN-8-mediated inhibition of LUBAC delayed BE-induced organoid collapse,

prevented the loss of luminal hPO morphology and extended hPO viability up to 24 h

(Figures 6A and 6B, Movie 1), although hPOs appeared shrunken and smaller in

size **(Figure 6A)**. HOIPIN-8 also blocked organoid collapse and cell death induced by

BiE and BZ **(Figures S6A and S6B, Figure S7, Movie 2)**. Of note, organoid collapse

and compaction, loss of hPO morphology and cell death triggered by BiE could be

blocked by NSA, indicating the appearance of MLKL-dependent necroptotic cell

death **(Figure S6, Movie 2)**.

In conclusion, we identify LUBAC-mediated M1 poly-Ub as novel checkpoint that

regulates necroptosis by controlling the cellular distribution of activated MLKL

between membrane hotspot formation and endo- and exocytosis in cell lines of

human origin and primary human organoids.

Discussion

Necroptosis relies on the highly regulated phosphorylation of RIPK1 and RIPK3 in the context of the death-inducing necrosome complex, leading to phoshorylation of the pseudokinase MLKL, followed by MLKL oligomerization, membrane accumulation and cell death [11-15,17]. By doing so, necroptotic cell death is involved in a wide variety of diseases, including infection [84-87], cancer [88,89] and ischemia-reperfusion injury [90,91]. TNFα-induced necroptosis, but also alternatively-induced forms of necroptosis, are controlled by checkpoints that include compartimentalization, complex formation and post-translational modification, like phosphorylation and ubiquitination, of the key necroptotic proteins RIPK1, RIPK3 and MLKL [24-26]. M1-linked poly-Ub, catalyzed by the LUBAC E3 ligase, serves as important regulator that balances pro-and anti-survival measures by modification of RIPK1. Here, we identify a novel regulatory function of M1 poly-Ub during necroptosis in facilitating the accumulation of activated MLKL in membrane hotspots to promote necroptotic cell death in cells of human origin.

In contrast to genetic models of LUBAC deficiency in mice, in which LUBAC-mediated M1 poly-Ub represses cell death [30,31,46-50,72], we found that pharmacological inhibition of LUBAC with either HOIPIN-8 or gliotoxin, as well as siRNA-mediated loss of LUBAC, prevents TNFα-induced necroptosis in human, but not murine, cell lines. Induction of necroptosis triggered M1 poly-Ub that was not dependent on the activity or the physical presence of RIPK1, RIPK3 and MLKL, suggesting that increases in M1 poly-Ub during necroptosis are not regulated by the core necroptotic machinery. In addition, loss of LUBAC activity reverts the cellular levels of necroptosis-induced M1 poly-Ub in human cell lines, but to a lesser degree in cell lines of murine origin. LUBAC inhibition abrogated TNFα-induced NF-κB signaling to the same extent in cell

lines from both species, suggesting species-, pathway- and context-specific modes of

LUBAC regulation during necroptosis. Importantly, structural studies revealed

important mechanistic and functional differences between human and murine

necroptotic signaling pathways [92-95]. These confirm that the mechanisms by which

MLKL is activated by RIPK3 differ substantially in humans and mice, as a result of

species-specific coevolution and divergence of RIPK3 and MLKL [92,94]. The activation

of MLKL is achieved through RIPK3-dependent phosphorylation of human MLKL at

T357 and S358 and murine MLKL at S345, located in the activation loop of MLKL

[14,15,96,97]. Intruigingly, human and murine MLKL are not able to complement

necroptotic signaling when expressed in cells of the opposing species [92-94]. One of

the most prominent differences in how human and murine MLKL are activated relies

on the transient interaction between murine RIPK3 and MLKL, which is in striking

contrast to the activation of human MLKL that requires recruitment and binding to

RIPK3 [92], indicating additional roles for T357/S358 phosphorylation in human MLKL

[92]. In addition, species-specific, yet unidentified, differences in necroptosis execution

downstream of activated MLKL may also explain the distinct functions of LUBAC and

M1 poly-Ub in the regulation of necroptotic cell death in mice and humans.

Our findings reveal that, apart from the core necroptosis machinery, LUBAC-

dependent signaling events might become activated in parallel that involve M1 poly-

Ub to eventually affect necroptosis. In line with this, loss of LUBAC activity did not

alter necroptotic phosphorylation of RIPK1, RIPK3 and MLKL, necrosome formation

or the oligomerization of MLKL during necroptosis induction. In addition, TNFα-

induced necroptosis has been linked to the cell-autonomous upregulation and

increased secretion of signaling molecules, inlcuding cytokines and chemokines [1,22].

Induction of this necroptotic cytokine profile involves two phases, the first of which

depends on TNFα-induced NF-κB-mediated upregulation of inflammatory signaling molecules. The second phase is a RIPK1-, RIPK3- and MLKL-mediated wave, that relies on necroptotic activation of RIPK1 and RIPK3, as well as MLKL [1,22]. Our experiments confirm the hypothesis of parallel signaling cascades activated during necroptosis and reveal that LUBAC-dependent regulation affects both necroptotic NF-κB- and MLKL-dependent production of inflammatory cytokines. Apart from necroptotic signaling, necroptosis-independent functions of MLKL include the control of ICAM1, VCAM1 and E-selectin expression via RNA-binding motif protein 6 (RBM6) [98]. Induction of necroptosis leads to a LUBAC- and MLKL-dependent upregulation of ICAM1 expression, suggesting that LUBAC also controls necroptosis-independent functions of MLKL. Taken together, these findings confirm parallel and independent LUBAC-dependent signaling events which most likely involve NF-κB and the core necroptotic machinery that converge downstream of activated MLKL to allow efficient membrane localization and necroptosis.

LUBAC inhibition prevents hotspot accumulation of phosphorylated MLKL as well as exocytosis in extracellular vesicles, suggesting that LUBAC-mediated M1 poly-Ub serves pro-necroptotic MLKL membrane thethering functions. Intruigingly, the M1 poly-Ub-dependent effects on MLKL mimic MLKL-specific monobodies that target the 4HB of human MLKL to inhibit necroptosis, but without affecting necrosome formation, MLKL activation and MLKL oligomerization [99]. The functional role of the MLKL 4HB in M1 poly-Ub-mediated necroptosis inhibition remains to be investigated. Pharmacological interference of MLKL trafficking with inhibitors of Golgi, microtubule and actin networks (e.g. with Brefeldin A, Nocodazole and Cytochalasin A, respectively), also prevented membrane hotspot formation of activated MLKL [21].

Intruigingly, loss of LUBAC triggered a similar phenotype, suggesting that LUBAC-

mediated M1 poly-Ub might be involved in the regulation of MLKL trafficking.

Both endocytic and exocytic events balance MLKL membrane accumulation and

involve the lipid raft-resided proteins FLOT1/2 [53], ESCRT complexes, ALIX, syntenin-

1 and Rab27 [22,53,74,75,100]. Recognition of ubiquitinated cargo by ESCRT complex

subunits, like hepatocyte growth factor-regulated tyrosine kinase substrate (Hrs),

signal-transducing adapter molecule (STAM) 1 and 2 and tumor susceptibility gene

(TSG) 101 via ubiquitin-binding domains (UBDs), such as the Vps27/Hrs/Stam (VHS)

domain, ubiquitin-interacting motif (UIM) domain and ubiquitin-conjugating enzyme

E2 variant (UEV) domain, are important for endo- or exocytosis [101-107]. Although the

binding specificity and affinity of these UBD-containing proteins for M1 poly-Ub

remains unclear, specific recognition of M1 poly-Ub might serve as docking platform

to position activated MLKL in hotspots and to allow necroptosis. Intruigingly, FLOT1

is implicated in the transfer of ubiquitinated cargo between different ESCRT

complexes by modulating the Ub-binding of Hrs upon epidermal growth factor (EGF)

stimulation and release of Ub-mediated autoinhibition of the Hrs UIM domain [108]. In

line with this notion, we detected a strong necroptosis-induced and LUBAC-

dependent enrichment of FLOT1/2 in M1 poly-Ub-based enrichments that was

dependent on the activation of RIPK1, RIPK3 and MLKL. Additionally, loss of

FLOT1/2 potentiated the HOIPIN-8-mediated decrease in necroptotic cell death.

Although the necroptosis-induced increase in cellular M1 poly-Ub levels is not

dependent on MLKL, inhibition of MLKL membrane translocation blocked the

association of FLOT1 with M1 poly-Ub, indicating a direct link for FLOT1/2 in M1

poly-Ub and MLKL membrane localization and further demonstrating independent

LUBAC-dependent signaling events during necroptosis.

Lipid rafts are important regulators of TNFα-mediated cell survival and death. Formation of the TNFR1 complex I in lipid rafts underlies RhoA and NF-κB activation, whereas activation of complex I outside lipid rafts promotes cell death [109,110]. Intruigingly, activated MLKL localizes to lipid rafts as well [18] and ALIX and ESCRT complexes regulate both MLKL endocytosis and TNFR1 internalization [21,111,112]. As LUBAC and M1 poly-Ub are prominent components of the TNFR1-initiated complex I, a potential proximation of complex I, LUBAC, M1 poly-Ub and activated MLKL in lipid rafts upon TNFα-induced necroptosis, as well as shared signaling functions of M1 poly-Ub, cannot be excluded.

MLKL itself has been reported to be modified by mono-Ub upon MLKL oligomerization at membranes that might direct MLKL for proteasomal and lysosomal degradation [113]. Intruigingly, although mutation of the identified mono-Ub sites in the MLKL 4HB did not prevent necroptosis, fusion of the non-specific DUB USP21 to MLKL did sensitize towards necroptosis [113]. In addition, K63 poly-Ub of MLKL at K219 increased sensitivity towards necroptosis [114]. Although the homologue residue in human MLKL, K230, is also ubiquitinated [115], it remains unclear if MLKL can be modified by M1 poly-Ub as well. Our findings further confirm that poly-Ub modification of MLKL or interacting proteins in close vicinity are important regulators of cell death.

We identified LUBAC-mediated M1 poly Ub as novel regulatory checkpoint that controls necroptosis in multicellular, primary human organoids *ex vivo*. The unique and first-time use of human primary organoids to systematically model necroptosis and the influence of LUBAC-mediated M1 poly-Ub in primary human organoids provides novel and relevant experimental models to study necroptosis in intact

multicellular, near-physiological settings. Apart from necroptosis-proficient cell lines,

experimental models to study necroptosis in human cells are extremely limited and

so far no multicellular human models have been described. Since necroptosis

underlies a wide variety of disease states, modeling necroptotic cell death in primary

organoids will significantly contribute to gain new fundamental insights and to

develop novel therapeutic options for emerging human diseases.

Acknowledgments

The authors thank Laura Zein, Rebekka Karlowitz and all members of the van Wijk lab for advice, discussions and support during the study, Christina Hugenberg for proofreading and administrative support and Lorenza Lazzari (Unit of Regenerative Medicine, Cell Factory Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico Milan, Italy) for providing the hPOs. Research in the lab of S.J.L.v.W. is supported by the Deutsche Forschungsgemeinschaft (DFG) (WI 5171/1-1, WI 5171/4-1, FU 436/20-1, FU 436/21-1 and project-ID 259130777 – CRC1177), the Deutsche Krebshilfe (70113680), Wilhelm Sander-Stiftung (2020.008.1), the Frankfurter Stiftung für krebskranke Kinder and the Dr. Eberhard and Hilde Rüdiger Foundation. Research in the group of F.P. is supported by EU FET-Open (828931), EU Transition Open (01 101057894), Wilhelm Sander-Stiftung (2020.008.1) and the German Aerospace Center (DLR) (IMMUNO3D-SHAPE). M.B. is supported by the DFG within the CRC1160 (Project ID 256073931-Z02), CRC/TRR167 (Project ID 259373024-Z01), CRC1453 (Project ID 431984000-S1), CRC1479 (Project ID: 441891347- S1). M.B. and G.A. also acknowledge funding from the German Federal Ministry of Education and Research (BMBF) within the Medical Informatics Funding Scheme - MIRACUM-FKZ 01ZZ1801B (M.B.) and EkoEstMed–FKZ 01ZZ2015 (G.A.).

Author Contributions

N.W. and S.J.L.v.W. designed the study. N.W. performed most of the experiments. K.N.W. and F.P. performed experimentation on primary hPOs and analyzed the data. J.R. generated KO cell lines. S.S. and B.J. supported with experiments. L.F. and B.R. performed MACE-seq-based transcriptome analysis. G.A., T.M. and M.B. analyzed and visualized the MACE-seq data. N.W. and S.J.L.v.W. analyzed the data and

557 prepared the manuscript. All authors have read, commented, and agreed on the

558 submitted version of the manuscript.

560 **Declaration of Interests**

561 The authors declare no competing interests.

 Figure titles and legends

Figure 1. LUBAC-mediated M1 poly-Ub is required for TNFα-induced

necroptosis in human cells. A. Quantification of cell death in untreated (UT) and

HOIPIN-8 (30 µM) pre-treated HT-29 cells after treatment with TBZ (10 ng/mL TNFα,

1 µM BV6, 20 µM zVAD.fmk) for the indicated time points. Mean and SEM of at least

three independent experiments are shown. ****P<0.0001. **B.** Quantification of cell

death in untreated (UT) and HOIPIN-8 (30 µM)-pre-treated THP-1 cells after

treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for the indicated

time points. Mean and SEM of at least three independent experiments are shown.

****P<0.0001. **C.** Quantification of cell death in control (siCtrl) and HOIP knockdown

(siHOIP) HT-29 cells treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk)

for the indicated time points. Mean and SEM of at least three independent

experiments are shown. ****P<0.0001. **D.** Western blot analysis of total M1 poly-Ub

levels in control or HOIPIN-8 (30 µM)-pre-treated HT-29 cells upon treatment with

TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for the indicated time points.

GAPDH was used as loading control. Representative blots of at least two

independent experiments are shown. **E.** Western blot analysis of total M1 poly-Ub

levels in control (siCtrl) and HOIP knockdown (siHOIP) HT-29 treated with TBZ (10

ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h. GAPDH was used as loading

control. Representative blots of at least two independent experiments are shown.

See also Figure S1.

Figure 2. Loss of LUBAC-catalyzed M1 poly-Ub does not affect necroptotic

signaling and necrosome formation. A. Western blot analysis of LUBAC subunit

(HOIP, HOIL-1 and Sharpin) expression levels in control *versus* HOIPIN-8 (30 µM)-

pre-treated HT-29 cells upon treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for 4 h. GAPDH was used as loading control. Representative blots of at

least two independent experiments are shown. **B.** Western blot analysis of

phosphorylated and total expression levels of RIPK1, RIPK3 and MLKL in control

versus HOIPIN-8 (30 µM)-pre-treated HT-29 cells upon treatment with TBZ (10

ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h. Vinculin was used as loading

control. Representative blots of at least two independent experiments are shown. **C.**

Immunoprecipitation of Strep-RIPK3 from control and HOIPIN-8 (30 µM)-pre-treated

RIPK3 KO HT-29 cells re-expressing PAM-mutated doxycycline (Dox)-inducible

Strep-tagged RIPK3 wild-type (WT) was performed in order to analyze necrosome

formation after treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for

3 h. Cells were incubated overnight with 1 µg/mL Dox prior to treatment to induce

expression of Strep-RIPK3. GAPDH was used as loading control. Representative

blots of at least two independent experiments are shown. **D.** Western blot analysis of

MLKL oligomerization upon treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for 4 h by non-reducing Western blotting of control, HOIPIN-8 (30 µM),

GSK'872 (20 µM) or NSA (10 µM)-pre-treated HT-29 cells. Vinculin or β-actin were

used as loading controls. Representative blots of at least two independent

experiments are shown. **E.** Western blot analysis of expression levels of

phosphorylated and total RIPK1, RIPK3 and MLKL and MLKL oligomerization (by

non-reducing Western blotting) upon 4 h TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) treatment of control (siCtrl) and HOIP knockdown (siHOIP) HT-29 cells.

GAPDH was used as loading control. Representative blots of at least two

independent experiments are shown. **F.** Western blot analysis of expression levels of

phosphorylated and total RIPK1, RIPK3, MLKL and M1 Ub upon 4 h TBZ (10 ng/mL

TNFα, 1 µM BV6, 20 µM zVAD.fmk) treatment of HT-29 cells pre-treated with

HOIPIN-8 (30 µM) or Nec-1s (30 µM), GSK'872 (20 µM) and NSA (10 µM). GAPDH

was used as loading control. Representative blots of at least two independent

experiments are shown. See also Figure S2.

Figure 3. LUBAC-mediated M1 poly-Ub limits necroptosis-induced and MLKL-

dependent production of inflammatory signaling molecules. A. mRNA

expression levels of CXCL1 and CXCL10 of control (UT), HOIPIN-8 (30 µM) or NSA

(10 µM)-pre-treated HT-29 cells upon treatment with TBZ (10 ng/mL TNFα, 1 µM

BV6, 20 µM zVAD.fmk) for 3 h. Gene expression was normalized against 18S and

RPII mRNA expression and is presented as x-fold mRNA expression compared to

the untreated (UT) control. Mean and SEM of at least three independent experiments

are shown. *P<0.05; ***P<0.001; ****P<0.0001. **B.** mRNA expression levels of

CXCL1 and CXCL10 of control (siCtrl) or HOIP knockdown (siHOIP) HT-29 cells

treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h. Gene

expression was normalized against 18S and RPII mRNA expression and is presented

as x-fold mRNA expression compared to the untreated (UT) control cells. Mean and

SEM of at least three independent experiments are shown. ****P<0.0001. **C.** ELISA-

based quantification of CXCL1 secretion in supernatants of control (UT) or HOIPIN-8

(30 µM)-pre-treated HT-29 cells after treatment with TBZ (10 ng/mL TNFα, 1 µM

BV6, 20 µM zVAD.fmk) for the indicated time points. Mean and SEM of at least three

independent experiments are shown. **P<0.01; ***P<0.001. **D.** ELISA-based

quantification of CXCL1 secretion in supernatants of control (siCtrl) or HOIP

knockdown (siHOIP) HT-29 cells treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20

µM zVAD.fmk) for the indicated time points. Mean and SEM of at least three

independent experiments are shown. **P<0.01; ***P<0.001. **E.** MACE-seq-

determined heatmap showing the row-wise scaled intensity (left panel) of the top-25

upregulated genes in HT-29 cells treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20

µM zVAD.fmk) for 3 h in comparison to HOIPIN-8 (30 µM)-pre-treated and TBZ-

treated HT-29 cells. Genes are sorted according to their log2 fold change values

(right panel). *$P<0.05$. See also Figure S3.

Figure 4. LUBAC and M1 poly-Ub segregate the subcellular distribution of

activated MLKL. A. Fractionation of phosphorylated and total MLKL in the micelle-

poor (aqueous) and micelle-rich (detergent) fractions upon phase separation using

Triton X-114 lysis buffer in untreated (UT) and HOIPIN-8 (30 µM) pre-treated HT-29

cells upon treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h.

GAPDH was used as loading control for soluble proteins and CD9 as loading control

for membrane proteins. Representative blots of at least two independent experiments

are shown. **B.** Representative fluorescence microscopy demonstrating the cellular

distribution of phosphorylated MLKL (green) in untreated (UT) and HOIPIN-8 (30 µM)

pre-treated HT-29 cells and after treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20

µM zVAD.fmk) for 3 h. Nuclei were stained with DAPI (blue). Arrows indicate

cytoplasmic clusters of phosphorylated MLKL. Representative images of at least two

independent experiments are shown. Scale bar 50 µm. **C.** Western blot analysis of

TBZ-induced release of MLKL in exosomes, isolated from the supernatants of

control, HOIPIN-8 (30 µM) or NSA (10 µM)-pre-treated HT-29 cells after treatment

with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 3 h. Exosome isolation

was confirmed with antibodies against ALIX, CD9 and Vinculin. Asterisk marks

unspecific band. Expression of phosphorylated and total MLKL was determined by

Western blotting of cell lysates from the same experiment. Vinculin was used as

loading control. Representative blots of at least two independent experiments are

shown. **D.** Western blot analysis of lysosomal degradation of phosphorylated MLKL

upon TBZ (10 ng/mL TNFα, 0.1 µM BV6, 20 µM zVAD.fmk) treatment of control,

HOIPIN-8 (30 µM) or NSA (10 µM)-pre-treated HT-29 cells. HT-29 cells were treated

with TBZ for 3 h and either harvested or the medium was exchanged with fresh

medium with or without BafA1 (100 nM) after a washing step with PBS. Washed-out

cells were harvested 12 h or 18 h after medium exchange. Vinculin was used as

loading control. Representative blots of at least two independent experiments are

shown. See also Figure S4.

Figure 5. FLOT1/2 license necroptosis in a M1 poly-Ub-dependent manner. A.

UBAN-mediated pulldown of M1 ubiquitinated proteins in control or HOIPIN-8 (30

µM)-pre-treated HT-29 cells treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for the indicated time points. GST and GAPDH were used as loading

controls. Representative blots of at least two independent experiments are shown. **B.**

Quantification of cell death in untreated (UT), HOIPIN-8 (30 µM) or NSA (10 µM)-pre-

treated control (siCtrl), FLOT1 (siFLOT1), FLOT2 (siFLOT2) or combined FLOT1 and

FLOT2 (siFLOT1+siFLOT2) knockdown HT-29 cells after treatment with TBZ (10

ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for the indicated time points. Mean and

SEM of at least three independent experiments are shown. $^{**}P$<0.001; $^{***}P$<0.001;

$^{****}P$<0.0001. **C.** Characterization of necrosome formation by immunoprecipitation of

Strep-RIPK3 from control and HOIPIN-8 (30 µM)-pre-treated control (siCtrl) and

combined FLOT1 and FLOT2 knockdown (siFLOT1+2) RIPK3 KO HT-29 cells re-

expressing PAM-mutated Dox-inducible Strep-tagged RIPK3 WT after treatment with

TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 3 h. Cell lines were incubated

overnight with 1 µg/mL Dox prior to treatment to induce expression of Strep-RIPK3.

GAPDH was used as loading control. Representative blots of at least two

independent experiments are shown. **D.** Western blot analysis of MLKL

oligomerization after treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for 4 h by non-reducing Western blotting of control (siCtrl) and combined

FLOT1 and FLOT2 knockdown (siFLOT1+2) HT-29 cells. GAPDH was used as

loading control. Representative blots of at least two independent experiments are

shown. **E.** UBAN-mediated pulldown of M1 ubiquitinated proteins in control, HOIPIN-

8 (30 µM), Nec-1s (30 µM), GSK'872 (20 µM) or NSA (10 µM)-pre-treated HT-29

cells treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h. GST

and GAPDH were used as loading controls. Representative blots of at least two

independent experiments are shown. See also Figure S5.

Figure 6. LUBAC-mediated M1 poly-Ub regulates necroptotic cell death in

primary hPOs. A. Representative images of time-lapse videos of untreated (UT),

BV6 (1 µM) and Emricasan (10 µM) (BE)-treated and HOIPIN-8 (30 µM)-pre-treated

and BE (BE+HOIPIN-8)-treated primary hPOs for a total period of 23.5 h. Scalebars:

250 µm. **B.** *Idem* as A., but treated primary hPOs were stained with Hoechst33342

(Hoe; blue), FDA (live; green) and PI (dead; red) after 24 h treatment with BE or

BE+HOIPIN-8 prior to imaging. Scalebars: 250 µm. See also Figures S6 and S7 and

Movies 1 and 2.

Materials and Methods

Cell lines, reagents and chemicals

The human HT-29 adenocarcinoma cell line was obtained from DMSZ (Braunschweig, Germany) and the human acute monocytic leukemia cell line THP-1 was kindly provided by Thomas Oellerich (Frankfurt am Main, Germany) and authenticated by DSMZ. MEFs were kindly provided by Joanne Hildebrand (Melbourne, Australia). The L-929 mouse fibroblast cell line was obtained from ATCC (Manassas, VA, USA). HT-29 cells were cultured in McCoy's 5A Medium +GlutaMAX^{TM}-I (Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 1% penicillin/streptomycin (Thermo Fisher Scientific) and 10% fetal bovine serum (FBS) (Thermo Fisher Scientific). THP-1 cells were maintained in RPMI1640 Medium +GlutaMAX^{TM}-I (Thermo Fisher Scientific) supplemented with 1% sodium pyruvate (Thermo Fisher Scientific), 1% penicillin/streptomycin and 10% FBS. MEFs were cultured in DMEM +GlutaMAX^{TM}-I (Thermo Fisher Scientific) supplemented with 1% sodium pyruvate, 1% penicillin/streptomycin and 10% FBS. L-929 cells were cultured in DMEM +GlutaMAX^{TM}-I (Thermo Fisher Scientific) supplemented with 1% sodium pyruvate, 1% penicillin/streptomycin, 1% MEM Non-Essential Amino Acids (NEAA) Solution (Thermo Fisher Scientific) and 10% FBS. HT-29 MLKL KO, HT-29 RIPK3 KO and HT-29 RIPK3 KO cells re-expressing PAM-mutated Dox-inducible Strep-tagged WT RIPK3 have been described previously [116]. All cell lines were cultured at 37°C with 5% carbon dioxide in a humidified atmosphere and regularly monitored for mycoplasma infection.

BV6 was kindly provided by Genentech Inc. (San Francisco, CA, USA), recombinant human TNFα was purchased from Biochrom Ltd. (Waterbeach, UK), Birinapant from Selleckchem (Houston, TX, USA), the pan-caspase inhibitors zVAD.fmk and

Emricasan from Bachem AG (Bubendorf, Switzerland) and Selleckchem, Nec-1s, GSK'872 and NSA from Merck KGaA (Darmstadt, Germany), HOIPIN-8 from Axon Medchem LLC (Reston, VA, USA), Gliotoxin from Tocris Biosciences (Bristol, UK), Doxycycline hydrochloride from Merck KGgA (Darmstadt, Germany) and BafA1 from Selleckchem. All other chemicals were purchased from Carl Roth (Karlsruhe, Germany) or Merck KGaA, unless stated otherwise.

Maintenance of primary hPOs

Primary hPOs are derived from the pancreatic duct of a male patient (age: 60, BMI: 26.3) and were provided by Lorenza Lazzari (Policlinico di Milano, Milan, Italy). The use of these hPOs is approved by the Institutional Review Board and the Pancreatic Islet Processing Unit, a National Transplant Center accredited facility (IT000679, https://webgate.ec.europa.eu/eucoding/reports/te/index.xhtml). hPOs were kept in an incubator at 37°C in a humidified atmosphere with 5% CO_2, embedded in Cultrex Reduced Growth Factor BME, Type 2 (BME-2) droplets (R&D Systems, Minneapolis, MN, USA) and covered under medium. Medium was fully changed every 2 to 3 days and hPOs were passaged every 7 to 14 days depending on the proliferation rate and organoid size.

The organoid culture protocol was adapted from described procedures [80]. Briefly, hPOs were maintained in basal medium, composed of Advanced DMEM/F-12 (Thermo Fisher Scientific) with 1x GlutaMAX™ Supplement (Thermo Fisher Scientific), 100 U/mL penicillin/streptomycin (Thermo Fisher Scientific) and 10 mM HEPES (Thermo Fisher Scientific), supplemented with 1x B-27™ (minus vitamin A, Thermo Fisher Scientific), 1x N-2 (Thermo Fisher Scientific), 2,5 mM *N*-acetylcysteine (MilliporeSigma, St. Louis, US), 1 µg/mL recombinant human R-

Spondin1 (PeproTech, Rocky Hill, US), 10 mM nicotinamide (MilliporeSigma),

50 ng/mL recombinant human EGF (PeproTech), 100 ng/mL recombinant human

FGF-10 (PeproTech), 25 ng/mL Noggin (PeproTech), 10 nM [Leu[15]]-Gastrin I

(MilliporeSigma), 5 µM A83-01 (Tocris Bioscience, Minneapolis, US), 10 µM

Forskolin (Tocris Bioscience) and 3 µM Prostaglandin E_2 (Tocris Bioscience) (hPO

medium).

For passaging of the primary hPOs, BME droplets were disrupted by pipetting and

organoids were retrieved by centrifugation in ice-cold hPO medium. Organoids were

sheared into fragments prior to resuspension in ice-cold liquified BME-2 and seeded

as droplets in pre-warmed non-treated CELLSTAR® 48-well suspension culture

plates (Greiner Bio-One GmbH, Kremsmünster, AUT). Organoid-containing BME-2

droplets were polymerized for 15 min at 37°C after which the droplets were overlaid

with hPO medium.

Induction and quantification of necroptotic cell death

The indicated cells were seeded in CELLSTAR® 96-well tissue culture plates

(Greiner Bio-One GmbH) at appropriate densities the day before treatment. For

necroptosis induction, cells were pretreated with 1 µM BV6 and 20 µM zVAD.fmk for

30 min (if not stated otherwise), followed by stimulation with 10 ng/mL recombinant

human TNFα for the indicated time (TBZ). The inhibitors HOIPIN-8 (30 µM), Gliotoxin

(1 µM), Nec-1s (30 µM), GSK'872 (20 µM) and NSA (10 µM) were added 1 h prior to

TBZ treatment. Cell death was determined by PI and Hoechst33342 (Hoe) staining

and fluorescence-based quantification of dead (PI-positive) cells compared to the

total cell number (Hoe-positive cells) using an ImageXpress® Micro XLS Widefield

High-Content Analysis System and MetaXpress® Software (Molecular Devices

Sunnyvale, CA, USA).

Induction and imaging of necroptotic cell death in primary hPOs

hPOs were seeded in 5 µL BME-2 droplets in sterile CELLSTAR® 96-well

suspension culture plates (Greiner Bio-One GmbH) and overlaid with 100 µL hPO

medium. hPOs were cultured for 3-5 days prior to experiments to assure organoid

formation and growth. For pre-treatment with HOIPIN-8, hPO medium was replaced

by hPO medium with the indicated concentrations of HOIPIN-8 and hPOs were

placed back in the incubator. After 30 min, hPO medium was again replaced with

medium supplemented with the indicated treatments and the plates were transferred

to the Zeiss AxioObserver Z1 microscope. Incubation was done at 37°C and 5% CO_2

during which 24 h time lapse imaging in 30 min intervals was performed. Each well

was imaged as tiles of two x two with eleven Z-slices of 60 µm thickness.

For live/dead assays, primary hPOs were stained using 10 µg/mL PI (Millipore

Sigma), 0.5 µg/mL FDA (Millipore Sigma) and 200 µg/mL Hoe (ThermoFisher

Scientific) for 30 min. Dyes were removed and the wells were washed with pre-

warmed PBS prior to imaging in PBS, using the same microscope settings as

described above.

Images were exported using the ZEN 2.6 lite software (Carl Zeiss Microscopy GmbH,

Oberkochen, Germany) and subsequently processed using the FiJi/ImageJ software.

RNAi-mediated silencing

Transient genetic silencing of HOIP and FLOT1/2 was performed by reverse

transfection of HT-29 cells with the following siRNAs: human siHOIP (RNF31)

(Silencer® Select siRNA #1 s30108, #2 s30109, #3 s30110, 40 nM) (Thermo Fisher

Scientific, Waltham, MA, USA), human FLOT1 (ON-TARGETplus Human FLOT1

siRNA SMART POOL, L-010636-00-0010) (Horizon Discovery Ltd., Waterbeach,

UK), human FLOT2 (ON-TARGETplus Human FLOT2 siRNA SMART POOL, L-

003666-01-0010) (Horizon Discovery Ltd.), non-targeting control siRNA (#4390843,

40 nM) (Thermo Fisher Scientific). In brief, the indicated cell lines were prepared in

penicillin/streptomycin-free medium and combined with the siRNA-lipid complexes,

prepared in OptiMEM (Thermo Fisher Scientific) using Lipofectamine™ RNAiMAX

Transfection Reagent (Thermo Fisher Scientific) according to the manufacturer's

protocol and incubated for 5 min at room temperature. The medium was changed to

full medium after 6 h and the cells were reseeded for the experiments 48 h post-

transfection.

Western blotting

For general cell lysis, RIPA lysis buffer (50 mM Tris-HCl pH 8, 150 mM NaCl, 1%

Nonidet P-40 (NP-40), 2 mM $MgCl_2$, 0.5% sodium deoxycholate) supplemented with

protease inhibitor complex (PIC) (Roche, Grenzach, Germany), 0.1% SDS, 1 mM

sodium orthovanadate, 5 mM sodium fluoride, 1 mM β-gylcerophosphate, 1 mM

phenylmethylsulfonyl fluoride and Pierce Universal Nuclease (Thermo Fisher

Scientific) was used. Indicated cell lines were incubated with lysis buffer for 30 min

on ice. The lysates were cleared by centrifugation at 18,000 x *g* and 4°C for 20 min

and protein concentration was determined using the BCA Protein Assay Kit from

Pierce™ (Thermo Fisher Scientific) according to the manufacturer's protocol.

Approximately 30-50 µg protein were boiled in 6x SDS loading buffer (350 mM Tris

Base pH 6.8, 38% glycerol, 10% SDS, 93 mg/ml dithiothreitol (DTT), 120 mg/ml bromophenol blue) followed by Western blot analysis. For Western blot analysis of MLKL oligomers, indicated cell lines were lysed as described above in 1% NP-40 lysis buffer (20 mM Tris-HCl pH 7.5, 50 mM NaCl, 5 mM EDTA, 10% glycerol, 1% NP-40) supplemented with PIC, 1 mM sodium orthovanadate, 5 mM sodium fluoride, 1 mM β-gylcerophosphate and Pierce Universal Nuclease. The cleared lysates were boiled in 6x SDS loading buffer or non-reducing 6x SDS loading buffer (350 mM Tris Base pH 6.8, 38% glycerol, 10% SDS, 120 mg/ml bromophenol blue), followed by Western blot analysis. The following antibodies were used in this study: human anti-M1 Ub (Genentech, San Francisco, CA, USA), rabbit anti-RNF31/HOIP (ab46322, Abcam, Cambridge, UK), mouse anti-HOIL-1 (MABC576, Merck KGaA, Darmstadt, Germany), rabbit anti-Sharpin (12541S, Cell Signaling Technology, Danvers, MA, USA), mouse anti-phospho-IκBα (S32/S36) (9246L, Cell Signaling Technology), rabbit anti-IκBα (9242S, Cell Signaling Technology), rabbit anti-phospho-RIPK1 (S166) (65746S, Cell Signaling Technology), rabbit anti-phospho-RIPK1 (S166) (31122S, Cell Signaling Technology), mouse anti-RIPK1 (551041, BD Biosciences, San Jose, CA, USA), mouse anti-RIPK1 (610459, BD Biosciences), rabbit anti-phospho-RIPK3 (S277) (ab209384, Abcam), rabbit anti-RIPK3 (13526S, Cell Signaling Technology), rabbit anti-phospho-MLKL (S358) (91689S, Cell Signaling Technology), rabbit anti-phospho-MLKL (S345) (ab196436, Abcam), rabbit anti-MLKL (14993S, Cell Signaling Technology), rabbit anti-MLKL (orb32399, Biorbyt Ltd., Cambridge, UK), mouse anti-ALIX (634502, BioLegend, San Diego, CA, USA), mouse anti-CD9 (sc-13118, Santa Cruz Biotechnologies, Santa Cruz, CA, USA), mouse anti-Flotillin-1 (610820, BD Biosciences), mouse anti-Flotillin-2 (610383, BD Biosciences), mouse anti-GAPDH (5G4cc, HyTest Ltd, Turku, Finland), mouse anti-Vinculin (V9131, Merck KGaA), mouse anti-β-actin (A5441, Merck KGaA). The

following horseradish peroxidase (HRP)-coupled secondary antibodies were used for

detection using Pierce™ ECL Western Blotting-Substrate (Thermo Fisher Scientific):

HRP-conjugated goat anti-mouse IgG (ab6789, Abcam), HRP-conjugated goat anti-

rabbit IgG (ab6721, Abcam), HRP-conjugated goat anti-human IgG (AP309P, Merck

KGaA). For stripping and re-probing of Western blot membranes, membranes were

incubated with 0.4 M NaOH solution followed by blocking for 1 h. Representative

blots of at least two independent experiments are shown. If the samples of one

experiment are detected on multiple Western blotting membranes only one

representative loading control is shown for clarity.

RIPK3 immunoprecipitation

For RIPK3 immunoprecipitations, HT-29 RIPK3 KO cells re-expressing PAM-mutated

Dox-inducible Strep-tagged RIPK3 WT [116] were seeded in full medium and RIPK3

expression was induced by adding 1 µg/mL Dox 6 h after seeding. Prior to treatment,

the medium was changed to fresh medium and the cells were lysed in 1% NP-40

lysis buffer after treatment (as described above). For Strep-RIPK3

immunoprecipitations, MagStrep "type3" XT beads (IBA Lifesciences GmbH,

Göttingen, Germany) were pre-washed twice in 1% NP-40 lysis buffer and incubated

with 1,500-1,800 µg protein over night at 4°C on a rotating wheel. On the next day,

the beads were washed five times with TBS-T (20 mM Tris, 150 mM NaCl,

0.1%Tween 20, pH 8.0) using a magnetic separator and then boiled in 2x SDS

loading buffer for 5 min at 96°C followed by Western blotting.

UBAN pulldowns

M1 ubiquitinated proteins were enriched using GST-tagged UBAN-coupled

Glutathione MagBeads (Genscript, Piscataway NJ, SA), as described previously [116].

Briefly, cell lysates were prepared in 1% NP-40 lysis buffer (as described above).

1,800-2,000 µg protein were incubated with pre-washed (NP-40 lysis buffer) GST-

UBAN beads overnight on a rotating wheel at 4°C. On the following day, the beads

were washed five times with TBS-T using a magnetic separator, then boiled in 2x

SDS loading buffer for 5 min at 96°C and analyzed by Western blotting. Equal

loading of the GST-UBAN beads was confirmed using ponceau S (Merck KGaA)

staining.

Isolation of exosomes

For the enrichment of exosomes, the indicated HT-29 cell lines were seeded in full

medium the day before treatment. Prior to the addition of compounds, the medium

was removed, cells were washed with PBS and fresh FBS-free medium was added.

After treatment, the cell culture supernatant was collected for exosome isolation and

cells were lysed in RIPA lysis buffer as described above. The supernatant was

cleared by centrifugation (2,000 x g, 30 min 4°C) in order to remove cell debris.

Exosome isolation was performed using the Total Exosome Isolation Reagent (from

cell culture media) (Thermo Fisher Scientific) according to the manufacturer's

instructions. The cleared lysates were mixed with 0.5 volumes of the Total Exosome

Isolation reagent and incubated over night at 4°C. The next day, the samples were

centrifuged for 1 h at 10,000 x g at 4°C. The supernatants were discarded and the

pelleted exosomes were lysed in RIPA lysis buffer, followed by incubation in an

appropriate amount of 6x SDS loading buffer at 96°C for 5 min. Exosome lysates and

cell lysates were analyzed by Western blotting.

Cell fractionation

Cell fractionation by phase separation using Triton X-114 lysis buffer was performed

as described before [17]. In brief, indicated cell lines were treated and lysed in Triton-

X114 (MilliporeSigma) lysis buffer (150 mM NaCl, 20 mM HEPES pH 7.4, 2% Triton

X-114), supplemented with PIC, 1 mM sodium orthovanadate, 5 mM sodium fluoride

and 1 mM β-gylcerophosphate, for 30 min on ice, followed by 10 min centrifugation at

15,000 x g at 4°C. The supernatant was warmed at 30°C for 4 min and centrifuged at

1,500 x g for 5 min. The aqueous layer of the two-phase solution was collected and

re-centrifuged at 1,500 x g for 5 min to remove contaminations from the detergent-

enriched phase. The detergent-enriched fraction was washed twice with basal buffer

(150 mM NaCl, 20 mM HEPES pH 7.4) by centrifugation 1,500 x g for 5 min and then

diluted in basal buffer. The protein concentrations of both the aqueous and

detergent-enriched fractions were determined and 40-50 µg of protein were

incubated in an appropriate amount of 6x SDS loading buffer at 96°C for 5 min,

followed by Western blotting.

RNA isolation, cDNA synthesis and quantitative real-time PCR

Total RNA was prepared using the peqGOLD total RNA isolation kit (VWR, Radnor,

PA, USA) according to manufacturer's protocol. In brief, cell lysates were prepared

using RNA lysis buffer, transferred to peqGOLD RNA Homogenizer Columns and

centrifuged at 13,000 x g for 1 min. The filtrates were mixed with an equal volume of

70% ethanol and loaded to a peqGOLD RNA Mini Column followed by centrifugation

at 10,000 xg for 1 min. After a washing step with RNA Wash Buffer I and two washing

steps with 80% ethanol (centrifugation at 10,000 x g), the columns were dried by

centrifugation at 12,000 x g for 2 min. The RNA was eluted in nuclease-free water by

centrifugation at 12,000 x g for 2 min. For cDNA synthesis, 1 µg of RNA was

transcribed using the RevertAid H Minus First Strand Kit (ThermoFisher Scientific)

according to the manufacturer's protocol. CXCL1, CXCL10, TNFα and ICAM1 gene

expression levels were determined using SYBR green-based quantitative real-time

PCR (Applied Biosystems, Darmstadt, Germany) using the 7900GR Fast Real-time

PCR system (Applied Biosystems). Data were normalized against 18S-rRNA and

RPII expression and relative gene expression levels were calculated using the ΔΔCt-

method. The primers used in this study are listed below and were obtained from

Eurofins (Hamburg, Germany). CXCL1: forward AACCGAAGTCATAGCCACAC,

reverse GTTGGATTTGTCACTGTTCAGC. CXCL10: forward

CTGAGCCTACAGCAGAGGAAC, reverse GATGCAGGTACAGCGTACAGT. TNFα:

forward ACAACCCTCAGACGCCACAT, reverse TCCTTTCCAGGGGAGAGAGG.

ICAM1: forward CTTCCTCACCGTGTACTGGAC, reverse

GGCAGCGTAGGGTAAGGTTC. 28S-rRNA: forward TTGAAAATCCGGGGGAGAG,

reverse ACATTGTTCCAACATGCCAG, RPII: forward GCACCACGTCCAATGACAT,

reverse GTGCGGCTGCTTCCATAA.

Quantification of CXCL1 cytokine secretion

To determine CXCL1 secretion from necroptotic cells, the indicated cell lines were

seeded in 96-well dishes and treated as described above. After treatment,

supernatants were collected and cleared by centrifugation (300 x *g*, 5 min, 4°C).

CXCL1 levels were measured after 3 h and 6 h using the Human CXCL1/GROα

DuoSet® ELISA kit (R&D Systems, Minneapolis, MN, USA) according to the

manufacturer's instructions. Absorbance at 450 nm was determined using the Infinite

M200 microplate reader (Tecan, Crailsheim, Germany).

Immunofluorescence staining

For immunofluorescence staining of phosphorylated MLKL, indicated cell lines were

seeded in CELLSTAR® 96-well tissue culture plates (Greiner Bio-One GmbH) at an

appropriate density the day before treatment. After treatment, the supernatant was

removed and the cells were fixed by incubation in 4% paraformaldehyde solution for

20 min at room temperature, followed by permeabilization for 15 min with 0.1%

Triton X-100. After three washing steps with PBS, the cells were blocked in antibody

dilution buffer (ADB) (0.9% NaCl, 10 mM Tris/HCl pH 7.5, 5 mM EDTA, 1 mg/mL

BSA) for 45 min, followed by incubation with rabbit anti-phospho-MLKL (S358)

(1:100, ab187091, Abcam) (diluted in ADB) overnight at 4°C. On the next day, the

cells were washed five times with PBS and incubated with Cy3 AffiniPure Donkey

Anti-Rabbit IgG (H+L) (1:800, 711-165-152, Jackson Immuno Research, West Grove,

PA, USA) and 0.1 µg/mL DAPI (diluted in ADB) for 90 min at room temperature. After

five washing steps with PBS the cells were analyzed using the ImageXpress® Micro

XLS Widefield High-Content Analysis System and MetaXpress® Software (Molecular

Devices).

Massive Analysis of cDNA Ends (MACE-seq)

Massive Analyses of cDNA Ends (MACE-Seq) was performed by GenXPro GmbH in

Frankfurt am Main using the MACE-Seq kit according to the manual of the

manufacturers. Briefly, cDNA was generated from fragmented RNA with barcoded

poly-A primers during reverse transcription. After second-strand synthesis and 5'

adapter integration, a PCR with minimum number of cylces was used to produce a

library that was sequenced on an Illumina NextSeq500 machine with 1x 75 bps [117].

MACE-seq analysis

Raw MACE-seq data was preprocessed using Cutadapt [118] to eliminate poly-A-tails

as well as bad-quality base reads. FastQC was used to assess the quality of

sequencing after trimming. Cleaned reads were mapped and quantified with a

reference genome using STAR. ENSEMBL-GTF data were used to provide genomic

locations for quantification as well as additional data for annotation (such as gene

name, gene description, Gene Ontology (GO)-Terms etc). Differential expression

analysis was performed using DESeq2 [119], which is based on negative binomial

generalized linear models. Results were compiled into a final table including

significance parameters (pvalue, FDR) and log2FoldChanges. BigWig track files were

generated by deepTools. Final data visualization of the significantly expressed and

the differentially regulated genes was performed by custom R-Scripts.

1003

1004 ***Statistical analysis***

1005 Statistical significance was determined with the GraphPad Prism 9 Software

1006 (GraphPad Software, La Jolla CA, USA) using 2-way ANOVA followed by Tukey's

1007 multiple comparisons test. *P* values < 0.05 are considered significant and depicted as

1008 follows: *P* ≤ 0.05: *, *P* ≤ 0.01: **, *P* ≤ 0.001: ***, *P* ≤ 0.0001: ****, ns: not significant.

1009

1010 **Data availability**

1011 MACE-seq data have been deposited at GEO and are publicly available as of the

1012 date of publication. All primary data will be shared by the lead contact upon request.

 Supplemental Information

 Figure S1. Related to Figure 1. A. Quantification of cell death in untreated (UT) and

 Gliotoxin (1 µM)-pre-treated HT-29 cells after treatment with TBZ (10 ng/mL TNFα, 1

 µM BV6, 20 µM zVAD.fmk) for the indicated time points. Mean and SEM of at least

 three independent experiments are shown. ****P<0.0001. **B.** Quantification of cell

 death in untreated (UT) and Gliotoxin (1 µM)-pre-treated THP-1 cells after treatment

 with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for the indicated time points.

 Mean and SEM of at least three independent experiments are shown. ***P<0.001;

 ****P<0.0001. **C.** Western blot analysis of HOIP levels in control (siCtrl) and HOIP

 knockdown (siHOIP) HT-29 cells. GAPDH was used as loading control.

 Representative blots of at least two independent experiments are shown. Related to

 Figure 1C. **D.** Western blot analysis of total M1 Ub levels in control or HOIPIN-8 (30

 µM)-pre-treated THP-1 cells treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

 zVAD.fmk) for 4 h. GAPDH was used as loading control. Representative blots of at

 least two independent experiments are shown. **E.** Amino acid sequence alignment

 showing conservation of human (aa841-aa960) and murine HOIP (RNF31) (aa835-

 aa954). Conserved residues interacting with HOIPIN-8 are highlighted and indicated

 by arrowheads. Sequence alignment was performed using the CLUSTAL O (1.2.4)

 algorithm. **F.** Western blot analysis of phosphorylated and total IκBα levels in control

 or HOIPIN-8 (30 µM)-pre-treated HT-29 cells upon treatment with 10 ng/mL TNFα for

 the indicated time points. Vinculin was used as loading control. Representative blots

 of at least two independent experiments are shown. **G.** Western blot analysis of

 phosphorylated and total IκBα levels in control or HOIPIN-8 (30 µM)-pre-treated

 MEFs upon treatment with 10 ng/mL TNFα for the indicated time. Vinculin was used

 as loading control. Representative blots of at least two independent experiments are

 shown. **H.** Quantification of cell death in untreated (UT) and HOIPIN-8 (30 µM) pre-

treated MEFs after treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for the indicated time points. Mean and SEM of at least three independent

experiments are shown. ***P<0.001; ****P<0.0001; ns not significant. **I.** Quantification

of cell death in untreated (UT) and HOIPIN-8 (30 µM)-pre-treated, GSK'872 (20 µM)-

pre-treated or HOIPIN-8 and GSK'872-pre-treated L-929 cells after treatment with

TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for the indicated time points.

Mean and SEM of at least three independent experiments are shown. ****P<0.0001;

ns not significant. **J.** Western blot analysis of total M1 Ub levels in control or HOIPIN-

8 (30 µM)-pre-treated MEFs treated with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for 3 h. GAPDH was used as loading control. Representative blots of at

least two independent experiments are shown.

1050

Figure S2. Related to Figure 2. A. Western blot analysis of LUBAC subunits HOIP

and HOIL-1 expression levels in control *versus* HOIPIN-8 (30 µM)-pre-treated MEFs

upon treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 3 h.

GAPDH was used as loading control. Representative blots of at least two

independent experiments are shown. **B.** Western blot analysis of expression levels of

phosphorylated and total RIPK1 and MLKL upon 4 h TBZ (10 ng/mL TNFα, 1 µM

BV6, 20 µM zVAD.fmk) treatment of THP-1 cells pre-treated with HOIPIN-8 (30 µM).

Vinculin was used as loading control. Representative blots of at least two

independent experiments are shown. **C.** Western blot analysis of expression levels of

phosphorylated and total RIPK1, RIPK3 and MLKL, and of M1 Ub in TBZ-treated (10

ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) control (nht), RIPK3 KO and MLKL KO

HT-29 cells pre-treated with HOIPIN-8 (30 µM). GAPDH was used as loading control.

Representative blots of at least two independent experiments are shown.

1064

Figure S3. Related to Figure 3. A. mRNA expression levels of CXCL1 and CXCL10

of control (nht) or MLKL KO HT-29 cells treated with TBZ (10 ng/mL TNFα, 1 µM

BV6, 20 µM zVAD.fmk) for 3 h. Gene expression was normalized against 18S and

RPII mRNA expression and is presented as x-fold mRNA expression compared to

the untreated (UT) control cells. Mean and SEM of at least three independent

experiments are shown. **$P<0.01$; ***$P<0.001$. **B.** mRNA expression levels of TNFα

of control (UT), HOIPIN-8 (30 µM) or NSA (10 µM)-pre-treated HT-29 cells upon

treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 3 h. Gene

expression was normalized against 18S and RPII mRNA expression and is presented

as x-fold mRNA expression compared to the untreated (UT) control. Mean and SEM

of at least three independent experiments are shown. ****$P<0.0001$. **C.** mRNA

expression levels of ICAM1 of control (UT), HOIPIN-8 (30 µM) or NSA (10 µM)-pre-

treated HT-29 cells upon treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM

zVAD.fmk) for 3 h. Gene expression was normalized against 18S and RPII mRNA

expression and is presented as x-fold mRNA expression compared to the untreated

(UT) control. Mean and SEM of at least three independent experiments are shown.

****$P<0.0001$.

1082

Figure S4. Related to Figure 4. A. Distribution of M1 poly-Ub in the micelle-poor

(aqueous) and micelle-rich (detergent) fraction after phase separation using Triton X-

114 lysis buffer in untreated (UT) and HOIPIN-8 (30 µM)-pre-treated HT-29 cells

upon treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h.

GAPDH was used as loading control for soluble proteins and CD9 as loading control

for membrane proteins. Representative blots of at least two independent experiments

are shown. **B.** Fluorescence microscopy images show localization of phosphorylated

MLKL (green) in untreated (UT) and NSA (10 µM)-pre-treated HT-29 cells and after

treatment with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 3 h. Nuclei

were stained with DAPI (blue). Representative images of at least two independent

experiments are shown. Scale bar 50 µm.

1094

Figure S5. Related to Figure 5. A. GST-UBAN-mediated pulldown of M1

ubiquitinated proteins in control or HOIPIN-8 (30 µM)-pre-treated THP-1 cells treated

with TBZ (10 ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h. GST and GAPDH

were used as loading controls. Representative blots of at least two independent

experiments are shown. **B.** Western blot analysis of FLOT1/2 levels in control (siCtrl)

and FLOT1 (siFLOT1), FLOT2 (siFLOT2) or combined FLOT1/2 (siFLOT1+2)

knockdown HT-29 cells. Vinculin was used as loading control. Representative blots of

at least two independent experiments are shown. Related to Figure 5B. **C.** Western

blot analysis of total M1 Ub levels of control or HOIPIN-8 (30 µM)-pre-treated control

(siCtrl) or FLOT1/2 (siFLOT1+2) knockdown HT-29 cells after treatment with TBZ (10

ng/mL TNFα, 1 µM BV6, 20 µM zVAD.fmk) for 4 h. GAPDH was used as loading

control. Representative blots of at least two independent experiments are shown.

1107

1108   **Figure S6. Related to Figure 6. A.** Representative images of time-lapse videos of

1109   untreated (UT), Birinapant (20 µM) and Emricasan (10 µM) (BiE)-treated, HOIPIN-8

1110   (30 µM)-pre-treated and BiE-treated (BiE+HOIPIN-8) and NSA (10 µM)-pre-treated

1111   and BiE-treated (BiE+NSA) primary hPOs for a total period of 23.5 h. Scalebars: 250

1112   µm. **B.** *Idem* as A., but treated primary hPOs were stained with Hoechst33342 (Hoe)

(blue), FDA (live; green) and PI (dead; red) after 24 h treatment with BiE, BiE+HOIPIN-8 or BiE+NSA prior to imaging. Scalebars: 250 µm.

Figure S7. Related to Figure 6. Representative images of untreated (UT), BV6 (5 µM) and zVAD.fmk (20 µM) (BZ)-treated and HOIPIN-8 (30 µM)-pre-treated and BZ-treated (BZ+HOIPIN-8) primary hPOs that were stained with Hoechst33342 (Hoe) (blue), FDA (live; green) and PI (dead; red) after 24 h treatment prior to imaging. Scalebars: 250 µm.

Movie 1. Related to Figure 6A. Time-lapse videos of untreated (UT), BV6 and Emricasan (BE)-treated and HOIPIN-8-pre-treated and BE (BE+HOIPIN-8)-treated primary hPOs for a total period of 23.5 h. Scalebars: 250 µm.

Movie 2. Related to Figure S6A. Time-lapse videos of untreated (UT), Birinapant and Emricasan (BiE)-treated, HOIPIN-8-pre-treated and BiE (BiE+HOIPIN-8)-treated and NSA-pre-treated and BiE (BiE+NSA)-treated primary hPOs for a total period of 23.5 h. Scalebars: 250 µm.

References

1. Zhu, K., Liang, W., Ma, Z., Xu, D., Cao, S., Lu, X., Liu, N., Shan, B., Qian, L., and Yuan, J. (2018). Necroptosis promotes cell-autonomous activation of proinflammatory cytokine gene expression. Cell Death Dis *9*, 500. 10.1038/s41419-018-0524-y.

2. Kaczmarek, A., Vandenabeele, P., and Krysko, D.V. (2013). Necroptosis: the release of damage-associated molecular patterns and its physiological relevance. Immunity *38*, 209-223. 10.1016/j.immuni.2013.02.003.

3. Weinlich, R., Oberst, A., Beere, H.M., and Green, D.R. (2017). Necroptosis in development, inflammation and disease. Nat Rev Mol Cell Biol *18*, 127-136. 10.1038/nrm.2016.149.

4. Pasparakis, M., and Vandenabeele, P. (2015). Necroptosis and its role in inflammation. Nature *517*, 311-320. 10.1038/nature14191.

5. Emmerich, C.H., Bakshi, S., Kelsall, I.R., Ortiz-Guerrero, J., Shpiro, N., and Cohen, P. (2016). Lys63/Met1-hybrid ubiquitin chains are commonly formed during the activation of innate immune signalling. Biochem Biophys Res Commun *474*, 452-461. 10.1016/j.bbrc.2016.04.141.

6. Emmerich, C.H., Ordureau, A., Strickson, S., Arthur, J.S., Pedrioli, P.G., Komander, D., and Cohen, P. (2013). Activation of the canonical IKK complex by K63/M1-linked hybrid ubiquitin chains. Proc Natl Acad Sci U S A *110*, 15247-15252. 10.1073/pnas.1314715110.

7. Bertrand, M.J., Milutinovic, S., Dickson, K.M., Ho, W.C., Boudreault, A., Durkin, J., Gillard, J.W., Jaquith, J.B., Morris, S.J., and Barker, P.A. (2008). cIAP1 and cIAP2 facilitate cancer cell survival by functioning as E3 ligases that promote RIP1 ubiquitination. Mol Cell *30*, 689-700. 10.1016/j.molcel.2008.05.014.

8.   Draber, P., Kupka, S., Reichert, M., Draberova, H., Lafont, E., de Miguel, D.,

Spilgies, L., Surinova, S., Taraborrelli, L., Hartwig, T., et al. (2015). LUBAC-

Recruited CYLD and A20 Regulate Gene Activation and Cell Death by

Exerting Opposing Effects on Linear Ubiquitin in Signaling Complexes. Cell

Rep *13*, 2258-2272. 10.1016/j.celrep.2015.11.009.

9.   Wertz, I.E., Newton, K., Seshasayee, D., Kusam, S., Lam, C., Zhang, J.,

Popovych, N., Helgason, E., Schoeffler, A., Jeet, S., et al. (2015).

Phosphorylation and linear ubiquitin direct A20 inhibition of inflammation.

Nature *528*, 370-375. 10.1038/nature16165.

10.   Mompean, M., Li, W., Li, J., Laage, S., Siemer, A.B., Bozkurt, G., Wu, H., and

McDermott, A.E. (2018). The Structure of the Necrosome RIPK1-RIPK3 Core,

a Human Hetero-Amyloid Signaling Complex. Cell *173*, 1244-1253 e1210.

10.1016/j.cell.2018.03.032.

11.   Li, J., McQuade, T., Siemer, A.B., Napetschnig, J., Moriwaki, K., Hsiao, Y.S.,

Damko, E., Moquin, D., Walz, T., McDermott, A., et al. (2012). The RIP1/RIP3

necrosome forms a functional amyloid signaling complex required for

programmed necrosis. Cell *150*, 339-350. 10.1016/j.cell.2012.06.019.

12.   Cho, Y.S., Challa, S., Moquin, D., Genga, R., Ray, T.D., Guildford, M., and

Chan, F.K. (2009). Phosphorylation-driven assembly of the RIP1-RIP3

complex regulates programmed necrosis and virus-induced inflammation. Cell

137, 1112-1123. 10.1016/j.cell.2009.05.037.

13.   He, S., Wang, L., Miao, L., Wang, T., Du, F., Zhao, L., and Wang, X. (2009).

Receptor interacting protein kinase-3 determines cellular necrotic response to

TNF-alpha. Cell *137*, 1100-1111. 10.1016/j.cell.2009.05.021.

14.   Sun, L., Wang, H., Wang, Z., He, S., Chen, S., Liao, D., Wang, L., Yan, J., Liu,

W., Lei, X., and Wang, X. (2012). Mixed lineage kinase domain-like protein

1183 mediates necrosis signaling downstream of RIP3 kinase. Cell *148*, 213-227.

1184 10.1016/j.cell.2011.11.031.

1185 15. Murphy, J.M., Czabotar, P.E., Hildebrand, J.M., Lucet, I.S., Zhang, J.G.,

1186 Alvarez-Diaz, S., Lewis, R., Lalaoui, N., Metcalf, D., Webb, A.I., et al. (2013).

1187 The pseudokinase MLKL mediates necroptosis via a molecular switch

1188 mechanism. Immunity *39*, 443-453. 10.1016/j.immuni.2013.06.018.

1189 16. Su, L., Quade, B., Wang, H., Sun, L., Wang, X., and Rizo, J. (2014). A plug

1190 release mechanism for membrane permeation by MLKL. Structure *22*, 1489-

1191 1500. 10.1016/j.str.2014.07.014.

1192 17. Wang, H., Sun, L., Su, L., Rizo, J., Liu, L., Wang, L.F., Wang, F.S., and Wang,

1193 X. (2014). Mixed lineage kinase domain-like protein MLKL causes necrotic

1194 membrane disruption upon phosphorylation by RIP3. Mol Cell *54*, 133-146.

1195 10.1016/j.molcel.2014.03.003.

1196 18. Chen, X., Li, W., Ren, J., Huang, D., He, W.T., Song, Y., Yang, C., Li, W.,

1197 Zheng, X., Chen, P., and Han, J. (2014). Translocation of mixed lineage

1198 kinase domain-like protein to plasma membrane leads to necrotic cell death.

1199 Cell Res *24*, 105-121. 10.1038/cr.2013.171.

1200 19. Hildebrand, J.M., Tanzer, M.C., Lucet, I.S., Young, S.N., Spall, S.K., Sharma,

1201 P., Pierotti, C., Garnier, J.M., Dobson, R.C., Webb, A.I., et al. (2014).

1202 Activation of the pseudokinase MLKL unleashes the four-helix bundle domain

1203 to induce membrane localization and necroptotic cell death. Proc Natl Acad

1204 Sci U S A *111*, 15072-15077. 10.1073/pnas.1408987111.

1205 20. Dondelinger, Y., Declercq, W., Montessuit, S., Roelandt, R., Goncalves, A.,

1206 Bruggeman, I., Hulpiau, P., Weber, K., Sehon, C.A., Marquis, R.W., et al.

1207 (2014). MLKL compromises plasma membrane integrity by binding to

phosphatidylinositol phosphates. Cell Rep *7*, 971-981.

10.1016/j.celrep.2014.04.026.

21.   Samson, A.L., Zhang, Y., Geoghegan, N.D., Gavin, X.J., Davies, K.A.,

Mlodzianoski, M.J., Whitehead, L.W., Frank, D., Garnish, S.E., Fitzgibbon, C.,

et al. (2020). MLKL trafficking and accumulation at the plasma membrane

control the kinetics and threshold for necroptosis. Nat Commun *11*, 3151.

10.1038/s41467-020-16887-1.

22.   Gong, Y.N., Guy, C., Olauson, H., Becker, J.U., Yang, M., Fitzgerald, P.,

Linkermann, A., and Green, D.R. (2017). ESCRT-III Acts Downstream of

MLKL to Regulate Necroptotic Cell Death and Its Consequences. Cell *169*,

286-300 e216. 10.1016/j.cell.2017.03.020.

23.   Cai, Z., Jitkaew, S., Zhao, J., Chiang, H.C., Choksi, S., Liu, J., Ward, Y., Wu,

L.G., and Liu, Z.G. (2014). Plasma membrane translocation of trimerized

MLKL protein is required for TNF-induced necroptosis. Nat Cell Biol *16*, 55-65.

10.1038/ncb2883.

24.   Seo, J., Nam, Y.W., Kim, S., Oh, D.B., and Song, J. (2021). Necroptosis

molecular mechanisms: Recent findings regarding novel necroptosis

regulators. Exp Mol Med *53*, 1007-1017. 10.1038/s12276-021-00634-7.

25.   Karlowitz, R., and van Wijk, S.J.L. (2021). Surviving death: emerging concepts

of RIPK3 and MLKL ubiquitination in the regulation of necroptosis. FEBS J.

10.1111/febs.16255.

26.   Samson, A.L., Garnish, S.E., Hildebrand, J.M., and Murphy, J.M. (2021).

Location, location, location: A compartmentalized view of TNF-induced

necroptotic signaling. Sci Signal *14*. 10.1126/scisignal.abc6178.

27.   Akutsu, M., Dikic, I., and Bremm, A. (2016). Ubiquitin chain diversity at a

glance. J Cell Sci *129*, 875-880. 10.1242/jcs.183954.

28.   Komander, D., and Rape, M. (2012). The ubiquitin code. Annu Rev Biochem

1235   *81*, 203-229. 10.1146/annurev-biochem-060310-170328.

1236   29.   Swatek, K.N., and Komander, D. (2016). Ubiquitin modifications. Cell Res *26*,

1237   399-422. 10.1038/cr.2016.39.

1238   30.   Gerlach, B., Cordier, S.M., Schmukle, A.C., Emmerich, C.H., Rieser, E., Haas,

1239   T.L., Webb, A.I., Rickard, J.A., Anderton, H., Wong, W.W., et al. (2011). Linear

1240   ubiquitination prevents inflammation and regulates immune signalling. Nature

1241   *471*, 591-596. 10.1038/nature09816.

1242   31.   Ikeda, F., Deribe, Y.L., Skanland, S.S., Stieglitz, B., Grabbe, C., Franz-

1243   Wachtel, M., van Wijk, S.J., Goswami, P., Nagy, V., Terzic, J., et al. (2011).

1244   SHARPIN forms a linear ubiquitin ligase complex regulating NF-kappaB

1245   activity and apoptosis. Nature *471*, 637-641. 10.1038/nature09814.

1246   32.   Tokunaga, F., Nakagawa, T., Nakahara, M., Saeki, Y., Taniguchi, M., Sakata,

1247   S., Tanaka, K., Nakano, H., and Iwai, K. (2011). SHARPIN is a component of

1248   the NF-kappaB-activating linear ubiquitin chain assembly complex. Nature

1249   *471*, 633-636. 10.1038/nature09815.

1250   33.   Kirisako, T., Kamei, K., Murata, S., Kato, M., Fukumoto, H., Kanie, M., Sano,

1251   S., Tokunaga, F., Tanaka, K., and Iwai, K. (2006). A ubiquitin ligase complex

1252   assembles linear polyubiquitin chains. EMBO J *25*, 4877-4887.

1253   10.1038/sj.emboj.7601360.

1254   34.   Stieglitz, B., Rana, R.R., Koliopoulos, M.G., Morris-Davies, A.C., Schaeffer, V.,

1255   Christodoulou, E., Howell, S., Brown, N.R., Dikic, I., and Rittinger, K. (2013).

1256   Structural basis for ligase-specific conjugation of linear ubiquitin chains by

1257   HOIP. Nature *503*, 422-426. 10.1038/nature12638.

1258   35.   Smit, J.J., Monteferrario, D., Noordermeer, S.M., van Dijk, W.J., van der

1259   Reijden, B.A., and Sixma, T.K. (2012). The E3 ligase HOIP specifies linear

ubiquitin chain assembly through its RING-IBR-RING domain and the unique

LDD extension. EMBO J *31*, 3833-3844. 10.1038/emboj.2012.217.

36.   Lechtenberg, B.C., Rajput, A., Sanishvili, R., Dobaczewska, M.K., Ware, C.F.,

Mace, P.D., and Riedl, S.J. (2016). Structure of a HOIP/E2~ubiquitin complex

reveals RBR E3 ligase mechanism and regulation. Nature *529*, 546-550.

10.1038/nature16511.

37.   Oikawa, D., Sato, Y., Ohtake, F., Komakura, K., Hanada, K., Sugawara, K.,

Terawaki, S., Mizukami, Y., Phuong, H.T., Iio, K., et al. (2020). Molecular

bases for HOIPINs-mediated inhibition of LUBAC and innate immune

responses. Commun Biol *3*, 163. 10.1038/s42003-020-0882-8.

38.   Katsuya, K., Oikawa, D., Iio, K., Obika, S., Hori, Y., Urashima, T., Ayukawa,

K., and Tokunaga, F. (2019). Small-molecule inhibitors of linear ubiquitin chain

assembly complex (LUBAC), HOIPINs, suppress NF-kappaB signaling.

Biochem Biophys Res Commun *509*, 700-706. 10.1016/j.bbrc.2018.12.164.

39.   Hrdinka, M., and Gyrd-Hansen, M. (2017). The Met1-Linked Ubiquitin

Machinery: Emerging Themes of (De)regulation. Mol Cell *68*, 265-280.

10.1016/j.molcel.2017.09.001.

40.   Hrdinka, M., Fiil, B.K., Zucca, M., Leske, D., Bagola, K., Yabal, M., Elliott,

P.R., Damgaard, R.B., Komander, D., Jost, P.J., and Gyrd-Hansen, M. (2016).

CYLD Limits Lys63- and Met1-Linked Ubiquitin at Receptor Complexes to

Regulate Innate Immune Signaling. Cell Rep *14*, 2846-2858.

10.1016/j.celrep.2016.02.062.

41.   Boisson, B., Laplantine, E., Dobbs, K., Cobat, A., Tarantino, N., Hazen, M.,

Lidov, H.G., Hopkins, G., Du, L., Belkadi, A., et al. (2015). Human HOIP and

LUBAC   deficiency   underlies   autoinflammation,   immunodeficiency,

amylopectinosis, and lymphangiectasia. J Exp Med *212*, 939-951.

10.1084/jem.20141130.

42. Boisson, B., Laplantine, E., Prando, C., Giliani, S., Israelsson, E., Xu, Z.,

Abhyankar, A., Israel, L., Trevejo-Nunez, G., Bogunovic, D., et al. (2012).

Immunodeficiency, autoinflammation and amylopectinosis in humans with

inherited HOIL-1 and LUBAC deficiency. Nat Immunol *13*, 1178-1186.

10.1038/ni.2457.

43. Oda, H., Manthiram, K., Chavan, P.P., Nakabo, S., Kuehn, H.S., Beck, D.B.,

Chae, J.J., Nehrebecky, M., Ombrello, A.K., Romeo, T., et al. (2022). Human

LUBAC deficiency leads to autoinflammation and immunodeficiency by

dysregulation in TNF-mediated cell death. medRxiv,

2022.2011.2009.22281431. 10.1101/2022.11.09.22281431.

44. Tokunaga, F., Sakata, S., Saeki, Y., Satomi, Y., Kirisako, T., Kamei, K.,

Nakagawa, T., Kato, M., Murata, S., Yamaoka, S., et al. (2009). Involvement

of linear polyubiquitylation of NEMO in NF-kappaB activation. Nat Cell Biol *11*,

123-132. 10.1038/ncb1821.

45. Seymour, R.E., Hasham, M.G., Cox, G.A., Shultz, L.D., Hogenesch, H.,

Roopenian, D.C., and Sundberg, J.P. (2007). Spontaneous mutations in the

mouse Sharpin gene result in multiorgan inflammation, immune system

dysregulation and dermatitis. Genes Immun *8*, 416-421.

10.1038/sj.gene.6364403.

46. Rickard, J.A., Anderton, H., Etemadi, N., Nachbur, U., Darding, M., Peltzer, N.,

Lalaoui, N., Lawlor, K.E., Vanyai, H., Hall, C., et al. (2014). TNFR1-dependent

cell death drives inflammation in Sharpin-deficient mice. Elife *3*.

10.7554/eLife.03464.

47. Kumari, S., Redouane, Y., Lopez-Mosqueda, J., Shiraishi, R., Romanowska,
M., Lutzmayer, S., Kuiper, J., Martinez, C., Dikic, I., Pasparakis, M., and Ikeda,
F. (2014). Sharpin prevents skin inflammation by inhibiting TNFR1-induced
keratinocyte apoptosis. Elife *3*. 10.7554/eLife.03422.

48. Peltzer, N., Rieser, E., Taraborrelli, L., Draber, P., Darding, M., Pernaute, B.,
Shimizu, Y., Sarr, A., Draberova, H., Montinaro, A., et al. (2014). HOIP
deficiency causes embryonic lethality by aberrant TNFR1-mediated
endothelial cell death. Cell Rep *9*, 153-165. 10.1016/j.celrep.2014.08.066.

49. Shimizu, Y., Peltzer, N., Sevko, A., Lafont, E., Sarr, A., Draberova, H., and
Walczak, H. (2017). The Linear ubiquitin chain assembly complex acts as a
liver tumor suppressor and inhibits hepatocyte apoptosis and hepatitis.
Hepatology *65*, 1963-1978. 10.1002/hep.29074.

50. Taraborrelli, L., Peltzer, N., Montinaro, A., Kupka, S., Rieser, E., Hartwig, T.,
Sarr, A., Darding, M., Draber, P., Haas, T.L., et al. (2018). LUBAC prevents
lethal dermatitis by inhibiting cell death induced by TNF, TRAIL and CD95L.
Nat Commun *9*, 3910. 10.1038/s41467-018-06155-8.

51. Katsuya, K., Hori, Y., Oikawa, D., Yamamoto, T., Umetani, K., Urashima, T.,
Kinoshita, T., Ayukawa, K., Tokunaga, F., and Tamaru, M. (2018). High-
Throughput Screening for Linear Ubiquitin Chain Assembly Complex (LUBAC)
Selective Inhibitors Using Homogenous Time-Resolved Fluorescence (HTRF)-
Based Assay System. SLAS Discov *23*, 1018-1029.
10.1177/2472555218793066.

52. Shimizu, Y., Taraborrelli, L., and Walczak, H. (2015). Linear ubiquitination in
immunity. Immunol Rev *266*, 190-207. 10.1111/imr.12309.

53. Fan, W., Guo, J., Gao, B., Zhang, W., Ling, L., Xu, T., Pan, C., Li, L., Chen, S.,
Wang, H., et al. (2019). Flotillin-mediated endocytosis and ALIX-syntenin-1-

mediated exocytosis protect the cell membrane from damage caused by necroptosis. Sci Signal *12*. 10.1126/scisignal.aaw3423.

54.   Bickel, P.E., Scherer, P.E., Schnitzer, J.E., Oh, P., Lisanti, M.P., and Lodish, H.F. (1997). Flotillin and epidermal surface antigen define a new family of caveolae-associated integral membrane proteins. J Biol Chem *272*, 13793-13802. 10.1074/jbc.272.21.13793.

55.   Lang, D.M., Lommel, S., Jung, M., Ankerhold, R., Petrausch, B., Laessing, U., Wiechers, M.F., Plattner, H., and Stuermer, C.A. (1998). Identification of reggie-1 and reggie-2 as plasmamembrane-associated proteins which cocluster with activated GPI-anchored cell adhesion molecules in non-caveolar micropatches in neurons. J Neurobiol *37*, 502-523. 10.1002/(sici)1097-4695(199812)37:4<502::aid-neu2>3.0.co;2-s.

56.   Lingwood, D., and Simons, K. (2010). Lipid rafts as a membrane-organizing principle. Science *327*, 46-50. 10.1126/science.1174621.

57.   Sezgin, E., Levental, I., Mayor, S., and Eggeling, C. (2017). The mystery of membrane organization: composition, regulation and roles of lipid rafts. Nat Rev Mol Cell Biol *18*, 361-374. 10.1038/nrm.2017.16.

58.   Morrow, I.C., Rea, S., Martin, S., Prior, I.A., Prohaska, R., Hancock, J.F., James, D.E., and Parton, R.G. (2002). Flotillin-1/reggie-2 traffics to surface raft domains via a novel golgi-independent pathway. Identification of a novel membrane targeting domain and a role for palmitoylation. J Biol Chem *277*, 48834-48841. 10.1074/jbc.M209082200.

59.   Solis, G.P., Hoegg, M., Munderloh, C., Schrock, Y., Malaga-Trillo, E., Rivera-Milla, E., and Stuermer, C.A. (2007). Reggie/flotillin proteins are organized into stable tetramers in membrane microdomains. Biochem J *403*, 313-322. 10.1042/BJ20061686.

60.   Frick, M., Bright, N.A., Riento, K., Bray, A., Merrified, C., and Nichols, B.J.

(2007). Coassembly of flotillins induces formation of membrane microdomains,

membrane curvature, and vesicle budding. Curr Biol *17*, 1151-1156.

10.1016/j.cub.2007.05.078.

61.   Glebov, O.O., Bright, N.A., and Nichols, B.J. (2006). Flotillin-1 defines a

clathrin-independent endocytic pathway in mammalian cells. Nat Cell Biol *8*,

46-54. 10.1038/ncb1342.

62.   Langhorst, M.F., Solis, G.P., Hannbeck, S., Plattner, H., and Stuermer, C.A.

(2007). Linking membrane microdomains to the cytoskeleton: regulation of the

lateral mobility of reggie-1/flotillin-2 by interaction with actin. FEBS Lett *581*,

4697-4703. 10.1016/j.febslet.2007.08.074.

63.   Rossy, J., Schlicht, D., Engelhardt, B., and Niggli, V. (2009). Flotillins interact

with PSGL-1 in neutrophils and, upon stimulation, rapidly organize into

membrane domains subsequently accumulating in the uropod. PLoS One *4*,

e5403. 10.1371/journal.pone.0005403.

64.   Affentranger, S., Martinelli, S., Hahn, J., Rossy, J., and Niggli, V. (2011).

Dynamic reorganization of flotillins in chemokine-stimulated human T-

lymphocytes. BMC Cell Biol *12*, 28. 10.1186/1471-2121-12-28.

65.   Solis, G.P., Schrock, Y., Hulsbusch, N., Wiechers, M., Plattner, H., and

Stuermer, C.A. (2012). Reggies/flotillins regulate E-cadherin-mediated cell

contact formation by affecting EGFR trafficking. Mol Biol Cell *23*, 1812-1825.

10.1091/mbc.E11-12-1006.

66.   Guillaume, E., Comunale, F., Do Khoa, N., Planchon, D., Bodin, S., and

Gauthier-Rouviere, C. (2013). Flotillin microdomains stabilize cadherins at cell-

cell junctions. J Cell Sci *126*, 5293-5304. 10.1242/jcs.133975.

67.   Amaddii, M., Meister, M., Banning, A., Tomasovic, A., Mooz, J., Rajalingam,

K., and Tikkanen, R. (2012). Flotillin-1/reggie-2 protein plays dual role in

activation of receptor-tyrosine kinase/mitogen-activated protein kinase

signaling. J Biol Chem *287*, 7265-7278. 10.1074/jbc.M111.287599.

68.   Pust, S., Klokk, T.I., Musa, N., Jenstad, M., Risberg, B., Erikstein, B.,

Tcatchoff, L., Liestol, K., Danielsen, H.E., van Deurs, B., and Sandvig, K.

(2013). Flotillins as regulators of ErbB2 levels in breast cancer. Oncogene *32*,

3443-3451. 10.1038/onc.2012.357.

69.   Damgaard, R.B., Elliott, P.R., Swatek, K.N., Maher, E.R., Stepensky, P.,

Elpeleg, O., Komander, D., and Berkun, Y. (2019). OTULIN deficiency in

ORAS causes cell type-specific LUBAC degradation, dysregulated TNF

signalling and cell death. EMBO Mol Med *11*, e9324.

10.15252/emmm.201809324.

70.   Damgaard, R.B., Walker, J.A., Marco-Casanova, P., Morgan, N.V.,

Titheradge, H.L., Elliott, P.R., McHale, D., Maher, E.R., McKenzie, A.N.J., and

Komander, D. (2016). The Deubiquitinase OTULIN Is an Essential Negative

Regulator of Inflammation and Autoimmunity. Cell *166*, 1215-1230 e1220.

10.1016/j.cell.2016.07.019.

71.   Sakamoto, H., Egashira, S., Saito, N., Kirisako, T., Miller, S., Sasaki, Y.,

Matsumoto, T., Shimonishi, M., Komatsu, T., Terai, T., et al. (2015). Gliotoxin

suppresses NF-kappaB activation by selectively inhibiting linear ubiquitin chain

assembly complex (LUBAC). ACS Chem Biol *10*, 675-681.

10.1021/cb500653y.

72.   Peltzer, N., Darding, M., Montinaro, A., Draber, P., Draberova, H., Kupka, S.,

Rieser, E., Fisher, A., Hutchinson, C., Taraborrelli, L., et al. (2018). LUBAC is

essential for embryogenesis by preventing cell death and enabling
haematopoiesis. Nature *557*, 112-117. 10.1038/s41586-018-0064-8.

73. Vanlangenakker, N., Bertrand, M.J., Bogaert, P., Vandenabeele, P., and
1415 Vanden Berghe, T. (2011). TNF-induced necroptosis in L929 cells is tightly
1416 regulated by multiple TNFR1 complex I and II members. Cell Death Dis *2*,
1417 e230. 10.1038/cddis.2011.111.

74. Zargarian, S., Shlomovitz, I., Erlich, Z., Hourizadeh, A., Ofir-Birin, Y., Croker,
1419 B.A., Regev-Rudzki, N., Edry-Botzer, L., and Gerlic, M. (2017).
1420 Phosphatidylserine externalization, "necroptotic bodies" release, and
1421 phagocytosis during necroptosis. PLoS Biol *15*, e2002711.
1422 10.1371/journal.pbio.2002711.

75. Yoon, S., Kovalenko, A., Bogdanov, K., and Wallach, D. (2017). MLKL, the
1424 Protein that Mediates Necroptosis, Also Regulates Endosomal Trafficking and
1425 Extracellular Vesicle Generation. Immunity *47*, 51-65 e57.
1426 10.1016/j.immuni.2017.06.001.

76. Eiraku, M., and Sasai, Y. (2012). Self-formation of layered neural structures in
1428 three-dimensional culture of ES cells. Curr Opin Neurobiol *22*, 768-777.
1429 10.1016/j.conb.2012.02.005.

77. Lancaster, M.A., and Knoblich, J.A. (2014). Organogenesis in a dish: modeling
1431 development and disease using organoid technologies. Science *345*,
1432 1247125. 10.1126/science.1247125.

78. Fatehullah, A., Tan, S.H., and Barker, N. (2016). Organoids as an in vitro
1434 model of human development and disease. Nat Cell Biol *18*, 246-254.
1435 10.1038/ncb3312.

79. Boj, S.F., Hwang, C.I., Baker, L.A., Chio, II, Engle, D.D., Corbo, V., Jager, M.,
1437 Ponz-Sarvise, M., Tiriac, H., Spector, M.S., et al. (2015). Organoid models of

human and mouse ductal pancreatic cancer. Cell *160*, 324-338. 10.1016/j.cell.2014.12.021.

80. Broutier, L., Andersson-Rolf, A., Hindley, C.J., Boj, S.F., Clevers, H., Koo, B.K., and Huch, M. (2016). Culture and establishment of self-renewing human and mouse adult liver and pancreas 3D organoids and their genetic manipulation. Nat Protoc *11*, 1724-1743. 10.1038/nprot.2016.097.

81. Dossena, M., Piras, R., Cherubini, A., Barilani, M., Dugnani, E., Salanitro, F., Moreth, T., Pampaloni, F., Piemonti, L., and Lazzari, L. (2020). Standardized GMP-compliant scalable production of human pancreas organoids. Stem Cell Res Ther *11*, 94. 10.1186/s13287-020-1585-2.

82. Georgakopoulos, N., Prior, N., Angres, B., Mastrogiovanni, G., Cagan, A., Harrison, D., Hindley, C.J., Arnes-Benito, R., Liau, S.S., Curd, A., et al. (2020). Long-term expansion, genomic stability and in vivo safety of adult human pancreas organoids. BMC Dev Biol *20*, 4. 10.1186/s12861-020-0209-5.

83. Jung, N., Moreth, T., Stelzer, E.H.K., Pampaloni, F., and Windbergs, M. (2021). Non-invasive analysis of pancreas organoids in synthetic hydrogels defines material-cell interactions and luminal composition. Biomater Sci *9*, 5415-5426. 10.1039/d1bm00597a.

84. Mocarski, E.S., Guo, H., and Kaiser, W.J. (2015). Necroptosis: The Trojan horse in cell autonomous antiviral host defense. Virology *479-480*, 160-166. 10.1016/j.virol.2015.03.016.

85. Kaiser, W.J., Upton, J.W., and Mocarski, E.S. (2013). Viral modulation of programmed necrosis. Curr Opin Virol *3*, 296-306. 10.1016/j.coviro.2013.05.019.

86. Pearson, J.S., Giogha, C., Muhlen, S., Nachbur, U., Pham, C.L., Zhang, Y., Hildebrand, J.M., Oates, C.V., Lung, T.W., Ingle, D., et al. (2017). EspL is a

bacterial cysteine protease effector that cleaves RHIM proteins to block necroptosis and inflammation. Nat Microbiol *2*, 16258. 10.1038/nmicrobiol.2016.258.

87.   Upton, J.W., and Kaiser, W.J. (2017). DAI Another Way: Necroptotic Control of Viral Infection. Cell Host Microbe *21*, 290-293. 10.1016/j.chom.2017.01.016.

88.   Seifert, L., Werba, G., Tiwari, S., Giao Ly, N.N., Alothman, S., Alqunaibit, D., Avanzi, A., Barilla, R., Daley, D., Greco, S.H., et al. (2016). The necrosome promotes pancreatic oncogenesis via CXCL1 and Mincle-induced immune suppression. Nature *532*, 245-249. 10.1038/nature17403.

89.   Najafov, A., Chen, H., and Yuan, J. (2017). Necroptosis and Cancer. Trends Cancer *3*, 294-301. 10.1016/j.trecan.2017.03.002.

90.   Pefanis, A., Ierino, F.L., Murphy, J.M., and Cowan, P.J. (2019). Regulated necrosis in kidney ischemia-reperfusion injury. Kidney Int *96*, 291-301. 10.1016/j.kint.2019.02.009.

91.   Muller, T., Dewitz, C., Schmitz, J., Schroder, A.S., Brasen, J.H., Stockwell, B.R., Murphy, J.M., Kunzendorf, U., and Krautwald, S. (2017). Necroptosis and ferroptosis are alternative cell death pathways that operate in acute kidney failure. Cell Mol Life Sci *74*, 3631-3645. 10.1007/s00018-017-2547-4.

92.   Petrie, E.J., Sandow, J.J., Jacobsen, A.V., Smith, B.J., Griffin, M.D.W., Lucet, I.S., Dai, W., Young, S.N., Tanzer, M.C., Wardak, A., et al. (2018). Conformational switching of the pseudokinase domain promotes human MLKL tetramerization and cell death by necroptosis. Nat Commun *9*, 2422. 10.1038/s41467-018-04714-7.

93.   Davies, K.A., Tanzer, M.C., Griffin, M.D.W., Mok, Y.F., Young, S.N., Qin, R., Petrie, E.J., Czabotar, P.E., Silke, J., and Murphy, J.M. (2018). The brace helices of MLKL mediate interdomain communication and oligomerisation to

regulate cell death by necroptosis. Cell Death Differ *25*, 1567-1580.

10.1038/s41418-018-0061-3.

94. Tanzer, M.C., Matti, I., Hildebrand, J.M., Young, S.N., Wardak, A.,

Tripaydonis, A., Petrie, E.J., Mildenhall, A.L., Vaux, D.L., Vince, J.E., et al.

(2016). Evolutionary divergence of the necroptosis effector MLKL. Cell Death

Differ *23*, 1185-1197. 10.1038/cdd.2015.169.

95. Petrie, E.J., Czabotar, P.E., and Murphy, J.M. (2019). The Structural Basis of

Necroptotic Cell Death Signaling. Trends Biochem Sci *44*, 53-63.

10.1016/j.tibs.2018.11.002.

96. Tanzer, M.C., Tripaydonis, A., Webb, A.I., Young, S.N., Varghese, L.N., Hall,

C., Alexander, W.S., Hildebrand, J.M., Silke, J., and Murphy, J.M. (2015).

Necroptosis signalling is tuned by phosphorylation of MLKL residues outside

the pseudokinase domain activation loop. Biochem J *471*, 255-265.

10.1042/BJ20150678.

97. Rodriguez, D.A., Weinlich, R., Brown, S., Guy, C., Fitzgerald, P., Dillon, C.P.,

Oberst, A., Quarato, G., Low, J., Cripps, J.G., et al. (2016). Characterization of

RIPK3-mediated phosphorylation of the activation loop of MLKL during

necroptosis. Cell Death Differ *23*, 76-88. 10.1038/cdd.2015.70.

98. Dai, J., Zhang, C., Guo, L., He, H., Jiang, K., Huang, Y., Zhang, X., Zhang, H.,

Wei, W., Zhang, Y., et al. (2020). A necroptotic-independent function of MLKL

in regulating endothelial cell adhesion molecule expression. Cell Death Dis *11*,

282. 10.1038/s41419-020-2483-3.

99. Petrie, E.J., Birkinshaw, R.W., Koide, A., Denbaum, E., Hildebrand, J.M.,

Garnish, S.E., Davies, K.A., Sandow, J.J., Samson, A.L., Gavin, X., et al.

(2020). Identification of MLKL membrane translocation as a checkpoint in

necroptotic cell death using Monobodies. Proc Natl Acad Sci U S A *117*, 8468-8475. 10.1073/pnas.1919960117.

100. Douanne, T., Andre-Gregoire, G., Trillet, K., Thys, A., Papin, A., Feyeux, M., Hulin, P., Chiron, D., Gavard, J., and Bidere, N. (2019). Pannexin-1 limits the production of proinflammatory cytokines during necroptosis. EMBO Rep *20*, e47840. 10.15252/embr.201947840.

101. Mizuno, E., Kawahata, K., Kato, M., Kitamura, N., and Komada, M. (2003). STAM proteins bind ubiquitinated proteins on the early endosome via the VHS domain and ubiquitin-interacting motif. Mol Biol Cell *14*, 3675-3689. 10.1091/mbc.e02-12-0823.

102. Lohi, O., and Lehto, V.P. (1998). VHS domain marks a group of proteins involved in endocytosis and vesicular trafficking. FEBS Lett *440*, 255-257. 10.1016/s0014-5793(98)01401-x.

103. Hirano, S., Kawasaki, M., Ura, H., Kato, R., Raiborg, C., Stenmark, H., and Wakatsuki, S. (2006). Double-sided ubiquitin binding of Hrs-UIM in endosomal protein sorting. Nat Struct Mol Biol *13*, 272-277. 10.1038/nsmb1051.

104. Bache, K.G., Raiborg, C., Mehlum, A., and Stenmark, H. (2003). STAM and Hrs are subunits of a multivalent ubiquitin-binding complex on early endosomes. J Biol Chem *278*, 12513-12521. 10.1074/jbc.M210843200.

105. Pornillos, O., Alam, S.L., Rich, R.L., Myszka, D.G., Davis, D.R., and Sundquist, W.I. (2002). Structure and functional interactions of the Tsg101 UEV domain. EMBO J *21*, 2397-2406. 10.1093/emboj/21.10.2397.

106. Sundquist, W.I., Schubert, H.L., Kelly, B.N., Hill, G.C., Holton, J.M., and Hill, C.P. (2004). Ubiquitin recognition by the human TSG101 protein. Mol Cell *13*, 783-789. 10.1016/s1097-2765(04)00129-7.

107.   Teo, H., Veprintsev, D.B., and Williams, R.L. (2004). Structural insights into
endosomal sorting complex required for transport (ESCRT-I) recognition of
ubiquitinated   proteins.   J   Biol   Chem   *279*,   28689-28696.
10.1074/jbc.M400023200.

108.   Meister, M., Banfer, S., Gartner, U., Koskimies, J., Amaddii, M., Jacob, R., and
Tikkanen, R. (2017). Regulation of cargo transfer between ESCRT-0 and
ESCRT-I complexes by flotillin-1 during endosomal sorting of ubiquitinated
cargo. Oncogenesis *6*, e344. 10.1038/oncsis.2017.47.

109.   Legler, D.F., Micheau, O., Doucey, M.A., Tschopp, J., and Bron, C. (2003).
Recruitment of TNF receptor 1 to lipid rafts is essential for TNFalpha-mediated
NF-kappaB activation. Immunity *18*, 655-664. 10.1016/s1074-7613(03)00092-
x.

110.   Hunter, I., and Nixon, G.F. (2006). Spatial compartmentalization of tumor
necrosis factor (TNF) receptor 1-dependent signaling pathways in human
airway smooth muscle cells. Lipid rafts are essential for TNF-alpha-mediated
activation of RhoA but dispensable for the activation of the NF-kappaB and
MAPK pathways. J Biol Chem *281*, 34705-34715. 10.1074/jbc.M605738200.

111.   Maminska, A., Bartosik, A., Banach-Orlowska, M., Pilecka, I., Jastrzebski, K.,
Zdzalik-Bielecka, D., Castanon, I., Poulain, M., Neyen, C., Wolinska-Niziol, L.,
et al. (2016). ESCRT proteins restrict constitutive NF-kappaB signaling by
trafficking cytokine receptors. Sci Signal *9*, ra8. 10.1126/scisignal.aad0848.

112.   Mahul-Mellier, A.L., Strappazzon, F., Petiot, A., Chatellard-Causse, C., Torch,
S., Blot, B., Freeman, K., Kuhn, L., Garin, J., Verna, J.M., et al. (2008). Alix
and ALG-2 are involved in tumor necrosis factor receptor 1-induced cell death.
J Biol Chem *283*, 34954-34965. 10.1074/jbc.M803140200.

113. Liu, Z., Dagley, L.F., Shield-Artin, K., Young, S.N., Bankovacki, A., Wang, X.,

1566   Tang, M., Howitt, J., Stafford, C.A., Nachbur, U., et al. (2021). Oligomerization-

1567   driven MLKL ubiquitylation antagonises necroptosis. bioRxiv,

1568   2021.2005.2001.442209. 10.1101/2021.05.01.442209.

1569 114. Garcia, L.R., Tenev, T., Newman, R., Haich, R.O., Liccardi, G., John, S.W.,

1570   Annibaldi, A., Yu, L., Pardo, M., Young, S.N., et al. (2021). Ubiquitylation of

1571   MLKL at lysine 219 positively regulates necroptosis-induced tissue injury and

1572   pathogen clearance. Nat Commun *12*, 3364. 10.1038/s41467-021-23474-5.

1573 115. Akimov, V., Barrio-Hernandez, I., Hansen, S.V.F., Hallenborg, P., Pedersen,

1574   A.K., Bekker-Jensen, D.B., Puglia, M., Christensen, S.D.K., Vanselow, J.T.,

1575   Nielsen, M.M., et al. (2018). UbiSite approach for comprehensive mapping of

1576   lysine and N-terminal ubiquitination sites. Nat Struct Mol Biol *25*, 631-640.

1577   10.1038/s41594-018-0084-y.

1578 116. Roedig, J., Kowald, L., Juretschke, T., Karlowitz, R., Ahangarian Abhari, B.,

1579   Roedig, H., Fulda, S., Beli, P., and van Wijk, S.J. (2021). USP22 controls

1580   necroptosis by regulating receptor-interacting protein kinase 3 ubiquitination.

1581   EMBO Rep *22*, e50163. 10.15252/embr.202050163.

1582 117. Muller, S., Rycak, L., Afonso-Grunz, F., Winter, P., Zawada, A.M., Damrath,

1583   E., Scheider, J., Schmah, J., Koch, I., Kahl, G., and Rotter, B. (2014). APADB:

1584   a database for alternative polyadenylation and microRNA regulation events.

1585   Database (Oxford) *2014*. 10.1093/database/bau076.

1586 118. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

1587   sequencing reads. 2011 *17*, 3. 10.14806/ej.17.1.200.

1588 119. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold

1589   change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

1590   10.1186/s13059-014-0550-8.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6