Explaining the primate extinction crisis: predictors of extinction risk and active threats

Maria J.A. Creighton[a] & Charles L. Nunn[bc]

[a]Department of Biology, Duke University, Durham, North Carolina, USA

[b]Department of Evolutionary Anthropology, Duke University, Durham, North Carolina, USA

[c]Global Health Institute, Duke University, Durham, North Carolina, USA

*Author for correspondence: maria.creighton@duke.edu; Department of Biology, Duke University, 130 Science Drive, Durham, NC 27708, USA; ORCID: https://orcid.org/0000-0003-1337-1946

ABSTRACT:

Explaining why some species are disproportionately impacted by the extinction crisis is of critical importance for conservation biology as a science and for proactively protecting species that are likely to become threatened in the future. Using the most current data on threat status, population trends, and threat types for 446 primate species, we advance previous research on the determinants of extinction risk by including a wider array of phenotypic traits as predictors, filling gaps in these trait data using multiple imputation, and considering more explicitly the mechanisms that connect organismal traits to extinction risk. Our Bayesian phylogenetically controlled analyses reveal that larger-bodied and insular species exhibit higher threat status, while those that are more omnivorous and live in larger groups have lower threat status. The same traits are not linked to risk when repeating our analyses with older IUCN data, suggesting that the traits that influence species risk are changing as anthropogenic effects continue to transform natural landscapes. We also show that larger-bodied and arboreal species are more susceptible to key threats responsible for primate population declines. Collectively, these results provide new insights to the determinants of primate extinction and identify the mechanisms (i.e., threats) that link traits to risk.

INTRODUCTION:

Anthropogenic activity is causing species to disappear at an alarming rate. However, not all species are affected equally: some species show no changes in abundance while others are increasing in abundance as human activity increases. Explaining why some species are more susceptible to extinction than others has become a major goal of conservation biologists as these contributions help to both explain current extinction patterns and allow for proactive protection of species possessing traits that could increase their probability of becoming imperiled. Previous studies have shown that phenotypic traits affect a species' susceptibility to extinction (Chichorro et al., 2019). Physical traits such as large body size and life history traits such as long generation lengths have been associated with increased risk of extinction (Purvis et al., 2000; Cardillo & Bromham, 2001; Cardillo et al., 2005; Lee & Jetz, 2011; Matthews et al., 2011). Behavioral traits have also been linked to increased extinction risk, including small group size and reduced innovativeness (Davidson et al., 2009; 2012; Ducatez et al., 2020).

While much effort has been put toward identifying how different traits covary with extinction risk, important knowledge gaps have limited the effectiveness of these analyses. First,

only a handful of studies have incorporated a broad range of traits in a single analysis. Chichorro
et al. (2019) reviewed studies investigating the correlates of extinction risk and found significant
variability in the traits that were investigated (or controlled for). In addition, some traits have only
recently been linked to extinction risk, such as behavioral flexibility (Ducatez et al., 2020), and
thus have not been widely investigated across clades.

Second, the relationship between the actual anthropogenic drivers of environmental change
that are responsible for extinction and species traits are understudied in many clades (e.g., in
primates; Estrada et al., 2017), limiting the impact of these comparative studies in applied
conservation (Cardillo & Meijaard, 2012). Identifying which threats are most impactful to species
with different trait types would enable actionable conservation steps. This could include mitigating
key threats in susceptible species' ranges and identifying which species are most likely to be
vulnerable to certain types of environmental change in the future. Despite this possible benefit of
considering specific threats, previous studies have been mostly limited to linking traits and species'
threat status, although a few exceptions have incorporated consideration of threats into these
analyses (e.g., Purvis et al., 2005; González-Suárez et al., 2013; Murray et al., 2014).

Lastly, we lack information on relevant traits for many species, resulting in incomplete
data. The species for which we lack data may be systematically biased towards those that are more
difficult to study, such as arboreal or nocturnal species. In addition to reducing statistical power,
removing these species from analyses has potential to bias observed relationships between
variables (Nakagawa & Freckleton, 2008) and can result in a loss of real information when some
traits included in an analysis have better data coverage than others.

Primates have been especially important in studies assessing predictors of extinction risk
(Purvis et al., 2000; Purvis et al., 2005; Matthews et al., 2011; Machado et al., 2022). Primates are
one of the most threatened animal clades, with ~65% of species at risk of extinction (IUCN, 2021).
The number of threatened primate species is on the rise (up ~5% in approximately 5 years), yet
the last comprehensive assessment of the major determinants of primate extinction risk was
published over 20 years ago (Purvis et al., 2000). The number of recognized primate species has
changed dramatically since earlier studies, having more than doubled from 180 to over 500 in the
past few decades (Rylands & Mittermeier, 2014; Creighton et al. 2022). As a result of these
taxonomic changes and limitations of older phylogenies, older studies focused on a relatively small
number of currently recognized primate species. More speciose and up-to-date phylogenies have

recently become available (Upham et al., 2019) in addition to a greater quantity and quality of trait

data for many species. These contributions create an opportunity for the inclusion of more

described primate species in comparative analyses, bringing us closer to capturing the true scope

of primate diversity. Primates are crucial components of tropical biodiversity, core players in the

function of ecosystems, and central to many cultures and religions (Estrada et al., 2017). It is thus

an urgent goal to determine which biological and behavioral traits contribute to primate extinction

vulnerability and how these traits interact with anthropogenic impacts to contribute to population

declines.

Here, we analyze the biological and behavioral determinants of primate extinction risk

using a phylogenetic comparative approach. We investigate the relationship between multiple

phenotypic traits and two measures of extinction risk reported by the International Union for

Conservation of Nature (IUCN): threat status and population trend. We then assess how these same

traits covary with vulnerability to the major threats facing primate species. This research addresses

the gaps above by including multiple traits in the analysis and using imputation approaches based

on phylogeny and phenotypic traits to fill in data for species with missing trait values. In addition,

by investigating population trends and specific threats, we improve understanding of the

connections between specific traits and the abundance of primates.

We focus on 10 key traits with proposed links to extinction risk (Table 1).

Table 1: The predicted direction of effect of biological and behavioral traits on extinction risk.

Trait:	Expected risk high when:	Reason:
Body mass (g)	Large body	Animals with large bodies have slow life histories and have been previously shown to be at increased extinction risk (Cardillo & Bromham, 2001; Cardillo et al., 2005).
Generation length (yrs)	Long generations	Slow life histories mean fewer generations to adapt to environmental changes (Purvis et al., 2000).
Home range size (ha)	Large home range size	Species that have individuals that maintain large home ranges are particularly vulnerable to habitat

		loss, degradation, and edge effects (Woodroffe & Ginsberg, 1998; Purvis et al., 2000).
Group size	Small group size	Small group size has been associated with heightened extinction risk, perhaps because large groups reduce predation and enhance foraging (Davidson et al., 2009; 2012).
Brain volume (cm³)	Small brain volume	Large relative brain size is a proxy of general intelligence and behavioral flexibility (Reader et al., 2011; Navarrete et al., 2016), which have been associated with reduced extinction risk in birds (Ducatez et al., 2020).
Omnivory (true or false)	False	Animals with a large dietary breadth can rely on a wider range of food types when resources become limited (Boyles & Storm, 2007).
Social system	Polygynandry	Species characterized by complex social organization are hypothesized to have larger critical population sizes (i.e., more individuals must persist to maintain what is considered to be a healthy population) and may therefore go extinct more quickly than species in simpler social systems (Höglund, 1996).
Lifestyle	Arboreal	Strictly arboreal species are disproportionately affected from losing habitat via deforestation (Munstermann et al., 2022).
Insularity (true or false)	True	Island ecosystems are particularly vulnerable to anthropogenic change due to small population sizes, low habitat availability, and low functional redundancy (Biber, 2002; Blackburn et al., 2004; Leclerc et al., 2018).
Nocturnal (true or false)	Diurnal	Diurnal species are more likely to be disturbed by human activity (e.g., traffic) and diurnal activity

has been connected to extinction risk (Purvis et al., 2000).

METHODS:

DATA:

We collected information on threat status (least concern = LC, near threatened = NT, vulnerable =

VU, endangered = EN, critically endangered = CR, data deficient = DD, and not evaluated = NE)

and population trend (increasing = I, stable = S, decreasing = D, and unknown = U), from the

IUCN (2021) for 446 primate species present in the ultrametric primate phylogeny published by

Upham et al. (2019) (Figure 1). We also collected a list of active threat types affecting each species

in the IUCN (2021) as defined by the Salafsky et al. (2008) threat classification system: 1 =

residential and commercial development, 2 = agriculture and aquaculture, 3 = energy production

and mining, 4 = transportation and service corridors, 5 = biological resource use, 6 = human

intrusions and disturbance, 7 = natural system modifications, 8 = invasive and other problematic

species and genes, 9 = pollution, 10 = geological events, and 11 = climate change and severe

weather.

Figure 1: Phylogenetic distribution of threat status and population trends (IUCN, 2021) for 446 primate species in the Upham et al. (2019) phylogeny. Images of representative species are presented next to family labels. Codes for threat status: data deficient = DD, not evaluated = NE, least concern = LC, near threatened = NT, vulnerable = VU, endangered = EN, and critically endangered = CR. Codes for population trend: unknown = U, increasing = I, stable = S, and decreasing = D (IUCN, 2021).

For each of the 446 species in our dataset, we recorded data on 10 different biological and behavioral traits that have been proposed to be associated with extinction risk from various sources: body mass (g) (Galán-Acedo et al., 2019), generation length (yrs) (IUCN, 2021), home range size (ha) (Galán-Acedo et al., 2019), group size (Rowe & Myers, 2011), brain volume (Powell et al., 2017), omnivory (true or false), social system (solitary, pair-living, harem polygyny and polygynandry) (DeCasien et al., 2017; Rowe & Myers, 2011 and other sources), lifestyle

(arboreal, terrestrial, or both) (Rowe & Myers, 2011 and other sources), insular (true or false)
(inferred from range data available in Rowe & Myers, 2011; IUCN, 2021), and nocturnal (true or
false) (Estrada et al 2017; IUCN, 2021). The full list of references for trait values is available in
the Supplementary Data. Table 1 summarizes how we expected each trait to be associated with
primate extinction risk. Previous studies have included geographic range size as a covariate in
similar analyses (e.g., Purvis et al., 2000; Machado et al., 2022). However, a species' geographic
range size is one of the main criteria used by the IUCN to assign threat status: species with small
population sizes that have small or restricted geographic ranges are considered to be more
imperiled (IUCN, 2021) (i.e., threatened species have small geographic ranges by definition).
Because we were interested in how biological and behavioral trait contribute to extinction risk,
including effects on what geographic ranges they are able to occupy, we did not include geographic
range size in our analysis. Notably, by including insularity in our analysis we controlled for the
fact that species on small islands may not be able to maintain geographic ranges large enough to
be considered healthy by the IUCN due to geographic barriers.
Following Powell et al. (2017), for sexually dimorphic clades (size difference > 10%) only
brain volume and body mass data from adult females were used in analysis. For all other species,
averages for all adults measured in the original source were used. Species found exclusively on
Madagascar, Borneo, or Sumatra were not scored as insular since these islands are large enough
to support large geographic ranges comparable to many mainland species. Further details on
operational definitions and trait coding are provided in the Supplementary Materials, along with a
correlation matrix of all traits and response variables (Figure S1), and a comparison of trait data
from different sources (Figures S2, S3, and S4).

ANALYSIS:

Trait imputation

The availability of data varied across the species in our dataset. Percentages of species with missing
trait data were: body mass (6%), omnivory (17%), generation length (23%), home range size
(25%), group size (39%), and brain volume (46%). Restricting the analysis to only species with
observed data on all traits reduced our sample size of species by over half (e.g., to from n=430 to
n=151 in our analysis of threat status). We thus opted to use multiple imputation to avoid losing
species from our analysis where one or more traits had missing observations. The main advantage

of imputation is that it avoids a loss of statistical power caused by reduced sample sizes due to
missing trait data and that it reduces potential biases in parameter estimates caused by eliminating
missing observations (see Nakagawa & Freckleton, 2008). For instance, if missing observations
are not missing completely at random, the relationship between two variables could change when
they are excluded from analysis (Nakagawa & Freckleton, 2008). In analyses with more than one
independent variable imputation also prevents the loss of real information if some variables have
better data coverage than others. For example, if an analysis has one independent variable with
100% data coverage and a second independent variable with only 80% data coverage, imputation
will allow all observations for the first trait to be kept in the analysis by substituting the 20% of
missing values for the second variable.

Multiple imputation was accomplished using phylogenetic information and data for other
traits. We started with phylogenetic multiple imputation using 100 randomly sampled trees
available from Upham et al. (2019). The imputation of traits was ordered so that imputed
information could be used to inform subsequent model fits along with phylogenetic information
(e.g., once body mass was imputed it was used to inform model fits for the imputation of other
variables). From each tree, we generated a variance-covariance matrix which we then dissolved
into 445 eigenvectors using the 'PVRdecomp' function from the R package PVR (Santos et al.,
2018). Using forward-backward model selection, we determined which phylogenetic eigenvectors
and traits had the best support for inclusion in models predicting each trait with missing data based
on Akaike information criterion (AICc) scores using the 'stepAIC' function from the MASS R
package (Ripley et al., 2013). We chose how many eigenvectors to include in model selection for
the imputation based on model performance in cross validation (Table S1). The top model for each
trait was used to impute values for each species missing data using the 'predict' function from the
car package in R (Fox et al., 2012). To propagate error, we then used the fits and standard
deviations associated with predicted values to take a randomly sampled trait value for each species
from the normal or binomial distribution (depending on whether traits were continuous or binary).

This imputation process was repeated once for each tree, resulting in 100 imputed datasets.
We performed a leave-one-out cross-validation of each imputation, where we removed observed
datapoints and used our imputation method to predict their value. When comparing these
predictions to the actual observed datapoints performance proved to be good in all cases (predictive
accuracy > 0.8 for continuous variables and area under the ROC curve > 0.8 for binary variables;

see Table S1).

Modelling threat status, population trends, and threat types

We ran multiple models to test predictors of three types of outcomes for primate populations. First,

we tested which biological and behavioral traits are associated with the threat status of a species.

Second, we tested the effects of species' behavioral and biological traits on population trends (i.e.,

population growth or decline). Third, we tested which traits were associated with species

susceptibility to the most important threats that primate species face.

To determine which biological and behavioral traits are associated with threat status, we

ran two phylogenetic generalized linear mixed models using Bayesian approximation as

implemented in the MCMCglmm R package (Hadfield, 2010). The first model had an ordinal error

structure, and the response variable was an ordinal measure of threat status scored as follows:

LC=0, NT=1, VU=2, EN=3, and CR=4 (Butchart et al., 2007). We ran the second model with a

threshold error structure; here the response variable was threat status scored as a binary outcome

where species were scored as either being threatened (VU, EN, or CR) (scored as 1) or not

threatened (NT or LC) (scored as 0).

To test the effects of traits on population trends we ran a third phylogenetic generalized

linear mixed model with a threshold error structure (Hadfield, 2010). In this model, each species

was assigned a binary outcome of either declining (scored as 1) or not declining (i.e., stable or

increasing) (scored as 0).

Finally, to determine which biological and behavioral traits were associated with species'

susceptibility to the most important threats that primate species face, we ran five phylogenetic

generalized linear mixed models with threshold error structures (Hadfield, 2010), one for each of

the top five threats to primates identified by the IUCN (2021). These top five threats identified for

primate species were: 1 = residential and commercial development (35% of species), 2 =

agriculture and aquaculture (80%), 3 = energy production and mining (22%), 5 = biological

resource use (82%), and 7 = natural system modifications (23%). Here, each of our five models

had a binary outcome of 1 (indicating that a species was affected by a particular threat) or 0

(indicating that a species was not affected by the threat).

Each model described above was run on 100 imputed datasets and phylogenies to account

for uncertainty in phylogeny and trait estimates (Nakagawa & De Villemereuil, 2019). Models ran

a total of 550,001 iterations, with a thinning interval of 500 and a burn-in of 50,000 to ensure
convergence had occurred. We confirmed convergence had occurred by assessing trace plots after
analyses finished running (Hadfield, 2010). Species were dropped from analyses when the true
value of a response variable was unknown by the IUCN (2021) (e.g., if the threat status was DD
or NE). Traits were included as fixed effects and continuous variables were ln-transformed,
centered with respect to the mean, and scaled by 2 standard deviations in all models to make their
effect sizes comparable to those reported for binary variables (Gelman, 2008). We used a weakly
informative gelman prior for fixed effects.

We also tested the hypothesis that some traits previously shown to be associated with
primate extinction risk are losing signal as more species become imperiled, for example, if
anthropogenic threats are becoming so overwhelming that all species are beginning to suffer
regardless of their attributes. This analysis involved repeating our analyses of threat status using
an older IUCN threat status data and species list (obtained from Harcourt & Parks, 2003).

To interpret the output from our Bayesian analyses, we provide (i) the distribution of
posterior means for tests from all 100 imputed datasets in graphical form (Figures 2 and 3), (ii) the
89% credible intervals (per McElreath, 2018) from the full posterior distribution of estimates in
graphical form (Figures 2 and 3), and (iii) the percentage of iterations from each set of 100 models
that were consistently positive or negative (Tables S2 to S13). We focused on results that were
most supported based on these outcomes. For the purposes of providing an estimate of the
magnitude of an effect in the main text, posterior means were pooled across datasets using Rubin's
rules (Nakagawa & De Villemereuil, 2019) (hereafter, "pooled posterior mean").

RESULTS:
Predictors of threat status and population trends
When scored as an ordinal outcome (LC=0, NT=1, VU=2, EN=3, and CR=4), threat status was
positively associated with insularity (pooled posterior mean = 1.217; 100% of 100,100 posterior
estimates > 0) and, albeit not as strongly, with body mass (pooled posterior mean = 0.995; 94%
estimates > 0) (Figure 2; Tables S2) across 430 species with known threat statuses. Ordinal threat
status was negatively associated with omnivory (pooled posterior mean = -0.475; 95% estimates
< 0) and group size (pooled posterior mean = -0.601; 95% estimates < 0) (Figure 2; Table S2). In
our analysis using older IUCN data, we found that only insularity (pooled posterior mean = 2.255;

100% estimates > 0) and home range size (pooled posterior mean = 1.532; 100% estimates > 0)

were associated with threat status (ordinal) in the predicted direction (Table S3).

Threat status was not strongly associated with any biological or behavioral traits when

scored as a binary response (Figure 2; Table S4). We repeated this analysis removing imputed

datapoints to see if their inclusion affected the lack of strong associations. We also ran three

separate models with body mass, generation length, and brain volume as sole predictors to

determine if correlations among these predictors (see Figure S1) in the full models affected the

results. Results from these additional analyses again consistently showed no strong effect of any

traits (Tables S5 and S6). In our analysis using older IUCN data, we found that insularity (pooled

posterior mean = 2.565; 100% estimates > 0) and home range size (pooled posterior mean = 1.116;

99% estimates > 0) were associated with threat status (binary) in the predicted direction (Table

S7). Population trend was not consistently associated with any biological or behavioral traits across

401 species with known population trends (Figure 2; Table S8).

Figure 2: Outcomes from three sets of models testing the relationship between traits and: threat

status scored as an ordinal variable (first panel), threat status scored as a binary variable (second

panel), and population trend scored as a binary variable (third panel). Each cell contains 100

posterior means (plotted as translucent diamonds) with their associated 89% credible intervals

(plotted as translucent horizontal lines) obtained from 100 MCMCglmm models run with 100

randomly sampled phylogenies across 100 trait datasets with missing datapoints obtained through

multiple imputation. Darker shading behind effects interpreted in text. Continuous variables were

ln-transformed, centered with respect to the mean, and scaled by 2 standard deviations.

Predictors of specific threat types

Our analyses of threat types included 404 species with known threats (Tables S9 to S13). Species

with larger body masses were more likely to be affected by threat 1 = residential and commercial

development (pooled posterior mean = 1.001; 96% estimates > 0; Table S9) and threat 3 = energy

production and mining (pooled posterior mean = 1.267; 99% estimates > 0; Table S11) (Figure 3).

Insularity was negatively associated with threat 1 = residential and commercial development

(pooled posterior mean = -0.755; 98% estimates < 0; Table S9), threat 3 = energy production and

mining (pooled posterior mean = -1.811; 100% estimates < 0; Table S11), and threat 7 = natural

system modifications (pooled posterior mean = -1.312; 100% estimates < 0; Table S13) (Figure

3). Species living a strictly arboreal lifestyle were more likely to be affected by threat 1 =

residential and commercial development than strictly terrestrial species (i.e., the baseline) (pooled

posterior mean = 1.023; 97% estimates > 0; Table S9) (Figure 3).

Figure 3: Outcomes from five sets of models testing the relationship between traits and species
susceptibility to: threat 1 = residential and commercial development (first panel), threat 2 =
agriculture and aquaculture (second panel), threat 3 = energy production and mining (third panel),
threat 5 = biological resource use (fourth panel), and threat 7 = natural system modifications (fifth
panel). Each cell contains 100 posterior means (plotted as translucent diamonds) with their
associated 89% credible intervals (plotted as translucent horizontal lines) obtained from 100
MCMCglmm models run with 100 randomly sampled phylogenies across 100 trait datasets with
missing datapoints obtained through multiple imputation. Darker shading behind effects
interpreted in text. Continuous variables were ln-transformed, centered with respect to the mean,
and scaled by 2 standard deviations.

DISCUSSION:
We investigated the correlates of extinction risk and threat susceptibility in primates using
phylogenetic comparative methods to analyze the most complete and up-to-date set of trait data
and IUCN data. One novelty of our approach involved the use of phylogenetic and trait-based
imputation of missing data. Our analyses revealed that larger-bodied and more insular species
exhibit higher threat status, while those that are more omnivorous or live in larger groups have
lower threat status. When looking at specific threats, we found that larger-bodied and arboreal
species are more vulnerable to key threats, while insular species are less vulnerable to these threats.

291   Our analyses with binary outcomes included threat status scored as threatened versus non-
threatened and population trend scored as declining versus not declining. These analyses revealed
that most traits are not strong predictors of whether a given primate species is considered
threatened (contrary to findings in other taxonomic groups, such as birds; Lee & Jetz, 2011). When
we scored threat status ordinally – with five ranked categories from least concern to critically
endangered – higher threat status was associated with insularity, large body mass, omnivory, and
small group size, consistent with our predictions for these traits. Therefore, primate species that
are most imperiled, and thus score highest in ordinal threat status, do tend to be those with
biological and behavioral predispositions to extinction.

300   We also considered how the effects of predictors changed over time. When applying the
same methods to analyze a 1999 IUCN dataset, we found that insularity and home range size
shared a positive relationship with both binary and ordinal threat status, but other traits were not

meaningfully associated with threat status. This pattern of results using newer versus older data
indicates that some traits (i.e., home range size) are becoming less important to extinction risk as
the effects of anthropogenic activity become increasingly overwhelming. The fact that traits
identified in our analysis of ordinal 2021 threat status (i.e., body mass, group size, and omnivory)
do not come out as strongly in our analysis with older data could indicate that these traits are
beginning to have a larger signal over time, or that the larger number of species in our 2021 dataset
(a consequence of taxonomic reevaluations in many clades; Creighton et al., 2022) enabled us to
more powerfully detect effects of these traits.

Previous studies have found an effect of variables not shown to explain meaningful
variation in our analysis. For example, Ducatez et al. (2020) investigated whether threat status
covaries with innovativeness, which is a known measure of behavioral flexibility associated with
general intelligence (Reader & Laland, 2002; Reader et al., 2011). We did not find an effect of
brain size, another known measure of behavioral flexibility and general intelligence (Reader et al.,
2011; Creighton et al. 2021), and correlate of innovativeness, in our analysis. The caveats
associated with each of these measures is discussed in Creighton et al. (2021), including the
difficultly of adequately controlling for literature effort when estimating innovativeness and the
small sample sizes used to estimate average brain sizes for many species. However, our difference
in results could point to the fact that different traits contribute to heightened extinction risk for
these two clades, and that the relationship between some traits and extinction risk may be far more
complicated than originally expected. Species with higher behavioral flexibility and general
intelligence may be less at risk of extinction because they are better equipped to find novel
solutions that allow them to overcome environmental challenges (Beever et al., 2017; Ducatez et
al., 2020). However, the relationship between behavioral flexibility and extinction may be more
complex, particularly in a clade like primates where innovations are frequent and human conflict
is common. Certain behavioral innovations can help species to avoid human conflict and endure
habitat modifications, such as novel approaches for accessing foods or shifting to new food
resources (Beever et al., 2017). However, other innovative behaviors can increase human-wildlife
conflict. In many primate species, crop-raiding and garbage eating have become common practice
and indicate their flexibility to survive in changing environments (e.g., chimpanzees and baboons;
Maples et al., 1976; Hahn et al., 2003; Hockings et al., 2009). These behaviors bring animals in
direct conflict with humans and, in some cases, attract them to lower quality habitats. Future

contributions should further address the paradox of how flexibility both helps and hinders survival,

and what this means for species' persistence.

In analyses assessing predictors of direct threats, we found that strictly arboreal species

were more likely to be threatened with residential and commercial development – a major driver

of deforestation in many regions. We also found that species with large body masses were more

likely to be affected by residential and commercial development and energy production and

mining. Thus, the tendency of larger bodied species to be severely threatened is likely influenced

by development and energy/mining activities. Finally, insular species were less likely to be

vulnerable to multiple threats (residential and commercial development, energy production and

mining, and natural system modifications) despite being more likely to be highly threatened. This

indicates that the high threat statuses of insular species may not be driven by anthropogenic

activity. Instead, their small geographic ranges enforced by geographic barriers could simply make

it impossible to maintain healthy population sizes, despite not being subject to major threats.

There are some limitations to our analyses. One issue involves correlated predictor

variables. Body mass, generation length, and brain volume were highly correlated in our dataset

(correlation coefficients range between 0.6 and 0.9). We included these predictors in the same

models to identify how they independently contributed to extinction risk and threat vulnerability

(Freckleton, 2002). However, this creates the possibility of collinearity in model estimates for

these variables. Increased uncertainty (i.e., wide credible intervals) around estimates for these three

traits indeed point to some collinearity among estimates. We therefore investigated the effects of

these variables individually in separate models, which yielded largely consistent results. Secondly,

like most previous studies, response variables in our analysis come from IUCN assessments. While

the IUCN maintains the largest global dataset on species extinction risk and threats useful for

comparative analyses, these measures are vulnerable to errors in empirical data and in models used

to estimate population declines and extinction risk (Rueda-Cediel et al., 2018). As a result, there

is likely to be uncaptured uncertainty associated with the measures of extinction risk used in our

analyses.

Information about how biological and behavioral traits contribute to species' susceptibility

to direct threats has been a persistent knowledge gap in the conservation literature, particularly for

primates (Estrada et al., 2017). We provide a first step in addressing this knowledge gap by linking

traits to key threats responsible for population declines. This approach has potential to be expanded

to other clades. Understanding the biological and behavioral predictors of threat susceptibility in

broader range of taxa could help groups like the IUCN to identify which threats pose the most

imminent harm to species with shared characteristics. There is also an opportunity to apply the

knowledge contributed here to projections about which species are next to become threatened: we

propose that factoring in traits and threat information could provide more powerful inference and

will lead to more proactive monitoring of species that face the greatest risks in the future. For

instance, strictly arboreal species living in regions with residential development may be expected

to decline at a faster rate than other species without these characteristics.

In summary, by taking advantage of statistical approaches for dealing with missing data

and Bayesian inference, we have shown that multiple traits contribute to primate threat status. Our

findings suggest that the effects of some traits, such as home range size, have weakened over the

past 20 years, indicating that the traits that influence a species' threat status are changing as

anthropogenic effects continue to transform natural landscapes. Other characteristics shown to

affect extinction risk in other clades, such as behavioral flexibility, do not appear to affect primate

extinction risk, suggesting that different processes likely govern extinction in different clades.

Focusing on mitigating key threats, as identified here, from susceptible species' geographic ranges

will be an important and necessary step for future recovery.

ACKNOWLEDGEMENTS: This work was funded by Duke University and The Triangle Center

for Evolutionary Medicine (TriCEM). We thank Dan Greenberg, Mark Janko, and Jarrod Hadfield

for crucial statistical advice. We also thank Susan Alberts, members of the Alberts Lab at Duke

University, and Brian Lerch for their comments on previous drafts of the manuscript.

Data and code for this manuscript will be made available on publication through Zenodo.

REFERENCES:

Beever, E. A., Hall, L. E., Varner, J., Loosen, A. E., Dunham, J. B., Gahl, M. K., ... & Lawler, J.

J. (2017). Behavioral flexibility as a mechanism for coping with climate change. *Frontiers*

in Ecology and the Environment, 15, 299-308.

Biber, E. (2002). Patterns of endemic extinctions among island bird species. *Ecography*, 25, 661-

676.

Blackburn, T. M., Cassey, P., Duncan, R. P., Evans, K. L., & Gaston, K. J. (2004). Avian extinction
and mammalian introductions on oceanic islands. *Science*, 305, 1955-1958.

Boyles, J. G., & Storm, J. J. (2007). The perils of picky eating: dietary breadth is related to
extinction risk in insectivorous bats. *PLoS One*, 2, e672.

Butchart, S. H., Resit Akçakaya, H., Chanson, J., Baillie, J. E., Collen, B., Quader, S., ... & Hilton-
Taylor, C. (2007). Improvements to the red list index. *PloS One*, 2, e140.

Cardillo, M., & Bromham, L. (2001). Body size and risk of extinction in Australian mammals.
Conservation Biology, 15, 1435-1440.

Cardillo, M., Mace, G. M., Jones, K. E., Bielby, J., Bininda-Emonds, O. R., Sechrest, W., ... &
Purvis, A. (2005). Multiple causes of high extinction risk in large mammal species.
Science, 309, 1239-1241.

Cardillo, M., & Meijaard, E. (2012). Are comparative studies of extinction risk useful for
conservation? *Trends in Ecology & Evolution*, 27, 167-171.

Chichorro, F., Juslén, A., & Cardoso, P. (2019). A review of the relation between species traits
and extinction risk. *Biological Conservation*, 237, 220-229.

Creighton, M. J., Greenberg, D. A., Reader, S. M., & Mooers, A. Ø. (2021). The role of
behavioural flexibility in primate diversification. *Animal Behaviour*, 180, 269-290.

Creighton, M. J., Luo, A. Q., Reader, S. M., & Mooers, A. Ø. (2022). Predictors of taxonomic
inflation and its role in primate conservation. *Animal Conservation*.

Davidson, A. D., Boyer, A. G., Kim, H., Pompa-Mansilla, S., Hamilton, M. J., Costa, D. P., ... &
Brown, J. H. (2012). Drivers and hotspots of extinction risk in marine mammals.
Proceedings of the National Academy of Sciences, 109, 3395-3400.

Davidson, A. D., Hamilton, M. J., Boyer, A. G., Brown, J. H., & Ceballos, G. (2009). Multiple
ecological pathways to extinction in mammals. *Proceedings of the National Academy of*
Sciences, 106, 10702-10705.

DeCasien, A. R., Williams, S. A., & Higham, J. P. (2017). Primate brain size is predicted by diet
but not sociality. *Nature Ecology & Evolution*, 1, 1-7.

Ducatez, S., Sol, D., Sayol, F., & Lefebvre, L. (2020). Behavioural plasticity is associated with
reduced extinction risk in birds. *Nature Ecology & Evolution*, 4, 788-793.

Estrada, A., Garber, P. A., Rylands, A. B., Roos, C., Fernandez-Duque, E., Di Fiore, A., ... & Li,
B. (2017). Impending extinction crisis of the world's primates: Why primates matter.
Science Advances, 3, e1600946.

Fox, J., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., ... & Monette, G. (2012).
Package 'car'. Retrieved from: https://cran.r-project.org/web/packages/car/index.html.

Freckleton, R. P. (2002). On the misuse of residuals in ecology: regression of residuals vs. multiple
regression. *Journal of Animal Ecology*, 71, 542–545.

Galán-Acedo, C., Arroyo-Rodríguez, V., Andresen, E., & Arasa-Gisbert, R. (2019). Ecological
traits of the world's primates. *Scientific Data*, 6, 1-5.

Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics in*
Medicine, 27, 2865–2873.

González-Suárez, M., Gómez, A., & Revilla, E. (2013). Which intrinsic traits predict vulnerability
to extinction depends on the actual threatening processes. *Ecosphere*, 4, 1-16.

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. *Journal of Statistical Software*. 33, 1–22.

Hahn, N. E., Proulx, D., Muruthi, P. M., Alberts, S., & Altmann, J. (2003). Gastrointestinal
parasites in free-ranging Kenyan baboons (*Papio cynocephalus and P. anubis*).
International Journal of Primatology, 24, 271-279.

Harcourt, A. H., & Parks, S. A. (2003). Threatened primates experience high human densities:
adding an index of threat to the IUCN Red List criteria. *Biological Conservation*, 109, 137-
149.

Hockings, K. J., Anderson, J. R., & Matsuzawa, T. (2009). Use of wild and cultivated foods by
chimpanzees at Bossou, Republic of Guinea: feeding dynamics in a human-influenced
environment. *American Journal of Primatology*, 71, 636-646.

Höglund, J. (1996). Can mating systems affect local extinction risks? Two examples of lek-
breeding waders. *Oikos*, 77, 184-188.

IUCN. (2021). The IUCN Red List of Threatened Species Version 2021-1. IUCN Red List of
Threatened Species. Retrieved from: https://www.iucnredlist.org/en.

Leclerc, C., Courchamp, F., & Bellard, C. (2018). Insular threat associations within taxa
worldwide. *Scientific Reports*, 8, 1-8.

Lee, T. M., & Jetz, W. (2011). Unravelling the structure of species extinction risk for predictive conservation science. *Proceedings of the Royal Society B: Biological Sciences*, 278, 1329-1338.

Machado, F. F., Jardim, L., Dinnage, R., Brito, D., & Cardillo, M. (2022). Diet disparity and diversity predict extinction risk in primates. *Animal Conservation*.

Maples, W. R., Maples, M. K., Greenhood, W. F., & Walek, M. L. (1976). Adaptations of crop-raiding baboons in Kenya. *American Journal of Physical Anthropology*, 45, 309-315.

Matthews, L. J., Arnold, C., Machanda, Z., & Nunn, C. L. (2011). Primate extinction risk and historical patterns of speciation and extinction in relation to body mass. *Proceedings of the Royal Society B: Biological Sciences*, 278, 1256-1263.

McElreath, R. (2018). Statistical rethinking: A Bayesian course with examples in R and Stan. Chapman and Hall/CRC, New York.

Munstermann, M. J., Heim, N. A., McCauley, D. J., Payne, J. L., Upham, N. S., Wang, S. C., & Knope, M. L. (2022). A global ecological signal of extinction risk in terrestrial vertebrates. *Conservation Biology*, 36, e13852.

Murray, K. A., Verde Arregoitia, L. D., Davidson, A., Di Marco, M., & Di Fonzo, M. M. (2014). Threat to the point: improving the value of comparative extinction risk analysis for conservation action. *Global Change Biology*, 20, 483-494.

Nakagawa, S., & De Villemereuil, P. (2019). A general method for simultaneously accounting for phylogenetic and species sampling uncertainty via Rubin's rules in comparative analysis. *Systematic Biology*, 68, 632-641.

Nakagawa, S., & Freckleton, R. P. (2008). Missing inaction: the dangers of ignoring missing data. *Trends in Ecology & Evolution*, 23, 592-596.

Navarrete, A. F., Reader, S. M., Street, S. E., Whalen, A., & Laland, K. N. (2016). The coevolution of innovation and technical intelligence in primates. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150186.

Powell, L. E., Isler, K., & Barton, R. A. (2017). Re-evaluating the link between brain size and behavioural ecology in primates. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20171765.

Purvis, A., Cardillo, M., Grenyer, R., & Collen, B. (2005). Correlates of extinction risk: phylogeny,
biology, threat and scale. In Purvis, A., Gittleman, J.L., & Brooks, T. (Eds.), *Phylogeny*
and Conservation (pp. 295-316). Cambridge University Press, Cambridge.

Purvis, A., Gittleman, J. L., Cowlishaw, G., & Mace, G. M. (2000). Predicting extinction risk in
declining species. *Proceedings of the Royal Society B: Biological Sciences*, 267, 1947-
1952.

Reader, S. M., Hager, Y., & Laland, K. N. (2011). The evolution of primate general and cultural
intelligence. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366,
1017-1027.

Reader, S. M., & Laland, K. N. (2002). Social intelligence, innovation, and enhanced brain size in
primates. *Proceedings of the National Academy of Sciences*, 99, 4436-4441.

Ripley, B., Venables, B., Bates, D. M., Hornik, K., Gebhardt, A., Firth, D., & Ripley, M. B. (2013).
Package   'MASS'.   CRAN   Repository,   See   http://cran.r-
projectorg/web/packages/MASS/MASS.pdf.

Rowe, N., & Myers, M. (2011). *All the world's primates*. Primate Conservation, Inc. Retrieved
from: https://www.alltheworldsprimates.org.

Rueda-Cediel, P., Anderson, K. E., Regan, T. J., & Regan, H. M. (2018). Effects of uncertainty
and variability on population declines and IUCN Red List classifications. *Conservation*
Biology, 32, 916-925.

Rylands, A. B., & Mittermeier, R. A. (2014). Primate taxonomy: species and conservation.
Evolutionary Anthropology, 23, 8-10.

Salafsky, N., Salzer, D., Stattersfield, A. J., Hilton-Taylor, C. R. A. I. G., Neugarten, R., Butchart,
S. H., ... & Wilkie, D. (2008). A standard lexicon for biodiversity conservation: unified
classifications of threats and actions. *Conservation Biology*, 22, 897-911.

Santos, T., Diniz-Filho, J. A., e Luis, T. R., Bini, M., & Santos, M. T. (2018). Package 'PVR'.
Retrieved from: https://cran.r-project.org/web/packages/PVR/index.html.

Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). Inferring the mammal tree: Species-level sets
of phylogenies for questions in ecology, evolution, and conservation. *PLoS Biology*, 17, 1-
44.

Woodroffe, R., & Ginsberg, J. R. (1998). Edge effects and the extinction of populations inside
protected areas. *Science*, 280, 2126-2128.