SMRT-AgRenSeq-d in potato (Solanum tuberosum) identifies candidates for the nematode resistance Gpa5

Yuhan Wang[1*], Lynn H Brown[1*], Thomas M Adams[2*], Yuk Woon Cheung[1], Jie Li[3], Vanessa Young[4], Drummond T Todd[4], Miles R Armstrong[1], Konrad Neugebauer[5], Amanpreet Kaur[1,6], Brian Harrower[2], Stan Oome[7], Xiaodan Wang[3], Micha Bayer[2], Ingo Hein[1,2,3t]

[1] The University of Dundee, Division of Plant Sciences at the Hutton, Errol Road, Invergowrie, Dundee, DD2 5DA, UK

[2] The James Hutton Institute, Errol Road, Invergowrie, Dundee, DD2 5DA, UK

[3] China Agricultural University, Haidian District, Beijing, 100083, China

[4] James Hutton Limited, Errol Road, Invergowrie, Dundee, DD2 5DA, UK

[5] Biomathematics and Statistics Scotland, Errol Road, Invergowrie, Dundee, DD2 5DA, UK

[6] Teagasc, Crop Research Centre, Oak Park, Carlow R93 XE12, Ireland

[7] HZPC Research B.V., Edisonweg 5, 8501 XG Joure, the Netherlands

[*] These authors contributed equally to this study

[t] Corresponding author: Ingo.Hein@hutton.ac.uk

Running Title:

SMRT-AgRenSeq-d for identifying PCN resistance

Keywords:
Potato, Potato Cyst Nematodes, *Globodera pallida*, *Gpa5*, RenSeq, SMRT-AgRenSeq-d

Abstract:

Potato is the third most important food crop in the world. Diverse pathogens threaten sustainable crop production but can be controlled, in many cases, through the deployment of disease resistance genes belonging to the family of nucleotide-binding, leucine-rich-repeat (NLR) genes.

To identify functional NLRs in established varieties, we have successfully established SMRT-AgRenSeq in tetraploid potatoes and have further enhanced the methodology by including dRenSeq in an approach that we term SMRT-AgRenSeq-d. The inclusion of dRenSeq enables the filtering of candidates after the association analysis by establishing a presence/absence matrix across resistant and susceptible potatoes that is translated into an F1 score. Using a SMRT-RenSeq based sequence representation of the NLRome from the cultivar Innovator, SMRT-AgRenSeq-d analyses reliably identified the late blight resistance benchmark genes *R1, R2-like, R3a* and *R3b* in a panel of 117 varieties with variable phenotype penetrations. All benchmark genes were identified with an F1 score of 1 which indicates absolute linkage in the panel.

When applied to the elusive nematode disease resistance gene *Gpa5* that controls the Potato Cyst Nematode (PCN) species *Globodera pallida* (pathotypes Pa2/3), SMRT-AgRenSeq-d identified nine strong candidates. These map to the previously established position on potato chromosome 5 and are potential homologs of the late blight resistance gene *R1*.

Assuming that NLRs are involved in controlling many types of resistances, SMRT-AgRenSeq-d can readily be applied to diverse crops and pathogen systems. In potato, SMRT-AgRenSeq-d lends itself, for example, to further study the elusive PCN resistances *H1* or *H3* for which phenotypic data exist.

Introduction:

Potato is the third most important global food crop and is both nutritionally and economically valuable [1]. Potato is a staple of many diets around the world, and diseases can thus have a significant impact on crop production and food security.

62 Diverse and phylogenetically unrelated pathogens ranging from oomycetes, fungi,
63 bacteria, viruses, nematodes, and insects can cause significant crop losses in potato
64 [2].
65
The Potato Cyst Nematode (PCN) species *Globodera rostochiensis* and *G. pallida*
are economically important pathogens of potato and are present in most potato
growing regions of the world [3]. PCN is difficult to eradicate once established as
cysts can survive for over 20 years in the soil, rendering even longer crop rotations
insufficient for clearing land [4,5]. Thus, the impact of PCN goes beyond the
immediate yield losses, which are estimated to be around 9% on average in
susceptible potatoes, with the presence of PCN rendering land unsuitable for high-
health potato seed production, which significantly impacts on the crop supply chain
[4,6].

The only cloned disease resistance genes effective against PCN are *Gpa2* and
Gro1-4 which encode for canonical nucleotide-binding, leucine-rich-repeat (NLR)
genes [7,8]. In current potato cultivars, the most effective resistance gene used to
control *G. pallida* is *Gpa5*, which has been introduced from the wild Solanaceae
species *Solanum vernei* [9]. The resistance has been mapped to potato
chromosome 5 [9,10] and a haplotype specific PCR marker, HC, was developed
which corresponds in its position to an NLR-rich locus on chromosome 5 [11]. This
locus is also associated with the late blight resistance gene *R1,* amongst other
functional NLRs such as *Grp1* and *H2* [3,12]. Intriguingly, Rouppe van der Voort *et*
al. [9] and an independent study aimed at mapping an *S. vernei* source of resistance
against *G. pallida*, independently revealed a further QTL, *Gpa6*, on chromosome 9
that complements the major effect mapped to chromosome 5 [13,14].
87
To specifically study potato NLRs, RenSeq was developed as a target enrichment
tool following the annotation of NLRs in the reference genome DM1-3 516 R44 (DM)
[15–17]. In combination with PacBio-based sequencing of long genomic DNA
fragments, described as SMRT-RenSeq, the technology enables the representation
of entire plant NLRomes [18,19]. A diagnostic version of RenSeq, dRenSeq, has
been used to ascertain the presence/absence of known NLRs in wild species or
cultivars. Thus, dRenSeq aids the identification of novel resistances in germplasm

collections as well as the deployment and pyramiding of complementary resistances
in cultivars [20,21].

RenSeq has also been adapted for association genetics (AgRenSeq) which takes
advantage of independent recombination events that occur during meiosis.
AgRenSeq, which determines the potential of candidate NLRs being responsible for
resistance traits, uses *k*-mer based association and has initially been implemented in
Aegilops tauschii, a wild progenitor of the D subgenome of hexaploid bread wheat
[22]. AgRenSeq has also been performed with PacBio HiFi reads to identify
candidate resistance genes in Rye controlling *Puccinia recondita* f. sp. *secalis* [23]*.*
This approach was termed SMRT-AgRenSeq.
105

Here we apply AgRenSeq for the first time in potatoes to identify candidate NLRs
that are associated with the *Gpa5* resistance linked to the HC marker. To study
Gpa5, we selected the tetraploid potato cultivar Innovator as a reference, since this
cultivar is known to exhibit high levels of *G. pallida* resistance based on the presence
of *Gpa5* [3]. As previously shown, Innovator also contains four known functional
NLRs effective against *P. infestans*: *R1*, *R2-like*, *R3a,* and a variant of *R3b* [20].
These four NLRs reside in three clusters of different sizes and thus function as ideal
benchmark genes to test the accuracy of the NLR assembly and subsequent
association studies.
115

116 We further developed SMRT-AgRenSeq by including a dRenSeq analysis to validate
117 the association of candidates and, importantly, identify candidate NLRs that are
118 100% linked to the resistance phenotype. We refer to this combination as SMRT-
119 AgRenSeq-d and suggest that this technology is well suited to identify resistances
120 against diverse pathogens that can be controlled through functional NLRs [24].
121

Results and Discussion:

122
SMRT-AgRenSeq-d encompasses a representation of the NLRs from a plant with
the desired resistance, a panel of plants with contrasting phenotypes to ascertain the
k-mer based association, and dRenSeq-based assessment of candidates. The

benchmark genes *Rpi-R1, Rpi-R2-like*, *Rpi-R3a* and *Rpi-R3b* in Innovator have been

utilised to provide a quality check throughout the analysis.

De-novo assembly of Illumina-based RenSeq reads fails to represent

benchmark NLRs

To assess the accuracy of the NLR representation from Illumina sequences, we

used RenSeq reads generated from the potato cultivar Innovator [20]. The *de novo*

assembly of Illumina-based RenSeq reads yielded 9,756 contigs with an average

length of 1,060.4 bp of which 1,111 contigs contained signatures of NLRs. NLR-

Annotator [25] predicted 1,150 NLRs in this assembly (Supplementary Table 1). The

integrity of the four benchmark genes was assessed in the assembled contigs (Table

1). Only *Rpi-R1* was accurately and completely represented in the contig

scf7180000044808. In contrast, *Rpi-R2-like*, and the *Rpi-R3b* variant were partially

and incorrectly represented in contigs scf7180000045065 and scf7180000045838,

respectively. For *Rpi-R2-like*, more than 7% sequence diversity was observed

compared to the reference, and for *Rpi-R3b*, 38% of the gene was missing.

Representation of *Rpi-R3a* was split over two contigs, scf7180000044942 and

scf7180000045673.

Table 1 Representation of four benchmark genes in the potato cultivar Innovator based on Illumina and PacBio RenSeq read assemblies.

Reference Gene	Representative Sequence	Identity (%)	Mismatch	Gap	Reference gene start	Reference gene end	Gene Length	Coverage of Gene (%)	Coverage of HSP** (%)
Illumina reference contigs									
Rpi-R1	scf7180000044808	100	0	0	1	4103	4103	100	100
Rpi-R2-like	scf7180000045065	92.635	170	4	1	2539	2545	99	99
Rpi-R3a	scf7180000044942	99.881	3	0	1	2531	3850	66	66
Rpi-R3a	scf7180000045673	99.937	1	0	2254	3850	3850	41	41
Rpi-R3b$^{G1696/G3111}$	scf7180000045838	99.958	1	0	698	3073	3853	62	62
PacBio reference contigs									
Rpi-R1	tig00000086	100	0	0	1	4103	4103	100	100
Rpi-R2-like	tig00001243	100	0	0	1	2545	2545	100	100
Rpi-R3a	tig00000180	100	0	0	1	3850	3850	100	100
Rpi-R3b$^{G1696/G3111}$	tig00000301	99.974	1	0	1	3853	3853	100	100

**A High-scoring Segment Pair (HSP) is a local alignment with no gaps that achieves one of the highest alignment scores in a given search.

Considering that the average full-length NLR in potato is approximately 3 kb in length

[16], the partial representation of *Rpi-R3a* and *Rpi-R3b* is consistent with the

relatively small average contig size observed. Importantly, the 150 bp Illumina reads

not only affect the assembly length of NLRs, but also their accuracy, the lack of

which is pronounced in *Rpi-R2-like* and *Rpi-R3a/R3b.* These genes originate in the

two largest NLR clusters in the potato genome [16]. The processes leading to larger

homogeneous clusters containing highly similar NLRs are typically governed by

tandem duplications [26]. It is likely that the assembly of the short reads collapsed

NLR gene clusters and insufficiently distinguished between alleles and paralogs.

Evidence for this was provided by comparing the number of predicted NLRs in the

doubled monoploid potato reference clone DM and Innovator. DM contains 755

NLRs [15] whereas the Illumina-based assembly of NLRs from the tetraploid cultivar

Innovator predicted 1,150, which suggests only a modest increase of 53% of the

total number of NLRs.

PacBio HiFi-based assembly of RenSeq reads generates a high-quality NLRome representation

The combination of RenSeq with single-molecule real-time (SMRT) sequencing (SMRT-RenSeq) has overcome the limitations of Illumina-based assemblies in the cloning of NLRs in wild diploid potato species [19]. The assembly of HiFi RenSeq reads from Innovator yielded 4,885 contigs with an average length of 11,279 bp. Amongst those, 1,760 contigs contained a total of 2,418 predicted NLRs (Supplementary Table 1).

In this assembly, the benchmark genes *Rpi-R1, Rpi-R2-like* and *Rpi-R3a* were represented completely and accurately in a single contig each (Table 1)

Acknowledgements

The authors thank John Jones for providing critical feedback on the manuscript. Further we acknowledge the Research/Scientific Computing teams at The James Hutton Institute and NIAB for providing computational resources and technical support for the "UK's Crop Diversity Bioinformatics HPC" (BBSRC grant BB/S019669/1), use of which has contributed to the results reported within this paper.

This work was supported by the Rural & Environment Science & Analytical Services (RESAS) Division of the Scottish Government through project JHI-B1-1, the Biotechnology and Biological Sciences Research Council (BBSRC) through award BB/S015663/1 and the Royal Society through award NAF\R1\201061. YW was supported through the CSC scholarship program, China. LB was supported through the East of Scotland Bioscience Doctoral Training Partnership (EASTBIO DTP), funded by the BBSRC award BB/T00875X/1. AK was supported through a Research Leaders 2025 fellowship funded by European Union's Horizon 2020 research and innovation programme under Marie Sklodowska-Curie grant agreement no. 754380.

Contributions

YW, LB, TMA, MA, AK conducted the computational analyses, developed the association panel and the F1 score. JL, XW, BH provided material for this study and

contributed towards the writing of the manuscript and the independent validation. VY,

DT and YW conducted the PCN score and HC marker analysis. TMA and KN

conducted the phylogenetic analysis. SO provided genomic data and recombination

analysis for this study. MB and IH conceived this work and secured funding. All

authors have read and endorsed the manuscript.

Data availability statement

BioProjects:

ERP141787 – Illumina Reads / contigs

ERP141789 – HiFi Reads / contigs

Conflict of interests

The authors declare that they have no competing interests.. A comparison between

the previously identified variant of *Rpi-R3b* in Innovator, *Rpi-R3b*$^{G1696/G3111}$, with the

well supported PacBio contig tig00000301 revealed 100% coverage of this gene and

99.974% identity to *Rpi-R3b*$^{G1696/G3111}$ with only one mismatch at position C918. We

referred to this variant as *Rpi-R3b*$^{C918/G1696/G3111}$. Interestingly, Illumina RenSeq reads

from Innovator and other cultivars, except for Brodie, could not distinguish *Rpi-*

R3b$^{G1696/G3111}$ from *Rpi-R3b*$^{C918/G1696/G3111}$ (Supplementary Table 2). This suggests

that highly similar sequences obscure the additional sequence polymorphism, or that

some cultivars contain both variants.

These results independently confirm and quantify the advances that SMRT-RenSeq

offers for the highly accurate representation of the complex NLR gene family that

cannot be achieved through Illumina-based assemblies [18,23,27]. In addition, these

contigs, which are larger in size compared to the described Illumina-derived contigs,

can contain multiple NLRs and their flanking regions on single contigs which aids

functional analyses such as expression of candidates with their native regulatory

elements.

A phylogenetic tree was constructed from the SMRT-RenSeq derived NLRs using

the NB domains of predicted NLRs from Innovator. Further included in the tree is a

selection of reference NLRs and two outgroup genes with NB domains from Humans

(*Apaf-1*) and *C. elegans (Ced4)* (Figure 1; Supplementary Figure 1). The resulting

tree shows clear groupings of NLRs into numerous distinct clades and emphasises
the scale of duplications that have occurred in tetraploid potatoes. In addition to
highlighting the NLRs related to *Rpi-R1, Rpi-R2-like, Rpi-R3a* and *Rpi-R3b*, a clear
NRC (NLR Required for Cell death) clade [28] is apparent (Figure 1; Supplementary
Figure 1).
229

Development of an association panel

The cultivar Innovator was released in 1999 and has since been extensively used in
breeding programs and is attributed as a direct parent to over 25 named varieties
[29] (Supplementary Table 3). To study *Gpa5* and to take advantage of the wide
genetic reach of Innovator, we developed an association panel of 117 potato
accessions with contrasting resistance/susceptible phenotypes against *G. pallida*,
associated with the presence/absence of *Gpa5.* Resistance scores were either
determined by replicated canister tests or, if readily available publicly, extracted from
databases (Supplementary Table 2). For *Gpa5*, the panel contained 23 highly
resistant clones with a *G. pallida* resistance score of between 7 and 9 all of which
tested positive for the presence of the HC marker, and 94 clones with a score of 1–3
(Supplementary Table 2).

Further, we assessed the same panel for the sequence representation of the late
blight resistance genes *Rpi-R1, Rpi-R2-like, Rpi-R3a* and *Rpi-R3b* through
dRenSeq, as they have been identified in Innovator previously [20]. The sequence
representation took into consideration known family members with sequence
polymorphisms. For example, in Armstrong *et al*. [20] we showed that Pentland Dell
contains the *Rpi-R2* family member *Rpi-abpt*[T86] which has also been identified in the
cultivar Bionica and the breeding clone 2573(2). Informed by the natural diversity of
Rpi-R2-like, which in the cultivar Pentland Dell has 97.4% dRenSeq coverage [20],
we set a minimum cut-off value of more than 97% sequence representation for a
gene to be classified as 'present'. This was converted into a presence/absence
matrix for the four genes, which form the benchmarks for this study, across the
association panel (Supplementary Table 2).
254

The representation of the benchmark genes in the panel was 10 clones for *Rpi-R2-*
like, 34 for *Rpi-R1*, 44 for *Rpi-R3a* and 60 for *Rpi-R3b*. Interestingly, all clones that
contained *Rpi-R3a* also contained *Rpi-R3b,* but not *vice versa*.

SMRT-AgRenSeq-d development for *R3a* as a proof of concept

Since potato is an autotetraploid crop with complex tetrasomic inheritance patterns,
we assessed first the suitability of SMRT-AgRenSeq-d on the benchmark genes *Rpi-*
R1, Rpi-R2-like, Rpi-R3a and *Rpi-R3b* (Figure 2; Supplementary Figures 2 – 4).

The SMRT-AgRenSeq-d study for *Rpi-R3a* yielded a maximum positive association
score of 44 (Figure 2a). To systematically assess the resulting candidates and to be
as inclusive as possible in the analysis, we selected half of the maximum association
score as an initial cut-off for candidate selection. At this lower association score of
22, 46 contigs were identified that contain 73 candidate NLRs. The sequences of
these 73 predicted NLRs were analysed by dRenSeq to independently validate their
prevalence in the 44 clones that contain the reference sequence of *Rpi-R3a* and
their absence in the 73 potato clones identified as missing *Rpi-R3a* (Supplementary
Table 4). Based on this graphical genotyping depicting the presence/absence of
candidate genes in the respective pools, an F1 score [30] was determined. An F1
score of 1 indicates the presence of the candidate gene in all clones from the
positive panel and the consistent absence in all negative clones. Importantly, the
Rpi-R3a containing contig tig00000180 (Table 1) was amongst these highly
positively associated candidates (Supplementary Table 4).

A score of less than 1 therefore indicates false positives (e.g., candidates that are
not present in all clones from the positive set) and/or false negatives (e.g.,
candidates that are present in the negative set). This conceptually equates to
identifying recombination events in potato clones that have, for example, lost a
specific candidate. Thus, the F1 score, which is determined through the dRenSeq
analysis, complements the association score by confirming and further prioritising
candidates. Interestingly, candidates that yielded an F1 score of less than 0.7, some
of which had an association score of 37, displayed unsystematic presence/absence
patterns that are inconsistent with robust association.

The analysis of candidates with an F1 score of larger than 0.7 but smaller than 1

highlighted examples of potential recombination and false positives, including two

NLRs with an association score of 44 from contig tig00000196 in the cultivar

Cammeo that does not contain *Rpi-R3a* (Supplementary Table 4). Consequently,

their F1 scores were lower than 1 at 0.9887. *Rpi-R3b* was also present in the *Rpi-*

R3a candidate gene list, on contig tig00000301 (Table 1, Supplementary Table 4).

This is consistent with the finding that all *Rpi-R3a* containing clones also contained

Rpi-R3b (Supplementary Table 2). However, as other potato clones also contain *Rpi-*

R3b, yet do not contain *Rpi-R3a*, the F1 score for this contig was again less than 1.

Similarly, five candidates from contigs tig00000247, tig00000325 and tig00000301

were missing in the cultivar Brodie that contains *Rpi-R3a* (Supplementary Table 4)

and hence also had F1 scores below 1. Consistent with this analysis, we found that

the number of candidates is inversely related to the F1 score (Figure 2b).

Only 11 putative NLRs had an F1 score of 1 and, as mentioned above, include

contig tig00000180 that contains the full-length *Rpi-R3a* gene sequence alongside a

closely related gene. We surmise that these 11 genes are physically linked and likely

represent paralogous sequences, as alleles would have been genetically randomly

distributed owing to the tetrasomic inheritance patterns of potato. Further, when

mapped to the DM reference genome, nine out of the 11 candidates resided in the

predicted *R3* locus on chromosome 11 (Figure 2c). The remaining two genes had

their best BLASTn hit on DM chromosome 9 which contains sequences that belong

to the *R3* family as demonstrated by Jupe *et al.* [16]. Consistent with this, all 11

candidates reside in the *Rpi-R3a/R3b* clade of the phylogenetic tree (Figure 1).

SMRT-AgRenSeq-d validation for benchmark genes *Rpi-R3b, Rpi-R2-like and*

Rpi-R1

Using the same SMRT-AgRenSeq-d approach for the benchmark genes *Rpi-R3b,*

Rpi-R2-like and *Rpi-R1* independently validated our methodology and successfully

identified the functional genes amongst the positively associated contigs and

specifically amongst those yielding an F1 score of 1. Thus, the integration of

dRenSeq to validate the association of NLRs and to determine the F1 score, is a

valuable step in reducing the number of candidates rapidly.

Following the SMRT-AgRenSeq-d analyses for *Rpi-R3b,* 83 NLR containing contigs

with 114 NLRs were identified with a minimum association score of 30

(Supplementary Figure 2). When filtered for an F1 score of larger than 0.7, 21

contigs that contained 31 NLRs were identified with an association score of between

56 and 60 (Supplementary Table 5). Owing to the absence of candidates in potatoes

that contain *Rpi-R3b* and the presence of candidates in potatoes that do not contain

Rpi-R3b, only three contigs remained with an F1 score of 1. These include

tig00000194 and tig00000301 of which the latter contains *Rpi-R3b* (Table 1).

Further, all genes map to the *R3* locus on potato chromosome 11 (Supplementary

Figure 2c) and reside in the phylogenetic clade associated with *Rpi-R3b* (Figure 1).

The *Rpi-R3a* containing contig tig00000180 yielded an association score of 59 but,

since this gene was absent in 16 of the 60 potatoes with *Rpi-R3b*, the F1 score was

reduced to approximately 0.85 which ruled out this contig as a candidate

(Supplementary Table 5).

Representation of the *Rpi-R2* gene family in the association set was the lowest of all

benchmark genes. Despite less than 10% penetration of the phenotype in the panel

(10 out of 117; Supplementary Table 2), SMRT-AgRenSeq-d identified ten candidate

NLRs with an F1 score of 1 which include the *Rpi-R2-like* containing contig

tig00001243 (Supplementary Figure 3; Supplementary Table 6, Table 1). These

candidates all map to the *Rpi-R2* locus on potato chromosome 4 (Supplementary

Figure 3c) and are members of the *Rpi-R2* clade in the phylogenetic tree (Figure 1).

The validation of SMRT-AgRenSeq-d on the *Rpi-R1* benchmark gene yielded the

lowest number of associated contigs (Supplementary Figure 4) which is most likely a

reflection of the smaller size of the *Rpi-R1* gene cluster compared to *Rpi-R2* and *Rpi-*

R3a/b. Indeed, in DM, *Rpi-R1* is one of the smallest clusters of NLRs [16]. At a

minimum association score of 17, 16 NLR containing contigs containing 20 NLRs

were identified (Supplementary Figure 4). After filtering for an F1 score of larger than

0.7, 11 contigs had association scores between 22 and 34 and those encode for 13

NLRs (Supplementary Table 7). The *Rpi-R1* containing contig, tig00000086 (Table 1)

was successfully identified with an F1 score of 1.

SMRT-AgRenSeq-d identified nine candidates for Gpa5:

Following the development and independent validation of SMRT-AgRenSeq-d on all four benchmark genes, the approach was applied to identify candidates for the resistance gene *Gpa5*. The association panel of 117 potatoes, as described above, includes 23 highly resistant clones with a *G. pallida* resistance score of between 7 and 9 and which tested positive for the HC marker [11], alongside 94 susceptible clones with a resistance score of 1 – 3 (Supplementary Table 2).

Conducting SMRT-AgRenSeq-d on this panel of highly resistant and highly susceptible potatoes identified 78 NLR-containing contigs with 139 predicted NLRs and with a minimum association score of 12 (Figure 3a). Following dRenSeq analysis and filtering of candidates that exceed an F1 score of 0.7, 46 contigs encoding for 79 predicted NLRs were identified with an association score between 13 and 23 (Supplementary Table 8). These NLRs mapped to three positions in the potato genome, chromosome 5, chromosome 7, and chromosome 9 (Supplementary Table 8). The position on chromosome 5 is consistent with previous mapping studies conducted by Rouppe van der Voort *et al.* [9] which also highlighted the position on chromosome 9 that was also observed by Bryan *et al.* [13].

Following dRenSeq analysis, nine NLRs with an F1 value of 1 were identified and were considered strong candidates for the resistance (Supplementary Table 8). Consistent with Sattarzadeh *et al.* [11] and van Eck *et al.* [10] these nine candidates map to the *Rpi-R1* locus on potato Chromosome 5 (Figure 3c; Supplementary Table 8) and phylogenetically cluster with the *Rpi-R1* gene family (Figure 1).

Genetically, this study provides evidence that *Gpa5* is in repulsion to *Rpi-R1* as the *Rpi-R1* containing contig tig00000086 (Table 1) was not associated with the nematode resistance (Supplementary Table 8). This is further supported by the presence/absence of *Rpi-R1* in the association panel, where more than 60% (14 out of 23) of clones that contained *Gpa5* did not contain *Rpi-R1* (Supplementary Table 2). None of the candidates on Chromosome 9 passed the dRenSeq validation, albeit their F1 score was relatively high and ranged from 0.90 to 0.76 mainly due to their absence in some resistant varieties.

Gpa5 might require additional genes to provide full resistance

As detailed above, SMRT-AgRenSeq-d identified nine potential candidates that are strongly associated with *Gpa5*-based resistance. To further validate their role in providing resistance, we independently assessed the presence/absence of these candidate genes in three additional potato cultivars (Tiger, Novano and Karelia) that all yielded a resistance score of between 7 and 8 against *G. pallida* Pa2/3 (Supplementary Table 9). We further included five breeding clones that tested positive for the HC marker, but only displayed intermediate resistance alongside the cultivar Alicante that also provided an intermediate resistance (score of 6 against Pa3; 9 against Pa2). As a negative control, we included the cultivar Vales Everest which displayed intermediate to high resistance against *G. pallida* Pa2/3. However this cultivar utilises the *H3* resistance instead of *Gpa5* [31].

Except for Vales Everest, all candidates with an F1 score of 1 showed consistent presence in this set, including the lines with intermediate phenotypes. Interestingly, the NLRs from chromosome 9 were observed in members of the high and intermediate resistance scoring group, making them candidates for the added response. None of the *Gpa5* candidates were present in Vales Everest, which is consistent with *H3*, introduced from *S. tuberosum* ssp. *andigena* (CPC 2802), being a distinct resistance from the *S. vernei*-derived *Gpa5* [3,9].

Based on previous studies, *Gpa6* on chromosome 9 exhibits an additive effect with *Gpa5* in terms of resistance to *G. pallida* Pa2/3 [9]. The loss of *Gpa6* may account for the observed intermediate resistance in some clones in our panel whilst its absence in some resistant clones could have been masked by the presence of other functional genes such as *Grp1* or *H2*. Thus, the NLRs on chromosome 9 could represent potential *Gpa6* candidates (Supplementary Table 8) and warrant further studies in the future.

It is further conceivable that *Gpa5*, a potential *Rpi-R1* homolog (Figure 1), requires additional genes such as NRCs to function. Indeed, *Rpi-R1* requires NRC4 to elicit a functional response [28]. Thus, the intermediate resistance response observed in some potatoes, despite the presence of the candidate NLRs, could be a consequence of NRC variations that are less conducive to triggering a full response.

Conclusion

In conclusion, we have further developed SMRT-AgRenSeq by including a dRenSeq analysis that significantly expedites the process of identifying candidates that are absolutely linked to the resistance phenotype. The F1 score that can be determined for all candidates in this combined approach is a highly informative measure of linkage and proved more informative than, for example, the association score alone. The resulting method, SMRT-AgRenSeq-d, successfully identified the benchmark genes *Rpi-R1, Rpi-R2-like, Rpi-R3a* and *Rpi-R3b* with the highest possible F1 score of 1 in the association panel. Importantly, SMRT-AgRenSeq-d correctly identified those genes even though their penetration in the panel varied between 8.5% for *Rpi-R2-like*, 29.3% for *Rpi-R1*, 37.6% for *Rpi-R3a*, and 51.3% for *Rpi-R3b*. When applied to *Gpa5,* which is currently the most effective and widespread resistance against *G. pallida* used in potato crop protection, SMRT-AgRenSeq-d identified strong candidate NLRs. All *Gpa5* candidates mapped to the previously established locus on potato chromosome 5 and were shown to be phylogenetically related to the late blight resistance gene *R1*. We propose that this refined method is suitable for studying NLR-based resistances in diverse crops and against diverse pathogens.

Materials and Methods:

Source of plant and pathogen material

G. pallida cysts that were used in the phenotyping test were of the Pa2/3 population known as 'Lindley' [32]. The population was multiplied on the susceptible cultivar Desiree in glasshouses in 2019. All potato accessions were grown in the glasshouse. In some cases, lyophilised DNA was received from collaborators.

Extraction of DNA

Bulk DNA was extracted from leaves of all potato accessions using the Qiagen DNEasy Plant Mini Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. DNA for HiFi sequencing was extracted using the Wizard® HMW DNA Extraction Kit (Promega, Madison, WI, USA), following the manufacturer's instructions.

RenSeq

453 The RenSeq bait library was as described in Armstrong *et al.* [20]. Enrichment
454 sequencing was performed by Arbor Biosciences (Ann Arbor, MI, USA) using the
455 Novaseq 6000 platform for Illumina reads and the PacBio Sequel II platform for HiFi
456 reads.

Assembly

Raw Illumina reads were mapped to the DM v6.1 genome assembly [33] with BWA-
mem v0.7.17-r1188 [34] using default parameters. Picard CollectInsertSizeMetrics
v2.25.1 [35] was used to calculate the mean and standard deviation of the read
insert size. This was used to inform the MASURCA [36] v3.4.2 assembly with PE=
pe 342 97 and CLOSE_GAPS=1 as non-default parameters. Assembly of HiFi reads
was conducted using HiCanu with the SMRT-RenSeq Assembly workflow within
HISS v2.0.0 [37,38].

Identification of known resistance genes

468 The CDS sequences of *R1, R2-like, R3a* and *R3b* [39–42] were searched for in both
469 sets of assembled contigs using BLASTN v2.11.0 [43,44] with an e-value of 1e-5.

Phylogenetics

NLR sequences from the SMRT-RenSeq assemblies were predicted using NLR-
Annotator [25] and Nucleotide Binding (NB) domains were identified with
Interproscan version 5.54-87.0 [45,46] with goterms, iprlookup and pathways options
enabled. The output XML file was parsed with a custom python script [47] to extract
the amino acid sequences of the NB domains (IPR002182). Reference NRC amino
acid sequences were retrieved from Adachi *et al.* [48] and submitted to Interproscan
and the output tsv file was parsed with a different custom python script [47]. The
BED files produced by these scripts were used to extract FASTA sequences with the
bedtools getfasta function (version 2.30.0) [49]. The known NLR and NRC gene NB
domain sequences were aligned with Clustal Omega version 1.2.4 [50] with 10
iterations. Following this, the NB domains of the automatically predicted NLRs were
added to the alignment with Clustal Omega using the previous alignment as a profile,
again with 10 iterations.

Tree construction was performed in R v4.1.3 [51] using the libraries ape v5.6-2 [52] and phangorn v2.9.0 [53]. Sites and sequences with 85% missing data, or more, were removed from the alignment. Following this, duplicated sequences were removed from the alignment. A phylogeny based on maximum likelihood was inferred from the final alignment. The Bayesian information criteria (BIC) obtained from the "modelTest" function of phangorn suggested a JTT+G model. The tree was bootstrapped with 1,000 replicates and its topology was optimised using nearest neighbour interchanges (NNI). The phylogeny produced was then rooted to the outgroup sequences from *Homo sapiens* and *Caenorhabditis elegans*. Clades were assigned with PhyloPart 2.1 [54] with a percentile threshold of 0.05. The figure was created in IToL [55], details on how to replicate the figure and text files for modifying the view are described in a markdown file [47].

Diversity panel formation and additional phenotyping

The resistance or susceptibility of clones within the association panel was determined by retrieving results from potato cultivar databases and/or through replicated PCN tests. Only resistant clones that also tested positive for the HC marker [11] were considered to ensure that the source of the resistance is consistent (Supplementary Table 3; Supplementary Table 9). The phenotyping experiment was based on the method published by [56]. Briefly, tubers of breeding clones and six controls (Desiree, Maris Piper, Vales Everest, Lady Balfour, Innovator and King Russet) were planted in canisters (one tuber per canister) with four replicates. All canisters were placed into four trays, each containing one replicate of all breeding clones and controls in a randomized design. Canisters were inoculated with approximately 10–15 cysts. After inoculation, canisters were stored in the dark at 18°C for seven weeks. After this time, the number of females observed on the root balls were counted, and the phenotypic scores were recorded. In these accessions and breeding clones, 117 lines whose scores were between 1 and 3 (susceptible accessions) and 7–9 (resistant accessions) were chosen for the diversity panel for *Gpa5*.

To determine the presence/absence of the late blight resistance genes *Rpi-R1, Rpi-R2-like, Rpi-R3a* and *Rpi-R3b* that are present in Innovator, dRenSeq was

performed as described previously by Armstrong *et al.* [20]. Based on the known

diversity within the *Rpi-R2* family [20], we set a minimum sequence coverage

threshold of 97% for a gene to be classified as present.

SMRT-AgRenSeq-d

SMRT-AgRenSeq-d was performed using the AgRenSeq and dRenSeq workflows in

HISS v2.0.0 [37]. AgRenSeq followed the workflow initially described for *A. tauschii*

[22]. Briefly, the Illumina reads of the previously described panel of samples were

assessed for the presence and absence of *k*-mers with a *k* value of 51 and arranged

into a presence/absence matrix. This matrix was then used alongside phenotypic

information to perform the association and the identities of contigs containing *k*-mers

highly associated with the phenotype were reported.

An initial threshold of 50% of the highest association score was used for selecting

candidate contigs and the NLR coordinates were determined with NLR-Annotator

[25] (*Rpi-R1*: 17; *Rpi-R2-like*: 5; *Rpi-R3a*: 22; *Rpi-R3b*: 30; *Gpa5*: 12). These

selected contigs were used for dRenSeq. This followed the pipeline previously

described [20,57] whilst specifically reporting on NLR regions linked to bait coverage.

Briefly, the Illumina reads of the panel were mapped to the CDS regions of candidate

genes. This mapping was then filtered to contain only reads with zero mismatches

within regions also represented by baits with at least 90% sequence identity. The

resulting coverage of the candidates was reported.

Refining of candidates

Candidates identified by SMRT-AgRenSeq-d were further refined through the

generation of precision and recall values and calculation of the F1 score of the

candidates [30]. The following formula was applied:

$$F1 = \frac{2 \times PPV \times TPR}{PPV + TPR}$$

548 PPV is the Positive Predictive Value or precision, calculated by the following formula:

$$PPV = \frac{TP}{TP + FP}$$

551 TPR is the True Positive Rate or sensitivity/recall, calculated by the following
552 formula:

$$TPR = \frac{TP}{TP + FN}$$

Here, TP represents true positive cases i.e., varieties/clones that are phenotypically
resistant and have complete coverage of the candidate in dRenSeq (within the
parameters specified above). FP (false positive) refers to varieties that are
phenotypically susceptible but still had complete coverage in dRenSeq. Conversely,
TN (true negative) refers to varieties that are phenotypically susceptible and had only
partial coverage in dRenSeq, whereas FN (false negative) refers to phenotypically
resistant varieties with partial coverage in dRenSeq. The individual F1 scores were
calculated for every candidate.

Acknowledgements

565
The authors thank John Jones for providing critical feedback on the manuscript.
Further we acknowledge the Research/Scientific Computing teams at The James
Hutton Institute and NIAB for providing computational resources and technical
support for the "UK's Crop Diversity Bioinformatics HPC" (BBSRC grant
BB/S019669/1), use of which has contributed to the results reported within this
paper.

This work was supported by the Rural & Environment Science & Analytical Services
(RESAS) Division of the Scottish Government through project JHI-B1-1, the
Biotechnology and Biological Sciences Research Council (BBSRC) through award
BB/S015663/1 and the Royal Society through award NAF\R1\201061. YW was
supported through the CSC scholarship program, China. LB was supported through
the East of Scotland Bioscience Doctoral Training Partnership (EASTBIO DTP),
funded by the BBSRC award BB/T00875X/1. AK was supported through a Research

Leaders 2025 fellowship funded by European Union's Horizon 2020 research and
innovation programme under Marie Sklodowska-Curie grant agreement no. 754380.

Contributions

YW, LB, TMA, MA, AK conducted the computational analyses, developed the
association panel and the F1 score. JL, XW, BH provided material for this study and
contributed towards the writing of the manuscript and the independent validation. VY,
DT and YW conducted the PCN score and HC marker analysis. TMA and KN
conducted the phylogenetic analysis. SO provided genomic data and recombination
analysis for this study. MB and IH conceived this work and secured funding. All
authors have read and endorsed the manuscript.

Data availability statement

591 BioProjects:
592 ERP141787 – Illumina Reads / contigs
593 ERP141789 – HiFi Reads / contigs

Conflict of interests

The authors declare that they have no competing interests.

Figure legends:

Figure 1: Phylogenetic tree displaying NB domains of Innovator NLRs following
assembly of SMRT-RenSeq reads. Expanded are the clades containing *NRCs, Rpi-*
R1, *Rpi-R2-like, Rpi-R3a* and *Rpi-R3b*. Collapsed clades either consist of automatic
predictions from NLR-Annotator or of NLR clades not investigated in this study.
Highlighted are the identified benchmark genes and candidates from each
association with F1=1. The NB containing proteins APAF1 from *H. sapiens* and
CED4 from *C. elegans* act as outgroups. The tree scale indicates substitutions/site.

Figure 2: SMRT-AgRenSeq-d of *Rpi-R3a*. a) Identification of NLR containing contigs
in the reference cultivar Innovator with association to *Rpi-R3a*. Columns on the X-
axis represents NLR containing contigs, with dots indicating mapped *k*-mers and
their association score to *Rpi-R3a* on the Y-axis. Dashed lines highlight highest
association score (44, red) and chosen threshold (22, Orange). b) F1 scores and
number of candidate NLRs for each association score group. c) The predicted
positions of *Rpi-R3a* candidates with F1=1 are shown (dots) in relation to all NLRs
from the reference DM v6.1 mapped over 12 chromosomes (peaks).

Figure 3: SMRT-AgRenSeq-d of *Gpa5*. a) Identification of NLR containing contigs in
the reference cultivar Innovator with association to *Gpa5*. Columns on the X-axis
represents contigs with signals of NLRs, with dots indicating mapped *k*-mers and
their association score to *Gpa5* on the Y-axis. Dashed lines highlight highest
association score (23, red) and chosen threshold (12, Orange). b) F1 score and
number of candidate NLRs for each association score group. c) The predicted
positions of *Gpa5* candidates with F1=1 are shown (dots) in relation to all NLRs from
the reference DM v6.1 mapped over 12 chromosomes (peaks).

Supplementary Tables:

Supplementary Table 1: Read and assembly statistics for RenSeq and SMRT-
RenSeq derived sequence presentation of the NLRome from the cultivar Innovator.

Supplementary Table 2: Description of the association panel. Highlighted are the
cultivars and breeding clones used in this study and their coverage of the benchmark
genes *Rpi-R1, Rpi-R2-like, Rpi-R3a* and *Rpi-R3b* as established through dRenSeq.
Coverage exceeding 97% of the full-length gene equates to the presence of the
gene. Presence/absence is transposed into a +1/-1 score used for the association
study. For *Gpa5*, only cultivars that yielded a disease resistance score between 7
and 9 or 1–3, were used to represent resistant/susceptible, respectively. Further, all
resistant plants had to also yield a positive PCR test for the HC marker to ensure the
source of the resistance is consistent. ENA read accession numbers are highlighted
for all plants in the panel.

Supplementary Table 3: Immediate descendent potato cultivars and breeding clones
from the cultivar Innovator (URL: http://www.plantbreeding.wur.nl/PotatoPedigree/).
Shown are the parents of the cross, the year, breeder and breeder code.

Supplementary Table 4: dRenSeq analysis of *Rpi-R3a.* Depicted is the dRenSeq-
based coverage of NLR candidates with an association score of larger than 22
across 117 potato clones representing the association panel. Based on the
presence/absence of candidate NLRs in potato clones associated with the *Rpi-R3a*
and those that do not contain *Rpi-R3a*, an F1 score was calculated. Only candidates
with an F1 score of 1 display absolute linkage to the phenotype.

Supplementary Table 5: dRenSeq analysis of *Rpi-R3b.* Depicted is the dRenSeq-
based coverage of NLR candidates with an association score of larger than 30 and
an F1 score of larger than 0.7 across 117 potato clones representing the association
panel. The F1 score was calculated based on the presence/absence of candidate
NLRs in potato clones associated with the *Rpi-R3b* and those that do not contain
Rpi-R3b.

Supplementary Table 6: dRenSeq analysis of *Rpi-R2-like*. Depicted is the dRenSeq-
based coverage of NLR candidates with an association score of larger than 5 and an
F1 score of larger than 0.7 across 117 potato clones representing the association
panel. The F1 score was calculated based on the presence/absence of candidate
NLRs in potato clones associated with the *Rpi-R2-like* and those that do not contain
Rpi-R2-like.

Supplementary Table 7: dRenSeq analysis of *Rpi-R1*. Depicted is the dRenSeq-
based coverage of NLR candidates with an association score of larger than 17 and
an F1 score of larger than 0.7 across 116 potato clones representing the association
panel. The F1 score was calculated based on the presence/absence of candidate
NLRs in potato clones associated with the *Rpi-R1* and those that do not contain *Rpi-*
R1.

Supplementary Table 8: dRenSeq analysis of *Gpa5.* Depicted is the dRenSeq-based
coverage of NLR candidates with an association score of larger than 12 and an F1
score of larger than 0.7 across 117 potato clones representing the association panel.
The F1 score was calculated based on the presence/absence of candidate NLRs in
potato clones associated with the *Gpa5* and those that do not contain *Gpa5.*

Supplementary Table 9: dRenSeq analysis of *Gpa5* candidate NLRs in additional
potato clones with high and intermediate resistance against *Globodera pallida* Pa2/3.
The cultivar Vales Everest is used as a negative control since the resistance is
known to be independent of *Gpa5* and utilises instead *H3*.

Supplementary Figures

Supplementary Figure 1: Phylogenetic tree displaying NB domains of Innovator NLRs following assembly of SMRT-RenSeq reads. All clades are expanded. Highlighted are the identified benchmark genes and candidates from each association with F1=1. The NB containing proteins APAF1 from *Homo sapiens* and CED4 from *Caenorhabditis elegans* act as outgroups. Bootstrap values are displayed as branch labels. The tree scale indicates substitutions/site.

Supplementary Figure 2: SMRT-AgRenSeq-d analysis for *Rpi-R3b*. a) Columns on the X-axis represents NLR containing contigs, with dots indicating mapped *k*-mers and their association score to *Rpi-R3b* on the Y-axis. Dashed lines highlight highest association score (60, red) and chosen threshold (30, Orange). b) F1 score and the number of candidate NLRs for each association score group. c) The predicted position of *Rpi-R3b* candidates with F1=1 are shown (dots) in relation to all NLRs from the reference DM v6.1 mapped over 12 chromosomes (peaks).

Supplementary Figure 3: SMRT-AgRenSeq-d analysis for *Rpi-R2-like*. a) Columns on the X-axis represents NLR containing contigs, with dots indicating mapped *k*-mers and their association score to *Rpi-R2-like* on the Y-axis. Dashed lines highlight highest association score (10, red) and chosen threshold (5, Orange). b) F1 score and the number of candidate NLRs for each association score group. c) The predicted position of *Rpi-R2-like* candidates with F1=1 are shown (dots) in relation to all NLRs from the reference DM v6.1 mapped over 12 chromosomes (peaks).

Supplementary Figure 4: SMRT-AgRenSeq-d analysis for *Rpi-R1*. a) Columns on the X-axis represents NLR containing contigs, with dots indicating mapped *k*-mers and their association score to *Rpi-R1* on the Y-axis. Dashed lines highlight highest association score (34, red) and chosen threshold (17, Orange). b) F1 score and the number of candidate NLRs for each association score group. c) The predicted position of *Rpi-R1* candidates with F1=1 are shown (dots) in relation to all NLRs from the reference DM v6.1 mapped over 12 chromosomes (peaks).

References

1. Birch PRJ, Bryan G, Fenton B *et al.* Crops that feed the world 8: Potato: are the trends of increased global production sustainable? *Food Sec* 2012;**4**:477–508.

2. Tiwari JK, Buckseth T, Zinta R *et al.* Germplasm, Breeding, and Genomics in Potato Improvement of Biotic and Abiotic Stresses Tolerance. *Frontiers in Plant Science* 2022;**13**.

3. Gartner U, Hein I, Brown LH *et al.* Resisting Potato Cyst Nematodes With Resistance. *Frontiers in Plant Science* 2021;**12**.

4. Jones JT, Haegeman A, Danchin EGJ *et al.* Top 10 plant-parasitic nematodes in molecular plant pathology. *Molecular Plant Pathology* 2013;**14**:946–61.

5. Turner SJ. Population decline of potato cyst nematodes (Globodera rostochiensis, G. pallida) in field soils in Northern Ireland. *Annals of Applied Biology* 1996;**129**:315–22.

6. Price JA, Coyne D, Blok VC *et al.* Potato cyst nematodes Globodera rostochiensis and G. pallida. *Molecular Plant Pathology* 2021;**22**:495–507.

7. Paal J, Henselewski H, Muth J *et al.* Molecular cloning of the potato Gro1-4 gene conferring resistance to pathotype Ro1 of the root cyst nematode Globodera rostochiensis, based on a candidate gene approach. *The Plant Journal* 2004;**38**:285–97.

8. Van Der Vossen EAG, Van Der Voort JNAMR, Kanyuka K *et al.* Homologues of a single resistance-gene cluster in potato confer resistance to distinct pathogens: a virus and a nematode. *The Plant Journal* 2000;**23**:567–76.

9. Rouppe van der Voort J, van der Vossen E, Bakker E *et al.* Two additive QTLs conferring broad-spectrum resistance in potato to Globodera pallida are localized on resistance gene clusters: *Theor Appl Genet* 2000;**101**:1122–30.

10. van Eck HJ, Vos PG, Valkonen JPT *et al.* Graphical genotyping as a method to map Ny(o,n)sto and Gpa5 using a reference panel of tetraploid potato cultivars. *Theor Appl Genet* 2017;**130**:515–28.

11. Sattarzadeh A, Achenbach U, Lübeck J *et al.* Single nucleotide polymorphism (SNP) genotyping as basis for developing a PCR-based marker highly diagnostic for potato varieties with high resistance to Globodera pallida pathotype Pa2/3. *Mol Breeding* 2006;**18**:301–12.

12. Finkers-Tomczak A, Danan S, van Dijk T *et al.* A high-resolution map of the Grp1 locus on chromosome V of potato harbouring broad-spectrum resistance to the cyst nematode species Globodera pallida and Globodera rostochiensis. *Theor Appl Genet* 2009;**119**:165–73.

13. Bryan G, McLean K, Bradshaw J *et al.* Mapping QTLs for resistance to the cyst
nematode Globodera pallida derived from the wild potato species Solanum vernei.
Theor Appl Genet 2002;**105**:68–77.

14. Dalton E, Griffin D, Gallagher TF *et al.* The effect of pyramiding two potato cyst
nematode resistance loci to Globodera pallida Pa2/3 in potato. *Mol Breeding*
2013;**31**:921–30.

15. Jupe F, Witek K, Verweij W *et al.* Resistance gene enrichment sequencing
(RenSeq) enables reannotation of the NB-LRR gene family from sequenced plant
genomes and rapid mapping of resistance loci in segregating populations. *The Plant*
Journal 2013;**76**:530–44.

16. Jupe F, Pritchard L, Etherington GJ *et al.* Identification and localisation of the
NB-LRR gene family within the potato genome. *BMC Genomics* 2012;**13**:75.

17. Xu X, Pan S, Cheng S *et al.* Genome sequence and analysis of the tuber crop
potato. *Nature* 2011;**475**:189–95.

18. Van de Weyer A-L, Monteiro F, Furzer OJ *et al.* A Species-Wide Inventory of
NLR Genes and Alleles in Arabidopsis thaliana. *Cell* 2019;**178**:1260-1272.e14.

19. Witek K, Jupe F, Witek AI *et al.* Accelerated cloning of a potato late blight–
resistance gene using RenSeq and SMRT sequencing. *Nature Biotechnology*
2016;**34**:656–60.

20. Armstrong MR, Vossen J, Lim TY *et al.* Tracking disease resistance deployment
in potato breeding by enrichment sequencing. *Plant Biotechnology Journal*
2019;**17**:540–9.

21. Chen X, Lewandowska D, Armstrong MR *et al.* Identification and rapid mapping
of a gene conferring broad-spectrum late blight resistance in the diploid potato
species Solanum verrucosum through DNA capture technologies. *Theoretical and*
Applied Genetics 2018;**131**:1287–97.

22. Arora S, Steuernagel B, Gaurav K *et al.* Resistance gene cloning from a wild
crop relative by sequence capture and association genetics. *Nat Biotechnol*
2019;**37**:139–43.

23. Vendelbo NM, Mahmood K, Steuernagel B *et al.* Discovery of Resistance Genes
in Rye by Targeted Long-Read Sequencing and Association Genetics. *Cells*
2022;**11**:1273.

24. Corrion A, Day B. Pathogen Resistance Signalling in Plants. *ELS*. John Wiley &
Sons, Ltd, 2015, 1–14.

25. Steuernagel B, Witek K, Krattinger SG *et al.* The NLR-Annotator Tool Enables
Annotation of the Intracellular Immune Receptor Repertoire. *Plant Physiol*
2020;**183**:468–82.

26. Baumgarten A, Cannon S, Spangler R *et al.* Genome-Level Evolution of
Resistance Genes in Arabidopsis thaliana. *Genetics* 2003;**165**:309–19.

27. Huang Z, Qiao F, Yang B *et al.* Genome-wide identification of the NLR gene
family in Haynaldia villosa by SMRT-RenSeq. *BMC Genomics* 2022;**23**:118.

28. Wu C-H, Abd-El-Haliem A, Bozkurt TO *et al.* NLR network mediates immunity to
diverse plant pathogens. *Proc Natl Acad Sci U S A* 2017;**114**:8113–8.

29. van Berloo R, Hutten RCB, van Eck HJ *et al.* An Online Potato Pedigree
Database Resource. *Potato Res* 2007;**50**:45–57.

30. Chinchor N, Diego S. MUC-4 EVALUATION METRICS. 1992, 22–9.

31. Adillah Tan MY, Park T-H, Alles R *et al.* GpaXItarloriginating from Solanum
tarijense is a major resistance locus to Globodera pallida and is localised on
chromosome 11 of potato. *Theor Appl Genet* 2009;**119**:1477–87.

32. Kaczmarek AM, Back M, Blok VC. Population dynamics of the potato cyst
nematode, Globodera pallida, in relation to temperature, potato cultivar and
nematicide application. *Plant Pathology* 2019;**68**:962–76.

33. Pham GM, Hamilton JP, Wood JC *et al.* Construction of a chromosome-scale
long-read reference genome assembly for potato. *GigaScience* 2020;**9**, DOI:
10.1093/gigascience/giaa100.

34. Li H. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. 2013, DOI: 10.48550/arXiv.1303.3997.

35. Broad Institute,. "Picard Toolkit." *GitHub repository*
https://broadinstitute.github.io/picard/, Broad Institute 2019.

36. Zimin AV, Marçais G, Puiu D *et al.* The MaSuRCA genome assembler.
Bioinformatics 2013;**29**:2669–77.

37. Adams T, Smith M, Bayer M *et al.* HISS: Snakemake-based workflows for
performing SMRT-RenSeq assembly, AgRenSeq and dRenSeq for the discovery of
novel plant disease resistance genes. 2022:2022.11.01.514708.

38. Nurk S, Walenz BP, Rhie A *et al.* HiCanu: accurate assembly of segmental
duplications, satellites, and allelic variants from high-fidelity long reads. *Genome Res*
2020;**30**:1291–305.

39. Ballvora A, Ercolano MR, Weiss J *et al.* The R1 gene for potato resistance to late
blight (Phytophthora infestans) belongs to the leucine zipper/NBS/LRR class of plant
resistance genes. *Plant J* 2002;**30**:361–71.

40. Huang S, Van Der Vossen EAG, Kuang H *et al.* Comparative genomics enabled
the isolation of the R3a late blight resistance gene in potato. *The Plant Journal*
2005;**42**:251–61.

41. Li G, Huang S, Guo X *et al.* Cloning and Characterization of R3b; Members of
the R3 Superfamily of Late Blight Resistance Genes Show Sequence and Functional
Divergence. *MPMI* 2011;**24**:1132–42.

42. Lokossou AA, Park T, van Arkel G *et al.* Exploiting Knowledge of R/Avr Genes to
Rapidly Clone a New LZ-NBS-LRR Family of Late Blight Resistance Genes from
Potato Linkage Group IV. *MPMI* 2009;**22**:630–41.

43. Altschul SF, Gish W, Miller W *et al.* Basic local alignment search tool. *J Mol Biol*
1990;**215**:403–10.

44. Zhang Z, Schwartz S, Wagner L *et al.* A greedy algorithm for aligning DNA
sequences. *J Comput Biol* 2000;**7**:203–14.

45. Blum M, Chang H-Y, Chuguransky S *et al.* The InterPro protein families and
domains database: 20 years on. *Nucleic Acids Research* 2021;**49**:D344–54.

46. Jones P, Binns D, Chang H-Y *et al.* InterProScan 5: genome-scale protein
function classification. *Bioinformatics* 2014;**30**:1236–40.

47. Adams T, Wang Y, Wangand K. TMAdams/gpa5_smrt_agrenseq_paper: v1.0.0.
2022.

48. Adachi H, Sakai T, Harant A *et al.* An atypical NLR protein modulates the NRC
immune receptor network. 2021:2021.11.15.468391.

49. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 2010;**26**:841–2.

50. Sievers F, Wilm A, Dineen D *et al.* Fast, scalable generation of high-quality
protein multiple sequence alignments using Clustal Omega. *Molecular Systems*
Biology 2011;**7**:539.

51. R Core Team. R: A language and environment for statistical computing. R: A
language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. URL https://www.R-project.org/. 2022.

52. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* 2019;**35**:526–8.

53. Schliep KP. phangorn: phylogenetic analysis in R. *Bioinformatics* 2011;**27**:592–3.

54. Prosperi MCF, Ciccozzi M, Fanti I *et al.* A novel methodology for large-scale
phylogeny partition. *Nat Commun* 2011;**2**:321.

55. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for
phylogenetic tree display and annotation. *Nucleic Acids Research* 2021;**49**:W293–6.

56. Phillips MS, Forrest JMS, Wilson LA. Screening for resistance to potato cyst
nematode using closed containers. *Annals of Applied Biology* 1980;**96**:317–22.

57. Van Weymers PSM, Baker K, Chen X *et al.* Utilizing "Omic" Technologies to
Identify and Prioritize Novel Sources of Resistance to the Oomycete Pathogen
Phytophthora infestans in Potato Germplasm Collections. *Front Plant Sci* 2016;**7**,
DOI: 10.3389/fpls.2016.00672.

Tree scale: 1

Benchmark Genes and Candidates

- R1
- R1 Candidates
- R2-like
- R2-like Candidates
- R3a
- R3a Candidates
- R3b
- R3b Candidates
- Gpa5 Candidates

R2-like

R1

R3

NRCs