Title Page

Title

Neuron- and microglia-specific immunoexpression in steroid-independent male sexual behaviour in

castrated B6D2F1 male mice

Authors

CHIANG, Vic Shao-Chih[1]; LIPKIN, Sophie [1]; MCCUSKER, Catherine D[2]; PARK, Jin Ho[1]

Affiliation

[1] Developmental and Brain Sciences, Department of Psychology, University of Massachusetts, Boston,

MA, United States

[2] Department of Biology, University of Massachusetts, Boston, MA, United States

Running title (40 characters maximum)

Steroid-independent sexual behaviour

Corresponding author

CHIANG, Vic Shao-Chih

Address

Developmental and Brain Sciences, Department of Psychology, University of Massachusetts, Boston,

MA, United States

E-mail address

V.chiang001@umb.edu

Conflict of interest

The authors declare no conflict of interest.

Total manuscript word count

11435 (including tables, figure legends, excluding references)

Page count (excluding references): 41

Number of figures: 5, and 5 supplementary figures

Number of tables: 4

Abstract (maximum 250 words)

Sexual behaviour is necessary for procreation for several species and is traditionally viewed to be

regulated by sex steroid hormones. However, several species exhibit steroid-independent sexual

behaviour, and its molecular understanding is only beginning to be uncovered. The main goal of our

experiment was to provide new insight into cell-specific roles that both neuronal and non-neuronal cells

may play in steroid-independent male sexual behaviour. Forty B6D2F1 hybrid male mice underwent

orchidectomy and were tested for reinstatement of steroid-independent male sexual behaviour after an

extended period of social isolation caused by the COVID-19-mandated laboratory shutdown. After 62

weeks post-orchidectomy, 20.59% demonstrated reinstatement of steroid-independent male sexual

behaviour (identified as 'steroid-independent persistent maters'), while 23.53% of the males did not

display steroid-independent male sexual behaviour (identified as 'steroid-independent non-maters').

Using flow cytometry, we compared the preoptic area immunoexpression in NeuN+ neurons and Iba1+

microglia between steroid-independent persistent maters and steroid-independent non-maters (n = 5-6

per group). We found neuronal immunoexpression up-regulated for amyloid precursor protein and

androgen receptor, as well as down-regulated for glucocorticoid receptor in steroid-independent

persistent maters compared to steroid-independent non-maters. In conjunction, microglial

immunoexpression of amyloid precursor protein was up-regulated in steroid-independent persistent

maters compared to steroid-independent non-maters. These data suggest there are cell-specific

immunoexpression differences, including the role of non-neuronal cells in steroid-independent male

sexual behaviour.

Keywords (maximum 6): Steroid independent; sexual behaviour; B6D2F1 hybrid male mice;

mechanism; microglia

Introduction

Sexual behaviour is integral to many organisms for procreation, social bonding, cooperation, dominance, and even bartering; additionally, it is not limited to the opposite sex and conspecifics (Pelé *et al.*, 2017; Monk *et al.*, 2019; Addessi & Bourgeois-Gironde, 2020; Steinberg, 2020; Garcia *et al.*, 2021; Lemoine *et al.*, 2022). The canonical view on how sexual behaviour is regulated involves sex steroid hormones (reviewed in (Wei *et al.*, 2021)), but several lines of evidence in various species demonstrate the prevalence of steroid-independent sexual behaviour (SIMSB; reviewed in (Chiang & Park, 2020)). One example is the B6D2F1 male mouse, which is a hybrid of C57BL/6J females and DBA/2J males. Approximately one-third of them (~33%) can exhibit gonadal SIMSB for up to four months after orchidectomy (McGill & Tucker, 1964; Manning & Thompson, 1976; Clemens *et al.*, 1988). Previous studies from our lab found molecular differences in the brain between 'steroid-independent (SI) maters' (those that demonstrated gonadal SIMSB long after orchidectomy) and 'SI non-maters' (Park *et al.*, 2010). For instance, increased amyloid precursor protein (*App)* gene expression and immunoexpression were uncovered using microarray and western blot, respectively. Follow-up experiments using transgenic mice that overexpressed APP revealed a higher proportion of mice exhibiting SIMSB compared to wildtype (Park *et al.*, 2010). These experiments microdissected samples from the medial preoptic area (POA) and the bed nucleus of the stria terminalis for downstream assays, as both are integral sites for MSB (reviewed in (Tsuneoka & Funato, 2021)). While these generated valuable insights, there were several limitations. One limitation was the heterogeneity of the sample could not be captured, which neglected expression differences in various cell types in the brain and neuron subtypes. With the development of single-cell omics in the past decade, the investigation into the specificity of cell types is much more feasible, substantially propelling our understanding of the neuroscience of social behaviour (Serrano-Saiz & Isogai, 2021; Stein *et al.*, 2021). For example, single-cell transcriptomics studies have discovered ~70 different neuronal subtypes in the POA, with specific cell types that are associated with sexual behaviour in males and females (Moffitt *et al.*, 2018; Zhang *et al.*, 2021). Additionally, an optogenetic study was able to control sexual behaviour

by targeting dopamine neurons that projected to the medial POA from the anteroventral and preoptic

periventricular area (Zhang *et al.*, 2021). However, there remains a lack of knowledge on cell-specific

regulations of SIMSB, especially of non-neuronal cells.

In addition to neurons, single-cell sequencing studies uncovered clusters of glial subtypes in

brain regions that are crucial for sexual behaviour (Moffitt *et al.*, 2018). Of particular interest are

microglia, which are resident immune cells in the brain (Lenz *et al.*, 2013; Paolicelli *et al.*, 2022).

Microglia are critical during development to wire neurocircuits required for sexual behaviour (Lenz *et*

al., 2013). There are other glial types in the brain such as astrocytes (Blanco-Suárez *et al.*, 2017), but

thus far, there is no evidence that supports their role in sexual behaviour. While existing studies do

support the involvement of microglia in sexual behaviour, much of this is not understood, especially in

terms of adult sexual behaviour. In light of previous findings on SIMSB transcriptomics, it would be of

interest to determine to what extent neurons and microglia contributed to the APP up-regulation.

Another line of inquiry would be to investigate sex steroid receptors in SIMSB, given how they are

paramount for sexual behaviour (reviewed in (Chiang & Park, 2020)). While the previous microarray

result did not find differential gene expression in AR, GR and ERa at the bulk level (Park *et al.*, 2010),

it remains to be determined whether there are cell-specific differences. Elucidating these cell-specific

differences would inform future studies that could test the causal mechanisms of SIMSB.

Here, using the B6D2F1 hybrid male mice, we replicated findings from previous studies that a

significant proportion exhibit SIMSB after long-term orchidectomy, but this was at 20.59%, compared

to the ~33% in previous studies (Park *et al.*, 2009; Templin *et al.*, 2019). However, it is important to

note that the present study differs from past iterations due to an inclusion of a COVID-19-mandated

laboratory shutdown. This shutdown led to an extended period of social isolation for all of our mice,

and social isolation is well-documented to impair rodent MSB (Liu *et al.*, 2019). For example, in

postweaning isolated CD-1 male mice, they exhibited shorter mating duration, longer mount latencies,

and longer intromission latencies relative to controls that were not socially isolated (Liu *et al.*, 2019).

While the effects of extended social isolation on SIMSB have not been directly tested in B6D2F1

hybrid male mice, a study which investigated the role of sexual experience on SIMSB may provide

some insight into the potential impact of extended social isolation on SIMSB. Immediately after

castration, B6D2F1 hybrid male mice were socially isolated for 12 weeks to withhold sexual

experience for these mice (Wee & Clemens, 1989). Interestingly, they found that a significant

proportion of the castrated B6D2F1 hybrid male mice were able to demonstrate reinstatement of

SIMSB; however, the latencies for mounting, intromission, and ejaculation were longer, as well as the

percentage of tests with mounts was lower when compared with males that were given weekly

exposure to sexually receptive females (Wee & Clemens, 1989). In our current study, the COVID-19-

mandated laboratory shutdown led to our mice being socially isolated for 23 weeks, and all of the mice

that had demonstrated SIMSB consistently prior to the shutdown did not demonstrate any MSB on the

first sexual behaviour test back. However, after they were provided social exposure, 20.59% of the

males demonstrated consistent SIMSB at least 62 weeks post-orchidectomy. Because this current study

differed from the previous SIMSB paradigms in our lab, we refer to the males that demonstrated

SIMSB in this study as 'SI persistent maters' and those that did not as 'SI non-maters' to differentiate

these mice from prior studies (David, Wyrosdic, & Park, 2022; David, Wyrosdic, Wan, *et al.*, 2022).

Once SIMSB was reinstated, we then set out to test our original experiment which was to investigate

neuron and microglia-specific immunoexpression using flow cytometry in the POA. We hypothesized

there would be neuron and microglia-specific expression profiles of APP and steroid receptors that

differ between SI persistent maters and SI non-maters.

Materials & Methods

Animals

Details of our animal model have been provided previously (Bharadwaj *et al.*, 2013). In brief, male B6D2F1 hybrid mice (*Mus musculus;* n = 40, RRID: MGI:5651959*)* were produced by crossing C57BL/6J females (RRID: IMSR_JAX:000664) with DBA/2J males (RRID: IMSR_JAX:000671). All males were bred and raised in the animal facility at the University of Massachusetts, Boston, weaned at 21 – 28 days, single-sex group-housed until the beginning of experiments (between 50 and 80 d of age), and individually housed afterwards for the rest of the experiments. All mice were housed in the same room, with cages individually ventilated by negative airflow to prevent inter-cage transmission of olfactory cues (Keller *et al.*, 2006; McCarthy *et al.*, 2017). Animals were given food (Teklad Global Rodent Diet 2016, Harlan Laboratories Inc.) and water *ad libitum* and maintained on a 12:12 light: dark cycle, with lights off at 1200 h EST. All procedures were performed according to the University of Massachusetts, Boston, IACUC (#IACUC2020). All surgeries and sacrifices that were performed were conducted while the animals were anaesthetized with isoflurane, and all efforts were made to minimize suffering including how the mice were handled (Guerra *et al.*, 2021). Meloxicam (LKT Labs; 0.5mg/mL in autoclaved distilled water) was given as a post-surgery analgesic.

Details of our sexual experience procedure have also been provided previously (McInnis *et al.*, 2016). In brief, C57BL/6J females were used as stimulus females for sexual experience and MSB testing. Females were ovariectomized in adulthood after at least one parturition and injected subcutaneously with oestradiol benzoate (MilliporeSigma; 10μg in 0.1 mL sesame oil) 48 h before testing followed by subcutaneous progesterone administration (Millipore Sigma; 400μg in 0.05 mL sesame oil) 3-6h before testing. The females were group-housed in the same colony room as the experimental males. B6D2F1 hybrid males were provided with sexual experience on four separate occasions before orchidectomy by introducing the stimulus female into the home cage and left overnight.

Assessment of male sexual behaviour

Details of our male sexual behaviour assessments have been provided previously (Templin *et al.*, 2019). In brief, after B6D2F1 hybrid males were surgically castrated with orchidectomy, they were allowed to recover for at least one week before assessing for MSB. The current paradigm differs distinctly from our previous iterations due to COVID-19 where any lab work was suspended for 23 weeks. In this paradigm, the B6D2F1 hybrid males were tested for MSB with stimulus females 11 times over 62 weeks after orchidectomy, and the suspension occurred between tests #2 and #3 (Figure 1). MSB was tested under dim red light at least one hour into the dark phase of a light/dark cycle. All males were habituated at least 30 min before the introduction of the stimulus female in a 17.8 ⬚ 25.4 ⬚ 17.8 cm clear Plexiglas testing cage containing the male's home cage bedding, which had not been changed for at least 48 hours. The test began with the introduction of a hormone-treated stimulus female and videotaped for more than 180 minutes. The videos were scored by an observer for the presence of ejaculatory reflexes. Due to the change in paradigm from COVID-19, the mice experienced long-term social isolation, and this may have played a significant role in the absence of MSB in the males that previously demonstrated MSB on the 2nd test after castration. Thus, upon returning to the lab at 37 weeks post-castration, each experimental male and stimulus female received social exposure in an attempt to determine if SIMSB could be reinstated after this extended social isolation. Details of the social exposure have been provided previously (Wersinger & Rissman, 2000). In brief, each experimental male and stimulus female received social exposure with a gonad-intact male and female mouse for three consecutive days before MSB. During each exposure, each social stimulus was individually placed in the subject's home cage for two min. After 11 MSB tests, males were considered as SI persistent maters if they demonstrated ejaculatory reflexes over 40% of all 11 tests, plus over 40% of the last five tests. Males were considered SI non-maters if they never demonstrated any MSB over the 11 tests.

Cell isolation

During the last MSB test, one of the six mice that were considered SI-maters did not exhibit

ejaculatory reflexes on the last test. This difference within the group would confound our molecular

results since we are investigating the sexual phenotype and not ejaculatory reflexes. Therefore, to

minimize this potential confounder, we collected brain samples two weeks after the last MSB test. All

SI non-maters never exhibited any ejaculatory reflexes post-castration.

Cells were isolated using the Adult Brian Dissociation kit (Miltenyi Biotec, cat # 130-107-677)

following the manufacturer's instructions with several modifications, based on recommendations

(Jaeger *et al.*, 2020) and from optimization. Given that only four mice could be tractably handled each

day, samples were pseudorandomly allocated on different days to include a mix of SI persistent maters

and SI non-maters, as well as one extra B6D2F1 hybrid male mouse to provide extra cells for flow

cytometry technical controls. Due to the small volume of the target brain region (POA), kidneys were

used as carrier cells to improve cell retention during the dissociation. We chose kidneys as it is an

easily dissociable organ, and does not substantially contaminate samples with cells expressing markers

of nervous system cells, which is common for other easily dissociable organs such as the prostate and

the spleen (Pislyagin *et al.*, 2017; Haga *et al.*, 2019). Mice ($n = 16$) were rapidly decapitated, and then

the brains and kidneys were rapidly extracted and fully immersed into ice-cold Dulbecco's PBS

(Thermo Fisher Scientific, cat# 21600010; free of Mg^{2+} & Ca^{2+}; filter-sterilized with 0.22μm filter).

The POA was isolated using a brain matrix (Kent Scientific, cat # RBMA-200C) and surgical blades,

immersed in ice-cold Dulbecco's PBS, under a dissection microscope. The POA sample was topped up

to 400 – 500 mg total sample weight with portions of the kidney. Since any sample that is >400mg

provides a sufficient bulk for processing the samples, the amount of kidney added to the POA would

not matter. Additionally, the single-cell resolution provided by flow cytometry meant that cells added

to the bulk sample would not affect the results, since flow cytometry could gate just the specific brain

cells for analysis.

Alongside with kidney, the POA was finely minced in enzyme mix 1. After adding enzyme mix

2, the mixture was incubated in a hybridization chamber (Fisher Scientific, # 13-247-10) with

maximum rotation (20min, $35^{\circ}C$). Ensuing that, the mixture was gently triturated with a 20G needle

(Becton Dickinson, cat # 305175) and then further incubated (40min, 35^{o}C). Enzymatic digestion was

stopped by topping up to 15mL with DMEM (Gibco, cat# 12800017; added 3.7g/L sodium bicarbonate;

adjusted to pH 7.2 – 7.4; filter-sterilized with 0.22μm filter) and added 0.1mg/mL ovomucoid protease

inhibitor (Sigma, cat# T2011). The mixture was centrifuged (300 ⮾ g, 8min, 4^{o}C) (Thermo Scientific,

cat # 50136519) and then filtered through a 70 μm cell strainer (Fisher Scientific, cat # 352350) into a

fresh 50mL conical tube, topped up to 20mL with DMEM. After centrifugation (300 ⮾ g, 8 min, 4^{o}C),

we carried out the debris removal procedures according to the kit. The suspension was centrifuged with

slow acceleration and deceleration rates (900 ⮾ g, 15 min, 4^{o}C), and then the supernatant was aspirated.

Subsequent to that, the cell pellets were washed by topping up to 15mL with ice-cold Dulbecco's PBS

and then centrifugation (300 ⮾ g, 10 min, 4^{o}C) with full acceleration and brake. Following that, red

blood cells were removed according to the kit with the incubation step done using the mini rotator

(speed 18, 4^{o}C) (Fisher Scientific, cat # 88861052). To get a sense of the cell viability after these steps,

cells were stained with trypan blue (Gibco, cat # 15250061), coverslipped onto a haemocytometer

(Hausser Scientific, cat # 02-671-51B), and visualized under a fluorescent cell imager (Bio-Rad, cat #

1450031).

Staining

Cells were resuspended to a concentration of ~4 – 6 ⮾ 10^{6} cells per ml and stained with 1μL of

fixable viability dye, Zombie Aqua (Biolegend; cat# 423101), then incubated in the dark with rotation

(room temperature, 30 minutes). The samples were washed with 1mL DMEM, centrifuged (300 ⮾ g, 8

min, room temperature) and aspirated until 100 μL remained. Subsequently, the cells were fixed using

the Foxp3 / transcription factor staining buffer kit (eBioscience, cat # 00-5521-00) according to the

manufacturer's protocol with several modifications based on the recommendation (Reichard &

Asosingh, 2019) and optimization. After fixation, the samples were centrifuged (600 ⮾ g, 5 min, room

temperature), supernatant aspirated, and then resuspended into 1:1 glycerol (Millipore, cat # 4750-1L):

Dulbecco's PBS (~1 – 2 ⮾ 10^{6} cells per ml). The samples were stored at -20 ^{o}C in the dark until

permeabilization within two weeks. Before permeabilization, the samples were diluted 10-fold into

autoclaved distilled H$_2$O and centrifuged (600 ⨯ g, 5 min, room temperature) followed by

permeabilization according to the manufacturer's instruction. The samples were then divided to include

technical controls of unstained control, compensation controls, crystallizable fragment receptor

blocking control, and fluorescence minus one control, as well as two subsamples from each sample to

accommodate two different antibody panels, with each sample and control resuspended into 100μL of

permeabilization buffer. Samples and relevant controls were added with 2 μL of crystallizable fragment

receptor blocker (Miltenyi Biotec, cat # 130-092-575) and incubated in the dark with rotation (12min,

4^{o}C). The cells were then incubated with the intracellular primary antibody, rabbit anti-Ionized

calcium-binding adaptor molecule 1 (Iba1; Fujifilm Wako; cat# 019-19741; RRID: AB_839504;

polyclonal; 1:100; 0.5 mg/mL) in the dark with rotation (30min, room temperature). Following that, the

cells were washed thrice with 1mL Dulbecco's DPBS and centrifugation (600 ⨯ g, 5 min, room

temperature)(Eppendorf, cat # 5424R). Afterwards, the samples and relevant controls were incubated

with the secondary antibody, Alexa Fluor 568-conjugated goat anti-rabbit IgG (H+L) (IgG-AF568;

Invitrogen; cat# A-11011; RRID:AB_143157; polyclonal; 1:100; 2mg/mL) and the following directly

conjugated antibodies: fluorescein isothiocyanate-conjugated rabbit anti-NeuN (NeuN-FITC; Abcam;

cat# ab223994; RRID:AB_2892735; monoclonal EPR12763; 1:100; 0.5mg/mL); phycoerythrin-

conjugated mouse anti-androgen receptor (AR-PE; R&D Systems; cat# IC5876P; RRID:AB_10890613;

monoclonal 523339; 1:10); allophycocyanin-conjugated mouse anti-oestrogen receptor alpha (ERα-

APC; Novus Biologicals; cat# NB300-560; RRID:AB_10001032; monoclonal 33; 1:50; 1mg/mL);

phycoerythrin-conjugated rabbit anti-amyloid precursor protein (APP-PE; Abcam; cat# ab208744;

RRID:AB_2892736; monoclonal Y188; 1:1000; 0.5mg/mL); and allophycocyanin-conjugated mouse

anti-glucocorticoid receptor (GR-APC; Invitrogen; cat# 17-6189-82; RRID:AB_2811764; monoclonal

BuGR2; 1:200; 0.2mg/mL). These antibodies were selected based on the intended instrument, its

spectral signatures, the antigen density to match fluorophore brightness, and available fluorophores for

that antibody (Ferrer-Font *et al.*, 2020). Supplementary to these, on the basis of prevalent issues of

antibodies in terms of specificity, reproducibility and selectivity (Hsu *et al.*, 2020), we also selected

antibodies from reputable vendors that validated the antibody and cited by existing studies. Anti-Iba1

was validated using mouse retinal whole-mount and cited in an erythro-myeloid progenitors study

(Mass *et al.*, 2017). Anti-rabbit IgG-AF568 was validated using rat cerebellum and cited in a wind

navigation study (Okubo *et al.*, 2020). Anti-NeuN-FITC was validated with flow cytometry using PC12

cells and cited in a viral encephalitis study (Michael *et al.*, 2020). Anti-AR-PE was validated with flow

cytometry using LNCaP cells and cited in a cortical thymic epithelial cell study (Rode & Boehm, 2012).

Anti-ERα-APC was validated using human breast tissue and cited in a follicular atresia study (Yao *et*

al., 2020). Anti-APP-PE was validated with flow cytometry using SH-SY5Y cells and cited in an

adrenergic receptor study (Chena *et al.*, 2014). Anti-GR-APC was validated with flow cytometry using

mouse splenocytes and cited in an endosomolytic peptide study (Akishiba *et al.*, 2017). After staining,

samples were subjected to procedures in the Foxp3 / transcription factor staining buffer kit according to

the manufacturer's protocol and then resuspended into Dulbecco's PBS for flow cytometry analysis.

Flow cytometry

The fluorescently stained populations (described above) were then analysed using a BD

FACSAria Fusion with a 70 μm nozzle. Data were collected using BD FACSDiva software version 6.0

(http://www.bdbiosciences.com/instruments/software/facsdiva/index.jsp; RRID: SCR_001456). A

compensation matrix was developed using singly stained and unstained controls of harvested cells.

Provisional gates were made using forward and side scatter of light to eliminate debris and non-singlet

cells. Dead cells were then excluded using Zombie Aqua as well as Iba1 and NeuN as secondary

markers to identify microglial and neuronal populations, respectively. AR, APP, GR and ERα

expressions were then analysed. Unstained and fluorescence minus one controls were used for the

correct placement of gates. As a benchmark for storage events, more than 30,000 NeuN+ neurons were

acquired.

Flow cytometry bioinformatics

Data were analysed using FlowJo software version 10.6.1

(https://www.flowjo.com/solutions/flowjo; RRID: SCR_008520) following the recommended

bioinformatics pipeline (Wang & Brinkman, 2019). The compensated data were quality assessed, and t-

transformed using the Logicle function. Subsequently, gating was carried out as described above using

contour plots with outliers that show axis labels stating the marker and fluorochrome used, including

numbers along the axes. A figure exemplifying the sequential flow cytometry gating strategy can be

found in Supplementary Figure 1. Data on AR, APP, GR, and ERα were analysed from the contour plot

showing the percentage of neurons or microglia, and from the histogram showing the distribution of the

number of neurons or microglia at each fluorescence intensity. The percentage of cells was normalised

by dividing the values from a SI persistent mater with a pseudorandomized paired SI non-mater.

Pseudorandomization was carried out by listing out all the sample codes for each of the SI persistent

maters and SI non-maters in order from the smallest to the largest number. Then, using

https://www.random.org/lists/, the sample codes for SI non-maters were pasted to generate the sample

codes in random order. This random order was used to pair with the sample codes of SI persistent

maters (Supplementary Table 1).

In conjunction with the percentage of cells, we needed a measure of expression intensity. One

concern was that flow cytometry fluorescence data were log-normally distributed, which meant that it

would have a long-skewed tail that confounds the mean. On the basis of this, we selected the median

fluorescent intensity (MdFI) for the measure of expression (O'Neill *et al.*, 2013). The MdFI of SI

persistent maters were normalized with the pseudorandomized paired SI non-maters (Supplementary

Table 1) by the resolution metric. The resolution metric is commonly adopted in flow cytometry

experiments (Romano *et al.*, 2021; Soh *et al.*, 2021; Özgün & Garipcan, 2021; Ditzig *et al.*, 2022),

because it is able to account for various sources of noise in flow cytometry data (Bhowmick *et al.*,

2020). In our experiment, the resolution metric is defined as $(\text{MdFI}_{\text{SI persistent maters}} - \text{MdFI}_{\text{SI non-}}$

$_{\text{maters}})/(\text{rSD}_{\text{SI persistent maters}} + \square\text{rSD}_{\text{SI non-maters}})$. A higher resolution metric indicates a better separation

between the two groups. This takes a similar approach to the ddCT method, which is widely used in

qPCR data analysis (Flatschacher *et al.*, 2022).

Preadsorption controls to validate antibodies

Cells were isolated in the same way described in "Cell isolation", except the samples were from

the forebrains of C57BL/6J male mice, without the need for carrier cells. After isolation, cells were

stained in the same way described in "Staining" until after the samples were blocked. Each antibody

was incubated with fivefold the amount of the respective recombinant protein antigen for one hour at

room temperature, and then proceeded with the staining protocol described in "Staining". Each

antibody was single-stained with or without the respective recombinant protein antigen (N=1 per

antibody-antigen combination since this is a validation experiment) (Supplementary Table 2). Flow

cytometry was used to collect data as described in "Flow cytometry", and antibody signals in cells were

analysed after dead cells were excluded. Unstained controls were used for the correct placement of

gates. As a benchmark for storage events, more than 50,000 cells were acquired. Data were analysed

according to "Flow cytometry bioinformatics" except no normalization to the percentage of cells and

MdFI were made since the N=1 for antibody validation did not require statistical analyses.

Statistical analyses

No statistical methods were used to predetermine sample sizes, but the sample size for flow

cytometry experiments are similar to those generally employed in the field (Esk *et al.*, 2020; Farmen *et*

al., 2021; Sanmarco *et al.*, 2021), and sample size for sexual behaviour testing was based on our

previous studies (McInnis *et al.*, 2016). Biologically independent samples represent six SI persistent

maters, and six SI non-maters were used for flow cytometry. Given that there were more than six SI

persistent maters and SI non-maters in our mice cohort, we randomly selected six SI persistent maters

and six SI non-maters for flow cytometry. Experiments were performed blinded during the annotation

of MSB videos to the history of the ejaculatory reflexes of each mouse. Experiments were also blinded

to mater status during flow cytometry sample preparation, flow cytometry and bioinformatics analyses.

All statistical analyses were performed using SPSS version 28 (https://www.ibm.com/products/spss-

statistics; RRID: SCR_019096) and graphs were made with GraphPad Prism version 9.0.0

(http://www.graphpad.com/; RRID: SCR_002798). Formal outlier tests were conducted using Grubb's

test to determine outlying data points, then excluded from the analysis. The data excluded from the

analysis is reflected in the changes in the sample size in each result from the original six per group. The

final sample size used for each analysis is shown in the figure legends. The outliers from the raw data

are provided for reference (Supplementary Table 3), but due to the design of our data analysis, outliers

were only excluded after the data were processed as above. Most of the data fulfilled the assumption of

Normality based on the Shapiro-Wilks test, without multivariate outliers based on the Mahalanobis

distance, but do not fulfil the assumption of the homogeneity of variance based on Levene's test.

Therefore, we proceeded with the non-parametric test, the Mann-Whitney U test to compare SI

persistent maters and SI non-maters with the threshold for significance set at $p < 0.05$. Statistical details

for specific analysis, including exact n values and what n represents, statistical tests used, central

tendency measure, U values, P values, effect sizes, and all plotted data points, can be found in the

figure legends.

To give a sense of the variability of the raw data for the immunoexpression of all molecules

measured, we obtained the standard deviation for each molecule (APP, AR, GR, ERa) in neurons and

microglia (total N = 8), and used Student's T-test to compare between SI persistent maters and SI non-

maters with the threshold for significance set at $p < 0.05$. Since the same animal samples were used for

different molecular analyses, we also ran correlation analyses of the immunoexpression of each

molecule in neurons and microglia, in terms of how well they correlated with the immunoexpression of

all other molecules.

Results

B6D2F1 male mice exhibit male sexual behaviour 62 weeks after orchidectomy

At five weeks post-orchidectomy, 50% of the B6D2F1 exhibited ejaculatory reflexes (Figure 1A). Following that, due to COVID-19, we had to suspend any lab work for an extended period. Upon returning to the lab at 28 weeks post-orchidectomy, none of the B6D2F1 exhibited ejaculatory reflexes (Figure 1A). We speculated this could be due to extended social isolation with the absence of physical exposure to other mice, minimal human handling, or ageing (Liu *et al.*, 2019; Cho *et al.*, 2021; Sokolov *et al.*, 2021). In accordance with our values in animal ethics to reduce the waste of animals (reviewed in (Smith, 2020)), we attempted to salvage this experiment. Social isolation is known to adversely affect MSB (Liu *et al.*, 2019), therefore we provided the experimental and stimulus mice with multiple social exposures prior to each sexual behaviour test after 0% of the males showed SIMSB. The repeated sexual behaviour testing and social exposure correlated with an increase in the percentage of males exhibiting ejaculatory reflexes back to levels quasi to previous studies with 28.13% at 43 weeks post-orchidectomy and 25% at 62 weeks post-orchidectomy (Figure 1A). Previous studies unearthed two distinct sexual phenotypes in these post-orchiectomized B6D2F1 males, namely, the SI maters that consistently exhibit MSB and SI non-maters that do not (Park *et al.*, 2009; Bharadwaj *et al.*, 2013; Templin *et al.*, 2019). In general, these studies defined SI maters as those who demonstrated ejaculatory reflex on at least three out of the last four tests, including during the last test, and SI non-maters as those not demonstrating ejaculatory reflex on the last four tests (Park *et al.*, 2009; Bharadwaj *et al.*, 2013; Templin *et al.*, 2019). Given changes to our paradigm due to COVID-19, our current cohort is distinctly different to previous iterations, and could not utilize the same definition from previous studies. To incorporate the SIMSB reinstatement aspect of this study paradigm, we have labelled the mice that exhibited ejaculatory reflex at least 40% across all 11 tests and at least 40% in the last 5 tests as 'SI persistent maters' (seven out of 34 mice were SI persistent maters, 20.59%). The definition of SI non-maters was the same as previous studies, which are those that never exhibited ejaculatory reflexes across all 11 tests (eight out of 34 mice were SI non-maters, 23.53%). We ran

statistical analysis on the number of ejaculatory reflexes on the SI persistent maters and SI non-maters

we chose for downstream molecular analysis. Our statistical analysis showed SI persistent maters

exhibited increased SIMSB compared to SI non-maters (Median ejaculatory reflex: SI persistent maters

= 7; SI non-maters = 0. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent mater,

$n = 6$ mice; SI non-mater, $n = 6$ mice, p-value = 0.002, U-value = 0.000, and effect size = 0.892)

(Figure 1B).

Steroid-independent persistent maters exhibited altered amyloid precursor protein expression

We next investigated how SIMSB may be regulated at the molecular level in the POA using

flow cytometry. Our first target was APP since a previous study found increased APP in SI persistent

maters compared to SI non-maters (Park *et al.*, 2010). Based on the APP MdFI from the flow

cytometry histogram of the SI persistent maters, which was normalized to the SI non-maters using the

resolution metrics, we found a decrease in APP immunoexpression in POA neurons (Median MdFI

resolution metrics: SI persistent maters neuron = -0.0232; SI non-maters neuron = 0.000. Mann-

Whitney U nonparametric statistical test, two-tailed, SI persistent maters neuron, $n = 5$ mice; SI non-

maters neuron, $n = 5$ mice, p-value = 0.003, U-value = 0.0000, and effect size = 0.9003) (Figure 2A,

2C). There were no changes to the percentage of neurons expressing APP from the flow cytometry

contour plot of SI persistent maters, which was normalized to the SI non-maters (Median normalized

percentage of cells: SI persistent maters neuron = 0.9113; SI non-maters neuron = 1.000. Mann-

Whitney U nonparametric statistical test, two-tailed, SI persistent maters neuron, $n = 6$ mice; SI non-

maters neuron, $n = 6$ mice, p-value = 0.305, U-value = 12.000, and effect size = 0.2962.) (Figure 2B,

2D). This indicates that existing neurons expressing APP reduced APP immunoexpression.

There is a paucity of studies investigating non-neuronal cells in the brain in relation to sexual

behaviour, therefore, we also investigated APP immunoexpression in Iba1+ microglia. We found an

increase in APP expression in Iba1+ microglia of SI persistent maters compared to SI non-maters

(Median MdFI resolution metrics: SI persistent maters microglia = 0.0124; SI non-maters microglia =

0.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent maters microglia, n = 6

mice; SI non-maters microglia, n = 6 mice, p-value = 0.04, U-value = 6.000, and effect size = 0.5921)

(Figure 2A, 2C). There were no changes in the percentage of APP-expressing microglia in SI persistent

maters compared to SI non-maters (Median normalized percentage of cells: SI persistent maters

microglia = 0.9072; SI non-maters microglia = 0.000. Mann-Whitney U nonparametric statistical test,

two-tailed, SI persistent maters microglia, n = 6 mice; SI non-maters microglia, n = 6 mice, p-value =

1.000, U-value = 18.000, and effect size = 0.000). This indicates that existing microglia expressing

APP increased APP immunoexpression.

We ran additional experiments using preadsorption controls as preliminary efforts to help

validate the antibodies. All the antibodies were incubated with their respective recombinant protein

antigens. For the percentage of cells in preadsorption controls, we found 1.4% NeuN+ cells, and 50.2%

with only antibody; 96.4% Iba1+ cells, and 99.5% with only antibody; 98.8% APP+ cells, and 98.9%

with only antibody (Supplementary Figure 2; Supplementary Table 4). In terms of the MdFI of the

preadsorption controls, these were 68.1 for NeuN, and 236 with only antibody; 88.7 for Iba1, and 232

with only antibody; 107 for APP, and 140 with only antibody (Supplementary Figure 2;

Supplementary Table 4). Due to the design of our normalization method, our result graphs did not

display the level of variability in the raw data. Therefore, we obtained the standard deviation for all of

the molecules measured (AR, APP, GR and ERa), and compared them between the SI persistent maters

and SI non-maters, to determine if there were differences in variability. There were no differences in

the standard deviation of the percentage of cells between SI persistent maters compared to SI non-

maters (Mean standard deviation for the percentage of expressing cells: SI persistent maters = 3.178; SI

non-maters = 4.397. Student's T statistical test, two-tailed, SI persistent maters, n = 8 molecules in

neurons and microglia; SI non-maters, n = 8 molecules in neurons and microglia, p-value =

0.238)(Supplementary Figures 3). For MdFI, there were no differences in the standard deviation of the

MdFI between SI persistent maters compared to SI non-maters (Mean standard deviation for MdFI: SI

persistent maters = 67.512; SI non-maters = 68.027. Student's T statistical test, two-tailed, SI persistent

maters, n = 8 molecules in neurons and microglia; SI non-maters, n = 8 molecules in neurons and

microglia, p-value = 0.990)(Supplementary Figures 3).

Steroid-independent persistent maters exhibit decreased androgen receptor expression in neurons

The sex steroid receptors, AR and ERa are the cognate receptors of sex steroid hormones, which are paramount for canonical sexual behaviour (reviewed in (Chiang & Park, 2020)). Thus, we compared AR and ERa expression between SI persistent maters and SI non-maters. We discovered a reduced AR expression in NeuN+ neurons of SI persistent maters compared to SI non-maters (Median MdFI resolution metrics: SI persistent maters neuron = -0.0115; SI non-maters neuron = 0.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent maters neuron, n = 6 mice; SI non-maters neuron, n = 6 mice, p-value = 0.04, U-value = 6.0000, and effect size = 0.5921) (Figure 3A, C). In addition, we discovered a reduction in the percentage of AR-expressing neuron in SI persistent maters compared to SI non-maters (Median normalized percentage of cells: SI persistent maters neuron = 0.9602; SI non-maters neuron = 1.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent maters neuron, n = 6 mice; SI non-maters neuron, n = 6 mice, p-value = 0.04, U-value = 6.000, and effect size = 0.5921) (Figure 3B, D). This indicates that there was an overall reduction of AR expression in SI persistent maters neurons, and this could be related to the reduced neuronal population expressing AR. In terms of non-neuronal cells, Iba1+ microglia did not exhibit differential AR expression in SI persistent maters compared to SI non-maters (Median MdFI resolution metrics: SI persistent maters microglia = -0.007; SI non-maters microglia = 0.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent maters microglia, *n* = 6 mice; SI non-maters microglia, *n* = 6 mice, *p*-value = 1.000, *U*-value = 18.000, and effect size = 0.000) (Figure 3A, C). In tandem with this, there were no changes in the percentage of Iba1+ neurons expressing AR in SI persistent maters compared to SI non-maters (Median normalized percentage of cells: SI persistent maters microglia = 0.9737; SI non-maters microglia = 0.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent maters microglia, *n* = 6 mice; SI non-maters microglia, *n* = 6 mice, *p*-value = 0.305, *U*-value = 12.000, and effect size = 0.2962) (Figure 3B, D). By the same token, we examined ERa, but there were no differences detected between SI persistent maters and SI non-maters (Supplementary Figure 2). For the percentage of cells in preadsorption controls, we found

86.6% AR+ cells, and 19.4% with only antibody; 39.9% ERa cells, and 12.9% with only antibody

(Supplementary Figure 2; Supplementary Table 4). In terms of the MdFI of the preadsorption controls,

these were 68.1 for AR, and 11.6 with only antibody; 585 for ERa, and 293 with only antibody

(Supplementary Figure 2; Supplementary Table 4).

Steroid-independent persistent maters exhibit increased glucocorticoid receptor expression in

neurons

Finally, we investigated the cell heterogeneous expression of GR, due to its homology with ERa

and AR (reviewed in (Spaanderman *et al.*, 2018). Furthermore, glucocorticoid acts through GR, leading

to various cellular changes including increasing the transcription of glucocorticoid-induced transcript 1

(Kiuchi *et al.*, 2019), which the previous SIMSB microarray study found were down-regulated in SI

maters compared to SI non-maters (Park *et al.*, 2010). On top of these, GR is known to regulate sexual

behaviour in the context of stress (reviewed in (Ralph *et al.*, 2016). We unearthed an increased GR

expression in NeuN+ neurons of SI persistent maters compared to SI non-maters (Median MdFI

resolution metrics: SI persistent maters neuron = 0.0698; SI non-maters neuron = 0.000. Mann-Whitney

U nonparametric statistical test, two-tailed, SI persistent maters neuron, n = 6 mice; SI non-maters

neuron, n = 6 mice, p-value = 0.002, U-value = 0.000, and effect size = 0.8883) (Figure 4A, C). At the

same time, there were no changes in the percentage of GR-expressing neurons in SI persistent maters

compared to SI non-maters (Median normalized percentage of cells: SI persistent maters neuron =

0.8824; SI non-maters neuron = 1.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI

persistent maters neuron, n = 5 mice; SI non-maters neuron, n = 5 mice, p-value = 0.550, U-value =

12.000, and effect size = 0.18). What this means is the existing neurons acquired higher GR expression.

Apropos to comparing SI persistent maters to SI non-maters microglia in the POA, GR expression was

unchanged in terms of expression level (Median MdFI resolution metrics: SI persistent maters

microglia = 0.0015; SI non-maters microglia = 0.000. Mann-Whitney U nonparametric statistical test,

two-tailed, SI persistent maters microglia, *n* = 6 mice; SI non-maters microglia, *n* = 6 mice, *p*-value =

0.305, *U*-value = 12.000, and effect size = 0.2962), and percentage of GR-expressing microglia

(Median normalized percentage of cells: SI persistent maters microglia = 0.9441; SI non-maters

microglia = 0.000. Mann-Whitney U nonparametric statistical test, two-tailed, SI persistent maters

microglia, n = 5 mice; SI non-maters microglia, n = 5 mice, p-value = 0.073, U-value = 6.000, and

effect size = 0.54030).

For the percentage of cells in preadsorption controls, we found 1.49% GR+ cells, and 1.53%

with only the antibody (Supplementary Figure 2; Supplementary Table 4). In terms of the MdFI of the

preadsorption controls, these were 585 for GR, and 293 with only the antibody. Aside from the

immunoexpression itself, we wanted to see if the immunoexpression of any of the molecules in either

the microglia or neurons correlated with other molecules. To do so, we ran correlation analyses for the

MdFI, but we did not observe any highly positive (correlation coefficient >0.8) or highly negative

(correlation coefficient < -0.8) correlations between different molecules (Supplementary Figure 5).

Discussion

We show for the first time differential immunoexpression profiles in a cell-specific manner between castrated B6D2F1 hybrid male mice that demonstrate SIMSB relative to those that ceased. The paradigm of this study differs from previous studies on SIMSB of B6D2F1 hybrid male mice due to a COVID-19 laboratory shutdown which led mice to be singly housed and socially isolated for 23 weeks. However, SIMSB was reinstated and maintained in 20.59% of the mice after repeated social exposure. In terms of neurons, SI persistent maters had down-regulated APP and AR immunoexpression as well as up-regulated GR immunoexpression compared to SI non-maters. For microglia, SI persistent maters demonstrated increased APP immunoexpression compared to SI non-maters.

Our mice model illustrated MSB post-orchidectomy, but the 20.59% of SI persistent maters is less than the one-third observed in previous studies (reviewed in (Chiang & Park, 2020)). The social isolation in our study may not explain this difference fully because a previous study with 12 weeks of social isolation immediately after castration still resulted in 38% of the hybrid males demonstrating SIMSB (Wee & Clemens, 1989). Several differences exist between our study and this previous study. Namely, we had an extended 23 weeks of social isolation, extended testing for 62 weeks post-orchidectomy, and social exposure introduced. On top of that, one notable difference was that the mice in our study were 17 – 18 months old on their last MSB test, which is older than the traditional breeding age (Moreira *et al.*, 2015). Comparatively, the mice in previous studies were around 4 – 6 months of age on their last MSB test (David, Wyrosdic, & Park, 2022; David, Wyrosdic, Wan, *et al.*, 2022). This is a potential confound as ageing is known to impair MSB (Wilson *et al.*, 2018; Jesús *et al.*, 2022). Future studies could independently test these variables (i.e social isolation, social exposure, and ageing) to determine how each may contribute to the altered percentage of SI persistent maters.

Steroid hormone receptors are pivotal for the development of neurocircuits essential for MSB (Karigo & Deutsch, 2022). We found down-regulated AR and up-regulated GR immunoexpression in POA neurons of SI persistent maters compared to SI non-maters. These results are discrepant with the previous microarray study finding no differences in *Ar* and *Gr* expression between these groups (Park

et al., 2010), and that AR blockers were ineffective to abate SIMSB (Park *et al.*, 2009). These

differences are not surprising, as our approach was cell-specific, as opposed to the bulk approach in

previous studies. However, one potential confound to note was the time of brain sample collection

because how long the brain was collected after the last MSB test could affect the measured molecular

profile. This is because one out of the six SI persistent maters we chose for molecular analysis, did not

display ejaculatory reflexes in the last MSB test. Therefore, in our experiments, brain samples were

collected two weeks after the last MSB test, to minimize the confound from the ejaculatory reflexes.

This was not a problem for SI non-maters, as they never presented any SIMSB post-castration.

There may also be issues with not immediately collecting brain samples after the last MSB test.

Since one of the six SI persistent maters did not display ejaculatory reflexes in the last MSB test, there

is a possibility that during the two-week interval, if MSB tests were conducted, we may find some mice

may no longer be considered as SI persistent maters under the current definition. The reason why this

quandary exists is related to how different studies have varying standards on which mice were

considered SI maters. Instead of an easy demarcation of SI maters and SI non-maters, there appears to

be a continuum of SI 'intermediate' maters with varying degrees of SIMSB shown across cohorts of

B6D2F1 hybrid male mice that are neither SI maters nor SI non-maters (Park *et al.*, 2009). Future

studies would need to think of robust ways to characterise and demarcate the SIMSB phenotypic

categories. One possible way could be to obtain more sexual behaviour measures to be taken into

account for the categorisation, instead of just the ejaculatory reflexes (Heijkoop *et al.*, 2018). SIMSB is

well-established to differ from the MSB of gonad-intact control without orchidectomy (McGill &

Tucker, 1964), but MSB of gonad-intact mice would similarly benefit from this approach to identify

nuances in their sexual phenotypes. Traditionally, obtaining multiple sexual behaviour measures is a

daunting task, but this is now more tractable with machine-learning behaviour methods (Blau *et al.*,

2022; Sun *et al.*, 2022)

Notwithstanding, we believe our comparison was still meaningful since we compared two

sexual phenotypes at two ends of extremities, with one end of the extremity with mice never exhibiting

SIMSB, compared to another end of the extremity with mice that consistently exhibited SIMSB, albeit

not on all MSB tests. Despite the two-week interval to brain sample collection may affect their SI

persistent maters status, mice with slightly less SIMSB exhibited would still provide insight into the

molecular profiles relevant to SIMSB, though possibly with a weaker signal.

During development, complementary to neuronal roles to shape sexual behaviour, non-neuronal

cells drive spinogenesis essential for sexual behaviour by increasing the production of prostaglandin E2

(Lenz *et al.*, 2013, 2018). For microglia, we found up-regulated APP in SI persistent maters compared

to SI non-maters, which is in concordance with the previous study that found APP up-regulation (Park

et al., 2010). However, for neurons, APP was down-regulated. If APP is indeed up-regulated at the

global level, what we observed could be that the microglial APP up-regulation may compensate for any

down-regulation of neuronal APP. To obtain a more complete picture, future studies can use a multi-

method approach to measure APP. This would include repeating the bulk transcriptomics approach for

our reinstatement paradigm. Our current study using flow cytometry allowed cell-specific assessment

of protein immunoexpression, but future studies can ascertain these results through fluorescence-

activated cell sorting followed by western blot on the sorted neurons and microglia. Furthermore, gene

expression can be explored using qPCR on the sorted neurons and microglia to validate the congruence

between the transcriptional and translational levels.

From these findings and supporting studies from the literature, we composed a putative

signalling pathway for SIMSB (Figure 5). Our targets APP, AR, and GR interact with molecules

related to sexual behaviour, namely BDNF (MacPherson, 2017; Brague *et al.*, 2018), ERK (Jean *et al.*,

2017; Kirouac *et al.*, 2017), CREB (Barrot *et al.*, 2005; Sun *et al.*, 2021), nitric oxide (Melis, 1998;

Will *et al.*, 2014; Gu *et al.*, 2020), glutamate (Numakawa *et al.*, 2009), and dopamine 1 receptor (Melis

et al., 2003; Will *et al.*, 2014; Wu *et al.*, 2016).They also interact with each other such that AR binds to

APP (Takayama *et al.*, 2019), AR increases GR transcription (Spaanderman *et al.*, 2018) and GR

induce secretase processing of APP (Baglietto-Vargas *et al.*, 2013). Our correlational analyses did not

support this putative signalling pathway, which was expected due to the experimental design being

cross-sectional and complicated by the reinstatement of SIMSB. To verify our putative signalling

pathway, future *in silico* studies are needed as well as empirical evidence from high-throughput omics

that could simultaneously measure millions of molecules at the single-cell resolution. This should

include the comparison between the intact control with SIMSB phenotypes, to decipher which

molecular changes occurred post-gonadectomy in a mating-dependent manner.

Several limitations should be addressed in our study which future studies may help address. The

first pertains to the prevalent issues of the specificity of antibodies (Hsu *et al.*, 2020). For Iba1, GR, and

APP, we found no differences in the signals between the antibody-only sample and the preadsorption

control. This may indicate these antibodies did not actually bind to the immunogens reported by the

vendors. While this may falsify the tested immunogen as the antibody target, our results may still be

valid, as it does not eliminate the possibility that other immunogens of the same protein bind to the

antibody as well (Holmseth *et al.*, 2012; Brownjohn & Ashton, 2014). For ERa and AR, the

preadsoprtion control actually exhibited more signal than the antibody-only sample. We suspect for

these specific antibodies, there may be heavy complexes formed between the antibody and immunogen,

that could not be washed away in our protocol following the commonly cited protocol from Synaptic

Systems and R&D Systems. Future studies may need to optimize the protocol for each specific

antibody and conduct additional antibody validation experiments to ascertain the validity of results

generated from these antibodies. This should be a concerted effort with vendors and a consortium of

multiple laboratories.

Another limitation was the normalization method we used. This normalization method has been

used in multiple flow cytometry studies (Bhowmick *et al.*, 2020; Romano *et al.*, 2021; Soh *et al.*, 2021;

Ditzig *et al.*, 2022), and follows a similar logic to the ddCt method used for qPCR (Pabinger *et al.*,

2014). In this way, each pair of the comparison group and the control group would combine into only

one data point to reflect only their difference. Some argue that this wipes away the variability, but

some think that the variability is still reflected in the one data point when the comparison and control

group is combined (Pabinger *et al.*, 2014). Despite this, we recognize this could be a limitation because

there is no consensus on how flow cytometry analysis should be conducted. There are values in future

consortiums to identify a gold standard for the field to follow, especially with the burgeoning of

bioinformatic techniques that focus on flow cytometry analysis (Cossarizza *et al.*, 2021).

In summary, we report for the first time cell-specific differential immunoexpression of AR, GR and APP in contrasting sexual phenotypes, defined according to whether or not they exhibited gonadal steroid-independent male sexual behaviour over the long term after orchidectomy and social isolation. Overall, this paves the way to disentangle the environmental context, the phenotypic profile, and the molecular pathways associated with this phenomenon.

Acknowledgements: We thank Kaylee Wells of the UMass Boston Flow Cytometry Core for support on the BD FACSAria Fusion. We thank Dr Caroline David for assistance with animal husbandry, Benny Wan for assistance with surgeries, and other Park Lab members for general assistance. We thank Renjie Stephen Hu for their help with figures production.

Author contributions: JHP and VSCC conceived and designed the experiments. VSCC performed the experiments and data analysis with help from SL. CDM supervised the flow cytometry. VSCC created the figures and wrote the manuscript with input from JHP. JHP, CDM and VSCC reviewed and edited the manuscript. JHP supervised the study, provided the resources, and acquired the funding.

Conflict of interest statement: The authors declare no conflict of interests

Abbreviations: AF568, Alexa Fluor 568; APC, allophycocyanin; APP, amyloid precursor protein; AR, androgen receptor; ERa, oestrogen receptor alpha; FITC, fluorescein isothiocyanate; GR, glucocorticoid receptor; Iba1, Ionized calcium-binding adaptor molecule 1; MdFI, median fluorescent intensity; MSB, male sexual behaviour; PE, phycoerythrin; POA, preoptic area; SI, steroid-independent

Data availability

Any data and materials that can be shared will be released via a material transfer agreement. Full access to biological data will be granted to collaborating scientists who are willing to complete a "Data Access

Agreement" and a "Data Access Application form" in accordance with General Data Protection

Regulation.

Additional Information

Supplementary Information: Supplementary material is available at the end of this manuscript.

Correspondence and requests for materials: Should be addressed to VSC.

Peer review information: Thanks in advance to the editorial team and reviewers for their contribution

to the peer review of this work.

References

Addessi, E. & Bourgeois-Gironde, S. (2020) Are the roots of human economic systems shared with non-human primates? *Neuroscience and Biobehavioral Reviews*, **109**, 1–15.

Akishiba, M., Takeuchi, T., Kawaguchi, Y., Sakamoto, K., Yu, H.H., Nakase, I., Takatani-Nakase, T., Madani, F., Gräslund, A., & Futaki, S. (2017) Cytosolic antibody delivery by lipid-sensitive endosomolytic peptide. *Nature Chemistry*, **9**, 751–761.

Baglietto-Vargas, D., Medeiros, R., Martinez-Coria, H., Laferla, F.M., & Green, K.N. (2013) Mifepristone alters amyloid precursor protein processing to preclude amyloid beta and also reduces tau pathology. *Biological Psychiatry*, **74**, 357–366.

Barrot, M., Wallace, D.L., Bolaños, C.A., Graham, D.L., Perrotti, L.I., Neve, R.L., Chambliss, H., Yin, J.C., & Nestler, E.J. (2005) Regulation of anxiety and initiation of sexual behavior by CREB in the nucleus accumbens. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 8357–8362.

Bharadwaj, P., McInnis, C., Madden, A.M.K., Bonthuis, P.J., Zup, S., Rissman, E.F., & Park, J.H. (2013) Increased Dendritic Spine Density and Tau Expression Are Associated with Individual Differences in Steroidal Regulation of Male Sexual Behavior. *PLoS ONE*, **8**.

Bhowmick, D., Sheridan, R.T.C., Bushnell, T.P., & Spalding, K.L. (2020) Practical Guidelines for Optimization and Characterization of the Beckman Coulter CytoFLEX^TM Platform. *Cytometry Part A*, **97**, 800–810.

Blanco-Suárez, E., Caldwell, A.L.M., & Allen, N.J. (2017) Role of astrocyte-synapse interactions in CNS disorders. *The Journal of Physiology*, **595**, 1903–1916.

Blau, A., Gebhardt, C., Bendesky, A., Paninski, L., & Wu, A. (2022) SemiMultiPose: A Semi-supervised Multi-animal Pose Estimation Framework. *arXiv*,.

Brague, J.C., Lenchur, C.N., Hayden, J.M., Davidson, R.H., Corrigan, K., Santini, G.T., & Swann, J.M. (2018) BDNF infusion into the MPN mag is sufficient to restore copulatory behavior in the castrated Syrian hamster. *Hormones and Behavior*, **102**, 69–75.

Brownjohn, P.W. & Ashton, J.C. (2014) What can be concluded from blocking peptide controls? *Appl Immunohistochem Mol Morphol*, **22**, 634.

Chena, Y., Penga, Y., Chea, P., Gannona, M., Liua, Y., Lib, L., Bu, G., Van Groena, T., Jiaod, K., & Wanga, Q. (2014) α2A adrenergic receptor promotes amyloidogenesis through disrupting APP-SorLA interaction. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 17296–17301.

Chiang, V.S.C. & Park, J.H. (2020) Glutamate in Male and Female Sexual Behavior: Receptors, Transporters, and Steroid Independence. *Frontiers in Behavioral Neuroscience*, **14**, 589882.

Cho, C., Chan, C., & Martin, L.J. (2021) Can Male Mice Develop Preference Towards Gentle Stroking by an Experimenter? *Neuroscience*, **464**, 26–32.

Clemens, L.G., Wee, B.E.F., Weaver, D.R., Roy, E.J., Goldman, B.D., & Rakerd, B. (1988) Retention of masculine sexual behavior following castration in male B6D2F1 mice. *Physiology and Behavior*, **42**, 69–76.

Cossarizza, A., Chang, H.D., Radbruch, A., Abrignani, S., Addo, R., Akdis, M., Andrä, I., Andreata, F., Annunziato, F., Arranz, E., Bacher, P., Bari, S., Barnaba, V., Barros-Martins, J., Baumjohann, D., Beccaria, C.G., Bernardo, D., Boardman, D.A., Borger, J., Böttcher, C., Brockmann, L., Burns, M., Busch, D.H., Cameron, G., Cammarata, I., Cassotta, A., Chang, Y., Chirdo, F.G., Christakou, E., Čičin-Šain, L., Cook, L., Corbett, A.J., Cornelis, R., Cosmi, L., Davey, M.S., De Biasi, S., De Simone, G., del Zotto, G., Delacher, M., Di Rosa, F., Santo, J.D., Diefenbach, A., Dong, J., Dörner, T., Dress, R.J., Dutertre, C.A., Eckle, S.B.G., Eede, P., Evrard, M., Falk, C.S., Feuerer, M., Fillatreau, S., Fiz-Lopez, A., Follo, M., Foulds, G.A., Fröbel, J., Gagliani, N., Galletti, G., Gangaev, A., Garbi, N., Garrote, J.A., Geginat, J., Gherardin, N.A., Gibellini, L., Ginhoux, F., Godfrey, D.I., Gruarin, P., Haftmann, C., Hansmann, L., Harpur, C.M., Hayday, A.C., Heine, G., Hernández, D.C., Herrmann, M., Hoelsken, O., Huang, Q., Huber, S., Huber, J.E., Huehn, J., Hundemer, M., Hwang, W.Y.K., Iannacone, M., Ivison, S.M., Jäck, H.M., Jani,

P.K., Keller, B., Kessler, N., Ketelaars, S., Knop, L., Knopf, J., Koay, H.F., Kobow, K., Kriegsmann, K., Kristyanto, H., Krueger, A., Kuehne, J.F., Kunze-Schumacher, H., Kvistborg, P., Kwok, I., Latorre, D., Lenz, D., Levings, M.K., Lino, A.C., Liotta, F., Long, H.M., Lugli, E., MacDonald, K.N., Maggi, L., Maini, M.K., Mair, F., Manta, C., Manz, R.A., Mashreghi, M.F., Mazzoni, A., McCluskey, J., Mei, H.E., Melchers, F., Melzer, S., Mielenz, D., Monin, L., Moretta, L., Multhoff, G., Muñoz, L.E., Muñoz-Ruiz, M., Muscate, F., Natalini, A., Neumann, K., Ng, L.G., Niedobitek, A., Niemz, J., Almeida, L.N., Notarbartolo, S., Ostendorf, L., Pallett, L.J., Patel, A.A., Percin, G.I., Peruzzi, G., Pinti, M., Pockley, A.G., Pracht, K., Prinz, I., Pujol-Autonell, I., Pulvirenti, N., Quatrini, L., Quinn, K.M., Radbruch, H., Rhys, H., Rodrigo, M.B., Romagnani, C., Saggau, C., Sakaguchi, S., Sallusto, F., Sanderink, L., Sandrock, I., Schauer, C., Scheffold, A., Scherer, H.U., Schiemann, M., Schildberg, F.A., Schober, K., Schoen, J., Schuh, W., Schüler, T., Schulz, A.R., Schulz, S., Schulze, J., Simonetti, S., Singh, J., Sitnik, K.M., Stark, R., Starossom, S., Stehle, C., Szelinski, F., Tan, L., Tarnok, A., Tornack, J., Tree, T.I.M., van Beek, J.J.P., van de Veen, W., van Gisbergen, K., Vasco, C., Verheyden, N.A., von Borstel, A., Ward-Hartstonge, K.A., Warnatz, K., Waskow, C., Wiedemann, A., Wilharm, A., Wing, J., Wirz, O., Wittner, J., Yang, J.H.M., & Yang, J. (2021) Guidelines for the use of flow cytometry and cell sorting in immunological studies (third edition). *European Journal of Immunology*, **51**, 2708–3145.

David, C.D., Wyrosdic, B.N., & Park, J.H. (2022) Strain differences in post-castration sexual and aggressive behavior in male mice. *Behavioural Brain Research*, **422**, 113747.

David, C.D., Wyrosdic, B.N., Wan, H.Y., Lapp, H.E., Bartlett, A.A., Yitbarek, S., & Park, J.H. (2022) B6D2F1 mice that retain sexual behavior long term after castration outperform those that cease in the radial arm maze. *Psychoneuroendocrinology*, **136**, 105627.

Ditzig, Z., Wilson, C.M., Salas, J., & Serve, K.M. (2022) Plasminogen Binding and Activation at the Mesothelial Cell Surface Promotes Invasion through a Collagen Matrix. *International Journal of Molecular Sciences*, **23**.

Esk, C., Lindenhofe, D., Haendeler, S., Wester, R.A., Pflug, F., Schroeder, B., Bagley, J.A., Elling, U., Zuber, J., Von Haeseler, A., & Knoblich, J.A. (2020) A human tissue screen identifies a regulator of ER secretion as a brain-size determinant. *Science*, **370**, 935–941.

Farmen, K., Nissen, S.K., Stokholm, M.G., Iranzo, A., Østergaard, K., Serradell, M., Otto, M., Svendsen, K.B., Garrido, A., Vilas, D., Borghammer, P., Santamaria, J., Møller, A., Gaig, C., Brooks, D.J., Tolosa, E., Pavese, N., & Romero-Ramos, M. (2021) Monocyte markers correlate with immune and neuronal brain changes in REM sleep behavior disorder. *Proceedings of the National Academy of Sciences of the United States of America*, **118**.

Ferrer-Font, L., Pellefigues, C., Mayer, J.U., Small, S.J., Jaimes, M.C., & Price, K.M. (2020) Panel Design and Optimization for High-Dimensional Immunophenotyping Assays Using Spectral Flow Cytometry. *Current Protocols in Cytometry*, **92**.

Flatschacher, D., Speckbacher, V., & Zeilinger, S. (2022) qRAT: an R-based stand-alone application for relative expression analysis of RT-qPCR data. *BMC Bioinformatics*, **23**.

Garcia, C., Bouret, S., Druelle, F., & Prat, S. (2021) Balancing costs and benefits in primates: Ecological and palaeoanthropological views. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **376**.

Gu, L., Lian, D., Zheng, Y., Zhou, W., Gu, J., & Liu, X. (2020) Echinacoside□induced nitric oxide production in endothelial cells: Roles of androgen receptor and the PI3K□Akt pathway. *International Journal of Molecular Medicine*, **45**, 1195–1202.

Guerra, S., Chesworth, R., Weickert, C.S., & Karl, T. (2021) Effects of handling on the behavioural phenotype of the neuregulin 1 type III transgenic mouse model for schizophrenia. *Behavioural Brain Research*, **405**.

Haga, N., Akaihata, H., Hata, J., Hiraki, H., Honda, R., Tanji, R., Onagi, A., Koguchi, T., Hoshi, S., Ogawa, S., Kataoka, M., Sato, Y., Ishibashi, K., & Kojima, Y. (2019) The association between local arteriosclerosis of the prostatic arteries and chronic inflammation in human benign prostatic enlargement. *Prostate*, **79**, 574–582.

Heijkoop, R., Huijgens, P.T., & Snoeren, E.M.S. (2018) Assessment of sexual behavior in rats: The potentials and pitfalls. *Behavioural Brain Research*, **352**, 70–80.

Holmseth, S., Dehnes, Y., Huang, Y.H., Follin-Arbelet, V.V., Grutle, N.J., Mylonakou, M.N., Plachez, C., Zhou, Y., Furness, D.N., Bergles, D.E., Lehre, K.P., & Danbolt, N.C. (2012) The Density of EAAC1 (EAAT3) Glutamate Transporters Expressed by Neurons in the Mammalian CNS. *J Neurosci*, **32**, 6000–6013.

Hsu, C.-N., Chang, C.-H., Poopradubsil, T., Lo, A., William, K.A., Lin, K.-W., Bandrowski, A., Ozyurt, I.B., Grethe, J.S., & Martone, M.E. (2020) Antibody Watch: Text Mining Antibody Specificity from the Literature. *PLOS Computational Biology*, **17**, e1008967.

Jaeger, B.N., Yánguez, E., Gesuita, L., Denoth-Lippuner, A., Kruse, M., Karayannis, T., & Jessberger, S. (2020) Miniaturization of Smart-seq2 for Single-Cell and Single-Nucleus RNA Sequencing. *STAR Protocols*, **1**, 100081.

Jean, A., Trouillet, A.C., Andrianarivelo, N.A., Mhaouty-Kodja, S., & Hardin-Pouzet, H. (2017) Phospho-ERK and sex steroids in the mPOA: Involvement in male mouse sexual behaviour. *Journal of Endocrinology*, **233**, 257–267.

Jesús, P.L. de, Arenas-Ríos, E., Ruíz-Ramos, M., Flores-Alonso, J.C., Mendoza-Núñez, V.M., Arrieta-Cruz, I., & Arteaga-Silva, M. (2022) Effect of Chronic Moderate Caloric Restriction on the Reproductive Function in Aged Male Wistar Rats. *Nutrients*, **14**, 1256.

Karigo, T. & Deutsch, D. (2022) Flexibility of neural circuits regulating mating behaviors in mice and flies. *Frontiers in Neural Circuits*, **16**.

Keller, M., Pierman, S., Douhard, Q., Baum, M.J., & Bakker, J. (2006) The vomeronasal organ is required for the expression of lordosis behaviour, but not sex discrimination in female mice. *European Journal of Neuroscience*, **23**, 521–530.

Kirouac, L., Rajic, A.J., Cribbs, D.H., & Padmanabhan, J. (2017) Activation of Ras-ERK signaling and GSK-3 by amyloid precursor protein and amyloid beta facilitates neurodegeneration in Alzheimer's disease. *eNeuro*, **4**.

Kiuchi, Z., Nishibori, Y., Kutsuna, S., Kotani, M., Hada, I., Kimura, T., Fukutomi, T., Fukuhara, D., Ito-Nitta, N., Kudo, A., Takata, T., Ishigaki, Y., Tomosugi, N., Tanaka, H., Matsushima, S., Ogasawara, S., Hirayama, Y., Takematsu, H., & Yan, K. (2019) GLCCI1 is a novel protector against glucocorticoid-induced apoptosis in T cells. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, **33**, 7387–7402.

Lemoine, S.R.T., Samuni, L., Crockford, C., & Wittig, R.M. (2022) Parochial cooperation in wild chimpanzees: A model to explain the evolution of parochial altruism. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **377**.

Lenz, K.M., Nugent, B.M., Haliyur, R., & McCarthy, M.M. (2013) Microglia are essential to masculinization of brain and behavior. *Journal of Neuroscience*, **33**, 2761–2772.

Lenz, K.M., Pickett, L.A., Wright, C.L., Davis, K.T., Joshi, A., & McCarthy, M.M. (2018) Mast cells in the developing brain determine adult sexual behavior. *Journal of Neuroscience*, **38**, 8044–8059.

Liu, Z.W., Yu, Y., Lu, C., Jiang, N., Wang, X.P., Xiao, S.Y., & Liu, X.M. (2019) Postweaning isolation rearing alters the adult social, sexual preference and mating behaviors of male CD-1 mice. *Frontiers in Behavioral Neuroscience*, **13**.

MacPherson, R.E.K. (2017) Filling the void: A role for exercise-induced BDNF and brain amyloid precursor protein processing. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology*, **313**, R585–R593.

Manning, A. & Thompson, M.L. (1976) Postcastration retention of sexual behaviour in the male BDF1 mouse: The role of experience. *Animal Behaviour*, **24**, 523–533.

Mass, E., Jacome-Galarza, C.E., Blank, T., Lazarov, T., Durham, B.H., Ozkaya, N., Pastore, A., Schwabenland, M., Chung, Y.R., Rosenblum, M.K., Prinz, M., Abdel-Wahab, O., & Geissmann, F. (2017) A somatic mutation in erythro-myeloid progenitors causes neurodegenerative disease. *Nature*, **549**, 389–393.

McCarthy, M.M., Vries, G.J., & Forger, N.G. (2017) Sexual Differentiation of the Brain: A Fresh Look at Mode, Mechanisms, and Meaning. In *Hormones, Brain and Behavior*. Elsevier, pp. 3–32.

McGill, T.E. & Tucker, G.R. (1964) Genotype and sex drive in intact and in castrated male mice. *Science*, **145**, 514–515.

McInnis, C.M., Venu, S., & Park, J.H. (2016) Steroid-independent male sexual behavior in B6D2F2 male mice. *Hormones and Behavior*, **85**, 26–29.

Melis, M.R. (1998) Nitric oxide production is increased in the paraventricular nucleus of the hypothalamus of male rats during non-contact penile erections and copulation. *European Journal of Neuroscience*, **10**, 1968–1974.

Melis, M.R., Succu, S., Mascia, M.S., Cortis, L., & Argiolas, A. (2003) Extra-cellular dopamine increases in the paraventricular nucleus of male rats during sexual activity. *European Journal of Neuroscience*, **17**, 1266–1272.

Michael, B.D., Bricio-Moreno, L., Sorensen, E.W., Miyabe, Y., Lian, J., Solomon, T., Kurt-Jones, E.A., & Luster, A.D. (2020) Astrocyte- and Neuron-Derived CXCL1 Drives Neutrophil Transmigration and Blood-Brain Barrier Permeability in Viral Encephalitis. *Cell Reports*, **32**.

Moffitt, J.R., Bambah-Mukku, D., Eichhorn, S.W., Vaughn, E., Shekhar, K., Perez, J.D., Rubinstein, N.D., Hao, J., Regev, A., Dulac, C., & Zhuang, X. (2018) Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic region. *Science*, **362**.

Monk, J.D., Giglio, E., Kamath, A., Lambert, M.R., & McDonough, C.E. (2019) An alternative hypothesis for the evolution of same-sex sexual behaviour in animals. *Nature Ecology and Evolution*, **3**, 1622–1631.

Moreira, V.B., Mattaraia, V.G.M., & Moura, A.S.A.M.T. (2015) Lifetime Reproductive Efficiency of BALB/c Mouse Pairs after an Environmental Modification at 3 Mating Ages. *Journal of the American Association for Laboratory Animal Science : JAALAS*, **54**, 29.

Numakawa, T., Kumamaru, E., Adachi, N., Yagasaki, Y., Izumi, A., & Kunugi, H. (2009) Glucocorticoid receptor interaction with TrkB promotes BDNF-triggered PLC-γ signaling for glutamate release via a glutamate transporter. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 647–652.

Okubo, T.S., Patella, P., D'Alessandro, I., & Wilson, R.I. (2020) A Neural Network for Wind-Guided Compass Navigation. *Neuron*, **107**, 924-940.e18.

O'Neill, K., Aghaeepour, N., Špidlen, J., & Brinkman, R. (2013) Flow Cytometry Bioinformatics. *PLoS Computational Biology*, **9**.

Özgün, A. & Garipcan, B. (2021) Magnetic field-induced Ca2+ intake by mesenchymal stem cells is mediated by intracellular Zn2+ and accompanied by a Zn2+ influx. *Biochimica et Biophysica Acta - Molecular Cell Research*, **1868**.

Pabinger, S., Rödiger, S., Kriegner, A., Vierlinger, K., & Weinhäusel, A. (2014) A survey of tools for the analysis of quantitative PCR (qPCR) data. *Biomolecular Detection and Quantification*, **1**, 23–33.

Paolicelli, R.C., Sierra, A., Stevens, B., Tremblay, M.E., Aguzzi, A., Ajami, B., Amit, I., Audinat, E., Bechmann, I., Bennett, M., Bennett, F., Bessis, A., Biber, K., Bilbo, S., Blurton-Jones, M., Boddeke, E., Brites, D., Brône, B., Brown, G.C., Butovsky, O., Carson, M.J., Castellano, B., Colonna, M., Cowley, S.A., Cunningham, C., Davalos, D., De Jager, P.L., de Strooper, B., Denes, A., Eggen, B.J.L., Eyo, U., Galea, E., Garel, S., Ginhoux, F., Glass, C.K., Gokce, O., Gomez-Nicola, D., González, B., Gordon, S., Graeber, M.B., Greenhalgh, A.D., Gressens, P., Greter, M., Gutmann, D.H., Haass, C., Heneka, M.T., Heppner, F.L., Hong, S., Hume, D.A., Jung, S., Kettenmann, H., Kipnis, J., Koyama, R., Lemke, G., Lynch, M., Majewska, A., Malcangio, M., Malm, T., Mancuso, R., Masuda, T., Matteoli, M., McColl, B.W., Miron, V.E., Molofsky, A.V., Monje, M., Mracsko, E., Nadjar, A., Neher, J.J., Neniskyte, U., Neumann, H., Noda, M., Peng, B., Peri, F., Perry, V.H., Popovich, P.G., Pridans, C., Priller, J., Prinz, M., Ragozzino, D., Ransohoff, R.M., Salter, M.W., Schaefer, A., Schafer, D.P., Schwartz, M., Simons, M., Smith, C.J., Streit, W.J., Tay, T.L., Tsai, L.H., Verkhratsky, A., von Bernhardi, R., Wake, H., Wittamer, V., Wolf, S.A., Wu, L.J., & Wyss-Coray, T. (2022) Microglia states and nomenclature: A field at its crossroads. *Neuron*, **110**, 3458–3483.

Park, J.H., Bonthius, P.J., Tsai, H.W., Bekiranov, S., & Rissman, E.F. (2010) Amyloid β precursor protein regulates male sexual behavior. *Journal of Neuroscience*, **30**, 9967–9972.

Park, J.H., Bonthuis, P., Ding, A., Rais, S., & Rissman, E.F. (2009) Androgen- and estrogen-independent regulation of copulatory behavior following castration in male B6D2F1 mice. *Hormones and Behavior*, **56**, 254–263.

Pelé, M., Bonnefoy, A., Shimada, M., & Sueur, C. (2017) Interspecies sexual behaviour between a male Japanese macaque and female sika deer. *Primates*, **58**, 1–4.

Pislyagin, E.A., Manzhulo, I.V., Gorpenchenko, T.Y., Dmitrenok, P.S., Avilov, S.A., Silchenko, A.S., Wang, Y.M., & Aminin, D.L. (2017) Cucumarioside A2-2 causes macrophage activation in mouse spleen. *Marine Drugs*, **15**.

Ralph, C.R., Lehman, M.N., Goodman, R.L., & Tilbrook, A.J. (2016) Impact of psychosocial stress on gonadotrophins and sexual behaviour in females: Role for cortisol? *Reproduction*, **152**, R1–R14.

Reichard, A. & Asosingh, K. (2019) Best Practices for Preparing a Single Cell Suspension from Solid Tissues for Flow Cytometry. *Cytometry Part A*, **95**, 219–226.

Rode, I. & Boehm, T. (2012) Regenerative capacity of adult cortical thymic epithelial cells. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 3463–3468.

Romano, A., Parrinello, N.L., Parisi, M., Del Fabro, V., Curtopelle, A., Leotta, S., Conticello, C., & Di Raimondo, F. (2021) Lenalidomide and pomalidomide improve function and induce fcγri/cd64 in multiple myeloma neutrophils. *Biomedicines*, **9**.

Sanmarco, L.M., Wheeler, M.A., Gutiérrez-Vázquez, C., Polonio, C.M., Linnerbauer, M., Pinho-Ribeiro, F.A., Li, Z., Giovannoni, F., Batterman, K.V., Scalisi, G., Zandee, S.E.J., Heck, E.S., Alsuwailm, M., Rosene, D.L., Becher, B., Chiu, I.M., Prat, A., & Quintana, F.J. (2021) Gut-licensed IFNγ+ NK cells drive LAMP1+TRAIL+ anti-inflammatory astrocytes. *Nature*, **590**, 473–479.

Serrano-Saiz, E. & Isogai, Y. (2021) Single-cell molecular and developmental perspectives of sexually dimorphic circuits underlying innate social behaviors. *Current Opinion in Neurobiology*, **68**, 159–166.

Smith, A.J. (2020) Guidelines for planning and conducting high-quality research and testing on animals. *Laboratory Animal Research*, **36**, 1–6.

Soh, K.T., Tario, J.D., Hahn, T., Hillengass, J., McCarthy, P.L., & Wallace, P.K. (2021) CD319 (SLAMF7) an alternative marker for detecting plasma cells in the presence of daratumumab or elotuzumab. *Cytometry Part B - Clinical Cytometry*, **100**, 497–508.

Sokolov, O.Y., Prokhorova, T.A., Tereshkina, E.B., Zozulya, S.A., Simonov, A.N., Kost, N.V., Dadayan, A.K., Bogachouk, A.P., & Zolotarev, Y.A. (2021) Neurotropic peptide HLDF-6-amide reduces age-related decline in sexual activity in old male rats. *Experimental Gerontology*, **149**.

Spaanderman, D.C.E., Nixon, M., Buurstede, J.C., Sips, H.H.C.M., Schilperoort, M., Kuipers, E.N., Backer, E.A., Kooijman, S., Rensen, P.C.N., Homer, N.Z.M., Walker, B.R., Meijer, O.C., & Kroon, J. (2018) Androgens modulate glucocorticoid receptor activity in adipose tissue and liver. *Journal of Endocrinology*, **240**, 51–63.

Stein, C.M., Weiskirchen, R., Damm, F., & Strzelecka, P.M. (2021) Single-cell omics: Overview, analysis, and application in biomedical science. *Journal of Cellular Biochemistry*, **122**, 1571–1578.

Steinberg, D. (2020) Immoral-like Behavior in Animals. In *The Multidisciplinary Nature of Morality and Applied Ethics*. Springer International Publishing, pp. 37–44.

Sun, J.J., Ryou, S., Goldshmid, R.H., Weissbourd, B., Dabiri, J.O., Anderson, D.J., Kennedy, A., Yue, Y., & Perona, P. (2022) Self-Supervised Keypoint Discovery in Behavioral Videos. *arXiv*,.

Sun, R.F., Zhao, C.Y., Chen, S., Yu, W., Zhou, M.M., & Gao, C.R. (2021) Androgen Receptor Stimulates Hexokinase 2 and Induces Glycolysis by PKA/CREB Signaling in Hepatocellular Carcinoma. *Digestive Diseases and Sciences*, **66**, 802–813.

Takayama, K. ichi, Fujiwara, K., & Inoue, S. (2019) Amyloid precursor protein, an androgen-regulated gene, is targeted by RNA-binding protein PSF/SFPQ in neuronal cells. *Genes to Cells*, **24**, 719–730.

Templin, J.S., Wyrosdic, J.C., David, C.D., Wyrosdic, B.N., Lapp, H.E., Bala, A., Bartlett, A., Khan, Z.,
Rokicki, A., & Park, J.H. (2019) Peripubertal gonadal steroids are necessary for steroid-
independent male sexual behavior in castrated B6D2F1 male mice. *Hormones and Behavior*,
113, 38–46.
Tsuneoka, Y. & Funato, H. (2021) Cellular Composition of the Preoptic Area Regulating Sleep,
Parental, and Sexual Behavior. *Frontiers in Neuroscience*, **15**.
Wang, S. & Brinkman, R.R. (2019) Data-driven flow cytometry analysis. In *Methods in Molecular*
Biology. Humana Press Inc., pp. 245–265.
Wee, B.E. & Clemens, L.G. (1989) Environmental influences on masculine sexual behavior in mice.
Physiol Behav, **46**, 867–872.
Wei, D., Talwar, V., & Lin, D. (2021) Neural circuits of social behaviors: Innate yet flexible. *Neuron*,
109, 1600–1620.
Wersinger, S.R. & Rissman, E.F. (2000) Dopamine activates masculine sexual behavior independent of
the estrogen receptor α. *Journal of Neuroscience*, **20**, 4248–4254.
Will, R.G., Hull, E.M., & Dominguez, J.M. (2014) Influences of dopamine and glutamate in the medial
preoptic area on male sexual behavior. *Pharmacology Biochemistry and Behavior*, **121**, 115–
123.
Wilson, E.N., Anderson, M., Snyder, B., Duong, P., Trieu, J., Schreihofer, D.A., & Cunningham, R.L.
(2018) Chronic Intermittent Hypoxia Induces Hormonal and Male Sexual Behavioral Changes:
Hypoxia as an Advancer of Aging. *Physiology & behavior*, **189**, 64.
Wu, B., Liang, Y., Dong, Z., Chen, Z., Zhang, G., Lin, W., Wang, S., Wang, B., Ge, R.S., & Lian, Q.
(2016) Glucocorticoid receptor mediated the propofol self-administration by dopamine D1
receptor in nucleus accumbens. *Neuroscience*, **328**, 184–193.
Yao, J., Ma, Y., Zhou, S., Bao, T., Mi, Y., Zeng, W., Li, J., & Zhang, C. (2020) Metformin prevents
follicular atresia in aging laying chickens through activation of PI3K/Akt and calcium signaling
pathways. *Oxidative Medicine and Cellular Longevity*, **2020**.
Zhang, S.X., Lutas, A., Yang, S., Diaz, A., Fluhr, H., Nagel, G., Gao, S., & Andermann, M.L. (2021)
Hypothalamic dopamine neurons motivate mating through persistent cAMP signalling. *Nature*
2021 597:7875, **597**, 245–249.

Sample Gating Strategy

Fluorescence Minus One Controls

	Mg ERa	Mg AR	Neu ERa	Neu AR	Mg GR	Mg APP	Neu GR		Legend
Mg ERa	1.000	-0.230	0.957	-0.223	-0.360	-0.047	-0.018		1.000
Mg AR	-0.230	1.000	-0.396	0.878	0.517	-0.138	-0.379		0.700
Neu ERa	0.957	-0.396	1.000	-0.308	-0.337	-0.197	0.087		0.350
Neu AR	-0.223	0.878	-0.308	1.000	0.407	-0.462	-0.534		0.000
Mg GR	-0.360	0.517	-0.337	0.407	1.000	-0.446	0.186		-0.350
Mg APP	-0.047	-0.138	-0.197	-0.462	-0.446	1.000	0.172		-0.700
Neu GR	-0.018	-0.379	0.087	-0.534	0.186	0.172	1.000		-1.000