Copepod life history evolution under high and low food regimes

Alexander Blake*, Dustin J. Marshall

alexander.gangur@monash.edu

dustin.marshall@monash.edu

School of Biological Sciences, Monash University, 25 Rainforest Walk. Clayton, VIC, 3800, Australia

*Corresponding author

Abstract

Copepods play a critical role in the carbon cycle of the planet – they mediate the sequestration of

carbon into the deep ocean, and are the trophic link between phytoplankton and marine foodwebs.

Global change stressors that decrease copepod productivity create the potential for catastrophic

positive feedback loops. Accordingly, a growing list of studies examine the evolutionary capacity of

copepods to adapt to the two primary stressors associated with global change: warmer

temperatures and lower pH. But the evolutionary capacity of copepods to adapt to changing food

regimes, the third major stressor associated with global change, remains unknown. We used

experimental evolution to explore how a 10-fold difference in food availability affects life history

evolution in the copepod, *Tisbe sp.* over two years, and spanning 30+ generations. Different food

regimes evoked evolutionary responses across the entire copepod life history: we observed

evolution in body size, size-fecundity relationships and offspring investment strategies. Our results

suggest that changes to food regimes reshape life histories and that cryptic evolution in traits such

as body size is likely. We demonstrate that evolution in response to changes in ocean productivity

will alter consumer life histories, and may distort trophic links in marine foodchains. Evolution in

response to changing phytoplankton productivity may alter the efficacy of the global carbon pump in

ways that have not been anticipated until now.

Introduction

The oceans account for around 50% of carbon sequestration (Sabine *et al.* 2004). Phytoplankton are

the primary producers that account for most marine carbon fixation (del Giorgio & Duarte 2002), but

only a small proportion is sequestered into the deep sea (Dunne *et al.* 2007). Copepods play a key

role here, they feed on phytoplankton, and excrete them in faecal pellets that much sink faster than

individual phytoplankton cells. Of the total biomass fixed every year, up to 15-58% is consumed by

copepods during some times of the year, and 7.4-29 gigatonnes of carbon are annually consumed on

average (Steinberg & Landry 2017). Given the massive biomass turnover between phytoplankton

and copepods, anything that affects the dynamics of this trophic link has significant consequences

for the efficacy global carbon pump. If global change affects copepods, then the carbon pump will

also be affected. Worse still, a positive feedback loop could be generated: increasing atmospheric

CO_2 increases stressors (e.g. higher temperatures, lower seawater pH), decreasing copepod

productivity, decreasing carbon sequestration rates, leading to more rapid increases in atmospheric

CO_2. The capacity for copepods to adapt to global change represents a key uncertainty in making

predictions for the future of the global carbon pump.

For organisms with relatively short lifespans, predicting their capacity to adapt to global change

requires an understanding of their evolutionary capacity to adapt to future conditions (Munday *et al.*

2013; Kelly & Griffiths 2021). Accordingly, a growing number of studies estimate how copepods

adapt to higher temperature regimes and lower pH – the two primary stressors associated with

global change in the sea (Brennan *et al.* 2021, 2022; deMayo *et al.* 2021; Sasaki & Dam 2021). While

temperature and pH are undoubtedly important, they are not the only factors that will change in

future oceans – food availability is likely to be dramatically different for copepods under future

regimes (Fu *et al.* 2016), and indeed is already changing (Capuzzo *et al.* 2018).

Phytoplankton depend on sunlight for photosynthesis, so they are most productive in the surface

waters of the ocean – the productivity of these surface waters are likely to alter with global change.

The upper, sunlit layers of the ocean are depleted of nutrients by growing phytoplankton and must

replenished by cooler, nutrient-rich water from deeper, darker layers. Unfortunately, global warming

can create more stratification in the water column, generating more intense thermoclines, and

reducing the rate of mixing between water layers (Hannon *et al.* 2001). This thermal shoaling is likely

to reduce phytoplankton productivity in some places by up to 40% (Jang *et al.* 2011). On the other

hand, some regions are predicted to experience more intense storms, which will actually enhance

mixing of surface waters with deeper layers, possibly increasing the productivity in those places

(Nicholson *et al.* 2016). Thus, copepods must not only evolve to cope with warmer, more acidic

oceans, they are also likely to experience massive changes to their food regimes. The capacity for

evolution in response to such changes remains unclear.

Food regimes have classically been recognised as key drivers of life history evolution across a broad

range of species (Macarthur & Levins 1967; Tilman 1982; Chesson 2000; Grant & Grant 2006), but

the direction and nature of such effects remain debated. For example, guppies in food-scarce

environments evolve larger body sizes (Felmy *et al.* 2022) while flies get smaller (Kolss *et al.* 2009).

As such, it is difficult to predict how, or even if, copepods will evolve in response to changes in food

based on studies in other systems. Biogeographical studies are similarly ambiguous. For example,

some imply that copepods should become larger under low productivity regimes but whether body

size is decreasing because of food availability *per se* or other covarying factors such as predation risk

remains unclear (Brun et al. 2016). As far as we are aware, no study has directly addressed the

capacity of copepods to evolve in response to different food regimes. We sought to address this

important knowledge gap here. We subjected copepod populations to either high-food or low-food

environments for 2 years and then used multigenerational common garden experiments to examine

how life histories have evolved independently of any cross-generational parental effects (Burgess &

Marshall 2014).

Materials and Methods

STUDY ORGANISM

Tisbe sp. is a littoral marine copepod from the Tisbidae family (Arthropoda: Harpacticoida) that has

not been resolved to species level in the Southern Ocean. Mothers brood clutches of approximately

10-40 eggs on abdominal egg-sacs for 1-3 days. The naupliar larvae are facultative planktotrophs,

and pass through 6 stages before metamorphosing into a juvenile copepodid approximately 3-5 days

after hatching with *ad libitum* food. Juveniles pass through a further 6 copepodid stages before

reaching the final adult and sexually mature stage at approximately 18-20 days old. We collected

copepods from Port Phillip Bay, Australia in 2017 and maintained these 'ancestral' populations in

gently oxygenated 500 ml mason jars in freshly pasteurised seawater (FSW). Ancestral populations

were reared on the marine microalga *Dunaliella tertiolecta* and fed at a rate of 2.475×10^9 algae cells

per litre of copepod culture per week.

Dunaliella cultures were maintained using F2 media (Guillard & Ryther 1962) and algae concentrate

for feeding copepod populations was prepared three times a week. Density of algal cultures was

determined spectroscopically using a SPECTROstar Nano and algal concentrations were adjusted to

approximately 1.1×10^{10} *D. tertiolecta* cells/L in FSW after centrifuging and removing media.

Ancestral populations were fed manually 3 times a week, while for experimental cultures housed in

flow-through culture vessels feeding was automated (see Experimental Evolution).

EXPERIMENTAL EVOLUTION

Experimental design

Experimental evolution commenced on the 13[th] of October 2018 at Monash University Clayton

Campus, Melbourne, Australia. Copepods from ancestral stocks were randomly assigned to either

high-food or low-food environments, which differed in their rate of food supply (of *D. tertiolecta*

cells) by an order of magnitude. A Folsom plankton splitter was used to ensure equal numbers of

copepods were assigned to each culture and the 1L glass culture vessels were topped up with FSW.

In total, 20 copepod cultures were subjected to experimental evolution, consisting of 10 high-food

and 10 low-food replicates reared in separate glass pressure-equalising dropping funnels. One low-

food replicate went extinct one year into the experiment due to bacterial contamination.

High and low food rates were determined through pilot experiments in 2018 and were set as

follows: high food replicates received 4.5 x 10^9 algae cells per litre of copepod culture per week, and

low food replicates received 4.5 x 10^8 algae cells per litre of copepod culture per week. An

intermediate feeding rate (2.475 x 10^9 algae cells per litre of copepod culture) was used for the

ancestral populations and the common garden. Differential food supply in experimental treatments

was ramped up gradually, with high and low food treatments receiving the same intermediate rate

of supply for the first week, partial treatments in week two (3.5 x 10^9 and 1.5 x 10^9 *D. tertiolecta* cells

per week, respectively), and final treatments of high or low food supply by week three after

initiating the experiment. These treatment differences were then maintained for the next 16

months.

Experimental cultures were organised into 5 feeding blocks of four (in a randomised sequence of 2

high-food and 2 low-food replicates) due to spatial constraints. Each block received food from a

separate algae reservoir using a Kamoer X4 peristaltic dosing pump. Both high-food and low-food

treatments received a total inflow of 80mL per litre of culture per day (on weekdays only, no dosing

on weekends). High-food treatments were dosed with 80mL of algae concentrate per day, while low-

food treatments received 8 mL of algae concentrate and 72mL of FSW (i.e. 10% of the high food

regime). Pumps provisioned algae concentrate semi-continuously by dosing 12 times a day. Pilot

work indicated that very few adult copepods were lost at this rate of inflow-outflow. Algae and FSW

reservoirs were covered to minimise evaporation and cleaned weekly to minimise build-up of waste

and pathogens, and algae was kept well-mixed using magnetic stirrers.

COMMON GARDENS

Experimental design

To disentangle genetic responses from plastic responses in experimental evolution, individuals need

to be sampled from divergent evolutionary lineages and reared in a common environment (Huey &

Rosenzweig 2009). Because environmental effects can persist between generations, such common

environment (or 'common garden') experiments must also be performed over multiple generations

to minimise any lingering parental and grandparental effects on offspring phenotypes (Burgess &

Marshall 2014). To evaluate the evolutionary response to high- and low-food environments, we

performed a common garden experiment wherein copepods were sampled from their treatment

cultures (G0) and their descendants were reared (separately) under the same environmental

conditions over two generations (G1 and G2). With a generation time of ~17 days, our *Tisbe sp.*

cultures had undergone approximately 30 generations of evolution prior to common gardening,

which commenced on 18 February 2020. Evolutionary responses in five traits were measured across

the three generations. Maternal size, mean egg size, and fecundity were measured in G0, G1, and

G2, while survival and age at maturity were measured in G1 and G2 only. Maternal size was

measured as length between end of urosome to tip of prosome. We used mean egg size from 10

randomly measured eggs within each clutch, and also estimated fecundity as number of eggs per egg

sac. Maternal body size, egg size, and fecundity were recorded with photographs using a Motic

Moticam 1080 camera mounted on an Olympus SZ61 dissecting microscope and digitally measured

using FIJI version 1.53c (Schindelin et al. 2012).

Paired high- and low-food cultures were randomly sampled between the 18th and 20th of February

2020. 10 gravid G0 mothers were collected from each culture, photographed, and transferred to

sterile plastic culture trays containing 4 mL FSW. 16uL of 10000 units mL^{-1} (approximately 6 mg mL^{-1})

penicillin G and 10 mg ml^{-1} streptomycin solution (Sigma-Aldrich) was added to each tray to inhibit

the growth of pathogens (Gangur & Marshall 2020). G0 mothers were monitored daily and returned

to their cultures after releasing their G1 eggs, until all eggs had hatched and all G0 mothers had been

removed (generally 3-5 days after initial collection). All common garden replicates commenced G1

with >100 larvae.

Throughout the experiment, *D. tertiolecta* was provisioned at an intermediate level of food supply.

For G0, G1, and G2 juveniles and adults, food was provisioned each weekday (5 times a week) in a

176 uL pulse from a 1.1×10^7 cells/mL stock representing an intermediate food supply of

approximately 2.475×10^9 cells per litre per week. Due to their lower feeding rate, larvae were

provisioned a single 176uL (at 1.1×10^7 cells/mL) pulse of food to achieve the same maximum

ambient food density experienced by adults under an intermediate feed regime (approx. 5×10^6 *D*

tertiolecta cells per mL each day).

Data collection

Freshly hatched G1 larvae within each replicate were counted and randomly allocated to individual

culture trays of 25 ± 5 individuals with 4 mL FSW, antibiotics, and food. When metamorphosis was

first observed within a replicate, the replicate was censused and individuals were transferred to

fresh trays with 4 mL FSW, antibiotics, and food. For juveniles, water was then changed and

censusing was conducted weekly until sexual maturity was first observed within a replicate. Then

subreplicates were censused, pooled, mixed, and reallocated into new adult subreplicates of 25 ± 5

with fortnightly water changes and censusing. All reproductive mothers were collected and

photographed with their first clutch. Mothers and their egg sacs were photographed, then pooled at

the replicate level into fresh culture trays containing 4 mL FSW, antibiotics, and food. At least 5 G1

mothers and 50 offspring were obtained for 17 of 19 replicates (8 high-food and 9 low-food

cultures), and these G2 offspring were collected for the final stage of the experiment. Replicates

containing copepods from the two remaining high-food cultures were accidentally dropped before

reaching G2.

G2 larvae were collected from culture trays containing gravid G1 mothers in a similar fashion to G1

larvae collection from G0 mothers, but we also accounted for temporal staggering. For each

replicate, G2 larval collection took place over a week after the first clutch hatched. Larvae collected

during this week were continually transferred to culture trays at a density of 25 ± 5 individuals with 4

mL FSW, antibiotics, and food. At the end of the collection week these larvae (some of which had

metamorphosed) were re-pooled, mixed, and randomly allocated to new culture trays of 25 ± 5

individuals in 4 mL FSW, antibiotics, and food. G2 larvae that hatched outside this initial collection

week were retained but reared in separate trays. G2 larvae were then reared to sexual maturity and

reproductive G2 mothers were measured following the same protocol used for G1.

STATISTICAL ANALYSES

All phenotypes were analysed using linear mixed effects models. Full models for size, survival, and

age at maturity included treatment and generation as fixed effects, as well as their interaction.

Fecundity and egg size were modelled separately for each generation due to complex interactions

and included mother size as a fixed covariate, as well as its interaction with treatment. G0 feeding

block was included as a fixed effect in all models due to insufficient replication to treat as a random

effect. All models also included culture nested within treatment as a random intercept term. Where

interaction terms were nonsignificant they were removed and the analysis was repeated. Models

were evaluated using type III tests due to imbalance of high-food and low-food replicates. *p* values

for relevant fixed effects were obtained with *F* tests using Sattertwaithe's approximation

(Kuznetsova *et al.* 2017).

Analyses were performed with R version 4.1.2 (R Core Team 2021) and RStudio version 2021.09.1

(RStudio Team 2021), using dplyr (Wickham et al. 2021) to prepare the data. Linear mixed effect

models were fitted with lme4 (Bates et al. 2015). The lmerTest package (Kuznetsova et al. 2017) was

used to perform Sattertwaithe's approximations and type III tests on fixed effects, and likelihood

ratio tests on random effects. Bootstrapped 95% confidence intervals were obtained between

cultures using merTools (Knowles & Frederick 2020), and plots were built using ggplot2 (Wickham

2016). Diagnostic residuals vs fits and QQ plots were visually assessed as per Keough & Quinn (2002),

and VIF calculated to check for collinearity. Predictors were plotted against each other to visually

assess acceptable domain/range overlap.

Results

SURVIVAL AND BODY SIZE

Copepod body sizes evolved in response to the food regime, but the direction of difference changed

from one generation to another. Prior to common gardening, G0 females were slightly larger in the

high-food environments but in the common-gardened generations (G1 and G2) females were

consistently larger in the low-food regimes relative to the high-food regimes (Table 1, Figure 1a).

There was also an increase in body size over the course of the common gardening relative to G0

(Figure 1a). Survival in the common garden was unaffected by food regime (Table S1 and Figure S1,

see Supporting Information).

AGE AT MATURITY

Age at maturity differed between food regimes in G1 copepods but converged in G2 (Table 2, Figure

1b). In G1, copepods that had evolved in high-food regimes matured later despite being smaller than

copepods that had evolved in low-food regimes, but by G2, these differences had disappeared.

REPRODUCTIVE OUTPUT

The relationships between maternal size and reproduction evolved in response to food regime but

these effects only manifested in the second generation of common gardening (Egg size: Table 3,

Fecundity: Table 4). Larger mothers from the low-food lineages produced larger (Figure 1c) but

slightly fewer offspring (Figure 1d), whereas from the high-food lineages larger mothers produced

smaller (Figure 1c) but many more offspring (Figure 1d). Combining these two components of

reproduction, mothers from the low-food lineages showed a steeper positive relationship between

body size and reproductive output than mothers from the high food lineages (Figure 2).

Discussion

Copepods under different resource regimes evolved different life histories: body size, fecundity, and

per-offspring investment all evolved, while age at maturity also changed but appeared to be driven

by a strong parental environment effects that dissipated across generations. Copepods evolved to be

slightly larger in low food regimes, and within those regimes, larger mothers invested more in their

individual offspring in this environment. In high food regimes, copepods evolved to be smaller and

within those regimes, larger mothers invested less in their individual offspring but were much more

fecund. Interestingly, we found evidence for differences in evolved responses to different resource

environments relative to the expressed phenotypes in those environments, indicating

countergradient evolution. Overall, our results suggest that the changes in food regimes predicted to

occur in future oceans will generate life history evolution in copepods but not in straightforward

ways. Our results also imply that biogeographical patterns in life history and covariance between

productivity and phenotypes may provide very little information or predictive power about the

underlying genetic clines in these traits due countergradient variation.

Copepods were smaller in the low food regime but their offspring grew to be larger when

transferred to common environment – indicating countergradient evolution in body size. Assuming

cubic scaling with length, copepods from low-food environments were only 5.9% smaller by volume.

Once released from the low-food conditions, they were 5.6% larger than copepods from high-food

lineages, suggesting that the impact of food scarcity was moderated by genetic compensation

(Grether 2005). Such countergradient variation is observed in field studies as well and seems

particularly common in fish (Arendt & Wilson 1999; Conover et al. 2009). We suggest that global

changes to phytoplankton productivity will evoke evolutionary change in copepod body sizes but

that these changes may be masked by countergradient evolution. Studies seeking to understand

how copepod body sizes have changed, and continue to change should consider common garden

experiments to disentangle phenotypic and genetic responses, which may counteract each other,

resulting in what is sometimes called 'cryptic evolution' (Grether 2005).

The relationship between body size and reproductive investment evolved, albeit in subtle ways. In

low food environments, egg size increased with maternal size at the expense of fecundity while,

while clutch size increased with maternal size at the expense of egg size in high-food lineages. These

results are in keeping with general offspring size theory whereby in poor environments, mothers

make larger offspring in order to buffer them from harsh conditions (Parker & Begon 1986). Such

phenotypic effects have been observed in other taxa (e.g. Allen et al. 2008; Fox & Czesak 2000; see

Marshall et al. 2018 for a review) but to our knowledge, ours is one of the few unequivocal

demonstrations that differences in reproductive investment strategies can rapidly evolve.

The way reproduction scaled with maternal size showed clear evolutionary responses to food

regime. Regardless of evolutionary lineage, larger mothers reproduced more than smaller mothers,

implying larger mothers have more resources to invest in offspring. Larger individuals may be better

competitors, able to acquire more resources for reproduction (e.g. Bassar et al. 2016). Larger

mothers may also alter their allocation of resources among fitness components, or may simply be

physically able to brood more or larger eggs (Bernardo 1996). We find that the way in which larger

mothers deploy this resource advantage relative to smaller mother depends on the resource regime

in which they evolved – in low resource regimes, they make better provisioned offspring, in high

resource regimes, they make more offspring. These different allocation patterns yield an evolved

difference in reproductive scaling – reproductive scaling was steeper in low resource environments

than high resource environments. Interestingly, low resource lineages also evolved larger body sizes

(at least genetically) – it may be that they evolved to be larger so as to gain the fitness advantages of

increased body size that come from steeper reproductive scaling but this remains speculative.

Nevertheless, to our knowledge our study is the first to demonstrate experimental evolution in

reproductive scaling, which has clear consequences for population dynamics (Marshall et al. 2022).

Generally theory predicts that when reproductive scaling is steeper, population replenishment and

productivity depends more strongly on having larger females in population (Marshall et al. 2022).

Given smaller body sizes are expected under higher temperatures (Atkinson & Sibly 1997), our

results suggest that combining high temperature regimes (the redaction of larger females) and low

food conditions (steeper reproductive scaling) could synergise to generate catastrophic reductions in

population productivity. An important next step will be to examine how combined stressor regimes

(e.g. temperature and food) affect evolutionary trajectories.

We found differences in the timing of maturity in G1 copepods but not G2, suggesting that age at

maturity is a transgenerational plasticity effect rather than an evolved response. It seems to us at

least that the low food parental environment programs offspring to mature sooner than offspring

whose parents experience high food levels. While transgenerational plasticity in key life history traits

is relatively common (Yin *et al.* 2019), we are unaware of studies that have shown such effects on

the timing of maturity specifically. That this effect disappears after a single generation suggests that

this parentally program trait has evolved to track environmental variation closely, with minimal

persistent lags as predicted by some theory (Burgess & Marshall 2014).

In summary, we show that differing food regimes induce rapid (in less than 50 generations)

evolutionary responses, affecting every aspect of their life history from offspring size, through to

growth and reproduction. These evolutionary responses may maximise the fitness of individuals in

their particular food regimes but will undoubtedly wreak changes to the productivity of whole

populations. None of the responses we observed were entirely predictable based on existing theory

or studies in other systems. Our findings emphasise that evolution will alter and complicate

biological responses to global change – with concomitant changes to global food webs that cannot

be anticipated based on ecological experiments alone. An important next step is to understand the

eco-evolutionary consequences of copepod life history evolution for higher trophic links – how do

planktivores that depend on copepods evolve in response to copepod evolution themselves?

References

Allen, R.M., Buckley, Y.M. & Marshall, D.J. (2008). Offspring Size Plasticity in Response to
Intraspecific Competition: An Adaptive Maternal Effect across Life-History Stages. *The*
American Naturalist, 171, 225–237.
Arendt, J.D. & Wilson, D.S. (1999). Countergradient selection for rapid growth in pumpkinseed
sunfish: disentangling ecological and evolutionary effects. *Ecology*, 80, 2793–2798.
Atkinson, D. & Sibly, R.M. (1997). Why are organisms usually bigger in colder environments? Making
sense of a life history puzzle. *Trends in Ecology & Evolution*, 12, 235–239.
Bassar, R.D., Childs, D.Z., Rees, M., Tuljapurkar, S., Reznick, D.N. & Coulson, T. (2016). The effects of
asymmetric competition on the life history of Trinidadian guppies. *Ecology Letters*, 19, 268–
278.
Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software*, 67, 1–48.
Bernardo, J. (1996). The particular maternal effect of propagule size, especially egg size: patterns,
models, quality of evidence and interpretations. *American Zoologist*, 36, 216–236.
Brennan, R.S., deMayo, J.A., Dam, H.G., Finiguerra, M.B., Baumann, H., Buffalo, V. & Pespeni, M.H.
(2022). Experimental evolution reveals the synergistic genomic mechanisms of adaptation to
ocean warming and acidification in a marine copepod. *Proceedings of the National Academy*
of Sciences, 119, e2201521119.
Brennan, R.S., deMayo, J.A., Dam, H.G., Finiguerra, M.B., Baumann, H. & Pespeni, M.H. (2022). Loss
of transcriptional plasticity but sustained adaptive capacity after adaptation to global change
conditions in a marine copepod. *Nature Communications*, 13, 1–13.
Brun, P., Payne, M.R. & Kiørboe, T. (2016). Trait biogeography of marine copepods – an analysis
across scales. *Ecology Letters*, 19, 1403–1413.
Burgess, S.C. & Marshall, D.J. (2014). Adaptive parental effects: the importance of estimating
environmental predictability and offspring fitness appropriately. *Oikos*, 123, 769–776.
Capuzzo, E., Lynam, C.P., Barry, J., Stephens, D., Forster, R.M., Greenwood, N., McQuatters-Gollop,
A., Silva, T., van Leeuwen, S.M. & Engelhard, G.H. (2018). A decline in primary production in
the North Sea over 25 years, associated with reductions in zooplankton abundance and fish
stock recruitment. *Global Change Biology*, 24, e352–e364.
Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review Ecology and*
Systematics, 31, 343–366.
Conover, D.O., Duffy, T.A. & Hice, L.A. (2009). The covariance between genetic and environmental
influences across ecological gradients: reassessing the evolutionary significance of
countergradient and cogradient variation. *Annals of the New York Academy of Sciences*,
1168, 100–129.
deMayo, J.A., Girod, A., Sasaki, M.C. & Dam, H.G. (2021). Adaptation to simultaneous warming and
acidification carries a thermal tolerance cost in a marine copepod. *Biology Letters*, 17,
20210071.
Dunne, J.P., Sarmiento, J.L. & Gnanadesikan, A. (2007). A synthesis of global particle export from the
surface ocean and cycling through the ocean interior and on the seafloor. *Global*
Biogeochemical Cycles, 21.
Felmy, A., Reznick, D.N., Travis, J., Potter, T. & Coulson, T. (2022). Life histories as mosaics: Plastic
and genetic components differ among traits that underpin life-history strategies. *Evolution*,
76, 585–604.
Fox, C.W. & Czesak, M.E. (2000). Evolutionary Ecology of Progeny Size in Arthropods. *Annual Review*
of Entomology, 45, 341–369.
Fu, W., Randerson, J.T. & Moore, J.K. (2016). Climate change impacts on net primary production
(NPP) and export production (EP) regulated by increasing stratification and phytoplankton
community structure in the CMIP5 models. *Biogeosciences*, 13, 5151–5170.

del Giorgio, P.A. & Duarte, C.M. (2002). Respiration in the open ocean. *Nature*, 420, 379–384.

Grant, P. & Grant, B.R. (2006). Evolution of Character Displacement in Darwin's Finches. *Science,*
347   *New Series*, 313, 224–226.

348   Grether, G.F. (2005). Environmental Change, Phenotypic Plasticity, and Genetic Compensation. *The*
349   *American Naturalist*, 166, E115–E123.

350   Hannon, E., Boyd, P., Silvoso, M. & Lancelot, C. (2001). Modeling the bloom evolution and carbon
351   flows during SOIREE: Implications for future in situ iron-enrichments in the Southern Ocean.
352   *Deep Sea Research Part II: Topical Studies in Oceanography*, 48, 2745–2773.

353   Hoffmann, A.A. & Harshman, L.G. (1999). Desiccation and starvation resistance in Drosophila:
354   patterns of variation at the species, population and intrapopulation levels. *Heredity*, 83,
355   637–643.

356   Huey, R.B. & Rosenzweig, F. (2009). Laboratory evolution meets catch-22: balancing simplicity and
357   realism. *Experimental evolution: Concepts, methods, and applications of selection*
358   *experiments*, 671–702.

359   Jang, C.J., Park, J., Park, T. & Yoo, S. (2011). Response of the ocean mixed layer depth to global
360   warming and its impact on primary production: a case for the North Pacific Ocean. *ICES*
361   *Journal of Marine Science*, 68, 996–1007.

362   Kelly, M.W. & Griffiths, J.S. (2021). Selection Experiments in the Sea: What Can Experimental
363   Evolution Tell Us About How Marine Life Will Respond to Climate Change? *The Biological*
364   *Bulletin*, 241, 30–42.

365   Knowles, J.E. & Frederick, C. (2020). *merTools: Tools for Analyzing Mixed Effect Regression Models*.

366   Kolss, M., Vijendravarma, R.K., Schwaller, G. & Kawecki, T.J. (2009). Life-history consequences of
367   adaptation to larval nutritional stress in Drosophila. *Evolution: International Journal of*
368   *Organic Evolution*, 63, 2389–2401.

369   Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2017). lmerTest Package: Tests in Linear Mixed
370   Effects Models. *Journal of Statistical Software*, 82, 1–26.

371   Macarthur, R. & Levins, R. (1967). The Limiting Similarity, Convergence, and Divergence of Coexisting
372   Species. *The American Naturalist*, 101, 377–385.

373   Marshall, D.J., Malerba, M., Lines, T., Sezmis, A.L., Hasan, C.M., Lenski, R.E. & McDonald, M.J. (2022).
374   Long-term experimental evolution decouples size and production costs in Escherichia coli.
375   *Proceedings of the National Academy of Sciences*, 119, e2200713119.

376   Marshall, D.J., Pettersen, A.K. & Cameron, H. (2018). A global synthesis of offspring size variation, its
377   eco-evolutionary causes and consequences. *Functional Ecology*, 32, 1436–1446.

378   Munday, P.L., Warner, R.R., Monro, K., Pandolfi, J.M. & Marshall, D.J. (2013). Predicting evolutionary
379   responses to climate change in the sea. *Ecology Letters*, 16, 1488–1500.

380   Nicholson, S.-A., Lévy, M., Llort, J., Swart, S. & Monteiro, P.M.S. (2016). Investigation into the impact
381   of storms on sustaining summer primary productivity in the Sub-Antarctic Ocean.
382   *Geophysical Research Letters*, 43, 9192–9199.

383   Parker, G.A. & Begon, M. (1986). Optimal Egg Size and Clutch Size: Effects of Environment and
384   Maternal Phenotype. *The American Naturalist*, 128, 573–592.

385   R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for
386   Statistical Computing, Vienna, Austria.

387   RStudio Team. (2021). *RStudio: Integrated Development Environment for R*. RStudio, PBC, Boston,
388   MA.

389   Sabine, C.L., Feely, R.A., Gruber, N., Key, R.M., Lee, K., Bullister, J.L., *et al.* (2004). The Oceanic Sink
390   for Anthropogenic CO_2. *Science*, 305, 367–371.

391   Sasaki, M.C. & Dam, H.G. (2021). Negative relationship between thermal tolerance and plasticity in
392   tolerance emerges during experimental evolution in a widespread marine invertebrate.
393   *Evolutionary Applications*, 14, 2114–2123.

394   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Rueden, C., Saalfeld,
395   S., Schmid, B., Tinevez, J, White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A.

(2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, 9, 676–

682.

Steinberg, D.K. & Landry, M.R. (2017). Zooplankton and the Ocean Carbon Cycle. *Annual Review of*

Marine Science, 9, 413–444.

Tilman, D. (1982). *Resource competition and community structure*. Princeton university press.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wickham, H., François, R., Henry, L. & Müller, K. (2021). *dplyr: A Grammar of Data Manipulation*.

Yin, J., Zhou, M., Lin, Z., Li, Q.Q. & Zhang, Y.-Y. (2019). Transgenerational effects benefit offspring

across diverse environments: a meta-analysis in plants and animals. *Ecology Letters*, 22,

1976–1986.

Table 1: Linear mixed effects model for the effect of evolutionary treatment on adult female body

size across three generations (G0, G1, G2), with ancestral G0 cultures reared within feeding blocks of

four. *p* values are provided for tests of interest, significant effects are specified in bold, and random

effect is specified in italics. Degrees of freedom (df) reported as numerator df, denominator df.

Effect	Df	Mean squares	F	*p*
Treatment	1, 11.67	5.69×10^{-1}	0.16	0.70
Generation	2, 691.43	3.20×10^{-4}	287.52	**$<10^{-15}$**
Block	4, 11.76	1.95×10^{-3}	0.98	0.45
Treat x Gen	2, 686.54	1.01×10^{-2}	5.10	**0.006**
Culture(Treat)		1.93×10^{-3}		

412 **Table 2:** Linear mixed effects model for the effect of evolutionary treatment on age at maturity

413 across two generations (G1, G2), with ancestral G0 cultures reared within feeding blocks of four. *p*

414 values are provided for tests of interest, significant effects are specified in bold, and random effect is

415 specified in italics. Degrees of freedom (df) reported as numerator df, denominator df.

Effect	Df	Mean squares	F	*p*
Treatment	1, 10.26	95.65	6.31	**0.03**
Generation	1, 517.99	2.39	0.16	0.69
Block	4, 10.44	15.20	1.00	0.45
Treat x Gen	1, 517.91	517.91	18.33	**$<10^{-4}$**
Culture(Treat)		14.74		

418   **Table 3:** Linear mixed effects models for the effects of evolutionary treatment and maternal size on

419   mean egg size across three generations (G0, G1, G2), with ancestral G0 cultures reared within

420   feeding blocks of four. *p* values are provided for tests of interest, significant effects are specified in

421   bold, and random effect is specified in italics. Degrees of freedom (df) reported as numerator df,

422   denominator df.

Effect	Df	Mean squares	F	*p*
G0				
Treatment	1, 12.98	1.36×10^{-4}	3.00	0.11
Maternal Size	1, 77.59	7.36×10^{-5}	1.62	0.21
Treat x Size	1, 95.48	3.12×10^{-5}	0.69	0.41
Block	4, 12.45	3.87×10^{-5}	0.85	0.52
Culture(Treat)		4.14×10^{-5}		
G1				
Treatment	1, 12.32	1.83×10^{-6}	0.07	0.80
Maternal Size	1, 184.00	2.24×10^{-5}	0.84	0.36
Treat x Size	1, 184.00	1.39×10^{-5}	0.52	0.47
Block	4, 13.11	8.57×10^{-6}	0.32	0.86
Culture(Treat)		2.48×10^{-5}		
G2				
Treatment	1, 232.57	6.11×10^{-4}	17.69	**$<10^{-4}$**
Maternal Size	1, 231.65	9.87×10^{-4}	2.85	0.09
Treat x Size	1, 233.32	5.99×10^{-3}	17.32	**$<10^{-4}$**
Block	4, 11.35	9.57×10^{-4}	2.77	0.08
Culture(Treat)		3.25×10^{-5}		

423 **Table 4:** Linear mixed effects models for the effects of evolutionary treatment and maternal size on

424 fecundity across three generations (G0, G1, G2), with ancestral G0 cultures reared within feeding

425 blocks of four. *p* values are provided for tests of interest, significant effects are specified in bold, and

426 random effect is specified in italics. Degrees of freedom (df) reported as numerator df, denominator

427 df.

Effect	Df	Mean squares	F	*p*
G0				
Treatment	1, 11.15	0.83	0.05	0.83
Maternal Size	1, 102.98	206.72	11.38	**0.001**
Treat x Size	1, 114.55	3.11	0.17	0.68
Block	4, 10.98	10.98	0.48	0.75
Culture(Treat)		16.10		
G1				
Treatment	1, 9.99	8.75	0.27	0.62
Maternal Size	1, 183.48	863.73	26.38	**$<10^{-6}$**
Treat x Size	1, 183.95	0.99	0.03	0.86
Block	4, 10.57	9.39	0.29	0.88
Culture(Treat)		30.30		
G2				
Treatment	1, 234	383.72	7.44	**0.007**
Maternal Size	1, 234	165.44	3.86	0.051
Treat x Size	1, 234	342.10	6.69	**0.01**
Block	4, 234	271.44	5.36	**$<10^{-3}$**
Culture(Treat)		49.71		

Figure 1: Key life history responses to high (red, circular points) and low (blue, triangular points) food

regimes. Panels show a: mean adult female body length across all three generations, b: mean cohort

age at first observation of sexual maturity (within each replicate) in G1 and G2, c: mean first-clutch

egg diameter in G2 mothers, d: mean first-clutch fecundity in G2 mothers. In a and b: errors bars

show between-culture bootstrapped 95% confidence intervals. In c and d: points show raw data at

the sub-replicate (individual female) level, lines show regressions at the culture level.

Figure 2: Predicted relationships between reproductive volume (product of mean egg volume and fecundity) and maternal size in G2 mothers from high-food (solid red lines, circular points) and low-food (dotted blue lines, triangular points) regimes. Points show data at the individual level, lines show regressions at the culture level, data obtained from G2 egg-size and G2 fecundity model coefficients.