m^{6}A modification of TERRA RNA is required for telomere maintenance and is a therapeutic target for ALT positive Neuroblastoma

Roshan Vaid[1][†], Ketan Thombare[1][†], Akram Mendez[1], Rebeca Burgos-Panadero[1], Anna Djos[1], Daniel Jachimowicz[2], Christoph Bartenhagen[3], Navinder Kumar[4], Carina Sihlbom[5], Susanne Fransson[1], John Inge Johnsen[6], Per Kogner[6], Tommy Martinsson[1], Matthias Fischer[3], Tanmoy Mondal[1,7][*]

[1]Department of Laboratory Medicine, Institute of Biomedicine, University of Gothenburg, Gothenburg, Sweden.

[2]Translational Genomics, Discovery Biology, Discovery Sciences, BioPharmaceuticals R&D, AstraZeneca, Gothenburg, Sweden.

[3]Department of Experimental Pediatric Oncology, University Children's Hospital of Cologne, Medical Faculty, Cologne, Germany, Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany, Department of Pediatric Oncology and Hematology, University of Cologne, Cologne, Germany.

[4]Department of Microbiology and Immunology, Institute of Biomedicine, University of Gothenburg.

[5]Proteomics Core Facility, Sahlgrenska Academy at the University of Gothenburg, Medicinaregatan 5, 41390 Gothenburg, Sweden.

[6]Childhood Cancer Research Unit, Department of Women's and Children's Health, Karolinska Institute, and Pediatric Oncology, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden.

[7]Department of Clinical Chemistry, Sahlgrenska University Hospital, Gothenburg, Sweden.

[†]**Equal contribution**

[*]Corresponding author, tanmoy.mondal@gu.se.

Department of Clinical Chemistry, Bruna Stråket 16, Sahlgrenska University Hospital, Gothenburg University, Gothenburg, 41345, Sweden.

Abstract:

Telomerase-negative tumors can maintain telomere length by alternative lengthening of telomeres (ALT) but the mechanism behind ALT is poorly understood. Aggressive Neuroblastoma (NB), in particular, relapsed tumors are positive for ALT (ALT+) which suggests that better dissection of the ALT mechanism could provide novel therapeutic opportunities. TERRA long non-coding RNA (lncRNA) which is derived from the telomere ends is localized to telomeres in R-loop dependent manner and is essential for telomere maintenance. In the present study, we provide evidence that RNA modification at the N^6 position of internal adenosine (m^{6}A) in TERRA RNA by methyltransferase METTL3 is essential for telomere maintenance in ALT+ cells and that loss of TERRA m^{6}A/METTL3 leads to telomere damage. We observed that R-loop enriched TERRA is abundantly m^{6}A modified and m^{6}A mediated recruitment of hnRNPA2B1 to TERRA RNA is essential for R-loop formation. Our data suggest that m^{6}A drives telomere targeting of TERRA via R-loop and this m^{6}A mediated R-loop formation could be a widespread mechanism utilized by other chromatin-interacting lncRNAs. Furthermore, treating ALT+ NB cells with METTL3 inhibitor leads to compromised telomere targeting of TERRA and accumulation of DNA damage over telomere, suggesting METTL3 inhibition could be a therapeutic opportunity for ALT+ NB.

Introduction:

RNA modification at the N^6 position of internal adenosine (m^{6}A) plays a critical role in regulating RNA splicing and many key steps in RNA metabolism. m^{6}A modification in RNA contributes to a wide range of cellular functions like stress response, DNA damage, cellular differentiation, and the regulation of oncogenes and tumor suppressor gene functions (1). METTL3 is the key enzyme in complex with METTL14 and WTAP, which deposits m^{6}A in RNA. Active chromatin marks H3K36me3 together with adaptor protein RBM15 and VIRMA (also known as KIAA1429) is thought to guide METTL3 complex to RNA substrates to perform m^{6}A modification (2-4). The "DRACH" like motif is the most preferred site for m^{6}A modification however, other non-canonical motifs can also be used for m^{6}A deposition by METTL3 (5). m^{6}A modification can also be dynamically regulated as m^{6}A demethylase like ALKBH5 can remove m^{6}A from the transcript (6). Once m^{6}A is deposited in RNA, reader proteins in the nucleus (i.e. YTHDC1, RBMX, hnRNPA2B1) and cytoplasmic (YTHDF1/2/3) identify m^{6}A residues to guide the functional complex to the modified RNA (1,7,8). Recently it has been shown that chromatin binding of METTL3 and METTL14 in both promoter and distal enhancer elements has been shown to regulate the transcriptional process (9). m^{6}A

modification in retrotransposon-derived transcripts modulates the local chromatin structure to

regulate transcription (5,10). RNA strand of the RNA:DNA hybrid containing R-loop structure

is enriched with m^{6}A modification and METTL3 depletion affects the global level of the R-

loop (11,12). How m^{6}A modification in the RNA modulates the R-loop level and the functional

significance of the presence of m^{6}A in the R-loop structure require further investigation.

The role of m^{6}A modification in long non-coding RNA (lncRNA) function is evident in several

studies (13,14). The *cis*-acting lncRNA *Xist* which localizes to inactive X-chromosome is m^{6}A

modified at several residues in both the "DRACH" and non- "DRACH" motif context. Xist is

associated with adaptor protein RBM15 which recruits the METTL3/METTL14 complex to

deposit m^{6}A on *Xist* RNA. The reader protein YTHDC1 has been shown to recognize m^{6}A in

the UCG tetra-loop of Repeat A in the *Xist* RNA and to be required for *Xist* mediated gene

silencing (3,15). The presence of m^{6}A modification is also detected in non-canonical GGAAU

motif in HSATIII lncRNA and has been shown to be critical for the sequestration of splicing

factors in nuclear stress bodies (16). Wide-spread m^{6}A modification is also detected in several

chromatin-bound RNAs including enhancer RNAs, and promoter-associated RNAs (17). m^{6}A

modification of enhancer RNAs has been shown to be required for YTHDC1 and BRD4

mediated transcriptional activation of neighboring genes (18). But how m^{6}A modification

influences the chromatin localization of lncRNAs to the specific genomic locus is not well

understood.

TERRA (telomeric repeat-containing RNA) is a lncRNA derived from the subtelomeric ends

that contain telomeric repeats in the mature transcript (Feretzaki et al., 2020). TERRA

promoters can be located within 1-2 kilobases (kb) from telomere ends to several kb away from

the telomere end. TERRA lncRNAs are transcribed by RNA polymerase II (RNA Pol II) that

moves towards the telomere ends and thus incorporates UUAGGG containing telomeric repeats

(19,20). TERRA RNA is shown to be localized to telomere both in *Cis* and *trans*-acting manner

and RAD51/BRCA1 driven R-loop formation is required for this process (21). TERRA

transcription is upregulated in tumor cells positive for alternative lengthening of telomeres

(ALT+) and is thought to contribute to the maintenance of long telomeres. It is unknown what

additional factors are involved in R-loop mediated TERRA targeting in ALT+ cells.

Understanding chromatin targeting of TERRA RNA might shed light on the mechanism

utilized by other *cis* and *trans*-acting lncRNA for their chromatin recruitment. As R-loop

formation can be regulated in m^{6}A dependent manner, we wanted to explore if m^{6}A

modification has any role in TERRA function. The telomere capping protein TRF2 has also

been found to be interacting with the m^{6}A related proteins (RBM15 and ALKBH5) but the

significance of this interaction is unknown (22,23). Recently it has been shown that G-

quadruplex derived from TERRA repeats and other G-quadruplex forming sequences has been

shown to bind METTL14, but if such sequences have any role to drive m^6A modification in

TERRA RNA is not known (24)

NB is the most common extracranial solid tumor in children, and it is highly heterogeneous in

its clinical and biological nature. The presence of telomere maintenance mechanisms in NB

has been proposed to in part explain the heterogeneity of NB tumors and disease outcomes.

The presence of telomere maintenance mechanisms are strongly associated with high-risk NB.

By contrast, telomere maintenance mechanisms are absent in spontaneously regressing low-

risk tumors. In the majority of high-risk cases, telomere maintenance is conferred by

telomerase, which is induced transcriptionally, e.g., through the oncogene MYCN, or by

genomic translocations leading to enhancer hijacking events (25). In roughly one-third of high-

risk NB, however, telomeres are maintained by the alternative lengthening of telomeres (ALT)

pathway, which is characterized by long telomeres and ALT-associated promyelocytic

leukemia nuclear bodies (APBs) (26,27). The molecular mechanisms underlying ALT are still

poorly understood, despite advancements over recent years (28). This suggests a deeper

understanding of the biology of ALT+ tumors and systematic investigation of the molecular

mechanism of ALT may help us to identify potential therapeutic targets for these clinically

unfavorable high-risk NB tumors with the significant need for new better treatment options.

In the present study, using well-characterized ALT+ U-2 OS cells we have dissected the role

of m^6A RNA modification in the TERRA RNA and characterized the significance of such

modification in R-loop formation. We also show that METTL3 mediated m^6A modification of

the TERRA is essential for telomere maintenance in ALT+ NB cells and inhibiting METTL3

could be a therapeutic option for ALT+ NB tumor

Results:

TERRA RNA is enriched with m^6A modification

To check if TERRA RNA is m^6A modified or not, we have performed m^6A RNA Immuno-

precipitation (m^6A RIP) with m^6A antibody in ALT+ U-2 OS cells prior slot blot with a probe

against telomeric repeats. Through this we observed the enrichment of TERRA RNA and that

m^6A enrichment of TERRA RNA was sensitive to RNase A treatment (**Figure 1A**). To

investigate if m^6A modification of is located in telomeric repeats or elsewhere in TERRA

transcript, we have performed m^6A RIP with both fragmented and non-fragmented RNA under

the assumption that if m^6A modifications are present in telomeric repeat sequences, extensive

fragmentation (<150 nt) will have a low impact in TERRA m^{6}A enrichment when detected

with a probe against telomeric repeats in the slot blot assay. On the other hand, if major m^{6}A

modification is located in non-repetitive regions of TERRA RNA, m^{6}A enrichment will have

a more profound effect after fragmentation. As no major difference in TERRA enrichment was

seen between fragmented and non-fragmented RNA, this indicated that the major fraction of

m^{6}A modifications in TERRA RNA is located in telomeric repeat sequences (**Figure 1B, C,**

and Supplementary Figure S1A). TERRA m^{6}A enrichment was compromised in METTL3-

KD cells in both the non-fragmented and fragmented RNA (**Figure 1C**).

To confirm the presence of m^{6}A in TERRA repeats we purified TERRA from total RNA using

biotin-labeled probe antisense to TERRA along with negative control (probe against Luciferase

RNA) (**Figure 1D**). We verified TERRA enrichment only with the TERRA antisense probe

but not with the negative control probe and this enrichment was sensitive to RNase A treatment

(**Figure 1E**). Further, purified TERRA RNA along with synthesized RNA oligo standard

containing TERRA repeats with or without m^{6}A modification were digested with ribonuclease

T1 (RNase T1) and were subjected to RNA mass spectrometry (RNA-MS) analysis (**Figure**

1D and F). Using the RNA-MS and higher-energy collisional dissociation (HCD) tandem MS

(MS2) data we could confirm the presence of UUAGp and UUm^{6}AGp fragments in our oligo

standards (**Figure 1F**). Utilizing this MS2 information from the standards, we could identify

the UUAGp tetramers in the complex mixture of other RNA fragments generated by RNase T1

cleavage. HCD MS2 analysis further confirmed m^{6}A modification in the UUAGp tetramer

(UUm^{6}AGp) and approximately 21% of UUAGp fragments were m^{6}A modified (**Figure 1F**).

To validate further if the telomeric repeat sequence in TERRA could be m^{6}A modified, we

have performed an *in vitro* m^{6}A methyltransferase assay using a purified METTL3/METTL14

enzyme complex. As a substrate, we have used either synthesized TERRA oligo (UUAGGG)$_3$

or DRACH oligo (GGACU)$_3$ followed by treatment with Nuclease P1 and liquid

chromatography-tandem mass spectrometry (LC-MS/MS)LC/MS analysis to detect A/m^{6}A

(**Supplementary Figure S1B**). Synthesized standard oligo with or without m^{6}A were digested

parallelly with Nuclease P1 and included in the LC-MS/MS. Employing the parallel reaction-

monitoring (PRM) mode, we screened LC-MS/MS profiles for A and m^{6}A which corresponds

to m/z 268.0–136.0 and m/z 282.0–150.1 respectively (29) (**Supplementary Figure S1B**). We

observed *in vitro* m^{6}A modification of both TERRA and DRACH oligo by

METTL3/METTL14 complex (**Supplementary Figure S1B**). Although the efficiency of m^{6}A

modification in TERRA oligo was less compared to the canonical "DRACH" oligo as detected

by the A/m^6A ratio (**Supplementary Figure S1B**), nevertheless our data support

METTL3/METTL14 complex can modify telomeric repeats in TERRA RNA**.**

We checked if METTL3/METT14 complex is localized over the telomeres in ALT+ U-2 OS

cells. We performed a proximity ligation assay (PLA) using TRF2 in combination with either

METTL3 or METTL14 (**Figure 1G**). We observed an overlap of METTL3/TRF2 and

METTL14/TRF2 PLA signal with telomeres detected by TRF1-mCherry. The negative control

PLA with a single antibody did not give any TRF1-mCherry overlapping signal (**Figure 1G**

Supplementary Figure S1C). Using Chromatin Immuno-precipitation (ChIP) we also

observed that METTL3 is recruited to telomere chromatin in U-2 OS cells (**Supplementary**

Figure S1D) suggesting METTL3 is the m^6A methyltransferase for TERRA RNA.

We performed RNA *in situ* hybridization to validate the TERRA RNA localization at telomeres

and we observed TERRA foci frequently overlapped with the TRF1-mCherry signal (**Figure**

1H). We have also verified the specificity of TERRA RNA foci using locked-nucleic acid

(LNA) mediated KD of TERRA RNA (**Supplementary Figure S1E**). Using a modified RNA-

situ hybridization protocol which reduces the background in m^6A immuno-staining (30) we

next checked if TERRA foci overlapped with m^6A, and we found a frequent overlap of m^6A

and TERRA foci further suggesting the presence of m^6A modification on TERRA RNA

(**Figure 1I**).

To explore further TERRA expression and m^6A modification we have performed short

(Illumina) and long reads (ONT: Oxford Nanopore Technologies) RNA-seq and m^6A RIP-seq

in Control-sh and METTL3-KD U-2 OS cells (**Figure 2A**). To better map sequencing reads to

telomere ends, we utilized the latest Telomere to telomere (T2T) human genome assembly (31)

for our RNA-seq data analysis specifically focusing on subtelomeric regions. Combine with

strand-specific Illumina short reads and ONT long-reads we could identify transcriptionally

active and inactive subtelomeres from both p and q arms of chromosomes (**Figure 2B**). As

TERRA RNAs are transcribed in the direction towards the telomere ends, consistent with this

strand-specific Illumina short reads were mainly derived from one of the strands depending on

from which chromosome arms (p and q) they are originating (**Figure 2B and Supplementary**

Figure 2A). By remapping available CTCF ChIP-seq data to T2T assembly, we observed both

proximal and distal CpG islands located over subtelomere, are flanked by CTCF sites as

reported earlier (**Figure 2B**) (32). Using available RNA Pol II ChIP-seq data we also find

occupancy of RNA Pol II at actively TERRA transcribing chromosome ends (**Figure 2B**).

Considering the repetitive nature of chromosome ends we employed mappability normalization

to RNA-seq data. Mappability normalized short and long reads RNA-seq data was used to

define high and low TERRA expressing subtelomeres using the reads covering 30 kb (distance

from the last consecutive Telomeric repeats) of subtelomeric regions which include both

proximal and distal CpG islands. We observed consistent activity of high TERRA expressing

subtelomere in both short and long-read RNA-seq datasets (**Figure 2C, Supplementary**

Figure S2B and C). Though METTL3-KD shows a trend of decreased TERRA RNA level

compared to Control-sh cells these differences were not statistically significant

(**Supplementary Figure S2B and C**). However, we observed a minor (~15%) but significant

decrease in TERRA expression in METTL3-KD cells compared to control-sh using the slot

blot assay (**Supplementary Figure S2D**).

Once we establish the subtelomeric TERRA expression profile in U-2 OS cells we wanted to

check if the telomeric repeats and subtelomeric sequences are enriched with m^{6}A

modifications. We employed a few changes in the m^{6}A RIP-seq protocol so that we could

profile m^{6}A with less amount of RNA (33) while preserving the strand information of the

RNA-seq data. We also used bacterial RNA as a spike-in control in our m^{6}A RIP-seq protocol

which allowed us to accurately normalize the data (**Figure 2A**). The m^{6}A distribution from

RIP-seq performed on U-2 OS nuclear RNA was very similar as described earlier (5), with

major m^{6}A peaks at 3$^\prime$UTR and enrichment of "DRACH" motif (**Supplementary Figure S2E**

and F). In the m^{6}A RIP-seq data from METTL3-KD cells, we observed an overall decrease in

m^{6}A peaks compared to the control suggesting the specificity of the m^{6}A RIP-seq

(**Supplementary Figure S2G**).

We have quantified the amount of telomeric repeat reads present in m^{6}A RIP-seq data and we

found higher enrichment of telomeric repeat reads in Control-sh compared to METTL3-KD

and IgG control (**Supplementary Figure S2H**). Further, we observed that TERRA RNA from

several subtelomeric regions is enriched with m^{6}A modifications (m^{6}A/input ratio) and this

enrichment was affected in METTL3-KD cells (**Figure 2D**). We detected m^{6}A peaks in the

subtelomeres and we see an overall decrease in subtelomere-associated peaks in METTL3-KD

cells (**Figure 2C**). To visualize m^{6}A enrichment across the several active and inactive

subtelomere together, we plotted mappability and spike-in normalized m^{6}A RIP-seq signal over

input (log$_2$ m^{6}A/input) obtained from Control-sh and METTL3-KD cells over the 30kb

subtelomeric regions as metagene plots. We observed m^{6}A enrichment over subtelomere in

Control-sh cell and this was significantly decreased after METTL3-KD (**Figure 2E and F**).

A recent report suggests that "DRACH" like sequences in TERRA can also contribute to m^{6}A

modification (34). We observe the presence of the DRACH sequence in the subtelomeric m^{6}A

peaks and m^{6}A enrichment (log$_2$ m^{6}A/input) over the DRACH sequence was modestly but

significantly decreased in METTL3-KD compared to Control-sh (**Supplementary Figure**

S2J).

We also used long-read RNA-seq data from Control-sh and METTL3-KD cells to identify

differentially modified m^{6}A sites. Among the 27125 significantly modified NNANN sites

identified by xPore (35) genome-wide, 26952 could be successfully translated to annotated

gene coordinates. The differentially modified "m^{6}A" sites follow similar 3′UTR enrichment

like m^{6}A RIP-seq peaks with considerable overlap between the two m^{6}A detection methods

(**Supplementary Figure S2E and I**). We also observed the enrichment of the "DRACH" like

motif in the long-read detected m^{6}A sites as well (**Supplementary Figure 2F**). Next, we refine

the search for possible m^{6}A sites at chromosome ends especially focusing on the non-repeated

part of subtelomeres given inherent base calling errors in Nanopore long-read sequencing at

telomeric repeats (36). To this end, we performed a transcriptome assembly based on the long-

reads mapped to the T2T reference genome and identified 92 NNANN sites that were

differentially m^{6}A modified between Control-sh and METTL3-KD and are located within the

last 30 kb regions of subtelomere ends (as defined above) with 6 of them having a DRACH

motif (Supplementary **Figure S2K**). Subtelomere with higher TERRA expression also contains

more differentially detected m^{6}A sites in ONT reads compared to low expressed ones and these

sites frequently overlap with m^{6}A RIP-seq detected peaks (**Figure 2C and D**).

Collectively, we demonstrate that TERRA RNA undergoes m^{6}A modification in both telomeric

repeats and subtelomeric non-repeated regions which frequently contain a "DRACH" like

motif.

METTL3 mediated m^{6}A modification of TERRA RNA is required for telomere

maintenance in ALT+ cells

We next wanted to explore the importance of m^{6}A modification on TERRA RNA. As evident

from the results above, TERRA RNA forms foci over telomeres in the U-2 OS cells and these

foci are sensitive to TERRA KD (**Figure 1H and Supplementary Figure S1E**). We first tested

the role of m^{6}A modification on TERRA RNA foci formation by performing RNA FISH on U-

2 OS cells that had transient KD (siRNA) of METTL3. We observed that TERRA RNA foci

formation was compromised in the METTL3 siRNA-treated cells compared to control siRNA

treatment (**Figure 3A, Supplementary Figure S3A**). As we observed METT3/METTL14 co-

localize over ALT telomeres (**Figure 1G**), so we also tested the effect of siRNA mediated

depletion of METTL14 on TERRA foci formation. Similar to METTL3, METTL14 depletion

by siRNA leads to compromised TERRA foci formation (**Supplementary Figure S3B**). The

loss of TERRA RNA foci was further validated in stable KD of METTL3 in the U-2 OS cells

(**Figure 3B**). TERRA RNA is thought to contribute to the maintenance of telomeres, therefore

we sought to address the significance of TERRA RNA m^{6}A modification on telomere

maintenance. To this end, we performed immuno-fluorescence staining with γ-H2AX, a DNA

damage response marker, and TRF2 (to mark the telomere) in control and METTL3-KD cells.

We observed that METTL3-KD leads to a higher accumulation of γ-H2AX over telomere

compared to control cells, indicating higher telomere damage in cells lacking METTL3 (**Figure**

3C). Higher γ-H2AX deposition over telomere was also noticed in the TERRA depleted cells

as well (**Supplementary Figure S3C**), suggesting TERRA is required for telomere

maintenance in ALT telomeres. Consistent with telomere damage in METTL3-KD cells we

also observed an increase in TRF2 overlapping Promyelocytic leukemia (PML) bodies

(**Supplementary Figure S3D**).

Recent publications have elucidated the role of METTL3 mediated m^{6}A modification in DNA

damage response (37,38). We reasoned if we create telomere DNA damage exogenously using

bleomycin (a well-characterized DNA damaging agent) in our U-2 OS cells (39), it will allow

us to dissect the role of TERRA m^{6}A modification in such repair processes. We treated control

U-2 OS cells with bleomycin and detected DNA damage by the accumulation of γ-H2AX over

the telomere within 4 hours of treatment. Interestingly if we allow the U-2 OS cells to recover

(20 hours) post bleomycin treatment, γ-H2AX accumulation was significantly reduced over

telomere (**Figure 3D**). However, in the METTL3-KD cells, this accumulation of γ-H2AX did

not cease, but rather increased, even after cells were off media containing bleomycin for 20

hours (**Figure 3D**). We observed a higher number of m^{6}A overlapping TERRA foci, during

recovery post bleomycin treatment in U-2 OS cells suggesting the role of TERRA m^{6}A in

telomere damage repair (**Figure 3E**).

To elucidate the importance of m^{6}A modification on TERRA RNA, we specifically removed

m^{6}A modification from TERRA RNA using RNA-guided CRISPR/dCasRx system (40).

Catalytically dead RNA CRISPR/dCasRx fused with m^{6}A demethylase- FTO (dCasRx-FTO)

was recruited to TERRA RNA to specifically remove m^{6}A from TERRA repeats in U-2 OS

cells (41) (**Figure 3F**). We verified the removal of m^{6}A from TERRA RNA by performing

m^{6}A RIP followed by TERRA slot blot, as evident from the blot, recruiting of dCasRx-FTO to

TERRA resulted in a lower enrichment of m^{6}A over TERRA RNA compared to non-template

control (NTC) (**Figure 3G**). Recruitment of dCasRx-FTO by TERRA gRNA did not result in

any change in TERRA RNA (**Supplementary Figure S3E**). We next evaluated the effect of

this specific removal of m^6A on the ability of TERRA RNA foci formation by performing

TERRA RNA-FISH. We observed a loss of TERRA foci in cells where dCasRx-FTO was

recruited to TERRA (**Figure 3H**). To test the specificity of guide RNA, instead of recruiting

dCasRx-FTO we deployed catalytically active CasRx and checked if TERRA RNA was

downregulated. As expected, we saw lower levels of TERRA RNA in cells where CasRX was

recruited at TERRA RNA compared to NTC (**Supplementary Figure S3F**).

Owing to the growing importance of METTL3 mediated m^6A modification in diseases, in

recent years two independent studies developed METTL3 inhibitors (42,43). In particular,

Yankova et al. extensively tested a small molecule METTL3 inhibitor in various AML mouse

models and concluded that METTL3 inhibition is a potential therapeutic strategy in AML (42).

We wanted to test the effect of these available METTL3 inhibitors on TERRA RNA foci

formation. Treatment with Actinomycin D (ActD) which blocks RNA Pol II thereby the

synthesis of new RNAs, suggests one hour of ActD treatment leads to a major depletion of

existing TERRA (**Supplementary Figure S3G**). Keeping the TERRA half-life in

consideration obtained from ActD data (**Supplementary Figure 3G**), we treated U-2 OS cells

for 6 hours with two METTL3 inhibitors (STM2457 and UZH1A), which will lead to

compromised deposition of m^6A in the newly formed TERRA molecules. We observed a

pronounced reduction in TERRA RNA foci following 6 hours of METTL3 inhibition compared

to DMSO control (**Figure 2I**). Whereas treatment with inefficient METTL3 inhibitor (UZH1B)

(43) at the same time had a relatively less pronounced effect on TERRA foci (**Figure 3I**).

Treatment with METTL3 inhibitor STM2457 which showed the highest effect on TERRA foci

did not alter TERRA RNA level as compared to DMSO control (**Supplementary Figure S3H**).

By modulating the m^6A using either the dCasRx-FTO system or by METTL3 inhibitor

treatment, we unravel the critical role of m^6A modification in the functionality of TERRA

RNA. Our data collectively suggest that METTL3 mediated m^6A modification of TERRA is

vital for TERRA RNA foci formation and m^6A modification in TERRA RNA is required for

the repair of the damaged telomeres.

hnRNPA2B1 binds TERRA RNA in m^6A dependent manner and is required for TERRA

function

Our result so far suggests that METTL3 mediated m^6A modification of TERRA RNA is

required for TERRA foci formation. Next, we wanted to evaluate if m^6A reader proteins are

involved in this process. We performed a small siRNA screen to knock down known m^6A

reader proteins such as YTHDC1, hnRNPA2B1, RBMX (Nuclear m^6A readers), and YTHDF2

(cytoplasmic reader) (**Figure 4A, Supplementary Figure S4A**). Among these hnRNPA2B1 and YTHDC1 have been shown to interact with TERRA RNA before (34,44). We observed that out of all m^{6}A reader proteins tested hnRNPA2B1 KD had a major effect on TERRA foci formation (**Figure 4A**). Though we observed the loss of TERRA RNA foci in hnRNPA2B1 KD cells but TERRA level was unchanged (**Supplementary Figure S4B**). We could also validate hnRNPA2B1 interaction with TERRA RNA using RNA Immunoprecipitation (RIP) followed by slot blot (**Figure 4B**). Next, we asked how a change in TERRA m^{6}A level contributes to hnRNPA2B1 interaction. To this end, we performed hnRNPA2B1 RIP in Control-sh and METTL3-KD cells followed by TERRA slot blot. We observed reduced hnRNPA2B1 interaction with TERRA in METTL3-KD cells, suggesting a role of m^{6}A modification in TERRA and hnRNPA2B1 interaction (**Figure 4C**). To check further if hnRNPA2B1 enriched TERRA was m^{6}A modified or not, we performed m^{6}A RIP using hnRNPA2B1 bound RNA and observed hnRNPA2B1 interacting TERRA RNA was modified with m^{6}A (**Figure 4C**). Consistent with our RIP experiment purified hnRNPA2B1 interacts efficiently with TERRA RNA derived from control U-2 OS cells but with reduced efficiency with TERRA derived from METTL3-KD cells (**Supplementary Figure S4C**). Finally, when we specifically demethylate TERRA RNA by deploying dCasRx-FTO, we see reduced occupancy of hnRNPA2B1 at telomere (**Figure 4D**). To know if hnRNPA2B1 KD phenocopy the increased DNA damage phenotype at telomers as observed in METTL3-KD cells, we immuno-stained γ-H2AX and TRF2 in U-2 OS cells following hnRNPA2B1 KD. KD of hnRNPA2B1 resulted in increased accumulation of γ-H2AX at telomeres (**Figure 4E**). Our results demonstrate that the binding of hnRNPA2B1 and m^{6}A co-operate with each other for TERRA dependent maintenance of telomere in ALT+ cells.

TERRA m^{6}A modification promotes R-loop formation

Recent studies reported that m^{6}A modification is present in the R-loop and might contribute to R-loop formation (11,12). R-loop formation by TERRA is also critical for targeting TERRA to the telomere (21). We observed lack of m^{6}A in TERRA RNA leads to a decrease in TERRA RNA foci and this could be attributed to compromised telomere targeting of TERRA RNA. We reasoned that a decrease in TERRA R-loop formation in the absence of m^{6}A might lead to inefficient targeting of TERRA to the telomere. We modified the published DNA-RNA immunoprecipitation (DRIP) protocol, where anti-RNA:DNA hybrid S9.6 antibody is used to identify RNA strand of R-loops, keeping two goals; (i) to measure accurately TERRA RNA

present in the R-loop structures and (ii) to identify if TERRA RNA in R-loop context is m^{6}A

modified. To obtain specific enrichment of R-loop associated TERRA RNA we treated the cell

lysate with RNase A briefly to remove free RNA from the subsequent genomic DNA isolation

step (**Figure 5A**). We also treated isolated genomic DNA before S9.6 IP with RNase III which

is shown to remove non-specific enrichment of double-stranded RNA fragments to identify

genuine R-loop enriched RNAs (**Figure 5A**)(45). We pre-treated genomic DNA with RNase

H to cross-check the specificity of S9.6 IP as a control (**Figure 5A**). Following S9.6 IP we

performed extensive DNase I treatment to obtain R-loop associated RNA (**Figure 5A**). We

further used R-loop associated RNA for slot blot assay and we could successfully detect

enrichment of R-loop associated TERRA (hereafter referred to as RL-TERRA) (**Figure 5B**).

RL-TERRA enrichment was decreased following RNase H treatment, and no enrichment was

observed with IgG control (used as negative control) suggesting the specificity of the obtained

RL-TERRA using our modified DRIP method. Enriched RL-TERRA was sensitive to RNase

A treatment as well (**Figure 5B**). Next, we used R-loop associated RNA obtained from our

modified DRIP method (as described above) as input for m^{6}A RIP, we refer to this complete

protocol as DRImR (**DR**IP followed by **m**^{6}A **R**IP) (**Figure 5A and B**). DRImR allows us to

detect m^{6}A modification present in R-loop structures. We observed that m^{6}A modified TERRA

was abundantly present in DRImR enriched RNA and this enrichment was sensitive to RNase

H treatment showing RL-TERRA is m^{6}A modified (**Figure 5B**). Further, METTL3-KD leads

to compromise RL-TERRA enrichment in U-2 OS cells suggesting TERRA m^{6}A contributes

to R-loop formation (**Figure 5C**).

To check the effect of R-loop removal in TERRA foci formation, we treated U-2 OS cells with

either RNase A alone or in combination with RNase H. We observed only RNase A had a

moderate effect on TERRA RNA foci in U-2 OS cells but combining RNase A and RNase H

led to the disappearance of TERRA foci suggesting the importance of R-loops in TERRA foci

formation (**Supplementary Figure S5A**). We reasoned if lack of TERRA m^{6}A in METTL3-

KD cells leads to compromised R-loop thereby disturbing TERRA foci formation, and whether

this can be rescued by removing RNase H which will allow R-loop accumulation. We could

rescue TERRA foci in METTL3-KD cells by further removing RNase H transiently with

siRNA, suggesting the role of m^{6}A in the R-loop driven TERRA foci formation (**Figure 5D,**

Supplementary Figure S5B).

Next, we went ahead to standardize a robust and fast PLA based method to check telomere R-

loop in different experimental conditions where we observed TERRA foci were compromised.

We used PLA assay with TRF2 and S9.6 antibodies (R-loop PLA) to determine the presence

of R-loop over telomere. Telomeres were marked by TRF1-mCherry to further confirm the

presence of R-loop PLA signal over telomere. We used wild-type RNase H (RNase H^{WT}) and

catalytically inactive mutant RNase H (RNase H^{Mut}) cells as a control in our experiment. As

expected, overexpression of RNase H^{WT} leads to a decrease in overall R-loop content

(**Supplementary Figure S5C**). RNase H^{WT} cells also show a decreased number of TERRA

RNA foci compared to RNase H^{Mut} (**Supplementary S5D**). We observe an overlap of R-loop

PLA signal over telomere in the cell expressing RNase H^{Mut} but this signal was considerably

decreased in RNase H^{WT} cells suggesting the specificity of our assay in detecting R-loop over

telomere (**Figure 5E**). We also did not detect any telomere overlapping PLA signal in single

antibody control of S9.6 and anti-TRF2 antibody (**Supplementary Figure S5E**).

We applied R-loop PLA to Control-sh and METTL3-KD cells and observed loss of PLA signal

over telomere in METTL3-KD cells and this observation is consistent with data obtained using

DRImR (**Figure 5C and F**). Further, the removal of TERRA m^{6}A using dCasRx-FTO and KD

of TERRA m^{6}A reader protein (hnRNPA2B1) also led to a loss of R-loop PLA signal

(**Supplementary S5F, Figure 5G**) suggesting TERRA m^{6}A and its reader hnRNPA2B1 is

required for R-loop formation over telomeres. Overall, we provide evidence that m^{6}A

modification of TERRA RNA is required for TERRA R-loop formation thereby TERRA

targeting to the telomere**.**

In ALT+ NB cells TERRA RNA is m^{6}A modified

A significant portion of the NB tumors are ALT+ and the prognosis of these patients is poor

and a recent study found that relapsed NB tumors are frequently ALT+ (26,27,46,47). The role

of TERRA RNA in ALT+ NB tumors has not been investigated in depth. We aim to

characterize if m^{6}A modification of TERRA plays a critical role in telomere maintenance in

ALT+ NB. First, we checked TERRA expression in ALT+ and MYCN amplified (ALT-

negative: ALT-) NB cell lines. TERRA expression was higher in ALT+ (CHLA-90 and SK-N-

F1) compared to ALT- (SK-N-BE(2) and IMR-32) NB cell lines (**Figure 6A**). ALT positivity

of NB cell lines was verified by C-circle assay (**Supplementary Figure S6A**). We performed

m^{6}A RIP in fragmented RNA from ALT+ and ALT- NB cell lines and we observed robust

enrichment of TERRA with m^{6}A antibody in ALT+ cells (**Figure 6B**). We also observe

prominent TERRA foci in ALT+ but not in ALT- NB cell lines (**Figure 6C**), probably higher

TERRA expression and m^{6}A enrichment in ALT+ cells contribute to TERRA foci formation.

TERRA foci in ALT+ NB cells are sensitive to LNA mediated KD and frequently overlap with

m^{6}A immuno-fluorescence signal in the nucleus (**Supplementary Figure S6B and Figure**

6D). To characterize TERRA expression and m^{6}A enrichment in ALT+ (SK-N-F1) and ALT-(SK-N-BE(2)) NB cells we performed RNA-seq and m^{6}A RIP-seq. We observed TERRA expression was low in ALT- cells compare to ALT+ and this is consistent with slot blot assay (**Figure 6A, Supplementary Figure S6C).** We observed enrichment of the "DRACH" motif in the m^{6}A peaks from both SK-N-F1 and SK-N-BE(2) m^{6}A RIP-seq data (**Supplementary Figure S6D**). Then we went ahead to check subtelomeric m^{6}A enrichment in these two cell lines, mappability and spike-in normalized subtelomeric m^{6}A enrichment (log$_2$ m^{6}A/input) was significantly higher in ALT+ SK-N-F1 cells compare to ALT- SK-N-BE(2) cells (**Figure 6E, F**).

To get a better view of TERRA expression in NB tumor samples we have compared TERRA expression in representative ALT+ and ALT- tumors for which we had access to RNA. Two of the NB tumors were previously reported by us as ALT+ (26) and for the rest of the used NB tumors, we verified ALT status by C-circle assay (**Supplementary Figure S6E**). We observed elevated levels of TERRA expression in ALT+ tumors and this is consistent with earlier observation (**Figure 6G**)(47). Next, we wanted to characterize TERRA expression and m^{6}A positivity using RNA-seq and m^{6}A RIP-seq in two ALT+ NB tumors (NB1 and NB2) which showed high TERRA expression in slot blot (**Figure 6G**). RNA-seq data from ALT+ NB tumors were used to define high and low TERRA expressing subtelomere in a similar way to U-2 OS cells. We observed consistent transcriptional activity from the subtelomeres in both tumors whereas others are silent or moderately transcribed (**Figure 6H, Supplementary Figure S6F**). Overall, actively TERRA transcribing subtelomeres are also similar in both NB tumors and U-2 OS cell lines suggesting the commonality in TERRA transcription (**Figure 6I and 2C**). We observed that "DRACH" like motifs are enriched in m^{6}A peaks obtained from both NB tumors (**Supplementary Figure S6G**). TERRA RNAs from several active subtelomeric regions are enriched with m^{6}A modifications in NB tumors and frequently contain m^{6}A peaks as well (**Figure 6I, J**). The number of m^{6}A peaks in NB1 was more compared to NB2 in subtelomere which could be contributed by tumor-specific variations (**Figure 6I**). Subtelomeric metagene profile of mappability and spike-in normalized m^{6}A enrichment (log$_2$ m^{6}A/input) suggests high TERRA expressing subtelomeres were more robustly m^{6}A modified compared to subtelomeres which were lowly expressed in both the ALT+ NB tumors (**Figure 6K**).

We find higher METTL3, METTL14, and hnRNPA2B1 expression in the NB tumor cohort (n=498) significantly associated with poor event free survival (EFS) (**Figure 6L** and

Supplementary Figure S6H). In a small cohort (n=21) of ALT+ NB tumors, we observed

higher METTL3 and METTL14 expression significantly correlated with worst EFS and for

hnRNPA2B1 though higher expression tend to show worst EFS, but it did not reach

significance (**Figure 6M** and **Supplementary Figure S6I**). We also observed frequent copy

number gain of METTL3, METTL14, and TERRA m^6A reader hnRNPA2B1 in ALT+ tumors

compared to MYCN amplified NB tumors (**Figure 6N**) suggesting the vital role of METTL3

and other m^6A related genes in NB.

m^6A modification in TERRA RNA is critical for telomere maintenance in ALT+ NB cells

To explore the role of METTL3 mediated m^6A modification in ALT+ NB, we depleted

METTL3 in ALT+ NB cells. We observed siRNA mediated transient removal of METTL3 in

ALT+ CHLA-90 and SK-N-F1 cells led to a reduction in the number of TERRA foci similar

to U-2 OS cells (**Figure 7A**). We next wanted to check the effect of stable depletion of

METTL3 in SK-N-F1 and CHLA-90 cells, but we failed to generate stable METTL3 depleted

cells after several attempts as these cells did not survive suggesting METTL3 was essential for

the survival of ALT+ NB cells. So, we went ahead to make doxycycline (Dox) inducible TetO

METTL3-KD cells (**Supplementary Figure S7A**), and consistent with the observation

following stable METTL3 depletion Dox inducible ALT+ cells show a decrease in cell viability

and reduction in colony formation compared to TetO Control cells (**Supplementary Figure**

S7B, Figure 7B). We observed TERRA foci loss in TetO METTL3-KD SK-N-F1 cells (**Figure**

7C) but TERRA expression was unchanged as visualized in the slot blot assay

(**Supplementary Figure S7C**). Dox-induced METTL3 depletion also induces telomere

damage in both SK-N-F1 and CHLA-90 cells with a frequent overlap of γ-H2AX with

telomeres (**Figure 7D and Supplementary Figure S7D**). METTL3 depletion also leads to a

decrease in C-circle formation in these ALT+ NB cells suggesting compromised ALT activity

(**Supplementary Figure 7E**). METTL3 depletion in ALT+ NB cells also leads to a decrease

in RL-TERRA enrichment like in U-2 OS cells, suggesting compromised TERRA R-loop

formation (**Supplementary Figure S7F**). We xenografted TetO Control and METTL3-KD

SK-N-F1 and CHLA-90 cells in immuno-compromised mice and we followed the effect on

tumor formation following Dox induction. We observed that METTL3-KD leads to reduced

xenografted tumor growth in both ALT+ cell lines with a more drastic effect in SK-N-F1 cells

(**Figure 7E and Supplementary Figure S7G**).

To test if TERRA foci formation in ALT+ NB cells is hnRNPA2B1 dependent, we have

depleted hnRNPA2B1, and this led to the loss of TERRA foci (**Figure 7F, G**). We also

observed TERRA foci are co-localized with hnRNPA2B1 in ALT+ NB cells (**Supplementary**

Figure S7H). Finally, we wanted to explore the therapeutic potential of inhibiting the METTL3

enzyme in ALT+ NB cell lines. We observed treatment of ALT+ NB cells with METTL3

inhibitors STM2457, and UZH1A for 6 hours led to a dramatic decrease in TERRA foci

formation compared to cells treated with UZH1B (less potent inhibitor) and DMSO control

(**Figure 7H**). We also observed 48 hours of treatment of STM2457 in ALT+ SK-N-F1 cells

leads to damaged telomeres which frequently overlapped with γ-H2AX (**Figure 7I**) but

STM2457 treatment did not induce any γ-H2AX accumulation over telomeres in MYCN

amplified SK-N-BE(2) cells (**Supplementary Figure S7I**). Treatment with STM2457 results

in reduced cell viability in SK-N-F1 and CHLA-90 cells (**Supplementary Figure S7J**).

Collectively, our data show that TERRA m^{6}A and METTL3 play a vital role in telomere

maintenance in ALT+ NB, and inhibiting METTL3 could be a potential therapeutic target for

ALT+ NB.

Discussion:

Our study has identified the role of m^{6}A RNA modification in the telomere-derived noncoding

RNA, TERRA, which has a role in the maintenance of telomeres in ALT+ cells. Our data

suggest that the METTL3/14 complex is recruited to telomeres in ALT+ U-2 OS cells. Recently

G-quadruplex derived from telomeric repeats has been shown to bind METTL14 (24), so it is

tempting to speculate that TERRA G-quadruplex might guide m^{6}A writer METTL3/14 to the

telomeres. Removal of METTL3 leads to a decrease in the m^{6}A modification in the TERRA

transcript and loss of TERRA RNA foci in the ALT+ cells. Modulating the m^{6}A level by

METTL3-KD leads to compromised repair of telomeric DNA which is visualized by increased

γ-H2AX accumulation over the telomere. Our data suggest that m^{6}A containing TERRA foci

were required for repairing bleomycin induced telomere damage and also that METTL3-KD

cells that lack TERRA m^{6}A failed to repair damaged telomere. In line with our observation, a

recent study also reported the role of TERRA m^{6}A modification in telomere maintenance in

ALT+ cells and METTL3-KD induces telomere shortening as well in ALT+ cells (34). They

also reported the presence of m^{6}A modification in TERRA RNA in the subtelomeric sequences.

Consistent with this, our m^{6}A RIP-seq and m^{6}A sites identified by xPore from long-read RNA-

seq data also show that subtelomere-derived TERRA transcripts are enriched with m^{6}A. Our

study using m^{6}A RIP with fragmented RNA (Figure 1B, C), *in vitro* m^{6}A methyltransferase

assay of telomeric repeats (Supplementary Figure S1B) and TERRA RNA-MS analysis (Figure

1D-F) hinted presence of m^{6}A modification over UUAGGG telomeric repeats in TERRA RNA.

Though the efficiency of m^{6}A deposition by METTL3/METTL14 on the telomeric repeats was

comparatively low to DRACH sequence *in vitro* however, presence of multiple copies of

telomeric repeats in single TERRA molecule could drive effective m^{6}A modification *in vivo*.

This m^{6}A modification of TERRA in the repeats could be further enhanced by G-quadruplex

like structures present in TERRA RNA (24). Modulating the m^{6}A specifically in the telomeric

repeats using dCasRx-FTO recapitulated METTL3-KD phenotype, further suggests the

importance of m^{6}A modification at telomeric repeat sequences in TERRA RNA. TERRA

subtelomeric sequences containing "DRACH" motif were differentially enriched in m^{6}A RIP-

seq data obtained from Control-sh and METTL3-KD cells. The m^{6}A sites detected by xPore

include a "DRACH" motif as well. The long-read RNA-seq data was generated using poly-A

selected RNA, so this data will not possibly include any m^{6}A sites present in a non-poly-A

fraction of TERRA RNA (48). Our m^{6}A RIP-seq and xPore data are consistent with the notion

as reported by Chen et al (34) that "DRACH" like sequences in TERRA could also be targeted

by METTL3 for m^{6}A modification. Taking together, we provide evidence that both telomeric

repeats and subtelomeric sequences contribute to the m^{6}A enrichment of TERRA RNA. Owing

to the complexity, that TERRA RNAs are derived from multiple chromosome ends and the

presence of telomeric repeats not only at the end of the chromosome but also interspersed in

the subtelomeric region (**Figure 2B**), it is challenging to precisely resolve the exact location of

m^{6}A modification in the TERRA RNA.

Our mechanistic study revealed that m^{6}A modification in TERRA promotes the association of

m^{6}A reader hnRNPA2B1. Recently another m^{6}A reader protein YTHDC1 was also shown to

interact with TERRA RNA in m^{6}A dependent manner to regulate TERRA RNA stability (34).

Though we observed a moderate decrease in TERRA RNA in METTL3-KD U-2 OS cells in

the slot blot assay, however, hnRNPA2B1 KD did not alter the TERRA level. TERRA specific

m^{6}A removal by an inducible dCasRx-FTO system in U-2 OS cells did not have any impact on

steady state level of TERRA RNA. Similarly, we saw unaltered TERRA level in inducible KD

of METTL3 in SK-N-F1 cells suggesting that lack of m^{6}A in TERRA might promote TERRA

degradation in cell type specific manner or depending on mode m^{6}A of depletion from TERRA

RNA. Short term treatment (6 hours) with the METTL3 inhibitor was sufficient to induce loss

of TERRA RNA foci without any change in TERRA RNA level, suggesting that m^{6}A

modification has a major role in driving telomere targeting of TERRA. But long-term depletion

of METTL3 leads to a lack of persistent TERRA targeting to telomeres and this might promote

degradation of TERRA RNA, where YTHDC1 mediated stabilization is impaired in the

absence of m^{6}A. Nevertheless, both hnRNPA2B1 and YTHDC1 KD induce telomere damage

in ALT+ cells suggesting an important role of the m^{6}A reader proteins in telomere maintenance

(34).

The m^{6}A reader protein hnRNPA2B1 does not have the YTH domain to identify m^{6}A and has

instead been proposed to identify RNA by a "m^{6}A switch" mechanism (49). It is interesting to

note that the motif which is recognized by hnRNPA2B1 is UAGG like sequences (50) whereas

the major motif for m^{6}A is "DRACH" like sequences (51). UUAGGG containing TERRA

repeats has been shown to bind hnRNPA2B1 *in vitro* and hnRNPA2B1 also recognise UAGGG

containing motifs *in vivo* (52,53). Our *in vitro* and *in vivo* observations suggest m^{6}A

modification in TERRA RNA is critical for hnRNPA2B1 association. We identified the

presence of m^{6}A in the non-canonical UUAGGG context in TERRA. The presence of m^{6}A in

a non-canonical context has also been observed in other lncRNAs such as in HSATIII (AAUG

repeats) and Xist (AUCG context in tetra loop of Repeat-A) lncRNAs (15,16). In case of Xist,

m^{6}A sites in AUCG context in Repeat-A create local unfolding which favors YTHDC1

recognition (15). "M^{6}A switch" mechanism also suggests the presence of m^{6}A creates local

folding in RNA thereby enhancing the association of RNA-binding proteins like hnRNPC and

RBMX to RNA molecules (8,54).

We observed both m^{6}A modification and hnRNPA2B1 interaction are required for TERRA

targeting to telomere by R-loop formation. R-loop associated TERRA (RL-TERRA) was found

to be abundantly m^{6}A modified as detected by DRImR which allows combined detection of

RNA strand of R-loop and m^{6}A modification. It is tempting to speculate that m^{6}A modification

in UUAGGG containing telomeric repeats might promote local unfolding of TERRA RNA

during transcription, resulting in efficient recruitment of hnRNPA2B1 and further other repair

proteins like RAD51 to promote R-loop formation. R-loop formation is essential for telomere

targeting of TERRA after transcription. (21). Chen et al reported compromised RAD51

recruitment to telomeres in METTL3-KD cells and defective RAD51 recruitment might affect

TERRA dependent R-loop formation (21,34). The mode of chromatin targeting of lncRNAs is

still a less understood phenomenon. Increasing evidence suggests that m^{6}A modification in

lncRNAs might promote chromatin association and m^{6}A modified nascent RNAs accumulate

at the site of transcription (55). Our observation from TERRA RNA provides evidence that

m^{6}A mediated R-loop formation might be a general mechanism utilized by other lncRNAs for

chromatin targeting (55). Compromise telomere targeting of TERRA in the absence of m^{6}A

might affect the recruitment of DNA repair proteins like BRCA2 which help in recombination-dependent telomere maintenance in ALT+ cells (21). TERRA has also been shown to frequently interact with epigenetic modifiers which are required for telomere maintenance (44,56). In METTL3-KD cells, lack of telomere targeting in TERRA RNA might results in the altered chromatin structure of ALT+ telomeres thereby resulting in a defect in telomere maintenance as well. But the extent of chromatin changes over telomere in METTL3 depleted/inhibited ALT+ cells require further study.

We observed that ALT+ NB cells are vulnerable to METTL3-KD or METTL3 inhibition. METTL3-KD or inhibition results in the accumulation of telomere damage suggesting METTL3 mediated m^6A modification of TERRA has a very critical role in Telomere maintenance in ALT+ NB cells. TERRA is upregulated in ALT+ NB tumors and ALT+ tumors frequently show a mutation in chromatin remodeler protein ATRX (47). It will be interesting to check if the loss of ATRX has any role in driving TERRA expression and m^6A modifications in ALT+ NB tumors. ALT+ NB tumors show protracted disease phenotype with frequent relapse (47) and higher METTL3 expression correlates with poor disease prognosis in ALT+ NB. Our data on METTL3 inhibitor treatment and telomere damage in ALT+ NB cells suggest inhibiting METTL3 could be a novel therapeutic approach for the unfavorable high-risk subset of ALT+ NB tumors.

Materials and methods:

Cell culture

U-2 OS (Sigma-Aldrich) cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% FBS and penicillin/streptomycin. ALT+ NB cell lines CHLA-90 (CCOG) and SK-N-F1 (Sigma-Aldrich) cells were cultured in DMEM media supplemented with 20% and 10% FBS respectively along with penicillin/streptomycin. SK-N-BE(2) cells were cultured in DMEM/F-12 media supplemented with 10% FBS, 4 mM L-glutamine, and penicillin/streptomycin. Cells used in the study were mycoplasma negative and tested routinely.

Plasmids, siRNAs, and molecular cloning:

METTL3 (sh1 and sh2) and Control shRNAs in pLKO.1-TRC cloning vector were purchased from Sigma-Aldrich to generate stable METTL3-KD cells. shRNA sequences are provided in Supplementary Table 1. To generate Dox induced METTL3-KD cells in ALT+ NB METTL3-sh1 sequences were cloned in the Tet-pLKO-puro vector (21915, Addgene). As a control for inducible shRNA KD, we used pLKO-Tet-On-shRNA-Control plasmid (98398, Addgene).

dCasRx-FTO-HA plasmid was received as a generous gift from Dr. Wenbo Li´s laboratory (University of Texas Health Science Center, Houston, USA) and CasRx (138149) was obtained from Addgene. The TERRA and negative control gRNAs were cloned in pLentiRNAGuide_002 (138151, Addgene). Sequences of guide RNAs are provided in Supplementary Table 1. RNase H WT and mutant overexpressing vectors were purchased from Addgene (111906 and 111905). TRF1-mCherry plasmid was received as a generous gift from Dr. Emilio Cusanelli (University of Trento, Italy). Pre-designed siRNAs were purchased from either Sigma-Aldrich or Thermo Fisher Scientific and details are provided in Supplementary Table 1. Lentiviral packaging plasmids psPAX2 (12260) and pMD2.G (12259) were obtained from Addgene.

Gene silencing and overexpression:

Lentiviral based vectors were packaged following instructions from Addgene using packaging plasmids as described above. To generate stable cells antibiotic selection was performed depending on the type of vector used following lentiviral transduction. To generate METTL3-KD cells U-2 OS and ALT+ NB cell lines were selected with media containing Puromycin (1 µg/ml). In NB cells Doxycycline (200 ng/ml) was used to induce METTL3-KD. To generate CasRx/dCasRx-FTO and gRNA expressing U-2 OS cells, dCasRx-FTO_HA containing lentivirus were transduced followed by selection with Puromycin (1 µg/ml). The puromycin selected cells were further transduced with gRNA lentivirus and selected with Blasticidin (5 µg/ml). To induce CasRx/dCasRx-FTO expression cells were cultured in Doxycycline (200 ng/ml) containing media. To generate stable RNase H WT/Mut cell lines corresponding plasmids were transfected using Lipofectamine 3000 reagent (Thermo Fisher Scientific) following manufacturer instruction followed by selection with media containing Hygromycin (400 µg/ml). siRNAs were transfected to induce transient in both U-2 OS and ALT+ NB cells using RNAiMAX (Thermo Fisher Scientific) reagents following the manufacturer's instructions.

RNA and DNA isolation

From Cell-lines

RNA was isolated from the cell line using TRIzol reagent (15596026, Thermo Fisher Scientific) and Direct-zol RNA Miniprep (R2050, ZYMO research) or using ReliaPrepTM RNA Cell Miniprep system (Z6011, Promega), according to the manufacturer's instruction. The DNase I treatment step was performed mandatorily, and the RNA was eluted in RNase/DNase-free water.

DNA was isolated from cells using the Wizard Genomic DNA purification Kit (Promega)
according to the manufacturer's instructions.

From Tumor material

Tumor tissue was collected after written or verbal informed consent was obtained from
parents/guardians according to ethical permits approved by the local ethics committees
(Karolinska Institute and Karolinska University Hospital Research Ethics Committee,
registration number 2009/1369-31/1 and 03-736). RNA was extracted from frozen tumor
samples using AllPrep DNA/RNA/Protein Mini Kit (Qiagen) according to the manufacturer's
instructions.

Genomic DNA was extracted from fresh/frozen tumors or blood using the DNeasy Blood &
Tissue kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. This study
was conducted per the Declaration of Helsinki.

cDNA synthesis and Quantitative PCR (qPCR

RNA was converted to cDNA using the High-Capacity RNA-to-cDNATM Kit (4387406,
Thermo Fisher Scientific). qPCR was performed with diluted cDNA (10-fold dilution) as a
template and gene-specific PCR primers (Supplementary Table S1) mixed with Power SYBR
Green Master Mix (4367659, Thermo Fisher Scientific) using a Quant Studio 3 thermocycler
(Thermo Fisher Scientific). The Expression values presented for each gene are normalized to
GAPDH using the delta-delta Ct method.

Slot blot assay

Nucleic acids (RNA or DNA) samples were loaded in Bio-Dot SF Microfiltration Apparatus
(Bio-Rad) and allowed to bind onto the membrane. Nucleic acids were UV-crosslinked to the
membrane and incubated with prehybridization buffer DIG Easy Hyb (11603558001, Roche,)
in a hybridization tube at 37° C for 30 min with rotation. Prehybridization buffer was replaced
by hybridization buffer (prehybridization buffer containing 20-50 nM 5′-(TAACCC)$_5$ DIG-
labeled probe) and the membrane was hybridized at 37°C overnight with rotation. After
hybridization, the membrane was washed with 2x SSC containing 01% SDS for 4x10 min at
37°C and equilibrated with washing buffer (0.1 M Maleic acid, 0.15 M NaCl, pH 7.5, 0.3%
Tween 20) for 2 min at RT. Next, the membrane was incubated in a blocking buffer (1% (w /
v) Roche Blocking Reagent, 0.1 M Maleic Acid, 0.15 M NaCl, pH 7.5) for 30 min at RT.
Blocking buffer was replaced with antibody solution (1: 20,000 dilution of Anti-DIG antibody
conjugated with alkaline phosphatase in blocking buffer) for 30 min at RT. The membrane was
washed twice 2x15 min with washing buffer. The membrane was equilibrated in a detection
buffer (0.1 M Tris-HCl, 0.1 M NaCl, pH 9.5) and incubated with CDP-Star Chemiluminescent

Substrate (Sigma-Aldrich) inside a hybridization bag. ChemiDoc XRS+ system (Bio-Rad) and

ImageLab software were used to detect the signal and quantify the bands.

TERRA RNA levels using slot blot assay

To check the level of TERRA RNA in different cell lines, tumors, and conditions, isolated total

RNA from different samples were first quantified on NanoDrop™ 2000 Spectrophotometer

(Thermo Fisher Scientific) and diluted to 30 ng/ul concentration. The diluted RNA was again

quantified on the Nanodrop to verify the concentration. The diluted RNA was then loaded onto

slot blot in amounts as mentioned in the Figures, and 1 µl of the diluted RNA which

corresponds to 30 ng was used as a loading control. The slot blots were probed with a TERRA

probe, developed as mentioned above, and normalized to the 28s rRNA signal from the loading

control. To obtain the 28s rRNA signal the RNA samples were subjected to electrophoresis on

a chip (ScreenTapes) using TapeStation 4200 (Agilent) according to manufactures instruction.

m^6A RIP followed by sequencing (m^6A RIP-seq)/ slot blot

m^6A RIP was performed on RNA as previously described by (33). In brief, for the m^6A RIP-

seq experiments the cellular (15 µg/IP) and tumor RNA (4-5 µg/IP) were supplemented with

10 ng or 3 ng of bacterial RNA respectively as a spike-in control. For m^6A RIP followed by

slot blot bacterial spike-in RNA was not added. The RNA was then subjected to fragmentation

using RNA Fragmentation Reagents (AM8740, Thermo Fisher Scientific). m^6A RIP was also

performed in one instance without the fragmentation step. RNA immunoprecipitation (RIP)

was performed using antibodies against m^6A modification (202003, Synaptic systems) or IgG

(sc-2027, Santa Cruz) in RIP buffer (150 mM NaCl, 10 mM Tris-HCl pH 7.5, 0.1% IGEPAL

CA-630 and supplemented with RNase inhibitor just before use) for 4 h at 4°C. IgG antibodies

served as a negative control for the immunoprecipitation experiments. The immunoprecipitated

RNA-antibody complex was captured by incubating with protein A and G magnetic beads

(10002D, 10004D, Thermo Fisher Scientific) which were washed twice in RIP buffer. After 2

h incubation, the magnetic beads were washed thrice with RIP buffer for 10 min.

Immunoprecipitated RNA was eluted either by competitive elution (for RIP-seq) as previously

described (33) or by direct extraction using TRIzol (for slot blot).

For the m^6A RIP slot blot, the input RNA and the immunoprecipitated RNA were directly

subjected to the slot blot assay described above.

For m^6A RIP and RNA-seq, sequencing libraries for total RNA-seq/input for m^6A RIP-seq

were prepared from 10 ng of the fragmented RNA parallelly along with m^6A RIP samples using

SMARTer Stranded Total RNA-seq Kit V2, Pico Input Mammalian (Takara Bio) according to

manufactures instructions. All the libraries were single-end sequenced (1x 88 bp) on an

Illumina NextSeq 2000 platform at the BEA core facility, in Stockholm, Sweden.

Processing of m^6A RIP-seq data

Raw Illumina short-reads data obtained from SMARTER-Stranded Total RNA-Seq Kit v2

sequencing was processed using Trim Galore v0.6.6

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with a minimum length

threshold of 20bp. Quality control of Illumina short-reads was performed using FastQC v0.11.9

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) with default parameters (-q 20).

Trimmed reads were mapped with HISAT2 v2.2.1 (57) preserving strand information (-U –

rna-strandness R) using the Telomere-to-Telomere (T2T) human reference genome assembly

(T2T-CHM13 v1.1) (31). The Escherichia coli K12 genome was concatenated to the T2T

reference assembly for further spike-in normalization. To control the systematic variation

across m^6A RIP experiments, the amount of spiked-in bacterial RNA was estimated by

counting the total number of reads uniquely mapping to the E. coli K-12 reference genome

using Sambamba v0.7.1 (58). The E. coli spike-in bacterial counts were further used to

calculate scaling factors for each batch of m^6A RIP-seq samples. Duplicate reads were labeled

using MarkDuplicates from Picard v2.23.4 (http://broadinstitute.github.io/picard/). Alignments

were downsampled according to calculated spike-in scaling factors using DownsampleSam

from Picard (--strategy HighAccuracy). Mapping files were separated by strand after duplicate

removal using Sambamba v0.7.1.

Analysis of m^6A RIP-seq and ChIP-seq data

Peak calling was performed using MACS v2.2.6 (https://github.com/macs3-project/MACS) on

m^6A RIP and input alignments normalized to spike-in content, with parameters --no-model, --

kee*p*-dup auto, --call-summits, –extsize 75, q-value cutoff of 0.05 and effective genome size

according to the T2T-CHM13 v1.1 UCSC table browser. Motif analysis on m^6A peaks was

carried out using findMotifsGenome.pl from HOMER v4.11 (59). m^6A RIP coverage tracks

were generated using bamCoverage from deepTools v3.3.2 (60).

Public CTCF and RNA Pol II data reported by Deng et al. 2012 (32) were obtained from GEO

(GSE1962, GSE19484). Visualization of m^6A RIP coverage tracks data was done in R using

the rtracklayer (61) and ggplot2 packages (https://www.R-project.org). Telomere coordinates,

GC% content, and CpG island coordinates were retrieved from the T2T CHM13 v1.1 reference

genome hub (http://t2t.gi.ucsc.edu/chm13/hub/t2t-chm13-v1.1). Telomeric repeat tracks were

generated using scanMotifGenomeWide.pl from HOMER v4.11 and bedToBigBed v2.8 from

UCSC utilities (62).

Direct RNA sequencing using Oxford Nanopore Technologies (ONT) long-read

sequencing

Total RNA isolated from Control-sh and METTL3-KD cells, 20-25 μg was directly sequenced

on the Oxford nanopore platform. In brief Total RNA quantification and quality control were

performed on Fragment Analyzer 5300 system and showed RNA integrity number (RIN) of 10

for both samples. Poly(A) RNA Selection Kit V1.5 (Lexogen) was used to enrich

polyadenylated transcripts with minor changes to the protocol. All the processing volumes were

doubled, and 4 μg of Control-sh or METTL3-KD RNA was used as an input in a single

reaction. Each sample was batch-processed in 5 separate reactions that were combined at the

final elution step. Control-sh and METTL3-KD poly(A)-enriched RNA was analyzed on

Fragment Analyzer showing remaining rRNA contamination to be 11% and 21% of mass for

each sample respectively. Enriched poly(A)-RNA was then purified using 1x of RNAClean XP

(Beckman Coulter) beads to concentrate the sample and remove very short RNA fragments and

eluted into 11ul of water. Qubit™ RNA High Sensitivity kit (Invitrogen) was used to quantify

the RNA before direct RNA sequencing. SQK-RNA002 (ONT) kit was used for library

preparation and followed according to the manufacturer's protocol. Briefly, 400 ng of poly(A)-

enriched RNA was used as an input, and recommended reverse-transcription step was

performed. Control-sh and METTL3-KD libraries were loaded on individual FLO-MIN106

(ONT) flowcells and sequenced on GridION Mk1 device (ONT) for 72 h yielding 1.65 and

1.35 million long-reads respectively.

Processing of direct-RNA Nanopore sequencing data

Direct RNA Oxford Nanopore Technology (ONT) sequencing data was base-called using

Guppy v5.0.16 (flowcell FLO-MIN106, kit SQK-RNA002). Quality control was performed

with PycoQC (https://github.com/a-slide/pycoQC) and NanoPlot

(https://github.com/wdecoster/NanoPlot) on base-called data. ONT reads were mapped to the

T2T-CHM13 v1.1 reference genome using the latest version of Minimap (v2.24) (-ma*p*-ont -

uf -k14) optimized for mapping long reads to highly repetitive regions (63,64).

Analysis of ONT data

Identification of m[6]A modifications from direct RNA ONT data was performed following the

xPore pipeline (https://xpore.readthedocs.io/)(35). xPore allows the identification of m[6]A

under multiple conditions without the need for unmodified control samples, enabling the

quantitative estimation of modification rates from Nanopore sequencing data. Briefly, raw

Nanopore signal-level events were aligned to the T2T CHM13 v1.1 reference transcriptome

using Nanopolish eventalign v0.12.0. Processed ONT data was further analyzed with xPore

(default parameters; minimum read count 5, pre-filtering t-test method, 500 maximum

iterations) to estimate the genome-wide RNA modification rates for each independent sample

and the differentially modified sites between control and METTL3-KD conditions. A

transcriptome assembly was performed using StringTie2 (65) on Nanopore reads, and

remapped reads were used with xPore to refine the search for m^{6}A modifications at

chromosome ends. Predicted m^{6}A sites were selected from differentially modified positions

with a base A within the NNANN context with a significance p-value < 0.05 and a differential

modification rate (DMR) above zero.

Metagene analysis

Genome-wide distribution of m^{6}A sites from Illumina and ONT data was analyzed using the

plyranges package(66) to map the m^{6}A coordinates to the genomic feature coordinates (5'

UTR, CDS, 3'UTR) obtained from the T2T-CHM13 v1.1 GTF annotation files. Mapped

coordinates were further translated to their relative metagene positions. The peak density

distribution for all m^{6}A sites along metagene coordinates was calculated using the density

function in R.

Mappability at chromosome ends

To account for the repetitive nature of chromosome ends, we calculated the mappability of

short and long reads using GenMap(67). This tool estimates the occurrence of k-mer sequences

of a given length across the genome and computes a probability score proportional to its

occurrence at different parts of the genome. A mappability score of 1 indicates that a sequence

could align to a unique genomic location, while a low mappability value indicates that a

sequence can map to one or many sites in the genome. The computed mappability scores for

Illumina (75 bp) and Nanopore (600 bp) reads were averaged over fixed-size windows using

BEDTools v2.30.0 (68) and further used to normalize the number of reads aligned at each

chromosome end to the likelihood of mapping at one or multiple locations.

Analysis of TERRA expression

To analyze the expression towards chromosome ends in a strand-specific manner, the reads

from Illumina alignments mapping to subtelomeric 30 kb regions from chromosome q and p

arms were split by strand and filtered from all T2T alignments after duplicate removal.

Alignments from direct RNA ONT data were filtered using positive strand information. The

number of reads mapping to subtelomeric regions on each chromosome was quantified using

the multicov function from BEDTools v2.30.0(68). Expression values were further converted

to counts per million (CPM) to normalize for differences in library sizes across samples and

normalized to the mappability scores computed using GenMap to adjust for the likelihood of

short and long reads to align at one or multiple repetitive sites at each chromosome end.

Chromosome ends were classified into two categories (active/inactive) according to their

relative CPM expression values, being classified as 'active' if their expression was greater than

the median normalized expression value across all chromosomes ends at the control condition

and 'inactive' otherwise. Enrichment of telomeric repeats in m^{6}A RIP-seq from Control-sh,

METTL3-KD, and IgG data was calculated using TelomereHunter v1.0.4 using a default repeat

threshold of at least 10 non-consecutive repeats to search for TTAGGG, TGAGGG, TCAGGG

and TTGGGG repeats (69). Filtered TelomereHunter read counts were normalized using CPM

values.

Immunofluorescence and Proximity ligation assay (PLA)

U-2 OS and NB cell lines (CHLA-90 and SK-N-FI) were seeded on a coverslip in a 24-well

plate at a density of 100,000 and 200,000 respectively. In case of NB cells, coverslips

(MENZCB00120RAC20, Epredia) were coated with 50 μg/ml collagen (5005, PureCol, Type

I Collagen Solution) at RT for 1 h. Cells were fixed using 4% formaldehyde for 10 min

followed by 2x washes with PBS and stored in PBS at 4°C until further use.

Immunofluorescence was performed using a general protocol. Briefly, cells were

permeabilized for 10 min with 0.25% Triton X-100 followed by 3x washes with PBS-T. Cells

were blocked with 3% BSA in PBS-T for 30 min and left with primary antibody overnight at

4°C. Immunofluorescence was continued the next day starting with 3X PBS-T washes followed

by incubation with respective secondary antibodies for 1 h at RT, and then 3X PBS-T washes.

Coverslips were dried and mounted on a slide using ProLong Gold Antifade Mountant with

DAPI (P36931, Thermo Fisher Scientific). Images were taken using EVOS M7000 microscope

(Thermo Fisher Scientific). Information on the used antibodies is provided in Supplementary

Table 1.

For PLA cell fixation and permeabilization are the same as above. PLA was performed using

a Duolink® PLA kit (DUO92014, Sigma-Aldrich) according to the manufacturer's protocol.

As a background control single antibody was used in this assay. For R-loop PLA with S9.6 and

TRF2 antibodies, cells were pre-treated with RNase A after permeabilization for 30 min at

37°C to reduce the backgrounds. Information on the used antibodies is provided in

Supplementary Table 1. Images were taken using EVOS M7000 microscope (Thermo Fisher

Scientific).

RNA-FISH

U-2 OS and NB cell lines were seeded on a coverslip in a 24-well plate at a density of 100,000 and 200,000 respectively. In case of NB cells, coverslips (MENZCB00120RAC20, Epredia) were coated with 50 μg/ml collagen (5005, PureCol, Type I Collagen Solution) at RT for 1 h. Cells were fixed using 4% formaldehyde for 10 min followed by 2x washes with PBS and stored in 70% ethanol at 4°C until further use. RNA- FISH was performed to detect TERRA RNA and Stellaris RNA FISH protocol for adherent cells was used with some modifications. Briefly, cells were permeabilized for 10 min with 0.25% Triton X-100 followed by a wash with PBS. Cells were then incubated for 2-5 min with wash buffer A (Stellaris RNA FISH Wash Buffer A, SMF- WA1-60, Biosearch Technologies). Hybridization was carried out at 37°C for 4 h by incubating cells with hybridization buffer (Stellaris RNA FISH Hybridization Buffer, SMF-HB1-10, Biosearch Technologies) containing TERRA probe (TAACCC)$_7$-Atto488 in a humidified dark chamber. Cells were then washed 2x with wash buffer A at 37°C for 15 min, followed by 2-3 min incubation at RT with wash buffer B (Stellaris RNA FISH Wash Buffer B, SMF- WB1-20, Biosearch Technologies). Coverslips were dried and mounted on a slide using ProLong Gold Antifade Mountant with DAPI (P36931, Thermo Fisher Scientific). Images were taken using EVOS M7000 microscope (Thermo Fisher Scientific).

Sequential RNA-FISH and immunofluorescence (IF)

For sequential IF staining followed by RNA-FISH as described above cells were fixed using 4% formaldehyde for 10 min followed by 2x washes with PBS and stored in 70% ethanol at 4°C until further use. Coverslips were rinsed with PBS and cells were permeabilized for 10 min using 0.25% Triton X-100 in 1x PBS at RT. Cells were then washed using PBS-T (0.1% Tween-20 in 1x PBS) and blocked in 3% BSA in PBS-T for 30 min. Primary antibodies were diluted in 3% BSA in PBS-T and incubated for 2 h after blocking. 3x5 min washes were performed using PBS-T and incubated for 1 h with secondary antibodies diluted in 3% BSA in PBS-T. 3x5 min washes were performed using PBS-T followed by 10 min fixation by 4% formaldehyde. Cells were rinsed with PBS twice, permeabilized with 0.25% Triton X-100 in 1x PBS for 5 min followed by a wash with PBS. TERRA RNA-FISH was performed as described above and proceeded for imaging. Details of antibodies are provided in Supplementary Table 1.

Simultaneous RNA-FISH and m⁶A IF staining

Simultaneous TERRA FISH and m⁶A IF were performed as described before (30) with the following modifications. Cells were fixed in MeOH at -20°C for 15 min and TERRA RNA FISH was performed as described above with some modification. After hybridization, during

2x30 min washing steps, cold wash buffer A (described above) was added to coverslips and

incubated at 37°C. This step allows refolding of rRNAs to reduce the nuclear RNA signal

coming from rRNAs as described by Fu et al (30). Coverslips were then washed with wash

buffer B (described above) for 2-3 min and then rinsed with 1x PBS. Coverslips were incubated

in wash buffer A containing 10% formamide for 5-10 min and followed by blocking for 1 h

(3% BSA in PBS-T) and the primary antibody against m^{6}A was incubated for 2 h at RT in 3%

BSA in PBS-T. 3x5 min PBS-T washes were done and the secondary antibody was incubated

for 1 h at RT. Coverslips were washed 3x5 min with PBS-T and mounted. Imaging was done

using EVOS M7000 microscope.

Western blot

Cell lysates were prepared using RIPA buffer (R0278, Sigma-Aldrich,) and total protein

content was quantified using Pierce BCA Protein Assay Kit (23225, Thermo Fisher Scientific)

as per manufacturer's instructions. Equal amounts and volume of samples were resolved by

SDS- PAGE on NuPAGE Bis-Tri Gels (4-12%) (Thermo Fisher Scientific), followed by

transfer onto 0.2 μm Nitrocellulose membrane using Trans-Blot Turbo Transfer System (Bio-

Rad). The membrane was blocked for 1 h at RT with a 5% blocking solution of non-fat dried

milk in PBS-T.

The membrane was incubated with the primary antibody in a 5% blocking solution overnight

at 4°C. Immunoblotting was continued the next day starting with 3x PBS-T washes followed

by incubation with respective secondary antibodies for 1 h at RT, and then 3X PBS-T washes.

Proteins were detected with SuperSignal West Pico PLUS Chemiluminescent Substrate

(34579, Thermo Fisher Scientific) using the ChemiDoc XRS+ system (Bio-Rad). ImageLab

software was used to quantify the bands. Antibodies used in Western blot analysis are provided

in Supplementary Table 1.

Proliferation assay (MTT)

5,000 cells/well were seeded in a 96-well plate to assess the proliferation of cells after METTL3

inhibitor STM2457 (10 μM) at indicated time points. CellTiter 96 Non-Radioactive Cell

Proliferation Assay kit (G4000, Promega) was used to determine cell growth and the

manufacturer's instructions were followed. Absorbance was measured using a microplate

reader Infinite 50 (Tecan, Austria).

Drug treatments:

U-2 OS cells were treated with Bleomycin 10 μg/ml containing media for 4 h. U-2 OS cells

were treated with Actinomycin D (10 μg/ml) for the indicated time points and RNA was

isolated from the treated cells to perform TERRA slot blot assay. METTL3 inhibitor STM2457

(10 μM, HY-134836), UZH1A (50 μM, HY-134673A), and UZH1B (50 μM, HY-134673B)

were treated for 6 h to perform TERRA RNA-FISH and 48 h for telomere DNA damage IF.

Colony formation assay

10,000 cells/well were seeded in a 6-well plate and allowed to attach overnight. Doxycycline

(200 ng/ml) was added to the media for induction of METTL3-KD and cells were left to grow

sparsely for 14 days with induction media which was replaced after every 3-4 days. Cells were

then washed 1x with PBS, fixed with 4% formaldehyde for 10 min at RT, washed 1x with PBS,

and then stained using 1% crystal violet solution at RT for 2 h. The plate was carefully washed

and allowed to dry. The image was taken using the ChemiDoc XRS+ system (Bio-Rad).

In-vitro METTL3/METTL14 methyltransferase assay

Methyl-transferase assay with RNA oligos (1 μm DRACH motif containing and TERRA repeat

containing oligo, Supplementary Table 1) was performed with recombinant

METTL3/METTL14 enzyme complex (31570, Active motif) according to manufacturer's

instructions, with 10 nM S-Adenosyl methionine (SAM) as methyl group donor. Oligos

without enzyme complex were used as a negative control. To detect if the RNA oligos

underwent m^6A modification LC-MS/MS-based quantification of m^6A was done, as previously

described (Liu et al. 2020). In brief oligos, post methyltransferase assay were digested by

Nuclease P1 (N8630, Sigma-Aldrich) followed by treatment with phosphatase (M0289S,

NEB). The sample was then filtered (0.22 μm pore size) and directly injected into the LC-MS.

We made triplicate injections of the RNA oligos (both experimental and negative controls) and

estimated the ratio of A/m^6A. LC-MS/MS profiles were monitored using the parallel reaction-

monitoring (PRM) mode for: m/z 268.0–136.0, m/z 282.0–150.1 which corresponds to A and

m^6A respectively as described before (Sun et al. 2021). DRACH oligos served as a positive

control.

Purification of TERRA RNA and RNA-MS

TERRA RNA pulldown: Capturing a specific RNA from total RNA was performed as

described previously (55). In brief, ~300 μg of DNase I treated RNA isolated from U2O-S cells

was mixed with 400 pmoles of biotinylated DNA probe (Supplementary Table S1) against

TERRA or Luciferase probe (negative control) in 5x SSC buffer and denatured at 90°C for 3

min. This mixture was immediately cooled down on the ice and again heated at 65°C for 10

min after which the samples were left to cool down at RT to allow hybridization of the

probe. To pulldown the RNA hybridized with biotinylated DNA probes, MyOne Streptavidin

C1 magnetic Dynabeads (Thermo Fisher Scientific) were added and incubated for 30 min at

room temperature. The magnetic beads were washed once with 1x SSC buffer and thrice with

0.1x SSC buffer. To elute the captured RNA, the magnetic beads were resuspended in nuclease-

free water, and heated at 75°C for 3 min. The eluted RNA was treated with RNase-free DNase

I for 30 min at 37°C. Finally, the RNA was precipitated with 3 volumes of 100% ethanol, 0.5

M ammonium acetate, and GlycoBlue (Thermo Fisher Scientific) at -80°C for 2 h. The

precipitated RNA was washed with 80% ethanol and resuspended in nuclease-free water.

The RNA captured using TERRA or Luciferase probe (Luciferase) was first checked for

specificity by a slot blot assay using TERRA probe (see slot blot assay).

Further, the captured TERRA RNA along with synthetic RNA oligo (Supplementary Table S1)

were digested with RNase T1 and analyzed on a mass spectrometer (MS).

RNA-MS analysis: RNA-MS was performed using an Easy-nLC 1200 LC system (Thermo

Fisher Scientific) coupled to Orbitrap Fusion™ Eclipse™ Tribrid™ mass spectrometer. The

analytes were trapped on a 2 cm x 100 μm Acclaim PepMap C18 precolumn (particle size

5 μm; Thermo Fisher Scientific) and separated on a 30 cm x 75 μm analytical column packed

in-house with 3 μm Reprosil-Pur C18 material (Dr. Maisch, Germany) at 300 nL/min flow

using a stepwise elution profile during 45 min. Solvent A was 5 mM di-n-butylamine and 8 mM

acetic acid in H_2O, and solvent B was 70% methanol, 5 mM di-n-butylamine, and 8 mM acetic

acid. The column oven temperature was 50°C. Nano-Flex ion source (Thermo Fisher

Scientific) was operated in negative ionization mode at 1.8 kV with the ion transfer capillary

temperature 275°C. The MS instrument was run in a targeted MS/MS setting with switching

quadrupole isolation between the two tetramers of interest. Each precursor was isolated with a

1.1 Da window and followed by HCD 30% recorded at 15,000 resolution in the Orbitrap with

the first m/z 130. The monoisotopic mass of UUAGp is 1304.1 Da equivalent to doubly charged

m/z 651.07 and UUm^{6}AGp has a monoisotopic mass of UUAGp is 1318.2 Da equivalent to

doubly charged m/z 658.08. The ratio of UUm^{6}AGp over UUAGp was calculated from the

average intensities of the respective fragment ions. The MS2 spectra over the targeted

precursors for 30 seconds were averaged and the total fragment intensities were calculated and

compared. The intensities for UUm^{6}AGp fragment ions were divided by the total sum of

UUAGp and UUm^{6}AGp fragment ions intensities.

METTL3 Chromatin Immunoprecipitation (ChIP)

ChIP with METTL3 antibody was performed as described before (70). U-2 OS cells were fixed

with formaldehyde (1% final concentration) for 15 min at RT. Fixed cells were washed with

cold PBS, lysed and the chromatin was sheared using a Bioruptor (Diagenode) until an average

fragment size range of 200-500 bp was achieved. Immunoprecipitation of solubilized

chromatin was carried out with 3 μg of METTL3/IgG antibody overnight at 4°C. The

immunoprecipitated complex was captured using a mix of Protein A and G Dynabeads (Invitrogen), washed and RNase A treated before eluting the DNA. The ChIP DNA was denatured at 95°C for 5 min before assaying on slot blot using TERRA probe.

Re-RIP (hnRNPA2B1-m⁶A)

Control-sh and METTL3-KD U-2 OS cells were harvested and fixed for 10 min with formaldehyde (1% final concentration) and quenched with Glycine (0.125 M, final concentration). Fixed cells were lysed in RIPA buffer (50 mM Tris pH7.4, 150 mM NaCl, 0.5% Sodium Deoxycholate, 0.2% SDS, 1% IGEPAL-CA630, Protease inhibitor, and RNase inhibitor) and sonicated for 10 cycles (30 secs on 30 secs off) on Bioruptor (Diagenode) and cleared lysate was used for RIP with 3 μg of hnRNPA2B1 (A73256, Gentek,) and IgG (sc-2027, Santa Cruz) antibodies. The RNA-protein-antibody complex was captured using protein A/G magnetic beads (Thermo Fisher Scientific). Magnetic beads were washed with low salt buffer (1x PBS, 0.1% SDS, and 0.5% IGEPAL-CA630) and high salt buffer (5x PBS, 0.1% SDS, and 0.5% IGEPAL-CA630) before eluting in elution buffer (10 mM Tris pH 7.4, 100 mM NaCl, 1 mM EDTA, 0.5% SDS) with Proteinase K followed by RNA extraction using TRIzol reagent. The RNA isolated after hnRNPA2B1/IgG RIP was subsequently used as input for Re-RIP with m⁶A antibody. m⁶A RIP was performed as described in the above section. The final Re-RIP RNA was directly assayed on slot blot.

Purification of hnRNPA2B1 protein and *in vitro* binding assay

His-tagged hnRNPA2_WT (98662, Addgene) was transformed in Rossetta DE3 bacterial strain. Cells were grown overnight and further diluted to 0.1 OD 600 nm and grown to about 0.6 OD 600 nm at 37°C and then cells were induced with 0.3 mM IPTG at 18°C for 14-16 h. Cells were pelleted and dissolved in 25 mM HEPES pH 7.5, 150 mM NaCl, 20 mM KCl, and 20 mM MgCl$_2$ along with 1x Protease inhibitor cocktail. Cells were lysed at 4°C in presence of lysozyme followed by brief sonication. Cells debris were removed by spinning cells lysate at 13000 rpm for 30 min at 4C and cleared lysate were loaded onto Co+ NTA resin for binding of His-tagged hnRNPA2B1 protein and resin were washed with 20 mM imidazole containing lysis buffer without 1x Protease inhibitor cocktail to remove non-specific proteins and His-tagged protein were eluted in 300 mM imidazole containing lysis buffer and eluted protein were further dialyzed in lysis buffer and proteins were stored in -80°C for further use.

The hnRNPA2B1 binding with RNA was performed following the protocol as described before (71) with few modifications. Total RNA from Control-sh or METTL3-KD U-2 OS cells were used instead of *in vitro* synthesized RNA. The RNA bound protein complex was captured by

HisPur Cobalt Resin (89965, Thermo Fisher Scientific) and TERRA enrichment was checked

by slot blot assay as described above.

Analysis associated with detection of Chromosomal aberration and ATRX status in NB

Swedish cohort

Tumor samples were analyzed with Applied biosystems CytoScan HD array (Thermo Fisher

Scientific, Waltham) according to the experimental procedure previously described (72,73).

The arrays detect both copy number alterations and allele-specific information at high

resolution. For primary data analysis, GDAS software (Thermo Fisher Scientific) was used.

Chromosome Analysis Suite (ChAS v.3.3; Thermo Fisher Scientific) was used for the

generation of genomic profiles and determination of chromosomal aberration, MYCN

amplification, and ATRX status. For Whole Genome Sequencing (WGS) DNA from tumor

material and matched constitutional DNA were subjected to sequencing and bioinformatical

handling as described previously (73).

C-circle assay

50 ng of genomic DNA was digested with 4 U/μg HinfI and RsaI restriction enzymes and 25

ng/μg RNase for 1 h at 37°C (prepare every sample twice to include no polymerase as negative

control). Digested DNA was incubated with 100ug/ml Albumin, 1 mM each dATP, dGTP, and

dTTP, 0.1% Tween, 1x Φ29 Buffer, and 7.5 U Φ29 DNA polymerase (NEB) at 30°C for 8 h

then 65°C for 20 min. The reaction products were slot blotted onto a 6x SSC soaked Biodyne

B nylon membrane (GE Healthcare) for quantification and UV-crosslinking the DNA onto the

membrane at 0.120 Joules. Prehybridization was done using DIG Easy Hyb (11603558001,

Roche) for 30 min at 37°C, and after adding 25 ng/ml DIG-labeled TERRA probe: 5´-

(TAACCC)$_5$-DIG (Sigma-Aldrich) at 37°C overnight with rotation. For hybridization, the DIG

Wash and Block Buffer set (11585762001, Roche) was used according to the manufacturer's

instructions. The membrane was incubated with an anti-DIG antibody conjugated with alkaline

phosphatase (1:20,000, 11093274910, Roche) for 1 h and CDP-Star Chemoluminescent

substrate (C0712, Sigma-Aldrich) to detect the C-circles. Images were taken using High-

resolution Chemiluminescence (ChemiDoc, Bio-Rad)

Telomere DNA FISH and Immunostaining

The cells were fixed with 4% formaldehyde for 10 min, followed two washes with PBS. Then,

cells were permeabilized using 0.25% Triton X-100 in PBS for 10 min, followed by two washes

with 0.1% PBS-Tween (PBS-T). Blocking step and RNase A treatment were performed by

adding RNase A (100 μg/ml, Sigma-Aldrich) in 3% bovine serum albumin (BSA) for 1h at RT,

followed by 2 washes with PBS-T. The cells were incubated overnight with primary antibody

γ-H2AX (1:500, Thermo Fisher Scientific) at 4°C, washed three times 5 min in PBS-T, and

subsequently incubated with secondary antibody conjugated with Alexa Fluor 555 (1:800,

A32727, Thermo Fisher Scientific) for a 1 h in the dark at room temperature. After incubation

with the antibody, cells were washed three times for 5 min in PBS-T. Following

immunofluorescence staining, the cells were refixed using 4% formaldehyde for 10 min. After

washing twice with PBS, the slides were dehydrated with 50, 70, and 100% ethanol for 3 min

and air dried for 5 min at RT. For hybridization, denature slide with probe (1:200, (TAACCC)$_7$-

Atto488) in hybridization buffer (20 mM Na$_2$HPO$_4$, pH 7.4, 20 mM Tris, pH 7.4, 60%

formamide, 2x SSC and 0.1 µg/ml salmon sperm DNA) for 10 min at 85°C, and incubated with

the probe in a humidified chamber for 2 h at RT. Following hybridization, wash twice in

washing solution (2x SCC/0.1% Tween) for 15 min each and MQ water twice for 2 min.

Prolong Gold with DAPI (Thermo Fisher Scientific) was added to each coverslip and air-dried

in the dark before imaging using 60X objectives in EVOS M7000 microscope.

Mouse xenograft experiments

TetO Control and METTL3-KD SK-N-F1 and CHLA-90 cells were subcutaneously injected

into the right dorsal flank (2 x 10^6 and 5 × 10^6 cells, respectively) of 5-week-old female nude

mice (Charles River) in 100 µL and 200 µL mixture of (1:3) Matrigel/PBS, (n=4 and 3 per

group, respectively). METTL3 knockdown was induced by adding 2 mg/mL Dox and 2%

sucrose to drinking water, after 4-5 days of cell injection. Mice weight was monitored weekly

and tumor volume was measured every 2–3 days using a digital calliper and calculated

according to the formula Volume (mm3) = (w2 x l x π)/6 (where w is width, shortest diameter,

and l is length, longest diameter). Mice were euthanized when tumors reached 1000 mm3 or

had weight loss ≥ 10% of initial weight. At the end of the experiment, tumors were collected,

weighed, and processed for further analysis. All experiments were carried out per the standards

approved by the Institutional Ethical Committee of Animal Experimentation, Gothenburg,

Sweden (ethical permit no 3722/21). METTL3-KD was verified before and after cell injection

by Western blot.

DNA-RNA immunoprecipitation (DRIP) and DRIP followed by m^{6}A RIP (DRImR)

DRIP was performed as described previously with few alterations. In brief, cells were lysed in

lysis buffer (1xTE supplemented with 0.5% SDS) for 15 min at RT. Brief RNase A treatment

at 37°C for 15 min was done to remove free RNA from the subsequent Proteinase K treatment

and genomic DNA isolation by Phenol chloroform isolation. The genomic DNA before S9.6

IP were digested with a cocktail of restriction enzymes (*Bsr*GI, *Eco*RI, *Hind*III, *Ssp*I, and *Xba*I)

and RNase III which is shown to remove non-specific enrichment of double-stranded RNA to

identify genuine R-loop enriched RNAs. IP reaction was carried out with fragmented genomic

DNA that was either pre-treated/non-pre-treated with RNase H to cross-check the specificity

of S9.6 IP. Following 9.6 IP we performed extensive DNase I treatment to obtain R-loop

associated RNA. For DRImR, DRIP enriched RNA was used as input for m^{6}A RIP as described

above with above without fragmentation.

Analysis of neuroblastoma next-generation sequencing data from a German cohort

1117

Copy number gains were estimated from a combination of published neuroblastoma paired-

end WGS (25) paired-end WES (26), and aCGH (74) datasets. ALT and MYCN status of the

patients were taken from sample overlaps with Roderweiser et al. (46), sequence alignment to

reference GRCh37/hg19 was done using BWA mem (v.0.7.13-r1126;

https://github.com/lh3/bwa) and copy number analysis was performed with Sclust(75).

Gene expression analysis was performed with the RNA-Seq dataset published by Zhang et al

(76). Expression values had been computed using the Magic-AceView pipeline and were given

as normalized log(sFPKM) values [see (46) for details]. Survival information, ALT, and

MYCN annotation were again taken from Roderweiser et al. (46). Survival analysis was done

with groups with high and low expression of selected genes in survival analysis were generated

by maximally selected rank statistics using the R package maxstat (version 0.7-25).

1129

Image quantification and statistical analysis:

Western and slot blots were quantified using ImageLab software (Bio-Rad). Images were

quantified using ImageJ software (77). DAPI was used to mark the nuclear border and TERRA

foci, PLA foci, or colocalization signals within the nucleus were counted using particle

analysis. More than 100 nuclei were counted for U-2 OS cells and at least 50 nuclei in the case

of NB cells for each experiment. Graphs were prepared, and statistical analysis was performed

using GraphPad Prism (version 9.4.1 for Windows, GraphPad Software, San Diego, California,

USA, www.graphpad.com). The data are displayed as Mean ± Standard Deviation (SD) of a

minimum of three independent experiments. Each box-and-whisker plot presents the median

value as a line, a box around the lower and higher quartiles, and whiskers extending to the

maximum and lowest values. Statistical significance between 2 groups was analyzed by

Student's 2-tailed unpaired t-test. Wilcoxon paired signed-rank test, Dunnett's test, Turkey's

test, and Sidak's test were used to perform multiple comparisons, and *p*-values were reported

in the Figures and the Figure legends.

1144

Data availability

The data underlying this article are available in NCBI Gene Expression Omnibus (GEO) at

https://www.ncbi.nlm.nih.gov/geo/ and can be accessed with the GSE219146 and GSE219212

accession numbers.

Acknowledgements

We would like to thank the core facility at Novum, BEA, Bioinformatics and Expression

Analysis, which is supported by the board of research at the Karolinska Institute and the

research committee at the Karolinska hospital for help with sequencing. Proteomics core

facility at GU for their support with LC-MS/MS. The computations and data handling were

enabled by resources in project SNIC-2022-22-85 provided by the Swedish National

Infrastructure for Computing (SNIC) at UPPMAX, partially funded by the Swedish Council

through grant agreement no. 2018-05973. We acknowledge Anders Sjölander for assistance

concerning technical and implementational aspects of the UPPMAX resources.

Funding:

This work was funded by grants from the Swedish Research Council (Vetenskapsrådet, 2018-

02224) and project grants from Svenska Läkaresällskapet, Barncancerfonden, Cancerfonden,

Åke Wibergs Stiftelse, and Kungl Vetenskaps- och Vitterhets-Samhället (KVVS) grant to TM,

4 years research position grant from Barncancerfonden to TM, Tore Nilsons Stiftelse and

Bollan scholarship to RV, Assar Gabrielsson Fond to RV and KT.

References:

1. Barbieri, I. and Kouzarides, T. (2020) Role of RNA modifications in cancer. *Nat Rev*
Cancer, **20**, 303-322.

2. Huang, H., Weng, H., Zhou, K., Wu, T., Zhao, B.S., Sun, M., Chen, Z., Deng, X., Xiao, G.,
Auer, F. *et al.* (2019) Histone H3 trimethylation at lysine 36 guides m(6)A RNA
modification co-transcriptionally. *Nature*, **567**, 414-419.

3. Patil, D.P., Chen, C.K., Pickering, B.F., Chow, A., Jackson, C., Guttman, M. and Jaffrey,
S.R. (2016) m(6)A RNA methylation promotes XIST-mediated transcriptional
repression. *Nature*, **537**, 369-373.

4. Yue, Y., Liu, J., Cui, X., Cao, J., Luo, G., Zhang, Z., Cheng, T., Gao, M., Shu, X., Ma, H. *et*
al. (2018) VIRMA mediates preferential m(6)A mRNA methylation in 3'UTR and near
stop codon and associates with alternative polyadenylation. *Cell Discov*, **4**, 10.

5. Xu, W., Li, J., He, C., Wen, J., Ma, H., Rong, B., Diao, J., Wang, L., Wang, J., Wu, F. *et al.*
(2021) METTL3 regulates heterochromatin in mouse embryonic stem cells. *Nature*,
591, 317-321.

6. Zheng, G., Dahl, J.A., Niu, Y., Fedorcsak, P., Huang, C.M., Li, C.J., Vagbo, C.B., Shi, Y.,
Wang, W.L., Song, S.H. *et al.* (2013) ALKBH5 is a mammalian RNA demethylase that
impacts RNA metabolism and mouse fertility. *Mol Cell*, **49**, 18-29.

7. Alarcon, C.R., Goodarzi, H., Lee, H., Liu, X., Tavazoie, S. and Tavazoie, S.F. (2015)
HNRNPA2B1 Is a Mediator of m(6)A-Dependent Nuclear RNA Processing Events. *Cell*,
162, 1299-1308.

8. Zhou, K.I., Shi, H., Lyu, R., Wylder, A.C., Matuszek, Z., Pan, J.N., He, C., Parisien, M. and
Pan, T. (2019) Regulation of Co-transcriptional Pre-mRNA Splicing by m(6)A through
the Low-Complexity Protein hnRNPG. *Mol Cell*, **76**, 70-81 e79.

9. Xu, W., He, C., Kaye, E.G., Li, J., Mu, M., Nelson, G.M., Dong, L., Wang, J., Wu, F., Shi,
Y.G. *et al.* (2022) Dynamic control of chromatin-associated m(6)A methylation
regulates nascent RNA synthesis. *Mol Cell*, **82**, 1156-1168 e1157.

10. Liu, J., Gao, M., He, J., Wu, K., Lin, S., Jin, L., Chen, Y., Liu, H., Shi, J., Wang, X. *et al.*
(2021) The RNA m(6)A reader YTHDC1 silences retrotransposons and guards ES cell
identity. *Nature*, **591**, 322-326.

11. Abakir, A., Giles, T.C., Cristini, A., Foster, J.M., Dai, N., Starczak, M., Rubio-Roldan, A.,
Li, M., Eleftheriou, M., Crutchley, J. *et al.* (2020) N(6)-methyladenosine regulates the
stability of RNA:DNA hybrids in human cells. *Nat Genet*, **52**, 48-55.

12. Yang, X., Liu, Q.L., Xu, W., Zhang, Y.C., Yang, Y., Ju, L.F., Chen, J., Chen, Y.S., Li, K., Ren,
J. *et al.* (2019) m(6)A promotes R-loop formation to facilitate transcription
termination. *Cell Res*, **29**, 1035-1038.

13. Porman, A.M., Roberts, J.T., Duncan, E.D., Chrupcala, M.L., Levine, A.A., Kennedy,
M.A., Williams, M.M., Richer, J.K. and Johnson, A.M. (2022) A single N6-
methyladenosine site regulates lncRNA HOTAIR function in breast cancer cells. *PLoS
Biol*, **20**, e3001885.

14. Wang, X., Liu, C., Zhang, S., Yan, H., Zhang, L., Jiang, A., Liu, Y., Feng, Y., Li, D., Guo, Y.
et al. (2021) N(6)-methyladenosine modification of MALAT1 promotes metastasis via
reshaping nuclear speckles. *Dev Cell*, **56**, 702-715 e708.

15. Jones, A.N., Tikhaia, E., Mourao, A. and Sattler, M. (2022) Structural effects of m6A
modification of the Xist A-repeat AUCG tetraloop and its recognition by YTHDC1.
Nucleic Acids Res, **50**, 2350-2362.

16. Ninomiya, K., Iwakiri, J., Aly, M.K., Sakaguchi, Y., Adachi, S., Natsume, T., Terai, G., Asai,
K., Suzuki, T. and Hirose, T. (2021) m(6) A modification of HSATIII lncRNAs regulates
temperature-dependent splicing. *EMBO J*, **40**, e107976.

17. Liu, J., Dou, X., Chen, C., Chen, C., Liu, C., Xu, M.M., Zhao, S., Shen, B., Gao, Y., Han, D.
et al. (2020) N (6)-methyladenosine of chromosome-associated regulatory RNA
regulates chromatin state and transcription. *Science*, **367**, 580-586.

18. Lee, J.H., Wang, R., Xiong, F., Krakowiak, J., Liao, Z., Nguyen, P.T., Moroz-Omori, E.V.,
Shao, J., Zhu, X., Bolt, M.J. *et al.* (2021) Enhancer RNA m6A methylation facilitates
transcriptional condensate formation and gene activation. *Mol Cell*, **81**, 3368-3385
e3369.

19. Azzalin, C.M., Reichenbach, P., Khoriauli, L., Giulotto, E. and Lingner, J. (2007)
Telomeric repeat containing RNA and RNA surveillance factors at mammalian
chromosome ends. *Science*, **318**, 798-801.

20. Schoeftner, S. and Blasco, M.A. (2008) Developmentally regulated transcription of
mammalian telomeres by DNA-dependent RNA polymerase II. *Nat Cell Biol*, **10**, 228-
236.

21. Feretzaki, M., Pospisilova, M., Valador Fernandes, R., Lunardi, T., Krejci, L. and Lingner,
J. (2020) RAD51-dependent recruitment of TERRA lncRNA to telomeres through R-
loops. *Nature*, **587**, 303-308.
22. Vohhodina, J., Goehring, L.J., Liu, B., Kong, Q., Botchkarev, V.V., Jr., Huynh, M., Liu, Z.,
Abderazzaq, F.O., Clark, A.P., Ficarro, S.B. *et al.* (2021) BRCA1 binds TERRA RNA and
suppresses R-Loop-based telomeric DNA damage. *Nat Commun*, **12**, 3542.
23. Giannone, R.J., McDonald, H.W., Hurst, G.B., Shen, R.F., Wang, Y. and Liu, Y. (2010)
The protein network surrounding the human telomere repeat binding factors TRF1,
TRF2, and POT1. *PLoS One*, **5**, e12407.
24. Yoshida, A., Oyoshi, T., Suda, A., Futaki, S. and Imanishi, M. (2022) Recognition of G-
quadruplex RNA by a crucial RNA methyltransferase component, METTL14. *Nucleic*
Acids Res, **50**, 449-457.
25. Peifer, M., Hertwig, F., Roels, F., Dreidax, D., Gartlgruber, M., Menon, R., Kramer, A.,
Roncaioli, J.L., Sand, F., Heuckmann, J.M. *et al.* (2015) Telomerase activation by
genomic rearrangements in high-risk neuroblastoma. *Nature*, **526**, 700-704.
26. Ackermann, S., Cartolano, M., Hero, B., Welte, A., Kahlert, Y., Roderwieser, A.,
Bartenhagen, C., Walter, E., Gecht, J., Kerschke, L. *et al.* (2018) A mechanistic
classification of clinical phenotypes in neuroblastoma. *Science*, **362**, 1165-1170.
27. Koneru, B., Lopez, G., Farooqi, A., Conkrite, K.L., Nguyen, T.H., Macha, S.J., Modi, A.,
Rokita, J.L., Urias, E., Hindle, A. *et al.* (2020) Telomere Maintenance Mechanisms
Define Clinical Outcome in High-Risk Neuroblastoma. *Cancer Res*, **80**, 2663-2675.
28. Macha, S.J., Koneru, B., Burrow, T.A., Zhu, C., Savitski, D., Rahman, R.L., Ronaghan,
C.A., Nance, J., McCoy, K., Eslinger, C. *et al.* (2022) Alternative Lengthening of
Telomeres in Cancer Confers a Vulnerability to Reactivation of p53 Function. *Cancer*
Res, **82**, 3345-3358.
29. Sun, H., Li, K., Zhang, X., Liu, J., Zhang, M., Meng, H. and Yi, C. (2021) m(6)Am-seq
reveals the dynamic m(6)Am methylation in the human transcriptome. *Nat Commun*,
12, 4778.
30. Fu, Y. and Zhuang, X. (2020) m(6)A-binding YTHDF proteins promote stress granule
formation. *Nat Chem Biol*, **16**, 955-963.
31. Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., Vollger,
M.R., Altemose, N., Uralsky, L., Gershman, A. *et al.* (2022) The complete sequence of
a human genome. *Science*, **376**, 44-53.
32. Deng, Z., Wang, Z., Stong, N., Plasschaert, R., Moczan, A., Chen, H.S., Hu, S.,
Wikramasinghe, P., Davuluri, R.V., Bartolomei, M.S. *et al.* (2012) A role for CTCF and
cohesin in subtelomere chromatin organization, TERRA transcription, and telomere
end protection. *EMBO J*, **31**, 4165-4178.
33. Zeng, Y., Wang, S., Gao, S., Soares, F., Ahmed, M., Guo, H., Wang, M., Hua, J.T., Guan,
J., Moran, M.F. *et al.* (2018) Refined RIP-seq protocol for epitranscriptome analysis
with low input materials. *PLoS Biol*, **16**, e2006092.
34. Chen, L., Zhang, C., Ma, W., Huang, J., Zhao, Y. and Liu, H. (2022) METTL3-mediated
m6A modification stabilizes TERRA and maintains telomere stability. *Nucleic Acids Res*.
35. Pratanwanich, P.N., Yao, F., Chen, Y., Koh, C.W.Q., Wan, Y.K., Hendra, C., Poon, P., Goh,
Y.T., Yap, P.M.L., Chooi, J.Y. *et al.* (2021) Identification of differential RNA
modifications from nanopore direct RNA sequencing with xPore. *Nat Biotechnol*, **39**,
1394-1402.

36.   Tan, K.T., Slevin, M.K., Meyerson, M. and Li, H. (2022) Identifying and correcting
1273   repeat-calling errors in nanopore sequencing of telomeres. *Genome Biol*, **23**, 180.
1274   37.   Zhang, C., Chen, L., Peng, D., Jiang, A., He, Y., Zeng, Y., Xie, C., Zhou, H., Luo, X., Liu, H.
1275   *et al.* (2020) METTL3 and N6-Methyladenosine Promote Homologous Recombination-
1276   Mediated Repair of DSBs by Modulating DNA-RNA Hybrid Accumulation. *Mol Cell*, **79**,
1277   425-442 e427.
1278   38.   Xiang, Y., Laurent, B., Hsu, C.H., Nachtergaele, S., Lu, Z., Sheng, W., Xu, C., Chen, H.,
1279   Ouyang, J., Wang, S. *et al.* (2017) RNA m(6)A methylation regulates the ultraviolet-
1280   induced DNA damage response. *Nature*, **543**, 573-576.
1281   39.   Liu, M., Hales, B.F. and Robaire, B. (2014) Effects of four chemotherapeutic agents,
1282   bleomycin, etoposide, cisplatin, and cyclophosphamide, on DNA damage and
1283   telomeres in a mouse spermatogonial cell line. *Biol Reprod*, **90**, 72.
1284   40.   Xiong, F., Wang, R., Lee, J.H., Li, S., Chen, S.F., Liao, Z., Hasani, L.A., Nguyen, P.T., Zhu,
1285   X., Krakowiak, J. *et al.* (2021) RNA m(6)A modification orchestrates a LINE-1-host
1286   interaction that facilitates retrotransposition and contributes to long gene
1287   vulnerability. *Cell Res*, **31**, 861-885.
1288   41.   Xu, M., Chigumira, T., Chen, Z., Tones, J., Zhao, R., Dahl, K.N., Chenoweth, D.M. and
1289   Zhang, H. (2021) CRISPR Cas13-Based Tools to Track and Manipulate Endogenous
1290   Telomeric Repeat-Containing RNAs in Live Cells. *Front Mol Biosci*, **8**, 785160.
1291   42.   Yankova, E., Blackaby, W., Albertella, M., Rak, J., De Braekeleer, E., Tsagkogeorga, G.,
1292   Pilka, E.S., Aspris, D., Leggate, D., Hendrick, A.G. *et al.* (2021) Small-molecule inhibition
1293   of METTL3 as a strategy against myeloid leukaemia. *Nature*, **593**, 597-601.
1294   43.   Moroz-Omori, E.V., Huang, D., Kumar Bedi, R., Cheriyamkunnel, S.J., Bochenkova, E.,
1295   Dolbois, A., Rzeczkowski, M.D., Li, Y., Wiedmer, L. and Caflisch, A. (2021) METTL3
1296   Inhibitors for Epitranscriptomic Modulation of Cellular Processes. *ChemMedChem*, **16**,
1297   3035-3043.
1298   44.   Chu, H.P., Cifuentes-Rojas, C., Kesner, B., Aeby, E., Lee, H.G., Wei, C., Oh, H.J., Boukhali,
1299   M., Haas, W. and Lee, J.T. (2017) TERRA RNA Antagonizes ATRX and Protects
1300   Telomeres. *Cell*, **170**, 86-101 e116.
1301   45.   Hartono, S.R., Malapert, A., Legros, P., Bernard, P., Chedin, F. and Vanoosthuyse, V.
1302   (2018) The Affinity of the S9.6 Antibody for Double-Stranded RNAs Impacts the
1303   Accurate Mapping of R-Loops in Fission Yeast. *J Mol Biol*, **430**, 272-284.
1304   46.   Roderwieser, A., Sand, F., Walter, E., Fischer, J., Gecht, J., Bartenhagen, C., Ackermann,
1305   S., Otte, F., Welte, A., Kahlert, Y. *et al.* (2019) Telomerase Is a Prognostic Marker of
1306   Poor Outcome and a Therapeutic Target in Neuroblastoma. *JCO Precis Oncol*, **3**, 1-20.
1307   47.   Hartlieb, S.A., Sieverling, L., Nadler-Holly, M., Ziehm, M., Toprak, U.H., Herrmann, C.,
1308   Ishaque, N., Okonechnikov, K., Gartlgruber, M., Park, Y.G. *et al.* (2021) Alternative
1309   lengthening of telomeres in childhood neuroblastoma from genome to proteome. *Nat
1310   Commun*, **12**, 1269.
1311   48.   Porro, A., Feuerhahn, S., Reichenbach, P. and Lingner, J. (2010) Molecular dissection
1312   of telomeric repeat-containing RNA biogenesis unveils the presence of distinct and
1313   multiple regulatory pathways. *Mol Cell Biol*, **30**, 4808-4817.
1314   49.   Wu, B., Su, S., Patil, D.P., Liu, H., Gan, J., Jaffrey, S.R. and Ma, J. (2018) Molecular basis
1315   for the specific and multivariant recognitions of RNA substrates by human hnRNP
1316   A2/B1. *Nat Commun*, **9**, 420.
1317   50.   Martinez, F.J., Pratt, G.A., Van Nostrand, E.L., Batra, R., Huelga, S.C., Kapeli, K., Freese,
1318   P., Chun, S.J., Ling, K., Gelboin-Burkhart, C. *et al.* (2016) Protein-RNA Networks

Regulated by Normal and ALS-Associated Mutant HNRNPA2B1 in the Nervous System.
Neuron, **92**, 780-795.

51.   Linder, B., Grozhik, A.V., Olarerin-George, A.O., Meydan, C., Mason, C.E. and Jaffrey,
S.R. (2015) Single-nucleotide-resolution mapping of m6A and m6Am throughout the
transcriptome. *Nat Methods*, **12**, 767-772.

52.   Petti, E., Buemi, V., Zappone, A., Schillaci, O., Broccia, P.V., Dinami, R., Matteoni, S.,
Benetti, R. and Schoeftner, S. (2019) SFPQ and NONO suppress RNA:DNA-hybrid-
related telomere instability. *Nat Commun*, **10**, 1001.

53.   Nguyen, E.D., Balas, M.M., Griffin, A.M., Roberts, J.T. and Johnson, A.M. (2018) Global
profiling of hnRNP A2/B1-RNA binding on chromatin highlights LncRNA interactions.
RNA Biol, **15**, 901-913.

54.   Liu, N., Dai, Q., Zheng, G., He, C., Parisien, M. and Pan, T. (2015) N(6)-
methyladenosine-dependent RNA structural switches regulate RNA-protein
interactions. *Nature*, **518**, 560-564.

55.   Duda, K.J., Ching, R.W., Jerabek, L., Shukeir, N., Erikson, G., Engist, B., Onishi-
Seebacher, M., Perrera, V., Richter, F., Mittler, G. *et al.* (2021) m6A RNA methylation
of major satellite repeat transcripts facilitates chromatin association and RNA:DNA
hybrid formation in mouse heterochromatin. *Nucleic Acids Res*, **49**, 5568-5587.

56.   Montero, J.J., Lopez-Silanes, I., Megias, D., M, F.F., Castells-Garcia, A. and Blasco, M.A.
(2018) TERRA recruitment of polycomb to telomeres is essential for histone
trymethylation marks at telomeric heterochromatin. *Nat Commun*, **9**, 1548.

57.   Kim, D., Paggi, J.M., Park, C., Bennett, C. and Salzberg, S.L. (2019) Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*, **37**, 907-
915.

58.   Tarasov, A., Vilella, A.J., Cuppen, E., Nijman, I.J. and Prins, P. (2015) Sambamba: fast
processing of NGS alignment formats. *Bioinformatics*, **31**, 2032-2034.

59.   Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
Singh, H. and Glass, C.K. (2010) Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B
cell identities. *Mol Cell*, **38**, 576-589.

60.   Ramirez, F., Dundar, F., Diehl, S., Gruning, B.A. and Manke, T. (2014) deepTools: a
flexible platform for exploring deep-sequencing data. *Nucleic Acids Res*, **42**, W187-
191.

61.   Lawrence, M., Gentleman, R. and Carey, V. (2009) rtracklayer: an R package for
interfacing with genome browsers. *Bioinformatics*, **25**, 1841-1842.

62.   Kent, W.J., Zweig, A.S., Barber, G., Hinrichs, A.S. and Karolchik, D. (2010) BigWig and
BigBed: enabling browsing of large distributed datasets. *Bioinformatics*, **26**, 2204-
2207.

63.   Li, H. (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*,
34, 3094-3100.

64.   Li, H. (2021) New strategies to improve minimap2 alignment accuracy. *Bioinformatics*,
37, 4572-4574.

65.   Kovaka, S., Zimin, A.V., Pertea, G.M., Razaghi, R., Salzberg, S.L. and Pertea, M. (2019)
Transcriptome assembly from long-read RNA-seq alignments with StringTie2. *Genome
Biol*, **20**, 278.

66.   Lee, S., Cook, D. and Lawrence, M. (2019) plyranges: a grammar of genomic data
transformation. *Genome Biol*, **20**, 4.

67. Pockrandt, C., Alzamel, M., Iliopoulos, C.S. and Reinert, K. (2020) GenMap: ultra-fast
computation of genome mappability. *Bioinformatics*, **36**, 3687-3692.
68. Quinlan, A.R. and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics*, **26**, 841-842.
69. Feuerbach, L., Sieverling, L., Deeg, K.I., Ginsbach, P., Hutter, B., Buchhalter, I.,
Northcott, P.A., Mughal, S.S., Chudasama, P., Glimm, H. *et al.* (2019) TelomereHunter
- in silico estimation of telomere content and composition from cancer genomes. *BMC*
Bioinformatics, **20**, 272.
70. Vaid, R., Wen, J. and Mannervik, M. (2020) Release of promoter-proximal paused Pol
II in response to histone deacetylase inhibition. *Nucleic Acids Res*, **48**, 4877-4890.
71. Mondal, T., Subhash, S., Vaid, R., Enroth, S., Uday, S., Reinius, B., Mitra, S.,
Mohammed, A., James, A.R., Hoberg, E. *et al.* (2015) MEG3 long noncoding RNA
regulates the TGF-beta pathway genes through formation of RNA-DNA triplex
structures. *Nat Commun*, **6**, 7743.
72. Caren, H., Kryh, H., Nethander, M., Sjoberg, R.M., Trager, C., Nilsson, S., Abrahamsson,
J., Kogner, P. and Martinsson, T. (2010) High-risk neuroblastoma tumors with 11q-
deletion display a poor prognostic, chromosome instability phenotype with later
onset. *Proc Natl Acad Sci U S A*, **107**, 4323-4328.
73. Fransson, S., Ostensson, M., Djos, A., Javanmardi, N., Kogner, P. and Martinsson, T.
(2016) Estimation of copy number aberrations: Comparison of exome sequencing data
with SNP microarrays identifies homozygous deletions of 19q13.2 and CIC in
neuroblastoma. *Int J Oncol*, **48**, 1103-1116.
74. Kocak, H., Ackermann, S., Hero, B., Kahlert, Y., Oberthuer, A., Juraeva, D., Roels, F.,
Theissen, J., Westermann, F., Deubzer, H. *et al.* (2013) Hox-C9 activates the intrinsic
pathway of apoptosis and is associated with spontaneous regression in
neuroblastoma. *Cell Death Dis*, **4**, e586.
75. Cun, Y., Yang, T.P., Achter, V., Lang, U. and Peifer, M. (2018) Copy-number analysis and
inference of subclonal populations in cancer genomes using Sclust. *Nat Protoc*, **13**,
1488-1501.
76. Zhang, W., Yu, Y., Hertwig, F., Thierry-Mieg, J., Zhang, W., Thierry-Mieg, D., Wang, J.,
Furlanello, C., Devanarayan, V., Cheng, J. *et al.* (2015) Comparison of RNA-seq and
microarray-based models for clinical endpoint prediction. *Genome Biol*, **16**, 133.
77. Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years
of image analysis. *Nat Methods*, **9**, 671-675.

1400

Figure legends:

Figure 1: (**A**) m^{6}A RNA Immunoprecipitation (m^{6}A RIP) on RNA isolated from U-2 OS cells

followed by slot blot, probed with DIG-labeled TERRA probe. RNase A was used as a control.

m^{6}A RIP in fragmented and non-fragmented RNA (**B**) Schematic diagram describing key steps

of m^{6}A RIP performed on either fragmented or non-fragmented RNA isolated from Control-sh

or METTL3-KD U-2 OS cells. (**C**) (top panel) m^{6}A RIP followed with fragmented and non-

fragmented RNA followed by slot blot with DIG-labeled TERRA repeat probe. (Middle panel)

Quantification of the slot blot. Unpaired *t*-test was used, **** $p < 0.0001$. (Lower panel)

Western blot showing METTL3-KD.

Detection of UUAG tetramer in TERRA RNA by RNA-MS analysis (**D**) Schematic diagram

describing key steps of TERRA pulldown and RNA-MS analysis (**E**) Slot blot showing the

specificity of TERRA pulldown**.** Probe against Luciferase (negative pulldown) and RNase A

was used as a control. (**F**) [**left panel**] Base peak chromatogram (BPC): first and second row

show BPC of Standard UUAGp, isolated m/z 651.07 (doubly charged) and Standard

UUm^{6}AGp, isolated m/z 658.08 (doubly charged) respectively. Third and fourth rows show

the BPC of the TERRA pulldown sample, isolated m/z 651.07 (doubly charged) and isolated

m/z 658.08 (doubly charged) respectively. [**Right panel**] MS2 spectrum: first and second rows

show the MS2 spectrum of Standard UUAGp, isolated m/z 651.07, and Standard UUm^{6}AGp,

isolated m/z 658.08 respectively with ions of mass difference 14 (methylation) starting from

m/z 673 and up. Y-axis represents relative abundance in percentage. Third and fourth rows

show the MS2 spectrum of the sample, isolated m/z 651.07 and isolated m/z 658.08

respectively with ions of mass difference 14 (methylation) starting from m/z 673 and up. Y-

axis represents absolute abundance.

(**G**) Proximity ligation assay (PLA) in U-2 OS cells expressing TRF1-mCherry depicting the

interaction of TRF2 with METTL3 (left) and METTL14 (right). TRF1-mCherry (Red), PLA

foci (green) in the nucleus (marked by DAPI in blue). Box plot shows the number of co-

localizations between TRF1-mCherry and PLA signal.

(**H**) TERRA foci (green) in U-2 OS cells expressing TRF1-mCherry (red). Box plot shows the

number of co-localizations between TRF1-mCherry and TERRA.

(**I**) m^{6}A immunofluorescence (IF) (red) was performed in U-2 OS along with TERRA foci

(green). Box plot shows the number of co-localizations between TERRA and m^{6}A.

Scale bar is 10 µm.

1433

Figure 2: (**A**) Flow chart depicting the major step of m^{6}A RIP-seq along with RNA-seq

performed either on Illumina platform or on Oxford Nanopore Technologies (ONT), to

generate short and long RNA-seq reads respectively. (**B**) Browser screenshot of 50 kb region

around the telomere ends in T2T assembly, showing TERRA RNA expression (CPM) from

two active chromosome ends (Chr12p, Chr20q) and one less active chromosome end (Chr3p)

using ONT long read and Illumina short-read RNA-seq. RNA Pol II (purple) and CTCF (pink)

enrichment at 50 kb region around the telomere ends from publicly available RNA Pol II and

CTCF ChIP-seq data which were re-analyzed with T2T human genome assembly. CpG islands

are marked with green bars and the telomeric repeats in this region are marked with black bars.

(**C**) Heatmap summarizing RNA-seq and m^{6}A RIP-seq data. Chromosomes are sorted based

on high (dark violet-pink) and low (light violet-pink) subtelomeric TERRA transcription. Black

dots denote m^{6}A RIP-seq peaks and black diamonds denote the differential modified m^{6}A sites

detected by xPore from ONT long reads at subtelomeres.

(**D**) Genome browser screenshots showing IP/input ratio tracks for m^{6}A RIP (Control-sh and

METTL3-KD) or IgG RIP (Control-sh) at two active chromosome ends (Chr12p, Chr20q) and

one inactive chromosome end (Chr3p). Red bars mark m^{6}A RIP peaks identified using MACS

(for short reads) or the differential m^{6}A sites identified using xPore (for long reads) for the

samples as denoted.

(**E**) The m^{6}A enrichment profiles across all subtelomeres in T2T assembly in Control-sh and

METTL3-KD. The m^{6}A RIP/input signals were normalized to spike-in and to the mappability

likelihood of each chromosome ends. Subtelomeres (high and low TERRA transcription) were

classified according to the normalized median expression are shown. (**F**) Box plot of spike-in,

mappability-normalized m^{6}A RIP/input enrichment profiles of all chromosome ends. Statistical

significance was calculated using the Wilcoxon test, **** $p < 0.0001$.

1458

Figure 3: (**A**) TERRA foci (green) in U-2 OS cells after siRNA mediated KD of METTL3.

Box plot showing the number of TERRA foci per nucleus. Dunnett's multiple comparisons test

was used, ** $p < 0.01$; **** $p < 0.0001$. (**B**) (top panel): Western blot showing METTL3 level

in stable shRNA-depleted U-2 OS cells. (Middle panel) TERRA foci (green) in U-2 OS cells

with stable KD of METTL3. (bottom panel): Box plot shows the number of TERRA foci per

nucleus. Dunnett's multiple comparisons test was used, **** $p < 0.0001$.

(**C**) Localization of the γ-H2AX (green) over telomere in Control-sh and METTL3-KD cells.

Telomere is detected by TRF2 staining (red). Box plot shows the number of overlaps of TRF2

and γ-H2AX. Dunnett's multiple comparisons test was used, **** $p < 0.0001$.

(**D**) Localization of γ-H2AX (green) over telomere (marked by TRF2) in bleomycin-treated

Control and METTL3-KD cells 4 h post-treatment and followed by 20 h of recovery without

bleomycin. Box plot shows the number of overlaps of TRF2 and γ-H2AX. Unpaired *t*-test were

used, * $p < 0.05$; ** $p < 0.01$. (**E**) m^{6}A IF (red) along with TERRA foci (green) was performed

in U-2 OS cells 4 h post-treatment and followed by 20 h of recovery without bleomycin. Box

plot shows the number of overlaps of TERRA and m^{6}A signal. Unpaired *t*-test was used, ** *p*

< 0.01.

(**F**) Cartoon demonstrating recruitment of dCasRx-FTO to target RNA. (**G**) m^{6}A RIP followed

by TERRA slot blot on RNA isolated from U-2 OS cells expressing dCasRx-FTO with either

non-target control (NTC) or TERRA guide RNA. IgG antibody was used as a negative control

for the RIP experiment. Bar graph shows the quantification of slot blot. Unpaired *t*-test was

used, **** *p* < 0.0001. (**H**) TERRA RNA-FISH detecting TERRA foci (green) was performed

on U-2 OS cells expressing dCasRx-FTO with either NTC or TERRA guide RNA. Box plot

shows the quantification of TERRA foci per nucleus. Unpaired *t*-test was used, **** *p* <

0.0001. (**I**) TERRA foci (green) in U-2 OS cells treated with various METTL3 inhibitors for 6

h. Box plot shows the quantification of TERRA foci per nucleus. Dunnett's multiple

comparisons test was used, **** *p* < 0.0001. Scale bar is 10 μm.

1485

Figure 4:

(**A**) TERRA foci (green) in U-2 OS cells with siRNA mediated transient KD of m^{6}A reader

proteins as denoted. Box plot shows the number of TERRA foci per nucleus. Dunnett's multiple

comparisons test was used, ** *p* < 0.01; **** *p* < 0.0001.

(**B**) hnRNPA2B1 RIP followed by TERRA slot blot demonstrates the interaction of

hnRNPA2B1 with TERRA RNA. IgG and RNase A treatment were used as control.

(**C**) Scheme and results of the re-RIP (hnRNPA2B1-m^{6}A) experiment in Control-sh or

METTL3-KD U-2 OS cells. First RIP was performed with hnRNPA2B1/IgG antibody, the

precipitated RNA was further subjected to RIP with m^{6}A antibody. Finally, the RNA after re-

RIP was blotted and probed with the TERRA probe. RIP with IgG served as a negative control.

Bar graph shows the quantification of the slot blot. Sidak's multiple comparisons test was used,

** *p* < 0.01; **** *p* < 0.0001.

(**D**) IF showing recruitment of dCasRx-FTO at TERRA but not to NTC leads to reduced

hnRNPA2B1(green) occupancy at telomeres (TRF2, red). Box plot shows the number of

overlaps of the hnRNPA2B1 and TRF2. Unpaired *t*-test was used, *** *p* < 0.001. (**E**) γ-H2AX

(green) accumulation at telomeres (TRF2, red) in U-2 OS cells with siRNA mediated KD of

hnRNPA2B1. Box plot shows the number of overlaps of the γ-H2AX and TRF2. Dunnett's

multiple comparisons test was used, * *p* < 0.05; ** *p* < 0.01.

Scale bar is 10 μm.

1505

Figure 5:

(**A**) Schematic diagram describing the DRIP and DRImR method to purify R-Loop bound RNA and R-loop bound m^{6}A modified RNA respectively. (**B**) Blot on the left is from RNA purified in the DRIP experiment and the blot on the right is from the DRImR experiment performed on U-2 OS cells. Experiments performed with IgG antibody or nucleic acid pre-treated with RNase H served as a control. RNase A treatment was done to affirm the presence of only RNA in the final elution in the lanes indicated. Blots were probed with a DIG-labeled TERRA probe. (**C**) RNA enriched in R-Loops from Control-sh or METTL3-KD U-2 OS cells were isolated by the DRIP method followed by slot blot assay. Blot was probed with a DIG-labeled TERRA probe. DRIP performed with IgG antibody or pre-treatment with RNase H served as a control. RNase A treatment was done to affirm the presence of only RNA in the final elution. Bar graph shows the quantification of the blots. Unpaired t-test was used, **** $p < 0.0001$.

(**D**) TERRA RNA-FISH detecting TERRA foci (green) in Control-sh or METTL3-KD U-2 OS cells that were transfected with either Control siRNA or siRNAs against RNase H. Box plot shows the quantification of TERRA foci per nucleus. Tukey's multiple comparisons test was used, **** $p < 0.0001$.

R-loop PLA assay (**E**) PLA in U-2 OS TRF1-mCherry cells with overexpression of either RNase H^{WT} (left) or RNase H^{Mut} (right) depicting the interaction of TRF2 with R-Loop (S9.6 antibody). TRF1-mCherry (Red), PLA foci (green) in the nucleus (marked by DAPI). Box plot shows the number of overlaps of PLA signal and TRF1-mCherry per nucleus. Unpaired t-test was used, **** $p < 0.0001$. (**F, G**) PLA depicting the interaction of TRF2 with R-Loop (S9.6 antibody) in U-2 OS TRF1-mCherry cells with (**F**) Control-sh or METTL3-KD (**G**) Control siRNA or hnRNPA2B1 siRNA2. TRF1-mCherry (Red), PLA foci (green) in the nucleus (marked by DAPI). Box plot shows the number of overlaps of PLA signal and TRF1-mCherry per nucleus. Unpaired t-test was used, **** $p < 0.0001$.

Scale bar is 10 μm.

Figure 6: (**A**) Profiling TERRA RNA expression in ALT+/- cell lines by slot blot assay performed with 100, 200 and 400 ng of RNA isolated from cell lines indicated. Blot probed with a DIG-labeled TERRA probe. TapeStation profile on the right shows 18s and 28s rRNA served as a loading control. Bar graph shows the TERRA level quantified from blots and normalized to loading control.

(**B**) m^{6}A RIP followed by TERRA slot blot on RNA isolated from SK-N-F1 (ALT+), SK-N-BE(2), and IMR-32 (ALT-) cell lines. RIP with IgG severed as a negative control. Bar graph

shows the quantification of the blots. Sidak's multiple comparisons test was used, **** $p <$ 0.0001.

(**C**) TERRA RNA foci (green) visualized by TERRA RNA-FISH in ALT+ (upper panel) and ALT- (lower panel) NB cell lines. Box plot shows the quantification of the TERRA foci per nucleus.

(**D**) TERRA foci (green) combined with m^{6}A (red) IF in CHLA-90 (left) and SK-N-F1 (right) cells. Box plot shows the number of overlaps of TERRA and m^{6}A per nucleus. Scale bar is 10 μm.

(**E**) The m^{6}A enrichment profiles across all subtelomeres using T2T assembly in SK-N-F1 and SK-N-BE(2) cells. The m^{6}A RIP/input signals were normalized to spike-in and to the mappability likelihood of each chromosome ends. (**F**) Box plot of spike-in, mappability-normalized m^{6}A RIP/input enrichment profiles of all subtelomeres in SK-N-F1 and SK-N-BE(2). Statistical significance was calculated using the Wilcoxon test, **** $p < 0.0001$.

(**G**) TERRA RNA expression in ALT+/- NB patient tumor samples were detected by slot blot assay performed with 100 ng of RNA. The tumor profile indicates genomic alteration. These tumors were also assayed for C-Circle (CCA).

(**H**) Representative browser screenshot of 50 kb region around the telomere ends, showing TERRA RNA expression (CPM) from two of the active chromosome ends (Chr9p, Chr16q) and one inactive chromosome end (Chr3p) using Illumina short read RNA-seq performed on NB1 and NB2 tumor samples. CpG islands are marked with green bars and the telomeric repeats in this region are marked with black bars.

(**I**) Heatmap summarizing RNA-seq and m^{6}A RIP-seq data of two NB tumor samples (NB1 and NB2). Chromosomes are sorted based on high (dark colored) and low (light colored) subtelomeric TERRA transcription. Black dots denote m^{6}A RIP-seq peaks identified using MACS peak caller in these NB tumor samples.

(**J**) Genome browser screenshots showing m^{6}A RIP/input ratio tracks for NB1 and NB2 tumor samples at two active chromosome ends (Chr12p, Chr20q) and one inactive chromosome end (Chr3p). Red bars mark m^{6}A RIP peaks identified using MACS peak caller.

(**K**) The m^{6}A enrichment profiles across subtelomeres using T2T assembly in NB1 and NB2 tumors. The m^{6}A RIP/input signals were normalized to spike-in and to the mappability likelihood of each chromosome ends. Subtelomeres (high and low TERRA transcription) were classified according to the normalized median expression are shown.

(**L**) Event-free survival of NB patients (n=498, SEQC cohort) with either low (blue) or high (red) expression of METTL3.

1574   (**M**) Event-free survival of ALT+ NB patients (n=21) with either low (blue) or high (red)

1575   expression of METTL3.

1576   (**N**) Bar graph showing percentage of ALT+ or MNA (MYCN amplified) tumors with copy

1577   number gain in METTL3, METTL14, or hnRNPA2B1 genes.

1578

Figure 7: TERRA foci following siRNA mediated METTL3 depletion in ALT+ NB cells (**A**)

(left panel) Western blot for METTL3 level after siRNA transfection in CHLA-90 cells.

(middle panel) TERRA RNA-FISH detecting TERRA foci (green) in CHLA-90 and SK-N-F1

cells after METTL3 knockdown as indicated. (right panel) Box plots show the quantification

of TERRA foci per nucleus. Dunnett's multiple comparisons test was used, * $p < 0.05$; ** $p <$

0.01; **** $p < 0.0001$.

(**B**) Representative images from colony formation assay performed on CHLA-90 (top) and SK-

N-F1(bottom) cells with TetO Control or METTL3-KD.

(**C**) TERRA RNA FISH in TetO Control and METTL3-KD SK-N-F1 and CHLA-90 cells. Box

plot shows the quantification of the TERRA foci in the indicated conditions. Unpaired *t*-test

was used, *** $p < 0.001$.

(**D**) Accumulation of γ-H2AX (red) over telomere after 48 h of Dox induction in SK-N-F1

TetO METTL3-KD cells. Telomeres were detected by telomere DNA FISH (green). Box plot

shows the overlap of telomere and γ-H2AX. Unpaired *t*-test was used, **** $p < 0.0001$.

(**E**) (top panel) Line graph showing average tumor growth (mm^3) of SK-N-F1 cells (TetO

Control and METTL3-KD) derived xenografts over time. Data are shown as Mean ± SD.

Sidak's multiple comparisons test was used, **** $p < 0.0001$. (bottom panel) plot showing

tumor weight post necropsies of xenografts derived from SK-N-F1 cells with either TetO

Control or METTL3-KD. Data are shown as Mean ± SD. Unpaired *t*-test was used, * $p < 0.05$

(n=4). One representative image of the tumor is shown per condition.

(**F**) Western blot to verify the hnRNPA2B1 KD in ALT+ NB cells. Vinculin was used as a

loading control. (**G**) TERRA foci (green) in CHLA-90 (top) and SK-N-F1 (bottom) cells after

transient KD of hnRNPA2B1. Box plot shows the quantification of TERRA foci per nucleus.

Unpaired *t*-test was used, ** $p < 0.01$; **** $p < 0.0001$.

(**H**) TERRA RNA-FISH detecting TERRA foci (green) in CHLA-90 (top panel) and SK-N-F1

(bottom panel) cells treated with METTL3 inhibitors (10 μM STM2457, 50 μM UZH1A, and

50 μM UZH1B) for 6 h. DMSO treatment served as a control. Box plot showing quantification

of the TERRA foci as indicated. Dunnett's multiple comparisons test was used, * $p < 0.05$; ***

$p < 0.001$; **** $p < 0.0001$.

(**I**) Telomere DNA FISH (green) and γ-H2AX IF (red) in SK-N-F1 cells treated with METTL3

inhibitor, STM2457 for 48 h. Box plot showing quantification of telomere and γ-H2AX

overlap. Unpaired *t*-test was used, **** $p < 0.0001$. Scale bar is 10 μm.

(**J**) Cell viability was measured by MTT assay after treating CHLA-90 and SK-N-F1 cells with

10 μM of METTL3 inhibitor (STM2457) at indicated time points. Data are normalized to

DMSO control. Sidak's multiple comparisons test was used, **** $p < 0.0001$.

1614

Supplementary Figure legends:

Supplementary Figure S1:

(**A**) TapeStation profile of RNA isolated from Control-sh or METTL3-KD U-2 OS cells which

is either fragmented or non-fragmented. Serving as a control for Figure 1C.

(**B**) Schematic diagram shows *In vitro* METTL3/METTL14 methyltransferase assay performed

with DRACH oligo or TERRA repeat containing oligo. A and m^{6}A were detected and

quantified by LC-MS/MS-based method. RNA oligo Standard with or without m^{6}A was used

as a control. Bar graph shows the ratio of A/m^{6}A in the indicated conditions. Unpaired *t*-tests

were used, * $p < 0.05$; **** $p < 0.0001$.

(**C**) Background control for PLA with only TRF2 and only METTL3 in U-2 OS cells

expressing TRF1-mCherry in the nucleus (marked by DAPI).

(**D**) Chromatin Immunoprecipitation (ChIP) using METTL3 antibody on U-2 OS cells

followed by slot blot. Blot probed with a DIG-labeled TERRA probe. IgG antibody was used

as a negative control.

(**E**) TERRA foci (green) in U-2 OS cells treated with either Scrambled or TERRA LNA. Box

plot shows the quantification of TERRA signal per nucleus. Unpaired *t*-test was used, ** $p <$

0.01.

Scale bar is 10 μm.

1633

Supplementary Figure S2:

(**A**) Genome browser tracks of normalized CPM coverage obtained from Strand-specific

Illumina short-reads in U-2 OS cells showing transcription away from telomere ends (direction

of transcription is opposite to TERRA). Reads were mapped to the positive strand on p arms,

and the negative strand on q arms, to match the direction of transcription, are shown for two

active chromosome ends (Chr12p, Chr20q) and one inactive chromosome end (Chr3p).

(**B-C**) Characterizing the high and low TERRA RNA expressing subtelomeres using

normalized read counts from either (**B**) Illumina short reads or (**C**) ONT-long reads. Read

counts were normalized using CPM and to the mappability likelihood of each chromosome

ends. Statistical significance was calculated using the two-sided paired *t*-test.

(**D**) Slot blot with total RNA isolated from Control-sh or METTL3-KD U-2 OS cells. Blot

probed with a DIG-labeled TERRA probe. TapeStation profile showing 18s and 28s rRNA

served as a loading control. Bar graph shows the quantification of the blot normalized to the

respective loading control. Unpaired *t*-test was used, ** $p < 0.01$.

(**E**) Metagene plot showing density distribution of m^{6}A peak or m^{6}A sites per bin from both

m^{6}A RIP peaks (red) and m^{6}A xPore sites (blue). (**F**) Identified top enriched motif logo from

de novo motif analysis of m^{6}A peaks from m^{6}A RIP-seq and m^{6}A xPore sites. (**G**) Bar plot

summarizing peaks identified genome-wide in RIP-seq experiment. (**H**) Telomeric repeat

reads enriched in m^{6}A/IgG RIP-seq using TelomereHunter.

(**I**) Venn diagram showing the overlap between m^{6}A RIP peaks (red) and m^{6}A xPore sites

(blue) at the 3' UTR.

(**J**) Enrichment of m^{6}A RIP/input signal at DRACH motif coordinates located within the

subtelomeric m^{6}A RIP-seq peaks in Control-sh and METTL3-KD U-2 OS cells. Statistical

significance was calculated using the two-sided paired *t*-test. ****: $p < 0.0001$.

(**K**) Scatter plot depicting differentially modified sites identified using xPore from Control-sh

and METTL3-KD U-2 OS long-reads. Predicted modified k-mers within the 30 kb

subtelomeres regions showing a positive modification rate in Control-sh within the NNANN

context being differentially modified as compared to METTL3-KD are shown.

1662

Supplementary Figure S3:

(**A**) IF showing METTL3-KD following METTL3 siRNA treatment. Bar graph shows the

quantification of the images as indicated. Dunnett's multiple comparisons test was used, ** *p*

< 0.01. (**B**) (left panel) Western blot to verify METTL14 siRNA mediated knockdown in U-2

OS cells. Vinculin was used as a loading control. (middle panel) TERRA foci (green) in U-2

OS cells within Control or METTL14 siRNA treated cells. Nucleus marked by DAPI. (right

panel) Box plot shows TERRA signal per nucleus in the condition indicated. Dunnett's multiple

comparisons test was used, *** $p < 0.001$; **** $p < 0.0001$.

(**C**) Localization of the γ-H2AX (green) over telomere in Scrambled or TERRA LNA cells.

Telomere is detected by TRF2 staining (red). Box plot shows the overlap of TRF2 and γ-H2AX.

Unpaired *t*-test was used, **** $p < 0.0001$.

(**D**) IF of TRF2 (red) and PML bodies (green) in control and METTL3-KD U-2 OS cells. Box

plot shows the overlap of PML and TRF2. Dunnett's multiple comparisons test was used, * *p*

< 0.05. Scale bar is 10 μm.

(**E**) (Top panel) Slot blot with total RNA isolated from U-2 OS cells expressing dCasRx-FTO

with either NTC or TERRA guide RNA. RNA was loaded in 3 different amounts100, 200, and

400 ng. Blot probed with a DIG-labeled TERRA probe. TapeStation profile showing 18s and

28s rRNA served as a loading control. Bar graph shows the quantification of the blot

normalized to loading control. Unpaired *t*-test was used. (Bottom panel) Western blot to verify

induction of HA-tagged dCasRx-FTO by Dox. GAPDH is loading control.

(**F**) (Top panel) Slot blot with total RNA isolated from U-2 OS cells expressing catalytically

active CasRx-HA with either NTC or TERRA guide RNA. RNA was loaded in 2 different

amounts100 ng and 200 ng. Blot probed with DIG-labeled TERRA repeat-containing probe.

TapeStation profile showing 18s and 28s rRNA served as a loading control. Bar graph shows

the quantification of the blot normalized to loading control. Unpaired *t*-test was used, *** *p* $<$

0.001. (Bottom panel) Western blot to verify induction of CasRx-HA by Dox. GAPDH is

loading control.

(**G**) TERRA slot blot with total RNA isolated from U-2 OS cells treated with Actinomycin D

for the indicated time points.

(**H**) TERRA slot blot with total RNA isolated from U-2 OS cells treated with METTL3

inhibitor, STM2457. DMSO served as a control for STM2457. TapeStation profile showing

18s and 28s rRNA served as a loading control. Bar graph shows the quantification of the blot

normalized to loading control. Unpaired *t*-test was used.

1696

Supplementary Figure S4:

(**A**) Relative mRNA expression of m^6A reader proteins- hnRNPA2B1, YTHDC1, RBMX, and

YTHDF2 that were knocked down with siRNA. GAPDH was used to normalize the qPCR data.

Dunnett's multiple comparisons test was used, ** $p < 0.01$; **** $p < 0.0001$.

(**B**) Slot blot with total RNA isolated from U-2 OS cells with either control or hnRNPA2B1

siRNA. RNA was loaded in 2 different amounts 100 and 200 ng. Blot probed with a DIG-

labeled TERRA probe. TapeStation profile showing 18s and 28s rRNA served as a loading

control. Bar graph shows the quantification of the blot normalized to loading control. Unpaired *t*-test was used.

(**C**) *In vitro* binding assay between total RNA isolated from Control-sh or METTL3-KD U-2 OS cells (2 different RNA concentrations) with purified His-tagged hnRNPA2B1. Left panel show commissive staining of purified hnRNPA2B1 bound to beads. Right panel, hnRNPA2B1 bound RNA fraction on slot blot probed with DIG-labeled TERRA probe. Bar graph shows the quantification of the blots. Unpaired *t*-test was used, ** $p < 0.01$.

Supplementary Figure S5:

(**A**) TERRA RNA-FISH detecting TERRA foci (green) in U-2 OS cells exogenously treated with either mock, RNase A, RNase H, or with both RNase A and RNase H. Box plot shows the number of TERRA foci per nucleus. Dunnett's multiple comparisons test was used, **** $p < 0.0001$.

(**B**) Relative RNase H expression following siRNA mediated KD. GAPDH was used to normalize the qPCR data. Dunnett's multiple comparisons test was used, **** $p < 0.0001$.

(**C**) Nucleic acid isolated from U-2 OS cells overexpressing RNase H^{WT} or RNase H^{Mut} were immobilized onto a nitrocellulose membrane and immunoblotted with S9.6 antibody. Bar graph is the quantification of the blot. Unpaired *t*-test was used, **** $p < 0.0001$.

(**D**) TERRA foci (green) in U-2 OS cells overexpressing RNase H^{WT} (left) or RNase H^{Mut} (right). Box plot shows the quantification of the TERRA foci per nucleus. Unpaired *t*-test was used, **** $p < 0.0001$.

(**E**) Background control for PLA with only TRF2 (left); or only S9.6 (right) antibody in RNase H^{Mut} U-2 OS cells expressing TRF1-mCherry.

(**F**) PLA depicting the interaction of TRF2 with R-loop (S9.6 antibody) in U-2 OS TRF1-mCherry cells expressing dCasRx-FTO with either NTC (left) or TERRA guide RNA (right). Box plot shows the quantification of the overlapping PLA and TRF1-mCherry signal. Unpaired *t*-test was used, ** $p < 0.01$. Scale bar is 10 μm.

Supplementary Figure S6:

(**A**) C-Circle assay (CCA) was performed on DNA from ALT+ and ALT- NB cells. Bar graph shows the quantification of the blots.

(**B**) TERRA foci (green) in ALT+ CHLA-90 (top) and SK-N-F1 (bottom) cells treated with either Scrambled or TERRA LNA. Scale bar is 10 μm.

(**C**) Subtelomeric TERRA expression (CPM) in SK-N-F1 and SK-N-BE(2) cells using
mappability normalized RNA-seq data. Statistical significance was calculated using the two-
sided paired *t*-test, **** $p < 0.0001$.
(**D**) Top enriched motif logo from de novo motif analysis of m^{6}A peaks from SK-N-F1 and
SK-N-BE(2) m^{6}A RIP-seq data.
(**E**) C-Circle assay (CCA) was performed on DNA from NB tumors.
(**F**) Subtelomeres are characterized as high and low on TERRA expression (CPM) in NB1 and
NB2 tumors using mappability normalized RNA-seq data. Statistical significance was
calculated using the two-sided paired *t*-test, *** $p < 0.001$.
(**G**) Top enriched motif logo from de novo motif analysis of NB1 and NB2 tumor m^{6}A peaks.
(**H**) Event-free survival of NB patients (n=498, SEQC cohort) with either low (blue) or high
(red) expression of METTL14 and hnRNPA2B1.
(**I**) Event-free survival of ALT+ NB patients (n=21) with either low (blue) or high (red)
expression of METTL14 and hnRNPA2B1.
1751
Supplementary Figure S7:
(A)Western blot showing METTL3 level in TetO Control and METTL3-KD ALT+ NB cells
following Dox induction for 48 h. GAPDH was loading control. (**B**) Bar graph displaying the
effect on cell viability after conditional KD of METTL3 (6 days post Dox treatment) in CHLA-
90 and SK-N-F1 TetO METTL3-KD cells. Data normalized to control and shown as Mean ±
SD. Unpaired *t*-test was used, **** $p < 0.0001$.
(**C**) Slot blot with total RNA isolated from SK-N-F1 cells with TetO Control or METTL3-KD.
RNA was loaded in 2 different amounts 200 and 400 ng. Blot probed with a DIG-labeled
TERRA probe. TapeStation profile showing 18s and 28s RNA served as a loading control. Bar
graph showing quantification normalized to loading control. Unpaired *t*-test was used.
(**D**) Accumulation of γ-H2AX (red) over telomere after 48 h of Dox induction in CHLA-90
cells with TetO METTL3-KD. Telomere was detected by telomere DNA FISH (green). Box
plot shows the quantification of γ-H2AX, and telomere overlap. Unpaired *t*-test was used, ****
$p < 0.0001$.
(**E**) C-circle assay results visualized on a slot blot. DNA isolated from CHLA-90 and SK-N-
F1 cells with TetO METTL3-KD (48 h Dox treatment) was subjected to a C-circle assay
with/without Phi29 polymerase (Pol+/Pol-). Bar graph shows the quantification of the blots.
Sídak's multiple comparisons test was used, ** $p < 0.01$; *** $p < 0.001$.

1770

(**F**) DRIP followed by slot blot assay and probed with DIG-labeled TERRA probe in TetO

Control or METTL3-KD SK-N-F1 cells, 72 h post-Dox induction. Experiments performed

with IgG antibody or nuclei acid pre-treated with RNase H served as a control.

(**G**) (Left panel) Line graph showing average tumor growth (mm^3) of CHLA-90 cells (TetO

Control and METTL3-KD) derived xenografts over time. Data are shown as Mean ± SD.

Sidak's multiple comparisons test was used, **** $p < 0.0001$. (Middle panel) Tumor weight

post necropsies of xenografts derived from CHLA-90 cells with either TetO Control or

METTL3-KD Data are shown as Mean ± SD. Unpaired t-test was used, ** $p < 0.01$ (n=3). One

representative image of the tumor is shown per condition. (Right panel) Western blot confirms

the METTL3-KD in xenograft tumors and GAPDH acted as a loading control.

(**H**) TERRA RNA-FISH detecting TERRA foci (green) combined with hnRNPA2B1 (red) IF

in CHLA-90 cells. Box plot shows the quantification of TERRA and m^6A overlap.

(**I**) Telomere DNA FISH (green) and γ-H2AX IF (red) in ALT- SK-N-BE(2) cells treated with

METTL3 inhibitor, STM2457 for 48 h. Box plot showing quantification of telomere and γ-

H2AX overlap. Unpaired t-test was used. Scale bar is 10 μm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

DRIP followed by m⁶A RIP (DRImR)

Cell lysis → RNase A → DNA isolation → RNase III / Restriction enzymes → RNase H → DRIP S9.6 antibody → R-loop purified → DNase I → R-loop associated RNA → m⁶A RIP

Legend:
- RNA
- dsRNA
- DNA
- R-loop
- R-loop associated RNA
- m⁶A
- RNase A
- Restriction enzymes
- RNase III
- RNase H
- DNase I

B

DRImR

DRIP | m⁶A RIP

DRIP Input (2.5%)
DRIP Input (5%)
DRIP S9.6 (RNA)
DRIP IgG (RNA)
DRIP S9.6 (RNA) +RNase H

DNase I treatment

RNase A: + / −
TERRA probe

RL-TERRA

DRIP S9.6 + m⁶A | DRIP S9.6 + RNase H + m⁶A | DRIP S9.6 + IgG

TERRA probe — RNase A: − / +

C

Control-sh | METTL3-KD | Control-sh | METTL3-KD

DRIP Input (2.5%)
DRIP Input (5%)
DRIP S9.6 (RNA)
DRIP IgG (RNA)
DRIP S9.6 (RNA) +RNase H

DNase I treatment

RNase A: − / − / + / +
TERRA probe

Relative levels of RL-TERRA
****
Control-sh / METTL3-KD

D

Control shRNA | METTL3-KD
Control siRNA | Control siRNA | RNase H siRNA-1 | RNase H siRNA-2
TERRA RNA-FISH

#TERRA foci/nucleus
**** **** ****
- Control-sh + Control siRNA
- METTL3-KD + Control siRNA
- METTL3-KD + RNase H siRNA-1
- METTL3-KD + RNase H siRNA-2

E

RNase H^WT | RNase H^Mut
Merge | Merge
S9.6-TRF2 PLA | S9.6-TRF2 PLA
TRF1-mCherry | TRF1-mCherry

S9.6-TRF2 PLA
PLA foci co-localized with TRF1-mCherry/nucleus
****
RNase H^WT / RNase H^Mut

F

Control-sh | METTL3-KD
Merge | Merge
S9.6-TRF2 PLA | S9.6-TRF2 PLA
TRF1-mCherry | TRF1-mCherry

S9.6-TRF2 PLA
PLA foci co-localized with TRF1-mCherry/nucleus
****
Control-sh / METTL3-KD

G

Control siRNA | hnRNPA2B1 siRNA-2
Merge | Merge
S9.6-TRF2 PLA | S9.6-TRF2 PLA
TRF1-mCherry | TRF1-mCherry

S9.6-TRF2 PLA
PLA foci co-localized with TRF1-mCherry/nucleus
****
Control siRNA / hnRNPA2B1 siRNA-2

Figure 6

Figure 7