Single cell guided deconvolution of bulk transcriptomics recapitulates differentiation

stages of acute myeloid leukemia and predicts drug response

E Onur Karakaslar[1,2,3], Jeppe Severens[1,2,3], Elena Sánchez-López[3,4], Peter A van Veelen[4], Mihaela Zlei[5,6],

Jacques JM van Dongen[6], Annemarie M. Otte[4], Constantijn JM Halkes[7], Peter van Balen[7], Hendrik

Veelken[3,7], Marcel JT Reinders[1,2,3], Marieke Griffioen[7], Erik B van den Akker[1,2,3]

Affiliations

[1]Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

[2]Pattern Recognition & Bioinformatics, Delft University of Technology, Delft, The Netherlands

[3]Leiden Center for Computational Oncology, Leiden University Medical Center, Leiden, The Netherlands

[4]Center for Proteomics and Metabolomics, Leiden University Medical Center, Leiden, The Netherlands

[5]Department of Flow Cytometry, Medical Laboratory, Regional Institute of Oncology, Iasi, Romania

[6]Department of Immunology, Leiden University Medical Center, Leiden, The Netherlands

[7]Department of Hematology, Leiden University Medical Center, Leiden, The Netherlands

Materials & Correspondence

Erik B. van den Akker, PhD; Department of Biomedical Data Sciences, Leiden University Medical Center,

Leiden, The Netherlands; Einthovenweg 20, 2333 ZC, Leiden, The Netherlands; Tel: +31 (0)71 526 85 57;

Fax: +31 (0)71 526 82 80; E-mail: e.b.van_den_akker@lumc.nl

Abstract

The diagnostic spectrum for AML patients is increasingly based on genetic abnormalities due to their

prognostic and predictive value. However, information on the AML blast phenotype regarding their

maturational arrest has started to regain importance due to its predictive power on drug responses. Here, we

deconvolute 1350 bulk RNA-seq samples from five independent AML cohorts on a single-cell healthy BM

reference and demonstrate that the morphological differentiation stage (FAB classification) could be

faithfully reconstituted using estimated immune compositions (ICs). Moreover, we show that the estimated

ICs reliably predict *ex-vivo* drug resistances as demonstrated for Venetoclax, a *BCL-2* inhibitor, resistance

specifically in AML with CD14+ monocyte phenotype. We further validate these predictions using in-house

proteomics data by showing that *BCL-2* protein abundance is split into two distinct clusters for NPM1-

mutated AML at the extremes of CD14+ monocyte percentages, which could be crucial for the Venetoclax

dosing for these patients. Our results suggest that Venetoclax resistance predictions can also be extended to

AML without recurrent genetic abnormalities (NOS), and possibly to MDS-related AML and secondary

AML. Collectively, we propose a framework for allowing a joint mutation and maturation stage modeling

that could be used as a blueprint for testing sensitivity for new agents across the various subtypes of AML.

Introduction

Acute myeloid leukemia (AML) is an aggressive hematological cancer of the myeloid lineage. AML is caused by oncogenic driver abnormalities that are predominantly somatically acquired and cooperate in inducing a maturation arrest and an uncontrolled proliferation of immature myeloid precursor cells. AML patients continue to experience high relapse and mortality rates, leading to a 5-year survival rate of 27% as of 2022[1]. However, the diagnosis of AML encompasses heterogeneous subtypes with widely varying outcomes, prompting attempts to characterize AML and its subtypes to provide guidelines for prognostication of patients and treatment stratification. The first framework was proposed by the French-American-British (FAB) cooperative groups in 1976 based on cytomorphologic criteria (M0-M7), later referred to as FAB classification [2]. Subsequently, large-scale genetic studies in AML led to the discovery of recurrent genetic abnormalities (RGA) and stimulated a gradual shift from morphology-based to genetic-based prognosis in the last decades[3–6]. The latest World Health Organization (WHO) classification distinguishes eleven RGA-based AML subtypes and a heterogeneous umbrella subtype comprising a highly diverse set of relatively rare RGAs[5].

In the WHO classification, AML differentiation stage is now only assessed as a generic "fallback" option in cases without detectable RGA. Recently, FAB classification regained interest as differentiation stages (or immune composition) of AML might influence sensitivity or resistance to new drugs [7–9]. AML differentiation stage can be measured by diagnostic flow cytometry. Yet, flow cytometry is limited to a relatively low number of differentiation markers. Therefore, gene expression profiling by bulk RNA-sequencing (RNA-seq) with fully-automated pipelines might be an attractive alternative technique as it allows both calling of genetic aberrations and estimation of cell subsets, i.e. immune composition (IC) from the same sample[10–12] . Reported attempts to estimate ICs by deconvoluting bulk AML samples utilizing single-cell RNA-seq as an *in-silico* reference, were mostly focused on detection of survival differences without performing thorough validation of the ICs and/or used leukemic samples as reference[13–15], which prevents assessing whether ICs are tissue- or sample- specific.

Here, we perform deconvolution of 1350 AML transcriptomic samples via a healthy single-cell reference

while validating our findings with our in-house flow cytometry data. Of note, we demonstrate that the ICs

recapitulate the entire FAB landscape (M0-M7). Then, using these ICs we predict *ex-vivo* drug resistance

data from literature and show the agreement of these results at protein level with the help of in-house

proteomics data in AML patients, for whom we also had acquired gene expression data. To conclude, we

hereby propose a transcription-based single-cell guided deconvolution framework to assess the drug

effectiveness to different maturational stages of AML. We also provide our framework as a CRAN R package

available at https://github.com/eonurk/seAMLess.

Results

Deconvolution pipeline recapitulates healthy and malignant hematopoiesis

To demonstrate that a deconvolution framework could capture immune phenotypes of AML patients, we

initially created a healthy bone marrow (BM) reference atlas from single-cell transcriptomics data (**Figure**

1A). Using three publicly available datasets (two full-transcriptome and one targeted) from two studies[16,17],

we integrated data from 69,101 cells covering 439 genes (**Figure 1B**, Methods). As expected, T-cells and B-

cells formed separate clusters in the UMAP plot, whereas the myeloid lineage cells clustered together. To

differentiate early myelopoiesis, we distinguished more than 3,349 hematopoietic stem cells (HSCs) and

1,432 erythro-myeloid progenitors (EMPs), 2,939 lymphoid multipotent progenitors (LMPPs), as well as

3,508 granulocytes-monocytes progenitors (GMPs) (**Figure 1C**). Inter-individual variability, possibly due to

age differences, mostly affected T- and B- cells and did not influence overall clustering (**Figure S1A**).

Moreover, the homogeneous distribution of the studies on the UMAP plot showed successful integration of

the different datasets (**Figure S1B**).

We next performed *in silico* experiments to validate our deconvolution set-up. We first used the

healthy BM reference to create *in silico* mixed bulk samples with known immune compositions.

Pseudobulk profiles were simulated with one abundant cell type (80% of the cells) with the remaining cells

being a mix created by random selection (Methods). Then, MuSiC[12] with the default settings was used to

deconvolute the simulated pseudobulk profiles into their respective cell types (**Figure S1C**). To validate on

an independent dataset, we also simulated pseudobulks from 40,000 healthy bone marrow cells of the

human cell atlas (HCA)[18] (**Figure S1D**) and performed deconvolution via MuSiC. All simulated pseudobulk

profiles were successfully deconvoluted into their respective cell types (to the most abundant cell type)

apart from EMPs (**Figure 1D**). For this profile, the annotations were shared mostly among EMPs (14%) as

well as HSCs (25%) and Early Erythrocytes (23%). This discrepancy might be explained by the

transcriptional similarity of these cell types as observed in **Figure 1B**.

Next, we analyzed our in-house diagnostic flow cytometry data (EuroFlow[19] panels; see Methods)

for 22 AML samples with matched bulk RNA-seq data. An overview of mean fluorescence intensity (MFI)

values for these samples' abnormal cells after staining with antibodies for 33 markers distributed over 7

tubes (1 Orientation + 6 AML assignment tubes) is shown in **Figure S1E** (Supplementary Table S1). In line

with our expectations, the most abundant cell types for all samples were in the myeloid lineage for both

flow cytometry analyses and immune compositions (ICs) (**Figure 1E**, Supplementary Table S4). To

investigate whether monocytic AML can be accurately distinguished from AML with more stem cell-like

phenotypes, we plotted CD14+ monocyte percentages as determined by deconvolution against MFI values

of antibodies of monocytic markers (CD11b, CD64, IREM2, and CD14), and observed statistically significant

correlations for 3 markers (CD11b, CD64, IREM2), with CD64b being most significant ($R^2 = 0.43$, $P < 0.001$)

(**Figure S1F**). Also, percentages of AML cells assigned to the monocytic subset by EuroFlow panels and ICs

showed statistically significant correlations ($R^2 = 0.64$, $P < 0.001$) (**Figure 1F**).

Lastly, we downloaded publicly available TARGET AML and ALL (B-ALL and T-ALL) data and

deconvoluted these samples (n=719) with the healthy BM reference (**Figure 1G,** Supplementary Table S1) to

show that different leukemic phenotypes could be captured by deconvolution. The estimates of cell type

abundances in the three immune lineages (myeloid, B- and T-lymphoid) were visualized using a ternary

plot. As expected, each type of acute leukemia was closer to their respective cell type of origin (ICs), most

compact being B-ALL (yellow samples on the right corner), showing the ICs' ability to capture the patients'

immune phenotypes at major cell type levels. Together, these benchmarking and validation analyses

demonstrated that given a healthy single cell BM transcriptomic atlas, the cell type proportions can be

recapitulated faithfully via deconvolution from bulk transcriptomic leukemia cases.

Deconvolution of bulk AML transcriptomics reveals the dominating immune fraction

To investigate heterogeneity in immune composition of AML, we next applied our framework to

deconvolute five independent bulk transcriptomic AML studies, i.e. TCGA-LAML[20] (n=151), BEAT-AML[21]

(n=460), TARGET-AML[22] (n=187), LEUCEGENE[23] (n=452) and our cohort LUMC[10] (n=100) totaling in 1350

samples from 1,267 patients (**Figure S2A**). The results are shown in **Figure 2A** as a heatmap where each

sample was decomposed into the 22 cell types from the healthy BM reference. We also added information

on patients' clinical blast counts, FAB[2], WHO 2016[4], and ELN 2017[6] classes for an overarching picture of

AML landscape, and we also calculated the stemness score[24], which is a gene expression signature for

patient prognosis trained on engraftment capacity of AML in immunodeficient mice (Supplementary Table

S2). Each sample was assigned to the most abundant cell type as determined by deconvolution (AML

phenotype) and arranged according to their fractions within each phenotype. The data showed a clear

dominance of one immune cell type for most of the cases, clearly indicating that maturational arrests and

lineage skewing are leukemic properties that can be readily assessed using transcriptome sequencing. The

majority of AML cases was typically estimated to be dominated by myeloid cells (**Figure 2A, S2A, S2B,**

S3D). As notable exceptions, a few pediatric AML samples from the TARGET cohort showed IC profiles

dominated by T-cells or B-cells, possibly caused by more eminent infiltration of these subsets within

pediatric AML. These cases have previously been reported in TARGET, and specifically T-cell dominated

patients were annotated as poor survival group[25]. Furthermore, in line with previous reports, patients with

acute promyelocytic leukemia (APL), classified as AML-M3 by FAB, were correctly assigned to have a cell

composition dominated by GMPs. Besides APL, there are also AML cases dominated by GMP and large

groups of AML assigned as monocytic AML or AML with an earlier HSC or EMP phenotype, again

demonstrating that bulk transcriptomics can be used to capture the stage of arrest of AML during

hematopoiesis by deconvolution.

Deconvolution of bulk AML transcriptomics agrees upon stemness score

To comprehensively visualize changes in ICs in relation to metadata, samples were visualized using a

UMAP. As expected, samples with similar ICs clustered, as shown after annotating the samples for their

most abundant cell type (**Figure 2B** – left panel). When superimposing the deconvoluted percentages of 6

major cell types on the same UMAP plot (**Figure 2B** – right panels), gradual shifts in immune composition

became apparent, with APL patients populating in the extremes of the GMP cluster. Moreover, a gradient

towards monocytic outgrowth and conversely to a high stemness was also observed. Comparisons with the

clinical blast percentage and previously published stemness score showed an overall agreement with ICs,

HSCs and EMPs having a higher score than CD14+ Monocytes and GMPs (**Figure S3A, S3B**). Late

Erythrocyte also had high stemness score, albeit with large variation. By dividing AML into cases with high

and low stemness scores, we observed similar compositional changes with abundance of HSCs and EMPs in

AML with high stemness scores in all cohorts. Late erythrocytes, however, were not consistently enriched

in all cohorts. Furthermore, we noticed that in TARGET, T cells are abundant in AML with high stemness

scores (**Figure S3C**). This raises the question whether the stemness score, which has been trained on adult

AML, is useful to predict prognosis of pediatric patients particularly considering that large subtypes of adult

AML such as APL and AML with mutated NPM1 are rare in pediatric AML. The consistent enrichment of

HSCs and EMPs in AML with high stemness scores in all cohorts, however, further supports correct

determination of the immune composition of AML by deconvoluting bulk transcriptomics.

Immune composition recapitulates FAB classes

Clear associations were also observed between cell type assignments by IC and FAB classification status (left

panel of **Figure 2C**). Plotting continuous values of deconvolutions rather than categorical assignments

yielded an even more eminent picture (right panel of **Figure 2C**): M0 (minimally differentiated AML) cases

174 had high levels of HSCs, whereas these levels decreased and EMPs and GMPs appeared in M1 (AML

175 without maturation) and M2 (AML with maturation). As expected, M3 (APL) cases almost entirely

176 consisted of GMPs, while M4 (Acute Myelomonocytic Leukemia) and M5 (Acute monoblastic/monocytic

177 leukemia) samples contained CD14+ Monocyte cells and GMPs. The small groups of M6 (Acute

178 Erythroleukemia) and M7 (Acute Megakaryoblastic Leukemia) AML were mostly composed of Late

179 Erythrocytes and Megakaryocyte Progenitors, respectively. These data demonstrated and confirmed that

180 our single cell guided deconvolution strategy successfully captures the maturational arrest of AML cells at

181 different differentiation stages of hematopoiesis.

183 **Immune composition captures genetic subtype-specific resistances to various drugs**

184 To explore whether ICs conveys information on drug resistances, *ex vivo* drug response data of 122 small

185 molecule inhibitors provided as area under the curves (AUC) for 363 AML samples from BEAT were

186 downloaded. A higher AUC indicates that cancer cells are relatively resistant since higher drug

187 concentrations are needed to induce cell death. To predict drug resistance for AML samples with calculated

188 ICs, we trained random forest (RF) models per drug via leave-one-out cross validation (LOOCV) setting and

189 then calculated the Spearman ρ values of these models' predictions (**Figure S4A**; Supplementary Table S5-

190 S7). We also stratified these results according to WHO classifications (**Figure 3A, S4B**, Supplementary Table

191 S8).

193 Drug responses to Erlotinib (Spearman $\rho = 0.376$), a lung and pancreatic cancer drug, and mTOR pathway

194 inhibitor Rapamycin (Spearman $\rho = 0.368$) showed high correlations with AML with mutated NPM1 (n =

195 77, $R^2 = 0.29$, P= 3.7×10^{-7}) and inv(3) (n = 6, $R^2 = 0.91$, P= 3.2×10^{-3}), respectively. Notably, responses to

196 Flavopiridol (Spearman $\rho = 0.325$), a CDK kinase inhibitor, showed strong correlations with IC within the

197 group of AML-NOS (n = 77, $R^2 = 0.31$, P= 3.7×10^{-7}) while showing higher resistance towards AML with

198 more stem cell like cell phenotypes (**Figure 3A, S5A, S5B**). The strongest correlation however was observed

199 for Venetoclax (ABT-199), a BCL2 inhibitor, (Spearman $\rho = 0.509$) and the predictions for this drug with RF

are shown in **Figure S5C**. Annotation with primary diagnosis from WHO classes (**Figure 3B**) revealed that

AML cases from APL patients with PML-RARA (n = 8) that are dominated by GMPs are sensitive for

Venetoclax, whereas AML with CBFB-MYH11 (n = 7) had the best fit (R^2 = 0.78, P = 0.009) for Venetoclax

resistance, although the small sample size of CBFB-MYH11 cases prohibits drawing robust conclusions.

Larger groups of AML, however, such as NPM1 mutated cases (n = 55, R^2 = 0.33, P < 0.001), and AML-NOS

(n = 42, R^2 = 0.17, P = 0.007), together accounting for approximately 58% of AML, also showed a clear trend

for Venetoclax resistance (**Figure 3B**). Annotating drug responses with predicted resistances via patient's

most abundant cell types by deconvolution (**Figure S5C**) revealed that CD14+ Monocytic AML are

relatively resistant to Venetoclax. This resistance of monocytic AML was also clear by overlaying the AUC

for Venetoclax onto the UMAP plot (**Figure 3C**) or boxplots (**Figure S5D**) (one-way ANOVA p = 6.31×10^{-13};

Supplementary Table S9). Also, within the group of NPM1 mutated samples, which is the largest class of

AML, CD14+ Monocyte dominated cases were most resistant to Venetoclax (**Figure S5E, S5F**; one-way

ANOVA p = 7.6×10^{-4}; Supplementary Table S10). Briefly, these findings suggest that information on the

immune composition of AML might yield therapeutic implications even within one genetic subtype.

Estimated CD14+ Monocyte percentages predict Venetoclax resistance better than *BCL-2* mRNA expression

Since Venetoclax is targeting *BCL-2* protein, we next checked *BCL-2* mRNA expression levels in AML and

overlaid the CD14+ Monocyte percentages for these cases (**Figure 3D**). The data showed that low BCL2

expression indeed correlated with strong resistance to Venetoclax. However, a few samples were resistant

despite relatively high *BCL-2* expression (samples at upper right corner in **Figure 3D**). As the majority of

these cases had a CD14+ Monocyte phenotype, we univariately associated *BCL-2* expression and CD14+

Monocyte percentage to investigate which factor best explains the Venetoclax response (**Figure 3E**). We

also associated metadata such as reported blast percentages, ELN status and primary diagnosis to determine

their effect sizes and significance (Supplementary Table S11). This analysis demonstrated that both CD14+

Monocyte percentages (adjusted P = 1.7×10^{-18}) and *BCL-2* mRNA expression (adjusted P = 4.8×10^{-14}) were

associated significantly with the Venetoclax response. Next, we created a multivariate model to investigate

whether the significance of CD14+ Monocyte percentages diminishes along with the presence of *BCL-2*

expression in the same model (**Figure 3F**, Multivariate Venetoclax Tab in Supplementary Table 12). In this

model, CD14+ Monocyte percentages remained significantly associated with Venetoclax resistance (P =

1.92×10^{-5}), while *BCL-2* expression was below the significance threshold of 0.01 (P = 0.015). In conclusion,

these results indicate that cellular composition is a more robust marker than *BCL-2* mRNA expression to

predict Venetoclax resistance, specifically for AML from NOS and NPM1 mutated patients.

Estimated CD14+ Monocyte percentages associates with BCL-2 protein abundance

Next, we compared the effects of *BCL-2* gene expression and CD14+ Monocyte percentages with *BCL-2*

protein abundance within each sample. For this purpose, we used our in-house produced proteomics data

after batch correction (**Figure S6A, S6B**) for matched AML cases (n = 39; LUMC) to correlate the abundance

of Apoptosis regulator *BCL-2* protein vs gene expression (**Figure S6C**) and CD14+ Monocyte percentages

(**Figure S6D**). The data showed that *BCL-2* gene expression and CD14+ Monocyte percentages both

correlated with *BCL-2* protein abundance to a similar extent (R^2 = 0.45, P < 0.001). We also overlaid plots

for *BCL-2* expression and CD14+ Monocyte percentages with information on patients' genetic abnormalities

(**Figure 4A**). As expected, based on their low AUC for Venetoclax response (**Figure 3B**), two AML cases with

PML-RARA had high *BCL-2* protein expression (n=2, dark green). We also confirmed the strong variability

in AUC for Venetoclax response within the group of AML with mutated NPM1 (n=18, green) by showing

two distinct groups at the extremes of CD14+ Monocyte percentages which correlated with *BCL-2* protein

expression (**Figure 4B**). Within the group of AML patients with mutated NPM1, CD14+ Monocyte

percentage associated stronger with both *BCL-2* protein abundance (R^2 = 0.56, P < 0.001) than *BCL-2* gene

expression (R^2 = 0.52) (**Figure 4C, 4D**). In conclusion, the data demonstrate the relevance of our

deconvolution approach on bulk RNA-Seq to separate AML, especially those with mutated NPM1, with

high and low monocyte percentages to predict the patient's response to Venetoclax.

Discussion

In this work, we utilized single cell guided deconvolution to decompose bulk transcriptomics data from five

independent AML cohorts and show that the obtained Immune Compositions (ICs) faithfully reconstitute

the FAB landscape (M0-M7). Moreover, using same-sample flow cytometry, we were able to validate our

deconvolution framework. Hence, following our previous work using deep transcriptomics to call various

types of leukemia-defining genetic aberrations[10,26], our current findings further underpin the power of

transcriptomic-based approaches as a comprehensive and versatile platform for AML diagnosis. We next

illustrated the potential use of our deconvolution framework for precision medicine applications, by

correlating the estimated IC to the results of an *ex vivo* drug resistance screening of 122 small-molecule

inhibitors in the BEAT-AML study. For the *BCL-2* inhibitor Venetoclax we show that higher levels of the

estimated IC subset 'CD14+ Monocyte' correspond to a higher resistance, and intriguingly, that estimated

CD14+ Monocyte levels is a better explanatory variable of resistance to Venetoclax than *BCL-2* expression

alone. Nevertheless, using same-sample in-house proteomics data in 39 patients, we show that the estimated

CD14+ Monocyte levels accurately mark BCL protein expression, and that for *NPM1*-mutated patients the

presence or absence of a CD14 monocytic outgrowth corresponds with a distinct NPM1 protein abundance.

As Venetoclax was approved by the US Food and Drug Administration (FDA) in October 2020 for its use in

clinics for untreated AML patients [27], other researchers also experimented with it and also observed a

similar phenomenon that we had observed. For instance, Kuusanmaki *et al.* reported that monocytic

differentiation of AML reduced sensitivity to Venetoclax ex vivo[8]. Similarly, Pei *et al.* reported that the

different monocytic subclones in vivo created resistance to Venetoclax treatment[9], and also recently, White

et al. showed that *BCL-2* inhibitor resistance could be predicted via the genes associated with monocytes.

Recently, using the flow cytometry data from untreated MDS patients Ganan-Gomez *et al.* showed that

after relapse and becoming secondary AML (sAML) patients, those patients with less matured cell types

(EMP) before treatment had a faster complete remission (CR) and longer relapse free status compared to

more matured cell type (GMP) with Venetoclax treatment, supporting our hypothesis that ICs could also

predict Venetoclax resistance for MDS and sAML patients[28]. Collectively, these studies suggest that

Venetoclax has different resistances at different maturational stages, and especially higher resistance for

patients with CD14+ Monocyte dominated phenotype as we have shown in this manuscript, and we provide

an open-source framework, *seAMLess*, for replicating our results or applying it to other clinically relevant

datasets.

Our findings may potentially have important implications on drug use especially for genetically

uncharacterized patients (AML, NOS) currently accounting for ~40% of all AML as well as other well-

characterized patients such as NPM1 mutated samples. For instance, one interesting case report was from

Korea [29], an elderly AML patient was treated with Venetoclax along with Azacitidine, and even though the

patient had a favorable NPM1 status this treatment lead into tumor lysis syndrome (TLS) and patient ended

up in ER 12 h into the treatment. Although the manuscript does not indicate the FAB status, AML cells

expressed CD33, CD117 and myeloperoxidase, demonstrating a more stem cell than monocytic phenotype,

thus one possible hypothesis could be the patient having a more sensitive cell types (e.g., EMP, GMP) to

Venetoclax and thus causing immediate apoptosis of these cells and leading to TLS.

One limitation of our deconvolution strategy is that it cannot distinguish the cancerous cell types as it uses a

healthy bone marrow as a reference. The reason we wanted to emphasize this is, for instance in **Figure 4D**

there is a relapsed NPM1 patient with 41% of CD14+ Monocytes, as deconvolution percentage might not be

exact, this patient could potentially have a healthy monocytic population but rather the reason for his

relapse could be due to a different cell type, possibly 17% EMP as it is the second most abundant cell type

for this patient. However, we have two rationales behind not using a single cell reference with cancerous

cells. First, without mutation calling for all cells, one cannot be sure whether a cell is cancerous or not.

Strategies like predicting cancer cells based on their transcriptional similarity of cells with mutation calling,

proposed by Van Galen *et al.*[16], adds another level of ambiguity to already not perfect deconvolution

pipelines. Secondly and more importantly, heterogeneity of AML causes further sub-clustering within

individual AML cases (e.g. UMAP plots of Triana *et al.*, 2021), therefore creating a not well-characterized

cell type signature but rather patient specific clusters[17]. Additionally, using healthy subsets as reference

allows our framework to provide more interpretable and intuitive results for clinicians and doctors, as it

reports immune phenotypes and percentages on contrary to score-based prognostic values[15,24,26]. To

summarize, we believe our proposed pipeline could be a blueprint for assessing new drugs' resistances on

different cell types of AML and along with our framework, they may provide better insights for clinicians

and help paving the way into precision medicine in AML.

Methodology

Creating the healthy BM single-cell reference

We downloaded three different healthy BM datasets from two different studies, namely Van Galen et. al.[16]

(full-transcriptome, n=6,915), and Triana et. al.[17] (full-transcriptome, n = 13,165; 462 targeted mRNA,

n=49,057). Then, all cell labels were uplifted up via *Seurat* package[31] (v4.0.3) default query annotation

pipeline to match with the Triana's full-transcriptome cell labels as it had the most recent and detailed

labels. Next, *Seurat*'s integration pipeline with *CCA* was run and the cells that were labeled as

doublets/multiplets were removed from the down-stream analyses, and this yielded a healthy BM atlas of

69,130 cells in total, covering 439 genes. Also, we used the 40,000 cell subset of HCA provided by

SeuratObject package (v4.0.2) for the validation analyses (**Figure 1C, 1D**).

Different schemes of creating pseudobulks

To create a cancerous-like pseudobulk profile from healthy BM reference and HCA subset, we selected a

total of 1,000 cells for each profile, majority (80%) of them coming from a cell-type (over-abundant) and

the rest of the cells were distributed according to inverse proportion of the numbers of cells for the

remaining cell types. To achieve this, first integrated counts were exponentiated to make them non-log

scale and *slice_sample* function from *dplyr* package (v1.0.7) was used with a replacement option. Then,

these non-log scale cell counts were summed up to create the pseudobulk profiles. For individual-based

pseudobulk, *AggregateExpression* from *Seurat* package was used.

Flow Cytometry data

AML cases were stained with fluorescent antibodies and analyzed by flow cytometry for diagnosis,

prognosis, and disease monitoring of AML in the diagnostic laboratory of the department of Hematology in

the Leiden University Medical Center. The flow cytometric test has been developed and performed

according to EuroFlow standard operating procedures (www.euroflow.org)[19]. EuroFlow antibody

combinations have been tested against references databases of normal cells from healthy individuals and

allow multidimensional identification and distinction of aberrant cells from normal cell populations. The

flow cytometric test includes 8 tubes with different antibody combinations, i.e. one ALOT tube (acute

leukemia orientation tube) and 7 AML tubes (AML1-7). The ALOT tube contains antibodies against CD3,

CD45, MPO, CD79, CD19 and CD7. The AML tubes contain antibodies against CD16, CD13, CD34, CD117,

CD11b, CD10, HLA-DR and CD45 (AML-1), CD35, CD64, CD34, CD117, CD300e/IREM2, CD14, HLA-DR

and CD45 (AML-2), CD36, CD105, CD34, CD117, CD33, CD71, HLA-DR and CD45 (AML-3), NuTdT,

CD56, CD34, CD117, CD7, CD19, HLA-DR and CD45 (AML-4), CD15, NG2, CD34, CD117, CD22, CD38,

HLA-DR and CD45 (AML-5), CD42a and CD61, CD203c, CD34, CD117, CD123, CD4, HLA-DR and CD45

(AML-6). AML tube 7 contains antibodies against CD41, CD25, CD34, CD117, CD42b, CD9, HLA-DR and

CD45, but this tube has not been used to stain AML cases analyzed in this study.

Obtaining bulk transcriptomic data and their preprocessing

We have downloaded the non-normalized count matrices (htseq-counts) and the meta files of the four

discovery cohorts (TCGA-LAML, BEAT-AML, TARGET-AML and TARGET-ALL) from

https://portal.gdc.cancer.gov. For LEUCEGENE, count data was downloaded from their dedicated site

(https://data.leucegene.iric.ca/) along with their provided meta data. All meta/count data were pre-

processed using *R* (v4.1.0). For the meta data, genomic aberration labels were relabeled to the main AML

WHO 2016 classes, non-AML samples were removed from the down-stream analyses, ELN-classes were

relabeled according to ELN 2017 recommendations. For the count data, ERCC spike-ins and mitochondrial

genes were removed, and the count matrix was then sorted according genes standard deviation in order to

remove the duplicated genes that had less variation thus providing less information, and lastly the gene

ensembl ids were converted to gene symbols. Before converting ensembl ids into gene symbols, the

stemness score for each patient was calculated via count-per-million (cpm) normalized libraries, and these

libraries were normalized using *cpm* function from *edgeR* package[32] (v.3.34.1).

Our 100 AML samples (LUMC) deposited to EGA with accession number EGAS00001003096 and they are

accessible upon request. QC benchmark analyses for these samples were done in our previous paper[10].

Therefore, we ran default *HT-SEQ* pipeline (v0.11.2) with paired-end option aligning fastq files to *hg38* to

obtain the count matrix. All above mentioned preprocessing steps (filtering, gene name conversion) were

also conducted for these samples as well before deconvolution.

Deconvolution pipeline

To deconvolute the simulated pseudobulks and bulk RNA-seq AML patients, we used *MuSiC*[12] package

(v0.1.1) as it benchmarked highly and consistently across different cell types at various settings[11] and had

an easy-to-use open-source (GPL-3) implementation (https://github.com/xuranw/MuSiC). We used non-log

scaled count values as inputs and set the normalization option to false. Patients were assigned to the groups

(e.g., GMP, CD14+ Monocyte etc.) according to their most abundant deconvoluted immune compositions.

In the heatmap (**Figure 2A**), patients were re-ordered according to their immune compositions within each

assignment.

UMAP of immune compositions

First, to obtain reproducible results with umap plots, we set a seed to 2 as UMAP procedure involves

random initialization. Then, we ran *umap* function with default parameters from *umap* package (v0.2.7) and

used the first two reduced dimensions to create the plots. All related figures were plotted using *ggplot2*

package (v3.3.5).

Drug resistance predictions via Random Forest

BEAT-AML has drug resistance data for 122 small-molecule inhibitors, we downloaded these from their

manuscript (Supplementary Table S5). Then, each drug response was min-max normalized, then matched

with their available RNA-seq samples. Next, drug resistances were predicted with the deconvoluted

immune compositions (ICs). Random forest algorithm from *randomForest* package (v.4.6-14) with default

parameters was used for the predictions. Each sample within each drug was predicted at leave-one-out

cross-validation settings. Then, for each drug, Spearman ρ values were calculated between predicted and

actual drug resistance values. To stratify the drugs according to their primary diagnosis of WHO

classification, samples within each diagnosis are selected and then each drugs predicted, and normalized

drug resistance were associated, then the correlation and p-values were calculated using *summary* function

in base R.

Venetoclax Association Analysis

First, each attribute was associated to standardized (min-max normalized) Venetoclax resistance from

BEAT-AML study at univariate settings using *lm* function in R environment (v4.1.0) (Supplementary Table

S11). Then, p-values and coefficients were calculated using *summary* function and then p-values were

multiple hypothesis corrected using Benjamini-Hochberg procedure. For multivariate models, FAB

classification, and immune composition levels (except CD14+ Monocytes) were excluded as only 76 out of

460 samples of BEAT-AML had FAB classifications and as other immune compositions are not independent

of CD14+ Monocyte percentages (as the question is whether CD14+ Monocyte levels are independently

predictive of Venetoclax resistance given *BCL-2* expression in the same model). Again, p-values and

coefficients were calculated with *summary* function and plotted in a volcano plot (Figure 3F,

Supplementary Table S12).

Proteomics Sample preparation

Cell lysis, digestion and TMT labeling was performed as described in Paula and et al[33]. Cell lysis was

performed using 5% SDS lysis buffer (100 mM Tris-HCl pH7.6) and 5 U benzonase nuclease (Thermo

Scientific) with incubation at 95 °C for 4 minutes. Protein concentration was determined using Pierce BCA

Gold protein assay (Thermo Fisher Scientific). 100 μg protein of each sample was then reduced with 5 mM

TCEP. Reduced disulfide bonds were alkylated using 15 mM iodoacetamide. Excess iodoacetamide was

quenched using 10 mM DTT. Protein lysates were precipitated using chloroform/methanol; resulting pellets

were re-solubilized in 40 mM HEPES pH 8.4 and digested using TPCK treated trypsin (1:12.5

enzyme/protein ratio) overnight at 37 °C. Peptide concentration was then determined using Pierce BCA

Gold protein assay.

The different samples, and reference samples, were arranged into five TMTpro 16plex sets. The peptides

were labeled with TMTpro Label Reagents (Thermo Fisher Scientific) in a 1:4 ratio by mass (peptides/TMT

reagents), total volume was 35 μL, for 1 h at RT. Excess TMT reagent was quenched with 5 μL 6%

hydroxylamine for 15 min at RT. Samples corresponding to a TMT set were pooled and lyophilized.

Each TMT sample (80 ug) was fractionated by high pH reverse phase chromatrography on a Zorbax RRHD

Eclipse Plus C18 2.1x150 mm 1.8-micron column, at 800 ul/min using an Agilent1200 binary HPLC system,

equipped with a UV detector. The mobile phases were 10 mM Ambic pH 8.4 (A) and 10 mM

Ambic/Acetonitrile 20/80 pH 8.4 (B). The gradient was from 2% to 90%B in 35 minutes. 20 fractions were

collected in a circular fashion, i.e., collection per vial for 20 sec before moving to the next collection vial.

After collection in the last vial collection is continued in the first vial. Fractions were subsequently freeze

dried.

Mass spectrometry

TMT-labeled peptide fractions were dissolved in water/formic acid (100/0.1 v/v) and analyzed by on line

C18 nanoHPLC MS/MS with a system consisting of an Ultimate3000nano gradient HPLC system (Thermo,

Bremen, Germany), and an Exploris480 mass spectrometer (Thermo) as in Rossi et al[34]. Fractions were

injected onto a cartridge precolumn (300 μm × 5 mm, C18 PepMap, 5 um, 100 A, and eluted via a

homemade analytical nano-HPLC column (50 cm × 75 μm; Reprosil-Pur C18-AQ 1.9 um, 120 A (Dr.

Maisch, Ammerbuch, Germany)). Solvent A was water/formic acid 100/0.1 (v/v). The gradient was run

from 2% to 40% solvent B (20/80/0.1 water/acetonitrile/formic acid (FA) v/v) in 120 min. The nano-HPLC

column was drawn to a tip of ~10 μm and acted as the electrospray needle of the MS source. The mass

spectrometer was operated in data-dependent MS/MS mode with a cycle time of 3 seconds, with the HCD

collision energy at 36 V and recording of the MS2 spectrum in the orbitrap, with a quadrupole isolation

width of 1.2 Da. In the master scan (MS1) the resolution was 120,000, the scan range 350-1200, at standard

AGC target @maximum fill time of 50 ms. A lock mass correction on the background ion m/z=445.12003

was used. Precursors were dynamically excluded after n=1 with an exclusion duration of 45 s, and with a

precursor range of 20 ppm. Charge states 2-5 were included. For MS2 the first mass was set to 110 Da, and

the MS2 scan resolution was 45,000 at an AGC target of 200% @maximum fill time of 60 ms.

Proteomics data processing and down-stream analysis

In a post-analysis process, raw data were first converted to peak lists using Proteome Discoverer version 2.4

(Thermo Electron), and submitted to the Uniprot database (Homo sapiens, 20596 entries), using Mascot v.

2.2.07 (www.matrixscience.com) for protein identification. Mascot searches were performed with 10 ppm

and 0.02 Da deviation for precursor and fragment mass, respectively, and the enzyme trypsin was specified.

Up to two missed cleavages were allowed. Methionine oxidation and acetyl on protein N-terminus were set

as variable modifications. Carbamidomethyl on Cys and TMTpro on Lys and N-terminus were set as fixed

modifications. Protein and peptide FDR were set to 1%. Normalization was on the total peptide amount.

The 5 TMT-16plex analyses were normalized to each other by the bridge samples.

First, the abundance data is log-cpm transformed to stabilize variance among samples and then to ensure

dealing with the technical batch effects, we ran *removeBatchEffect* function from *limma* package (v3.48.3)

providing batch information. Then, we overlaid the transformed protein abundances onto a PCA plot to

observe the clearance of batch related clustering. Then, the cell lines (positive-control) and controls

(negative-control) were removed from the down-stream analyses, selecting only primary AML (n=39).

Next, the transformed BCL-2 protein abundance was associated with the BCL-2 expression from LUMC

cohort (**Figure 4A**) and CD14+ Monocyte percentage within each sample (**Figure 4B**). R^2 and p-values were

calculated using *stat_poly_eq* function from *ggpmisc* (v0.4.5) package and lines were drawn with

geom_smooth function from ggplot2 package (v3.3.5) via method option set to linear model.

Data analysis and reproducibility

All analyses performed in this study are provided with the following link: https://github.com/eonurk/lumc-

sc-aml. We have also implemented an GPL-3 licensed R package at https://github.com/eonurk/seAMLess,

which deconvolutes a given bulk RNA-seq count matrix to 22 cell types from our single-cell reference and

predicts drug resistances via RF algorithm.

Acknowledgements

Authors want to thank Daniele Bizzarri for his fruitful critics on the project. This study was funded by a

strategic investment of the Leiden University Medical Center, embedded within the Leiden Oncology

Center, and executed within the Leiden Center for Computational Oncology. EvdA was funded by a

personal grant of the Dutch Research Council (NWO; VENI: 09150161810095). The funding bodies had no

role in the study design, the collection, analysis, and interpretation of data, the writing of the manuscript,

and the decision to submit the manuscript for publication.

Authors Contribution

Conceptualization: MJR, EvdA, MG

Resources: EvdA

Methodology: EOK, EvdA, PvA, AMO

Investigation: EOK, JS, MG, EvdA, PvA

Visualization: EOK

Funding acquisition: EvdA

Project administration: EvdA

Supervision: MJR, MG, EvdA

Writing – original draft: EOK, MG, EvdA

Writing – review & editing: All authors

Figure Captions

Figure 1

A The overview of the study. Integrated cells for healthy bone marrow (BM) dataset were collected from

two studies[16,17]. Five independent bulk transcriptomic AML cohort (TCGA-LAML, BEAT-AML, TARGET-

AML, LEUCEGENE and LUMC) samples (n=1,350) were deconvoluted. Drug resistance data (n=122) from

BEAT-AML were predicted using immune compositions (ICs). Additional ALL samples (n=532) from

TARGET study were used for validating deconvolution framework. Proteomics (n=39) and flow cytometry

samples (n=22) from LUMC cohort were used for further validation of the framework. **B** UMAP plot of

integrated healthy BM (n=69,101) cells. Annotations are lifted via Azimuth framework. T and B cell subset

were merged, resulting into 22 cell types (abbreviations; GMP: Granulocyte Monocyte Progenitors, LMPP:

Lymphoid Primed Multipotent Progenitors, NK: Natural Killer, EMP: Erythroid Megakaryocyte Progenitor,

pDC: Plasmacytoid Dendritic Cell, CLP: Common Lymphoid Progenitor, HSC: Hematopoietic Stem Cell,

BaEoMA: Basophil Eosinophil Mast Progenitor, Prog Mk: Progenitor Megakaryocyte, pre-mDC: Precursor

Myeloid Dendritic Cell, ASDC: AXL+ Dendritic Cell). **C** Barplot for cells per cell type of the healthy BM. X-

axis is in log10 scale. **D** Heatmap showing the fraction of deconvoluted cell types for simulated pseudobulks

from HCA subset. Each row indicates a simulated pseudobulk with an overabundant cell type (80%) and

adds up to 1, and each column is a cell type from the healthy BM reference. **E** Cellular composition of the

same samples via flow gating of orientation tube (ALOT) and immune compositions (RNA-seq). Abnormal

assignment within myeloid cells is determined by flow gating, thus only exists for flow cytometry. **F**

Percentage of monocytic subsets measured by flow cytometry and ICs. **G** Ternary plot showing the ICs of

TARGET AML and ALL cohorts (n=719), each dot represents a sample and corners indicate a major cell

type and colors indicate the primary diagnosis of each leukemic sample (Supplementary Table S1).

Figure S1

A Healthy BM reference overlaid on UMAP separated by samples **B** and by different studies. **C** Similar to

Figure 1D; heatmap showing the deconvolution results of simulated pseudobulks from healthy BM

reference via an over-abundant cell type (80%). **D** UMAP for HCA subset with the cell type annotations,

which were lifted via Azimuth framework. **E** Dot plot of MFI values of all EuroFlow markers for all

samples, size of the dot represents the abnormal cell percentages per sample assigned with EuroFlow

analyses. **F** MFI values of four EuroFlow monocytic markers (CD11b, CD64, IREM2 and CD14) and CD14+

Monocyte %. Colors indicate the abnormal percentage of cells by flow cytometry assignments.

Figure 2

A Heatmap showing the deconvoluted fractions for of five AML cohorts (TCGA-LAML, BEAT-AML,

TARGET-AML, LEUCEGENE and LUMC). Each column represents a sample, which is deconvoluted into

22 cell types (bottom part with blue) via healthy single cell BM reference. Top part of the annotations

shows the provided meta data (FAB, WHO and ELN classifications, origin, sex and reported blast

percentages) for these cohorts. Each patient was assigned to its most abundant deconvoluted cell type (AML

phenotype) and samples within each assignment were sorted according to the assigned phenotype. **B** UMAP

plot of deconvolution levels annotated via most abundant cell type as the left panel; and on the right,

immune compositions of 6 out of 22 cell types were shown in continuous scale (HSC, CD14+ Monocytes,

Late Erythrocytes, GMP, EMP and cDC). **C** Similar to UMAP plot in (B) but colored with FAB

classifications (left panel), and the fractions of cell types for each of these FAB classifications (right panel).

Figure S2

A Detailed table for sample characteristics for age, sex, FAB, ELN, and primary diagnosis (WHO) for each

cohort. Missing data is labeled as 'Unknown' for each attribute. *AML, NOS patients represents %39 of

WHO classes. **B** Ternary plots like Figure 1G for all five AML cohorts.

Figure S3

A Box plots of blast percentage vs AML phenotype and **B** Stemness score vs AML phenotype. Boxplot were

sorted according to median values. **C** Compositional changes of each cohort (except LEUCEGENE as it lacks

entire list of genes to calculate stemness score) for low and high stemness score (split by median). Violin

plots show the deconvoluted cellular percentages for 22 cell type and heatmap on the right panel

summarizes the changes. A linear model is fit to calculate the slope and significance per cell type vs.

stemness, and a red slope indicates more of that cell type with higher stemness score, and blue is the vice

versa. Asterix indicates a statistically significant trend ($P < 0.05$). **D** UMAP plots of deconvoluted

percentages (ICs) annotated via age, study, reported blast level, sex, ELN and stemness score.

Figure 3

A Correlation of predicted AUC and normalized AUC of *ex-vivo* drug resistances split by WHO classes (top

10 drugs). Bigger bubble corresponds to higher correlation, and colors indicate statistical significance. Only

groups with >5 samples were fit linear models thus R^2 and p values are shown. **B** Random Forest predictions

of Venetoclax resistance via immune compositions with LOOCV ($\rho = 0.509$; Supplementary Table - BEAT

AUC RF Predictions). Each dot represents a patient. Higher resistance (AUC) means more cells are viable

after administering the drug. **C** Samples with normalized Venetoclax resistance data are shown on the

UMAP plot of ICs, along with phenotype map from (**Figure 2B**). Patients without available data were

shown as gray dots for the sake of completeness. **D** Venetoclax resistance vs *BCL-2* expression; also, CD14+

Monocyte percentages for these patients were annotated. Expression data was count-per-million (cpm)

normalized. **E** Univariate analyses of different attributes (i.e., CD14+ Monocyte %, reported blast

percentages etc.) vs normalized Venetoclax resistance. P values were corrected using Benjamini-Hochberg

procedure and the names of the attributes are shown for FDR > 0.01. **F** Similar to **E**; A multivariate model is

associated with different attributes, and the names of the attributes with P> 0.01 is shown. Full list of

attributes' p values and estimates are provided in Supplementary Table for both **E** and **F**.

Figure S4

A Spearman ρ values for the predictions of the drug resistance of 122 small molecule inhibitors from BEAT-

AML (see Supplementary Table S7). For each drug, the resistance per sample is predicted via random forest

algorithm using immune compositions at leave-one-out cross validation setting. **B** Dot plot showing the

stratified associations with predicted resistance values. Colors of the dot show the significance and sizes

indicate the correlation strength. Groups with more than 5 samples were included in the plot.

Figure S5

A Flavopiridol resistance predictions with random forest annotated with patients' AML phenotype. **B**

Flavopiridol resistance predictions are faceted and highlighted by WHO classes, classes with >5 samples

were fit linear models thus R^2 and p values are shown. **C** Venetoclax resistance predictions with random

forest annotated with patients' AML phenotype. **D** Venetoclax resistance split by AML phenotypes. All

drugs' one-way anova p values are provided in Supplementary Table S9 **E** UMAP plot showing Venetoclax

resistance only for NPM1 mutated samples **F** Venetoclax resistance split by AML phenotype only for NPM1

mutated samples.

Figure 4

A *BCL-2* expression vs protein abundance (n = 39) annotated with genetic abnormalities (R^2 = 0.45, P <

0.001). Each dot represents a sample. **B** CD14+ Monocytes percentage vs *BCL-2* protein abundance

annotated with genetic abnormalities (R^2 = 0.45, P < 0.001). **C** Similar to A; but only limited to NPM1

mutated samples (n = 18) and annotated with patients' AML phenotypes (most abundant cell type) (R^2 =

0.52, P < 0.001) **D** Similar to B; but only limited to NPM1 mutated samples (n = 18) and annotated with

patients' AML phenotypes (most abundant cell type) (R^2 = 0.56, P < 0.001).

Figure S6

A All-proteomics data (n = 39) without batch correction and **B** batch corrected (with *limma*) log-cpm

normalized proteomics data. All abundances are log-cpm normalized. **C** *BCL-2* expression vs protein

abundance annotated with patients' AML phenotypes (R^2 = 0.45, P < 0.001). **D** CD14+ Monocytes

percentage vs *BCL-2* protein abundance annotated with patients' AML phenotypes (R^2 = 0.45, P < 0.001).

Supplementary Tables

Tab 1 TARGET deconvolutions

Deconvolutions of TARGET AML and ALL cohorts

Tab 2 Deconvolution Results

Immune compositions for 22 cell types along with meta data of 1,350 samples

Tab 3 EuroFlow MFI values

EuroFlow MFI values of abnormal populations of LUMC samples

Tab 4 EuroFlow cell type percentages

EuroFlow gating results for different cell types (abnormals are not annotated) of LUMC samples

Tab 5 BEAT AUC

Min-max normalized AUC values for 122 small-molecule inhibitors from BEAT-AML

Tab 6 BEAT AUC RF Predictions

Predictions of each drug resistance with random forest at LOOCV

Tab 7 Drug Spearman Rho

Spearman rho values for drug resistance predictions of 122 small-molecule inhibitors

Tab 8 Drug Associations stratified with WHO

Predicted drug and normalized resistance associations within each primary diagnosis

Tab 9 Anova - Drug vs Cell Types (All)

One-way anova test results with sample phenotype assignments for drug resistances

Tab 10 Anova - Drug vs Cell Types (NPM1)

One-way anova test results with sample phenotype assignments for drug resistances (NPM1 only)

Tab 11 Univariate Venetoclax

The results of linear model associations of Venetoclax resistance to different attributes

Tab 12 Multivariate Venetoclax

The results of linear model association of Venetoclax resistance with different attributes

References

1. *Cancer Facts & Figures 2022.* https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-

and-statistics/annual-cancer-facts-and-figures/2022/2022-cancer-facts-and-figures.pdf.

2. Bennett, J. M. *et al.* Proposals for the Classification of the Acute Leukaemias French-American-British

(FAB) Co-operative Group. *Br. J. Haematol.* **33**, 451–458 (1976).

3. Vardiman, J. W. *et al.* The 2008 revision of the World Health Organization (WHO) classification of

myeloid neoplasms and acute leukemia: rationale and important changes. *Blood* **114**, 937–951 (2009).

4. Arber, D. A. *et al.* The 2016 revision to the World Health Organization classification of myeloid

neoplasms and acute leukemia. *Blood* **127**, 2391–2405 (2016).

5. Khoury, J. D. *et al.* The 5th edition of the World Health Organization Classification of

Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia* **36**, 1703–1719

(2022).

6. Döhner, H. *et al.* Diagnosis and management of AML in adults: 2017 ELN recommendations from an

international expert panel. *Blood* **129**, 424–447 (2017).

7. Canaani, J. *et al.* Impact of FAB classification on predicting outcome in acute myeloid leukemia, not

otherwise specified, patients undergoing allogeneic stem cell transplantation in CR1: An analysis of 1690

patients from the acute leukemia working party of EBMT: CANAANI ET AL. *Am. J. Hematol.* **92**, 344–

350 (2017).

8. Kuusanmäki, H. *et al.* Phenotype-based drug screening reveals association between venetoclax response

and differentiation stage in acute myeloid leukemia. *Haematologica* **105**, 708–720 (2020).

9. Pei, S. *et al.* Monocytic Subclones Confer Resistance to Venetoclax-Based Therapy in Patients with

Acute Myeloid Leukemia. *Cancer Discov.* **10**, 536–551 (2020).

10. Arindrarto, W. *et al.* Comprehensive diagnostics of acute myeloid leukemia by whole transcriptome

RNA sequencing. *Leukemia* **35**, 47–61 (2021).

11. Avila Cobos, F., Alquicira-Hernandez, J., Powell, J. E., Mestdagh, P. & De Preter, K. Benchmarking of

cell type deconvolution pipelines for transcriptomics data. *Nat. Commun.* **11**, 5650 (2020).

12. Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution with multi-

subject single-cell expression reference. *Nat. Commun.* **10**, 380 (2019).

13. Dai, C., Chen, M., Wang, C. & Hao, X. Deconvolution of Bulk Gene Expression Profiles with Single-Cell

Transcriptomics to Develop a Cell Type Composition-Based Prognostic Model for Acute Myeloid

Leukemia. *Front. Cell Dev. Biol.* **9**, 762260 (2021).

14. Li, H., Sharma, A., Ming, W., Sun, X. & Liu, H. A deconvolution method and its application in analyzing

the cellular fractions in acute myeloid leukemia samples. *BMC Genomics* **21**, 652 (2020).

15. Zeng, A. G. X. *et al.* A cellular hierarchy framework for understanding heterogeneity and predicting

drug response in acute myeloid leukemia. *Nat. Med.* **28**, 1212–1223 (2022).

16. van Galen, P. *et al.* Single-Cell RNA-Seq Reveals AML Hierarchies Relevant to Disease Progression and

Immunity. *Cell* **176**, 1265-1281.e24 (2019).

17. Triana, S. *et al.* Single-cell proteo-genomic reference maps of the hematopoietic system enable the

purification and massive profiling of precisely defined cell states. *Nat. Immunol.* **22**, 1577–1589 (2021).

18. Hay, S. B., Ferchen, K., Chetal, K., Grimes, H. L. & Salomonis, N. The Human Cell Atlas bone marrow

single-cell interactive web portal. *Exp. Hematol.* **68**, 51–61 (2018).

19. on behalf of the EuroFlow Consortium (EU-FP6, LSHB-CT-2006-018708) *et al.* EuroFlow antibody

panels for standardized n-dimensional flow cytometric immunophenotyping of normal, reactive and

malignant leukocytes. *Leukemia* **26**, 1908–1975 (2012).

20. The Cancer Genome Atlas Research Network. Genomic and Epigenomic Landscapes of Adult De Novo
Acute Myeloid Leukemia. *N. Engl. J. Med.* **368**, 2059–2074 (2013).

21. Tyner, J. W. *et al.* Functional genomic landscape of acute myeloid leukaemia. *Nature* **562**, 526–531
(2018).

22. Farrar, J. E. *et al.* Genomic Profiling of Pediatric Acute Myeloid Leukemia Reveals a Changing
Mutational Landscape from Disease Diagnosis to Relapse. *Cancer Res.* **76**, 2197–2205 (2016).

23. MacRae, T. *et al.* RNA-Seq Reveals Spliceosome and Proteasome Genes as Most Consistent Transcripts in
Human Cancer Cells. *PLoS ONE* **8**, e72884 (2013).

24. Ng, S. W. K. *et al.* A 17-gene stemness score for rapid determination of risk in acute leukaemia. *Nature*
540, 433–437 (2016).

25. Jiang, F. *et al.* An Immune Checkpoint-Related Gene Signature for Predicting Survival of Pediatric
Acute Myeloid Leukemia. *J. Oncol.* **2021**, 1–14 (2021).

26. Docking, T. R. *et al.* A clinical transcriptome approach to patient stratification and therapy selection in
acute myeloid leukemia. *Nat. Commun.* **12**, 2474 (2021).

27. *FDA grants regular approval to venetoclax in combination for untreated acute myeloid leukemia.*
https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-regular-approval-
venetoclax-combination-untreated-acute-myeloid-leukemia (2020).

28. Ganan-Gomez, I. *et al.* Stem cell architecture drives myelodysplastic syndrome progression and predicts
response to venetoclax-based therapy. *Nat. Med.* **28**, 557–567 (2022).

29. Kim, M. *et al.* Venetoclax with Azacitidine Induced Tumor Lysis Syndrome in an Elderly Patient with
Acute Myeloid Leukemia: A Case Report. *Electrolytes Blood Press.* **19**, 46 (2021).

30. Ngai, L. L., Kelder, A., Janssen, J. J. W. M., Ossenkoppele, G. J. & Cloos, J. MRD Tailored Therapy in
AML: What We Have Learned So Far. *Front. Oncol.* **10**, 603636 (2021).

31. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

32. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinforma. Oxf. Engl.* **26**, 139–140 (2010).

33. Paulo, J. A. & Gygi, S. P. Nicotine-induced protein expression profiling reveals mutually altered proteins

across four human cell lines. *Proteomics* **17**, (2017).

34. Rossi, M. *et al.* PHGDH heterogeneity potentiates cancer cell dissemination and metastasis. *Nature* **605**,

747–753 (2022).

A Overview of the pipeline

B Integrated single-cell healthy BM reference

C Number of cells in each cell type (Healthy BM reference)

D Deconvolution of in-silico pseudobulks from Human Cell Atlas (HCA)

E Comparison of flow cytometry and immune composition

F Monocytic levels with flow cytometry and immune composition

G Deconvolution of TARGET-AML and -ALL

A Heatmap of deconvolution results for all patients

A Single cell atlas annotated with samples

B Single cell atlas annotated with study

C Deconvolution of healthy BM reference

D HCA annotated with new cell types

E MFI values for EuroFlow Markers

F Monocytic Markers' MFI vs CD14+ Monocyte percentages

A Patient meta summary from all studies

Characteristic	Overall, N = 1,267[1]	BEAT, N = 412[1]	LEUCE, N = 452[1]	LUMC, N = 96[1]	TARGET, N = 156[1]	TCGA, N = 151[1]
Age	55 (0, 89)	62 (2, 87)	57 (17, 87)	NA (Inf, -Inf)	9 (0, 23)	56 (22, 89)
Unknown	97	1	0	96	0	0
Sex						
female	567/1,267 (45%)	183/412 (44%)	195/452 (43%)	44/96 (46%)	77/156 (49%)	68/151 (45%)
male	700/1,267 (55%)	229/412 (56%)	257/452 (57%)	52/96 (54%)	79/156 (51%)	83/151 (55%)
FAB						
M0	51/761 (6.7%)	5/90 (5.6%)	27/381 (7.1%)	0/0 (NA%)	4/140 (2.9%)	15/150 (10%)
M1	189/761 (25%)	7/90 (7.8%)	129/381 (34%)	0/0 (NA%)	17/140 (12%)	36/150 (24%)
M2	138/761 (18%)	10/90 (11%)	54/381 (14%)	0/0 (NA%)	37/140 (26%)	37/150 (25%)
M3	56/761 (7.4%)	11/90 (12%)	30/381 (7.9%)	0/0 (NA%)	0/140 (0%)	15/150 (10%)
M4	155/761 (20%)	24/90 (27%)	59/381 (15%)	0/0 (NA%)	43/140 (31%)	29/150 (19%)
M5	145/761 (19%)	31/90 (34%)	69/381 (18%)	0/0 (NA%)	30/140 (21%)	15/150 (10%)
M6	14/761 (1.8%)	0/90 (0%)	10/381 (2.6%)	0/0 (NA%)	2/140 (1.4%)	2/150 (1.3%)
M7	13/761 (1.7%)	2/90 (2.2%)	3/381 (0.8%)	0/0 (NA%)	7/140 (5.0%)	1/150 (0.7%)
Unknown	506	322	71	96	16	1
ELN						
Adverse	197/686 (29%)	148/390 (38%)	0/0 (NA%)	0/0 (NA%)	9/148 (6.1%)	40/148 (27%)
Favorable	203/686 (30%)	107/390 (27%)	0/0 (NA%)	0/0 (NA%)	64/148 (43%)	32/148 (22%)
Intermediate	286/686 (42%)	135/390 (35%)	0/0 (NA%)	0/0 (NA%)	75/148 (51%)	76/148 (51%)
Unknown	581	22	452	96	8	3
Blast	72 (0, 100)	60 (0, 98)	77 (1, 99)	75 (15, 99)	74 (14, 100)	71 (0, 100)
Unknown	144	109	31	0	4	0
Primary Diagnosis						
Acute myeloid leukemia with inv(3)(q21q26.2) or t(3;3)(q21;q26.2); RPN1-EVI1	17/1,267 (1.3%)	8/412 (1.9%)	7/452 (1.5%)	2/96 (2.1%)	0/156 (0%)	0/151 (0%)
Acute myeloid leukemia with mutated CEBPA	37/1,267 (2.9%)	28/412 (6.8%)	0/452 (0%)	0/96 (0%)	9/156 (5.8%)	0/151 (0%)
Acute myeloid leukemia with mutated NPM1	163/1,267 (13%)	96/412 (23%)	0/452 (0%)	27/96 (28%)	7/156 (4.5%)	33/151 (22%)
Acute myeloid leukemia with myelodysplasia-related changes	213/1,267 (17%)	88/412 (21%)	114/452 (25%)	11/96 (11%)	0/156 (0%)	0/151 (0%)
Acute myeloid leukemia with t(10;11)(p12;q23.3); MLLT10-KMT2A	9/1,267 (0.7%)	0/412 (0%)	0/452 (0%)	0/96 (0%)	9/156 (5.8%)	0/151 (0%)
Acute myeloid leukemia with t(6;11)(q27;q23.3); MLLT4-KMT2A	5/1,267 (0.4%)	0/412 (0%)	0/452 (0%)	0/96 (0%)	5/156 (3.2%)	0/151 (0%)
Acute myeloid leukemia with t(6;9)(p23;q34); DEK-NUP214	8/1,267 (0.6%)	3/412 (0.7%)	3/452 (0.7%)	1/96 (1.0%)	1/156 (0.6%)	0/151 (0%)
Acute myeloid leukemia with t(8;21)(q22;q22); RUNX1-RUNX1T1	60/1,267 (4.7%)	9/412 (2.2%)	20/452 (4.4%)	3/96 (3.1%)	21/156 (13%)	7/151 (4.6%)
Acute myeloid leukemia with t(9;11)(p22;q23); MLLT3-MLL	35/1,267 (2.8%)	9/412 (2.2%)	8/452 (1.8%)	4/96 (4.2%)	12/156 (7.7%)	2/151 (1.3%)
Acute myeloid leukemia, CBF-beta/MYH11	101/1,267 (8.0%)	26/412 (6.3%)	29/452 (6.4%)	10/96 (10%)	28/156 (18%)	8/151 (5.3%)
Acute myeloid leukemia, NOS	492/1,267 (39%)	91/412 (22%)	215/452 (48%)	35/96 (36%)	64/156 (41%)	87/151 (58%)
Acute promyelocytic leukaemia, PML-RAR-alpha	62/1,267 (4.9%)	17/412 (4.1%)	29/452 (6.4%)	2/96 (2.1%)	0/156 (0%)	14/151 (9.3%)
Myeloid sarcoma	3/1,267 (0.2%)	3/412 (0.7%)	0/452 (0%)	0/96 (0%)	0/156 (0%)	0/151 (0%)
Therapy related myeloid neoplasm	62/1,267 (4.9%)	34/412 (8.3%)	27/452 (6.0%)	1/96 (1.0%)	0/156 (0%)	0/151 (0%)

[1] Median (Range) or Frequency (%)

B AML studies on ternary plot

A Drug resistance predictions for all drugs

Leave one out cross validation within each drug

B Drug resistance predictions stratified for primary diagnosis

A
Predicted Flavopiridol resistance by AML phenotype

B
Flavopiridol predicted resistance stratified with primary diagnosis

C
Predicted Venetoclax resistance by AML phenotype

D
Venetoclax resistance for each AML phenotype (raw AUC values)

E
Venetoclax resistance on AML with mutated NPM1

F
Venetoclax resistance for each AML phenotype on AML with mutated NPM1

Figure S5

A Proteomics data (non-batch corrected)

B Proteomics data (batch corrected)

Batch ● TMT#1 ● TMT#2 ● TMT#3 ● TMT#4 ● TMT#5   Tissue ● PRIM

C BCL2 expression vs protein abundance with IC assignments

$R^2 = 0.45$ $P < 0.001$

D CD14+ Mono% vs protein abundance with IC assignments

$R^2 = 0.45$ $P < 0.001$

Assignment ● CD14 Mono ● EMP ● HSC ● GMP ● pro B ● BaEoMa