Multiple genomically independent genotypes of *Ophiocordyceps sinensis* may not be targets of RIP mutagenesis causing C:G to T:A transitions

Xiu-Zhang Li[a], Yu-Ling Li[a], Ya-Nan Wang[b], Jia-Shi Zhu[a,c,*]

[a] State Key Laboratory of Plateau Ecology and Agriculture, Qinghai Academy of Animal and Veterinary Sciences, Qinghai University, Xining, Qinghai 810016, China

[b] State Key Laboratory of Breeding Base of Dao-di Herbs, National Resource Center for Chinese Materia Medica, China Academy of Chinese Medical Sciences, Beijing 100700, China

[c] Institute of Biopharmaceutical and Health Engineering, Shenzhen International Graduate School, Tsinghua University, Shenzhen 518055, China

[*] Corresponding author:

E-mail address: zhujosh@163.com (J-S Zhu)

Short title:

H. sinensis genomic ITS repeats are not generated by RIP mutagenesis

Abstract

It has been hypothesized that AT-biased genotypes of *Ophiocordyceps sinensis* are generated through repeat-induced point mutation (RIP) and coexist as ITS pseudogenes in the *Hirsutella sinensis* genome. This study examined the *H. sinensis* genome, which contains multiple GC-biased repetitive ITS copies with multiple insertion/deletion and transversion alleles; these alleles are not generated through RIP, which causes cytosine-to-threonine transitions. The repetitive ITS copies are genetically and phylogenetically distinct from the AT-biased genotypes of *O. sinensis*, which possess multiple transition alleles. Genotypes #2–17 are absent from the *H. sinensis* genome; these genotypes belong to independent *O. sinensis* fungi and occur in various combinations in different compartments of natural *Cordyceps sinensis*. Their abundances are dynamically altered in an asynchronous and disproportional manner during *C. sinensis* maturation. Metatranscriptomic analyses of natural *C. sinensis* revealed the transcriptional silencing of 5.8S genes in all *C. sinensis*-colonizing fungi, including *H. sinensis*. The transcription assay reported by Li et al. (2013) has provided incomplete, controversial evidence to support the notion that the 5.8S genes of AT-biased *O. sinensis* genotypes are permanently nonfunctional pseudogenes. In parallel with GC-biased genotypes, AT-biased genotypes might have been generated from a common ancestor during a long evolutionary history.

KEY WORDS: Multiple genotypes of *Ophiocordyceps sinensis*; genomic independence; repeat-induced point mutations (RIP); insertion/deletion, transversion and transition point mutations; 5.8S gene transcript; transcriptional silencing.

I. Introduction

Natural *Cordyceps sinensis* is one of most valued therapeutic agents in traditional Chinese medicine and has a rich history of clinical use for health maintenance, disease amelioration, postdisease/surgery recovery and antiaging therapy (Zhu et al. 1998a, 1998b; Li et al. 2001; Tan et al. 2001). The Chinese Pharmacopeia defines natural *C. sinensis* as an insect-fungal complex, which includes the remains of Hepialidae moth larva and >90 fungal species from >37 genera, with differential co-occurrence of some of 17 genotypes of *Ophiocordyceps sinensis*, *i.e.*, natural *C. sinensis* insect-fungal complex ≠ the entomopathogenic *O. sinensis* fungus (Engh 1999; Kinjo and Zang 2001; Chen et al. 2004, 2011; Stensrud et al. 2005, 2007; Wei et al. 2006, 2016; Xiao et al. 2009; Zhang et al. 2009, 2010; Zhu et al. 2010, 2012, 2018, 2019; Gao et al. 2011, 2012; Yao et al. 2011; Mao et al. 2013; Xia et al. 2015; Zhu and Wu 2015; Li et al. 2016b, 2020a, 2022a, 2022c, 2023b, 2023c; Zhu and Li 2017). However, the Latin name *C. sinensis* has been indiscriminately used for both the wild insect-fungal complex and the teleomorph/holomorph of the *C. sinensis* fungus since the 1840s (Yao & Zhu 2016; Li et al. 2022a). The fungus was renamed *Ophiocordyceps sinensis* using *Hirsutella sinensis* Strain EFCC 7287 (GC-biased Genotype #1 of *O. sinensis*) as the nomenclature reference (Sung et al. 2007), while the name of the wild and cultivated insect-fungal complex has remained unchanged. Since then, especially after the improper implementation of IMA's One Fungus=One name nomenclature rule in *O. sinensis* research (Hawksworth et al. 2011; Zhang et al. 2013b; Yao & Zhu 2016; Li et al. 2022a), the name *O. sinensis* has been used to refer to multiple teleomorphic and anamorphic fungi that have distinct genomes and has been indiscriminately applied to the natural insect-fungal complex (Zhu and Wu 2015; Yao & Zhu 2016; Li et al. 2016b, 2022a, 2022c; Zhu and Li 2017). The confusion surrounding Latin name use continues and has even spread from the scientific community to the public media and mass markets, causing a significant decrease in the wholesale and retail prices of natural *C. sinensis*. Because a consensus Latin name for the wild insect-fungal product has not been identified by taxonomists and multiple genotypes of *O. sinensis* with undetermined taxonomic positions are referred to by the same Latin name (Zhang et al. 2012; Ren et al. 2013; Yao & Zhu 2016; Li et al. 2022a, 2022c, 2023c), we temporarily refer to the multiple genotype fungi as *O. sinensis*, including Genotype #1 *H. sinensis*, and we continue to refer to the wild and cultivated insect-fungal complexes as *C. sinensis*; however, this practice only partially ameliorates the academic confusion arising from the indiscriminate use of Latin names and will likely be replaced by the exclusive use of distinct Latin names in the future.

Wei et al. (2006) hypothesized that GC-biased Genotype #1 *H. sinensis* is the sole anamorph of *O. sinensis*. However, numerous studies have identified 17 mutant genotypes of *O. sinensis* in natural *C. sinensis*, which are indiscriminately referred to by the same Latin name (Zhu et al. 2010, 2018, 2019; Yao & Zhu 2016; Li et al. 2020a, 2022a, 2022c, 2023b, 2023c). Stensrud et al. (2007) and Zhang et al. (2013a) analyzed 71 and 397 ITS sequences, respectively, that were annotated as *C. sinensis* or *O. sinensis* in GenBank under taxid:72228 and confirmed the existence of 3 phylogenetic clades of mutant *O. sinensis* sequences: Group A (GC-biased Genotype #1) and Groups B–C (AT-biased Genotypes #4–5). Stensrud et al. (2007) reported that the variation in the 5.8S gene sequences of *O. sinensis* in Groups A–C "far exceeds what is normally observed in fungi … even at higher taxonomic levels (genera and family)" and that the genotypes "share a common ancestor" and originated from "accelerated nrDNA evolution". Kinjo and Zang (2001) and Mao et al. (2013) reported that the *O. sinensis* genotypes shared the same *H. sinensis*-like morphological and growth characteristics. Xiao et al. (2009) concluded that the genetically variable genotypic sequences likely belonged to independent *O. sinensis* fungi based on the differential coexistence of multiple *O. sinensis* mutants in the stroma and caterpillar body of natural *C. sinensis* in different maturational stages. In the debate regarding "cryptic species" and independent fungi of *O. sinensis*, Zhang et al. (2013a) supported the latter and disproved the former based on the results of K2P analysis. Wei et al. (2016) reported the existence of GC- and AT-biased genotypic teleomorphs of *O. sinensis* in natural and cultivated *C. sinensis*, respectively, and the species contradiction between anamorphic inoculants of GC-biased *H. sinensis* strains and a sole teleomorph of AT-biased Genotype #4 in cultivated *C. sinensis*. Li et al. (2023a) reported the differential occurrence of the mating-type genes of the *MAT1-1* and *MAT1-2* idiomorphs in the *H. sinensis* genome and the differential transcription and silencing of the mating-type genes, which invalidated the "self-fertilization" hypothesis under homothallism and pseudohomothallism (Hu et al. 2013; Bushley et al. 2013) and instead suggested the self-sterilization of *O. sinensis* requiring sexual partners to accomplish reproduction via physiological heterothallism or hybridization. In addition to the identification of molecular heterogeneity of genotypic *O. sinensis* fungi and the requirement of sexual partners for *O. sinensis* reproduction, Figure 3 of (Bushley et al. 2013) illustrates the multicellular heterokaryotic structure of *C. sinensis* hyphae and ascospores, containing multiple mono-, bi-, and tri-nucleated cells. Zhang and Zhang (2015) believed that the various heterokaryotic cells of the *C. sinensis* hyphae and ascospores likely contain different genetic materials.

The molecular heterogeneity and self-sterilization features of *O. sinensis* and the multicellular heterokaryotic structure of natural *C. sinensis* challenge the sole anamorph hypothesis for *H. sinensis* (GC-biased Genotype #1 of *O. sinensis*). Responding to the academic challenge, Li et al. (2013) proposed the "ITS pseudogene" hypothesis for all AT-biased genotypes "in a single genome" of GC-biased Genotype #1 *H. sinensis*, based on (1) the identification of heterogeneous ITS sequences of both GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis* in 8 out of 15 clones following 25 days of liquid incubation of *C. sinensis* mono-ascospores and (2) the identification of the 5.8S gene cDNA of Genotype #1 but not Genotype #5 in a cDNA library of Strain 1220, one of 8 genetically heterogeneous clones. In this "ITS pseudogene" hypothesis, the simultaneous identification of homogenous Genotype #1 in 7 other clones derived from the same mono-ascospores was apparently ignored. The authors overgeneralized the "ITS pseudogene" hypothesis to all AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* based on insufficient evidence. Li et al. (2020b) further postulated that all AT-biased *O. sinensis* genotypes "could have emerged either before or after" a new GC-biased "ITS haplotype was generated" through "repeat-induced point mutations (RIP)" that cause C-to-T (cytosine→threonine) transitions and cooccur as repetitive ITS copies in "a single genome" of GC-biased *H. sinensis*. These hypotheses proposed by Li et al. (2013, 2020b) attempted to justify the coexistence of AT-biased genotypes in natural *C. sinensis* with GC-biased *H. sinensis* under the sole *O. sinensis* anamorph hypothesis postulated by Wei et al. (2006).

The study presented in this paper examined relevant genomic and transcriptomic sequences in the GenBank database to examine the "ITS pseudogene" hypothesis and the extended hypothesis that AT-biased genotypes formed after a new GC-biased "ITS haplotype was generated" in "a single genome" of *H. sinensis* through "RIP" mutagenesis. We discuss the genetic and phylogenetic characteristics of multiple repetitive ITS copies in the genome of Genotype #1 of *O. sinensis*, the genomic independence of AT-biased genotypes, and whether sufficient evidence exists to reach a conclusion that the 5.8S genes of AT-biased genotypes are permanently nonfunctional pseudogenes.

II. Materials and Methods

II-1. Reagents

Common laboratory reagents such as agarose, electrophoresis reagents, nylon membranes, and blocking solution were purchased from Beijing Bioland Technology Company. The endonucleases *Ava*I, *Dra*I, and *Eco*RI are products of New England BioLabs, United States. Mercuric chloride (0.1%)

for surface sterilization of freshly collected *C. sinensis* specimens was a gift from the Institute of Microbiology, Chinese Academy of Sciences. The Universal DNA Purification Kit was a product of TIANGEN BIOTECH Company, China. The DNeasy Plant Mini Kit was a product of Qiagen Company, Germany. The Gel Extraction Kit was a product of Omega Bio-Tek, United States. The *Taq* PCR Reagent Kit and Vector NTI Advance 9 software were purchased from Invitrogen, United States. All primers were synthesized by Invitrogen Beijing Lab. or Beijing Bomaide Technology Co. The DIG High Prime DNA Labeling and Detection Kit was purchased from Roche Diagnostics, Indiana, USA.

II-2. Information on fungal species in natural *C. sinensis*

Information regarding intrinsic fungal species in natural *C. sinensis* was summarized by Jiang and Yao (2013), and subsequent discoveries of additional species among natural *C. sinensis* specimens were reviewed by Li et al. (2016b, 2022a, 2022c). The sequences of 17 genotypes of *O. sinensis* have been obtained and uploaded to GenBank by many molecular mycology groups since 1999. A BLAST search of the GenBank database revealed 668 ITS sequences of *O. sinensis*, which were uploaded by many *O. sinensis* research groups worldwide. These genotypic sequences were analyzed genetically and phylogenetically by Stensrud et al. (2007), Zhu et al. (2010, 2012), Zhang et al. (2013a), Zhu & Wu (2015), Li et al. (2016b, 2022a. 2022c, 2023c), and Zhu & Li (2017).

II-3. Genetic, genomic and transcriptomic sequences from GenBank

Five genome assemblies, ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000, for *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively, were uploaded to GenBank (Hu et al. 2013; Shu et al. 2020; Li et al. 2016a; Jin et al. 2020; Liu et al. 2020) and used for the analysis of repetitive ITS copies and multiple GC- and AT-biased genotypes.

A Basic Local Alignment Search Tool (BLAST) search of the GenBank database revealed 668 *O. sinensis* ITS sequences under GenBank taxid:72228, 440 of which belong to GC-biased Genotypes #1–3 and #7–14 and 228 belong to AT-biased Genotypes #4–6 and #15–17 (Stensrud et al. 2005,2007; Zhang et al. 2013a; Li et al. 2016b, 2022c, 2023b, 2023c).

mRNA transcriptome assembly GCQL00000000 for *H. sinensis* Strain L0106 was uploaded to GenBank by Liu et al. (2015). Two metatranscriptome studies of natural *C. sinensis* have been

reported. Xiang et al. (2014) uploaded the GAGW00000000 metatranscriptome assembly to GenBank, which was obtained from natural specimens (unknown maturational stages) collected from Kangding County of Sichuan Province, China. Xia et al. (2017) uploaded the unassembled SRR5428527 Shotgun Read Archive (SRA) to GenBank for mature *C. sinensis* specimens collected from Deqin County of Yunnan Province, China, but the assembled metatranscriptome sequences were uploaded to www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas, which is currently inaccessible, but a previously downloaded file is used for analysis.

II-4. Genetic and phylogenetic analyses

The ITS sequences of multiple genotypes of *O. sinensis* and multiple genomic repetitive ITS copies as well as metatranscriptomic/transcriptomic sequences were analyzed using the GenBank BLAST algorithms (https://blast.ncbi.nlm.nih.gov/blast/).

A Bayesian phylogenetic tree of the ITS sequences was inferred using MrBayes v3.2.7 software (Markov chain Monte Carlo [MCMC] algorithm) based on 3×10^4 samples, with a sampling frequency of 10^3 iterations after discarding the first 25% of samples from a total of 4×10^6 iterations (Huelsenbeck and Ronquist 2001; Li et al. 2022a, 2023c); this phylogenetic analysis was performed at Nanjing Genepioneer Biotechnologies Co.

II-5. Collection and on-site processing of fresh *C. sinensis*

Fresh *C. sinensis* specimens were purchased in a local market in Kangding County of Sichuan Province, China (3800–4600 m' altitude; latitude 30°049'N, longitude 101°959'E) (Zhu et al. 2010; Yao et al. 2014). Immature *C. sinensis* specimens were characterized by a plump caterpillar body and very short stroma (1.0–2.0 cm). Maturing *C. sinensis* specimens were characterized by a plump caterpillar body and stroma of 2.5–5.0 cm in length, without the formation of an expanded fertile portion close to the stromal tip. Mature *C. sinensis* specimens were characterized by a plump or slightly less plump caterpillar body and long stroma (>5.0 cm) and by the formation of an expanded fertile portion close to the stromal tip, which was densely covered with numerous ascocarps (Zhu et al. 2010; Li et al. 2023a). Governmental permission was not required for *C. sinensis* purchases in local markets, and the collections of *C. sinensis* specimens from sales by local farmers fall under the governmental regulations for traditional Chinese herbal products.

Freshly collected *C. sinensis* specimens were washed thoroughly in running water with gentle

brushing and surface sterilized in 0.1% mercuric chloride for 10 min, followed by washing 3 times with sterile water (Zhu et al. 2010; Li et al. 2023c). The thoroughly cleaned and sterilized *C. sinensis* specimens were immediately frozen in liquid nitrogen and kept frozen during transportation and storage prior to further processing.

II-6. Extraction of genomic DNA and preparation of *C. sinensis* DNA

Caterpillar bodies and stromata of sterilized, frozen *C. sinensis* were separately ground into powder in liquid nitrogen. Genomic DNA was individually extracted from the powder samples using a DNeasy Plant Mini Kit.

II-7. PCR amplification and amplicon sequencing

Genomic DNA of immature, maturing and mature stromata of natural *C. sinensis* was used as the template for PCR amplification. *H. sinensis*-specific primers were designed (a forward primer *Hsprp1*: 5' ATTATCGAGTCACCACTCCCAAACCCCC 3'; and a reverse primer *Hsprp3*: 5' AGGTTCTCAGCGAGCTA 3') based on alignment analysis of the ITS sequences (AB067721, AB067740, and AB067744 representing Genotypes #1 and #4–5 of *O. sinensis*, respectively).

A touch-down PCR protocol was used, in which the annealing temperature was initially set at 70 °C and was then decreased stepwise with a decrement of 0.3 °C every cycle, for a total of 36 cycles (Zhu et al. 2007, 2010). PCR amplicons were subjected to agarose gel electrophoresis, and DNA bands were recovered, purified using a gel extraction kit, and sequenced either directly or after cloning of the amplicons.

II-8. Preparation of fungus-specific ITS1 probe and 18S internal control probe

An *H. sinensis*-specific probe (158 bp) was designed based on the ITS1 sequence of GC-biased Genotype #1 *H. sinensis* and generated through PCR using the forward primer *Hsprp1* described above and the reverse primer *Hsprp2* (5' ATTTGCTTGCTTCTTGACTGAGAGATGCC 3').

A nonspecific 18S internal control probe was designed based on alignments of GenBank 18S rDNA sequences (DQ838795, AB187268, AB067735, AB067736, and AB080090) of several *C. sinensis*-associated fungi. A highly homologous region of 18S rDNA, located over 700 bp upstream of the ITS1 regions of *C. sinensis*-associated fungi, was selected. The 18S control probe was generated through PCR using the forward primer *Icprp1* (5' ACCGTCGTAGTCTTAACCAT 3') and the reverse primer *Icprp2* (5' ATCGGCTTGAGCCGATAGTC 3').

219 Both amplicons were confirmed by sequencing, and the probes were labeled using the DIG High
220 Prime DNA Labeling and Detection Kit.

II-9. Southern blotting analysis

The stromata and caterpillar bodies of immature and mature *C. sinensis* were used for Southern
blotting. Genomic DNA was extracted from the samples and prepared by endonuclease digestion
using *Ava*I, *Dra*I, and *Eco*RI. The digested genomic DNA (5 µg of each sample) was subjected to
agarose gel electrophoresis and transferred onto a nylon membrane through capillary transfer,
followed by baking the membrane at 58 °C for 4 hours for fixation, blocking the binding sites on the
membrane in blocking solution, and hybridizing with the labeled *H. sinensis*-specific probe at 40 °C
overnight. The hybridized transblotted membrane was viewed by colorimetric photography using a
gel imaging system (Bio-Rad) and then stripped and re-probed with the labeled 18S internal control
probe.

III. The results

III-1. Multiple GC-biased repetitive ITS copies in the *H. sinensis* genome

Generally, fungal genomes contain multiple repetitive copies of nrDNA ITS1-5.8S-ITS2
sequences. However, a single copy of the ITS1-5.8S-ITS2 sequences was identified in genome
assemblies ANOV01021709, LKHE01000582, and LWBQ01000008 of *H. sinensis* Strains Co18,
1229, and ZJB12195, respectively (Hu et al. 2013; Li et al. 2016a; Jin et al. 2020), probably because
the repetitive ITS copies were discarded during the assembly of the genome shotgun sequences. On
the other hand, multiple repetitive ITS copies were identified in 2 additional genome assemblies,
JAAVMX000000000 and NGJJ00000000, for *H. sinensis* Strains IOZ07 and CC1406-203,
respectively (Liu et al. 2020; Shu et al. 2020) (Table 1), where 17 ITS1-5.8S-ITS2 copies within
JAAVMX000000000 and 7 ITS1-5.8S-ITS2 copies were found, as well as 2 partial ITS segments
within NGJJ00000000 that were not included in our analysis.

Most of the repetitive ITS copies listed in Table 1 were highly homologous (>97%) to the
reference sequence AB067721 of GC-biased Genotype #1 *H. sinensis*. Two repetitive copies
(6233→6733 and 44729→45251) within JAAVMX010000019 showed 94.5% and 90.8% similarities
to AB067721, respectively (Table 1). Two other copies, 18702095→18702586 within
JAAVMX010000002 and 700→1186 within JAAVMX010000018, showed similarities of 97.4% and

97.0% to AB067721, respectively. All the repetitive ITS copies showed 80.1%–89.9% similarity to

the sequences of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* (Table 1).

Table 1. *GC contents of the repetitive ITS1-5.8S-ITS2 sequences from the genome assemblies of 2 H.*
sinensis strains and percent similarities of the repetitive ITS sequences to the sequences of GC-biased
Genotype #1 (AB067721) and AT-biased Genotypes #4–6 and #15–17 of O. sinensis.

Genome segment	Sequence range & direction	GC content	% Similarity of the repetitive sequence	
			vs. Genotype #1 AB067721	*vs.* AT-biased Genotypes #4–6 and #15–17
JAAVMX010000002	18688917→18689407	64.8%	100%	86.7% – 89.9%
	18702095→18702586	64.5%	97.4%	82.8% – 88.0%
JAAVMX010000008	13823→14313	64.8%	100%	85.5% – 89.9%
	1199→1687	64.8%	99.2%	85.2% – 89.3%
JAAVMX010000017	9147←9637	64.8%	100%	85.5% – 89.9%
	21791←22281	64.8%	100%	85.5% – 89.9%
	34435←34925	64.8%	100%	85.5% – 89.9%
	47079←47569	64.8%	100%	85.5% – 89.9%
JAAVMX010000018	13381→13871	64.8%	100%	85.5% – 89.9%
	26076→26566	64.8%	100%	85.5% – 89.9%
	38771→39261	64.8%	100%	85.5% – 89.9%
	51467→51958	64.6%	98.8%	85.5% – 88.8%
	700→1186	64.5%	97.0%	82.6% – 87.6%
JAAVMX010000019	19404→19894	64.8%	100%	85.5% – 89.9%
	32048→32538	64.8%	100%	85.5% – 89.9%
	6233→6733	65.3%	94.5%	81.3% – 85.5%
	44729→45251	63.3%	90.8%	80.1% – 84.6%
NGJJ01000573	13133←13623	64.8%	99.4%	85.2% – 89.6%
	14624←15113	64.7%	99.4%	84.8% – 89.6%
NGJJ01000582	10703←11192	64.7%	99.8%	85.5% – 89.8%
NGJJ01000796	1514→2004	64.8%	100%	85.5% – 89.9%
NGJJ01000798	1177→1665	64.6%	99.6%	84.8% – 89.6%
NGJJ01000799	3295→3785	64.8%	100%	85.5% – 89.9%
	16666→17156	64.8%	100%	85.5% – 89.9%

253 **III-2. Genetic characteristics of the multiple repetitive ITS sequences in the *H.**
254 ***sinensis* genome**

The repetitive ITS copies within genome assemblies JAAVMX000000000 and NGJJ00000000 for *H. sinensis* Strains IOZ07 and CC1406-203, which are listed in Table 1, and the single genomic ITS copies ANOV01021709, LKHE01000582, and LWBQ01000008, for *H. sinensis* Strains Co18, 1229, and ZJB12195, respectively, which are listed in Table 2 (Hu et al. 2013; Jin et al. 2020; Li et al. 2016a; Liu et al. 2020; Shu et al. 2020), were GC biased; their average GC content was 64.7±0.33% (Table 1), similar to the GC content (64.8%) of the AB067721 Genotype #1 sequence. Apparently, these genetic features were distinct from those of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis*, which included multiple scattered transition alleles and contained 46.9%–55.2% AT bases and 44.8%–53.1% GC bases.

Table 2. *Percent similarities between nrDNA ITS1-5.8S-ITS2 sequences of 5 genome assemblies of Genotype #1 H. sinensis and 17 O. sinensis genotype sequences. (Modified from Table 1 of Li et al. 2023c).*

Genotype # and GenBank accession #	% Similarity between the *H. sinensis* genomic ITS segments and 17 *O. sinensis* genotypes				
	ANOV01021709 838→1504	LKHE01000582 2074→2740	LWBQ01000008 991739→992405	JAAVMX010000017 9029←9695	NGJJ01000799 3237→3903
#1 AB067721	99.7%	100%	99.6%	100%	100%
#2 MG770309	94.4%	94.7%	94.7%	94.7%	94.7%
#3 HM595984	95.9%	96.1%	95.6%	95.3%	95.3%
#7 AJ488254	94.7%	95.0%	95.3%	95.0%	95.0%
#8 GU246286	89.5%	89.5%	89.2%	89.5%	89.5%
#9 GU246288	95.0%	95.2%	94.6%	95.2%	95.2%
#10 GU246287	83.0%	83.2%	83.0%	83.2%	83.2%
#11 JQ695935	78.9%	79.1%	78.5%	79.1%	79.1%
#12 GU246296	94.3%	94.5%	93.9%	94.5%	94.5%
#13 KT339190	87.5%	87.5%	87.2%	87.5%	87.5%
#14 KT339178	89.3%	89.6%	89.1%	89.6%	89.6%
#4 AB067744	89.1%	89.4%	89.3%	89.4%	89.4%
#5 AB067740	86.3%	86.7%	86.2%	86.7%	86.7%
#6 KJ720572	85.5%	85.5%	85.5%	85.5%	85.5%
#15 KT232017	89.6%	89.9%	89.4%	89.9%	89.9%
#16 KT232019	87.4%	87.7%	87.6%	87.7%	87.7%
#17 KT232010	87.7%	88.1%	87.5%	88.1%	88.1%

Note: "GT" represents genotype. GC-biased genotypes are listed in blue in the upper panel. AT-

biased genotypes are listed in red in the lower panel.

The sequences of GC-biased repetitive ITS copies that were variable (94.5% and 90.8% similarities) and slightly less variable (97.4% and 97.0% similarities) in comparison to the reference sequence AB067721 of Genotype #1 predominantly contained insertion/deletion alleles and included less frequent transversion alleles, while no or only a few transition alleles were found. The variable ITS repeats 6233→6733 and 44729→45251 within JAAVMX010000019 (*cf.* Table 1) contained 22 and 32 insertion/deletion alleles and 4 and 11 transversion alleles, respectively (Table 3; in blue in Figure 1); they contained only 1 and 5 transition mutant alleles, having allelic ratios of insertion/deletion/transversion *vs.* transition of 26:1 and 43:5, respectively. The slightly less variable ITS repeats, 18702095→18702586 within JAAVMX010000002 and 700→1186 within JAAVMX010000018 (*cf.* Table 1), contained no transition alleles and predominantly included 13 and 15 insertion, deletion and transversion alleles, with allelic ratios of insertion/deletion/transversion *vs.* transition of 13:0 and 15:0, respectively (Table 3; in green in Figure 1). These genetic characteristics of the variable and slightly less variable repetitive ITS copies indicate that they might not have been generated through RIP mutagenesis, which theoretically causes C-to-T (cytosine→threonine) transition mutations. Other repetitive ITS copies with 98.1%–99.8% similarity to AB067721 listed in Table 1 contained only a few insertion/deletion alleles.

Table 3. *Types of mutations in the variable and slightly less variable genomic repetitive ITS sequences compared with GC-biased Genotype #1 (AB067721).*

	Sequence range & direction	% Similarity	Number of mutant alleles			Ins./Del./Transv. *vs.* Transit.
			Ins./Del.	Transv.	Transit.	
JAAVMX010000002	18702095→18702586	97.4%	13	0	0	13 : 0
JAAVMX010000018	700→1186	97.0%	10	5	0	15 : 0
JAAVMX010000019	6233→6733	94.5%	22	4	1	26 : 1
	44729→45251	90.8%	32	11	5	43 : 5

Note: "Ins./Del." indicates insertion/deletion point mutations. "Transv." denotes Transversion point mutations. "Transit." refers to transition point mutations.

```
GT#1  AB067721   59       TCGAGTCACCACTCCC AAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCCGGCG CCCCAGGGGCCCG 136
JAAVMX010000019 19404     ---------------- -------------------------------------------------- ------------ 19481
JAAVMX010000019 32048     ---------------- -------------------------------------------------- ------------ 32125
JAAVMX010000002 18702095  ---------------- --------------------------------------------T----------- 18702172
JAAVMX010000018 700       ---------  ------ -----------------------------T----------- ----------T 776
JAAVMX010000019 6233      ----------------C- -------------------------------------------------- ------------ 6311
JAAVMX010000019 44729     ---------------- -------------------------------------------------- ------------ 44806
GT#4  AB067744   55       ------T------T --------------T--T---------------------------T------- 132
GT#5  AB067740   59       ------T--T----TT --------ATA-T--T-T----------TA------------TA--- ---T-A---- 136
GT#6  KJ720572   11       ------A---T--T-T----------TA------------TA--- --TT-A----- 70
GT#15 KT232017   37       ------T------T --------------T--T---------------------------T------- 114
GT#16 KT232019   37       ------T--T-----T -------T--T-T----------TA------------TA--- ---T-A---- 114
GT#17 KT232010   37       ------T--T-----T -------T--T-T----------TA------------TA--- ---T-A---- 114

GT#1  AB067721  137       GACCAGG GCGCCC GCCGGAGGACCCCCAGAC CCTCCTGTCGCA GTGGCATCT CTCAGTCAAGAA GCAAGCAAAT 210
JAAVMX010000019 19482     ------- ------ ------------------ ------------ --------- ------------ ---------- 19555
JAAVMX010000019 32126     ------- ------ ------------------ ------------ --------- ------------ ---------- 32199
JAAVMX010000002 18702173  ------- ------A------------- ------------ -----A--------- ---- ------- ---------- 18702247
JAAVMX010000018 777       ------- ------ ---------------- ------------ ---------- --C------- -------- 850
JAAVMX010000019 6312      ------- ------ ------T----------- ---C-------- --------C------------G--------- 6388
JAAVMX010000019 44807     -------C------ ---------------- ------------ --------- ------------ ---------- 44881
GT#4  AB067744  133       ---T--- ------ ------------TT---- ------A---T- --A------ --T---T----- ---------- 206
GT#5  AB067740  137       ---T-AA ------ ------------TT---- ------A-TAT- --A-T---- --T---T----- ---------- 213
GT#6  KJ720572   71       ---T--- ------ ------------TT---- ------A---T- --A-T---- --T---T----- ---------- 182
GT#15 KT232017  115       ---T--- ------ ------------TT---- ------A----- --A------ --T---T----- ---------- 188
GT#16 KT232019  115       ---T-AA ------ ------------TT---- ------A----- --A-T---- --T---T----- ---------- 188
GT#17 KT232010  115       ---T-AA ------ ------------TT---- ------A-TAT- --A-T---- --T---T----- ---------- 188

GT#1  AB067721  211       GAATCAAAACTTTCAACAACGGATCTCTTGGTTCTGGCATC GATG AAGAACGCAGCGAAATGCGATAAGTAATGTGAA 288
JAAVMX010000019 19556     ----------------------------------------- ---- -------------------------------- 19633
JAAVMX010000019 32200     ----------------------------------------- ---- -------------------------------- 32277
JAAVMX010000002 18702248  -- ------------------------------------C---- -------------------------------- 18702325
JAAVMX010000018 851       ----------------------------------- ----A------------------- ---------- 928
JAAVMX010000019 6389      -------------- ---CAC---------------- ---- ---------- ---------- ------- 6463
JAAVMX010000019 44882     ----------------------------------------- ---- -------------------------------- 44959
GT#4  AB067744  207       A---T--------T--T-----------A---T-A----T A--A ----------------A----------A-- 284
GT#5  AB067740  214       A---T-------T---------A-----A----T A--A --A----------------A-------A-A-- 288
GT#6  KJ720572  183       A---T------T-----------A-----A----- ---A -------T--T----A----------A-A- 222
GT#15 KT232017  189       ----T------T--T-----------A---T-A----T A--A ----------------A----------A-- 266
GT#16 KT232019  189       ----T------T----------A-----A----- ---A --A----------------A-------A-A- 266
GT#17 KT232010  189       A---T-------T----------A-----A----- ---A --A----------------A-------A-A-- 266

GT#1  AB067721  289       TTGCAGAATTCAGTGAACCA TCGAATCTTT GAACG CACATTG CGCCCGCCAGCACTCTGGCGGGCATGC CTGTC 362
JAAVMX010000019 19634     -------------------- ---------- ----- ------- --------------------------- ----- 19707
JAAVMX010000019 32278     -------------------- ---------- ----- ------- --------------------------- ----- 32351
JAAVMX010000002 18702326  -------------------- ----------T----- ----- - ------ --------------------- ----- 18702398
JAAVMX010000018 929       ------------------ C---------- -----C---------C-------------------- ------- ----- 1000
JAAVMX010000019 6464      -------------------- ---C------- ----- ----- -------C----------------- ----- 6539
JAAVMX010000019 44960     -------------------- ---------- ----- ----- -------------------------CT----- 45035
GT#4  AB067744  285       ---T-------T---A---T- ---------- A----- T-T-C-A -------T--------A--------- ----- 358
GT#5  AB067740  289       --A-------T---A---T- ---------- A----- --T---A -------T--------A--------- --A-- 362
GT#6  KJ720572  223       --AT-------T---A---- ---------- A----- T-T---A -------T--------A--------- ----- 296
GT#15 KT232017  267       ---T-------T---A---T- ---------- A----- T-T---A -------T--------A--------- ----- 340
GT#16 KT232019  267       --A-------T---A---T- ---------- A----- T-T---A -------T--------A--------- ----- 340
GT#17 KT232010  267       --A-------T---A---T- ---------- A----- --T---- -------T--------A--------- --A-- 340

GT#1  AB067721  363       CGAGCGTCATCTCAACCCTCGAG CCCCCCGCCTCGCGG CGGCGGGGCCCGGCCTT GGGGGTCACGGCC CCGCGCC 437
JAAVMX010000019 19708     ----------------------- --------------- ----------------- ------------- ------- 19781
JAAVMX010000019 32352     ----------------------- --------------- ----------------- ------------- ------- 32425
JAAVMX010000002 18702399  ----------------------- ----- --------- ----------------- ------------- ------- 18702472
JAAVMX010000018 1001      ----------------------- ------ -------- ----------------G------------ --- -- 1074
JAAVMX010000019 6540      ----------------------- --------------- ----------------- ---- -------C------- 6613
JAAVMX010000019 45036     ----------------------C---------A-----TA--CG---C-------- ------TC-----AA------ 45119
GT#4  AB067744  359       -------T-C--T------TA-- --------------- ----------------- A---------- --A---- 432
GT#5  AB067740  363       -------T----T---------- --------------- ---------T------- A-----T------ ------- 436
GT#6  KJ720572  297       -------T-C--T---- 313
GT#15 KT232017  341       -------T----T---------- --------------- ---------T------- A-----T------ ------- 414
GT#16 KT232019  341       -------T-C--T------TA-- --------------- ---------T------- A---------- --A---- 414
GT#17 KT232010  341       -------T----T---------- --------------- ---------T------- A-----T------ ------- 414

GT#1  AB067721  438       G   CCCCCTAAACGC AG     TGGCGACC CCGCCGCGGCTCC CCTGCGCAGTAGCT  CG CTGAGAACCT 499
JAAVMX010000019 19782     -   ------------ --     -------- ------------- ---------------  -- ---------- 19844
JAAVMX010000019 32426     -   ------------ --     -------- ------------- ---------------  -- ---------- 32488
JAAVMX010000002 18702473  -   ------------ --     -------- ------------- ---------------  -- ---------- 18702534
JAAVMX010000018 1075      -   ------------ --     -------- ------------- ---------------  -- ---------- 1136
JAAVMX010000019 6614      -   -----------G--G     --------GC------------- -------------  --GCT--------- 6680
JAAVMX010000019 45120     -TCC-T---GT----- --CCTAAGTG------- ----GA-------A----T-A-------CA-- ---------- 45189
GT#4  AB067744  433       -   ------------ --     -A-T---- ------------- ---A---A-----  TA --A------ 495
GT#5  AB067740  437       -   ----T------T --     -A------ ------------- ---A--TG-----  -A -C------- 529
GT#6  KJ720572  313
GT#15 KT232017  415       -   -----------T --     -A------ ------------- ---A--T-A-----  -A -C------- 477
GT#16 KT232019  415       -   ----------- --     -A------ ------------- ---A---A-----  TA --A------ 477
GT#17 KT232010  415       -   -----------T --     -A------ ------------- ---A--T-A-----  -A -C------- 477

GT#1  AB067721  500       CGCACCGGGAGCGCGGA    GGCGG TCACGCC  TGG    AA ACCACC  ACACCCTCCA          549
JAAVMX010000019 19845     -----------------    ----- ----- - ---    -- ------  ----------          19894
JAAVMX010000019 32489     -----------------    ----- ----- - ---    -- ------  ----------          32537
JAAVMX010000002 18702535  -----------------    ----- -------GTGT-  -- ------  ----------          18702586
JAAVMX010000018 1137      -----------------    ----- ----- - ---    -- ------  ----------          1186
JAAVMX010000019 6681      C----------------    ----- ----- - ---    -- ------ACA----------          6733
JAAVMX010000019 45190     ----------------CCCAT-----C------- ---GATG--C------ACC----------          45251
GT#4  AB067744  496       -----TA----------    ----- -------  --A    -- -----T  -T------          543
GT#5  AB067740  530       --------A--------    ----- -------  --A    -- -----T  --G--          544
GT#6  KJ720572  313
GT#15 KT232017  478       --------A--------    ----- -------  --A    -- -----T  --G--          522
GT#16 KT232019  478       -----TA----------    ----- -------  --A    -- -----T  -T------          525
GT#17 KT232010  478       --------A--------    ----- -------  --A    -- -----T  --G--          522
```

Figure 1. ITS sequence alignment of AB067721, repetitive ITS copies within genome JAAVMX000000000 of H. sinensis Strain IOZ07, and AT-biased genotypes of O. sinensis.

The ITS sequences contain complete or partial ITS1-5.8S-ITS2 nrDNA segments. "GT" denotes the O. sinensis genotype. The underlined sequence in black represents the 5.8S gene of GC-biased Genotype #1 H. sinensis. AB067721 is the ITS sequence of GC-biased Genotype #1 H. sinensis. Genome assemblies JAAVMX010000002, JAAVMX010000018, and JAAVMX010000019 were obtained from the H. sinensis Strain IOZ07 (Shu et al. 2020). One copy each within JAAVMX010000002 and JAAVMX010000018 indicated in green shows 97.4% or 97.0% similarity to AB067721. JAAVMX010000019 contains 4 repetitive ITS copies, including 2 copies in black (19404→19894 and 32048→32537), which are 100% identical to AB067721, and 2 other copies in blue (6233→6733 and 44729→45251), with 94.5% and 90.8% similarities to AB067721. Sequences in red (AB067744, AB067740, KJ720572, KT232017, KT232019, and KT232010) represent AT-biased Genotypes #4–6 and #15–17 of O. sinensis, respectively. The underlined "GAATTC" sequences in purple represent the EcoRI endonuclease cleavage sites in the sequences of GC-biased Genotype #1 and GC-biased genome assembly JAAVMX000000000. EcoRI endonuclease cleavage sites are absent in AT-biased ITS sequences due to a single-base C-to-T (cytosine→threonine) transition. Hyphens indicate identical bases, and spaces denote unmatched sequence gaps.

III-3. Phylogenetic features of multiple repetitive ITS copies in the *H. sinensis* genome

As shown in Tables 1–2 and Figure 1, the sequences of GC-biased Genotypes #2–3 and #7–14 and of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* are absent in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000 of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively, and instead belong to the genomes of independent *O. sinensis* fungi (Stensrud et al. 2007; Xiao et al. 2009; Hu et al. 2013; Zhu & Wu 2015; Li et al. 2016a, 2016b, 2020a, 2022a, 2022c, 2023c; Zhu & Li 2017; Zhu et al. 2018, 2019; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). The genomic repetitive ITS copies were phylogenetically clustered into the GC-biased clades in blue alongside the Bayesian majority rule consensus tree in Figure 2 and were phylogenetically distant from the AT-biased clades containing Genotypes #4–6 and #15–17 of *O. sinensis* shown in red alongside in the tree, similar to what has previously been demonstrated in phylogenetic trees (Kinjo & Zang 2001; Stensrud et al. 2007; Zhang et al. 2013a; Zhu & Wu 2015; Li et al. 2016b, 2020a, 2022a, 2022c, 2023c; Wei et al. 2016; Zhu & Li 2017).

 Figure 2. A Bayesian majority rule consensus phylogenetic tree.
GT represents genotype. Twenty-eight ITS segments of the genome assemblies (ANOV01021709,
LKHE01000582, LWBQ01000008, JAAVMX010000002, JAAVMX010000008, JAAVMX0100000017,
JAAVMX0100000018, JAAVMX010000019, NGJJ01000573, NGJJ01000582, NGJJ01000796,
NGJJ01000798, and NGJJ01000799) of H. sinensis strains (Co18, 1229, ZJB12195, IOZ07, and CC1406-203,
respectively) and 25 ITS sequences of GC-biased Genotypes #1–3 and #7–14 (in blue *alongside the tree) and*
AT-biased Genotypes #4–6 and #15–17 of O. sinensis (in red *alongside the tree) were analyzed*
phylogenetically using MrBayes v3.2.7 software (cf. Methods Section II-4). Genome assemblies
JAAVMX000000000 and NGJJ00000000 contain multiple repetitive ITS copies (cf. Table 1). The percent
similarities of the genomic sequences of repetitive ITS copies with the representative Genotype #1 sequence
(AB067721) are shown in green *alongside the tree.*

Figure 2 includes the 2 genomic repetitive ITS copies with low similarities (94.5% and 90.8%) and the 2 repetitive ITS copies with relatively low similarities (97.4% and 97.0%) that are shown in Tables 1 and 3. The ITS copies are individually scattered in the GC-biased clades. The 2 low-similarity genomic repetitive ITS copies, 6233→6733 (94.5% similarity) and 44729→45251 (90.8% similarity), within genome assembly JAAVMX010000019 (*cf*. Tables 1 and 3), showed greater phylogenetic distances (horizontal lines) from other GC-biased ITS copies within the GC-biased clades. GC-biased Genotypes #2–3 and #7–14 of *O. sinensis* showed greater phylogenetic distances from the sequences of GC-biased Genotype #1 and the repetitive ITS copies. These results indicate that genetic variations in the GC-biased repetitive ITS sequences shown in blue in Figure 1 had an impact on Bayesian phylogenetic clustering analysis.

III-4. Genetic characteristics of the multiple repetitive ITS sequences in the *H. sinensis* genome

The Bayesian tree in Figure 2 shows that GC-biased Genotype #7 was placed within the Genotype #1 clade with a great phylogenetic distance, although it showed 94.7%–95.3% similarity to Genotype #1 *H. sinensis* (*cf*. Table 1). The Genotype #7 sequence (AJ488254) was identified in the stroma of a *C. sinensis* specimen (H1023) collected from Qinghai Province and uploaded to GenBank by Chen et al. (2011). The authors identified GC-biased Genotype #1 ITS sequence (AJ488255) from the tail of the caterpillar body of the same specimen, giving rise to the question of whether the sequences of AJ488254 and AJ488255 coexisted in a single genome of GC-biased Genotype #1 *H. sinensis* as 2 of repetitive ITS copies or belonged to the genomes of independent *O. sinensis* fungi cooccurring in different compartments of the same *C. sinensis* specimen. The GC-biased genotype sequences were further analyzed, as shown in Figure 3, and compared with the repetitive ITS copies

in the genome.

```
GT#1   AB067721    59    TCGAGTC ACCACTCCC AAA CCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCCGGCG CCCCAGGGCCCG 136
JAAVMX010000019 19404      ------- --------- --- -------------------------------------------- ------------ 19481
JAAVMX010000019 32048      ------- --------- --- -------------------------------------------- ------------ 32125
JAAVMX010000002 18702095   ------- --------- --- ---------------------------------------T---- ------------ 18702172
JAAVMX010000018 700        ------- -- ------- --- -----------------------------------T----------- ----------T 776
JAAVMX010000019 6233        ------- --------C--- -------------------------------------------- ------------ 6311
JAAVMX010000019 44729      ------- --------- --- -------------------------------------------- ------------ 44806
GT#2   MG770309
GT#3   HM595984    33    CT----T G--------- --- A----C-T------------------------------------ ------------ 110
GT#7   AJ488254   464    ------- -A----T-- G--G------------------------------------T-T---- -A----TG---- 380
GT#8   GU246286     4    ------T --T------ -- ---------------------------T------T---- --G---T---- 83
GT#9   GU246288     5    ------TT--------- -- -----------------------C---G------------- ------------ 84
GT#10  GU246287     6    ------T -G ------ T-- ---------------T----T----TTTG---------- -----A------ 82
GT#11  JQ695935    12    ------T --------- --- -------------------------------------------- ------------ 78
GT#12  GU246296     5    ------T --------- --- -------------------------------------------- ------------ 83

GT#1   AB067721   137    GACCAGG GCGCCC GCCGGAGGACCCCCAGAC CCTCCTGTCGCA GTGGCATCT CTCAGTCAAGAA GCAAGCAAATGA 213
JAAVMX010000019 19482      ------- ------ ----------------- ------------ --------- ------------ ------------ 19557
JAAVMX010000019 32126      ------- ------ ----------------- ------------ --------- ------------ ------------ 32201
JAAVMX010000002 18702173   ------- ------A---------------- ----------A- ---- ------- ------------ 18702249
JAAVMX010000018 777        ------- ------ ----------------- ------------ ---C------- ------------ 852
JAAVMX010000019 6312        ------- ------ -------T--------- ---C-------- --------C-- ------G------ 6390
JAAVMX010000019 44807      -------C------ ----------------- ------------ --------- ------------ 44883
GT#2   MG770309     1          ----G AT--ATTATT-A----C- -----C-A---- -------T- ------------ 68
GT#3   HM595984   111    ---T--- ------ ----------------- ------------ ---G--- ------------ 186
GT#7   AJ488254   379    A------- ------ ----------------- ------------ --------- ------------ 295
GT#8   GU246286    84    -C-G--- -----T -GAAAT-TT--AA-T-- --------C--- ------------ 159
GT#9   GU246288    85    ------- ------ ----------------- ------------ ------G- -----G------ 160
GT#10  GU246287    83    ------- ------ --------G-------- --------G ---GT-T- T---------- AGG--G------ 157
GT#11  JQ695935    79    ------- ------ ----------------- ------------ --------- ------------ 154
GT#12  GU246296    84    ------- ------ ----------------- ------------ --------- ------------ 159

GT#1   AB067721   214    ATCAAAACTTTCAACAACGGATCTCTTGGTTCTGGCATC GATG AAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGC 292
JAAVMX010000019 19557      --------------------------------------- ---- ------------------------------------ 19637
JAAVMX010000019 32201      --------------------------------------- ---- ------------------------------------ 32281
JAAVMX010000002 18702249   -------------------------------C---- ------------------------------------ 18702329
JAAVMX010000018 852        ----------------------------A--- ----------------------------------- 932
JAAVMX010000019 6390        ----------- --CAC--------------- ---- ------------- -------- ---------- 6467
JAAVMX010000019 44883      --------------------------------------- ---- ------------------------------------ 44963
GT#2   MG770309    68    --------------------- --- --- -----GA--------C ------------------- 147
GT#3   HM595984   186    --------------------------------------- ---- ------------------------------------ 266
GT#7   AJ488254   295    --------------------------------------- ---- ----A--------------------------- 224
GT#8   GU246286   159    --------------T-------------T---- ---- -G------------G----------------- 239
GT#9   GU246288   161    --------------------------T------- ---- ----------------------------------- 240
GT#10  GU246287   158    ------T--------------T----------- ---- -----C---C---------------G---------- 237
GT#11  JQ695935   155    --------------------------------------- ---- ------------------------------------ 234
GT#12  GU246296   160    --------------------------------------- ---- ---------------------------------G-- 239

GT#1   AB067721   293    AGAATTCAGTGAACCA TCGAATCTTT GAACG CACATT G CGCCCGCCAGCACTCTGGCGGGCATGC  CTGTCCGAGC 367
JAAVMX010000019 19638      ---------------- ---------- ----- ------ - ------------------------- ---------- 19712
JAAVMX010000019 32282      ---------------- ---------- ----- ------ - ------------------------- ---------- 32356
JAAVMX010000002 18702330   ---------------- ----------T----- - ------ - ------------------------- ---------- 18702403
JAAVMX010000018 933        ---------------- C---------- -----C------ --C--------------- ------- ---------- 1005
JAAVMX010000019 6468        ----------------C----------- ----- - --C----------------- ---------- 6544
JAAVMX010000019 44964      ---------------- ---------- ----- ------ - ----------------------CT---------- 45040
GT#2   MG770309   148    ---------------- ---------- ----- ------ - ------------------------- ---------- 222
GT#3   HM595984   267    ---------------- ---------- ----- ------ - ---------A--------- ---------- 341
GT#7   AJ488254   223    ---------------- ---------- -----A---- - ------------------------- ---------- 147
GT#8   GU246286   240    ---------------- --T-T-T--- ----- ------ - ----------------A---- ---------- 314
GT#9   GU246288   241    ---------------- ---------- ----- ------ - -------------C----------- ---------- 315
GT#10  GU246287   238    ---------------- ---------- ----- ------G- G---G--------------------- T-------- 313
GT#11  JQ695935   235    ---------------- ---------- ----- ------ - ------------------------- ---------- 309
GT#12  GU246296   240    ---------------- ---------- ----- ------ - ----------------------G--- ---------- 314

GT#1   AB067721   368    GTCATCTCAACCCTCGAG CCCCCCCGCCTC GCGG  C GGCGGGG CCCGGCCTT GGGGGTCACGGCC CCGC   GCCG 438
JAAVMX010000019 19713      ------------------ ---------- ---- - ------- --------- ------------- ---- ---- 19782
JAAVMX010000019 32357      ------------------ ---------- ---- - ------- --------- ------------- ---- ---- 32426
JAAVMX010000002 18702404   ------------------ ----- ----- ---- - ------- --------- ------------- ----- ---- 18702473
JAAVMX010000018 1006       ------------------ ----- ----- ---- - ------- --------G--------- ----- --- 1075
JAAVMX010000019 6545       ------------------ ---------- ---- - ------- --------- ----------C--- ---- 6614
JAAVMX010000019 45041      ------------------C----------A ----TA- -CG---C ------TC----AA--- ----T 45123
GT#2   MG770309   223    ------------------ ---------- ---- - ------- --------- ------------- ---- ---- 293
GT#3   HM595984   342    ------------------ ---------- ---- - -A------- --------- -------T-- ---- 412
GT#7   AJ488254   148    ------------------ ---A-A----T--A -T-------- --------- ------------- ---- 76
GT#8   GU246286   315    ------------------ ----- - -A----- ------- --------G----- ---- C--C 387
GT#9   GU246288   316    ------------------ ---------- -G-- - ------- --------- ------------- ---- 388
GT#10  GU246287   314    ----A-A----A------ ---------T -G-- T -------- ----T---C--- --C- C--C 384
GT#11  JQ695935   310    ------------------ ---------- ---- - ------- --TTCT--- --TT---CTTTAT TGAT -A-- 381
GT#12  GU246296   315    ------------------ ---------- ---- - ------- --------- ------------- --C-CGC--- 388

GT#1   AB067721   439    CCCCC  T AAACGC AG       TGGCGACC  CC  GCCGCGGCTCC CC TGCGCAGTAGCT  CG  CTGAGAAC 497
GT#1   AB067721   439    CCCCC  T AAACGC AG       TGGCGACC  CC  GCCGCGGCTCC CC TGCGCAGTAGCT  CG  CTGAGAAC 497
JAAVMX010000019 19783      -----  - ------ --       --------  --  ----------- -- ------------  --  -------- 19842
JAAVMX010000019 32427      -----  - ------ --       --------  --  ----------- -- ------------  --  -------- 32486
JAAVMX010000002 18702474   -----  - ------ --       --------  --  ----------- -- ------------  --  -------- 18702532
JAAVMX010000018 1076       -----  - ---G--G        --------GC--  ----------- -- ------------  --GCT------- 1134
JAAVMX010000019 6614       -----  - ---G--G        --------GC--  ----------- -- ------------  --GCT------- 6678
JAAVMX010000019 45125      CCT--- G - ------ --CCTAAGTG------- -- --GA-------A-- --T-A-------CA-- -------- 45195
GT#2   MG770309   294    -----  - --       --------  --  ----------- -- ------------  --  -------- 301
GT#3   HM595984   413    -----  - ---T-- --       --------  --  ----------- -- ------------  --  -C---G-- 479
GT#7   AJ488254    75    -----  - ------ --       --------  --  ----------- -T -----C-----  G-  -------- 18
GT#8   GU246286   388    ---T-  - ------ --       G-------  --  ----- ---T- A- ----------  -A  -C------ 443
GT#9   GU246288   389    -----  - ------ --       G-------  --  ---------- -- --- -C-G-A--  -A  -C------ 452
GT#10  GU246287   385    ----CC-A------ G-       G--G-C--  --  --T--GC-----TT- AA  -C-G----- 417
GT#11  JQ695935   382    G-TTG  - -CTT-G --       G-C-T-  --GG-T  -TCG- ----GG- -- --TC-GA--  T-  -ATTAGTG 445
GT#12  GU246296   361    -----  - -A------ G-       G-------  --CG--G-------- -- -------G------  -AAC-GA----- 420

GT#1   AB067721   498    CT CGCACCGG  GA GCGCGGA       GGCGG TCACGCC GTG   AA ACCACC   ACACCC TCCA       549
JAAVMX010000019 19843      -- --------  -- -------       ----- ------ - --G    -- -------   ------ ----       19894
JAAVMX010000019 32487      -- --------  -- -------       ----- ------ - --G    -- -------   ------ ----       32537
JAAVMX010000002 18702533   -- --------  -- -------       ----- ------ - --TG   -- -------   ------ ----       18702586
JAAVMX010000018 1135       -- --------  -- -------       ----- ------ - --G    -- -------   ------ ----       1186
JAAVMX010000019 6679       --C-------  -- -------       ----- ------ - --G    -- ------ACA------ ----       6733
JAAVMX010000019 45196      -- --------  -- -------CCCAT-----C------- --GGATG--C------ACC------ ----       45251
GT#2   MG770309   301                                                                                       301
GT#3   HM595984   480    -C --------  A- ------G       --T-- ------- -C-    -- ------  ---    ---       519
GT#7   AJ488254    17    --T------                                                                          9
GT#8   GU246286   444    T- -------A  A- ----- --       A---- -T----- ---    -- -A-T--  --G--T ----       494
GT#9   GU246288   453    -- -------  -- -G-------       ------ -----  -G-    -- ------  --G--- -- --       496
GT#10  GU246287   418    -C ---C----AA--  -G-G---       --G--GG- --    -- ------  ---    ---       478
GT#11  JQ695935   446    -- AA-GT     T- C-- -CC       --TTCGC-GT-AT AAT    T-TCTG  --    TTG-  -G-G       485
GT#12  GU246296   421    -- --------  --G-G-----       ---G--------G--- A-- ------   --G---T----       480
```

Figure 3. Alignment of repetitive ITS copies within genome JAAVMX000000000 of H. sinensis Strain IOZ07 and ITS sequences of GC-biased genotypes of O. sinensis.

*Genome assemblies JAAVMX010000002, JAAVMX010000018, and JAAVMX010000019 were obtained from H. sinensis Strain IOZ07 (Shu et al. 2020). One copy within each of JAAVMX010000002 and JAAVMX010000018, indicated in green, shows 97.4% or 97.0% similarity to AB067721. JAAVMX010000019 contains 4 repetitive ITS copies, including 2 copies in **black** (19404→19894 and 32048→32537), which are 100% identical to AB067721, and 2 other copies in blue (6233→6733 and 44729→45251), with 94.5% and 90.8% similarities to AB067721. "GT" denotes the O. sinensis genotype. AB067721, MG770309, HM595984, AJ488254, GU246286, GU246288, GU246287, JQ695935, and GU246296 are the ITS sequences of GC-biased Genotypes #1–3 and #7–12 of O. sinensis in pink, respectively. The underlined sequence in black represents the 5.8S gene of GC-biased Genotype #1 H. sinensis. Hyphens indicate identical bases, and spaces denote unmatched sequence gaps.*

GC-biased Genotypes #13–14 were characterized by large DNA segment substitutions and recombination of genetic material between the genomes of the 2 parental fungi, Genotype #1 *H. sinensis* and an AB067719-type fungus, and are not included in the sequence analysis in Figure 3.

In addition to the 94.7%–95.3% sequence similarities between the Genotype #1 genomic repetitive ITS copies and Genotype #7 (*cf.* Table 2), the genetic features of Genotype #7, with numerous mutant alleles, are unlike any of the repetitive ITS copies. The study conducted by Chen et al. (2011) did not involve the purification of the *O. sinensis* fungus or fungi within the *C. sinensis* specimen. Thus, there is no evidence to date for determining whether the sequence of Genotype #7 indeed resides in the genome of Genotype #1 *H. sinensis* as a repetitive ITS copy, although Genotype #7 may be phylogenetically closer to Genotype #1 than other GC-biased genotypes.

The Bayesian tree in Figure 2 shows that GC-biased Genotypes #2–3 and #8–14 were placed outside of the Genotype #1 phylogenetic clade with great phylogenetic distances (horizontal bars) and were genetically distinct from the genomic repetitive ITS copies of Genotype #1 *H. sinensis* (*cf.* Figure 3). Notably, phylogenetically different Genotypes #1 and #2 were simultaneously detected in the stromata of the immature, maturing and mature *C. sinensis* collected from Kangding County of Sichuan Province. These 2 genotypes showed completely different developmental patterns in an asynchronous, disproportional manner in the *C. sinensis* stromata during maturation, indicating that the genotypic sequences do not reside in the single *O. sinensis* genome and are therefore not represented as repetitive ITS copies.

III-5. Transcriptional silencing of 5.8S genes

The above genetic, genomic, and phylogenetic analyses demonstrate GC-biased characteristics

of the multiple repetitive ITS copies in the *H. sinensis* genome and genomic independence of all GC- and AT-biased genotypes of *O. sinensis*. Li et al. (2013) identified ITS sequences for both Genotypes #1 and #5 from genomic DNA that was isolated from 8 (1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) of 15 cultures of natural *C. sinensis* multicellular heterokaryotic mono-ascospores. However, the authors reported the identification of 5.8S cDNA only for Genotype #1 and not for Genotype #5 in a cDNA library constructed from total ascosporic RNA extracted from Clone 1220, one of the 8 genetically heterogeneous clones.

Li et al. (2013) did not detect the ITS sequences of AT-biased Genotypes #4, #6, and #15–17 in the genomic DNA isolated from any of the 15 ascosporic clones. Thus, there has been insufficient evidence to conclude that the 5.8S genes of all AT-biased genotypes of *O. sinensis* are permanently dysfunctional pseudogenic components coexisting in "a single genome" with a functional copy of GC-biased Genotype #1. Li et al. (2019) provided the following logical reasoning underlying the dysfunctional pseudogene hypothesis: "If AT-biased ITS sequences are not pseudogenes, 5.8S cDNA should have been detected because rRNAs are critical and essential for protein synthesis in all living organisms", which touches on 2 scientific issues. First, are the 5.8S genes of all genotypes of *O. sinensis* actively transcribed in *C. sinensis* ascospores (discussed in this section)? Second, is it correct to conclude that "5.8S cDNA should have been detected" by the techniques designed and used by Li et al. (2013) (discussed in Section III-6)?

Many research groups have reported differential coexistence of several AT-biased genotypes in various combinations in different compartments of natural *C. sinensis* (Table 4) (Xiao et al. 2009; Zhu et al. 2010, 2012; Gao et al. 2011, 2012; Li et al. 2023b, 2023c). Genotypes #4 and #15 of the AT-biased Cluster B of *O. sinensis* (*cf*. Figure 2) were reported to exist in the stroma and stromal fertile portion (densely covered with numerous ascocarps) of natural *C. sinensis* but not in the ascospores (Li et al. 2023b, 2023c), consistent with what Li et al. (2013) have reported. Thus, there has been no evidence to demonstrate that AT-biased Genotypes #4 and #15 occur in the *C. sinensis* ascospores, and there is no ground for claiming that the 5.8S genes of Genotypes #4 and #15 are dysfunctional pseudogenes in the genome of GC-biased Genotype #1 based on Genotypes #4 and #15 cDNA not being detected in the *C. sinensis* ascosporic cultures.

Table 4*. Differential occurrence of AT-biased genotypes of O. sinensis in the different compartments of natural C. sinensis.*

Genotype # and GenBank accession #	Phylogenetic AT-biased Cluster [§]	Stroma		Stromal Fertile Portion (covering with numerous ascocarps)	Ascospores
		Immature	Mature		
#4 AB067744	B	√	√	√	
#5 AB067740	A	√	√	√	√
#6 EU555436	A	√	√	√	√
#15 KT232017	B	√	√	√	
#16 KT232019	A	√			√
#17 KT232010	A	√			

Note: [§]*, Phylogenetic AT-biased Clusters A and B are shown in Figure 2 (the Bayesian majority rule consensus phylogenetic tree) and are similarly shown in the phylogenetic trees generated by using the Bayesian, maximum likelihood, and minimum evolution algorithms by Stensrud* et al. *(2007), Zhang* et al. *(2013a) and Li* et al. *(2016b, 2022a, 2022c, 2023b, 2023c).*

Xiang et al. (2014) reported a metatranscriptome study of natural *C. sinensis* specimens (unknown maturation stage) collected from Kangding County of Sichuan Province, China. A GenBank BLAST search confirmed that the GAGW00000000 metatranscriptome sequences do not contain any 5.8S gene transcripts of *C. sinensis*-associated fungi (including all genotypes of *O. sinensis*), indicating transcriptional silencing of 5.8S genes of all cocolonized fungi in natural settings.

Xia et al. (2017) reported another metatranscriptome study of mature natural *C. sinensis* specimens collected from Deqin County of Yunnan Province, China. The metatranscriptome assembly was not uploaded to GenBank but instead to another depository server www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas. Similarly, this assembled metatranscriptome contains no 5.8S gene transcripts. However, the authors uploaded the unassembled shotgun reads (SRR5428527) to GenBank. A BLAST search of the SRR5428527 Shotgun Read Archive (SRA) using the MegaBLAST algorithm identified 1894 unassembled metatranscriptome shotgun reads (100 bp each) that were 100% homologous to the 5.8S gene sequences (156 bp) of GC-biased Genotype #1 *H. sinensis*.

Fungal genomes contain multiple repetitive copies of rDNA ITS1-5.8S-ITS2 sequences (*cf.* Tables 1 and 3; Figures 1–3). Because the assembled metatranscriptome contained no 5.8S gene transcripts (Xia et al. 2017) and no segments of the rDNA 18S and 28S gene of *H. sinensis* were

homologous to the 5.8S gene, the 1894 unassembled shotgun reads (SRR5428527) of natural *C.*

sinensis may likely be derived from highly homologous non-rRNA genes that evolved in a

conservative manner within fungal metagenomes. Figure 4 shows the GenBank BLAST graphic

summary of the alignment of the 5.8S gene (the query sequence; segment 218–373 of Genotype #1

H. sinensis AB067721) and the first 75 of the overlapping 1894 metatranscriptome shotgun reads

(100 bp each) of mature natural *C. sinensis*.

Figure 4. Graphic alignment summary of the query sequence (in blue–green), the 5.8S gene (218–373

segment of Genotype #1 H. sinensis AB067721) and the first 75 in the list of 1894 overlapping shotgun

reads (pink bars; GenBank SRR5428527) of mature natural C. sinensis.

Various degrees of similarity were found when the 1894 unassembled shotgun reads were aligned

with the 5.8S gene sequences of the 17 genotypes of *O. sinensis*. Table 5 shows 6 examples of shotgun

reads of SRX2718557 aligned with the 5.8S sequences of *O. sinensis* genotypes, *Paecilomyces*

hepiali, *Tolypocladium sinensis* (Group D), and an AB067719-type Group-E fungus (Stensrud et al.

2007). The high homology between the shotgun reads and the various 5.8S gene sequences of *O.*

sinensis genotypes may not conclusively identify their fungal sources in natural *C. sinensis*, which

contains >90 species spanning at least 37 fungal genera (Zhang et al. 2010, 2018; Xiang et al. 2014;

Xia et al. 2015, 2017).

Table 5. *Percent similarity between the unassembled metatranscriptome shotgun reads (SRA:*
SRR5428527) of natural C. sinensis and the 5.8S genes of multiple genotypes of O. sinensis and P.
hepiali, T. sinensis *(Group D) and an AB067719-type fungus (Group-E).*

Geno-type	GenBank accession #	% Similarity between 5.8S genes and *C. sinensis* shotgun sequences					
		SRR5428527.17546327.1	SRR5428527.17554054.1	SRR5428527.17565575.2	SRR5428527.17679995.2	SRR5428527.17911498.2	SRR5428527.17929569.2
#1	AB067721	**100%**	**100%**	**100%**	**100%**	**100%**	**100%**
#2	MG770309	96%	96%	**100%**	96%	**98%**	96%
#3	HM595984	**100%**	**99%**	**99%**	**100%**	**99%**	**100%**
#7	AJ488254	**98%**	**98%**	**99%**	**98%**	**98%**	**98%**
#8	GU246286	94%	96%	95%	94%	**97%**	94%
#9	GU246288	**99%**	**99%**	**99%**	**100%**	**99%**	**100%**
#10	GU246287	96%	94%	93%	95%	93%	**97%**
#11	JQ695935	**100%**	**100%**	**100%**	**100%**	**100%**	**100%**
#12	GU246296	**99%**	**99%**	**99%**	**99%**	**99%**	**99%**
#13	KT339190	**98%**	93%	93%	**98%**	93%	**98%**
#14	KT339178	**98%**	93%	93%	**98%**	93%	**98%**
#4	AB067744	86%	87%	87%	83%	87%	83%
#5	AB067740	85%	87%	87%	84%	87%	84%
#6	KJ720572	85%	85%	87%	84%	86%	84%
#15	KT232017	85%	88%	88%	84%	88%	84%
#16	KT232019	86%	86%	87%	85%	87%	85%
#17	KT232010	87%	88%	88%	87%	88%	87%
P.h.	EF555097	**99%**	**98%**	**97%**	**99%**	**98%**	**99%**

T.s.	AB067720	**98%**	**97%**	**96%**	**98%**	**97%**	**98%**
	AB067719	**98%**	93%	93%	**98%**	93%	**98%**

Note: GC-biased genotypes are listed in blue*, and AT-biased genotypes are listed in* red*. GenBank sequence ID SRA: SRR5428527 fragments (17546327.1, 17554054.1, 17565575.2, 17679995.2, 17911498.2, and 17929569.2) are the unassembled transcriptome shotgun sequences obtained from the mature natural C. sinensis specimens collected from Deqin County of Yunnan Province by Xia* et al. *(2017). P.h. refers to P. hepiali; T.s. refers to Group-D* T. sinensis*, and AB067719 refers to an AB067719-type Group-E fungus (Stensrud* et al. *2007). Highly homologous percent similarities (97%–100%) are in bold* **green** *font.*

As shown in Table 5, the 5.8S gene sequence of Genotype #1 *H. sinensis* showed 98%–99% homology to the 5.8S gene sequence of *P. hepiali*. A GenBank BLAST search revealed the alignment of the 5.8S gene sequence (segment 218→373 of AB067721) of *H. sinensis* to 56–438 unassembled transcriptome shotgun sequence reads, which were identified in 5 RNA-Seq experimental data files (GenBank assessment #SRX3083435–SRX3083439) of *P. hepiali* Strain Feng (SRA Study: SRP115168; Pang et al. 2018) (Table 6). Pang et al. (2018) uploaded the 5 transcriptomic experimental datasets to GenBank for the total RNA samples extracted from the cultures of *P. hepiali* Strain Feng under different culture conditions and durations.

Table 6. *Summary of BLAST searches of the transcriptome shotgun reads of P. hepiali Strain Feng against the 5.8S gene sequence (218→373 of AB067721) of Genotype #1 H. sinensis.*

P. hepiali sample	Culture duration (days)	Additive in culture medium	GenBank transcriptome data file	*vs.* the sequence of the *H. sinensis* 5.8S gene		
				# Of the total BLAST hits	(% Of total filtered reads)	% Similarity
P4	4	no	SRX3083438	219	$(3.7\% \times 10^{-4})$	93%–99%
P9	9	no	SRX3083437	56	$(3.6\% \times 10^{-4})$	95%–100%
P15	15	no	SRX3083436	438	$(6.12\% \times 10^{-4})$	96%–100%
Ps	4	Adenosine	SRX3083435	356	$(4.13\% \times 10^{-4})$	96%–100%
Pa	4	Adenine	SRX3083439	207	$(6.12\% \times 10^{-4})$	96%–99%

The extremely high similarity between the 5.8S gene sequences of *H. sinensis*, *P. hepiali* and *T. sinensis,* as shown in Table 5, indicates conservative evolution of the 5.8S gene in many fungal species. Table 6 shows significant differences in the number of BLAST hits and percent similarities of the 5.8S gene sequence of *H. sinensis* aligned with the transcriptome shotgun reads obtained from the culture experiments of *P. hepiali* Strain Feng under different culture conditions and durations

(Pang et al. 2018). The results indicate the experimental switching on-or-off of *P. hepiali* genes,
including the 5.8S gene and many other non-rRNA genes, in response to the different experimental
settings.

Liu et al. (2015) reported the mRNA transcriptome assembly GCQL00000000 derived from *H.*
sinensis Strain L0106, which contained no transcript of rRNA or tRNA. Continuous *in vitro* liquid
fermentation performed in this study induced a nonlinear reduction in the total number of
transcriptomic unigenes, first from 25,511 (after 3 days of fermentation) to 25,214 (after 6 days of
fermentation) and then more sharply to 16,245 (after 9 days of fermentation). Table 7 summarizes the
dynamic alterations in the total number of transcriptomic unigenes, average length and GC content
of transcripts of *H. sinensis* Strain L0106, indicating the nonnatural switching on-or-off and
differential transcriptional activation and silencing of numerous *H. sinensis* genes in response to
continued liquid fermentation of *H. sinensis* Strain L0106. As shown in Table 7, more dramatic
changes might conceivably occur after 9 days of fermentation. The prolonged 25-day liquid
incubation period adopted by Li et al. (2013) might have a greater impact on the unbalanced
transcriptional activation (switching on) and silencing (switching off) of long and short fungal genes
with various GC content biases, in addition to the changes in the culture medium as an important
factor influencing the transcriptional activation and silencing of many fungal genes (*cf*. Table 6).

Table 7. *Dynamic alterations of transcripts in the mRNA transcriptome of the H. sinensis Strain*
L0106 during continuous in vitro fermentation.

Transcriptomic unigene	After 3 days of fermentation	After 6 days of fermentation (% change)	After 9 days of fermentation (% change)
Total number	25,511	25,214 (-1.2%)	16,245 (-36.3%)
Average length (nt)	681	682 (+0.1%)	994 (+46.0%)
GC content (%)	58.2%	57.9% (-0.5%)	57.0% (-2.1%)

III-6. 5.8S cDNA detected using 5.8S-F/R primers

In addition to the transcriptional silencing of 5.8S genes in natural settings and the significant
impact of continuous *in vitro* culture and altered experimental settings on unbalanced gene
transcription in experimental settings, as discussed above, the second issue related to the logic
described by Li et al. (2019) is whether the methods designed by Li et al. (2013) were suitable for
scientifically identifying 5.8S gene cDNAs of multiple *O. sinensis* genotypes (Li et al. 2020a).

For instance, Li et al. (2019) retrospectively explained the research design logic of 5.8S-F/R primers by Li et al. (2013): "The 5.8S-F/R primers … located in the most conserved region." No doubt such a design would largely reduce the primers' specificity for 5.8S cDNA because of the conservative evolution of the 5.8S gene and many other fungal genes. Table 8 and Figure 5 show similarities between the sequences of primers 5.8S-F (5' ACTTTTAACAACGGATCTCTT 3') and 5.8S-R (5' AAGATAACGCTCGGATAAGCAT 3') and the 5.8S gene of Genotypes #1, #4, and #5.

Table 8. *Percent similarity of the 5.8S-F/R primers and the 5.8S gene sequences of Genotypes #1, #4 and #5 of O. sinensis.*

Genotype # and GenBank accession #	% Similarity of the primer *vs*. the 5.8S gene sequence	
	Primer 5.8S-F	Primer 5.8S-R[§]
#1 AB067721	95% (20/21 nt)	85% (17/20 nt)
#4 AB067744	95% (20/21 nt)	90% (18/20 nt)
#5 AB067740	100% (21/21 nt)	95% (19/20 nt)

Note: [§], Primer 5.8S-R has 2 flanking residues downstream in the ITS2 region.

The primers 5.8S-F/R are 100% and 95% homologous to the 5.8S gene sequence of AT-biased Genotype #5, as shown in Table 8. This indicates that in contrast to the 95% and 85% similarities to the 5.8S gene sequence of GC-biased Genotype #1, the 5.8S-F/R primers may be favored for amplifying the 5.8S cDNA sequence of Genotype #5, rather than of Genotype #1. The primers showed variable similarity, ranging from 61.9%–100% and 31.8%–95.5%, respectively, among all 17 genotypes of *O. sinensis*, preferentially amplifying the most easily amplified cDNA sequence in the cDNA library considering the primary structure and secondary conformation of PCR templates and consequently causing false-negative results for other 5.8S cDNA sequences.

Section III-5 describes the 100% homologies between the 1894 unassembled metatranscriptome shotgun reads (100 bp) of mature natural *C. sinensis* and the 5.8S gene sequence (156 bp) of GC-biased Genotype #1 *H. sinensis*, including many non-rRNA transcripts among the highly homologous 1894 shotgun reads in natural settings. A BLAST search of *O. sinensis* transcriptomic sequences (taxid:72228) in the GenBank database showed that the 5.8S-F/R primers were 100% homologous to the mRNA transcriptome assemblies GCQL01001191, GCQL01006777, GCQL01007818, and GCQL01013443 of Genotype #1 *H. sinensis* Strain L0106 (Liu et al. 2015), which contain no rRNA or tRNA transcripts, and to assembled metatranscriptome sequences GAGW01003691, GAGW01007670, and GAGW01007671 of natural *C. sinensis* (Xiang et al. 2014). Figure 6 shows

that metatranscriptome GAGW01003691 is 99.8% homologous to a segment of mRNA transcriptome

GCQL01006777 of *H. sinensis* Strain L0106; metatranscriptomes GAGW01007670 and

GAGW01007671 are 100% identical to mRNA transcriptome segments of GCQL01007818 and

GCQL01001191, respectively (Li et al. 2020a). Because mRNA transcriptome assembly

GCQL00000000 contains no rRNA or tRNA transcripts, the extremely high homology between the

metatranscriptome and transcriptome sequences indicates that the 5.8S-F/R primers designed by Li

et al. (2013) may be able to recognize and nonspecifically amplify non-rRNA components in the

cDNA library derived from the genetically heterogeneous Clone 1220, one of the 8 heterogenous

clones (1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) of the *C. sinensis* mono-ascospore

cultures obtained by Li et al. (2013).

```
GT1 AB067721 1    TAGAGGAAGTAAAAGTCGTAACAAGGTCTCCGTTGGTGAACCAGCGGAGGGATCATTATC  60
GT4 AB067744 1    ----------------------------A--A---T-----------T------        56
GT5 AB067740 1    --A----------------T---------A--A---T-----------T------        60

GT1 AB067721 61   GAGTCACCACTCCCAAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCC  120
GT4 AB067744 57   -----------C-------------A--C-------------------------------  116
GT5 AB067740 61   ----T--T----TT---------ATA--T--T-T-----------TA------------T  120

GT1 AB067721 121  GGCGCCCCAGGGCCCGGACCAGGGCGCCCGCCGGAGGACCCCCAGACCCTCCTGTCGCAG  180
GT4 AB067744 117  -------T-----------T-------------------TT---------A---T--     176
GT5 AB067740 121  A------T-A---------T-AA------------------TT---------A-TAT--   180

GT1 AB067721 181  TGGCATCTCTCAGTCAAGAAGCAAGCAAATGAATCAAAACTTTCAACAACGGATCTCTTG  240
GT4 AB067744 177  -A---------T---T--------------A---T-A-ACTTTTAATAACGGATCTCTTA  236
GT5 AB067740 181  -A-T------T---T--------------A---T---ACTTTTAACAACGGATCTCTTA  240
Primer 5.8S-F                                     ACTTTTAACAACGGATCTCTT

GT1 AB067721 241  GTTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCA  300
GT4 AB067744 237  ---T-A----TA--A---------------A------------A-----T------T-   296
GT5 AB067740 241  -----A----TA--A--A------------A---------A-A----A-------T-   300

GT1 AB067721 301  GTGAACCATCGAATCTTTGAACGCACATTGCGCCCGCCAGCACTCTGGCGGGCATGCCTG  360
GT4 AB067744 297  --A---T-----------A----T-T-C-A-------T--------A------ATGCCTG  356
GT5 AB067740 301  --A---T-----------A------T---A-------T--------A------ATGCCTA  360
Primer 5.8S-R                                                ATGCTTA

GT1 AB067721 361  TCCGAGCGTCATCTCAACCCTCGAGCCCCCCGCCTCGCGGCGGCGGGGCCCGGCCTTGGG  420
GT4 AB067744 357  TCCGAGCGTTACCTT------TA----------------------------------A--  416
GT5 AB067740 361  TCCGAGCGTTATCTT-----------------------------T-------A--  420
Primer 5.8S-R     TCCGAGCGTTATCTT

GT1 AB067721 421  GGTCACGGCCCCGCGCCGCCCCCTAAACGCAGTGGCGACCCCGCCGCGGCTCCCCTGCGC  480
GT4 AB067744 417  -----------A-------------------A-T-------------------A---   476
GT5 AB067740 421  ---T----------------T------T---A--------------------A--T   480

GT1 AB067721 481  AGTAGCTCGCTGAGAACCTCGCACCGGGAGCGCGGAGGCGGTCACGCCGTGAAACCACCA  540
GT4 AB067744 477  -A-----TA--A-----------TA----------------------A-------T-   536
GT5 AB067740 481  G-------A-C--------------A-----------------A-------T-   540

GT1 AB067721 541  CACCCTCCAGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACTTAAGCATATCAATAA  600
GT4 AB067744 537  T------TG---A---------T-----------------A----------------   596
GT5 AB067740 541  -G--T-TT----A---------T----------------A------------T-----  600

GT1 AB067721 601  GCGGAGGAAAAGAAACCAACAGGGATTGCCCCAGTAACGGCGAGTGAAGCGGCAACAGCT  660
GT4 AB067744 597  --------                                                604
GT5 AB067740 601  ----------------T--T----------TT------------A------T--T----  660
```

Figure 5. ITS sequence alignment of AB067721 of GC-biased Genotype #1, AB067744 and AB067740 of AT-biased Genotypes #4–5 of O. sinensis and the 5.8S-F/R primers.

"GT1", "GT4", and "GT5" represent Genotypes #1 and #4–5 of O. sinensis, respectively. The sequences in blue are 18S and 28S genes (5' and 3' end regions, respectively). Those in pink represent sequences of 5.8S genes. Those in black are ITS1 between the 18S and 5.8S genes and ITS2 between the 5.8S and 28S genes. The sequences underlined in green are for primers 5.8S-F/R, while primer 5.8S-R was converted to the sense chain sequence. Hyphens indicate identical bases.

Unfortunately, Li et al. (2013) did not disclose the sequences of the identified 5.8S cDNA but

instead reported that "Of these clones, 28 showed identical sequences (104 bp excluding the primers)

to the most frequent sequence in group A (JQ900148). The remaining 14 sequences … had single

substitutions at various positions", where group A = Genotype #1 *H. sinensis*. GenBank annotates the

location of the 5.8S gene of JQ900148 as nucleotides 192→346, covering 155 nt, which is not a

multiple of three and is not in accord with the open reading frame of the gene.

Alternatively, the 5.8S genes were annotated with nucleotide positions 218→373 in AB067721

of Genotype #1 and AB067740 of Genotype #5 and 214→369 in AB067744 of Genotype #4, covering

156 nt and 52 codons in the 3 cases (Zhu & Li 2017; Li et al. 2020a, 2022c). Excluding the 5.8S-F/R

primer regions (*cf*. Figure 5), the remaining sequence of the 5.8S gene consists of 114 bp, with 10

additional nucleotides (mismatches) beyond the 104 bp of all cDNA clones in the 28 clones reported

by Li et al. (2013). Adding "single substitutions at various positions" in 14 other clones, as reported

by Li et al. (2013), the 42 clones presumably contained either 10 or 11 mismatches (8.8%–9.6%) in

the 114-bp segment of the 5.8S gene cDNA, corresponding to 90.4%–91.2% sequence similarity to

the 5.8S gene sequence of GC-biased Genotype #1 *H. sinensis*.

On the other hand, Li et al. (2013) did not provide any information about to the sequence

alignment of the detected 5.8S-F/R PCR amplicons with the 5.8S gene sequence of AT-biased

Genotype #5 of *O. sinensis*.

IV. Discussion

IV-1. The repetitive ITS copies in the genome of GC-biased Genotype #1 *H. sinensis* are unrelated to RIP mutagenesis

Selker and his colleague (1985, 1990, 1997, 2002) reported frequent C-to-T (cytosine→thymine)

transitions and concomitant epigenetic methylation of nearly all remaining cytosine residues in fungal

5S RNA genes associated with transcriptional silencing. Gladyshev (2017) summarized that RIP

"occurs specifically in haploid parental nuclei that continue to divide by mitosis in preparation for

karyogamy and ensuing meiosis;" and "mutates cytosines on both strands of each DNA duplex in a

pairwise fashion (e.g., preferentially affecting none or both repeated sequences)." Hane et al. (2017)

stated that "the RIP process selectively mutated duplicated sequences in both DNA strands by

inducing single-nucleotide point (SNP) mutations that converted C:G base pairs to T:A. This often

led to the introduction of nonsense or missense mutations which affected the expression of these

sequences." The sequence-specific targets of RIP mutations for de novo DNA methylation are under

debate (Hood et al. 2005; Taranto et al. 2014; Hane et al. 2017).

Table 3 and Figure 1 demonstrate that the variable repetitive ITS copies were genetically

characterized by multiple scattered insertion, deletion and transversion point mutations; these

mutations might not be generated through RIP mutagenesis, which theoretically causes

cytosine→threonine (C-to-T) transitions. On the other hand, the AT-biased genotypes of *O. sinensis*

characterized by multiple scattered transition point mutations were generated in parallel with other

GC-biased genotypes, through accelerated nuclear rDNA evolution from a common ancestor during

a long evolutionary history (Stensrud et al. 2007), regardless of whether this occurred *via* RIP

mutagenesis or other evolutionary mechanisms. Thus, Li et al. (2020b) reported that RIP mutagenesis

induces the formation of mutant AT-biased repetitive ITS copies of *O. sinensis* within the single

genome of GC-biased Genotype #1 *H. sinensis* lacking supporting evidence. Even if it were to be

proven that AT-biased genotypes were indeed induced by RIP mutagenesis during a long history of

evolution, there has been no evidence that AT-biased genotypes are genomic components of the GC-

biased Genotype #1 genome that emerged after a new GC-biased Genotype #1 "ITS haplotype was generated" as "ITS pseudogenes … in a single genome", as assumed by Li et al. (2013, 2019, 2020b).

Taranto et al. (2014) and Hane et al. (2015) reported that RIP "is observed only in certain fungal taxa; the Pezizomycotina (filamentous Ascomycota) and some species of the Basidiomycota." Tables 1 and 3 and Figures 1–3 show that all the multiple genomic repetitive ITS copies were genetically GC-biased, with no or only a few transition alleles, and were phylogenetically distinct from AT-biased genotypes, which possess numerous scattered transition alleles. It is then reasonable to question whether Genotype #1 *H. sinensis* is susceptible to epigenetic methylation attack through RIP mutagenesis.

IV-2. Genetic and phylogenetic differences of AT-biased genotypes of *O. sinensis* and GC-biased repetitive ITS copies in the genome of Genotype #1 *H. sinensis*

The sequences of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* shown in red in Figure 1 contain numerous scattered transition alleles. They are genetically distinct and phylogenetically distant from GC-biased repetitive ITS copies containing the predominant insertion, deletion, and transversion point mutations within the genome of GC-biased Genotype #1 *H. sinensis* (*cf*. Figures 1–2 and Tables 1–3). The sequences of AT-biased genotypes, as well as GC-biased Genotypes #2–3 and #7–14, of *O. sinensis* are absent in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000 of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively (*cf*. Tables 1–2 and Figures 1 and 3) but instead belong to the genomes of independent *O. sinensis* fungi (Stensrud et al. 2007; Xiao et al. 2009; Hu et al. 2013; Zhu and Wu 2015; Li et al. 2016a, 2016b; 2020a, 2022a, 2022c, 2023c; Zhu and Li 2017; Zhu et al. 2018, 2019; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). The nonresidency measure of AT-biased *O. sinensis* genotypes does not support the hypothesis that the AT-biased sequences are "ITS pseudogene" components of "a single genome" of GC-biased Genotype #1 *H. sinensis*. Stensrud et al. (2007) concluded that the variation in the 5.8S gene sequences of AT-biased genotypes "far exceeds what is normally observed in fungi … even at higher taxonomic levels (genera and family)". Xiao et al. (2009) concluded that the sequences of AT-biased genotypes likely belong to independent *O. sinensis* fungi.

IV-3. The previously proposed *O. sinensis* center of origin and the geography of *O. sinensis* evolution

GC-biased genotypes, including the GC-biased genomic repetitive ITS copies, might have evolved at accelerated rates in parallel with AT-biased genotypes from an ancient common ancestor during long-term evolution, as assumed by Stensrud et al. (2007). Zhang et al. (2009) detected GC-biased Genotype #3 in natural *C. sinensis* specimens collected from the Nyingchi District of Tibet and hypothesized that "Nyingchi might have acted as a center of origin of *O. sinensis* and *O. sinensis* might have been transmitted from Nyingchi to other locations". Accordingly, GC-biased Genotype #3 might represent a modern immediate ancestor of Genotype #1 *H. sinensis* that has been consistently identified in other production locations as the result of a relatively modern phase during long-term evolution. The authors described an evolution path from GC-biased Genotype #3 to Genotype #1 and transmission of *O. sinensis* from the "center of origin … spread eastward and westward first among southern regions on the Tibetan Plateau" and then "spread from south to north … at a later time".

Unfortunately, this evolutionary geography for *O. sinensis* is missing an evolutionary branch or branches related to AT-biased genotypes of *O. sinensis*. Perhaps the evolution of AT-biased genotypes of *O. sinensis* might have occurred in more ancient historical evolutionary phases, if not in modern evolutionary phases. Undoubtedly, the discovery of AT-biased genotypes of *O. sinensis* will trigger further research on the history of *O. sinensis* evolution and fungal-insect co-evolution [Zhang et al. 2014].

IV-4. Multicellular heterokaryons of natural *C. sinensis* hyphae and ascospores

Bushley et al. (2013) reported that *C. sinensis* ascospores are multicellular heterokaryons with mono-, bi-, and tri-nucleated cellular structures. Zhang and Zhang (2015) wondered whether the bi-nucleated cells (also tri-nucleated cells) of *C. sinensis* hyphae and ascospores contain homogenous or heterogeneous genetic materials, meaning that these polykaryotic cells may contain heterogeneous sets of chromosomes and genomes. Li et al. (2023b, 2023c) confirmed the genetic heterogeneity in the multicellular heterokaryotic ascospores of natural *C. sinensis*.

Li et al. (2013) identified heterogeneous ITS sequences of both GC-biased Genotype #1 and AT-biased Genotype #5 in 8 of 15 clones (1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) and simultaneously identified 7 other clones (1207, 1218, 1219, 1221, 1222, 1225, and 1229) that contained only homogenous sequences of Genotype #1 after 25 days of inoculation with natural *C. sinensis* mono-ascospores. These culture-dependent molecular mycological findings supported the morphological observations of multicellular heterokaryons discovered by their collaborators

(Bushley et al. 2013). Apparently, the 2 groups of the reported ascosporic clones might be derived from different cells of the multicellular heterokaryotic ascospores of natural *C. sinensis*. In addition to the information for the 15 clones of the 2 groups, the authors did not disclose any information of other possible clones, namely, clones 1210, 1211, 1212, 1213, 1215, 1216, 1217, 1223, and 1226, for which the clone serial numbers are missing among those of the 2 groups (*cf*. Table S1 of Li et al. 2013), nor did the authors state whether these 9 possible clones were simultaneously obtained in the same study, why they were not published along with the 15 reported clones, and whether they contained different genetic materials, although these questions should have been asked during the journal's peer review process. Li et al. (2023b, 2023c), from a different research group, used culture-independent techniques and reported the cooccurrence of GC-biased Genotypes #1 and #13–14, AT-biased Genotypes #5–6 and #16 of *O. sinensis*, *P. hepiali*, and an AB067719-type fungus in *C. sinensis* ascospores, probably indicating that many of these fungi may not be culturable in the *in vitro* experimental setting set up by Li et al. (2013).

IV-5. Genomic variations of *H. sinensis* strains

By aggregating the information from molecular and genomic studies (Hu et al. 2013; Li et al. 2013, 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020), different genomic DNA samples of "pure" *H. sinensis* strains were examined and published and can be grouped as follows.

(1) Group 1 of *H. sinensis* strains consisted of pure, homokaryotic anamorphic *H. sinensis* strains (1229, CC1406-203, Co18, IOZ07, and ZJB12195) (Hu et al. 2013; Li et al. 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). Total genomic DNA was isolated from these strains and subjected to genome-wide sequencing. The genomes contained no AT-biased genotype sequences.

(2) Group 2 of *H. sinensis* strains consisted of 7 clones (Strains 1207, 1218, 1219, 1221, 1222, 1225, and 1229) among 15 clones derived from 25 days of incubation of natural *C. sinensis* mono-ascospores (Li et al. 2013). Total genomic DNA was isolated from these strains and shown to contain the homogenous ITS sequence of Genotype #1 *H. sinensis* but included no AT-biased sequences of *O. sinensis* genotypes.

(3) Group 3 of "*H. sinensis*" strains consisted of 8 other clones (Strains 1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) obtained from the incubation of *C. sinensis* mono-ascospores for 25 days (Li et al. 2013). Total genomic DNA was isolated from these strains and showed genetic heterogeneity, with the coexistence of GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis*.

The *H. sinensis* strains of Groups 1 and 2 were similar, with each strain possessing a homogenous
GC-biased Genotype #1 genome, although multiple GC-biased repetitive ITS copies were identified
in genome assemblies JAAVMX000000000 and NGJJ00000000, for *H. sinensis* Strains IOZ07 and
CC1406-203, respectively (*cf.* Tables 1 and 3 and Figures 1–2) (Hu et al. 2013; Li et al. 2013, 2016a;
Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). Although a single copy of ITS sequences was
identified in genome assemblies ANOV00000000, LKHE00000000, and LWBQ00000000 of *H.*
sinensis Strains Co18, 1229, and ZJB12195, respectively (Hu et al. 2013; Li et al. 2016a; Jin et al.
2020), the genomes of these strains may theoretically contain more repetitive ITS copies that could
have been discarded during the assembly of genome shotgun reads. Conceivably, reassembling the
genome shotgun reads, if technically possible, may very likely identify additional repetitive ITS
copies in the genomes of Group 1 Strains Co18, 1229, and ZJB12195, among which Strain 1229 and
its genome assembly LKHE00000000 are also a Group-2 strain derived from mono-ascospores by Li
et al. (2013, 2016a). Similarly, the genome sequencing of Group-2 strains may very likely identify
multiple repetitive ITS copies.

The *H. sinensis* strains in Groups 2 and 3 were derived after 25 days of liquid culture of the same
C. sinensis mono-ascospores (Li et al. 2013); however, they differed genetically. The Group-3 strains
may likely have multicellular heterokaryotic mycelial structures, similar to the mycelia derived from
microcycle conidiation of the *H. sinensis* ascospores initially cultured on the solid medium plate for
30 days and subsequently in liquid medium for 10–53 days, as shown in Figures 2–3 of (Xiao et al.
2011). Li et al. (2013) reported that the Group 3 strains included cocultured GC-biased Genotype #1
and AT-biased Genotype #5 of *O. sinensis* fungi, which were genomically independent and coexisted
in heterokaryotic mycelial cells, indicating either diploid/polyploid mycelial cells containing
heterogeneous sets of chromosomes/genomes, or impure mycelia, each containing a different
chromosome/genome. However, the authors did not identify any haploid clones containing only AT-
biased Genotype #5, indicating that this AT-biased *O. sinensis* fungus may not exhibit independent
culturability under the *in vitro* experimental settings, although other studies have reported the
identification of single AT-biased Genotype #4 or #5 of *O. sinensis*, without coculture with GC-biased
Genotype #1, in cultures of natural and cultivated *C. sinensis* under different experimental settings
(Mao et al. 2013; Wei et al. 2016). Unfortunately, Li et al. (2013, 2019, 2020b) neglected the
differences in genetic material between the strains in Group 2 and Group 3, which were presumably
derived from the same or different cells of multicellular heterokaryotic ascospores (Bushley et al.

2013; Zhang and Zhang 2015; Li et al. 2020a, 2022c, 2023c).

In addition to the genome sequencing of the homogenous Group-2 Strain 1229 and the generation

of genome assembly LKHE00000000 by Li et al. (2016a), the genome sequencing of one of the

heterogeneous Group-3 strains will provide critical genomic evidence to validate the hypothesis that

AT-biased Genotype #5 of *O. sinensis* coexists as an "ITS pseudogene … in a single genome" of GC-

biased Genotype #1. Unfortunately, the authors elected not to do so since 2012 after obtaining the 8

Group-3 strains, or perhaps failed in such an endeavor due to the technical difficulties in the assembly

of shotgun reads because of the heterogeneous sequences from more than one set of genomes of

independent *O. sinensis* fungi having close evolutionary relations. Consequently, they elected

willingly or unwillingly to pass on the opportunity to experimentally confirm their "ITS pseudogene"

hypothesis that AT-biased Genotype #5 and GC-biased Genotype #1 physically coexist "in a single

genome". Studies from other research groups have confirmed that the sequences of all AT-biased

genotypes belong to the genomes of independent fungi (Stensrud et al. 2007; Xiao et al. 2009; Zhu et

al. 2010, 2012; Yao et al. 2011; Li et al. 2016, 2020a, 2022a, 2022c, 2023c) and that heterokaryotic

ascospores and hyphae with multiple mono-, bi-, and tri-nucleated cells contain heterogeneous sets

of chromosomes and genomes (Bushley et al. 2013; Zhang and Zhang 2015).

Of course, in addition to the 3 groups summarized above, many other *H. sinensis* strains have

been studied and presumably belong to Groups 1–3 or other groups. For instance, Li et al. (2016c)

isolated 2 *H. sinensis* strains, CH1 and CH2, from the intestine of healthy larvae of *Hepialus lagii*

Yan. These strains showed *H. sinensis*-like morphological and growth characteristics but contained

GC-biased Genotype #1, AT-biased Genotypes #4–5 and *P. hepiali*. The strains greatly enhanced the

inoculation potency of *H. sinensis* in the larvae (n=100 each experimental group) of *Hepialus*

armoricanus by 15- to 39-fold (*P*<0.001). Additionally, as mentioned above, Li et al. (2013) did not

discuss whether 9 other ascosporic clones (1210, 1211, 1212, 1213, 1215, 1216, 1217, 1223, and

1226) simultaneously obtained in the study constituted additional fungal group(s) containing different

genetic materials.

IV-6. Genomic independence of AT-biased genotypes of *O. sinensis*

IV-6-1 Differential occurrence of AT-biased genotypes in different compartments of natural *C. sinensis*

759   A BLAST search of the GenBank database revealed at least 668 ITS sequences of *O. sinensis*

(GenBank taxid:72228), 228 of which belong to AT-biased Genotypes #4–6 and #15–17 (Li et al. 2016b, 2022c, 2023c; Zhu et al. 2018, 2019; Zhu & Li 2017). Stensrud et al. (2007) and Zhang et al. (2013a) analyzed the phylogeny of 71 and 397 ITS sequences of *O. sinensis*, respectively, and revealed 3 phylogenetic clades, *i.e.*, Groups A–C (GC-biased Genotype #1 and AT-biased Genotypes 4–5), which consisted of correct *O. sinensis* sequences. Notably, the Genotype #1 clade (Group A) in the phylogenetic tree constructed by Zhang et al. (2013a) using the minimum evolution algorithm consisted of the ITS sequences of Genotype #1 and some of the other GC-biased genotypes of *O. sinensis*. Additional GC- and AT-biased *O. sinensis* genotypes have been successively discovered thereafter (*cf*. Table 2) (Xiao et al. 2009; Zhang et al. 2009; Zhu et al. 2010, 2012; Chen et al. 2011; Zhu and Wu 2015; Li et al. 2013, 2016, 2022a, 2022c, 2023c; Zhu and Li 2017).

Many early studies conducted since 1999 used a single pair of "universal" primers (*e.g.*, *ITS1–ITS5*) for PCR and identified one of the *O. sinensis* genotypes in natural *C. sinensis* specimens collected from different production areas or cultivated *C. sinensis* (Engh 1999; Zhao et al. 1999; Kinjo and Zang 2001; Liu et al. 2001; Stensrud et al. 2005; Wei et al. 2006, 2016; Chen et al. 2011; Mao et al. 2013). Importantly, Wei et al. (2016) reported an industrial artificial cultivation project and identified a single teleomorphic AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis* and a single teleomorphic GC-biased Genotype #1 in natural *C. sinensis*, indicating the existence of at least 2 *O. sinensis* teleomorphs. Unfortunately, this study revealed a species contradiction between the anamorphic inoculants (reported to be 3 GC-biased Genotype #1 *H. sinensis* strains) and teleomorphic AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis*, inconsistent with the hypothesis of a sole anamorph of *O. sinensis* for *H. sinensis* proposed by the same key authors 10 years ago (Wei et al. 2006).

Other studies optimized and combined the use of a variety of molecular techniques, such as several pairs of genotype-specific primers, touchdown PCR, nested PCR, restriction fragment length polymorphism (RFLP), cloning-based amplicon sequencing, and single-nucleotide polymorphism (SNP) mass spectrometry genotyping techniques, etc. These research approaches greatly improved the specificity and efficiency of detection and facilitated the identification of differential coexistence and dynamic alterations of AT-biased genotypes as well as GC-biased genotypes of *O. sinensis* in different combinations in the caterpillar body, stroma, stromal fertile portion with numerous ascocarps, and ascospores of natural *C. sinensis* during maturation (*cf*. Table 4) (Xiao et al. 2009; Zhu et al. 2010, 2012, 2018, 2019; Gao et al. 2011, 2012; Yao et al. 2011; Li et al. 2020a, 2022a, 2023b,

2023c).

Xiao et al. (2009) used pairs of genotype-specific primers for the first time and adopted a nested

PCR approach to amplify mutant sequences of *O. sinensis* from the stroma and caterpillar body of

immature and mature *C. sinensis* specimens collected from geographically remote production areas.

The authors reported the consistent occurrence of GC-biased Genotype #1 and *P. hepiali* and the

inconsistent occurrence of AT-biased Genotypes #4–5 of *O. sinensis*. The authors also detected the

first sequence (EU555436) of AT-biased Genotype #6 and uploaded it to GenBank. The authors then

concluded that the AT-biased genotypes likely belonged to independent fungal species.

Li et al. (2023b, 2023c) used several pairs of genotype-specific primers, touch-down PCR,

cloning-based PCR amplicon sequencing and MassARRAY SNP mass spectrometry genotyping

techniques and identified Genotypes #5–6 and #16 of AT-biased Cluster-A (*cf*. Figure 2) in the

ascospores of natural *C. sinensis*, co-occurring with GC-biased Genotype #1, *P. hepiali*, and an

AB067719-type fungus. The sequences of Genotypes #4 and #15 of AT-biased Cluster-B were

identified in the stromal fertile portion that was densely covered with numerous ascocarps but not in

ascospores obtained from the same *C. sinensis* specimens (*cf*. Table 4). Li et al. (2023c) also identified

GC-biased Genotype #13 in semiejected ascospores and GC-biased Genotype #14 in fully ejected

ascospores collected from the same samples of natural *C. sinensis*.

Li et al. (2013) made additional efforts to identify Genotype #4 of AT-biased Cluster B in 25-day

cultures of mono-ascospores but failed, consistent with the finding shown in Table 4 (Li et al. 2023c).

The authors did not manage to identify the ITS sequences of AT-biased Genotypes #4, #6, and #15–

17 of *O. sinensis* in genomic DNA preparations extracted from the 15 ascosporic strains in both

Groups 2 and 3 of *H. sinensis* (*cf*. Section IV-5). Thus, the expansion of the "ITS pseudogene"

hypothesis from the identified Genotype #5 from the mono-ascospore cultures to all AT-biased

genotypes was apparently an overly generalized interpretation with no supporting evidence. As shown

in Tables 1–3 and Figures 1–2, genomic and phylogenetic analyses did not support the hypothesis

that the "diverged ITS paralogs" (*i.e.*, AT-biased Genotypes #4–6 and #15–17) as the "ITS

pseudogenes" coexist "in a single genome" of Genotype #1 *H. sinensis*.

IV-6-2 Dynamic alterations of multiple genotypes of *O. sinensis* in different compartments of

natural *C. sinensis*

Molecular physiological studies have revealed that the biomasses/abundances of genotypes are

dynamically altered in an asynchronous, disproportional manner during *C. sinensis* maturation,

indicating the genomic independence of multiple genotypes of *O. sinensis* from different perspectives (Zhu et al. 2010, 2012; Gao et al. 2011, 2012; Yao et al. 2011; Li et al. 2016b, 2016c, 2022a, 2022c, 2023c).

Zhu et al. (2010) and Gao et al. (2011, 2012) reported that Genotype #4 of AT-biased Cluster-B of *O. sinensis* predominated in the immature stroma of natural *C. sinensis,* which was in its asexual growth stage; however, its abundance dramatically declined in the mature stroma*,* which was in its sexual reproductive stage. The maturational decline in Genotype #4 abundance in the *C. sinensis* stroma was accompanied by a significant reciprocal elevation in the abundance of Genotype #5 of AT-biased Cluster-A of *O. sinensis*, indicating that the multiple AT-biased genotypes belong to independent *O. sinensis* fungi.

Li et al. (2023b) used the MassARRAY SNP mass spectrometry genotyping technique and found dynamic alterations of GC-biased Genotype #1 and several AT-biased genotypes of *O. sinensis* in the stromal fertile portion (densely covered with numerous ascocarps) before and after ejection of *C. sinensis* ascospores and in the fully and semiejected ascospores collected from the same *C. sinensis* specimens. A highly abundant SNP peak that contained both AT-biased Genotypes #6 and #15 was shown in the preejection stromal fertile portion of natural *C. sinensis,* but the abundance dramatically declined in the stromal fertile portion after ascospore ejection, while the coexisting AT-biased Genotype #5 maintained high abundance.

The ITS sequences of the GC-biased genotypes contained an *Eco*RI endonuclease cleavage site (G▾AATTC in purple in Figure 1) at the position of nucleotides 294→299 of AB067721 (a representative sequence of Genotype #1 *H. sinensis*). The *Eco*RI cleavage site was lost (GAATTT) in all AT-biased genotypes due to a single base C-to-T (cytosine→threonine) transition mutation. To examine dynamic alterations of the natural biomasses of GC- and AT-biased genotypes of *O. sinensis* without PCR manipulation that causes nonlinear, exponential amplification of DNA sequences, the Southern blotting technique was applied to examine genomic DNA after enzymolysis with endonucleases *Ava*I, *Dra*I, and *Eco*RI (Zhu et al. 2010). This non-PCR approach identified a single *Eco*RI-sensitive DNA moiety in genomic DNA isolated from mycelia of *H. sinensis* (*cf.* Figure 4B of Zhu et al. 2010) and a doublet (both *Eco*RI-sensitive GC-biased Genotype #1 and *Eco*RI-resistant AT-biased genotypes) in genomic DNA isolated from natural *C. sinensis* (Figure 7). The lack of *Eco*RI-resistant AT-biased genotypes in pure *H. sinensis* shown in Figure 4B of (Zhu et al. 2010) indicates that the sequences of AT-biased genotypes of *O. sinensis* are absent in the genome of a GC-

biased Genotype #1 *H. sinensis* strain, similar to the *H. sinensis* strains in Groups 1 and 2 summarized
above in Section IV-5. Because Li et al. (2013) incorrectly explained biochemical findings obtained
by Southern blotting (Zhu et al. 2010), we provide help to the mycologists by reproducing the results
(with permission) below in Figure 7 with detailed explanations. The upper panel of Figure 7 shows
the Southern blot using an *H. sinensis*-specific probe that was designed based on the ITS1 sequence
of Genotype #1 *H. sinensis* and demonstrates the following:

Figure 7. Southern blot of C. sinensis nrDNA in the stroma and caterpillar body of natural C. sinensis
during maturation. *[Reproduced with permission from AJBMS (www.nwpii.com/ajbms) Am J Biomed Sci 2010;*
2(3): 217-238] [Zhu et al. *2010].*
Caterpillar refers to caterpillar body. Genomic DNA templates were isolated from the stromata or caterpillar
bodies of immature (Im) or mature (M) C. sinensis specimens collected from Sichuan Province in China and
*prepared using the restriction enzymes Ava*I*, Dra*I*, and Eco*RI*. **The upper panel** was probed with an H.*
*sinensis-specific probe that was designed based on the ITS1 sequence of Genotype #1 H. sinensis. **The lower***
***panel** was probed with a nonspecific 18S internal control probe that was designed based on the 18S gene*
sequence of Genotype #1 H. sinensis, which is >700 bp upstream of the ITS1 segment.

(1) <u>*O. sinensis* genotypes identified in the stromata</u>: the left 2 lanes of the upper panel of Figure 7
show that the slower-migrating DNA moieties (*Eco*RI-resistant AT-biased genotypes)
predominate naturally without PCR amplification in the stromata of immature (IM) and mature
(M) *C. sinensis*. The faster-migrating moieties (*Eco*RI-sensitive GC-biased *H. sinensis*) are
always minor moieties in the stromata during *C. sinensis* maturation.

(2) <u>*O. sinensis* genotypes identified in the caterpillar bodies</u>: although the second lane from the right
in the upper panel of Figure 7 seems to show a single *Eco*RI-sensitive moiety in the immature
(IM) caterpillar body but a doublet in the mature (M) caterpillar body shown in the far right lane,
PCR experiments using genotype-specific primers and touch-down and nested PCR protocols
were able to identify *Eco*RI-resistant AT-biased genotypes in the immature caterpillar body.
These results suggest that the natural quantity of *Eco*RI-resistant AT-biased genotypes in the
immature caterpillar body was below the detection limit of the Southern blotting technique,
which is much less sensitive than PCR-based techniques. The quantity of AT-biased genotypes

was dramatically and disproportionally increased from a microquantity in the immature

caterpillar body to a level comparable to that of *Eco*RI-sensitive GC-biased *H. sinensis* in the

mature caterpillar body.

(3) <u>*O. sinensis* genotypes identified in the stroma and caterpillar body of immature *C. sinensis*</u>: the

first and third lanes from the left in Figure 7 show that the abundances and density ratios of

*Eco*RI-resistant AT-biased genotypes and *Eco*RI-sensitive GC-biased *H. sinensis* differed

significantly in the stroma and caterpillar body of immature *C. sinensis*, which were sampled

from the same specimens.

(4) <u>*O. sinensis* genotypes identified in the stroma and caterpillar body of mature *C. sinensis*</u>: the first

and third lanes from the right in Figure 7 show that the abundances and density ratio of *Eco*RI-

resistant and *Eco*RI-sensitive genotypes differed in the stroma and caterpillar body of mature *C.*

sinensis, which were sampled from the same specimens.

The disproportional distribution of GC- and AT-biased *O. sinensis* genotypes in the stromata and

caterpillar bodies of natural *C. sinensis* and their asynchronous, disproportional alterations of the

natural biomasses without PCR amplification during *C. sinensis* maturation indicate that the

sequences of AT-biased *O. sinensis* genotypes do not reside in the single genome of Genotype #1 *H.*

sinensis. Instead, the genomically independent AT-biased *O. sinensis* genotypes are interindividually

accompanied by GC-biased Genotype #1 *H. sinensis* in the stroma and caterpillar body of natural *C.*

sinensis.

Zhu et al. (2010, 2012) also used the *Eco*RI enzymolysis approach and a restriction fragment

length polymorphism (RFLP) technique and examined GC-biased Genotypes #1 and #2 of *O. sinensis*

in the *C. sinensis* stromata. The authors demonstrated dynamic but asynchronous maturational

alterations of the cooccurring Genotypes #1 and #2 in the stromata of immature, maturing, and mature

C. sinensis specimens, indicating genomic independence of these 2 GC-biased genotypes of *O.*

sinensis. Figures 2 and 3 above show that Genotype #2 of *O. sinensis* is genetically and

phylogenetically distinct from the repetitive ITS copies identified in the genome of Genotype #1 *H.*

sinensis, further confirming the genomic independence of the 2 GC-biased genotypes and ruling out

the possibility that they represent 2 of the repetitive ITS copies in the single *O. sinensis* genome.

If "Different types of ITS sequences were thus proved to coexist in a single genome" as assumed

by Li et al. (2013), the distribution of GC- and AT-biased *O. sinensis* genotypes should be proportional

in the *C. sinensis* stroma and caterpillar body and any biomass alterations of the co-occurring

genotypes should be synchronous and proportional during *C. sinensis* maturation. Southern blotting, as well as RFLP after pretreatment with *Eco*RI enzymolysis and MassARRAY SNP mass spectrometry genotyping assays, clearly demonstrated asynchronous and disproportional alterations of the GC- and AT-biased *O. sinensis* genotypes (Zhu et al. 2010, 2012; Gao et al. 2011, 2012; Yao et al. 2011; Li et al. 2023b, 2023c), and Figures 1–3 and Tables 2 and 4 in this paper further confirmed the genomic independence of the multiple *O. sinensis* genotypes, which belong to independent *O. sinensis* fungi (Stensrud et al. 2007; Xiao et al. 2009).

After the hypothesis of "ITS pseudogenes … in a single genome" was published and subjected to academic discussion, Li et al. (2013) had sufficient time to validate their hypothesis by conducting genome sequencing on one of the Group 3 strains (*cf*. Section IV-5). However, scientific papers addressing this have yet to appear. Instead, Li et al. (2013, 2019, 2020b) stated repeatedly that "Different types of ITS sequences were thus proved to coexist in a single genome", "from a single genome, and not from different genomes", and "The coexistence of diverged ITS paralogs in a single genome". These scientists either did not conduct genome sequencing of one of the Group 3 strains (*cf*. Section IV-5) or may have failed the experiments due to high similarity of the genomic sequences of multiple genotypes of *O. sinensis* that evolved from a common ancestor during a long evolutionary history, which causes difficulties in assembly of shotgun reads to verify their hypothesis that Genotypes #1 and #5 coexist in "a single genome", although this genome sequencing strategy may not be able to confirm the coexistence of AT-biased Genotypes #4, #6, and #15–17 as the genomic components of "a single genome" of GC-biased Genotype #1 *H. sinensis*.

IV-7. Possible synergistic effects of several genotypic fungi in *O. sinensis* sexual reproduction

Hu et al. (2013) and Bushley et al. (2013) hypothesized self-fertilization as the reproductive mode of Genotype #1 *H. sinensis* under homothallism or pseudohomothallism based on the identification of mating-type genes of both *MAT1-1* and *MAT1-2* idiomorphs in 2 *H. sinensis* strains. Li et al. (2020b) stated accordingly that "*O. sinensis* is one of the only known homothallic species in *Ophiocordycipitaceae* … the homothallic life-style may have increased the likelihood and frequency of sex in *O. sinensis*, leading to increased RIP in this species." However, Zhang and Zhang (2015) revealed differential occurrence of the mating-type genes of *MAT1-1* and *MAT1-2* idiomorphs in numerous *H. sinensis* strains, destabilizing the foundation of the self-fertilization hypothesis, and they alternatively hypothesized facultative hybridization as the mode of *O. sinensis* reproduction.

In addition to the analysis of general intraspecific genetic variations and differential expression of *H. sinensis* genes (Li et al. 2022b, 2022c), Li et al. (2021, 2023a) focused on the differential occurrence of mating-type genes of the *MAT1-1* and *MAT1-2* idiomorphs of 236 *H. sinensis* strains: 22 strains (9.3%) contained only MAT1-1-1 genes, 65 (27.5%) possessed only MAT1-2-1 genes, and 149 (63.1%) had both genes. The differential occurrence of mating-type genes underlies the genetic control of *O. sinensis* reproduction. The authors also identified differential but reciprocal transcription of mating-type genes in two *H. sinensis* strains. The MAT1-1-1 transcript was absent, but the MAT1-2-1 transcript was present in transcriptome assembly GCQL00000000 of *H. sinensis* Strain L0106 (Liu et al. 2015; Li et al. 2021, 2023a). The MAT1-1-1 transcript was present in *H. sinensis* Strain 1229, but the MAT1-2-1 transcript included an unspliced intron I, which contains 3 stop codons (Bushley et al. 2013; Li et al. 2021, 2023a). Bushley et al. (2013) specifically commented that a "comparison of DNA and cDNA sequences of MAT1-2-1 revealed only one spliced intron of 55 bp" in *H. sinensis* Strain 1229, which "did not result from contamination of genomic DNA". Thus, the results indicated the production of a largely truncated and dysfunctional MAT1-2-1 protein lacking the majority of the protein encoded by exons II and III. The transcriptional silencing of one of the mating-type genes and the MAT1-2-1 transcript with an unspliced intron I constitute *O. sinensis* reproduction controls at the transcriptional level and the coupled transcriptional-translational level.

Accordingly, Li et al. (2021, 2023a) demonstrated the self-sterility of GC-biased Genotype #1 *H. sinensis*, suggesting a requirement of mating partners for *O. sinensis* sexual reproduction. These partners could be either intraspecific species, such as *H. sinensis* Strains L0106 and 1229, for physiological heterothallism regardless of whether *H. sinensis* is monoecious or dioecious, or heterospecific fungal species that may undergo hybridization if they are able to overcome interspecific reproductive barriers (Zhang and Zhang 2015; Li et al. 2021, 2022c, 2023a, 2023c). The taxonomic positions of Genotypes #2–17 of *O. sinensis* have not been determined, especially for AT-biased genotypes belonging to different *O. sinensis* fungi "even at higher taxonomic levels (genera and family)" (Stensrud et al. 2007) and GC-biased Genotypes #2 and #7 of *O. sinensis*, which cooccur with Genotype #1 in natural *C. sinensis* (Zhu et al. 2010, 2012; Chen et al. 2011). The coexistence of several AT-biased genotypes with GC-biased Genotype #1 in the ascocarps and ascospores of natural *C. sinensis* may constitute a genetic basis for sexual reproduction of *O. sinensis*, involving fungal hybridization (Li et al. 2023b, 2023c).

In addition, Barseghyan et al. (2012) reported the findings of a macro- and micromycology

study and concluded that both *H. sinensis* and *T. sinensis* were the anamorphs of *O. sinensis*. Engh (1999) hypothesized that *Cordyceps* and *Tolypocladium* form a fungal complex in natural *C. sinensis*, and the *Cordyceps* AJ786590 sequence was reported and uploaded to the GenBank database by Stensrud et al. (2005). Stensrud et al. (2007) clustered AJ786590 and other sequences into Group-B of *O. sinensis*, which belong to the AT-biased Genotype #4 and are genetically and phylogenetically distinct from GC-biased Genotype #1 *H. sinensis* (*cf.* Figures 1–2).

Li et al. (2023c) discovered GC-biased Genotypes #13–14 of *O. sinensis*, which exhibit the genetic characteristics of large DNA segment reciprocal substitutions and genetic material recombination between the genomes of the 2 parental fungi, *H. sinensis* and an AB067719-type fungus. The discovery of Genotypes #13–14 of *O. sinensis* likely indicates the possibility of fungal hybridization or parasexuality between 2 heterospecific fungi.

IV-8. Transcription of the 5.8S gene and PCR amplicons of 5.8S-F/R primers

The analysis in Section III-5 indicates that Li et al. (2013, 2019) have neglected the evidence of transcriptional silencing of 5.8S genes demonstrated in metatranscriptome studies in natural settings and the evidence of dynamic and unbalanced transcriptional activation and silencing of numerous fungal genes during the continuous *in vitro* fermentation or liquid incubation of Genotype #1 *H. sinensis* (Xiang et al. 2014; Liu et al. 2015; Xia et al. 2017). Changes in culture conditions and durations have significant impacts on the transcription of fungal genes (*cf.* Tables 6–7), which involves the transcriptional activation and silencing of numerous fungal genes. Because of the conservative evolution of 5.8S genes and many other nonrDNA genes and the incomplete study design and controversial findings reported by Li et al. (2013), it needs to be verified using other sophisticated transcriptional expression methods whether the cDNA identified by Li et al. (2013) was truly derived from the 5.8S gene of Genotype #1 *H. sinensis*, from mutant genotypes of *O. sinensis* or other colonized fungi, or even from non-rRNA genes before confidently declaring that the 5.8S genes of the multiple AT-biased genotypes of *O. sinensis* are truly permanently nonfunctional pseudogenes under the "ITS pseudogene … in a single genome" hypothesis.

Li et al. (2013) did not disclose the amplicon sequences amplified using 5.8SF/R primers and the PCR template of the cDNA library constructed through reverse transcription of total RNA derived from *H. sinensis* Clone 1220. Li et al. (2019) confirmed that Li et al. (2013) designed 5.8SF/R primers based on the sequence "in the most conserved region" of *H. sinensis* rRNA, meaning low specificity of the primers and resulting in uncertain genetic sources of the PCR amplicons. According to the

sequence analysis in Section III-6, the amplicon sequences showed only 90.4%–91.2% similarity to the 5.8S gene sequence of Genotype #1 of *O. sinensis*. The 8.8–9.6% dissimilarity rates may not be convincingly attributed to "mismatches during reverse transcription or PCR amplification", as explained by Li et al. (2013). Thus, solid evidence is lacking to conclude that the amplicon sequences were truly derived from the 5.8S gene of Genotype #1 *H. sinensis* and to claim the *H. sinensis* 5.8S gene as the functional gene and the 5.8S gene of Genotype #5 of *O. sinensis* as the "ITS pseudogene".

Prior to declaring that the 5.8S genes of all AT-biased *O. sinensis* genotypes are permanently dysfunctional pseudogenes co-occurring with a functional 5.8S gene in the genome of GC-biased Genotype #1, several other concerns should be addressed.

(1) Selker and his colleague (1985, 1990, 1997, 2002), Taranto et al. (2014), Hane et al. (2015) and Gladyshev (2017) have repeatedly reported epigenetic methylation of nearly all cytosine residues in 5.8S RNA genes and subsequent transcriptional silencing of GC-biased genes, which occur only in certain fungal species. As shown in Table 3 and Figures 1 and 3 above, the multiple genomic repetitive ITS copies of GC-biased Genotype #1 *H. sinensis* contain mainly insertion, deletion and transversion alleles but no or only a few transition alleles. The repetitive ITS copies of Genotype #1 are genetically and phylogenetically distinct from the AT-biased genotypes of *O. sinensis* (Figures 1–3). Although the 5.8S gene of GC-biased Genotype #1 contains a large number of cytosine residues and is potentially more susceptible to epigenetic methylation attack, introducing nonsense or missense mutations that subsequently cause translational silencing of 5.8S genes, the evidence presented in this paper indicates that *H. sinensis* species may not be the target of RIP mutagenesis and epigenetic methylation attack. It is unlikely that the AT-biased genotypes of *O. sinensis* emerged after the generation of a new *H. sinensis* genome; instead, they are likely genomically independent, existing in different *O. sinensis* fungi.

(2) Li et al. (2013) did not detect the ITS sequences of Genotypes #2–4 and #6–17 of *O. sinensis* in the genomic DNA pool of mono-ascosporic cultures. Thus, it is an apparent overgeneralization to infer that the 5.8S genes of these genotypes are dysfunctional "ITS pseudogene" components of the genome of Genotype #1 *H. sinensis*.

(3) The sequences of Genotypes #2–17 are absent in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000 of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-2031229, respectively (Hu et al. 2013; Li et al. 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020), indicating genomic independence

1037   of *O. sinensis* genotypes.

(4)   The culture-dependent approach used by Li et al. (2013) might have missed some fungal species that are nonculturable or difficult to culture under the *in vitro* experimental settings of the study. The culture-dependent strategy might have a significant impact on the transcription of many fungal genes, with many genes being switched on-or-off nonnaturally, as evidenced by the nonlinear reduction in the total number of transcriptomic unigenes and increases in the average length but decreases in the GC content of transcripts during 3–9 days of liquid fermentation of *H. sinensis* Strain L0106 (*cf*. Table 7) (Liu et al. 2015). Table 6 also demonstrates that changes in the culture duration from 4–15 days and the use of culture medium with or without the addition of adenosine or adenine had dramatic impacts on fungal gene transcription (Pang et al. 2018). A much greater impact on differential transcriptional activation and silencing of many genes may be expected after the prolonged 25-day liquid incubation period adopted by Li et al. (2013).

(5)   Three distinct types of secondary steric conformations of 5.8S rRNA were predicted for Groups A–C (*i.e.*, Genotypes #1 and #4–5) of *O. sinensis* by Li et al. (2013). The possibility of producing circular RNA through back splicing-exon circularization (Dong et al. 2014a) should be considered during study design because these distinct steric structures may have a considerable impact on reverse transcription PCR and cDNA sequencing. In addition, the ITS1-5.8S-ITS2 sequences of Genotypes #2 and #6 may adopt different types of steric conformations (Zhu & Wu 2015; Li et al. 2016b, 2022c; Zhu & Li 2017); thus, the design of other genotype-specific primers and the combined use of other molecular techniques should be considered.

(6)   Wei et al. (2016) reported the identification of a single teleomorph of AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis* and a single teleomorph of GC-biased Genotype #1 in natural *C. sinensis* (*cf*. Figure 6 of Wei et al. 2016). If AT-biased Genotype #4 represents a nonfunctional ITS pseudogene, as believed by Li et al. (2013, 2019, 2020b), the single *O. sinensis* teleomorph in cultivated *C. sinensis* might have disturbed teleomorphic functions, leading to reproductive sterility and an abnormal lifecycle of cultivated *C. sinensis*. Wei et al. (2016) did not share any information about ascospore production in cultivated *C. sinensis*, which is the critical sign of sexual reproduction of *O. sinensis* (Jiang and Yao 2003).

(7)   Li et al. (2022c, 2023b) reported the identification of teleomorphic Genotypes #4 and #15 of *O. sinensis* of AT-biased Cluster-B in the stromal fertile portion densely covered with numerous

ascocarps but not in ascospores collected from the same pieces of natural *C. sinensis*. AT-biased
Genotypes #6 and #15 were found at high abundance in the stromal fertile portion prior to
ascospore ejection, and their abundance drastically declined after ascospore ejection, while
teleomorphic Genotype #5 maintained its high abundance in the stromal fertile portion before
and after ascospore ejection (Li et al. 2022a, 2022c).

(8) The 5.8S genes of multiple *O. sinensis* genotypes may be transcriptionally activated or silenced
in different developmental and maturational stages of natural *C. sinensis*. Significant changes in
the proteomic expression profiles of the stroma and caterpillar body of natural *C. sinensis* during
maturation constitute evidence of such dynamic alterations in epigenetic, transcriptional,
translational and posttranslational modifications of numerous genes (Dong et al. 2014b).

Apparently, Li et al. (2013) improperly designed the study and provided insufficient and
nonconclusive evidence to support their "ITS pseudogene" hypothesis. The whole package of gene
expression in *O. sinensis* fungi and natural *C. sinensis* needs to be further explored before the 5.8S
genes of AT-biased genotypes of *O. sinensis* can be defined as permanently nonfunctional
pseudogenes.

V. Conclusion

This study analyzes the multiple repetitive ITS copies in the genome of Genotype #1 *H. sinensis*
and multiple genotypes of *O. sinensis* fungi and discusses the "ITS pseudogene … in a single genome"
hypothesis for all AT-biased *O. sinensis* genotypes (Li et al. 2013, 2019, 2020b). The repetitive ITS
copies in the genome of Genotype #1 contain multiple scattered insertion, deletion and transversion
point mutations and may not be generated through RIP mutagenesis because RIP mutagenesis
theoretically causes C-to-T transitions. The GC-biased repetitive ITS copies are genetically and
phylogenetically distinct from AT-biased *O. sinensis* genotypes that contain multiple scattered
transition alleles. GC-biased Genotype #1 *H. sinensis* may not be the fungal species target of RIP
mutagenesis and epigenetic methylation attack. The AT-biased genotypes are independent from the
genome of Genotype #1 and might have been generated from a common ancestor through certain
mutagenic and evolutionary mechanisms in the long history of evolution in response to the extremely
harsh environment of the Qinghai-Tibet Plateau and become independent *O. sinensis* fungi with
independent genomes coexisting in parallel with GC-biased genotypes in natural *C. sinensis*. Li et al.
(2013) provided incomplete and controversial information to support their "ITS pseudogene … in a

single genome" hypothesis, which needs to be further validated.

VI. Acknowledgments:

1099 This research was supported by a grant from the Science and Technology Department of Qinghai
1100 Province, China, grant number 2021-SF-A4 "Study on key technologies of conservation of natural
1101 resource and industrial upgrading of *Cordyceps sinensis*", the major science and technology projects
1102 in Qinghai Province.

References

[1] Barseghyan GS, Holliday JC, Price TC, Madison LM, Wasser SP. Growth and cultural-morphological
characteristics of vegetative mycelia of medicinal caterpillar fungus *Ophiocordyceps sinensis* G.H. Sung et al.
(Ascomycetes) isolates from Tibetan Plateau (P.R.China). Intl J Med Mushrooms, 2011, 13(6): 565−581.

[2] Bushley KE, Li Y, Wang W-J, Wang X-L, Jiao L, Spatafora JW, Yao Y-J. Isolation of the MAT1-1 mating type
idiomorph and evidence for selfing in the Chinese medicinal fungus *Ophiocordyceps sinensis*. Fungal Biol, 2013,
117, 599–610. (PMID: 22181845 DOI: 10.1016/j.funbio.2013.06.001)

[3] Chen C-S, Hseu R-S, Huang C-T. Quality control of *Cordyceps sinensis* teleomorph, anamorph, and Its products.
Chapter 12, *in* (Shoyama Y, Ed.) *Quality Control of Herbal Medicines and Related Areas*. InTech, Rijeka, Croatia.
2011. (www.intechopen.com)

[4] Chen Y-Q, Hu B, Xu F, Zhang W, Zhou H, Qu L-H. Genetic variation of *Cordyceps sinensis*, a fruit-body-producing
entomopathogenic species from different geographical regions in China. FEMS Microbiol Lett, 2004, 230, 153–158.
(PMID: 14734179 DOI: 10.1016/S0378-1097(03)00889-9)

[5] Dong R, Chen L-L, Yang L. Circular RNA studies in the post-genomic era. Chin J Cell Biol, 2014a, 36 (11), 1455–
1459. (DOI: 10.11844/cjcb.2014.11.9001)

[6] Dong Y-Z, Zhang L-J, Wu Z-M, Gao L, Yao Y-S, Tan N-Z, Wu J-Y, Ni L-Q, Zhu J-S. Altered proteomic
polymorphism in the caterpillar body and stroma of natural *Cordyceps sinensis* during maturation. PLoS ONE,
2014b, 9 (10), e109083. (PMID: 25310818 doi:10.1371/journal.pone.0109083)

[7] Engh IB. Molecular phylogeny of the *Cordyceps-Tolypocladium* complex. Candidate scientific thesis, Department
of Biology, University of Oslo, Oslo. 1999.

[8] Gao L, Li X-H, Zhao J-Q, Lu J-H, Zhu J-S. Detection of multiple *Ophiocordyceps sinensis* mutants in premature
stroma of *Cordyceps sinensis* by MassARRAY SNP MALDI-TOF mass spectrum genotyping. Beijing Da Xue Xue
Bao, 2011, 43 (2), 259–266. (PMID: 21503123 doi: 10.3969/j.issn.1671-167X.2011.02.019)

[9] Gao L, Li X-H, Zhao J-Q, Lu J-H, Zhao J-G, Zhu J-S. Maturation of *Cordyceps sinensis* associates with alterations
of fungal expressions of multiple *Ophiocordyceps sinensis* mutants with transition and transversion point mutations
in stroma of *Cordyceps sinensis*. Beijing Da Xue Xue Bao, 2012, 44 (3), 454–463. (PMID: 22692321 doi:
10.3969/j.issn.1671-167X.2012.03.025)

[10] Gladyshev E. Repeat-Induced Point Mutation (RIP) and Other Genome Defense Mechanisms in Fungi. Microbiol
Spectr, 2017, 5(4): 10.1128. (PMID: 28721856 doi: 10.1128/microbiolspec.FUNK-0042-2017)

[11] Hane JK, Williams AH, Taranto AP, Solomon PS, Oliver RP. Repeat-induced point mutation: A fungal-specific,

endogenous mutagenesis process. In: van den Berg M, Maruthachalam K (eds) Genetic transformation systems in

fungi, Volume 2. Fungal Biology. Springer, Cham. 2015. https://doi.org/10.1007/978-3-319-10503-1_4

[12] Hawksworth DL, Crous PW, Redhead SA, Reynolds DR, Samson RA, Seifert KA, Taylor JW, Wingfield MJ, Abaci

O, Aime C, Asan A, Bai F-Y, de Beer ZW, Begerow D, Berikten D, Boekhout T, K Buchanan P, Burgess T, Buzina

W, Cai L, Cannon PF, Crane JL, Damm U, Daniel H-M, van Diepeningen AD, Druzhinina I, Dyer PS, Eberhardt

U, Fell JW, Frisvad JC, Geiser DM, Geml J, Glienke C, Gräfenhan T, Groenewald JZ, Groenewald M, de Gruyter J,

Guého-Kellermann E, Guo L-D, Hibbett DS, Hong S-B, de Hoog GS, Houbraken J, Huhndorf SM, Hyde K D,

Ismail A, Johnston PR, Kadaifciler DG, Kirk PM, Kõljalg U, Kurtzman CP, Lagneau P-E, Lévesque CA, Liu X,

Lombard L, Meyer W, Miller A, Minter DW, Najafzadeh MJ, Norvell L, Ozerskaya SM, Oziç R, Pennycook SR,

Peterson SW, Pettersson OV, Quaedvlieg W, Robert VA, Ruibal C, Schnürer J, Schroers H-J, Shivas R, Slippers B,

Spierenburg H, Takashima M, Taşkın E, Thines M, Thrane U, Uztan AH, van Raak M, Varga J, Vasco A, Verkley G,

Videira SIR, de Vries RP, Weir BS, Yilmaz N, Yurkov A, Zhang N. The Amsterdam declaration on fungal

nomenclature. IMA Fungus 2011, 2: 105−112 (PMID: 22679594 DOI: 10.5598/imafungus.2011.02.01.14).

[13] Hood ME, Katawczik M, Giraud T. Repeat-induced point mutation and the population structure of transposable

elements in Microbotryum violaceum. Genetics. 2005, 170(3): 1081-1089. doi: 10.1534/genetics.105.042564. Epub

2005 May 23. PMID: 15911572; PMCID: PMC1451165

[14] Hu X, Zhang Y, Xiao G, Zheng P, Xia Y, Zhang X, St Leger RJ, Liu X, Wang C. Genome survey uncovers the secrets

of sex and lifestyle in caterpillar fungus. Chin Sci Bull, 2013, 58, 2846–2854. (doi: 10.1007/s11434-013-5929-5)

[15] Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogeny. Bioinformat, 2001, 17: 754−755. (PMID:

11524383 DOI: 10.1093/bioinformatics/17.8.754)

[16] Jiang Y, Yao Y-J. Anamorphic fungi related to *Cordyceps sinensis*. Mycosystema, 2003, 22(1): 161−176.

[17] Jin L-Q, Xu Z-W, Zhang B, Yi M, Weng C-Y, Lin S, Wu H, Qin X-T, Xu F, Teng Y, Yuan S-J, Liu Z-Q, Zheng Y-G.

Genome sequencing and analysis of fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*. AMB Expr.

2020, 10:105. (PMID: 32494871 DOI: 10.1186/S13568-020-01039-x)

[18] Kinjo N, Zang M. Morphological and phylogenetic studies on *Cordyceps sinensis* distributed in southwestern China.

Mycoscience, 2001, 42, 567–574. (https://doi.org/10.1007/BF02460956)

[19] Li C-L, Tan N-Z, Berger JL, Ferguson SB, Zhang Y, Prolla TA, Zhu J-S. The Combined use of ageLOC whole-gene

expression profiling technology and mouse lifespan test in anti-aging herbal product study. Proc. 2011 New TCM

Prod. Innovat. Indust. Devel. Summit, Hangzhou, China, 2011, pp. 443–448.

(https://xueshu.baidu.com/usercenter/paper/show?paperid=08341c17fa58c8f85584b92572b90f75&site=xueshu_se)

[20] Li X-Z, Li Y-L, Zhu J-S. *Differential expression of mating-type genes in Hirsutella sinensis and natural Cordyceps*

sinensis. Proc. 2021 Intl Conf Info Technol Biomed Engineering, 2021, pp. 318−333.

(https://conferences.computer.org/icitbepub/pdfs/ICITBE2021-

6JNtJiThlp10s9N5Sodvuy/009900a318/009900a318.pdf)

[21] Li X-Z, Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. Further discussion with Li et al. (2013, 2019) regarding the "ITS

pseudogene hypothesis" for *Ophiocordyceps sinensis*. Mol Phylogenet Evol, 2020a, 146: 106728. (PMID: 31904509

DOI: 10.1016/j.ympev.2019.106728)

[22] Li X-Z, Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. Molecular identification of *Ophiocordyceps sinensis* genotypes and the indiscriminate use of the Latin name for the multiple genotypes and the natural insect-fungi complex. Am J Biomed Sci, 2022a, 14(3): 115–135. (doi:10.5099/aj220300115)

[23] Li X-Z, Li Y-L, Yao Y-S, Zhu J-S. Differential expressions of the *Hirsutella sinensis* genes and intraspecific genetic variations among *H. sinensis* strains. Biochem Mol Biol J. 2022b, 8(2): 58. (DOI: 10.36648/2471-8084.22.8.58)

[24] Li X-Z, Li Y-L, Zhu J-S. Differential transcription of the mating-type genes during the sexual life of natural *Cordyceps sinensis* insect-fungi complex. Chin J Chin Mater Med, 2023a, 48(xx): xxx (in press).

[25] Li Y, Hsiang T, Yang R-H, Hu X-D, Wang K, Wang W-J, Wang X-L, Jiao L, Yao Y-J. Comparison of different sequencing and assembly strategies for a repeat-rich fungal genome, *Ophiocordyceps sinensis*. J Microbiol Methods, 2016a, 128, 1–6. (http://dx.doi.org/10.1016/j.mimet.2016.06.025)

[26] Li Y, Jiang L, Wang K, Wu H-J, Yang R-H, Yan Y-J, Bushley KE, Hawksworth DL, Wu Z-J, Yao Y-J. RIP mutated ITS genes in populations of *Ophiocordyceps sinensis* and their implications for molecular systematics. IMA Fungus, 2020b, 11, 18. (PMID: 27343682 DOI: 10.1016/j.mimet.2016.06.025)

[27] Li Y, Jiao L, Yao Y-J. Non-concerted ITS evolution in fungi, as revealed from the important medicinal fungus *Ophiocordyceps sinensis*. Mol Phylogenetics Evol, 2013, 68, 373–379. (http://dx.doi.org/10.1016/j.ympev.2013.04.010)

[28] Li Y, Wang Y-H, Wang K, Yang R-H, Jiao L, Yao Y-J. Response to "The multiple genotypes of *Ophiocordyceps sinensis* and the ITS pseudogene hypothesis". Mol Phylogenetics Evol, 2019, 139, 106522. (PMID: 23618625 DOI: 10.1016/j.ympev.2013.04.010)

[29] Li Y-L, Gao L, Yao Y-S, Li X-Z, Wu Z-M, Tan N-Z, Lou Z-Q, Xie W-D, Wu J-Y, Zhu J-S. Altered GC- and AT-biased genotypes of *Ophiocordyceps sinensis* in the stromal fertile portions and ascospores of natural *Cordyceps sinensis*. PLOS ONE, 2023b. (in peer review). BIORXIV Preprint 2022-11-12. (https://www.biorxiv.org/content/10.1101/2022.10.07.511361v1.full.pdf)

[30] Li Y-L, Li X-Z, Yao Y-S, Wu Z-M, Gao L, Tan N-Z, Lou Z-Q, Xie W-D, Wu J-Y, Zhu J-S. Differential cooccurrence of multiple genotypes of *Ophiocordyceps sinensis* in the stromata, stromal fertile portion (ascocarps) and ascospores of natural *Cordyceps sinensis*. PLOS ONE, 2023c. (decision in process). BIORXIV Preprint 2022-6-20. (https://www.biorxiv.org/content/10.1101/2022.06.20.496766v2.full.pdf)

[31] Li Y-L, Li X-Z, Zhu J-S. *Cordyceps sinensis* Fungal Heterokaryons. Chemistry Industry Press Co., Beijing, China. 2022c. (ISBN: 978-7-122-40189-2 http://www.cip.com.cn/Book/Index/48939)

[32] Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. The molecular heterogeneity of natural *Cordyceps sinensis* with multiple *Ophiocordyceps sinensis* fungi challenges the anamorph-teleomorph connection hypotheses. Am J Biomed Sci, 2016b, 8, 123–159. (doi: 10.5099/aj100300217)

[33] Li Y-L, Yao Y-S, Zhang Z-H, Xu H-F, Liu X, Ma S-L, Wu Z-M, Zhu J-S. Synergy of fungal complexes isolated from the intestines of *Hepialus lagii* larvae in increasing infection potency. J Fungal Res, 2016c, 14, 96–112. (DOI: 10.13341/j.jfr.2014.1067)

[34] Liu J, Guo L-N, Li Z-W, Zhou Z, Li Z, Li Q, Bo X-C, Wang S-Q, Wang J-L, Ma S-C, Zheng J, Yang Y. Genomic analyses reveal evolutionary and geologic context for the plateau fungus *Ophiocordyceps sinensis*. Clin Med, 2020, 15:107–119. (PMID: 33042212 DOI: 10.1186/s13020-020-00365-3)

1208   [35] Liu Z-Q, Lin S, Baker P, Wu L-F, Wang X-R, Wu H, Xu F, Wang H-Y, Brathwaite M, Zheng Y-G. Transcriptome
1209   sequencing and analysis of the entomopathogenic fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*.
1210   BMC Genome, 2015, 16, 106. (PMID: 25765329 DOI 10.1186/s12864-015-1269-y)

1211   [36] Liu Z-Y, Yao Y-J, Liang Z-Q, Liu A-Y, Pegler DN, Chase MW. Molecular evidence for the anamorph-teleomorph
1212   connection in *Cordyceps sinensis*. Mycol Res, 2001, 105: 827−832. (https://doi.org/10.1017/S095375620100377X)

1213   [37] Mao X-M, Zhao S-M, Cao L, Yan X, Han R-X. The morphology observation of *Ophiocordyceps sinensis* from
1214   different origins. J Environ Entomol, 2013, 35, 343–353. (doi: 10.3969/j.issn.1674-0858.2013.03.11)

1215   [38] Pang F, Wang LP, Jin Y, Guo LP, Song LP, Liu GM, Feng CQ. Transcriptome analysis of *Paecilomyces hepiali* at
1216   different growth stages and culture additives to reveal putative genes in cordycepin biosynthesis. Genomics 2018,
1217   110, 162–170. http://dx.doi.org/10.1016/j.ygeno.2017.09.008.

1218   [39] Ren Y, Wan D-G, Lu X-M, Guo J-L. The study of scientific name discussion for TCM *Cordyceps*. Lishizhen Med
1219   Materia Medica Res, 2013, 24: 2211−2212 (Doi: 10.3969/j.issn.1008-0805.2013.09.075).

1220   [40] Selker EU. Premeiotic instability of repeated sequences in Neurospora crassa. Annu Rev Genet, 1990, 24:579–613.
1221   (PMID: 2150906 DOI: 10.1146/annurev.ge.24.120190.003051)

1222   [41] Selker EU. Epigenetic phenomena in filamentous fungi: useful paradigms or repeat-induced confusion? Trends
1223   Genet, 1997, 13:296–301. (PMID: 9260514 DOI: 10.1016/S0168-9525(97)01201-8)

1224   [42] Selker EU. Repeat-induced gene silencing in fungi. Adv Genet, 2002, 46:439–450. (PMID: 11931234 DOI:
1225   10.1016/S0065-2660(02)46016-6)

1226   [43] Selker EU, Stevens JN. DNA methylation at asymmetric sites is associated with numerous transition mutations.
1227   Proc Natl Acad Sci USA, 1985, 82:8114–8118. (PMID: 2415981 DOI: 10.1073/pnas.82.23.8114)

1228   [44] Shu R-H, Zhang J-H, Meng Q, Zhang H, Zhou G-L, Li M-M, Wu P-P, Zhao Y-N, Chen C, Qin Q-L. A new high-
1229   quality draft genome assembly of the Chinese cordyceps *Ophiocordyceps sinensis*. Genome Biol Evol, 2020, 12(7):
1230   1074−1079. (PMID: 32579174 DOI: 10.1093/gbe/evaa112)

1231   [45] Stensrud Ø, Hywel-Jones NL, Schumacher T. Towards a phylogenetic classification of Cordyceps: ITS nrDNA
1232   sequence data confirm divergent lineages and paraphyly. Mycol Res, 2005, 109 (1), 41–56. (PMID: 15736862 DOI:
1233   10.1017/S095375620400139X)

1234   [46] Stensrud Ø, Schumacher T, Shalchian-Tabrizi K, Svegården IB, Kauserud H. Accelerated nrDNA evolution and
1235   profound AT bias in the medicinal fungus *Cordyceps sinensis*. Mycol Res, 2007, 111, 409–415. (PMID: 17509850
1236   doi:10.1016/j.mycres.2007.01.015)

1237   [47] Sung G-H, Hywel-Jones NL, Sung J-M, Luangsa-ard JJ, Shrestha B, Spatafora JW. Phylogenetic classification of
1238   *Cordyceps* and the clavicipitaceous fungi. Stud Mycol, 2007, 57, 5–59. (PMID: 18490993 DOI:
1239   10.3114/sim.2007.57.01)

1240   [48] Tan N-Z, Berger JL, Zhang Y, Prolla TA, Weindruch R, Zhao C-S, Bartlett M, Zhu J-S. The lifespan-prolonging
1241   effect of *Cordyceps sinensis* Cs-4 in normal mice and its molecular mechanisms. FASEB J, 2011, 25(1):599.1.
1242   (http://www.fasebj.org/cgi/content/meeting_abstract/25/1_MeetingAbstracts/599.1)

1243   [49] Taranto AP, Hane JK, Williams AH, Solomon PS, Oliver RP. Repeat-induced point mutation: A fungal-specific
1244   endogenous mutagenesis process. In: van den Berg M, Maruthachalam K (eds) Genetic transformation systems in
1245   fungi, Vol. 2. Fungal Biology, Chapter 4. Springer, Cham. 2014. https://doi.org/10.1007/978-3-319-10503-1_4

1246   [50] Wei J-C, Wei X-L, Zheng W, Guo W, Liu R. Species identification and component detection of *Ophiocordyceps*
1247   *sinensis* cultivated by modern industry. Mycosystema, 2016, 35, 404–410. (doi: 10.13346/j.mycosystema.160045)

1248   [51] Wei X-L, Yin X-C, Guo Y-L, Shen N-Y, Wei, J.-C. Analyses of molecular systematics on *Cordyceps sinensis* and its
1249   related taxa. Mycosystema, 2006, 25(2): 192–202.

1250   [52] Xia E-H, Yang D-R, Jiang J-J, Zhang Q-J, Liu Y, Liu Y-L, Zhang Y, Zhang H-B, Shi C, Tong Y, Kim C, Chen H,
1251   Peng Y-Q, Yu Y, Zhang W, Eichler E-E, Gao L-Z. The caterpillar fungus, *Ophiocordyceps sinensis*, genome provides
1252   insights into highland adaptation of fungal pathogenicity. Sci Rep, 2017, 7, 1806. (PMID: 28496210 doi:
1253   10.1038/s41598-017-01869-z)
1254   (www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas for the cDNA
1255   file)

1256   [53] Xia F, Liu Y, Shen G-R, Guo L-X, Zhou X-W. Investigation and analysis of microbiological communities in natural
1257   *Ophiocordyceps sinensis*. Can J Microbiol, 2015, 61: 104−111. (PMID: 25578897 http://dx.doi.org/10.1139/cjm-
1258   2014-0610)

1259   [54] Xiang L, Li Y, Zhu Y-J, Luo H, Li C-F, Xu X-L, Sun C, Song J-Y, Shi L-C, He L, Sun W, Chen S-L. Transcriptome
1260   analysis of the *Ophiocordyceps sinensis* fruiting body reveals putative genes involved in fruiting body development
1261   and cordycepin biosynthesis. Genomics, 2014, 103, 154–159. (PMID: 24440419
1262   http://dx.doi.org/10.1016/j.ygeno.2014.01.002)

1263   [55] Xiao Y-Y, Chen C, Dong J-F, Li C-R, Fan M-Z. Morphological observation of ascospores of *Ophiocordyceps*
1264   *sinensis* and its anamorph in growth process. J Anhui Agricult Univ, 2011, 38(4): 587–591.
1265   (http://www.cnki.net/kcms/detail/34.1162.S.20110624.1628.030.html)

1266   [56] Xiao W, Yang J-L, Zhu P, Cheng K, He H, Zhu H, Wang Q. Non-support of species complex hypothesis of *Cordyceps*
1267   *sinensis* by targeted rDNA-ITS sequence analysis. Mycosystema, 2009, 28, 724–730. (DOI:
1268   10.13346/j.mycosystema.2009.05.028)

1269   [57] Yao Y-S, Gao L, Li Y-L, Ma S-L, Wu Z-M, Tan N-Z, Wu J-Y, Ni L-Q, Zhu J-S. Amplicon density-weighted
1270   algorithms for analyzing dissimilarity and dynamic alterations of RAPD polymorphisms of *Cordyceps sinensis*.
1271   Beijing Da Xue Xue Bao Yi Xue Ban 2014, 46, 618−628. (http://xuebao.bjmu.edu.cn/EN/Y2014/V46/I4/618)

1272   [58] Yao Y-S, Zhou Y-J, Gao L, Lu J-H, Wu Z-M, Zhu J-S. Dynamic alterations of the differential fungal expressions of
1273   *Ophiocordyceps sinensis* and its mutant genotypes in stroma and caterpillar during maturation of natural *Cordyceps*
1274   *sinensis*. J Fungal Res, 2011, 9, 37–49,53. (PMID: 22692321 Doi: 10.3969/j.issn.1672-3538.2011.01.007)

1275   [59] Yao Y-S, Zhu J-S. Indiscriminative use of the Latin name for natural *Cordyceps sinensis* insect-fungi complex and
1276   multiple *Ophiocordyceps sinensis* fungi. Chin J Chin Mater Med, 2016, 41 (7), 1316–1366. (PMID: 28879757 DOI:
1277   10.4268/cjcmm20160733)

1278   [60] Zhang S, Zhang Y-J. Molecular evolution of three protein-coding genes in the Chinese caterpillar fungus
1279   *Ophiocordyceps sinensis*. Microbiology China. 2015, 42(8): 1549−1560. (DOI: 10.13344/j.microbiol.china.150279)

1280   [61] Zhang S, Zhang Y-J, Liu X-Z, Zhang H, Liu D-S. On the reliability of DNA sequences of *Ophiocordyceps sinensis*
1281   in public data bases. J Ind Microbiol Biotechnol, 2013a, 40: 365−378. DOI: 10.1007/s10295-012-1228-4.

1282   [62] Zhang S, Zhang Y-J, Shrestha B, Xu J-P, Wang C-S, Liu X-Z. *Ophiocordyceps sinensis* and *Cordyceps militaris*:
1283   research advances, issues and perspectives. Mycosystema, 2013b, 32: 577−597 (DOI: 10.3969/j.issn.1673-

1284 [5110.2014.18.061](#)).

[63] Zhang S-W, Cem K, Liu Y, Zhou X-W, Wang C-S. Metatranscriptomics analysis of the fruiting caterpillar fungus collected from the Qinghai-Tibetan plateau. Scientia Sinica Vitae, 2018, 48(5): 562. (doi: [10.1360/N052017-00253](#))

[64] Zhang Y-J, Li E-W, Wang C-S, Li Y-L, Liu X-Z. *Ophiocordyceps sinensis*, the flagship fungus of China: terminology, life strategy and ecology. Mycology, 2012, 3: 2−10. (DOI: [10.1080/21501203.2011.654354](#))

[65] Zhang Y-J, Sun B-D, Zhang S, Wang M, Liu X-Z, Gong W-F. Mycobiotal investigation of natural *Ophiocordyceps sinensis* based on culture-dependent investigation. Mycosystema, 2010, 29, 518−527.

[66] Zhang Y-J, Xu L, Zhang S, Liu X-Z, An Z, Wang M, Guo Y. Genetic diversity of *Ophiocordyceps sinensis*, a medicinal fungus endemic to the Tibetan Plateau: implications for its evolution and conservation. BMC Evolut Biol, 2009,. 9, 290. (PMID: [20003548](#) DOI: [10.1186/1471-2148-9-290](#))

[67] Zhang Y-J, Zhang S, Li Y-L, Ma S-L, Wang C-S, Xiang M-C, Liu X-Z, An Z-Q, Xu J-P. Phylogeography and evolution of a fungal–insect association on the Tibetan Plateau. Mol Ecol, 2014, 23, 5337−5355. (doi: [10.1111/mec.12940](#))

[68] Zhao J, Wang N, Chen Y-Q, Li T-H, Qu L-H. Molecular identification for the asexual stage of *Cordyceps sinensis*. Acta Scientiarum Naturalium Universitatis Sunyatseni 1999, 38: 122−123.

[69] Zhu J-S, Gao L, Li X-H, Yao Y-S, Zhao J-Q. Maturational alteration of oppositely orientated rDNA and differential proliferation of GC- and AT-biased genotypes of *Ophiocordyceps sinensis* and *Paecilomyces hepiali* in natural *Cordyceps sinensis*. Am J Biomed Sci, 2010, 2, 217–238. (doi: [10.5099/aj100300217](#))

[70] Zhu J-S, Guo Y-L, Yao Y-S, Zhou Y-J, Lu J-H, Qi Y, Liu X-Z, Wu Z-M, Chen W, Zhang L, Yin W-T, Zheng T-Y, Zhang L-J. Maturation of *Cordyceps sinensis* associates with co-existence of *Hirsutella sinensis* and *Paecilomyces hepiali* DNA and dynamic changes in fungal competitive proliferation predominance and chemical profiles. J. Fungal Res 2007, 5(4), 214–224.

[71] Zhu J-S, Halpern GM, Jones K. The scientific rediscovery of a precious ancient Chinese herbal regimen: *Cordyceps sinensis*: Part I. J Altern Complem Med, 1998a, 4(3): 289–303. (PMID: [9764768](#) DOI: [10.1089/acm.1998.4.3-289](#))

[72] Zhu J-S, Halpern GM, Jones K. The scientific rediscovery of an ancient Chinese herbal medicine: *Cordyceps sinensis*: Part II. J Altern Complem Med, 1998b, 4(4): 429–457. (PMID: [9884180](#) DOI: [10.1089/acm.1998.4.429](#))

[73] Zhu J-S, Li Y-L. A Precious Transitional Chinese Medicine, *Cordyceps sinensis*: Multiple Heterogeneous *Ophiocordyceps sinensis* in the Insect-Fungi Complex. Lambert Academic Publishing, Germany, 2017. ([ISBN-10: 9783330046689](#), [ISBN-13: 978-3330046689](#))

[74] Zhu J-S, Li Y-L, Yao Y-S, Xie W-D. Molecular evidence not supporting the use of *Hirsutella sinensis* Strain EFCC 7287 as the taxonomic standard for multiple *Ophiocordyceps sinensis* fungi. Int J Mol Biol, 2018, 3, 114–115. (DOI: [10.15406/ijmboa.2018.03.00062](#))

[75] Zhu J-S, Li Y-L, Yao Y-S, Xie W-D. The multiple genotypes of *Ophiocordyceps sinensis* and the ITS pseudogene hypothesis. Mol Phylogenetics Evol, 2019, 139, 106322. (PMID: [30394296](#) DOI: [10.1016/j.ympev.2018.10.034](#))

[76] Zhu J-S, Wu J-Y. Genetic heterogeneity of natural *Cordyceps sinensis* with co-existence of multiple fungi. Chin J Cell Biol, 2015, 37 (2), 284–298. (DOI: [10.11844/cjcb.2015.02.0284](#))

[77] Zhu J-S, Zhao J-G, Gao L, Li X-H, Yao Y-S, Zhao J-Q. Dynamically altered expressions of at least 6 *Ophiocordyceps*

1321   *sinensis* mutants in the stroma of *Cordyceps sinensis*. J Fungal Res, 2012, 10, 100–112. (DOI:

1322   10.13341/j.jfr.2012.02.007)