AT-biased genotypes of *Ophiocordyceps sinensis* may not be generated through

RIP mutagenesis in a single genome of GC-biased *Hirsutella sinensis*

Xiu-Zhang Li [a], Yu-Ling Li [a], Ya-Nan Wang [b], Jia-Shi Zhu [a,c,*]

[a] State Key Laboratory of Plateau Ecology and Agriculture, Qinghai Academy of Animal and

Veterinary Sciences, Qinghai University, Xining, Qinghai 810016, China

[b] State Key Laboratory of Breeding Base of Dao-di Herbs, National Resource Center for Chinese

Materia Medica, China Academy of Chinese Medical Sciences, Beijing 100700, China

[c] Institute of Biopharmaceutical and Health Engineering, Shenzhen International Graduate School,

Tsinghua University, Shenzhen 518055, China

[*] Corresponding author:

E-mail address: zhujosh@163.com (J-S Zhu)

Short title:

AT-biased *O. sinensis* genotypes are not generated through RIP

Abstract

It has been hypothesized that AT-biased genotypes of *Ophiocordyceps sinensis* are generated through repeat-induced point mutations (RIP) and coexist as permanently nonfunctional internal transcribed spacer (ITS) pseudogenes in the genome of GC-biased Genotype #1 of *O. sinensis* (*Hirsutella sinensis*). This study examined the *H. sinensis* genome, which contains multiple repetitive ITS copies (GC content: 64.7±0.33%) with multiple insertion/deletion and transversion alleles. Apparently, these alleles were not generated through RIP, which theoretically causes cytosine-to-thymine and guanine-to-adenine transitions. The repetitive ITS copies in the *H. sinensis* genome were found to be genetically and phylogenetically distinct from the AT-biased *O. sinensis* genotypes (GC content: 51.1±1.69%), which possess multiple transition alleles. The sequences of Genotypes #2–17, both GC- and AT-biased, are absent from the *H. sinensis* genome; these genotypes belong to interindividual *O. sinensis* fungi and differentially occur in different compartments of natural *Cordyceps sinensis,* with dynamic alterations in abundance occurring in an asynchronous, disproportional manner during *C. sinensis* maturation. Metatranscriptomic analyses of natural *C. sinensis* revealed the transcriptional silencing of 5.8S genes in all *C. sinensis*-colonizing fungi, including *H. sinensis*. The transcription assay reported by Li *et al*. (2013) provided unsound, controversial evidence that the 5.8S genes of AT-biased genotypes are nonfunctional pseudogenes. In addition to the single ITS-locus analysis, repetitive genomic copies were also examined at multiple loci in the *H. sinensis* genome, and approximately 8.27% of the authentic genes had repetitive copies, including various transitions, transversions, insertions, and deletions. The transcripts for the repetitive copies were identified in the *H. sinensis* transcriptome, regardless of the decreases, increases, or bidirectional changes in AT contents of the repetitive copies. These results are inconsistent with those of RIP, which generates pseudogenic, nonfunctional, repetitive copies. AT-biased genotypes of *O. sinensis* might have evolved through evolutionary mechanisms from a common ancestor in parallel with GC-biased Genotype #1 *H. sinensis* over a long evolutionary history.

KEY WORDS: Multiple genotypes of *Ophiocordyceps sinensis*; genomic independence; repeat-induced point mutations (RIP); insertion/deletion, transversion and transition point mutations; 5.8S gene transcript; transcriptional silencing; repetitive copies of authentic genes in multiple genomic loci.

I. Introduction

Natural *Cordyceps sinensis* is one of most valued therapeutic agents in traditional Chinese medicine and has a rich history of clinical use for health maintenance, disease amelioration, postdisease and postsurgery recovery, and antiaging therapy (Zhu *et al*. 1998a, 1998b; Li *et al*. 2011; Tan *et al*. 2011). The Chinese Pharmacopeia defines natural *C. sinensis* as an insect-fungal complex that includes the remains of an Hepialidae moth larva (Zhu and Wu 2015; Li *et al*. 2016b, 2022a, 2022c; Zhu and Li 2017) and >90 fungal species from >37 genera (Zhang *et al*. 2010; Xia *et al*. 2015; Zhang *et al*. 2018), with differential co-occurrence of some of 17 *Ophiocordyceps sinensis* genotypes in different compartments of natural *C. sinensis* (Engh 1999; Kinjo and Zang 2001; Chen *et al*. 2004, 2011; Stensrud *et al*. 2005, 2007; Wei *et al*. 2006, 2016; Xiao *et al*. 2009; Zhang *et al*. 2009; Zhu *et al*. 2010, 2012, 2018, 2019; Gao *et al*. 2011, 2012; Yao *et al*. 2011; Mao *et al*. 2013; Li *et al*. 2020b, 2022c, 2023b, 2023c), *i.e.*, the natural *C. sinensis* insect-fungal complex ≠ the entomopathogenic *O. sinensis* fungus. However, since the 1840s, the Latin name *C. sinensis* has been indiscriminately used for both the wild insect-fungal complex and the teleomorph/holomorph of the *C. sinensis* fungus (Yao & Zhu 2016; Li *et al*. 2022a). The fungus was renamed *Ophiocordyceps sinensis* using *Hirsutella sinensis* Strain EFCC 7287 (GC-biased Genotype #1 of *O. sinensis*) as the nomenclature reference (Sung *et al*. 2007), while the indiscriminately used name of the wild and cultivated insect-fungal complex has remained unchanged (Zhu *et al*. 2018). Since then, especially after the improper implementation of the One Fungus=One Name nomenclature rule in *O. sinensis* research (Hawksworth *et al*. 2011; Zhang *et al*. 2013b), the name *O. sinensis* has been used to refer to multiple teleomorphic and anamorphic fungi that have distinct genomes and has been indiscriminately applied to the natural insect-fungal complex (Zhu and Wu 2015; Yao & Zhu 2016; Li *et al*. 2016b, 2022a, 2022c; Zhu and Li 2017). The confusion surrounding Latin name use continues and has even spread from the scientific community to the public media and mass markets, causing a significant decrease in the wholesale and retail prices of natural *C. sinensis*. Because a consensus Latin name for the wild insect-fungal product has not been identified by taxonomists and multiple genotypes of *O. sinensis* with undetermined taxonomic positions are referred to by the same Latin name (Zhang *et al*. 2012; Ren *et al*. 2013; Yao & Zhu 2016; Li *et al*. 2022a, 2022c, 2023c), we temporarily refer to the multiple genotype fungi as *O. sinensis*, including Genotype #1 *H. sinensis*, and we continue to refer to the wild and cultivated insect-fungal complexes as *C. sinensis*; however, this practice only partially ameliorates the academic confusion arising from the

indiscriminate use of Latin names and will likely be replaced by the exclusive use of distinct Latin names in the future.

Wei *et al*. (2006) hypothesized that the GC-biased Genotype #1 *H. sinensis* is the sole anamorph of the sole teleomorphic *O. sinensis*. However, numerous studies have identified 17 mutant genotypes of *O. sinensis* in natural *C. sinensis*, which are indiscriminately referred to by the same Latin name (Kinjo and Zang 2001; Stensrud *et al*. 2007; Xiao *et al*. 2009; Zhang *et al*. 2009, 2013a; Zhu *et al*. 2010; Yao & Zhu 2016; Li *et al*. 2020b, 2022a, 2022c, 2023b, 2023c). Stensrud *et al*. (2007) analyzed 71 ITS sequences that were annotated as *C. sinensis* (renamed *O. sinensis* by Sung *et al*. 2007) in GenBank under taxid:72228 and confirmed the existence of 3 phylogenetic clades of mutant *O. sinensis* sequences using a Bayesian phylogenetic algorithm: Group A (GC-biased Genotype #1) and Groups B□C (AT-biased Genotypes #4□5). The authors reported that the variation in the 5.8S gene sequences of *O. sinensis* in Groups A□C "far exceeds what is normally observed in fungi … even at higher taxonomic levels (genera and family)" and that the genotypes "share a common ancestor" and originated from "accelerated nrDNA evolution". Zhang *et al*. (2013a) conducted a GenBank data mining study involving 397 ITS sequences under the name *C. sinensis* or *O. sinensis*, followed by phylogenetic analysis using a minimum evolutionary algorithm, and confirmed the existence of 3 phylogenetic clades of mutant *O. sinensis* sequences: Clade A (GC-biased genotypes) and Clade B□C (AT-biased Genotypes #4□5). In contrast to the homogenous GC-biased Group-A sequences containing only Genotype #1 reported by Stensrud *et al*. (2007), the GC-biased Clade A reported by Zhang *et al*. (2013a) was quite divergent, containing the sequences of Genotype #1 *H. sinensis* and several other mutant GC-biased sequences that exhibited long phylogenetic distances to GC-biased Genotype #1, making it questionable whether they truly belonged to Genotype #1.

Kinjo and Zang (2001) and Mao *et al*. (2013) reported that *O. sinensis* genotypes share the same *H. sinensis*-like morphological and growth characteristics. Xiao *et al*. (2009) concluded that the genetically variable genotypic sequences likely belonged to independent *O. sinensis* fungi based on the differential coexistence of multiple *O. sinensis* mutants in the stroma and caterpillar body of natural *C. sinensis* in different maturation stages. In the debate regarding "cryptic species" (Stensrud *et al*. 2007) and independent fungi of *O. sinensis* (Xiao *et al*. 2009), Zhang *et al*. (2013a) supported the latter and disproved the former based on the results of Kimura 2-parameter (K2P) analysis. Wei *et al*. (2016) reported the existence of the sole GC-biased Genotype #1 teleomorph of

O. sinensis in natural *C. sinensis*. However, they reported the sole AT-biased Genotype #4

teleomorph of *O. sinensis* in cultivated *C. sinensis*, contradicting the anamorphic inoculants of GC-

biased *H. sinensis* strains. Li *et al*. (2023a) reported the differential occurrence of the mating-type

genes of the *MAT1-1* and *MAT1-2* idiomorphs in the *H. sinensis* genome and the differential

transcription and silencing of the mating-type genes, which invalidated the "self-fertilization"

hypothesis under homothallism (Hu *et al*. 2013) and pseudohomothallism (Bushley *et al*. 2013) and

instead suggested that the self-sterility of *O. sinensis* requires sexual partners to accomplish sexual

reproduction via physiological heterothallism or hybridization. In addition to identifying the

molecular heterogeneity of genotypic *O. sinensis* fungi and the requirement of sexual partners for *O.*

sinensis reproduction, Figure 3 of (Bushley *et al*. 2013) illustrates the multicellular heterokaryotic

structure of *C. sinensis* hyphae and ascospores, which include multiple mononucleated, binucleated,

trinucleated, and tetranucleated cells. Zhang and Zhang (2015) suggested that the various

heterokaryotic cells of *C. sinensis* hyphae and ascospores likely contain different genetic materials.

The molecular heterogeneity and self-sterility features of *O. sinensis* and the multicellular

heterokaryotic structure of natural *C. sinensis* challenge the sole anamorph hypothesis for *H.*

sinensis (GC-biased Genotype #1 of *O. sinensis*). Responding to this academic challenge, Li *et al*.

(2013) proposed the "ITS pseudogene" hypothesis for all AT-biased genotypes "in a single genome"

of GC-biased Genotype #1 *H. sinensis* based on (1) the identification of heterogeneous ITS

sequences of both GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis* in 8 out of 15

clones following 25 days of liquid incubation of *C. sinensis* mono-ascospores and (2) the

identification of the 5.8S gene cDNA of Genotype #1 but not of Genotype #5 in a cDNA library of

Strain 1220, one of 8 genetically heterogeneous, ascosporic clones. According to this "ITS

pseudogene" hypothesis, the simultaneous identification of homogenous Genotype #1 in 7 other

clones derived from the same mono-ascospores was apparently ignored. The authors

overgeneralized the "ITS pseudogene" hypothesis to all AT-biased Genotypes #4, #6 and #15□17 of

O. sinensis based on insufficient evidence while attempting to justify AT-biased genotypes as

pseudogenes in the single genome of GC-biased *H. sinensis* under the sole anamorph hypothesis for

H. sinensis (Wei *et al*. 2006). Li *et al*. (2020c) further postulated that all AT-biased *O. sinensis*

genotypes "could have emerged either before or after" a new GC-biased "ITS haplotype was

generated" through "repeat-induced point mutations (RIP)" that cause C-to-T (cytosine-to-thymine)

and G-to-A (guanine-to-adenine) transitions and cooccur as AT-biased, nonfunctional repetitive ITS

copies in "a single genome" of GC-biased *H. sinensis*.

Because of the unavailability of pure cultures of AT-biased *O. sinensis* genotypes for taxonomic

determination, which prevents genomic, transcriptomic, and multigene/multilocus analyses of

hypothetical pseudogenes, the present study included analysis of relevant genomic and

transcriptomic sequences of GC-biased Genotype #1 available in GenBank to examine the "ITS

pseudogene" hypothesis and the extended hypothesis that AT-biased genotypes formed after a new

GC-biased "ITS haplotype was generated" in "a single genome" of *H. sinensis* through "RIP"

mutagenesis. We also extended the single ITS locus analysis to genomic multilocus analysis for the

repetitive copies of 104 of 1271 authentic *H. sinensis* genes. We discuss the genetic and

phylogenetic characteristics of multiple repetitive copies (including repetitive ITS copies) in the

genome of Genotype #1 of *O. sinensis*, the genomic independence of AT-biased genotypes, and

whether sufficient evidence exists to conclude that the 5.8S genes of AT-biased genotypes are

permanently nonfunctional pseudogenes.

II. Materials and Methods

II-1. Reagents

Common laboratory reagents such as agarose, electrophoresis reagents, nylon membranes, and

blocking solution were purchased from Beijing Bioland Technology Company. The endonucleases

*Ava*I, *Dra*I, and *Eco*RI are products of New England BioLabs, United States. Mercuric chloride

(0.1%) for surface sterilization of freshly collected *C. sinensis* specimens was a gift from the

Institute of Microbiology, Chinese Academy of Sciences. The Universal DNA Purification Kit was

obtained from TIANGEN BIOTECH Company (China). The DNeasy Plant Mini Kit was obtained

from the Qiagen Company, Germany. The Gel Extraction Kit was obtained from Omega Bio-Tek

(United States). The *Taq* PCR Reagent Kit and Vector NTI Advance 9 software were purchased

from Invitrogen (United States). All primers used were synthesized by Invitrogen Beijing Lab. or

Beijing Bomaide Technology Co. The DIG High Prime DNA Labeling and Detection Kit was

purchased from Roche Diagnostics (Indiana, USA).

II-2. Information on fungal species in natural *C. sinensis*

171   Information regarding intrinsic fungal species in natural *C. sinensis* was summarized by Jiang

172   and Yao (2013), and subsequent discoveries of additional species among natural *C. sinensis*

173   specimens were reviewed by Li *et al*. (2016b, 2022a, 2022c). The sequences of 17 genotypes of *O.*

sinensis have been obtained and uploaded to GenBank by many *O. sinensis* research groups worldwide since 2001. These genotypic sequences were analyzed genetically and phylogenetically by Kinjo and Zang (2001); Stensrud *et al*. (2007); Xiao *et al*. (2009); Zhang *et al*. (2009, 2013a); Zhu *et al*. (2010, 2012); Chen *et al*. (2011); Zhu & Wu (2015); Li *et al*. (2016b, 2022a. 2022c, 2023c); and Zhu & Li (2017).

II-3. Genetic, genomic, and transcriptomic sequences from GenBank

Five genome assemblies, ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000, for *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively, are available in GenBank (Hu *et al*. 2013; Shu *et al*. 2020; Li *et al*. 2016a; Jin *et al*. 2020; Liu *et al*. 2020) and were used for the analysis of repetitive copies of authentic genes in the ITS locus and multiple loci and multiple GC- and AT-biased *O. sinensis* genotypes.

The GenBank database includes at least 668 *O. sinensis* ITS sequences of *O. sinensis* under the GenBank taxid:72228, 440 of which belong to GC-biased Genotypes #1□3 and #7□14 and 228 of which belong to AT-biased Genotypes #4□6 and #15□17 (Stensrud *et al*. 2007; Zhang *et al*. 2009, 2013a; Chen *et al*. 2011; Li *et al*. 2016b, 2022c, 2023b, 2023c).

The mRNA transcriptome assembly GCQL00000000 for the *H. sinensis* Strain L0106 was uploaded to GenBank by Liu *et al*. (2015). Two metatranscriptome studies of natural *C. sinensis* have been reported. Xiang *et al*. (2014) uploaded the GAGW00000000 metatranscriptome assembly to GenBank, which was obtained from natural specimens (unknown maturational stages) collected from Kangding County of Sichuan Province, China. Xia *et al*. (2017) uploaded the unassembled SRR5428527 Shotgun Read Archive (SRA) to GenBank for mature *C. sinensis* specimens collected from Deqin County of Yunnan Province, China; however, the assembled metatranscriptome sequences were uploaded to www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas, which is currently inaccessible, but a previously downloaded file was used for analysis.

GenBank does not provide annotations for authentic genes in the genome assemblies (ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000) of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively. One (JAACLJ010000002) of the 13 genome contigs for *Ophiocordyceps camponoti-floridani* Strain EC05 contained 1271 authentic genes (Will *et al*. 2020), which was used to cross-reference, position,

205 and annotate the authentic *H. sinensis* genes and their repetitive copies in the *H. sinensis* genome.

206 The transcription of the authentic *H. sinensis* genes and their repetitive genomic copies was
207 explored against the mRNA transcriptome assembly GCQL00000000 for the *H. sinensis* Strain
208 L0106 using the GenBank BLAST algorithms (https://blast.ncbi.nlm.nih.gov/blast/).

II-4. Genetic and phylogenetic analyses

The ITS sequences of multiple genotypes of *O. sinensis* and multiple repetitive genomic ITS
copies as well as metatranscriptomic/transcriptomic sequences were analyzed using the GenBank
BLAST algorithms (https://blast.ncbi.nlm.nih.gov/blast/). The DNA sequences and transcripts of
authentic *H. sinensis* genes and their repetitive copies in the *H. sinensis* genome identified during
multilocus analysis were also analyzed using BLAST algorithms.

A Bayesian phylogenetic tree of the ITS sequences and their repetitive copies was inferred
using MrBayes v3.2.7 software (Markov chain Monte Carlo [MCMC] algorithm) based on 3×10^4
samples, with a sampling frequency of 10^3 iterations after discarding the first 25% of samples from
a total of 4×10^6 iterations (Huelsenbeck and Ronquist 2001; Li *et al*. 2022a, 2023c). This
phylogenetic analysis was performed at Nanjing Genepioneer Biotechnologies Co.

II-5. Collection and on-site processing of fresh *C. sinensis*

Fresh *C. sinensis* specimens were purchased from a local market in Kangding County of
Sichuan Province, China (3800□4600 m' altitude; latitude 30°049'N, longitude 101°959'E) (Zhu *et*
al. 2010; Yao *et al*. 2014). Immature *C. sinensis* specimens were characterized by a plump
caterpillar body and very short stroma (1.0□2.0 cm). Maturing *C. sinensis* specimens were
characterized by a plump caterpillar body and stroma 2.5□5.0 cm in length, without the formation
of an expanded fertile portion close to the stromal tip. Mature *C. sinensis* specimens were
characterized by a plump or slightly less plump caterpillar body and long stroma (>5.0 cm) and by
the formation of an expanded fertile portion close to the stromal tip, which was densely covered
with numerous ascocarps (Zhu *et al*. 2010; Li *et al*. 2023a). Governmental permission was not
required for *C. sinensis* purchases in local markets, and the collections of *C. sinensis* specimens
from sales by local farmers fall under governmental regulations for traditional Chinese herbal
products.

Freshly collected *C. sinensis* specimens were washed thoroughly in running water with gentle
brushing and surface sterilized in 0.1% mercuric chloride for 10 min, followed by washing 3 times
with sterile water (Zhu *et al*. 2010; Li *et al*. 2023c). The thoroughly cleaned and sterilized *C.*

sinensis specimens were immediately frozen in liquid nitrogen and kept frozen during transportation
and storage prior to further processing.

II-6. Extraction of genomic DNA and preparation of *C. sinensis* DNA

239 Caterpillar bodies and stromata of sterilized, frozen *C. sinensis* were separately ground into
240 powder in liquid nitrogen. Genomic DNA was individually extracted from the powder samples
241 using a DNeasy Plant Mini Kit.

II-7. PCR amplification and amplicon sequencing

The genomic DNA of immature, maturing and mature stromata of natural *C. sinensis* was used
as the template for PCR amplification. *H. sinensis*-specific primers were designed (forward primer,
Hsprp1: 5' ATTATCGAGTCACCACTCCCAAACCCCC 3'; reverse primer, *Hsprp3*: 5'
AGGTTCTCAGCGAGCTA 3') based on alignment analysis of the ITS sequences (AB067721,
AB067740, and AB067744, representing Genotypes #1 and #4□5 of *O. sinensis*, respectively).

A touch-down PCR protocol was used, in which the annealing temperature was initially set at
70°C and was subsequently decreased stepwise with a decrement of 0.3°C every cycle, for a total of
36 cycles (Zhu *et al*. 2007, 2010). The PCR amplicons were subjected to agarose gel electrophoresis,
and the DNA bands were recovered, purified using a gel extraction kit, and sequenced either
directly or after cloning the amplicons.

II-8. Preparation of the fungus-specific ITS1 probe and 18S internal control probe

An *H. sinensis*-specific probe (158 bp) was designed based on the ITS1 sequence of GC-biased
Genotype #1 *H. sinensis* and generated through PCR using the forward primer *Hsprp1* described
above and the reverse primer *Hsprp2* (5' ATTTGCTTGCTTGACTGAGAGATGCC 3').

A nonspecific 18S internal control probe was designed based on alignments of GenBank 18S
rDNA sequences (DQ838795, AB187268, AB067735, AB067736, and AB080090) of several *C.*
sinensis-associated fungi. A region highly homologous to the 18S rDNA region, located more than
700 bp upstream of the ITS1 region of *C. sinensis*-associated fungi, was selected. The 18S control
probe was generated through PCR using the forward primer *Icprp1* (5'
ACCGTCGTAGTCTTAACCAT 3') and the reverse primer *Icprp2* (5'
ATCGGCTTGAGCCGATAGTC 3').

Both amplicons were confirmed by sequencing, and the probes were labeled using the DIG

High Prime DNA Labeling and Detection Kit.

II-9. Southern blotting analysis

The stromata and caterpillar bodies of immature and mature *C. sinensis* were subjected to
Southern blotting. Genomic DNA was extracted from the samples and prepared by endonuclease
digestion using *Ava*I, *Dra*I, and *Eco*RI. The digested genomic DNA (5 µg of each sample) was
subjected to agarose gel electrophoresis and transferred onto a nylon membrane through capillary
transfer, followed by baking the membrane at 58°C for 4 hours for fixation, blocking the binding
sites on the membrane in blocking solution, and hybridizing with the labeled *H. sinensis*-specific
probe at 40°C overnight. The hybridized transblotted membrane was viewed by colorimetric
photography using a gel imaging system (Bio-Rad) and then stripped and reprobed with the labeled
18S internal control probe.

III. The results

III-1. Multiple GC-biased repetitive ITS copies in the *H. sinensis* genome

In general, fungal genomes contain multiple repetitive copies of the nrDNA ITS1-5.8S-ITS2
sequences. Li *et al*. (2020c) submitted their paper to the IMA Fungus in January 2020, when 3
genome assemblies (ANOV00000000, LKHE00000000, and LWBQ00000000) of *H. sinensis*
Strains Co18, 1229, and ZJB12195, respectively (Hu *et al*. 2013; Li *et al*. 2016a; Jin *et al*. 2020),
were available in GenBank. However, only single copies of the ITS1-5.8S-ITS2 sequences could be
identified in each of the genomes (ANOV01021709, LKHE01000582, and LWBQ01000008) of the
strains, probably because the repetitive ITS copies were discarded during the assembly of the
genome shotgun reads, which indicated that Li *et al*. (2013, 2020c) might not have been able to
obtain real genomic data for repetitive ITS copies before paper submission as the basis for the
maternal backup of their "ITS pseudogene" hypotheses. On the other hand, multiple repetitive ITS
copies were identified in 2 additional genome assemblies, JAAVMX000000000 and NGJJ00000000,
for *H. sinensis* Strains IOZ07 and CC1406-203, respectively (Liu *et al*. 2020; Shu *et al*. 2020),
which were released in GenBank after June 2020. Seventeen repetitive ITS1-5.8S-ITS2 copies were
found within JAAVMX000000000, 7 ITS1-5.8S-ITS2 copies within NGJJ00000000 (Table 1), and
2 partial ITS segments within NGJJ00000000 were not included in our analysis.

Most of the repetitive ITS sequences listed in Table 1 were highly homologous (98.8–100%) to
AB067721, the reference sequence of GC-biased Genotype #1 *H. sinensis*. Two repetitive copies

(6233→6733 and 44729→45251) within JAAVMX010000019 showed 94.5% and 90.8% similarity

to AB067721, respectively (Table 1). Two other copies, 18702095→18702586 within

JAAVMX010000002 and 700→1186 within JAAVMX010000018, showed similarities of 97.4%

and 97.0%, respectively, to AB067721. The repetitive ITS sequences showed 80.1□89.9%

similarity to the AT-biased Genotype #4□6 and #15□17 sequences of *O. sinensis* (Table 1).

Table 1. GC contents of the repetitive ITS1-5.8S-ITS2 sequences from the genome assemblies of 2 H. sinensis *strains and percent similarities of the repetitive ITS sequences to the sequences of GC-biased Genotype #1 (AB067721) and AT-biased Genotypes #4□6 and #15□17 of* O. sinensis.

Genome segment	Sequence range & direction	GC content	% Similarity of the repetitive sequence	
			vs. Genotype #1 AB067721	*vs*. AT-biased Genotypes #4□6 and #15□17
JAAVMX010000002	18688917→18689407	64.8%	100%	86.7 □ 89.9%
	18702095→18702586	64.5%	97.4%	82.8 □ 88.0%
JAAVMX010000008	13823→14313	64.8%	100%	85.5 □ 89.9%
	1199→1687	64.8%	99.2%	85.2 □ 89.3%
JAAVMX010000017	9147←9637	64.8%	100%	85.5 □ 89.9%
	21791←22281	64.8%	100%	85.5 □ 89.9%
	34435←34925	64.8%	100%	85.5 □ 89.9%
	47079←47569	64.8%	100%	85.5 □ 89.9%
JAAVMX010000018	13381→13871	64.8%	100%	85.5 □ 89.9%
	26076→26566	64.8%	100%	85.5 □ 89.9%
	38771→39261	64.8%	100%	85.5 □ 89.9%
	51467→51958	64.6%	98.8%	85.5 □ 88.8%
	700→1186	64.5%	97.0%	82.6 □ 87.6%
JAAVMX010000019	19404→19894	64.8%	100%	85.5 □ 89.9%
	32048→32538	64.8%	100%	85.5 □ 89.9%
	6233→6733	65.3%	94.5%	81.3 □ 85.5%
	44729→45251	63.3%	90.8%	80.1 □ 84.6%
NGJJ01000573	13133←13623	64.8%	99.4%	85.2 □ 89.6%
	14624←15113	64.7%	99.4%	84.8 □ 89.6%
NGJJ01000582	10703←11192	64.7%	99.8%	85.5 □ 89.8%
NGJJ01000796	1514→2004	64.8%	100%	85.5 □ 89.9%
NGJJ01000798	1177→1665	64.6%	99.6%	84.8 □ 89.6%
NGJJ01000799	3295→3785	64.8%	100%	85.5 □ 89.9%
	16666→17156	64.8%	100%	85.5 □ 89.9%

Note: AB067721 represents GC-biased Genotype #1. The representative sequences for AT-biased genotypes are Genotype #4 AB067744, Genotype #5 AB067740, Genotype #6 KJ720572, Genotype #15 KT232017, Genotype #16 KT232019, Genotype #17 KT232010.

III-2. Genetic characteristics of the multiple repetitive ITS sequences in the *H. sinensis* genome

The repetitive ITS copies within genome assemblies JAAVMX000000000 and NGJJ00000000

for the *H. sinensis* Strains IOZ07 and CC1406-203, which are listed in Table 1, and the single

genomic ITS copies within ANOV01021709, LKHE01000582, and LWBQ01000008 for the *H.*

sinensis Strains Co18, 1229, and ZJB12195, respectively, which are listed in Table 2 (Hu *et al*. 2013;

Jin *et al*. 2020; Li *et al*. 2016a; Liu *et al*. 2020; Shu *et al*. 2020), were GC biased; their average GC

content was 64.7±0.33% (Table 1), nearly identical to the GC content (64.8%) of the AB067721

Genotype #1 sequence but significantly greater (*P*<0.001) than 51.1±1.69% (44.8□53.1%) for the

AT-biased Genotypes #4□6 and #15□17 (Li *et al*. 2023c). Apparently, these genetic features of the

variable repetitive ITS copies were distinct from those of AT-biased *O. sinensis* genotypes, which

included multiple scattered transition alleles.

Table 2. *Percent similarities between nrDNA ITS1-5.8S-ITS2 sequences of 5 genome assemblies of*
Genotype #1 H. sinensis *and 17* O. sinensis *genotype sequences. (Modified from Table 1 of* Li et al.
2023c).

Genotype # and GenBank accession #	Percent similarity between the *H. sinensis* genomic ITS segments and 17 *O. sinensis* genotypes				
	ANOV01021709 838→1504	LKHE01000582 2074→2740	LWBQ01000008 991739→992405	JAAVMX010000017 9029←9695	NGJJ01000799 3237→3903
#1 AB067721	99.7%	100%	99.6%	100%	100%
#2 MG770309	94.4%	94.7%	94.7%	94.7%	94.7%
#3 HM595984	95.9%	96.1%	95.6%	95.3%	95.3%
#7 AJ488254	94.7%	95.0%	95.3%	95.0%	95.0%
#8 GU246286	89.5%	89.5%	89.2%	89.5%	89.5%
#9 GU246288	95.0%	95.2%	94.6%	95.2%	95.2%
#10 GU246287	83.0%	83.2%	83.0%	83.2%	83.2%
#11 JQ695935	78.9%	79.1%	78.5%	79.1%	79.1%
#12 GU246296	94.3%	94.5%	93.9%	94.5%	94.5%
#13 KT339190	87.5%	87.5%	87.2%	87.5%	87.5%
#14 KT339178	89.3%	89.6%	89.1%	89.6%	89.6%
#4 AB067744	89.1%	89.4%	89.3%	89.4%	89.4%
#5 AB067740	86.3%	86.7%	86.2%	86.7%	86.7%
#6 KJ720572	85.5%	85.5%	85.5%	85.5%	85.5%
#15 KT232017	89.6%	89.9%	89.4%	89.9%	89.9%
#16 KT232019	87.4%	87.7%	87.6%	87.7%	87.7%
#17 KT232010	87.7%	88.1%	87.5%	88.1%	88.1%

323   *Note: "GT" represents the genotype. GC-biased genotypes are listed in blue in the upper panel. AT-*
324   *biased genotypes are listed in red in the lower panel. The GenBank accession numbers are*
325   *hyperlinked to the GenBank database.*

The sequences of GC-biased repetitive ITS copies that were variable (94.5% and 90.8% similarity) in comparison to the reference sequence AB067721 of Genotype #1 predominantly contained insertion/deletion alleles and included less frequent transversion alleles, while only a few transition alleles were found (Table 3). The variable ITS repeats 6233→6733 and 44729→45251 within JAAVMX010000019 (*cf*. Table 1) contained 22 and 32 insertion/deletion alleles and 4 and 11 transversion alleles, respectively (Table 3; in blue in Figure 1); they contained only 1 and 5 transition mutant alleles, having allelic ratios of insertion/deletion/transversion *vs*. transition of 26:1 and 43:5, respectively. The sequences of repetitive ITS copies, 18702095→18702586 within JAAVMX010000002 and 700→1186 within JAAVMX010000018 (*cf*. Table 1), were less variable (97.4% and 97.0% similarity to AB067721) and contained no transition alleles and predominantly included 13 and 15 insertion, deletion and transversion alleles, with allelic ratios of insertion/deletion/transversion *vs*. transition of 13:0 and 15:0, respectively (Table 3; in green in Figure 1). These genetic characteristics of the variable and less variable repetitive ITS copies indicate that they might not have been generated through RIP mutagenesis, which theoretically causes cytosine-to-thymine (C-to-T) and guanine-to-adenine (G-to-A) transition mutations. Other repetitive ITS copies with 98.1□99.8% similarity to AB067721 listed in Table 1 contained only a few insertion/deletion alleles.

Table 3. *Types of mutations in the variable and slightly less variable repetitive ITS sequences compared with those of GC-biased Genotype #1 (AB067721).*

	Sequence range & direction	% Similarity	Number of mutant alleles			Ins./Del./Transv. *vs*. Transit.
			Ins./Del.	Transv.	Transit.	
JAAVMX010000002	18702095→18702586	97.4%	13	0	0	13 : 0
JAAVMX010000018	700→1186	97.0%	10	5	0	15 : 0
JAAVMX010000019	6233→6733	94.5%	22	4	1	26 : 1
	44729→45251	90.8%	32	11	5	43 : 5

Note: "Ins./Del." indicates insertion/deletion point mutations. "Transv." denotes Transversion point mutations. "Transit." refers to transition point mutations.

```
GT#1  AB067721  59       TCGAGTCACCACTCCC AAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCCGGCG CCCCAGGGCCCG 136
JAAVMX010000019 19404    ---------------- ------------------------------------------------- ------------ 19481
JAAVMX010000019 32048    ---------------- ------------------------------------------------- ------------ 32125
JAAVMX010000002 18702095 ---------------- ------------------------------------------T------------ 18702172
JAAVMX010000018 700      ---------- ------ -----------------------------T----------- ----------T 776
JAAVMX010000019 6233     ----------------C- ----------------------------------------- ------------ 6311
JAAVMX010000019 44729    ---------------- ------------------------------------------------- ------------ 44806
GT#4  AB067744  55       ------T-------T --------------T--T------------------------- ---T------ 132
GT#5  AB067740  59       ------T--T----TT ---------ATA-T--T-T-----------TA-----------TA--- ---T-A------ 136
GT#6  KJ720572  11              -------A----T--T-T-----------TA------------TA--- --TT-A------ 70
GT#15 KT232017  37       ------T-------T --------------T--T------------------------- ---T------- 114
GT#16 KT232019  37       ------T--T-----T --------------T--T-T-----------TA-----------TA--- ---T-A------ 114
GT#17 KT232010  37       ------T--T-----T --------------T--T-T-----------TA------------TA--- ---T-A------ 114

GT#1  AB067721  137      GACCAGG GCGCCC GCCGGAGGACCCCCAGAC CCTCCTGTCGCA GTGGCATCT CTCAGTCAAGAA GCAAGCAAAT 210
JAAVMX010000019 19482    ------- ------ ----------------- ------------ --------- ------------ ---------- 19555
JAAVMX010000019 32126    ------- ------ ----------------- ------------ --------- ------------ ---------- 32199
JAAVMX010000002 18702173 ------- ------A---------------------------A--------- ---- ------------ ---------- 18702247
JAAVMX010000018 777      ------- ------ ----------------- ------------ ---C--------- ---------- 850
JAAVMX010000019 6312     ------- ------ ------T----------C------------ ---------C-----------G--------- 6388
JAAVMX010000019 44807    -------C----- ----------------- ------------ --------- ------------ ---------- 44881
GT#4  AB067744  133      ---T--- ------ -----------TT---- ------A---T- --A------ --T---T----- ---------- 206
GT#5  AB067740  137      ---T-AA ------ -----------TT---- ------A-TAT- --A-T---- --T---T----- ---------- 213
GT#6  KJ720572  71       ---T--- ------ -----------TT---- ------A---T- --A-T---- --T---T----- ---------- 182
GT#15 KT232017  115      ---T--- ------ -----------TT---- ------A----- --A------ --T---T----- ---------- 188
GT#16 KT232019  115      ---T-AA ------ -----------TT---- ------A----- --A-T---- --T---T----- ---------- 188
GT#17 KT232010  115      ---T-AA ------ -----------TT---- ------A-TAT- --A-T---- --T---T----- ---------- 188

GT#1  AB067721  211      GAATCAAAACTTTCAACAACGGATCTCTTGGTTCTGGCATC GATG AAGAACGCAGCGAAATGCGATAAGTAATGTGAA 288
JAAVMX010000019 19556    ----------------------------------------- ---- -------------------------------- 19633
JAAVMX010000019 32200    ----------------------------------------- ---- -------------------------------- 32277
JAAVMX010000002 18702248 -- -------------------------------------C-- ---- -------------------------------- 18702325
JAAVMX010000018 851      ----------------------------------------- ----A--------------------- --------- 928
JAAVMX010000019 6389     ------------- ---CAC----------------------- --------------------------- ------- 6463
JAAVMX010000019 44882    ----------------------------------------- ---- -------------------------------- 44959
GT#4  AB067744  207      A---T--------T--T-----------A---T-A----T A--A -----------------A-------------A-- 284
GT#5  AB067740  214      A---T--------T----------A----A----T A--A --A---------------A--------A-A-- 288
GT#6  KJ720572  183      A---T--------T-----------A----A--- ---A ------T--T-----A-----------A-A-- 222
GT#15 KT232017  189      ----T--------T--T-----------A---T-A----T A--A -----------------A----------A-- 266
GT#16 KT232019  189      ----T--------T----------A----A--- ---A --A------------------A--------A-A-- 266
GT#17 KT232010  189      A---T--------T-----------A----A--- ---A --A---------------A--------A-A-- 266

GT#1  AB067721  289      TTGCAGAATTCAGTGAACCA TCGAATCTTT GAACG CACATTG CGCCCGCCAGCACTCTGGCGGGCATGC  CTGTC 362
JAAVMX010000019 19634    -------------------- ---------- ----- ------- -------------------------- ----- 19707
JAAVMX010000019 32278    -------------------- ---------- ----- ------- -------------------------- ----- 32351
JAAVMX010000002 18702326 -------------------- ----------T----- ----- - ------- --------------- ----- 18702398
JAAVMX010000018 929      -------------------- C----------- -----C-------C-------------- ------- ----- 1000
JAAVMX010000019 6464     -------------------C----------- ----- -------C------------------ ------- ----- 6539
JAAVMX010000019 44960    -------------------- ---------- ----- ------- -------------------------CT----- 45035
GT#4  AB067744  285      ---T------T---A---T- ---------- A---- T-T-C-A -------T--------A--------- ----- 358
GT#5  AB067740  289      --A--------T---A---T- ---------- A---- --T---A -------T--------A--------- --A-- 362
GT#6  KJ720572  223      --AT------T---A----- ---------- A---- T-T---A -------T--------A--------- ----- 296
GT#15 KT232017  267      ---T------T---A--T- ---------- A---- T-T---A -------T--------A--------- ----- 340
GT#16 KT232019  267      --A--------T---A---T- ---------- A---- T-T---A -------T--------A--------- ----- 340
GT#17 KT232010  267      --A--------T---A--T- ---------- A---- --T---- -------T--------A--------- --A-- 340

GT#1  AB067721  363      CGAGCGTCATCTCAACCCTCGAG CCCCCCGCCTCGCGG  CGGCGGGGCCCGGCCTT GGGGGTCACGGCC CCGCGCC 437
JAAVMX010000019 19708    ----------------------- ---------------  ----------------- ------------- ------- 19781
JAAVMX010000019 32352    ----------------------- ---------------  ----------------- ------------- ------- 32425
JAAVMX010000002 18702399 ----------------------- ----- ---------  ----------------G------------- ------- 18702472
JAAVMX010000018 1001     ----------------------- ------ --------  ----------------G------------- ---- 1074
JAAVMX010000019 6540     ----------------------- ---------------  ----------------- ---------C----- 6613
JAAVMX010000019 45036    --------------------C----------A-----TA--CG---C--------- ------TC-----AA------ 45119
GT#4  AB067744  359      -------T-C--T------TA-- ---------------  ----------------- A---------- --A---- 432
GT#5  AB067740  363      -------T----T------- ---------------  ----------T------- A-----T------ ------- 436
GT#6  KJ720572  297      -------T-C--T---- 313
GT#15 KT232017  341      -------T----T---------- ---------------  ----------T------- A-----T------ ------- 414
GT#16 KT232019  341      -------T-C--T-----TA-- ---------------  ----------------- A---------- --A---- 414
GT#17 KT232010  341      -------T----T---------- ---------------  ----------T------- A-----T------ ------- 414

GT#1  AB067721  438      G  CCCCCTAAACGC AG     TGGCGACC  CCGCCGCGGCTCC CCTGCGCAGTAGCT  CG  CTGAGAACCT 499
JAAVMX010000019 19782    -  ------------ --     --------  ------------- ------------ --  ---------- 19844
JAAVMX010000019 32426    -  ------------ --     --------  ------------- ------------ --  ---------- 32488
JAAVMX010000002 18702473 -  ------------ --     --------  ------------- ------------ --  ---------- 18702534
JAAVMX010000018 1075     -  ------------ --     --------  ------------- ------------ --  ---------- 1136
JAAVMX010000019 6614     -  -----------G--G     --------GC------------ ------------ --GCT--------- 6680
JAAVMX010000019 45120    -TCC-T---GT----- --CCTAAGTG------- ----GA-------A----T-A-------CA-- ---------- 45189
GT#4  AB067744  433      -  ------------ --     -A-T----  ------------ ---A---A----- TA  --A------- 495
GT#5  AB067740  437      -  ----T------T --     -A------  ------------ ---A--TG------  -A  -C------- 529
GT#6  KJ720572  313      -                                                                         313
GT#15 KT232017  415      -  -----------T --     -A------  ------------ ---A--T-A-----  -A  -C------- 477
GT#16 KT232019  415      -  ------------ --     -A------  ------------ ---A--A------ TA  --A------- 477
GT#17 KT232010  415      -  -----------T --     -A------  ------------ ---A--T-A----  -A  -C------- 477

GT#1  AB067721  500      CGCACCGGGAGCGCGGA     GGCGG TCACGCC  TGG   AA ACCACC   ACACCCTCCA           549
JAAVMX010000019 19845    -----------------     ----- ----- - ---   -- ------   ----------           19894
JAAVMX010000019 32489    -----------------     ----- ----- - ---   -- ------   ----------           32537
JAAVMX010000002 18702535 -----------------     ----- -------GTGT-   -- ------   ----------           18702586
JAAVMX010000018 1137     -----------------     ----- ----- - ---   -- ------   ----------           1186
JAAVMX010000019 6681     C----------------     ----- ----- - ---   -- ------ACA----------           6733
JAAVMX010000019 45190    -----------------CCCAT-----C--------   ---GATG--C-----ACC---------           45251
GT#4  AB067744  496      -----TA----------     ----- -------  --A   -- -----T   -T------           543
GT#5  AB067740  530      --------A---------     ----- -------  --A   -- -----T   --G--             544
GT#6  KJ720572  313                                                                                313
GT#15 KT232017  478      --------A---------     ----- -------  --A   -- -----T   --G--             522
GT#16 KT232019  478      -----TA----------     ----- -------  --A   -- -----T   -T------           525
GT#17 KT232010  478      --------A---------     ----- -------  --A   -- -----T   --G--             522
```

Figure 1. ITS sequence alignment of AB067721, repetitive ITS copies within the genome JAAVMX000000000 of H. sinensis Strain IOZ07, and AT-biased genotypes of O. sinensis.

The ITS sequences contained complete or partial ITS1-5.8S-ITS2 nrDNA segments. "GT" denotes the O. sinensis *genotype. The underlined sequence in black represents the 5.8S gene of GC-biased Genotype #1 H. sinensis. AB067721 is the ITS sequence of GC-biased Genotype #1 H. sinensis. The genome assemblies JAAVMX010000002, JAAVMX010000018, and JAAVMX010000019 were obtained from the* H. sinensis *Strain IOZ07 (Shu et al. 2020). One copy each within JAAVMX010000002 and JAAVMX010000018, indicated in* **green***, shares 97.4% or 97.0% similarity with AB067721. JAAVMX010000019 contains 4 repetitive ITS copies, including 2* **black** *sequences (19404→19894 and 32048→32537), which are 100% identical to AB067721, and 2 other* **blue** *sequences (6233→6733 and 44729→45251), with 94.5% and 90.8% similarity to AB067721. The sequences in* **red** *(AB067744, AB067740, KJ720572, KT232017, KT232019, and KT232010) represent AT-biased Genotypes #4□6 and #15□17 of* O. sinensis*, respectively. The underlined "GAATTC" sequences in* **purple** *represent the EcoRI endonuclease cleavage sites in the sequences of GC-biased Genotype #1 and the GC-biased genome assembly JAAVMX000000000. EcoRI endonuclease cleavage sites are absent in AT-biased ITS sequences due to a single-base cytosine-to-thymine (C-to-T) transition. Hyphens indicate identical bases, and spaces denote unmatched sequence gaps.*

III-3. Phylogenetic features of multiple repetitive ITS copies in the *H. sinensis* genome

As shown in Tables 1□2 and Figure 1, the sequences of GC-biased Genotypes #2–3 and #7–14 and of AT-biased Genotypes #4□6 and #15□17 of *O. sinensis* are absent in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000 of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively, and instead belong to the genomes of independent *O. sinensis* fungi (Kinjo & Zang 2001; Stensrud *et al.* 2007; Xiao *et al.* 2009; Zhang *et al.* 2009, 2013a; Chen *et al.* 2011; Hu *et al.* 2013; Zhu & Wu 2015; Li *et al.* 2016a, 2016b, 2020b, 2022a, 2022c, 2023c; Zhu & Li 2017; Zhu *et al.* 2018, 2019; Jin *et al.* 2020; Liu *et al.* 2020; Shu *et al.* 2020). The genomic repetitive ITS copies were phylogenetically clustered into the GC-biased Genotype #1 clade in blue alongside the Bayesian majority rule consensus tree in Figure 2 and were phylogenetically distant from the AT-biased clades containing Genotypes #4□6 and #15□17 of *O. sinensis* shown in red alongside the tree, similar to what has previously been demonstrated in phylogenetic trees that were conferred using a variety of phylogenetic algorithms (Kinjo & Zang 2001; Stensrud *et al.* 2007; Zhang *et al.* 2013a; Zhu & Wu 2015; Li *et al.* 2016b, 2020b, 2022a, 2022c, 2023c; Wei *et al.* 2016; Zhu & Li 2017).

Figure 2. A Bayesian majority rule consensus phylogenetic tree.

"GT" represents genotype. Twenty-eight ITS segments within the genome assemblies (ANOV01021709, LKHE01000582, LWBQ01000008, JAAVMX010000002, JAAVMX010000008, JAAVMX0100000017, JAAVMX0100000018, JAAVMX010000019, NGJJ01000573, NGJJ01000582, NGJJ01000796, NGJJ01000798, and NGJJ01000799) of H. sinensis *strains (Co18, 1229, ZJB12195, IOZ07, and CC1406-203, respectively) and 25 ITS sequences of GC-biased Genotypes #1–3 and #7–14 (in blue alongside the tree) and AT-biased Genotypes #4–6 and #15–17 of* O. sinensis *(in red alongside the tree) were analyzed phylogenetically using MrBayes v3.2.7 software (cf. Methods Section II-4). Genome assemblies JAAVMX000000000 and NGJJ00000000 contain multiple repetitive ITS copies (cf. Table 1). The percent similarities of the genomic sequences of repetitive ITS copies with the representative Genotype #1 sequence (AB067721) are shown in green alongside the tree.*

Figure 2 includes the 2 genomic repetitive ITS copies with low similarity (94.5% and 90.8%) and the 2 other repetitive ITS copies with relatively low similarity (97.4% and 97.0%), which are shown in Tables 1 and 3. The repetitive ITS copies are individually scattered within the GC-biased Genotype #1 clade. The 2 low-similarity genomic repetitive ITS sequences, 6233→6733 (94.5% similarity) and 44729→45251 (90.8% similarity), within the JAAVMX010000019 genome assembly (*cf*. Tables 1 and 3), showed greater phylogenetic distances (horizontal lines) from other GC-biased ITS copies within the GC-biased clades. GC-biased Genotypes #2□3 and #7□14 of *O. sinensis* showed greater phylogenetic distances from the sequences of GC-biased Genotype #1 and the repetitive ITS copies. These results indicate that genetic variations in the GC-biased repetitive ITS sequences shown in blue in Figure 1 had a significant impact on the Bayesian phylogenetic clustering analysis.

III-4. Multiple GC-biased genotypes of *O. sinensis* in relation to the number of repetitive genomic ITS copies

In addition to the phylogenetic analysis of multiple genotypes of *O. sinensis* shown in Figure 2, the sequences of GC-biased Genotypes #1□3 and #7□12 were further analyzed (Figure 3) and compared with the genomic repetitive ITS copies. GC-biased Genotypes #13□14 were characterized by large DNA segment reciprocal substitutions and recombination of genetic material between the genomes of the 2 parental fungi, Genotype #1 *H. sinensis* and an AB067719-type fungus, and they were not included in the analysis in Figure 3. Figure 2 shows that GC-biased Genotypes #2□3 and #8□14 formed several GC-biased phylogenetic branches outside the Genotype #1 phylogenetic clade with great phylogenetic distances (horizontal lines), and they were phylogenetically and genetically distinct from the genomic repetitive ITS copies of Genotype #1 *H. sinensis* (*cf*. Table 2 and Figure 3).

```
GT#1   AB067721      59        TCGAGTC ACCACTCCC AAA CCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCCGGCG CCCCAGGGCCCG 136
JAAVMX010000019 19404         ------- --------- --- ---------------------------------------------- ------------ 19481
JAAVMX010000019 32048         ------- --------- --- ---------------------------------------------- ------------ 32125
JAAVMX010000002 18702095      ------- --------- --- --------------------------------------T------- ------------ 18702172
JAAVMX010000018 700           ------- -- ------ --- --------------------T-------------------------- ----------T 776
JAAVMX010000019 6233          ------- --------C-- --- ---------------------------------------------- ------------ 6311
JAAVMX010000019 44729         ------- --------- --- ---------------------------------------------- ------------ 44806
GT#2   MG770309
GT#3   HM595984      33        CT----T G------- --- A----C-T------------------------------------- ----------- 110
GT#7   AJ488254      464       ------- -A----T- G--G------------------------------T-T----- -A---TG--- 380
GT#8   GU246286      4         -----T- --T----- --- ------------------------------T------- --G---T--- 83
GT#9   GU246288      5         ------TT -------- --- -----C---G--------------------------- ----------- 84
GT#10  GU246287      6         -----T- -G ------ T-- ---------------------T----T-----TTTG---------- ----A------ 82
GT#11  JQ695935      12        ------- --------- --- ---------------------------------------------- ----------- 78
GT#12  GU246296      5         ------T --------- --- ---------------------------------------------- ----------- 83

GT#1   AB067721      137       GACCAGG GCGCCC GCCGGAGGACCCCCAGAC CCTCCTGTCGCA GTGGCATCT CTCAGTCAAGAA GCAAGCAAATGA 213
JAAVMX010000019 19482         ------- ------ ----------------- ------------ --------- ------------ ------------ 19557
JAAVMX010000019 32126         ------- ------ ----------------- ------------ --------- ------------ ------------ 32201
JAAVMX010000002 18702173      ------- ------A ---------------- ------------A ---------- ------------ ------------ 18702249
JAAVMX010000018 777           ------- ------ ----------------- ------------ --C------ ------------ ------------ 852
JAAVMX010000019 6312          ------- ------ ------T---------- ---C-------- --------- ------G----- ------------ 6390
JAAVMX010000019 44807         -------C----- ------ ----------------- ------------ --------- ------------ ------------ 44883
GT#2   MG770309      1         ----G AT--ATTATT-A----C- -----C-A----- -------T-- ------------ 68
GT#3   HM595984      111       ---T--- ------ ----------------- ------------ --G------ ------------ ------------ 186
GT#7   AJ488254      379       A------ ------ ----------------- ------------ --------- ------------ ------------ 295
GT#8   GU246286      84        -C-G--- ------T -GAAAT-TT--AA-T-- ------------ --C------ ------------ ------------ 159
GT#9   GU246288      85        ------- ------ ----------------- ------------ --------- ------G- -----G----- 160
GT#10  GU246287      83        ------- ------ ----G------------ ------------G ---GT-T- T--------- AGG--G----- 157
GT#11  JQ695935      79        ------- ------ ----------------- ------------ --------- ------------ ------------ 154
GT#12  GU246296      84        ------- ------ ----------------- ------------ --------- ------G----- ------------ 159

GT#1   AB067721      214       ATCAAAACTTTCAACAACGGATCTCTTGGTTCTGGCATC GATG AAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGC 292
JAAVMX010000019 19557         --------------------------------------- ---- ------------------------------------ 19637
JAAVMX010000019 32201         --------------------------------------- ---- ------------------------------------ 32281
JAAVMX010000002 18702249      ----------------------------------C---- ---- ------------------------------------ 18702329
JAAVMX010000018 852           --------------------------------------- ---- ---A-------------------------------- 932
JAAVMX010000019 6390          ----------- --CAC----------------------- ---- ------------------------------------ 6467
JAAVMX010000019 44883         --------------------------------------- ---- ------------------------------------ 44963
GT#2   MG770309      68        --------------------------------------- ---- ---GA--------C ---------------------- 147
GT#3   HM595984      186       --------------------------------------- ---- ------------------------------------ 266
GT#7   AJ488254      295       --------------------------------------- ---- ----A------------------------------- 224
GT#8   GU246286      159       -----------T----------------T--------- ---- --G--------------G----------------- 239
GT#9   GU246288      161       --------------------------T------------ ---- ------------------------------------ 240
GT#10  GU246287      158       ------T-----------------T--------------- ---- --------C---C----------G----------- 237
GT#11  JQ695935      155       --------------------------------------- ---- ------------------------------------ 234
GT#12  GU246296      160       --------------------------------------- ---- -------------G--------------------- 239

GT#1   AB067721      293       AGAATTCAGTGAACCA TCGAATCTTT GAACG CACATT G CGCCCGCCAGCACTCTGGCGGGCATGC  CTGTCCGAGC 367
JAAVMX010000019 19638         ---------------- ---------- ----- ------ - ------------------------- ---------- 19712
JAAVMX010000019 32282         ---------------- ---------- ----- ------ - ------------------------- ---------- 32356
JAAVMX010000002 18702330      ---------------- --------T----- ------ - ------------------------- ---------- 18702403
JAAVMX010000018 933           ---------------- -C--------- -----C------ --C--------------- ------------ ---------- 1005
JAAVMX010000019 6468          ----------------C--------- ----- --C--- - ------------------------- ---------- 6544
JAAVMX010000019 44964         ---------------- ---------- ----- ------ - ----------------------CT--------- 45040
GT#2   MG770309      148       ---------------- ---------- ----- ------ - --------------A---------- ---------- 222
GT#3   HM595984      267       ---------------- ---------- ----- ------ - ------------------------- ---------- 341
GT#7   AJ488254      223       ---------------- ----A----- ----- ------ - ------------------------- ---------- 147
GT#8   GU246286      240       ---------------- --T-T-T--- ----- ------ - ----------------A-------- ---------- 314
GT#9   GU246288      241       ---------------- ---------- ----- ------ - ----------C-------------- ---------- 315
GT#10  GU246287      238       ---------------- ---------- ----- --G- G---- - ------------------T------ ---------- 313
GT#11  JQ695935      235       ---------------- ---------- ----- ------ - ------------------------- ---------- 309
GT#12  GU246296      240       ---------------- ---------- ----- ------ - ----------G-------------- ---------- 314

GT#1   AB067721      368       GTCATCTCAACCCTCGAG CCCCCCGCCTC GCGG C GGCGGGG CCCGGCCTT GGGGGTCACGGCC CCGC GCCG 438
JAAVMX010000019 19713         ------------------ ----------- ---- - ------- --------- ------------- ---- ---- 19782
JAAVMX010000019 32357         ------------------ ----------- ---- - ------- --------- ------------- ---- ---- 32426
JAAVMX010000002 18702404      ------------------ ----------- ---- - ------- --------- ------------- ---- ---- 18702473
JAAVMX010000018 1006          ------------------ ----------- ---- - ------- --------- ----------G-- ---- --- 1075
JAAVMX010000019 6545          ------------------ ----------- ---- - ------- --------- ----------C--- ---- ---- 6614
JAAVMX010000019 45041         ------------------C---------- --A ----TA- -CG---C ----------- ------TC-----AA--- ----T 45123
GT#2   MG770309      223       ------------------ ----------- ---- - ------- --------- ------------- ---- ---- 293
GT#3   HM595984      342       ------------------ -----A----- ---- - ------- --------- -------T----- ---- ---- 412
GT#7   AJ488254      148       ------------------ ---A-A---T--A -T--- --------- --------- ---- ---- 76
GT#8   GU246286      315       ------------------ ----A------ ---- - ------- -------G------ ---- C--C 387
GT#9   GU246288      316       ------------------ -------G--- ---- - ------- --------- ------------- ---- ---- 388
GT#10  GU246287      314       ----A--A----A----- --------T -G-- T ----------T- T----T--C-- --- C--C 384
GT#11  JQ695935      310       ------------------ ----------- ---- - --------TTCT--- --TT---CTTTAT TGAT -A--- 381
GT#12  GU246296      315       ------------------ ----------- ---- - ------C---------- ------------- --C-CGC--- 388

GT#1   AB067721      439       CCCCC T AAACGC AG TGGCGACC CC GCCGCGGCTCC CC TGCGCAGTAGCT CG CTGAGAAC 497
JAAVMX010000019 19783         ----- - ------ -- -------- -- ----------- -- ------------ -- -------- 19842
JAAVMX010000019 32427         ----- - ------ -- -------- -- ----------- -- ------------ -- -------- 32486
JAAVMX010000002 18702474      ----- - ------ == -------- -- ----------- -- ------------ -- -------- 18702532
JAAVMX010000018 1076          ----- - ------ -- -------- -- ----------- -- ------------ -- -------- 1134
JAAVMX010000019 6614          ----- - ------G--G -------GC-- ----------- -- -----GCT------ 6678
JAAVMX010000019 45125         CCT--- G - ------ --CCTAAGTG------- -- --GA-------A-- --T-A----CA-- -------- 45195
GT#2   MG770309      294       ----- - -- 301
GT#3   HM595984      413       ----- - ----T-- -- -------- -- ----------- -- ------------ -- -C---G-- 479
GT#7   AJ488254      75        ----- - ------ -- -------- -- -------T --- -----C---- G- -------- 18
GT#8   GU246286      388       ----T -- ------ -- G------- -- ---T- A- -A -C---- 443
GT#9   GU246288      389       ----- -- ------ -- G------- -- ---C-G-A-- -A -C----- 452
GT#10  GU246287      385       -----CC-A----- G- G-G-C-- -- C--C--C-C-- --T--GC----TT- AA -C-G---- 417
GT#11  JQ695935      382       G-TTG - -CTT-G -- G-C-T- --GG-T -TCG- ----GG- - - -TC-GA-- T- -ATTAGTG 445
GT#12  GU246296      361       ----- -A----G- G- G------- ---- G----- --- --- --- --AAC-GA--- 420

GT#1   AB067721      498       CT CGCACCGG GA GCGCGGA GGCGG TCACGCC GTG AA ACCACC ACACCC TCCA 549
JAAVMX010000019 19843         -- -------- -- ------- ----- ------ - --G -- ------ ------ ---- 19894
JAAVMX010000019 32487         -- -------- -- ------- ----- ------ - --G -- ------ ------ ---- 32537
JAAVMX010000002 18702533      -- -------- -- ------- ----- ------ ---TG -- ------ ------ ---- 18702586
JAAVMX010000018 1135          -- -------- -- ------- ----- ------ - --G -- ------ ------ ---- 1186
JAAVMX010000019 6679          -- --C----- -- ------- ----- ------ - --G -- ---ACA----- ---- 6733
JAAVMX010000019 45196         -- -------- -- -------CCCAT-----C------ --GGATG--C-----ACC----- ---- 45251
GT#2   MG770309      301
GT#3   HM595984      480       -C -------- A- ----G --T- ------- -C- -- ------ --- --- 519
GT#7   AJ488254      17        --T----- 9
GT#8   GU246286      444       T- ------A A- A---- -T----- --- -- -A-T-- --G--T ---- 494
GT#9   GU246288      453       -- ------- --G------ ----- ------- -G- -- --- --G--- - -- 496
GT#10  GU246287      418       -C ---C----AA-- -G-G-- --G-G--GG- -- 478
GT#11  JQ695935      446       -- AA-GT T- C-- -CC --TTCGC-GT-AT AAT T-TCTG -- TTG- -G-G 485
GT#12  GU246296      421       -- -------- --G-G----- ---G-------G--- A-- ------ --G--T---- 480
```

Figure 3. Alignment of repetitive ITS copies within the genome JAAVMX000000000 of the **H. sinensis** *Strain IOZ07 and ITS sequences of GC-biased genotypes of* **O. sinensis.**
The genome assemblies JAAVMX010000002, JAAVMX010000018, and JAAVMX010000019 were obtained from the H. sinensis *Strain IOZ07 (Shu* et al. *2020). One copy each within JAAVMX010000002 and JAAVMX010000018, indicated in* **green**, *shares 97.4% or 97.0% similarity with AB067721. JAAVMX010000019 contains 4 repetitive ITS copies, including 2* **black** *sequences (19404→19894 and 32048→32537), which are 100% identical to AB067721, and 2 other* **blue** *sequences (6233→6733 and 44729→45251), with 94.5% and 90.8% similarity to AB067721. "GT" denotes the* O. sinensis *genotype. AB067721, MG770309, HM595984, AJ488254, GU246286, GU246288, GU246287, JQ695935, and GU246296 are the* **pink** *ITS sequences of GC-biased Genotypes #1□3 and #7□12 of* O. sinensis, *respectively. The underlined sequence in black represents the 5.8S gene of GC-biased Genotype #1* H. sinensis. *Hyphens indicate identical bases, and spaces denote unmatched sequence gaps.*

Genotype #7 was unique among the GC-biased genotypes and was placed within the Genotype #1 clade with great phylogenetic distance (horizontal line) in the Bayesian tree in Figure 2. Genotype #7 was 94.7□95.3% similar to Genotype #1, which had repetitive ITS sequences within the genome assemblies JAAVMX000000000 and NGJJ00000000, which had numerous mutant alleles (*cf*. Table 2, Figure 3).

The Genotype #7 sequence (AJ488254) was originally identified in the stroma of a *C. sinensis* specimen (H1023) collected from Qinghai Province and uploaded to GenBank by Chen *et al*. (2011). The authors identified the GC-biased Genotype #1 ITS sequence (AJ488255) from the tail of the caterpillar body of the same specimen, leading to the question of whether the sequences of AJ488254 and AJ488255 coexisted in a single genome of GC-biased Genotype #1 *H. sinensis* as 2 repetitive ITS copies or belonged to the genomes of independent *O. sinensis* fungi cooccurring in different compartments of the same *C. sinensis* specimen. The study conducted by Chen *et al*. (2011) did not involve the purification of the *O. sinensis* fungus or fungi within the same natural *C. sinensis* specimen. Thus, there is no evidence to date for determining whether the sequence (AJ488254) of Genotype #7 indeed resides in the genome of Genotype #1 *H. sinensis* as one of the repetitive ITS copies, although Genotype #7 was placed within the Genotype #1 phylogenetic clade shown in Figure 2 and was phylogenetically distant from GC-biased Genotypes #2–3 and #8–14, which were placed outside the Genotype #1 clade in the Bayesian tree shown in Figure 2.

Notably, GC-biased Genotypes #1 and #2 were simultaneously detected in the stromata of the immature, maturing and mature *C. sinensis* collected from Kangding County of Sichuan Province. These 2 genotypes showed completely different developmental patterns in an asynchronous, disproportional manner in *C. sinensis* stromata during maturation (Zhu *et al*. 2012; Li *et al*. 2023c),

indicating the genomic independence of the 2 GC-biased genotypes. The Bayesian tree shown in Figure 2 confirms that the sequences of Genotypes #1 and #2 were placed in separate GC-biased phylogenetic clades and unlikely represented as repetitive ITS copies in a single *O. sinensis* genome.

III-5. Multilocus analysis of repetitive copies of authentic genes in the *H. sinensis* genome and transcriptome

To explore the hypothesis of repetitive AT-biased pseudogenes in a single genome of GC-biased *H. sinensis* and the postulated cause of "RIP" mutagenesis, we extended the analysis of the single ITS locus to multiple loci of the *H. sinensis* genome and transcriptome. Unfortunately, GenBank does not provide annotations for authentic genes in the genome assemblies (ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000) of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively. Cross-referencing based on gene annotations for one (JAACLJ010000002) of the 13 genome contigs of *O. camponoti‐floridani* Strain EC05 (Will *et al.* 2020), which contains 5126525 bp for 1271 authentic genes, the *H. sinensis* genes and their repetitive genomic copies were then positioned and annotated in the genomes of 5 *H. sinensis* strains, and their transcription was analyzed in the mRNA transcriptome of *H. sinensis* Strain L0106. Table 4 outlines results of the multilocus analysis of 1271 *H. sinensis* genes, 1167 of which had no genomic repetitive copies, 37 of which had repetitive copies in the genome of only one *H. sinensis* strain, and 67 of which had multiple repetitive copies in the genomes of 5 *H. sinensis* strains.

Table 4. *Summary of 1271 authentic genes and their repetitive copies in the genomes of* H. sinensis *Strains 1229, CC1406-203, Co18, IOZ07, and ZJB12195.*

	# of genes		Change in AT content	# of genes having repetitive copies
No genomic repetitive copies	1167			
Single repetitive copy in only one of 5 *H. sinensis* genomes	37	Homologous (≥97% similarity)		36
		96.5% similarity	↓ 0.2%	1
Multiple repetitive copies in genomes of 5 *H. sinensis* strains	67	Essentially no change	↑ or ↓ within ±1%	6
		Slight decreases	↓ ≤5%	11
		Large decreases	↓ >5%	7
		Slight increases	↑ ≤5%	17
		Large increases	↑ >5%	11
		Bidirectional changes		15

Total:	1271

Note: "↑" and "↓" indicate an increase or decrease in AT content, respectively.

The 37 authentic genes that had repetitive copies in only one *H. sinensis* genome were normally transcribed, and the transcripts were found in the mRNA transcriptome of the *H. sinensis* Strain L0106. Only one gene (32562→34036 within LWBQ01000158) for the PNGase family protein (GenBank accession #EQL03505 containing 462 amino acids) had a single repetitive copy (414324←415851 within LWBQ01000021) in the genome of *H. sinensis* Strain ZJB12195, which showed 96.5% similarity to the authentic gene with a 53-nt insertion but no transition or transversion point mutations. The AT content did not change substantially from 35.1% in the authentic gene to 34.9% in the repetitive copy. The authentic gene with an 87-nt intron (33112→33198 within LWBQ01000158) was normally transcribed (the transcript 610→1996 within GCQL01000547 in the transcriptome of the *H. sinensis* Strain L0106). The repetitive copy was also transcribed, and the transcript 610←1996 within GCQL01000547 had an extra deletion (53 nt) in addition to the 87-nt intron (Figure 4).

The remaining 67 authentic genes in the genomes of *H. sinensis* Strains 1229, CC1406-203, Co18, IOZ07, and ZJB12195 had multiple repetitive copies in the *H. sinensis* genome assemblies with various changes (decreases, increases, or bidirectional changes) in AT content.

III-5.1 Decreases in AT content of repetitive genomic copies

Two of the 18 authentic genes (the query sequences) (*cf*. Table 4) had multiple repetitive genomic copies (the subject sequences) with slight decreases (↓ ≤5) or large decreases (↓ >5) in AT content in comparison with the sequences of authentic genes for the triose-phosphate transporter (Table 5) and heavy metal tolerance protein precursor (Table 6), respectively.

Table 5 shows that repetitive copies (124108←124983 within LKHE01003657; 751126→752000 within NGJJ01000759; 27035→27910 within ANOV01000747; 9078524←9079399 within JAAVMX010000005; and 268003→268878 within LWBQ01000030) of the authentic gene for the triose-phosphate transporter (71694←72581 within LKHE01003487; 202613→203498 within NGJJ01001434; 5923→6447 and 5261→5919 within ANOV01001461; 1710624→1711511 within JAAVMX010000003; and 1755599→1756257 and 1756277→1756590 within LWBQ01000001), respectively, had slightly reduced AT contents (↓ 2.1%), which was attributed to more T-to-C and A-to-G transitions (70) and T-to-G and A-to-C transversions (58) than did the postulated RIP mutagenesis-related transitions C-to-T and G-to-A (62) and transversions of

501   G-to-T and C-to-A (21). Overall, the number (128) of A or T to G or C was much greater than the

502   number (83) of G or C to A or T in the repetitive copies in the 5 *H. sinensis* genomes, resulting in a

503   68.1% sequence similarity and a reduction in AT content from 44.2% in the authentic genes to 42.1%

504   in the repetitive copies.

```
LWBQ01000021  414324   GGGTCGCGTGGTGGTTGAAACTCCCGATTCCGAGCTTGCTGCTGTGCCAGCATCCACTCC   414383
GCQL01000547  1996     ------------------------------------------------------------   1937

LWBQ01000021  414384   TCGGACCCTGTTTGCCGCCCGGGTCGTTCGGCGGGCAGCTTCACGTCGTGACCAGGGGGC   414443
GCQL01000547  1936     ------------------------------------------------------------   1877

LWBQ01000021  414444   GCCGGCGCGCGGCGCGTAGGAGCCGAGCCCGAGGCCACAGGGACGAGGTTCGTGACGGCC   414503
GCQL01000547  1876     ------------------------------------------------------------   1817

LWBQ01000021  414504   TGCGCGATGGATGCCACCACGTATCCCCGCAGCTCCTGGTCCTCCCGCTGGTCCTCCTTC   414563
GCQL01000547  1816     ------------------------------------------------------------   1757

LWBQ01000021  414564   TCGAGGGCGAACCGCTGGTCCTTGTCCATGTTGGCGCGGCGGAGGCCCTTGATTTCCTGC   414623
GCQL01000547  1756     ------------------------------------------------------------   1697

LWBQ01000021  414624   GTAATGTGGAGCAGCACCTCCTCGGGACATCGGTTGCGGTCGTTGGCATGCTCGGTCTTG   414683
GCQL01000547  1696     ------------------------------------------------------------   1637

LWBQ01000021  414684   CGCACATATCGCCGGGTGACGTCCGTGGCCCCGTCAATGGAGAATGCGATGCAATACGAC   414743
GCQL01000547  1636     ------------------------------------------------------------   1577

LWBQ01000021  414744   ATCTTCTTGCCCCAACCTTCCGTGTACAGCCGCGGGTTGTCCCAGGCCTCTTCGCACGCG   414803
GCQL01000547  1576     ------------------------------------------------------------   1517

LWBQ01000021  414804   TCGACGTGGATCCATCGTCGGCGGTGCTCCGAGTACACCTCGGTCCACACGTGGTCCTCG   414863
GCQL01000547  1516     ------------------------------------------------------------   1457

LWBQ01000021  414864   GCGTTCCACACCCACCGCACCCTTCCGCCGACGGCGCGGCACAGCATGCTGAAGCAGTTG   414923
GCQL01000547  1456     ------------------------------------------------------------   1397

LWBQ01000021  414924   GCCCACTCGCCGACTCGGCCACGGCGCGTCTGGAGCAGCCACCACACGTCGCCGTAGCGC   414983
GCQL01000547  1396     ------------------------------------------------------------   1337

LWBQ01000021  414984   GGAAACCTCTCGTAGACACCGCAGTGTTGGGCAGAGCAGCGGTAGAGCTCGACGCGAAGC   415043
GCQL01000547  1336     ------------------------------------------------------------   1277

LWBQ01000021  415044   GCTCCGCGCGCGCTCTCTTCCGGCGTCGGCGCCGTCATGCCCTGGTAGATGCTTGGCGAC   415103
GCQL01000547  1276     ------------------------------------------------------------   1217

LWBQ01000021  415104   AGGCAGACGGGGCATGGAGGGTTGTTGACCCAGGTGAAGAATGATCGCTTGTACCATCTG   415163
GCQL01000547  1216     ---------------------------------------------------------    1159

LWBQ01000021  415164   GACTTTGGTTAGACTTTGGTACTTTGCGGGAATGGGTAGCCGCAGACGGGTAGCCGCAGA   415223
GCQL01000547

LWBQ01000021  415224   CGACATGGCTGGAGTAGAAACATACCGAAGCAATGCCCTGATGACACAGTCCTGGTGTCC   415283
GCQL01000547  1159                               ----------------------------------    1125

LWBQ01000021  415284   CCACTCGGGTTTGCGCCCATCGCCCATGCTCTCTGCCTCGGCCTGGAGCACCTGAACCTC   415343
GCQL01000547  1124     ------------------------------------------------------------   1065

LWBQ01000021  415344   CTCCTCGGCCTCAGCGTAGATGCGATCGAGAGGAATTTTCAAGAGGGCCTCGTCGAGGAG   415403
GCQL01000547  1064     ------------------------------------------------------------   1005

LWBQ01000021  415404   GCCGGGGTTTTCGTACTTTGTGGGCGTCAACGACAGGCTGATGAGGAGGTTGCGGAACTT   415463
GCQL01000547  1039     ------------------------------------------------------------   945

LWBQ01000021  415464   TTGGGAGTCGCGGTCGGCGGCGTCTGGCGCAGTCGGGATCATGGGAAGATGGCGCATGCA   415523
GCQL01000547  944      ------------------------------------------------------------   885

LWBQ01000021  415524   CGAGTAGGATGGAGGCGGAGGAGTGGCTTGACCGTGGGCGGTGGCAGGCCGGCATGCACG   415583
GCQL01000547  884      -----------------------------------------------------------    835

LWBQ01000021  415584   AATGGAGGCGGAGGAGTGGCTTGACCGTGGGCGGTGGCAGGCCGAGCGTCGTTGGCGGCG   415643
GCQL01000547  834      --------------------------------------------    -----------   818

LWBQ01000021  415644   GGACGGGCGCGGAGGTTTGCGCTCGATGACCGCTCCGTGACGCCGGGAGAAGCATGGCTC   415703
GCQL01000547  817      ------------------------------------------------------------   758

LWBQ01000021  415704   GAGGTGGTGCGCAAGTCGGTCATGCGCTTGTCGCGCAGCAGGCCCTCGAACTTGATGCGC   415763
GCQL01000547  757      ------------------------------------------------------------   698

LWBQ01000021  415764   AGGTCCTGGGCCCATTCCTCGCCGTACTGCGAATCAGACGGGTGTCCGGGTCCCGAGGGG   415823
GCQL01000547  697      ------------------------------------------------------------   638

LWBQ01000021  415824   GTGCTTCGCCCCGCCATGATGGGGGAGG                                  415851
GCQL01000547  637      ---------------------------                                  610
```

505
506   *Figure 4. Alignments of the repetitive copy of the PNGase family protein (414324←415851 within*

LWBQ01000021) and the transcript (610←1996 within GCQL01000547) of **H.** **sinensis** *Strains ZJB12195* *and L0106 (*Liu **et al.** 2015, 2020*) were performed. Hyphens indicate identical bases, and spaces denote unmatched sequence gaps.*

The authentic genes for the triose-phosphate transporter (the genome segments within ANOV01001461, JAAVMX010000003, LKHE01003487, LWBQ01000001, and NGJJ01001434) shown in Table 5, were 98.4–100% homologous and were normally transcribed. The transcript (451→1271 within GCQL01008460 of the *H. sinensis* Strain L0106) in the transcriptome of the *H. sinensis* Strain L0106 was 100% homologous to the gene (GenBank accession #EQL01658 for the protein containing 440 aa residues), with query coverages of 79–99%. EQL01658 was defined as the triose-phosphate transporter according to the GenBank protein annotation for the genome of the *H. sinensis* Strain Co18 and showed 61.6–80.8% similarity to the triose-phosphate transporter of 30 other fungi.

The repetitive copies (the genome segments within ANOV01000747, LKHE01003657, JAAVMX010000005, LWBQ01000030, and NGJJ01000759; *cf.* Table 5) were 68.1% similar to the authentic gene and were also normally transcribed. The transcript (1←821 within GCQL01008629 of the *H. sinensis* Strain L0106) for the triose-phosphate transporter (accession #EQL02567 containing 330 aa residues) is 94.3–94.5% similar to the repetitive copy sequences with a 48-nt deletion with a query coverage of 99%. Both EQL01658 and EQL02567, which were encoded by the authentic genes and their repetitive copies, respectively, were annotated as the triose-phosphate transporter; however, these 2 protein sequences were only 65.4% similar to each other with a query coverage of 89%. EQL02567 encoded by the repetitive copies showed 59.7–68.4% similarity to 26 other protein sequences of the triose-phosphate transporter of many fungi (*Amylocarpus encephaloides*, *Colletotrichum* sp., *Coniochaeta ligniaria*, *Diplogelasinospora grovesii*, *Phialemonium atrogriseum*, *Rhexocercosporidium* sp., *Thozetella* sp., *Xylariaceae* sp., *Zalerion maritima*) and 61.3–67.4% similarity to 20 protein sequences of the solute carrier family 35 member E3 of many fungi (*Colletotrichum* sp., *Coniochaeta* sp., *Diaporthaceae* sp., *Hymenoscyphus varicosporioides*, *Lasiosphaeria hispida*, *Pyricularia oryzae*, *Tolypocladium ophioglossoides*, *Tolypocladium paradoxum*).

Table 5: The authentic H. sinensis *genes for triose-phosphate transporter (the query sequence) and the repetitive genomic copies with slight decrease (≤ 5%) in AT content.*

vs. The query sequence (202613→203498 of NG.110001434 of the authentic gene for the triose-phosphate transporter)		Mutation in the subject sequence comparing with the query sequence						Transcriptomic fragment
		Transitions		Transversions		Total point mutations		
Similarity	Change in AT content	C-to-T & G-to-A	T-to-C & A-to-G	C-to-A & G-to-T	A-to-C & T-to-G	G or C to A or T	A or C to G or C	
99.9%	Nearly no change (↓ 44.2% to 44.1%)							GCQI01008460 (451—
68.1%	44.2% to 42.1%	62	70	21	58	83	128	GCQL01008629 (1—8: with a 48-nt deletion
100%	No change (44.2%)							GCQI01008460 (451—
68.1%	44.2% to 42.1%	62	70	21	58	83	128	GCQL01008629 (1—8: with a 48-nt deletion
100%	No change (43.5%)							GCQI01008460 (810—
99.8%	Nearly no change (↓ 43.9% to 43.7%)							GCQI01008460 (451—
68.1%	44.2% to 42.1%	62	70	21	58	83	128	GCQL01008629 (1—8: with a 48-nt deletion
99.9%	Nearly no change (↓ 44.2% to 44.1%)							GCQI01008460 (451—
68.1%	44.2% to 42.1%	62	70	21	58	83	128	GCQL01008629 (1—8: with a 48-nt deletion
99.8%	Nearly no change (43.9%)	0	1	0	0	0	1	GCQI01008460 (451—
98.4%	Nearly no change (↓ 43.5% to 43.3%)	2	1	1	0	3	1	GCQI01008460 (1022—
68.1%	44.2% to 42.1%	62	70	21	58	83	128	GCQL01008629 (1—8: with a 48-nt deletion

Note: "→" and "←" indicate the sequence directions of the sense and antisense chains, respectively. "↓" indicates a decrease in AT content.

Among the 11 authentic genes outlined in Table 4 that had multiple repetitive copies with slightly reduced AT content (↓ ≤5), the repetitive copies of 4 authentic genes had mutagenetic patterns similar to that for triose-phosphate transporter with a combination of more RIP-related C-to-T than T-to-C transitions but more A-to-G than RIP-related G-to-A transitions, or *vice versa*. The repetitive copies of 6 other authentic genes had consistently more T-to-C and A-to-G transitions than did the postulated RIP-related C-to-T and G-to-A transitions, in addition to numerous transversion, insertion, and deletion point mutations.

Table 6 shows the authentic genes for heavy metal tolerance protein precursor (2772←5944 within LKHE01001285; 195005→198178 within NGJJ01001482; 2485←5657 within ANOV01008830; 1309808←1312980 within JAAVMX010000003; and 1338987← 1342159 within LWBQ01000001), which had multiple repetitive copies with large decreases (↓ >5) in AT

content. The repetitive copies of the authentic gene could be divided into 2 groups.

(1) The first group of repetitive copies (3819←4233 within LKHE01002445; 150283→150697

within NGJJ01001243; 3175→3589 within ANOV01010958; 833901←834315 within

JAAVMX010000002; and 468456←468870 within LWBQ01000010) in the genomes of the *H.*

sinensis Strains 1229, CC1406-203, Co18, IOZ07, and ZJB12195, respectively, were 65.4% similar

to the authentic gene. These repetitive copies had many more T-to-C and A-to-G transitions (26) and

A-to-C and T-to-G transversions (27) than postulated RIP-related C-to-T and G-to-A transitions (14)

and C-to-A and G-to-T transversions (14). Overall, the total number (28) of G or C to A or T point

mutations was much less than the number (53) of A or T to G or C mutations in the repetitive copies,

resulting in largely reduced AT contents (↓ 6.2%) from 39.5% in the authentic gene to 33.3% in the

repetitive copies.

Table 6: The authentic H. sinensis *genes for the heavy metal tolerance protein precursor (the*
query sequence) and the repetitive genomic copies with major decrease (> 5%) in AT content.

vs. The query sequence (19500S→198178 of NGJJ01001482 of the authentic gene for the heavy metal tolerance protein precursor)		Mutation in the subject sequence comparing with the query sequence						Transcript in mRNAtr	
		Transitions		Transversions		Total point mutations			
Similarity	Change in AT content	C-to-T & G-to-A	T-to-C & A-to-G	C-to-A & G-to-T	A-to-C & T-to-G	G or C to A or T	A or T to G or C	Transcriptomic fragment	Simi
99.97%	Nearly no Δ (41.2%)							GCQL01012210 (442←1471); GCQL01006458 (2←683); GCQL01000615 (1←403)	100%; 99.9%; 100%
65.4%	↓39.5% to 33.3%	14	26	14	27	28	53	GCQL01017466 (726←1140)	100%
67.4%	↓38.3% to 34.6%	24	34	21	34	45	68	GCQL01013313 (1←426); GCQL01016959 (1←271)	99.8%; 100%
100%	No Δ (40.2%)							GCQL01012210 (442←1471); GCQL01006458 (2←683); GCQL01000615 (1←403)	100%; 99.9%; 100%
65.4%	↓39.5% to 33.3%	14	26	14	27	28	53	GCQL01017466 (726←1140)	100%
67.4%	↓38.3% to 34.6%	24	34	21	34	45	68	GCQL01013313 (1←426); GCQL01016959 (1←271)	99.8%; 100%
99.97%	Nearly no Δ (40.2%)							GCQL01012210 (442←1471); GCQL01006458 (2←683); GCQL01000615 (1←403)	100%; 99.9%; 100%
65.4%	↓39.5% to 33.3%	14	26	14	27	28	53	GCQL01017466 (726←1140)	100%
67.4%	↓38.3% to 34.6%	24	34	21	34	45	68	GCQL01013313 (1←426); GCQL01016959 (1←271)	99.8%; 100%
99.97%	Nearly no Δ (40.2%)							GCQL01012210 (442←1471); GCQL01006458 (2←683); GCQL01000615 (1←403)	100%; 99.9%; 100%
65.4%	↓39.5% to 33.3%	14	26	14	27	28	53	GCQL01017466 (726←1140)	100%
67.4%	↓38.3% to 34.6%	24	34	21	34	45	68	GCQL01013313 (1←426); GCQL01016959 (1←271)	99.8%; 100%
99.97%	Nearly no Δ (41.2%)							GCQL01012210 (442←1471); GCQL01006458 (2←683); GCQL01000615 (1←403)	100%; 99.9%; 100%

Note: "→" and "←" indicate the sequence directions of the sense and antisense chains, respectively. "↓" indicates a decrease in AT content.

(2) The second group of repetitive copies (128317←128839 and 128885←129014 within LKHE01001747; 461774←461903 and 461949←462471 within NGJJ01001310; 4224←4746 and 4792←4921 within ANOV01005573; 85112←85634 and 85680←85809 within JAAVMX010000012; and 141513→142035 and 142081→142210 within LWBQ01000044) of the authentic gene in the 5 *H. sinensis* genome assemblies showed 67.4% sequence similarity to the authentic genes. There were more T-to-C and A-to-G transitions (34) and A-to-C and T-to-G transversions (34) than postulated RIP-related C-to-T and G-to-A transitions (24) and C-to-A and G-to-T transversions (21). Overall, this group of repetitive copies had a total of 45 G or C to A or T

point mutations, which was much less than the number (68) of A or T to G or C point mutations in the repetitive copies, resulting in slightly reduced AT contents (↓ 3.7%) from 38.3% in the authentic gene to 34.6% in the repetitive copies.

Also shown in Fig. 6, the authentic genes (the genome segments within ANOV01008830, JAAVMX010000003, LKHE01001285, LWBQ01000001, and NGJJ01001482) were normally transcribed. The fragmented transcripts (GCQL01012210, GCQL01006458, and GCQL01000615 of the *H. sinensis* Strain L0106) were 99.9–100% identical to the authentic gene for the heavy metal tolerance protein precursor (accession #EQK98634 containing 1029 amino acids), with query coverages of 12–32%.

The first group of repetitive copies (the genome segments within ANOV01010958, JAAVMX010000002, LKHE01002445, LWBQ01000010, and NGJJ01001243) were 65.4% similar to the authentic gene and were normally transcribed (*cf*. Table 6). The transcript GCQL01017466 was 100% homologous to the repetitive copies with a query coverage of 100%; however, it presumptively encoded a different protein, the transporter ATM1 (accession #EQK98286, which contains 598 amino acids), according to the GenBank protein annotation for the genome of the *H. sinensis* Strain Co18. EQK98286 was 84.8–89.4%% similar to 57 protein sequences of the transporter ATM1 or mitochondrial iron-sulfur clusters transporter ATM1 of different fungi.

The second group of repetitive copies (the genome segments within ANOV01005573, JAAVMX010000012, LKHE01001747, LWBQ01000044, and NGJJ01001310) were 67.4% similar to the authentic genes and were also normally transcribed (*cf*. Table 6). The fragmented transcripts (GCQL01013313 and GCQL01016959) were 99.8–100% homologous to the repetitive copies with query coverages of 38–61%; however, they presumably encoded a different protein, the ABC transporter (accession #EQK99315 containing 844 amino acids), according to the GenBank protein annotation for the genome of the *H. sinensis* Strain Co18. EQK99315 was 50.2–89.0% similar to 22 other protein sequences of the ABC transporter of many different fungi.

Similar to the mutagenetic pattern of the authentic gene encoding the heavy metal tolerance protein precursor, the repetitive copies of 6 other genes whose AT content decreased substantially (↓ >5%) (*cf*. Table 4) were associated with consistently more T-to-C and A-to-G transitions than were the postulated RIP-related C-to-T and G-to-A transitions, in combination with numerous transversion, insertion and deletion point mutations.

III-5.2 Increases in AT content of repetitive genomic copies

The authentic genes for the maltose permease and lipase/serine esterase are shown in Tables 7–8 as representatives of the 28 authentic genes that had multiple repetitive copies with slight increases (↑ ≤5%) and large increases (↑ >5%) in AT content, respectively.

Table 7 shows the authentic genes for the maltose permease in 5 *H. sinensis* genome assemblies (47693→49623 within LKHE01002657; 77988→79917 within NGJJ01000203; 1725←3655 within ANOV01009266; 4614206→4616136 within JAAVMX010000006; and 66747→68677 within LWBQ01000157).

The repetitive sequences (6824→7385 within LKHE01000336; 782871←782913 and 1076399←1076960 within NGJJ01001217; 4433←4994 and 5254←5290 within ANOV01000509; 12448464→12449025 and 12779815→12779864 within JAAVMX010000005; and 292175→292217 and 570642→571203 within LWBQ01000003) of the authentic gene showed 69.6–70.4% sequence similarities to the gene. These repetitive sequences had more T-to-C and A-to-G transitions (35–37) than did the postulated RIP-related C-to-T and G-to-A transitions (30–31), and more C-to-A and G-to-T transversions (28) than did the A-to-C and T-to-G transversions (10–11). Overall, there was a greater number of G or C to A or T point mutations (58–59) than of A or T to G or C point mutations (46–47) in the repetitive copies, resulting in slight increases (↑ ≤5%) in AT content from 36.8–37.9% in the authentic genes to 38.5–39.9% in the repetitive copies.

Table 7: The authentic H. sinensis *genes for the maltose permease (query sequence) and the repetitive genomic copies with minor increase (≤ 5%) in AT content.*

Table (rotated): *...permease) and the repetitive genomic copies with minor increase (≤5%) in the AT content.*

vs. The query sequence (47693→49623 of LKHE01002657 of the authentic gene for the maltose permease)		Mutation in the subject sequence comparing with the query sequence							Transcriptome
		Transitions		Transversions		Total point mutations			
Similarity	Change in AT content	C-to-T & G-to-A	T-to-C & A-to-G	C-to-A & G-to-T	A-to-C & T-to-G	G or C to A or T	A or T to G or C		
100%	No ↓ (+0.1%)								GCQL01010488
69.6%	↑ 37.9% to 39.3%	30	35	28	11	58	46		GCQL01015773; GCQL01019033 with a 25-nt del
99.8%	Nearly no ∆ (-0.1% to +0.2%)	1	0	0	0	1	0		GCQL01010488
70.4%	↑ 36.8% to 38.5%	31	37	28	10	59	47		GCQL01015773; GCQL01019033 with a 25-nt del
99.8%	Nearly no ∆ (-0.1% to +0.0%)	1	2	0	0	1	2		GCQL01010488
70.1%	↑ 37.9% to 39.9%	31	37	28	10	59	47		GCQL01015773; GCQL01019033 with a 25-nt del
99.9%	Nearly no ∆ (+0.1%)	1	0	0	0	1	0		GCQL01010488
70.4%	↑ 37.6% to 39.2%	31	37	28	10	59	47		GCQL01015773; GCQL01019033 with a 25-nt del
99.9%	Nearly no ∆ (+0.1%)	1	0	0	0	1	0		GCQL01010488
70.4%	↑ 36.8% to 38.5%	30	35	28	11	58	46		GCQL01015773; GCQL01019033 with a 25-nt del

...sense chains, respectively. "↑" and "↓" indicate increases and decreases in AT content.

Note: "→" and "←" indicate the sequence directions of the sense and antisense chains, respectively. "↑" and "↓" indicate increases and decreases in AT content, respectively.

Table 7 also shows that the authentic genes (the genome segments within ANOV01009266, JAAVMX010000006, LKHE01002657, LWBQ01000157, and NGJJ01000203) were normally transcribed. The transcript GCQL01010488 was 99.8–100% homologous to the authentic genes for the maltose permease (accession #EQK98556 according to the GenBank protein annotation for the genome of the *H. sinensis* Strain Co18), with a query coverage of 80%.

The repetitive copies (the genome segments within ANOV01000509, JAAVMX010000005, LKHE01000336, LWBQ01000003, and NGJJ01001217) were transcribed, and 2 transcripts (GCQL01015773 and GCQL01019033) were 98.7% identical with a query coverage of 100%. GCQL01015773 was 100% homologous to the repetitive copies, and GCQL01019033 was 95.6% similar to the repetitive copies with a 25-nt deletion. These 2 transcripts encoded a different protein (MRT, a raffinose family of oligosaccharide transporters; 152→338 of EQL02995 containing 550 amino acids according to the GenBank protein annotation for the genome of the *H. sinensis* Strain Co18); however, the protein encoded by GCQL01019033 had a deletion of 8 amino acids.

In addition to the authentic *H. sinensis* genes for the maltose permease shown in Table 7, there were 16 other authentic genes outlined in Table 4 with slight increases in AT content. Seven of them exhibited a mutagenetic pattern with contradictory transitions (T-to-C and A-to-G *vs*. RIP-related C-to-T and G-to-A) and transversions (T-to-G and A-to-C *vs*. C-to-A and G-to-T) in combination with various numbers of insertion and deletion point mutations, similar to the mutagenetic pattern for the maltose permease. Nine other genes had consistently more RIP-related C-to-T and G-to-A transitions than T-to-C and A-to-G transitions, and more C-to-A and G-to-T transversions than T-to-G and A-to-C transversions; and many insertion and deletion point mutations.

Table 8 shows the sequences of the authentic genes for the lipase/serine esterase (22031→22947 within LKHE01002746; 235828→236719 within NGJJ01000647; 9368→10284 within ANOV01000049; 10692268←10693159 within JAAVMX010000005; and 203574←204490 within LWBQ01000026) in the 5 *H. sinensis* genomes. The repetitive copies (39965→40763 and 40848→40982 within LKHE01001863; 1146814→1147612 and 1147697→1147831 within NGJJ01001576; 6319←6453 and 6538←7335 within ANOV01000484; 22221540←22222338 within JAAVMX010000001; and 513365→514163 and 514248→514382 within LWBQ01000013) were 67.0–68.5% similar to the authentic genes in the 5 *H. sinensis* genomes. There were more postulated RIP-related C-to-T and G-to-A transitions (63) than T-to-C and A-to-G transitions (30) and more C-to-A and G-to-T transversions (41) than A-to-C and T-to-G transversions (20) in the repetitive copies. Overall, there were more point mutations of G or C to A or T (104) in the repetitive copies than the point mutations of A or T to G or C (50), resulting in large increases (↑ >5%) in AT content from 34.9–35.0% in the authentic genes to 41.5–41.6% in the repetitive copies.

Table 8 also shows that the highly homologous (99.8–100%) authentic genes (the segments within ANOV01000049, JAAVMX010000005, LKHE01002746, LWBQ01000026, and NGJJ01000647,) were transcribed. Two transcripts (GCQL01000519 and GCQL01015314) were 100% homologous to the authentic gene and encoded partially overlapping protein sequences of lipase/serine esterase (accession #EQL04141) with query coverages of 68–100%.

The repetitive copies (the segments within ANOV01000484, JAAVMX010000001, LKHE01001863, LWBQ01000013, and NGJJ01001576) were 67.0–68.5% similar to the authentic gene and were normally transcribed. The transcript (GCQL01000532) was 99.7–100% homologous to the repetitive copies with a query coverage of 100%; however, it presumptively encoded a

different protein, the hypothetical protein OCS_01262 (accession #EQL03018 containing 383 amino acids), according to the protein annotation for the genome of the *H. sinensis* Strain Co18. This protein sequences (EQL03018) and EQL04141 that was encoded by the authentic gene were only 46.7% similar to each other with a query coverage of 94%, indicating the 2 sequences represented different proteins. EQL03018 and another protein sequence KAF4513393 were obtained from the *H. sinensis* Strains Co18 and IOZ07 and annotated as the hypothetical proteins OCS_01262 and G6O67_000668 in GenBank, respectively. They were 61.5–82.3% similar to 15 GenBank protein sequences for the lipase/serine esterase of *Pochonia chlamydosporia* and Purpureocillium sp., the putative serine esterase of *Hirsutella rhossiliensis*, *Ilyonectria robusta*, *Metarhizium acridum*, *Thelonectria olida*, and *Trichoderma breve*; and the putative lipase-like protein of *Fusarium albosuccineum*, *Hapsidospora chrysogenum*, *Metarhizium anisopliae*, *Tolypocladium* sp., and *Trichoderma lentiforme*. Thus, EQL03018 and EQL04141% might probably represent 2 different proteins.

The repetitive copies of 10 other *H. sinensis* genes with largely increased AT content (*cf*. Table 4) had consistent point mutations with more RIP-related C-to-T and G-to-A transitions than T-to-C and A-to-G transitions, in addition to transversion, insertion, and deletion point mutations.

Table 8: The authentic H. sinensis *genes for the lipase/serine esterase (query sequence) and the repetitive genomic copies with major increase (> 5%) in AT content.*

Table caption (partial): vs. The query sequence (235828—236719 of NGJJ01000647 of the authentic gene for the lipase/serine esterase)

Similarity	Change in AT content	Transitions		Transversions		Total point mutations		Transcriptomic
		C-to-T & G-to-A	T-to-C & A-to-G	C-to-A & G-to-T	A-to-C & T-to-G	G or C to A or T	A or T to G or C	
100%	No Δ (34.5%)							GCQL01000519, GCQL01015311
68.5%	↑35.0% to 41.5%	63	30	41	20	104	50	GCQL01000532
100%								GCQL01000519, GCQL01015311
68.5%	↑35.0% to 41.6%	63	30	41	20	104	50	GCQL01000532
99.8%	No Δ (34.5%)							GCQL01000519, GCQL01015314
67.0%	↑34.9% to 41.5%	63	30	41	20	104	50	GCQL01000532
100%	No Δ (34.5%)							GCQL01000519, GCQL01015314
67.1%	↑34.9% to 41.6%	63	30	41	20	104	50	GCQL01000532
100%	No Δ (34.5%)							GCQL01000519, GCQL01015314
68.5%	↑35.0% to 41.6%	63	30	41	20	104	50	GCQL01000532

...nse and antisense chains respectively. "↑" indicates an increase in AT content

Note: "→" and "←" indicate the sequence directions of the sense and antisense chains, respectively. "↑" indicates an increase in AT content.

III-5.3 Bidirectional changes in AT content of repetitive genomic copies

Table 9 shows the authentic gene for β-lactamase/transpeptidase-like protein (19330→20932 within LKHE01000540; 70191→71142 within NGJJ01001580; 2857←4459 within ANOV01000528; 238583←240185 within JAAVMX010000002; and 168259←169259 within LWBQ01000052) of the 15 authentic genes that had multiple repetitive copies with bidirectional changes in AT content in 5 *H. sinensis* genome assemblies. The multiple repetitive copies could be divided into 3 groups, 2 of which had reduced AT contents but slightly different patterns of point mutations, and the other group had increased AT contents.

The first group of repetitive copies with slight decreases (↓ ≤5%) in AT content (54230←54311 and 54418←55104 within LKHE01001740 and 883241→883927 and 884034→884115 within JAAVMX010000009) had an equal number (68) of RIP-related transitions (C-to-T and G-to-A) and opposite transitions (T-to-C and A-to-G) but more A-to-C and T-to-G transversions (40) than G-to-T and C-to-A transversions (26). Overall, the repetitive copies contained more A or T to G or C point

mutations (108) than G or C to A or T point mutations (94), resulting in a sequence similarity of 67.9% to the authentic gene (β-lactamase/transpeptidase-like protein) and a 5% decrease in AT content from 38.0% in the authentic genes to 33.0% in the repetitive copies.

The second group of repetitive copies with slightly reduced AT content (697372→698058 and 698165→698246 within NGJJ01000083; 136→549 and 656→737 within ANOV01000033; and 1407095←1407176 and 1407283←1407696 within LWBQ01000002) were 67.9–70.4% similar to the authentic gene and had more T-to-C and A-to-G transitions (110 and 27) than the postulated RIP-related C-to-T and G-to-A transitions (85 and 16) and more A-to-C and T-to-G transversions (71 and 15) than C-to-A and G-to-T transversions (40 and 14). Overall, there were fewer total point mutations of G or C to A or T (125 and 30) in the repetitive copies than the point mutations of A or T to G or C (181 and 44), which was attributed to slight decreases in AT content from 35.4–38.0% in the authentic gene sequences to 32.7–33.0% in the repetitive copies.

Table 9: The authentic H. sinensis *genes for the β-lactamase/transpeptidase-like protein (the query sequence) and the repetitive genomic copies with bidirectional changes in AT content.*

Table (rotated): es for the β-lactamase/transpeptidase-like protein (the query sequence) and the repetitive genomic copies with bidirectional changes in AT content.

vs. The query sequence (70191→71142 of NGJJ01001580 of the authentic gene for the β-lactamase/transpeptidase-like protein)			Mutation in the subject sequence comparing with the query sequence						Transcript in mRNA tran	
			Transitions		Transversions		Total point mutations			
(sequence range)	Similarity	Change in AT content	C-to-T & G-to-A	T-to-C & A-to-G	C-to-A & G-to-T	A-to-C & T-to-G	G or C to A or T	A or T to G or C	Transcriptomic fragment	Simila
0932)	99.9%	No Δ (38.7%)							GCQL01006585 (201→504); GCQL01001125 (1→411); GCQL01011658 (1152→1882) with a 61-nt deletion	100%; 99.8%; 92.1%
4311; 54418→55104)	67.9%	↓38.0% to 33.0%	68	68	26	40	94	108	GCQL01006426 (997→1188); GCQL01002069 (3→392); GCQL01008668 (979→1256)	100%; 99.7%; 98.2%
1185)	64.2%	↑38.0% to 42.0%	74	46	48	43	122	89	GCQL01012138 (368→1264) with a 51-nt intron	94.6%
142)	100%	No Δ (38.7%)							GCQL01006585 (228→504); GCQL01001125 (1→411); GCQL01011658 (1718→1882)	99.6%; 99.8%; 99.4%
'00058; 698165→698246)	67.9%	↓38.0% to 33.0%	85	110	40	71	125	181	GCQL01006426 (997→1188); GCQL01002069 (3→392); GCQL01008668 (979→1256)	100%; 99.7%; 98.2%
51549)	64.2%	↑38.0% to 41.9%	74	46	48	43	122	89	GCQL01012138 (368→1264) with a 51-nt intron	94.5%
59)	99.8%	Nearly no Δ (↑38.7% to 38.9%)	0	0	1	0	1	0		99.7%; 99.8%; 92.0%
656→737)	70.4%	↓35.7% to 32.7%	16	27	14	15	30	44	GCQL01006426 (997→1188); GCQL01002069 (3→392)	100%; 99.7%
6246)	64.2%	↑38.0% to 41.9%	74	46	48	43	122	89	GCQL01012138 (368→1264) with a 51-nt intron	94.6%
83→240185)	100%	No Δ (38.7%)							GCQL01006585 (201→504); GCQL01001125 (1→411); GCQL01011658 (1152→1882) with a 61-nt deletion	99.7%; 99.8%; 92.0%
41→883597; 884034→884115)	67.9%	↓38.0% to 33.0%	68	68	26	40	94	108	GCQL01006426 (997→1188); GCQL01002069 (3→392); GCQL01008668 (979→1256)	100%; 99.7%; 98.2%
9894→16800841)	64.2%	↑38.0% to 42.0%	74	46	48	43	122	89	GCQL01012138 (368→1264) with a 51-nt intron	94.6%
-169259)	100%	No Δ (40.2%)							GCQL01001125 (1→196); GCQL01011658 (1149→1882)	100%; 92.2%

720
Note: "→" and "←" indicate the sequence directions of the sense and antisense chains,
respectively. "↑" and "↓" indicate increases and decreases in AT content, respectively.

The third group of repetitive copies with increased AT contents shown in Table 9

(20238←21185 within LKHE01002381; 250601→251549 within NGJJ01000732; 15299→16246

within ANOV01000652; 16799894←16800841 within JAAVMX010000003; and 210989←211936

within LWBQ01000004) were 64.2% similar to the authentic gene. There were more postulated

RIP-related C-to-T and G-to-A transitions (74) than T-to-C and A-to-G transitions (46) and more C-

to-A and G-to-T transversions (48) than A-to-C and T-to-G transversions (43) in the repetitive

sequences. Overall, the repetitive copies had more G or C to A or T point mutations (122) than A or

T to G or C point mutations (89), resulting in 3.9–4.0% increases in AT content from 38.0% in the
authentic genes to 41.9–42.0% in the repetitive copies.

Also shown in Table 9, the authentic genes (the genome segments within ANOV01000528,
JAAVMX010000002, LKHE01000540, LWBQ01000052, and NGJJ01001580) for the β-
lactamase/transpeptidase-like protein were transcribed. Among the 3 fragmented transcripts,
GCQL01006885 and GCQL01011125 were 99.6–100% homologous to the authentic gene with
query coverages of 17–43%; while the other transcript, GCQL01011658, was 92.1% similar (with a
58-nt deletion) to the authentic genes with query coverages of 29–49%. The fragmented transcripts
encoded the non-overlapping sequences of the β-lactamase/transpeptidase-like protein with
sequence gaps (accession #EQL02970, containing 531 amino acids).

The repetitive copies with reduced AT contents (the genome segments within ANOV01000033,
JAAVMX010000009, LKHE01001740, LWBQ01000002, and NGJJ01000083) were transcribed.
The transcripts (GCQL01006426, GCQL01008668, and GCQL01020269) were 98.2–100%
homologous to the sequences of repetitive copies, with query coverages of 21–65% and short
overlaps and gaps in protein sequences. The transcripts presumably encoded a different protein, the
hypothetical protein G6O67_008083 (accession #KAF4504658 containing 542 amino acids),
according to the protein annotation for the *H. sinensis* Strain IOZ07.

The repetitive copies with increased AT content (the genome segments within ANOV01000652,
JAAVMX010000003, LKHE01002381, LWBQ01000004, and NGJJ01000732) were transcribed.
The GCQL01012138 transcript was 94.6% similar (with a 51-nt deletion) to the repetitive
sequences and also encoded the β-lactamase/transpeptidase-like protein (accession #EQL02706
containing 558 amino acids) with 100% query coverage.

Both EQL02706 and KAF4504658 that were encoded by the repetitive sequences, and their
sequences were only 50% and 46.2% similar to EQL02970 encoded by the authentic gene for the β-
lactamase/transpeptidase-like protein of the *H. sinensis* Strain Co18 with query coverages of 96%.
The protein sequences were 47.9–80.4% similar to the β-lactamase/transpeptidase-like proteins of
different fungi: *Akanthomyces lecanii*, *Dactylonectria* sp., *Fusarium* sp., *Hirsutella rhossiliensis*,
Ilyonectria sp., *Mariannaea* sp., *Metarhizium* sp., *Microthyrium microscopicum*, *Thelonectria* sp.,
and *Trichoderma* sp.; and 49.3–63.3% similar to the penicillin-binding protein of different fungi:
Drechmeria coniospora, *Fusarium* sp., *Pochonia chlamydosporia*, *Purpureocillium* sp., and
Trichoderma arundinaceum.

III-6. Transcriptional silencing of 5.8S genes

The above genetic, genomic, and phylogenetic analyses demonstrated GC-biased characteristics of the multiple repetitive ITS copies in the *H. sinensis* genome and that all GC- and AT-biased genotypes of *O. sinensis* were genome independent. Li *et al*. (2013) identified ITS sequences for both Genotypes #1 and #5 from genomic DNA that was isolated from 8 (1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) of 15 cultures of natural *C. sinensis* multicellular heterokaryotic mono-ascospores (Bushley *et al*. 2013). However, Li *et al*. (2013) found 5.8S cDNA only for Genotype #1 and not for Genotype #5 in a cDNA library constructed from total ascosporic RNA extracted from Clone 1220, one of the 8 genetically heterogeneous clones. This paper did not report cDNA library constructions of 7 other genetically heterogeneous, ascosporic clones (1206, 1208, 1209, 1214, 1224, 1227, and 1228) or subsequent cDNA examinations for Genotypes #1 and #5 of *O. sinensis*.

Li *et al*. (2013) did not detect the ITS sequences of AT-biased Genotypes #4, #6, or #15□17 in the genomic DNA isolated from any of the 15 ascosporic clones. Thus, there is insufficient evidence to conclude that the 5.8S genes of all AT-biased genotypes of *O. sinensis* are permanently nonfunctional pseudogenic components coexisting in "a single genome" with a functional copy of GC-biased Genotype #1. Li *et al*. (2019) provided the following logical reasoning underlying the nonfunctional pseudogene hypothesis: "If AT-biased ITS sequences are not pseudogenes, 5.8S cDNA should have been detected because rRNAs are critical and essential for protein synthesis in all living organisms", which critically touches on 2 serious scientific issues. First, are the 5.8S genes of all the *O. sinensis* genotypes actively transcribed in *C. sinensis* ascospores (discussed in this section)? Second, is it correct to conclude that "5.8S cDNA should have been detected" by the techniques designed by Li *et al*. (2013) (discussed in Section III-7)?

Many research groups have reported the differential coexistence of several AT-biased genotypes in various combinations in different compartments of natural *C. sinensis*, as summarized in Table 10 (Xiao *et al*. 2009; Zhu *et al*. 2010, 2012; Gao *et al*. 2011, 2012; Li *et al*. 2023b, 2023c). Genotypes #4 and #15 of the AT-biased Cluster B of *O. sinensis* (*cf*. Figure 2) were reported to exist in the stroma and stromal fertile portion (densely covered with numerous ascocarps) of natural *C. sinensis* but not in the ascospores (Li *et al*. 2023b, 2023c), consistent with the failure to identify Genotype #4 of AT-biased Cluster B in mono-ascospores as reported by Li *et al*. (2013) after additional specific efforts. Thus, there is no evidence demonstrating that AT-biased Genotypes #4 and #15

occur in *C. sinensis* ascospores, and there is no ground for claiming that the 5.8S genes of Genotypes #4 and #15 are nonfunctional pseudogenes in the genome of GC-biased Genotype #1.

Table 10. *Differential occurrence of AT-biased genotypes of* O. sinensis *in different compartments of natural* C. sinensis.

Genotype # and GenBank accession #	Phylogenetic AT-biased Cluster [§]	Stroma		Stromal Fertile Portion (covering with numerous ascocarps)	Ascospores
		Immature	Mature		
#4 AB067744	B	√	√	√	
#5 AB067740	A	√	√	√	√
#6 EU555436	A	√	√	√	√
#15 KT232017	B	√	√	√	
#16 KT232019	A	√			√
#17 KT232010	A	√			

Note: This table was adapted and altered from Li et al. *(2023c* https://doi.org/10.1371/journal.pone.0270776*). [§], Phylogenetic AT-biased Clusters A and B are shown in Figure 2 (the Bayesian majority rule consensus phylogenetic tree) and are similarly shown in the phylogenetic trees generated by using the Bayesian, maximum likelihood, and minimum evolution algorithms by Stensrud* et al. *(2007), Zhang* et al. *(2013a) and Li* et al. *(2016b, 2022a, 2022c, 2023b, 2023c).*

Xiang *et al*. (2014) reported a metatranscriptome study of natural *C. sinensis* specimens (unknown maturation stage) collected from Kangding County of Sichuan Province, China. A GenBank BLAST search confirmed that the metatranscriptome GAGW00000000 sequences do not contain any 5.8S gene transcripts of *C. sinensis*-associated fungi (including all genotypes of *O. sinensis*), indicating transcriptional silencing of 5.8S genes of all cocolonized fungi in natural settings.

Xia *et al*. (2017) reported another metatranscriptome study of mature natural *C. sinensis* specimens collected from Deqin County of Yunnan Province, China. The metatranscriptome assembly was not uploaded to GenBank but rather to another depository server, www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas.

Similarly, this assembled metatranscriptome contains no 5.8S gene transcripts. However, the authors uploaded the unassembled shotgun reads (SRR5428527) to GenBank. A BLAST search of the SRR5428527 Shotgun Read Archive (SRA) using the MegaBLAST algorithm identified 1894 unassembled metatranscriptome shotgun reads (100 bp each) that were 100% homologous to the 5.8S gene sequence (156 bp) of GC-biased Genotype #1 *H. sinensis*.

In general, fungal genomes contain multiple repetitive copies of rDNA ITS1-5.8S-ITS2

sequences (*cf*. Tables 1 and 3; Figures 1□3). Because the assembled metatranscriptome contained

no 5.8S gene transcripts (Xia *et al*. 2017) and no segments of the rDNA 18S or 28S gene of *H.*

sinensis were homologous to the 5.8S gene, the 1894 unassembled shotgun reads (SRR5428527) of

natural *C. sinensis* may be derived from highly homologous non-rRNA genes that evolved in a

conservative manner within fungal metagenomes. Figure 5 shows the GenBank BLAST graphic

summary of the alignment of the 5.8S gene (the query sequence; segment 218□373 of Genotype #1

H. sinensis AB067721) and the first 75 of the overlapping 1894 metatranscriptome shotgun reads

(100 bp each) of mature natural *C. sinensis*.

Figure 5. Graphic alignment summary of the query sequence (in blue□green), the 5.8S gene (218□373

segment of Genotype #1* H. sinensis *AB067721) and the first 75 in the list of 1894 overlapping shotgun

reads (pink bars; GenBank SRR5428527) of mature natural* C. sinensis*.

Various degrees of sequence similarity were found when the 1894 unassembled

metatranscriptome shotgun reads were aligned with the 5.8S gene sequences of Genotypes #2–17 of

O. sinensis. Table 11 shows 6 examples of shotgun reads of SRX2718557 aligned with the 5.8S

sequences of *O. sinensis* genotypes, *P. hepiali*, *T. sinensis* (Group D), and a Group-E AB067719-

type fungus (Stensrud *et al.* 2007). The high homology between the shotgun reads and the various

5.8S gene sequences of *O. sinensis* genotypes may not conclusively define the fungal sources in

natural *C. sinensis*, which contains >90 species spanning at least 37 fungal genera (Zhang *et al.*

2010, 2018; Xiang *et al.* 2014; Xia *et al.* 2015, 2017).

Table 11. Similarity between the unassembled metatranscriptome shotgun reads (SRA:

SRR5428527) of natural C. sinensis *and the 5.8S genes of multiple genotypes of* O. sinensis *and* P.

hepiali, T. sinensis *(Group D) and a Group-E AB067719-type fungus.*

Geno-type	GenBank accession #	Percent similarity between 5.8S genes and *C. sinensis* shotgun sequences					
		SRR5428527. 17546327.1	SRR5428527.1 7554054.1	SRR5428527.1 7565575.2	SRR5428527.1 7679995.2	SRR5428527.1 7911498.2	SRR5428527.1 7929569.2
#1	AB067721	100%	100%	100%	100%	100%	100%
#2	MG770309	96%	96%	100%	96%	98%	96%
#3	HM595984	100%	99%	99%	100%	99%	100%
#7	AJ488254	98%	98%	99%	98%	98%	98%
#8	GU246286	94%	96%	95%	94%	97%	94%
#9	GU246288	99%	99%	99%	100%	99%	100%
#10	GU246287	96%	94%	93%	95%	93%	97%
#11	JQ695935	100%	100%	100%	100%	100%	100%
#12	GU246296	99%	99%	99%	99%	99%	99%
#13	KT339190	98%	93%	93%	98%	93%	98%
#14	KT339178	98%	93%	93%	98%	93%	98%
#4	AB067744	86%	87%	87%	83%	87%	83%
#5	AB067740	85%	87%	87%	84%	87%	84%
#6	KJ720572	85%	85%	87%	84%	86%	84%
#15	KT232017	85%	88%	88%	84%	88%	84%
#16	KT232019	86%	86%	87%	85%	87%	85%
#17	KT232010	87%	88%	88%	87%	88%	87%
P.h.	EF555097	99%	98%	97%	99%	98%	99%
T.s.	AB067720	98%	97%	96%	98%	97%	98%
	AB067719	98%	93%	93%	98%	93%	98%

Note: GenBank sequence ID SRA: SRR5428527 fragments (17546327.1; 17554054.1; 17565575.2;

17679995.2; 17911498.2; and 17929569.2) are the unassembled transcriptome shotgun sequences

obtained from the mature natural C. sinensis *specimens collected from Deqin County of Yunnan*

Province by Xia et al. *(2017). GC-biased genotypes are listed in* blue, *and AT-biased genotypes are*

listed in red. *P.h. refers to* Paecilomyces hepiali; *T.s. refers to Group-D* Tolypocladium sinensis;

and AB067719 refers to an AB067719-type Group-E fungus (Stensrud et al. 2007). *Highly*

homologous percent similarities (97□100%) are underlined.

As shown in Table 11, the representative unassembled metatranscriptome shotgun reads of
natural *C. sinensis* were 97□99% homologous to the 5.8S gene sequence of *P. hepiali*. Some of the
shotgun reads were 97□98% homologous to the 5.8S gene sequences of *T. sinensis* and the
AB067719-type fungus. A GenBank BLAST search revealed the alignment of the 5.8S gene
sequence (segment 218→373 within AB067721) of *H. sinensis* to 56□438 unassembled
transcriptome shotgun sequence reads, which were identified in 5 RNA-Seq experimental data files
(GenBank assessment #SRX3083435□SRX3083439) of *P. hepiali* Strain Feng (SRA Study:
SRP115168; Pang *et al*. 2018) (Table 12). Pang *et al*. (2018) uploaded 5 transcriptomic
experimental datasets to GenBank for total RNA samples extracted from cultures of *P. hepiali* Strain
Feng under different culture conditions and durations.

Table 12. *Summary of BLAST searches of the transcriptome shotgun reads of* P. hepiali *Strain Feng*
against the 5.8S gene sequence (218→373 within AB067721) of Genotype #1 H. sinensis*.*

P. hepiali sample	Culture duration (days)	Additive in culture medium	GenBank transcriptome data file	*vs*. the 5.8S gene sequence of *H. sinensis*		
				# of the total BLAST hits	(% of total filtered reads)	% Similarity
P4	4	no	SRX3083438	219	(3.7% x 10^{-4})	93□99%
P9	9	no	SRX3083437	56	(3.6% x 10^{-4})	95□100%
P15	15	no	SRX3083436	438	(6.12% x 10^{-4})	96□100%
Ps	4	Adenosine	SRX3083435	356	(4.13% x 10^{-4})	96□100%
Pa	4	Adenine	SRX3083439	207	(6.12% x 10^{-4})	96□99%

Note: Data are adapted from Pang et al*. 2018 http://dx.doi.org/10.1016/j.ygeno.2017.09.008.*

The extremely high sequence similarities between the 5.8S gene sequences of *H. sinensis*, *P.*
hepiali and *T. sinensis,* as shown in Table 11, indicate conservative evolution of the 5.8S gene in
many fungal species. Table 12 shows significant differences in the number of BLAST hits and
percent similarities of the 5.8S gene sequence of *H. sinensis* aligned with the transcriptome shotgun
reads obtained from culture experiments of *P. hepiali* Strain Feng under different culture conditions
and durations (Pang *et al*. 2018). The results indicate the experimental switching on or off of *P.*
hepiali genes, including the 5.8S gene and many other non-rRNA genes, in response to the different
experimental settings.

Liu *et al*. (2015) reported the mRNA transcriptome assembly GCQL00000000, which was
derived from the *H. sinensis* Strain L0106 and contained no rRNA or tRNA transcripts. Table 13
summarizes the dynamic, nonlinear alterations during continuous *in vitro* liquid fermentation in

terms of the total number of transcriptomic unigenes, average length and GC content of transcripts of *H. sinensis* Strain L0106, indicating the nonnatural switching on or off and differential transcriptional activation and silencing of numerous *H. sinensis* genes in response to continued liquid fermentation of *H. sinensis* Strain L0106. As shown in Table 13, more dramatic changes might conceivably occur after 9 days of fermentation. The prolonged 25-day liquid incubation period adopted by Li *et al*. (2013) might have a greater impact on the unbalanced transcriptional activation (switching on) and silencing (switching off) of long and short fungal genes with various GC content biases, in addition to the changes in the culture medium as an important factor influencing the transcriptional activation and silencing of many fungal genes (*cf*. Table 12).

Table 13. *Dynamic alterations of transcripts in the mRNA transcriptome of the* H. sinensis *Strain L0106 during continuous in vitro fermentation.*

Transcriptomic unigene	After 3 days of fermentation	After 6 days of fermentation (% change)	After 9 days of fermentation (% change)
Total number	25,511	25,214 (↓ 1.2%)	16,245 (↓ 36.3%)
Average length (nt)	681	682 (↑ 0.1%)	994 (↑ 46.0%)
GC content (%)	58.2%	57.9% (↓ 0.5%)	57.0% (↓ 2.1%)

Note: Data are adapted from Liu et al. *2015 DOI 10.1186/s12864-015-1269-y and summarized.*

III-7. 5.8S cDNA detected using 5.8S-F/R primers

In addition to the transcriptional silencing of 5.8S genes in natural settings and the significant impact of continuous *in vitro* culture and altered experimental settings on unbalanced gene transcription in experimental settings, as discussed above, the second issue related to the logic described by Li *et al*. (2019) is whether the methods designed by Li *et al*. (2013) were suitable for scientifically identifying 5.8S gene cDNAs of multiple *O. sinensis* genotypes (Li *et al*. 2020b).

For instance, Li *et al*. (2019) retrospectively explained the research design logic of 5.8S-F/R primers by Li *et al*. (2013): "The 5.8S-F/R primers … located in the most conserved region." Accordingly, there is no doubt that such a design would largely reduce the specificity of the primers for 5.8S cDNA because of the conservative evolution of the 5.8S gene and many other fungal genes. Table 14 and Figure 6 show the similarities and alignments between the sequences of primers 5.8S-F (5' ACTTTTAACAACGGATCTCTT 3') and 5.8S-R (5' AAGATAACGCTCGGATAAGCAT 3') and the 5.8S gene of Genotypes #1, #4, and #5.

Table 14. *Similarity of the 5.8S-F/R primers and the 5.8S gene sequences of Genotypes #1, #4 and #5 of* O. sinensis.

Genotype # and GenBank accession #	Percent similarity of the primer *vs*. the 5.8S gene sequence	
	Primer 5.8S-F	Primer 5.8S-R[§]
#1 AB067721	95% (20/21 nt)	85% (17/20 nt)
#4 AB067744	95% (20/21 nt)	90% (18/20 nt)
#5 AB067740	100% (21/21 nt)	95% (19/20 nt)

Note: [§], *Primer 5.8S-R has 2 flanking residues downstream in the ITS2 region.*

```
GT1 AB067721 1    TAGAGGAAGTAAAAGTCGTAACAAGGTCTCCGTTGGTGAACCAGCGGAGGGATCATTATC   60
GT4 AB067744 1    ----------------------------A--A---T-----------T------   56
GT5 AB067740 1    --A----------------T----------A--A---T-----------T------   60

GT1 AB067721 61   GAGTCACCACTCCCAAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCC   120
GT4 AB067744 57   ------------C--------------A--C----------------------------   116
GT5 AB067740 61   ----T--T----TT---------ATA--T--T-T-----------TA------------T   120

GT1 AB067721 121  GGCGCCCCAGGGCCCGGACCAGGGCGCCCGCCGGAGGACCCCCAGACCCTCCTGTCGCAG   180
GT4 AB067744 117  -------T-----------T-------------------TT---------A---T--   176
GT5 AB067740 121  A------T-A---------T-AA-----------------TT---------A-TAT--   180

GT1 AB067721 181  TGGCATCTCTCAGTCAAGAAGCAAGCAAATGAATCAAAACTTTCAACAACGGATCTCTTG   240
GT4 AB067744 177  -A--------T---T--------------A---T-A-ACTTTTAATAACGGATCTCTTA   236
GT5 AB067740 181  -A-T------T---T--------------A---T---ACTTTTAACAACGGATCTCTTA   240
Primer 5.8S-F                                          ACTTTTAACAACGGATCTCTT

GT1 AB067721 241  GTTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCA   300
GT4 AB067744 237  ---T-A----TA--A---------------A------------A-----T------T-   296
GT5 AB067740 241  -----A----TA--A--A-------------A---------A-A----A-------T-   300

GT1 AB067721 301  GTGAACCATCGAATCTTTGAACGCACATTGCGCCCGCCAGCACTCTGGCGGGCATGCCTG   360
GT4 AB067744 297  --A---T-----------A----T-T-C-A-------T--------A------ATGCCTG   356
GT5 AB067740 301  --A---T-----------A------T---A-------T--------A------ATGCCTA   360
Primer 5.8S-R                                                    ATGCTTA

GT1 AB067721 361  TCCGAGCGTCATCTCAACCCTCGAGCCCCCCGCCTCGCGGCGGCGGGGCCCGGCCTTGGG   420
GT4 AB067744 357  TCCGAGCGTTACCTT------TA--------------------------------A--   416
GT5 AB067740 361  TCCGAGCGTTATCTT-----------------------------------T-------A--   420
Primer 5.8S-R     TCCGAGCGTTATCTT

GT1 AB067721 421  GGTCACGGCCCCGCGCCGCCCCCTAAACGCAGTGGCGACCCCGCCGCGGCTCCCCTGCGC   480
GT4 AB067744 417  -----------A-------------------A-T------------------A---   476
GT5 AB067740 421  ---T-----------------T------T---A--------------------A--T   480

GT1 AB067721 481  AGTAGCTCGCTGAGAACCTCGCACCGGGAGCGCGGAGGCGGTCACGCCGTGAAACCACCA   540
GT4 AB067744 477  -A-----TA--A------------TA-----------------------A-------T-   536
GT5 AB067740 481  G-------A-C---------------A--------------------A-------T-   540

GT1 AB067721 541  CACCCTCCAGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACTTAAGCATATCAATAA   600
GT4 AB067744 537  T------TG---A---------T----------------A------------------   596
GT5 AB067740 541  -G--T-TT----A---------T----------------A------------T-----   600

GT1 AB067721 601  GCGGAGGAAAAGAAACCAACAGGGATTGCCCCAGTAACGGCGAGTGAAGCGGCAACAGCT   660
GT4 AB067744 597  --------                                                    604
GT5 AB067740 601  ----------------T--T----------TT-------------A------T--T----   660
```

Figure 6. ITS sequence alignment of AB067721 of GC-biased Genotype #1, AB067744 and AB067740 of AT-biased Genotypes #4□5 of **O. sinensis** *and the 5.8S-F/R primers.*

"GT1", "GT4", and "GT5" represent Genotypes #1 and #4□5 of O. sinensis, *respectively. The sequences in* *blue* *are from the 18S and 28S genes (5' and 3' end regions, respectively). The sequences in* *pink* *represent*

the sequences of 5.8S genes. ITS1 is shown in black between the 18S and 5.8S genes, and ITS2 is shown in black between the 5.8S and 28S genes. The sequences underlined in green are for primers 5.8S-F/R, while primer 5.8S-R was converted to the sense chain sequence. The hyphens indicate identical bases.

The primer pair 5.8S-F/R was 100% and 95% homologous to the 5.8S gene sequence of AT-biased Genotype #5, as shown in Table 14. This indicates that in contrast to the 95% and 85% similarity to the 5.8S gene sequence of GC-biased Genotype #1, the 5.8S-F/R primer pair may be favored for amplifying the 5.8S cDNA sequence of Genotype #5 rather than for amplifying the 5.8S cDNA sequence of Genotype #1. The favorability of the primers toward the 5.8S cDNA sequence of Genotype #5 was clearly shown in Figure 6. The primers showed variable sequence similarity, ranging from 61.9□100% and 31.8□95.5%, respectively, among all 17 genotypes of *O. sinensis*, preferentially amplifying the most easily amplified cDNA sequence in the cDNA library considering the primary structure and secondary conformation of the PCR templates and consequently causing false-negative results for the other 5.8S cDNA sequences.

Section III-6 describes the 100% homology between the 1894 unassembled metatranscriptome shotgun reads (100 bp) of mature natural *C. sinensis* and the 5.8S gene sequence (156 bp) of GC-biased Genotype #1 *H. sinensis*, including many non-rRNA transcripts among the highly homologous 1894 shotgun reads in natural settings. A BLAST search of *O. sinensis* transcriptomic sequences (taxid:72228) in the GenBank database showed that the 5.8S-F/R primer pair was 100% homologous to the mRNA transcriptome assemblies GCQL01001191, GCQL01006777, GCQL01007818, and GCQL01013443 of Genotype #1 *H. sinensis* Strain L0106 (Liu *et al*. 2015), which do not contain rRNA or tRNA transcripts, and to the assembled metatranscriptome sequences GAGW01003691, GAGW01007670, and GAGW01007671 of natural *C. sinensis* (Xiang *et al*. 2014). Figure 7 shows that the metatranscriptome GAGW01003691 is 99.8% homologous to a segment of the mRNA transcriptome GCQL01006777 of the *H. sinensis* Strain L0106; the metatranscriptomes GAGW01007670 and GAGW01007671 are 100% identical to the mRNA transcriptome segments of GCQL01007818 and GCQL01001191, respectively (Li *et al*. 2020b). Because the mRNA transcriptome assembly GCQL00000000 does not contain rRNA or tRNA transcripts, the extremely high homology between the segments of the metatranscriptome GAGW00000000 and transcriptome sequences GCQL00000000 indicates that the 5.8S-F/R primer pair designed by Li *et al*. (2013) may be able to recognize and nonspecifically amplify non-rRNA components in the cDNA library derived from the genetically heterogeneous Clone 1220, one of the 8 heterogeneous clones (1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) of the *C. sinensis*

937   mono-ascospore cultures (Li *et al*. 2013).

938
Figure 7. Graphic alignment summary of the mRNA transcriptome assembly sequence (GCQL00000000)
of the **H. sinensis** *Strain L0106 and the metatranscriptome assembly sequence (GAGW00000000) of*
natural **C. sinensis.**
The graphic alignment summary was generated using GenBank BLAST. GCQL01001191, GCQL01006777,
and GCQL01007818 are sequences of transcriptome segments that were obtained from the mRNA pool of H.
sinensis Strain L0106 and contained no rRNA or tRNA transcripts. GAGW01003691, GAGW01007670, and
GAGW01007671 are segment metatranscriptome sequences obtained from the total RNA of natural C.
sinensis specimens collected from Kangding County of Sichuan Province, China.

Unfortunately, Li *et al*. (2013) did not disclose the sequences of the identified 5.8S cDNA but
instead reported that "Of these clones, 28 showed identical sequences (104 bp excluding the primers)
to the most frequent sequence in group A (JQ900148). The remaining 14 sequences … had single
substitutions at various positions", where group A = Genotype #1 *H. sinensis*. The GenBank
annotation of the 5.8S gene of JQ900148 is at 192→346, covering 155 nt, which is not a multiple of
three and is not in accordance with the open reading frame of the gene.

Alternatively, the 5.8S genes were annotated at the following nucleotide positions: 218→373
within AB067721 of Genotype #1 and AB067740 of Genotype #5, and 214→369 within AB067744
of Genotype #4, covering 156 nt and 52 codons in the 3 cases (Zhu & Li 2017; Li *et al*. 2020b,
2022c). Excluding the 5.8S-F/R primer regions (*cf*. Figure 6), the remaining sequence of the 5.8S
gene consisted of 114 bp, with 10 additional nucleotides (mismatches) beyond the 104 bp in the 28
cDNA clones reported by Li *et al*. (2013). Adding "single substitutions at various positions" to 14
other clones, as reported by Li *et al*. (2013), the 42 clones presumably contained either 10 or 11
mismatches (8.8□9.6%) in the 114-bp segment of the 5.8S gene cDNA, corresponding to
90.4□91.2% sequence similarity to the 5.8S gene of *H. sinensis*. Such low similarities indicated
unlikelihood that the detected 5.8S gene cDNA by Li *et al*. (2013) truly belonged to GC-biased

Genotype #1. Alternatively speaking, nondisclosure of the detected cDNA sequence did not exclude the possibility that the cDNA belonged to AT-biased Genotype #5 of *O. sinensis*.

IV. Discussion

IV-1. The GC-biased repetitive ITS copies in the genome of Genotype #1 *H. sinensis* are unrelated to RIP mutagenesis

Selker and his colleague (1985, 1990, 1997, 2002) reported frequent C-to-T (cytosine-to-thymine) transitions and concomitant epigenetic methylation of nearly all remaining cytosine residues in fungal 5S RNA genes associated with transcriptional silencing. Gladyshev (2017) summarized that RIP "occurs specifically in haploid parental nuclei that continue to divide by mitosis in preparation for karyogamy and ensuing meiosis" and "mutates cytosines on both strands of each DNA duplex in a pairwise fashion (e.g., preferentially affecting none or both repeated sequences)." Hane *et al*. (2017) stated that "the RIP process selectively mutated duplicated sequences in both DNA strands by inducing single-nucleotide point (SNP) mutations that converted C:G base pairs to T:A. This often led to the introduction of nonsense or missense mutations which affected the expression of these sequences." The sequence-specific targets of RIP mutations for *de novo* DNA methylation are under debate (Hood *et al*. 2005; Taranto *et al*. 2014; Hane *et al*. 2017).

Table 3 and Figure 1 demonstrate that the variable repetitive ITS copies were genetically characterized by multiple scattered insertion, deletion and transversion point mutations. These mutations were not generated through RIP mutagenesis, which theoretically causes cytosine-to-thymine (C-to-T) and guanine-to-adenine (G-to-A) transitions. On the other hand, the AT-biased genotypes of *O. sinensis* characterized by multiple scattered transition point mutations (*cf*. Figure 1) were generated in parallel with other GC-biased genotypes through accelerated nuclear rDNA evolution from a common ancestor during a long evolutionary history (Stensrud *et al*. 2007), regardless of whether this occurred *via* RIP mutagenesis or other evolutionary mechanisms. Thus, the evidence is lacking to support the hypothesis that RIP mutagenesis induces the formation of mutant AT-biased repetitive ITS copies within the single genome of GC-biased Genotype #1 *H. sinensis*, postulated by Li *et al*. (2020c). Even if it was proven that AT-biased genotypes were indeed induced by RIP mutagenesis during a long history of evolution, there has been no evidence that AT-biased genotypes "could have emerged either before or after" a new GC-biased "ITS haplotype was generated" through "repeat-induced point mutations" (RIP) and became components

of the single genome of GC-biased Genotype #1, as assumed by Li *et al*. (2013, 2019, 2020c).

Taranto *et al*. (2014) and Hane *et al*. (2015) reported that RIP "is observed only in certain

fungal taxa; the Pezizomycotina (filamentous Ascomycota) and some species of the Basidiomycota."

Tables 1 and 3 and Figures 1□3 show that all the multiple genomic repetitive ITS copies were GC

biases genetically, with no or only a few transition alleles, and were phylogenetically distinct from

AT-biased genotypes, which possess numerous scattered transition alleles. It is thus reasonable to

question whether Genotype #1 *H. sinensis* is susceptible to RIP mutagenesis and epigenetic

methylation attack.

IV-2. Genetic and phylogenetic differences in the AT-biased genotypes of *O. sinensis* and GC-biased repetitive ITS copies in the genome of Genotype #1 *H. sinensis*

The sequences of AT-biased Genotypes #4□6 and #15□17 of *O. sinensis* shown in red in

Figure 1 contain numerous scattered transition alleles. These sequences are genetically distinct and

phylogenetically distant from GC-biased repetitive ITS copies containing the predominant insertion,

deletion, and transversion point mutations within the genome of GC-biased Genotype #1 *H. sinensis*

(*cf*. Figures 1□2 and Tables 1□3). The sequences of AT-biased Genotypes #4□6 and #15□17 of *O.*

sinensis, as well as GC-biased Genotypes #2□3 and #7□14, are absent in the genome assemblies

ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000 of

H. sinensis Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively (*cf*. Tables 1□2

and Figures 1 and 3), but instead belong to the genomes of independent *O. sinensis* fungi (Stensrud

et al. 2007; Xiao *et al*. 2009; Hu *et al*. 2013; Zhu and Wu 2015; Li *et al*. 2016a, 2016b; 2020b,

2022a, 2022c, 2023c; Zhu and Li 2017; Zhu *et al*. 2019; Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*.

2020). The nonresidency measure of AT-biased *O. sinensis* genotypes does not support the

hypothesis that the AT-biased sequences are "ITS pseudogene" components of "a single genome" of

GC-biased Genotype #1 *H. sinensis*. Stensrud *et al*. (2007) concluded that the variation in the 5.8S

gene sequences of AT-biased genotypes "far exceeds what is normally observed in fungi … even at

higher taxonomic levels (genera and family)". Xiao *et al*. (2009) concluded that the sequences of

AT-biased genotypes likely belong to the genomes of independent *O. sinensis* fungi.

IV-3. The previously hypothesized the center of origin and the evolutionary geography for *O. sinensis*

GC-biased genotypes might have evolved at accelerated rates in parallel with AT-biased genotypes from an ancient common ancestor during long-term evolution, as hypothesized by Stensrud *et al*. (2007). Zhang *et al*. (2009) detected GC-biased Genotype #3 in natural *C. sinensis* specimens collected from the Nyingchi District of Tibet and hypothesized that "Nyingchi might have acted as a center of origin of *O. sinensis* and *O. sinensis* might have been transmitted from Nyingchi to other locations". Accordingly, GC-biased Genotype #3 might represent a modern "immediate" ancestor of Genotype #1 *H. sinensis* that has been consistently identified in other production locations as the result of a relatively modern phase of the long-term evolution. The authors described an evolutionary path from GC-biased Genotype #3 to Genotype #1 and the transmission of *O. sinensis* from the "center of origin … spread eastward and westward first among southern regions on the Tibetan Plateau" and then "spread from south to north … at a later time".

Unfortunately, this evolutionary geography for *O. sinensis* lacks an evolutionary branch or branches related to AT-biased genotypes of *O. sinensis*. Perhaps the generations of AT-biased genotypes of *O. sinensis* might have occurred during more ancient historical evolutionary phases. Undoubtedly, the discovery of AT-biased genotypes of *O. sinensis* will trigger further research on the history of *O. sinensis* evolution and fungus-insect coevolution [Zhang *et al*. 2014].

The evolutionary geography hypothesis for *O. sinensis* also overlooks an evolutionary path or paths extending south to (or going north from) the production areas in Bhutan, Sikkim, and Nepal. Biswa *et al*. (2022) identified a mutant sequence (MW990119) from a *C. sinensis* specimen (CBUS3) collected in Yumesamdong (27°84'88" N 88°69'04" E), North Sikkim, which is far from Nyingchi in Tibet (29°38′56″ N, 94 21′41″ E). MW990119 was 98.8% homologous to HM595984 of the representative sequence of GC-biased Genotype #3, which had 3 insertions and 2 transition alleles but was 95.2% similar to AB067721 of Genotype #1 *H. sinensis*. The discovery of the MW990119 sequence may challenge the "center of origin" hypothesis for Nyingchi, Tibet, and the "evolutionary geography" for *O. sinensis*, which may need to be amended, supplemented, or reassessed.

IV-4. Multicellular heterokaryons of natural *C. sinensis* hyphae and ascospores

Bushley *et al*. (2013) reported that *C. sinensis* ascospores are multicellular heterokaryons with mononucleated, binucleated, trinucleated, and tetranucleated cellular structures. Zhang and Zhang (2015) wondered whether the binucleated cells (and other mono- and polykaryotic cells) of *C. sinensis* hyphae and ascospores contained homogenous or heterogeneous genetic materials, meaning

that these mono- and polykaryotic cells may contain heterogeneous sets of chromosomes and genomes.

Li *et al*. (2013) identified heterogeneous ITS sequences of both GC-biased Genotype #1 and AT-biased Genotype #5 in 8 of 15 ascosporic clones (1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) and simultaneously identified 7 other clones (1207, 1218, 1219, 1221, 1222, 1225, and 1229) that contained only homogenous sequences of Genotype #1 after 25 days of *in vitro* inoculation with natural *C. sinensis* mono-ascospores. These culture-dependent molecular mycological findings support the morphological observations of multicellular heterokaryons discovered by their collaborators (Bushley *et al*. 2013), although molecular mycological strategies may overlook nonculturable fungal components. Apparently, the 2 groups of reported ascosporic clones were derived from different cells of multicellular heterokaryotic mono-ascospores of natural *C. sinensis*, confirming the inference of Zhang and Zhang (2015) that there are heterogeneous sets of chromosomes and genomes in mono- and polykaryotic cells.

In addition to the 15 clones reported by Li *et al*. (2013), the authors did not disclose any information on other possible ascosporic clones, namely, clones 1210, 1211, 1212, 1213, 1215, 1216, 1217, 1223, and 1226, for which the clone serial numbers are missing among those of the reported 2 groups (*cf*. Table S1 of Li *et al*. 2013); nor did the authors state whether these 9 possible clones were simultaneously obtained in the same study, why they were not published along with the 15 reported clones, and whether they contained different fungal genetic materials, although these questions and other relevant questions should have been asked during the journal's peer review process and should have been answered by the authors. The Editor-in-Chief of Molecular Phylogenetics and Evolution, Drs. Wildman and Zimmer encouraged discussions of the relevant scientific issues in the journal with the authors (Zhu *et al*. 2019; Li *et al*. 2020a).

In contrast, Li *et al*. (2023c), from a different research group, used culture-independent techniques and reported the cooccurrence of GC-biased Genotypes #1 and #13□14, AT-biased Genotypes #5□6 and #16 of *O. sinensis*, *P. hepiali*, and an AB067719-type fungus in *C. sinensis* ascospores. These findings from nonculture experiments confirmed the genetic heterogeneity of the multicellular heterokaryotic ascospores of natural *C. sinensis*. Zhu *et al*. (2010, 2012), Gao *et al*. (2011, 2012), and Li *et al*. (2023b) demonstrated that the *O. sinensis* genotypes underwent dynamic alterations in the immature, maturing, and mature stromata, the stromal fertile portion (SFP) that is densely covered with numerous ascocarps, and ascospores of natural *C. sinensis*, indicating the

genomic independence of the *O. sinensis* genotypes and that many of the genotypic fungi may not be culturable in the *in vitro* experimental settings established by Li *et al*. (2013).

IV-5. Genomic variations of *H. sinensis* strains

By aggregating the information from molecular and genomic studies (Hu *et al*. 2013; Li *et al*. 2013, 2016a; Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*. 2020), different genomic DNA samples of "pure" *H. sinensis* strains were examined and published and can be summarized and grouped as follows.

(1) Group 1 of *H. sinensis* strains consisted of pure, homokaryotic anamorphic *H. sinensis* strains (1229, CC1406-203, Co18, IOZ07, and ZJB12195) (Hu *et al*. 2013; Li *et al*. 2016a; Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*. 2020). Total genomic DNA was isolated from these strains and subjected to genome-wide sequencing. The genomes contained no AT-biased genotype sequences.

(2) Group 2 of *H. sinensis* strains consisted of 7 clones (Strains 1207, 1218, 1219, 1221, 1222, 1225, and 1229) among 15 clones derived from 25 days of incubation of natural *C. sinensis* mono-ascospores (Li *et al*. 2013). Total genomic DNA was isolated from these strains and shown to contain the homogenous ITS sequence of Genotype #1 *H. sinensis* but included no AT-biased sequences of *O. sinensis* genotypes.

(3) Group 3 of "*H. sinensis*" strains consisted of 8 other clones (Strains 1206, 1208, 1209, 1214, 1220, 1224, 1227, and 1228) obtained from the incubation of *C. sinensis* mono-ascospores for 25 days (Li *et al*. 2013). Total genomic DNA was isolated from these strains, which exhibited genetic heterogeneity, with the coexistence of GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis*.

The *H. sinensis* strains of Groups 1 and 2 were similar, with each strain possessing a homogenous GC-biased Genotype #1 genome, although multiple GC-biased repetitive ITS copies were identified in the genome assemblies JAAVMX000000000 and NGJJ00000000 for *H. sinensis* Strains IOZ07 and CC1406-203, respectively (*cf*. Tables 1 and 3 and Figures 1–3) (Hu *et al*. 2013; Li *et al*. 2013, 2016a; Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*. 2020). Although a single copy of the ITS sequence was identified in the genome assemblies ANOV00000000, LKHE00000000, and LWBQ00000000 of the *H. sinensis* Strains Co18, 1229, and ZJB12195, respectively (Hu *et al*. 2013; Li *et al*. 2016a; Jin *et al*. 2020), the genomes of these strains may theoretically contain many repetitive ITS copies that could have been discarded during the assembly of genome shotgun reads.

Conceivably, if technically possible, reassembling the genome shotgun reads may very likely identify additional repetitive ITS copies in the genomes of Group 1 Strains Co18, 1229, and ZJB12195, among which Strain 1229 is also a Group-2 strain, which was derived from mono-ascospores by Li *et al*. (2013, 2016a). Similarly, the genome sequencing of other Group-2 strains may very likely reveal multiple repetitive ITS copies.

The *H. sinensis* strains in Groups 2 and 3 were derived after 25 days of liquid culture of the same *C. sinensis* mono-ascospores (Li *et al*. 2013); however, they differed genetically. The Group-3 strains may have heterokaryotic mycelial structures, similar to the mycelia derived from microcycle conidiation of *H. sinensis* ascospores initially cultured on solid medium plates for 30 days and subsequently in liquid medium for 10□53 days, as shown in Figures 2□3 of (Xiao *et al*. 2011). Li *et al*. (2013) reported that the Group 3 strains included cocultured GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis* fungi, which were genomically independent and coexisted in heterokaryotic mycelial cells, indicating either diploid/polyploid mycelial cells containing heterogeneous sets of chromosomes/genomes or impure mycelia, each containing a different chromosome/genome. However, the authors did not identify or report any haploid clones containing only AT-biased Genotype #5, indicating that this AT-biased *O. sinensis* fungus may not exhibit independent culturability under *in vitro* experimental settings, although other studies have reported the identification of single AT-biased Genotype #4 or #5 of *O. sinensis* without coculture with GC-biased Genotype #1 in cultures of natural and cultivated *C. sinensis* under different experimental settings (Kinjo and Zang 2001; Mao *et al*. 2013; Wei *et al*. 2016). Unfortunately, Li *et al*. (2013, 2019, 2020c) neglected the differences in genetic material between the strains in Group 2 and Group 3, which were presumably derived from the same or different cells of multicellular heterokaryotic ascospores (Bushley *et al*. 2013; Zhang and Zhang 2015; Li *et al*. 2020b, 2022c, 2023c).

In addition to the genome sequencing of the homogenous Group-2 Strain 1229 and the genome assembly LKHE00000000 by Li *et al*. (2016a), the genome sequencing of one of the heterogeneous Group-3 strains will provide critical genomic evidence to validate the hypothesis that AT-biased Genotype #5 of *O. sinensis* coexists as an "ITS pseudogene … in a single genome" of GC-biased Genotype #1. The authors may also consider a multilocus approach to explore pseudogenes in the heterogeneous genomes of Group-3 strains and broaden the current single locus "ITS pseudogene" hypothesis to multiple metagenomic loci. Unfortunately, the authors elected not to do so since year 2012 after obtaining the 8 Group-3 strains or perhaps failed in such an endeavor due to technical

difficulties in the assembly of shotgun reads because of heterogeneous sequences from more than one set of genomes of independent *O. sinensis* fungi having close evolutionary relationships. Consequently, they elected willingly or unwillingly to forgo the opportunity to obtain and publish solid evidence to validate their "ITS pseudogene" hypothesis that AT-biased Genotype #5 and GC-biased Genotype #1 physically coexist "in a single genome". Studies from other research groups have confirmed that the sequences of all AT-biased genotypes belong to the genomes of independent fungi (Stensrud *et al*. 2007; Xiao *et al*. 2009; Zhu *et al*. 2010, 2012; Yao *et al*. 2011; Li *et al*. 2016b, 2020b, 2022a, 2022c, 2023c) and that heterokaryotic ascospores and hyphae with multiple mononucleated, binucleated, trinucleated, and tetranucleated cells contain heterogeneous sets of chromosomes and genomes (Bushley *et al*. 2013; Zhang and Zhang 2015).

In addition to the 3 groups of *H. sinensis* strains summarized above, many other *H. sinensis* strains have been studied and presumably belong to Groups 1□3 or other groups. For instance, Li *et al*. (2016c) isolated 2 *H. sinensis* strains, CH1 and CH2, from the intestine of healthy larvae of *Hepialus lagii* Yan. These strains exhibited *H. sinensis*-like morphological and growth characteristics but contained GC-biased Genotype #1, AT-biased Genotypes #4□5, and *P. hepiali*. The strains strongly enhanced the inoculation potency of *H. sinensis* by 15- to 39-fold in the larvae (n=100 in each experimental group) of *Hepialus armoricanus* (*P*<0.001). Additionally, as mentioned above, Li *et al*. (2013) did not discuss whether 9 other ascosporic clones (1210, 1211, 1212, 1213, 1215, 1216, 1217, 1223, and 1226) probably simultaneously obtained in the study constituted additional fungal group(s) containing different genetic materials.

IV-6. Genomic independence of AT-biased genotypes of *O. sinensis*

IV-6.1 Differential occurrence of AT-biased genotypes in different compartments of natural *C. sinensis*

A BLAST search of the GenBank database revealed at least 668 ITS sequences of *O. sinensis* (GenBank taxid:72228), 228 of which belong to AT-biased Genotypes #4□6 and #15□17 (Li *et al*. 2016b, 2022c, 2023c; Zhu *et al*. 2018, 2019; Zhu & Li 2017). Stensrud *et al*. (2007) and Zhang *et al*. (2013a) analyzed the phylogeny of 71 and 397 ITS sequences of *O. sinensis*, respectively, and revealed 3 phylogenetic clades, *i.e.*, Groups A□C (GC-biased Genotype #1 and AT-biased Genotypes 4□5), which consisted of correct *O. sinensis* sequences. In contrast to the homogenous Group-A sequences reported by Stensrud *et al*. (2007), the ITS sequences in Group A (Clade A) reported by Zhang *et al*. (2013a) appeared to be heterogeneous and diversified in a minimum

evolution phylogenetic tree, consisting of Genotype #1 and other GC-biased genotypes of *O. sinensis*. Additional GC- and AT-biased *O. sinensis* genotypes have been successively discovered thereafter (*cf*. Table 2) (Xiao *et al*. 2009; Zhang *et al*. 2009; Zhu *et al*. 2010, 2012; Chen *et al*. 2011; Zhu and Wu 2015; Li *et al*. 2013, 2016; 2022a, 2022c, 2023c; Zhu and Li 2017).

Many early studies conducted since 1999 used a single pair of "universal" primers (*e.g*., *ITS1□ITS5*) for PCR and identified one of the *O. sinensis* genotypes in natural *C. sinensis* specimens collected from different production areas or cultivated *C. sinensis* (Engh 1999; Zhao *et al*. 1999; Kinjo and Zang 2001; Liu *et al*. 2001; Stensrud *et al*. 2005; Wei *et al*. 2006, 2016; Chen *et al*. 2011; Mao *et al*. 2013). Importantly, Wei *et al*. (2016) reported an industrial cultivation project and identified a single teleomorphic AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis* and a single teleomorphic GC-biased Genotype #1 in natural *C. sinensis*, indicating the existence of at least 2 *O. sinensis* teleomorphs. Unfortunately, this study revealed a species contradiction between the anamorphic inoculants (reported to be 3 GC-biased Genotype #1 *H. sinensis* strains) and the teleomorphic AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis*. These findings are inconsistent with the hypotheses of a sole anamorph of *O. sinensis* for *H. sinensis* and the sole teleomorph hypothesis of *O. sinensis* proposed by the same key authors 10 years ago (Wei *et al*. 2006).

Other studies optimized and combined the use of a variety of molecular techniques, such as several pairs of genotype-specific primers, touchdown PCR, nested PCR, restriction fragment length polymorphism (RFLP), cloning-based amplicon sequencing, single-nucleotide polymorphism (SNP) mass spectrometry genotyping techniques, etc. These research approaches greatly improved the specificity and efficiency of DNA amplification and detection, and facilitated the identification of differential coexistence and dynamic alterations of AT-biased genotypes as well as GC-biased genotypes of *O. sinensis* in different combinations in the caterpillar body, stroma, stromal fertile portion with numerous ascocarps, and ascospores of natural *C. sinensis* during maturation (*cf*. Table 10) (Xiao *et al*. 2009; Zhu *et al*. 2010, 2012, 2018, 2019; Gao *et al*. 2011, 2012; Yao *et al*. 2011; Li *et al*. 2020b, 2022a, 2023b, 2023c).

Xiao *et al*. (2009) used pairs of genotype-specific primers for the first time and adopted a nested PCR approach to amplify mutant sequences of *O. sinensis* from the stroma and caterpillar body of immature and mature *C. sinensis* specimens collected from geographically remote production areas. The authors reported the consistent occurrence of GC-biased Genotype #1 and *P. hepiali* and the

inconsistent occurrence of AT-biased Genotypes #4□5 of *O. sinensis*. The authors also detected the first sequence (EU555436) of AT-biased Genotype #6 and uploaded it to GenBank. The authors then concluded that the AT-biased genotypes likely belonged to independent fungal species.

Li *et al*. (2023b, 2023c) used several pairs of genotype-specific primers, touch-down PCR, cloning-based PCR amplicon sequencing and MassARRAY SNP mass spectrometry genotyping techniques and identified Genotypes #5□6 and #16 of AT-biased Cluster A (*cf*. Figure 2) in the ascospores of natural *C. sinensis*, cooccurring with GC-biased Genotype #1, *P. hepiali*, and an AB067719-type fungus. The Genotypes #4 and #15 sequences of AT-biased Cluster-B were identified in the stromal fertile portion, which was densely covered with numerous ascocarps, but not in ascospores obtained from the same *C. sinensis* specimens (*cf*. Table 10). Li *et al*. (2023c) also identified GC-biased Genotype #13 in semiejected ascospores and GC-biased Genotype #14 in fully ejected ascospores collected from the same specimens of natural *C. sinensis*. Although Li *et al*. (2020a) amended their inoculant information, the inoculant used in the industrial cultivation project was *O. sinensis*, which was isolated from the ascospore of Chinese cordyceps (*i.e.*, Natural *C. sinensis*) collected from an alpine meadow of Nyingchi, Tibet Autonomous Region, China, but not the 3 pure GC-biased Genotype #1 *H. sinensis* strains reported by Wei *et al*. (2016), Li *et al*. (2023b, 2023c) reported that the fully ejected ascospores of natural *C. sinensis* contained GC-biased Genotypes #1 and #14, AT-biased *O. sinensis* Genotypes #5–6 and #16 within AT-biased Cluster A (*cf*. Figure 2), *P. hepiali* and an AB067719-type fungus. These studies reported that the ascospores did not contain Genotypes #4 and #15 of AT-biased Cluster B. Li *et al*. (2013) made additional efforts to design Genotype #4-specific primers for identifying Genotype #4 of AT-biased Cluster B in mono-ascospores but failed. Thus, Wei *et al*. (2016) and Li *et al*. (2020a) actually left an unsolved puzzle regarding the fungal source of the sole teleomorphic AT-biased Genotype #4 in cultivated *C. sinensis*.

Li *et al*. (2013) did not identify the ITS sequences of AT-biased Genotypes #4, #6, and #15□17 of *O. sinensis* in genomic DNA preparations extracted from the 15 ascosporic strains in both Groups 2 and 3 of *H. sinensis* (*cf*. Section IV-5). Thus, the expansion of the "ITS pseudogene" hypothesis from the identified AT-biased Genotype #5 from the mono-ascospore cultures to all AT-biased genotypes was apparently an overly generalized interpretation with no supporting genomic evidence. As shown in Tables 1□3 and Figures 1□2, genomic and phylogenetic analyses did not support the hypothesis that the "diverged ITS paralogs" (*i.e.*, AT-biased Genotypes #4□6 and #15□17), as the

"ITS pseudogene" components, coexist "in a single genome" of Genotype #1 *H. sinensis*.

IV-6.2 Dynamic alterations of multiple *O. sinensis* genotypes in different compartments of

natural *C. sinensis*

Molecular physiological studies have revealed that the biomasses/abundances of genotypes are

dynamically altered in an asynchronous, disproportional manner during *C. sinensis* maturation,

indicating the genomic independence of multiple genotypes of *O. sinensis* from different

perspectives (Zhu *et al*. 2010, 2012; Gao *et al*. 2011, 2012; Yao *et al*. 2011; Li *et al*. 2016b, 2016c,

2022a, 2022c, 2023c).

Zhu *et al*. (2010) and Gao *et al*. (2011, 2012) reported that Genotype #4 of AT-biased Cluster B

of *O. sinensis* (*cf*. Figure 2) predominated in the immature stroma of natural *C. sinensis,* which was

in its asexual growth stage; however, its abundance dramatically declined in the mature stroma*,*

which was in its sexual reproductive stage. The maturational decline in Genotype #4 abundance in

the *C. sinensis* stroma was accompanied by a significant reciprocal increase in the abundance of

Genotype #5 of *O. sinensis*, which was placed in the AT-biased Cluster A, indicating that the

sequences of the multiple AT-biased genotypes belong to the genomes of independent *O. sinensis*

fungi.

Li *et al*. (2023b) used the MassARRAY SNP mass spectrometry genotyping technique and

found dynamic alterations of GC-biased Genotype #1 and several AT-biased genotypes of *O.*

sinensis in the stromal fertile portion (densely covered with numerous ascocarps) before and after

the ejection of *C. sinensis* ascospores and in fully and semiejected ascospores collected from the

same *C. sinensis* specimens. A highly abundant SNP peak that contained both AT-biased Genotypes

#6 and #15 was shown in the preejection stromal fertile portion of natural *C. sinensis,* but the

abundance dramatically declined in the stromal fertile portion after ascospore ejection, while the

coexisting AT-biased Genotype #5 maintained high abundance.

The ITS sequences of the GC-biased genotypes contained an *Eco*RI endonuclease cleavage site

(G▾AATTC in purple in Figure 1) at the nucleotides 294→299 within AB067721 (a representative

sequence of Genotype #1 *H. sinensis*). The *Eco*RI cleavage site was lost (GAATTT) in all AT-biased

genotypes due to a single-base cytosine-to-thymine (C-to-T) transition mutation (*cf*. Figure 1). To

examine dynamic alterations of the natural biomasses of GC- and AT-biased genotypes of *O.*

sinensis without PCR manipulation that causes nonlinear, exponential amplification of DNA

sequences, the Southern blotting technique was applied to examine genomic DNA after enzymolysis

with the endonucleases *Ava*I, *Dra*I, and *Eco*RI (Zhu *et al.* 2010). This non-PCR approach identified a single *Eco*RI-sensitive DNA moiety in genomic DNA isolated from mycelia of *H. sinensis* (*cf.* Figure 4B of Zhu *et al.* 2010) and a doublet (both *Eco*RI-sensitive GC-biased Genotype #1 and *Eco*RI-resistant AT-biased genotypes) in genomic DNA isolated from natural *C. sinensis* (Figure 8). The lack of *Eco*RI-resistant AT-biased genotypes in pure *H. sinensis* shown in Figure 4B of (Zhu *et al.* 2010) indicates that the sequences of AT-biased genotypes of *O. sinensis* are absent in the genome of a GC-biased Genotype #1 *H. sinensis* strain (a gift from Prof. Guo Y-L), similar to the *H. sinensis* strains in Groups 1 and 2 summarized above in Section IV-5. Because Li *et al.* (2013) incorrectly explained the biochemical findings obtained by Southern blotting (Zhu *et al.* 2010), we provided help to mycologists by reproducing the results (with permission) shown in Figure 8 with detailed explanations. The upper panel of Figure 8 shows the Southern blot using an *H. sinensis*-specific probe that was designed based on the ITS1 sequence of Genotype #1 *H. sinensis* and demonstrates the following:

***Figure 8. Southern blot of* O. sinensis *nrDNA in the stroma and caterpillar body of natural* C. sinensis *during maturation.** [Reproduced with permission from AJBMS (www.nwpii.com/ajbms) Am J Biomed Sci 2010; 2(3): 217-238] [*Zhu* et al. *2010*].*

Caterpillar refers to the caterpillar body. Genomic DNA templates were isolated from the stromata or caterpillar bodies of immature (Im) or mature (M) C. sinensis *specimens collected from Sichuan Province in China and prepared using the restriction enzymes Ava*I, *Dra*I, *and Eco*RI. **_The upper panel_** *was probed with an* H. sinensis-*specific probe that was designed based on the ITS1 sequence of Genotype #1* H. sinensis. **_The lower panel_** *was probed with a nonspecific 18S internal control probe that was designed based on the 18S gene sequence of Genotype #1* H. sinensis, *which is >700 bp upstream of the ITS1 segment.*

(1) The *O. sinensis* genotypes identified in the stromata: the left 2 lanes of the upper panel of Figure 8 show that the slower-migrating DNA moieties (*Eco*RI-resistant AT-biased genotypes) predominate naturally without PCR amplification in the stromata of immature (IM) and mature (M) *C. sinensis*. The faster-migrating moieties (*Eco*RI-sensitive GC-biased *H. sinensis*) are always minor moieties in the stromata during *C. sinensis* maturation.

(2) The *O. sinensis* genotypes identified in the caterpillar bodies: although the second lane from the

right in the upper panel of Figure 8 seems to show a single *Eco*RI-sensitive moiety in the immature (IM) caterpillar body but a doublet in the mature (M) caterpillar body, as shown in the far right lane, PCR experiments using genotype-specific primers and touch-down and nested PCR protocols were able to identify *Eco*RI-resistant AT-biased genotypes in the immature caterpillar body. These results suggest that the natural quantity of *Eco*RI-resistant AT-biased genotypes in the immature caterpillar body was less than the detection limit of the Southern blotting technique, which is much less sensitive than PCR-based techniques. The quantity of AT-biased genotypes dramatically and disproportionally increased from a microquantity in the immature caterpillar body to a level comparable to that of *Eco*RI-sensitive GC-biased *H. sinensis* in the mature caterpillar body.

(3) The *O. sinensis* genotypes identified in the stroma and caterpillar body of immature *C. sinensis*: the first and third lanes from the left in Figure 8 show that the abundances and density ratios of *Eco*RI-resistant AT-biased genotypes and *Eco*RI-sensitive GC-biased *H. sinensis* differed significantly in the stroma and caterpillar body of immature *C. sinensis*, which were sampled from the same specimens.

(4) The *O. sinensis* genotypes identified in the stroma and caterpillar body of mature *C. sinensis*: the first and third lanes from the right in Figure 8 show that the abundances and density ratio of *Eco*RI-resistant and *Eco*RI-sensitive genotypes differed in the stroma and caterpillar body of mature *C. sinensis*, which were sampled from the same specimens.

The disproportional distribution of GC- and AT-biased *O. sinensis* genotypes in the stromata and caterpillar bodies of natural *C. sinensis* and their asynchronous, disproportional alterations of the natural biomass without PCR amplification during *C. sinensis* maturation indicate that the sequences of AT-biased *O. sinensis* genotypes do not reside in the single genome of Genotype #1 *H. sinensis*. Instead, the genomically independent AT-biased *O. sinensis* genotypes are interindividually accompanied by GC-biased Genotype #1 *H. sinensis* in the stroma and caterpillar body of natural *C. sinensis*.

Zhu *et al*. (2010, 2012) also used the *Eco*RI enzymolysis approach and restriction fragment length polymorphism (RFLP) technique and examined GC-biased Genotypes #1 and #2 of *O. sinensis* in *C. sinensis* stromata. The authors demonstrated dynamic but asynchronous maturational alterations of cooccurring GC-biased Genotypes #1 and #2 in the stromata of immature, maturing, and mature *C. sinensis* specimens, indicating genomic independence of these 2 GC-biased

genotypes of *O. sinensis*. Figures 2 and 3 above show that *O. sinensis* Genotype #2 is genetically and phylogenetically distinct from the repetitive ITS copies identified in the genome of Genotype #1 *H. sinensis*, further confirming the genomic independence of the 2 GC-biased genotypes and ruling out the possibility that Genotype #2 represents a repetitive ITS copy in the single *O. sinensis* genome.

If "Different types of ITS sequences were thus proved to coexist in a single genome", as assumed by Li *et al*. (2013), the distributions of GC- and AT-biased *O. sinensis* genotypes should be proportional in the *C. sinensis* stroma and caterpillar body, and any biomass alterations of the cooccurring genotypes should be synchronous and proportional during *C. sinensis* maturation. Southern blotting, as well as RFLP after pretreatment with *Eco*RI for enzymolysis and MassARRAY SNP mass spectrometry genotyping assays, clearly demonstrated asynchronous and disproportional alterations of the GC- and AT-biased *O. sinensis* genotypes (Zhu *et al*. 2010, 2012; Gao *et al*. 2011, 2012; Yao *et al*. 2011; Li *et al*. 2023b, 2023c), and Figures 1□3 and Tables 2 and 11 in this paper further confirmed the genomic independence of the multiple *O. sinensis* genotypes, the sequences of which belong to independent *O. sinensis* fungi (Stensrud *et al*. 2007; Xiao *et al*. 2009).

After the hypothesis of "ITS pseudogenes … in a single genome" was published and subjected to academic discussion, Li *et al*. (2013) had sufficient time to validate their hypothesis by conducting genome sequencing on one of the Group 3 strains (*cf*. Section IV-5). However, scientific papers addressing this topic have yet to appear. Instead, Li *et al*. (2013, 2019, 2020c) repeatedly stated that "Different types of ITS sequences were thus proved to coexist in a single genome", "from a single genome, and not from different genomes", and "The coexistence of diverged ITS paralogs in a single genome". These scientists either did not conduct genome sequencing of one of the Group 3 strains (*cf*. Section IV-5) or may have failed the experiments due to the high degree of similarity of the genomic sequences of multiple genotypes of *O. sinensis* that evolved from a common ancestor during a long evolutionary history, which caused difficulties in assembling shotgun reads to verify their hypothesis that Genotypes #1 and #5 coexist in "a single genome", although this genome sequencing strategy may not be able to confirm the coexistence of AT-biased Genotypes #4, #6, and #15□17 as the genomic components of "a single genome" of GC-biased Genotype #1 *H. sinensis*.

IV-7. Possible synergistic effects of several genotypic fungi on *O. sinensis* sexual

reproduction

Based on the identification of mating-type genes of both *MAT1-1* and *MAT1-2* idiomorphs in 2 *H. sinensis* strains, Hu *et al*. (2013) and Bushley *et al*. (2013) proposed self-fertilization as the reproductive mode of Genotype #1 *H. sinensis* under homothallism or pseudohomothallism. Accordingly, Li *et al*. (2020c) stated that "*O. sinensis* is one of the only known homothallic species in *Ophiocordycipitaceae* … the homothallic lifestyle may have increased the likelihood and frequency of sex in *O. sinensis*, leading to increased RIP in this species." However, Zhang and Zhang (2015) revealed differential occurrence of the mating-type genes of *MAT1-1* and *MAT1-2* idiomorphs in numerous *H. sinensis* strains or isolates, destabilizing the foundation of the self-fertilization hypothesis for *O. sinensis*, and the authors alternatively hypothesized facultative hybridization as the mode of *O. sinensis* reproduction.

In addition to the analysis of general intraspecific genetic variations and differential expression of *H. sinensis* genes (Li *et al*. 2022b, 2022c), Li *et al*. (2023a) focused on the differential occurrence of mating-type genes of the *MAT1-1* and *MAT1-2* idiomorphs of 236 *H. sinensis* strains: 22 strains (9.3%) contained only the MAT1-1-1 gene, 65 (27.5%) possessed only the MAT1-2-1 gene, and 149 (63.1%) had both genes. The differential occurrence of mating-type genes underlies the genetic control of *O. sinensis* sexual reproduction. The authors also identified differential but reciprocal transcription of mating-type genes in two *H. sinensis* strains. The MAT1-1-1 transcript was absent, but the MAT1-2-1 transcript was present in the transcriptome assembly GCQL00000000 of the *H. sinensis* Strain L0106 (Liu *et al*. 2015; Li *et al*. 2023a). The MAT1-1-1 transcript was present in the *H. sinensis* Strain 1229, but the MAT1-2-1 transcript included an unspliced intron I, which contains 3 stop codons (Bushley *et al*. 2013; Li *et al*. 2023a). Bushley *et al*. (2013) specifically commented that a "comparison of DNA and cDNA sequences of MAT1-2-1 revealed only one spliced intron of 55 bp (*i.e.*, intron II)" in *H. sinensis* Strain 1229, which "did not result from contamination of genomic DNA". Thus, the results indicated the production of a largely truncated and dysfunctional MAT1-2-1 protein lacking the majority of the protein encoded by exons II and III. The transcriptional silencing of one of the mating-type genes and the MAT1-2-1 transcript with an unspliced intron I constitute *O. sinensis* reproduction controls at the transcriptional and coupled transcriptional-translational levels.

Accordingly, Li *et al*. (2023a) demonstrated the self-sterility of GC-biased Genotype #1 *H. sinensis*, suggesting that mating partners are required for *O. sinensis* sexual reproduction. These

mating partners could be intraspecific species, such as *H. sinensis* Strains L0106 and 1229, for physiological heterothallism, regardless of whether *H. sinensis* is monoecious or dioecious, or heterospecific fungal species that may undergo hybridization if they are able to overcome interspecific reproductive barriers (Zhang and Zhang 2015; Li *et al*. 2022c, 2023a, 2023c). The taxonomic positions of Genotypes #2–17 of *O. sinensis* have not been determined, especially for AT-biased genotypes belonging to different *O. sinensis* fungi "even at higher taxonomic levels (genera and family)" (Stensrud *et al*. 2007) and GC-biased Genotypes #2 and #7 of *O. sinensis*, which cooccur with Genotype #1 in the same specimens of natural *C. sinensis* (Zhu *et al*. 2010, 2012; Chen *et al*. 2011). The coexistence of several AT-biased genotypes with GC-biased Genotype #1 in the ascocarps and ascospores of natural *C. sinensis* may constitute a genetic basis for sexual reproduction of *O. sinensis* involving fungal hybridization (Li *et al*. 2023a, 2023b, 2023c).

In addition, Barseghyan *et al*. (2011) reported the findings of macro- and micromycology studies and concluded that both *H. sinensis* and *T. sinensis* were anamorphs of *O. sinensis*. Engh (1999) hypothesized that *Cordyceps* and *Tolypocladium* form a fungal complex in natural *C. sinensis*, and the *Cordyceps* sequence AJ786590 was discovered by Engh (1999) and reported and uploaded to the GenBank database by Stensrud *et al*. (2005), a Norwegian molecular mycology group. Using a Bayesian phylogenetic algorithm, Stensrud *et al*. (2007) clustered AJ786590 and other ITS sequences (AB067741□AB067749, BD167325) into the Group-B clade of *O. sinensis*, which belong to AT-biased Genotype #4 and are genetically and phylogenetically distinct from GC-biased Genotype #1 *H. sinensis* in the Group-A clade (*cf*. Figures 1□2).

Li *et al*. (2023c) discovered GC-biased Genotypes #13□14 of *O. sinensis*, which exhibit the genetic characteristics of large DNA segment reciprocal substitutions and genetic material recombination between the genomes of the 2 parental fungi, *H. sinensis* and an AB067719-type fungus. The discovery of Genotypes #13□14 of *O. sinensis* likely indicates the possibility of fungal hybridization or parasexuality between the 2 heterospecific parental fungi.

IV-8. Transcription of the 5.8S gene and PCR amplicons of the 5.8S-F/R primers

The analysis in Section III-6 indicates that Li *et al*. (2013, 2019) have neglected the evidence of transcriptional silencing of 5.8S genes demonstrated in metatranscriptome studies in natural settings and the evidence of dynamic and unbalanced transcriptional activation and silencing of numerous fungal genes during the continuous *in vitro* fermentation or liquid incubation of Genotype #1 *H. sinensis* (Xiang *et al*. 2014; Liu *et al*. 2015; Xia *et al*. 2017; Li *et al*. 2022b, 2022c). Changes in *in*

vitro culture conditions and durations have significant impacts on the transcription of fungal genes (*cf*. Tables 12 and 13), which involves the transcriptional activation and silencing of numerous fungal genes. Because of the conservative evolution of 5.8S genes and many other non-rDNA genes, the incomplete study design and controversial findings reported by Li *et al*. (2013) need to be verified using other sophisticated transcription methods whether the cDNA identified by Li *et al*. (2013) was truly derived from the 5.8S gene of GC-biased Genotype #1 *H. sinensis*, from mutant AT-biased Genotype #5 of *O. sinensis*, from other colonized fungi, or even from non-rRNA genes before confidently declaring that the 5.8S genes of the multiple AT-biased genotypes of *O. sinensis* are truly permanently nonfunctional pseudogenes under the "ITS pseudogene … in a single genome" hypothesis.

Li *et al*. (2013) did not disclose the amplicon sequences amplified using 5.8SF/R primers and the PCR template of the cDNA library constructed through reverse transcription of total RNA derived from *H. sinensis* Clone 1220. Li *et al*. (2019) confirmed that Li *et al*. (2013) designed 5.8SF/R primers based on the sequence "in the most conserved region" of *H. sinensis* rRNA, meaning low specificity of the primers and resulting in uncertain genetic sources of the PCR amplicons. According to the sequence analysis in Section III-7, the amplicon sequences showed only 90.4□91.2% similarity to the 5.8S gene sequence of Genotype #1 of *O. sinensis*, and such high dissimilarity rates (8.8□9.6%) may not be convincingly attributed to "mismatches during reverse transcription or PCR amplification", as explained by Li *et al*. (2013). Thus, solid evidence is lacking to conclude that the amplicon sequences were truly derived from the 5.8S gene of Genotype #1 *H. sinensis,* and that the *H. sinensis* 5.8S gene was considered the functional gene, while the 5.8S gene of Genotype #5 of *O. sinensis* was considered the nonfunctional "ITS pseudogene".

Prior to declaring that the 5.8S genes of all AT-biased *O. sinensis* genotypes are permanently nonfunctional pseudogenes cooccurring with a functional 5.8S gene in the genome of GC-biased Genotype #1, several other concerns should be addressed.

(1) Selker and his colleague (1985, 1990, 1997, 2002), Taranto *et al*. (2014), Hane *et al*. (2015) and Gladyshev (2017) have repeatedly reported epigenetic methylation of nearly all cytosine residues in 5.8S RNA genes and subsequent transcriptional silencing of GC-biased genes, which occur only in certain fungal species. As shown in Table 3 and Figures 1 and 3 above, the multiple genomic repetitive ITS copies of GC-biased Genotype #1 *H. sinensis* contained mainly insertion, deletion and transversion alleles but no or only a few transition alleles. The repetitive

ITS copies of Genotype #1 are genetically and phylogenetically distinct from the AT-biased genotypes of *O. sinensis* (Figures 1–3). Although the 5.8S gene of GC-biased Genotype #1 contains many cytosine residues and is potentially more susceptible to epigenetic methylation attack, introducing nonsense or missense mutations that subsequently cause translational silencing of 5.8S genes, the evidence presented in this paper indicates that *H. sinensis* species may not be the target of RIP mutagenesis and epigenetic methylation attack. It is unlikely that the AT-biased genotypes of *O. sinensis* emerged right before or after the generation of a new *H. sinensis* genome; instead, they are likely genomically independent and exist in different *O. sinensis* fungi.

(2) Li *et al*. (2013) did not detect the ITS sequences of Genotypes #2–4 and #6–17 of *O. sinensis* in the genomic DNA pool of mono-ascosporic cultures. Thus, it is an apparent overgeneralization to infer that the 5.8S genes of these genotypes are nonfunctional "ITS pseudogene" components of the genome of Genotype #1 *H. sinensis*.

(3) The sequences of Genotypes #2–17 are absent in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000 of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-2031229, respectively (Hu *et al*. 2013; Li *et al*. 2016a; Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*. 2020), indicating genomic independence of multiple *O. sinensis* genotypes.

(4) The culture-dependent approach used by Li *et al*. (2013) might have overlooked some fungal species that are nonculturable or difficult to culture under the *in vitro* experimental settings of the study. The culture-dependent strategy might have a significant impact on the transcription of many fungal genes, with many genes being switched on or off nonnaturally, as evidenced by the nonlinear reduction in the total number of transcriptomic unigenes and increases in the average length but decreases in the GC content of transcripts during 3□9 days of liquid fermentation of the *H. sinensis* Strain L0106 (*cf*. Table 13) (Liu *et al*. 2015). Table 12 also demonstrates that changes in the culture duration from 4□15 days and the use of culture medium supplemented with or without adenosine or adenine had dramatic impacts on fungal gene transcription (Pang *et al*. 2018). A much greater impact on differential transcriptional activation and silencing of many *O. sinensis* genes may be expected after the prolonged 25-day liquid incubation period adopted by Li *et al*. (2013).

(5) Three distinct types of secondary steric conformations of 5.8S rRNA were predicted for Groups

A–C (*i.e.*, Genotypes #1 and #4–5) of *O. sinensis* by Li *et al*. (2013). The possibility of producing circular RNA through back-splicing-exon circularization (Dong *et al*. 2014a) should be considered during study design because these distinct steric structures may have a considerable impact on reverse transcription PCR and cDNA sequencing. In addition, the ITS1-5.8S-ITS2 sequences of Genotypes #2 and #6 may adopt different types of steric conformations (Zhu & Wu 2015; Li *et al*. 2016b, 2022c; Zhu & Li 2017); thus, the design of other genotype-specific primers and the combined use of other molecular techniques should be considered.

(6) Wei *et al*. (2016) reported the identification of a single teleomorph of AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis* and a single teleomorph of GC-biased Genotype #1 in natural *C. sinensis* (*cf*. Figure 6 of Wei *et al*. 2016). If AT-biased Genotype #4 represents a nonfunctional ITS pseudogene, as believed by Li *et al*. (2013, 2019), the single AT-biased *O. sinensis* teleomorph in cultivated *C. sinensis* might have disturbed teleomorphic functions, leading to reproductive sterility and an abnormal, disturbed lifecycle of cultivated *C. sinensis*. Wei *et al*. (2016) did not share any information about ascospore production in cultivated *C. sinensis*, which is the critical feature of *O. sinensis* sexual reproduction in natural and cultivated *C. sinensis* (Jiang and Yao 2003).

(7) Li *et al*. (2022c, 2023c) reported the identification of teleomorphic Genotypes #4 and #15 of *O. sinensis* of AT-biased Cluster B in the stromal fertile portion densely covered with numerous ascocarps but not in ascospores collected from the same pieces of natural *C. sinensis*. AT-biased Genotypes #6 and #15 were found at high abundance in the stromal fertile portion prior to ascospore ejection, and their abundance drastically declined after ascospore ejection, while teleomorphic Genotype #5 maintained its high abundance in the stromal fertile portion before and after ascospore ejection (Li *et al*. 2022a, 2023b).

(8) The 5.8S genes of multiple *O. sinensis* genotypes may be transcriptionally activated or silenced in different developmental and maturational stages of natural *C. sinensis*. Significant changes in the proteomic expression profiles of the stroma and caterpillar body of natural *C. sinensis* during maturation provide evidence of such dynamic alterations in epigenetic, transcriptional, translational, and posttranslational modifications of numerous genes (Dong *et al*. 2014b).

Apparently, Li *et al*. (2013) improperly designed the study and provided insufficient and inconclusive evidence to support their "ITS pseudogene" hypothesis. The whole package of gene expression in *O. sinensis* fungi and natural *C. sinensis* needs to be further explored before the 5.8S

genes of AT-biased genotypes of *O. sinensis* can be defined as permanently nonfunctional pseudogenes.

IV-9. Repetitive copies of multiple loci in the genome of Genotype #1 *H. sinensis*

In addition to the reassessment of the "ITS pseudogene" hypothesis and the underlying "RIP" as the root cause of pseudogenic mutagenesis by Li *et al*. (2013, 2020c) through single ITS-locus analysis of 5 *H. sinensis* genomes, we extended the genomic repetitive copy analysis to multiple authentic *H. sinensis* genes and their transcripts at multiple genomic loci in the *H. sinensis* genome assemblies, namely, ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000. Based on the high degrees of sequence similarity between the fungi in the *Ophiocordyceps* genus, authentic *H. sinensis* genes were positioned and annotated in the *H. sinensis* genome through cross-reference annotation of the 1271 authentic genes in the JAACLJ010000002 genome, which is one of the 13 genomic contigs of the *O. camponoti-floridani* Strain EC05 (Will *et al*. 2020).

Of the 1271 genes in the *H. sinensis* genome, 104 had repetitive copies (*cf*. Table 4), of which 37 had repetitive genomic copies in only one *H. sinensis* genome, and 67 had multiple repetitive copies in all 5 *H. sinensis* genomes. Tables 5–9 show 5 representative *H. sinensis* genes with a variety of point mutations, including the postulated RIP-related transitions (C-to-T & G-to-A) and opposite transitions (T-to-C & A-to-G), as well as transversions (C-to-A & G-to-T) and opposite transversions (A-to-C & T-to-G), in addition to insertions and deletions. The point mutations caused low sequence similarity of repetitive copies compared with the sequences of authentic genes and increases, decreases or bidirectional changes in the total AT conten.

In general, the repetitive copies of most authentic genes (83 of 104) at multiple *H. sinensis* genomic loci contained fewer postulated RIP-related transitions (C-to-T & G-to-A) than opposite transitions (T-to-C & A-to-G). In contrast, the repetitive copies with large increases in AT content contained more RIP-related transitions (C-to-T & G-to-A) than opposite transitions (T-to-C & A-to-G) in only 11 of 104 authentic genes (*cf*. Table 8). The slight increases in AT content in the repetitive copies of 8 authentic genes (*cf*. Table 7) were the result of transversion mutations; however, the postulated RIP-related transitions (C-to-T & G-to-A) were less frequent than opposite transitions (T-to-C & A-to-G).

As shown in Tables 5–9, the transcripts for the genomic repetitive copies of the authentic genes could be identified in the mRNA transcriptome GCQL00000000 for *H. sinensis* Strain L0106,

regardless of the increases, decreases, or bidirectional changes in AT content and low sequence
similarity of the repetitive copies compared with the authentic genes. The repetitive genomic copies
likely encode different proteins rather than the proteins encoded by the authentic genes. The
transcription feature rules out the possibility that the nonsense and missense point mutations in
repetitive copies cause concurrent epigenetic methylation and lead to dysfunction of the repetitive
copies, regardless of whether the RIP-related point mutations involved in the long-history
evolutionary processes.

Thus, the results of multilocus analysis further invalidated the pseudogene hypothesis and its
root cause of "RIP" mutagenesis, as proposed by Li *et al*. (2013, 2020c). Thus, *O. sinensis*,
especially GC-biased Genotype #1 *H. sinensis*, may not be the mutagenetic target of "RIP", which
causes the nonsense and missense mutations, concurrent epigenetic methylation attack, and
subsequent dysfunction of the authentic *O. sinensis* genes.

V. Conclusion

This study analyzed multiple repetitive ITS copies in the genome of GC-biased Genotype #1 *H.*
sinensis and multiple genotypes of *O. sinensis* fungi and evaluated the "ITS pseudogene … in a
single genome" hypothesis for all AT-biased *O. sinensis* genotypes (Li *et al*. 2013, 2019, 2020c).
The repetitive ITS copies in the genome of Genotype #1 contained multiple scattered insertion,
deletion and transversion point mutations and may not be generated through RIP mutagenesis
because RIP mutagenesis theoretically causes C-to-T and G-to-A transitions. The GC-biased
repetitive ITS copies are genetically and phylogenetically distinct from AT-biased *O. sinensis*
genotypes that contain multiple scattered transition alleles. Genotype #1 *H. sinensis* may not be the
fungal species target of RIP mutagenesis or concurrent epigenetic methylation attack. The AT-biased
genotypes are independent of the genome of Genotype #1 and might have been generated from a
common ancestor through certain mutagenic and evolutionary mechanisms during the long course
of evolution in response to the extremely harsh environment of the Qinghai-Tibet Plateau and
become independent *O. sinensis* fungi with independent genomes coexisting in parallel with GC-
biased genotypes in natural *C. sinensis*. The analysis of multiple genomic loci revealed numerous
repetitive copies of 104 of 1271 authentic *H. sinensis* genes. The repetitive copies contained various
transition, transversion, insertion, and deletion point mutations, causing decreases, increases, or
bidirectional changes in AT content. However, the repetitive copies were functionally transcribed,
regardless of whether the point mutations were caused by RIP-related or -nonrelated mutagenic,

evolutionary mechanisms. Thus, Li *et al*. (2013, 2020c) provided unsound and controversial information to support their "ITS pseudogene … in a single genome" and the causative "RIP" mutagenesis hypotheses, which have not be validated by genomic and transcriptomic evidence.

VI. Acknowledgments:

This research was supported by a grant from the Science and Technology Department of Qinghai Province, China, grant number 2021-SF-A4 "Study on key technologies of conservation of natural resource and industrial upgrading of *Cordyceps sinensis*", the major science and technology projects in Qinghai Province.

References

[1] Barseghyan, G.S., Holliday, J.C., Price, T.C., Madison, L.M., Wasser SP., 2011. Growth and cultural-morphological characteristics of vegetative mycelia of medicinal caterpillar fungus *Ophiocordyceps sinensis* G.H. Sung *et al*. (Ascomycetes) isolates from Tibetan Plateau (P.R.China). Intl. J. Med. Mushrooms, 13(6),.565–581. doi: 10.1615/intjmedmushr.v13.i6.90

[2] Biswa, V., Ali1, R., Pradhan, D.K., Boro, N., Baro, T., Das, D., Subba, R., Das, P., Das, S., 2021. Molecular characterization, microscopic characteristics, and phylogenetic analysis of Ophiocordyceps sinensis from Sikkim, India. Indian J. Hill. Farming, 34,.268☐275.

[3] Bushley,. K.E., Li, Y., Wang, W.-J., Wan,g X.-L., Jiao, L., Spatafora, J.W., Yao, Y.-J., 2013. Isolation of the MAT1-1 mating type idiomorph and evidence for selfing in the Chinese medicinal fungus *Ophiocordyceps sinensis*. Fungal Biol. 117, 599–610. http://dx.doi.org/10.1016/j.funbio.2013.06.001

[4] Chen, C.-S., Hseu, R.-S., Huang, C.-T., 2011. Quality control of *Cordyceps sinensis* teleomorph, anamorph, and Its products. Chapter 12, *in* (Shoyama Y, Ed.) *Quality Control of Herbal Medicines and Related Areas*. InTech, Rijeka, Croatia. www.intechopen.com

[5] Chen, Y.-Q., Hu, B., Xu, F., Zhang, W., Zhou, H., Qu, L.-H., 2004. Genetic variation of *Cordyceps sinensis*, a fruit-body-producing entomopathogenic species from different geographical regions in China. FEMS Microbiol. Lett. 230, 153–158. Doi: 10.1016/S0378-1097(03)00889-9

[6] Dong, R., Chen, L.-L., Yang, L., 2014a. Circular RNA studies in the post-genomic era. Chin. J. Cell Biol. 36(11), 1455–1459. DOI: 10.11844/cjcb.2014.11.9001

[7] Dong, Y.-Z., Zhang, L.-J., Wu, Z.-M., Gao, L., Yao, Y.-S., Tan, N.-Z., Wu, J.-Y., Ni, L.-Q., Zhu, J.-S., 2014b. Altered proteomic polymorphism in the caterpillar body and stroma of natural *Cordyceps sinensis* during maturation. PLoS ONE*,* 9(10), e109083. doi:10.1371/journal.pone.0109083

[8] Engh, I.B., 1999. Molecular phylogeny of the *Cordyceps-Tolypocladium* complex. Candidate scientific thesis, Department of Biology, University of Oslo, Oslo. 1999.

[9] Gao, L., Li, X.-H., Zhao, J.-Q., Lu, J.-H., Zhu, J.-S., 2011. Detection of multiple *Ophiocordyceps sinensis* mutants in premature stroma of *Cordyceps sinensis* by MassARRAY SNP MALDI-TOF mass

spectrum genotyping. Beijing Da Xue Xue Bao, 43(2), 259–266. doi: 10.3969/j.issn.1671-
167X.2011.02.019

[10] Gao, L., Li, X.-H., Zhao, J.-Q., Lu, J.-H., Zhao, J.-G., Zhu, J.-S., 2012. Maturation of *Cordyceps*
sinensis associates with alterations of fungal expressions of multiple *Ophiocordyceps sinensis* mutants
with transition and transversion point mutations in stroma of *Cordyceps sinensis*. Beijing Da Xue Xue
Bao, 44(3), 454–463. doi: 10.3969/j.issn.1671-167X.2012.03.025

[11] Gladyshev, E., 2017. Repeat-Induced Point Mutation (RIP) and Other Genome Defense Mechanisms in
Fungi. Microbiol Spectr 5(4), 10.1128. doi: 10.1128/microbiolspec.FUNK-0042-2017

[12] Hane, J.K., Williams, A.H., Taranto, A.P., Solomon, P.S., Oliver, R.P., 2015. Repeat-induced point
mutation: A fungal-specific, endogenous mutagenesis process. In: van den Berg M, Maruthachalam K
(eds) Genetic transformation systems in fungi, Volume 2. Fungal Biology. Springer, Cham.
https://doi.org/10.1007/978-3-319-10503-1_4

[13] Hawksworth, D.L., Crous, P.W., Redhead, S.A., Reynolds, D.R., Samson, R.A., Seifert, K.A., Taylor,
J.W., Wingfield, M.J., Abaci, O., Aime, C., Asan, A., Bai, F.-Y., de Beer, Z.W., Begerow, D., Berikten,
D., Boekhout, T., K Buchanan, P., Burgess, T., Buzina, W., Cai, L., Cannon, P.F., Crane, J.L., Damm, U.,
Daniel, H.-M., van Diepeningen, A.D., Druzhinina, I., Dyer, P.S., Eberhardt, U., Fell, J.W., Frisvad, J.C.,
Geiser, D.M., Geml, J., Glienke, C., Gräfenhan, T., Groenewald, J.Z., Groenewald, M., de Gruyter, J.,
Guého-Kellermann, E., Guo, L.-D., Hibbett, D.S., Hong, S.-B., de Hoog, G.S., Houbraken, J., Huhndorf,
S.M., Hyde, K.D., Ismail, A., Johnston, P.R., Kadaifciler, D.G., Kirk, P.M., Kõljalg, U., Kurtzman, C.P.,
Lagneau, P.-E., Lévesque, C.A., Liu, X., Lombard, L., Meyer, W., Miller, A., Minter, D.W., Najafzadeh,
M.J., Norvell, L., Ozerskaya, S.M., Oziç, R., Pennycook, S.R., Peterson, S.W., Pettersson, O.V.,
Quaedvlieg, W., Robert, V.A., Ruibal, C., Schnürer, J., Schroers, H.-J., Shivas, R., Slippers, B.,
Spierenburg, H., Takashima, M., Taşkın, E., Thines, M., Thrane, U., Uztan, A.H., van Raak, M., Varga,
J., Vasco, A., Verkley, G., Videira, S.I.R., de Vries, R.P., Weir, B.S., Yilmaz, N., Yurkov, A., Zhang, N.,
2011. The Amsterdam declaration on fungal nomenclature. IMA Fungus, 2, 105–112. DOI:
10.5598/imafungus.2011.02.01.14

[14] Hood, M.E., Katawczik, M., Giraud, T., 2005. Repeat-induced point mutation and the population
structure of transposable elements in Microbotryum violaceum. Genetics, 170(3), 1081-1089. doi:
10.1534/genetics.105.042564

[15] Hu, X., Zhang, Y., Xiao, G., Zheng, P., Xia, Y., Zhang, X., St Leger, R.J., Liu, X., Wang, C., 2013.
Genome survey uncovers the secrets of sex and lifestyle in caterpillar fungus. Chin. Sci. Bull, 58, 2846–
2854. doi: 10.1007/s11434-013-5929-5

[16] Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogeny. Bioinformat, 17,
754–755. DOI: 10.1093/bioinformatics/17.8.754

[17] Jiang, Y., Yao, Y.-J., 2003. Anamorphic fungi related to *Cordyceps sinensis*. Mycosystema 22(1),
161–176.

[18] Jin, L.□Q., Xu, Z.□W., Zhang, B., Yi, M., Weng, C.□Y., Lin, S., Wu, H., Qin, X.□T., Xu, F., Teng, Y.,

Yuan, S.□J., Liu, Z.-Q., Zheng, Y.-G., 2020. Genome sequencing and analysis of fungus *Hirsutella*
sinensis isolated from *Ophiocordyceps sinensis*. AMB Expr. 10, 105. DOI: 10.1186/S13568-020-01039-
x

[19] Kinjo, N., Zang, M., 2001. Morphological and phylogenetic studies on *Cordyceps sinensis* distributed in
southwestern China. Mycoscience, 42, 567–574.

[20] Li, C.-L., Tan, N.-Z., Berger, J.L., Ferguson, S.B., Zhang, Y., Prolla, T.A., Zhu, J.-S., 2011. The
Combined use of ageLOC whole-gene expression profiling technology and mouse lifespan test in anti-
aging herbal product study. Proc. 2011 New TCM Prod. Innovat. Indust. Devel. Summit, Hangzhou,
China.   pp.   443–448.
https://xueshu.baidu.com/usercenter/paper/show?paperid=08341c17fa58c8f85584b92572b90f75&site=x
ueshu_se

[21] Li, W.-J., Xia, J.-M., Li, Q.-P., Zhang, Z.-Y., Zhang, W.-W., Dong, C.-H., Wei, J.-C., Liu, X.-Z., 2020a.
Developmental recording of the ghost-moth larvae after *ex situ* infection by *Ophiocordyceps sinensis*.
Sci Chin (Life Sci), 663(7), 1093–1095. https://doi.org/10.1007/s11427-020-1686-7

[22] Li, X.-Z., Li, Y.-L., Yao, Y.-S., Xie, W.-D., Zhu, J.-S., 2020a. Further discussion with Li *et al.* (2013,
2019) regarding the "ITS pseudogene hypothesis" for *Ophiocordyceps sinensis*. Mol. Phylogenet. Evol,
146, 106728. DOI:10.1016/j.ympev.2019.106728

[23] Li, X.-Z., Li, Y.-L., Yao, Y.-S., Xie, W.-D., Zhu, J.-S., 2022a. Molecular identification of
Ophiocordyceps sinensis genotypes and the indiscriminate use of the Latin name for the multiple
genotypes and the natural insect-fungi complex. Am J Biomed Sci, 14(3), 115□135. doi:
10.5099/aj220300115

[24] Li, X.-Z., Li, Y.-L., Yao, Y.-S., Zhu, J.-S., 2022b. Differential expressions of the *Hirsutella sinensis*
genes and intraspecific genetic variations among *H. sinensis* strains. Biochem Mol Biol J, 8(2), 58. DOI:
10.36648/2471-8084.22.8.58

[25] Li, X.-Z., Li, Y.-L., Zhu, J.-S., 2023a. Differential transcription of the mating-type genes during the
sexual reproduction of natural *Cordyceps sinensis*. Chin J Chin Mater Med, 48(10), 2829-2840. DOI:
10.19540/j.cnki.cjcmm.20230213.102

[26] Li, Y., Hsiang, T., Yang, R.-H., Hu, X.-D., Wang, K., Wang, W.-J., Wang, X.-L., Jiao, L., Yao, Y.-J.,
2016a. Comparison of different sequencing and assembly strategies for a repeat-rich fungal genome,
Ophiocordyceps sinensis. J. Microbiol. Methods,. 128, 1–6.
http://dx.doi.org/10.1016/j.mimet.2016.06.025

[27] Li, Y., Jiang, L., Wang, K., Wu, H.-J., Yang, R.-H., Yan, Y.-J., Bushley, K.E., Hawksworth, D.L., Wu, Z.-
J., Yao, Y.-J., 2020c. RIP mutated ITS genes in populations of *Ophiocordyceps sinensis* and their
implications for molecular systematics. IMA Fungus, 11, 18. DOI:10.1016/j.mimet.2016.06.025

[28] Li, Y., Jiao, L., Yao, Y.-J., 2013. Non-concerted ITS evolution in fungi, as revealed from the important
medicinal fungus *Ophiocordyceps sinensis*. Mol. Phylogenetics Evol, 68, 373–379.
http://dx.doi.org/10.1016/j.ympev.2013.04.010

[29] Li, Y., Wang, Y.-H., Wang, K., Yang, R.-H., Jiao, L., Yao, Y.-J., 2019. Response to "The multiple genotypes of *Ophiocordyceps sinensis* and the ITS pseudogene hypothesis". Mol. Phylogenetics Evol, 139, 106522. https://doi.org/10.1016/j.ympev.2019.106522

[30] Li, Y.-L., Gao, L., Yao, Y.-S., Li, X.-Z., Wu, Z.-M., Tan, N.-Z., Lou, Z.-Q., Xie, W.-D., Wu, J.-Y., Zhu, J.-S., 2023b. Altered GC- and AT-biased genotypes of *Ophiocordyceps sinensis* in the stromal fertile portions and ascospores of natural *Cordyceps sinensis*. PLoS ONE, 18(6), e0286865. https://doi.org/10.1371/journal.pone.0286865

[31] Li, Y.-L., Li, X.-Z., Yao, Y.-S., Wu, Z.-M., Gao, L., Tan, N.-Z., Lou, Z.-Q., Xie, W.-D., Wu, J.-Y., Zhu, J.-S., 2023c. Differential cooccurrence of multiple genotypes of *Ophiocordyceps sinensis* in the stromata, stromal fertile portion (ascocarps) and ascospores of natural *Cordyceps sinensis*. PLoS ONE, 18(3), e0270776. https://doi.org/10.1371/journal.pone.0270776

[32] Li, Y.-L., Li, X.-Z., Zhu, J.-S., 2022c. *Cordyceps sinensis* Fungal Heterokaryons. Chemistry Industry Press Co., Beijing, China. ISBN: 978-7-122-40189-2. http://www.cip.com.cn/Book/Index/48939

[33] Li, Y.-L., Yao, Y.-S., Xie, W.-D., Zhu, J.-S., 2016b. The molecular heterogeneity of natural *Cordyceps sinensis* with multiple *Ophiocordyceps sinensis* fungi challenges the anamorph-teleomorph connection hypotheses. Am J Biomed Sci, 8, 123–159. doi: 10.5099/aj160100123

[34] Li, Y.-L., Yao, Y.-S., Zhang, Z.-H., Xu, H.-F., Liu, X., Ma, S.-L., Wu, Z.-M., Zhu, J.-S., 2016c. Synergy of fungal complexes isolated from the intestines of *Hepialus lagii* larvae in increasing infection potency. J. Fungal Res, 14, 96–112.

[35] Liu, J., Guo, L.-N., Li, Z.-W., Zhou, Z., Li, Z., Li, Q., Bo, X.-C., Wang, S.-Q., Wang, J.-L., Ma, S.-C, Zheng, J., Yang, Y., 2020. Genomic analyses reveal evolutionary and geologic context for the plateau fungus *Ophiocordyceps sinensis*. Clin Med, 15, 107⎕119. DOI: 10.1186/s13020-020-00365-3

[36] Liu, Z.-Q., Lin, S., Baker, P., Wu, L.-F., Wang, X.-R., Wu, H., Xu, F., Wang, H.-Y., Brathwaite, M., Zheng, Y.-G., 2015. Transcriptome sequencing and analysis of the entomopathogenic fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*. BMC Genome, 16, 106. DOI 10.1186/s12864-015-1269-y

[37] Liu, Z.-Y., Yao, Y.-J., Liang, Z.-Q., Liu, A.-Y., Pegler, D.N., Chase, M.W., 2001. Molecular evidence for the anamorph-teleomorph connection in *Cordyceps sinensis*. Mycol Res, 105, 827–832. https://doi.org/10.1017/S095375620100377X

[38] Mao, X.-M., Zhao, S.-M., Cao, L., Yan, X., Han, R.-X., 2013. The morphology observation of *Ophiocordyceps sinensis* from different origins. J. Environ. Entomol, 35, 343–353. doi: 10.3969/j.issn.1674-0858.2013.03.11

[39] Pang, F., Wang, L.P., Jin, Y., Guo, L.P., Song, L.P., Liu, G.M., Feng, C.Q., 2018. Transcriptome analysis of *Paecilomyces hepiali* at different growth stages and culture additives to reveal putative genes in cordycepin biosynthesis. Genomics, 110, 162–170. http://dx.doi.org/10.1016/j.ygeno.2017.09.008

[40] Ren, Y., Wan, D.-G., Lu, X.-M., Guo, J.-L., 2013. The study of scientific name discussion for TCM *Cordyceps*. Lishizhen Med Materia Medica Res, 24, 2211–2212. Doi: 10.3969/j.issn.1008-

0805.2013.09.075

[41] Selker, E.U., 1990. Premeiotic instability of repeated sequences in Neurospora crassa. Annu Rev Genet,
24, 579–613. DOI: 10.1146/annurev.ge.24.120190.003051

[42] Selker, E.U., 1997. Epigenetic phenomena in filamentous fungi: useful paradigms or repeat-induced
confusion? Trends Genet, 13, 296–301. DOI: 10.1016/S0168-9525(97)01201-8

[43] Selker, E.U., 2002. Repeat-induced gene silencing in fungi. Adv Genet, 46, 439–450. DOI:
10.1016/S0065-2660(02)46016-6

[44] Selker, E.U., Stevens, J.N., 1985. DNA methylation at asymmetric sites is associated with numerous
transition mutations. Proc Natl Acad Sci USA, 82, 8114–8118. DOI: 10.1073/pnas.82.23.8114

[45] Shu, R.-H., Zhang, J.-H., Meng, Q., Zhang, H., Zhou, G.-L., Li, M.-M., Wu, P.-P., Zhao, Y.-N., Chen, C.,
Qin, Q.-L., 2020. A new high-quality draft genome assembly of the Chinese cordyceps *Ophiocordyceps*
sinensis. Genome Biol. Evol, 12(7), 1074–1079. DOI: 10.1093/gbe/evaa112.

[46] Stensrud, Ø., Hywel-Jones, N,L., Schumacher, T., 2005. Towards a phylogenetic classification of
Cordyceps: ITS nrDNA sequence data confirm divergent lineages and paraphyly. Mycol Res, 109(1),
41–56. DOI: 10.1017/S095375620400139X

[47] Stensrud, Ø., Schumacher, T., Shalchian-Tabrizi, K., Svegården, I.B., Kauserud, H., 2007. Accelerated
nrDNA evolution and profound AT bias in the medicinal fungus *Cordyceps sinensis*. Mycol. Res, 111,
409–415. doi:10.1016/j.mycres.2007.01.015

[48] Sung, G.-H., Hywel-Jones, N,L,, Sung, J.-M., Luangsa-ard, J.J., Shrestha, B., Spatafora, J.W., 2007.
Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. Stud. Mycol, 57, 5–59.

[49] Tan, N.-Z., Berger, J.L., Zhang, Y., Prolla, T.A., Weindruch, R., Zhao, C.-S., Bartlett, M., Zhu, J.-S.,
2011. The lifespan-prolonging effect of *Cordyceps sinensis* Cs-4 in normal mice and its molecular
mechanisms.   FASEB   J,   25(1),   599.1.
http://www.fasebj.org/cgi/content/meeting_abstract/25/1_MeetingAbstracts/599.1

[50] Taranto, A.P., Hane, J.K., Williams, A.H., Solomon, P.S., Oliver, R.P., 2014. Repeat-induced point
mutation: A fungal-specific endogenous mutagenesis process. In: van den Berg M, Maruthachalam K
(eds) Genetic transformation systems in fungi, Vol. 2. Fungal Biology, Chapter 4. Springer, Cham.
https://doi.org/10.1007/978-3-319-10503-1_4

[51] Wei, J.-C., Wei, X.-L., Zheng, W., Guo, W., Liu, R., 2016. Species identification and component
detection of *Ophiocordyceps sinensis* cultivated by modern industry. Mycosystema, 35, 404–410.

[52] Wei, X.-L., Yin, X.-C., Guo, Y.-L., Shen, N.-Y., Wei, J.-C., 2006. Analyses of molecular systematics on
Cordyceps sinensis and its related taxa. Mycosystema, 25(2), 192–202.
https://europepmc.org/article/cba/618538

[53] Will I, Das B, Trinh T, Brachmann A, Ohm RA, de Bekker C. 2020. Genetic underpinnings of host
manipulation by *Ophiocordyceps* as revealed by comparative transcriptomics. G3 (Bethesda) 10, 2275–
2296. doi: 10.1534/g3.120.401290

[54] Xia, E.-H., Yang, D.-R., Jiang, J.-J., Zhang, Q.-J., Liu, Y., Liu, Y.-L., Zhang, Y., Zhang, H.-B., Shi, C.,

Tong, Y., Kim, C., Chen, H., Peng, Y.-Q., Yu, Y., Zhang, W., Eichler, E.-E., Gao, L.-Z., 2017. The caterpillar fungus, *Ophiocordyceps sinensis*, genome provides insights into highland adaptation of fungal pathogenicity. Sci Rep, 7, 1806. doi: 10.1038/s41598-017-01869-z

[55] Xia, F., Liu, Y., Shen, G.-R., Guo, L.-X., Zhou, X.-W., 2015 Investigation and analysis of microbiological communities in natural *Ophiocordyceps sinensis*. Can J Microbiol, 61, 104–111. http://dx.doi.org/10.1139/cjm-2014-0610

[56] Xiang, L., Li, Y., Zhu, Y.-J., Luo, H.-M., Li, C.-F., Xu, X.-L., Sun, C., Song, J.-Y., Shi, L.-C., He, L., Sun, W., Chen, S.-L., 2014. Transcriptome analysis of the *Ophiocordyceps sinensis* fruiting body reveals putative genes involved in fruiting body development and cordycepin biosynthesis. Genomics, 103, 154–159. http://dx.doi.org/10.1016/j.ygeno.2014.01.002

[57] Xiao, Y.-Y., Chen, C., Dong, J.-F., Li, C.-R., Fan, M.-Z., 2011 Morphological observation of ascospores of *Ophiocordyceps sinensis* and its anamorph in growth process. J Anhui Agricult Univ, 38(4), 587–591. DOI: 10.13610/j.cnki.1672-352x.2011.04.022

[58] Xiao, W., Yang, J.-L., Zhu, P., Cheng, K., He, H., Zhu, H., Wang, Q., 2009. Non-support of species complex hypothesis of *Cordyceps sinensis* by targeted rDNA-ITS sequence analysis. Mycosystema, 28, 724–730.

[59] Yao, Y.-S., Gao, L., Li, Y.-L., Ma, S.-L., Wu, Z.-M., Tan, N.-Z., Wu, J.-Y., Ni, L.-Q., Zhu, J.-S., 2014. Amplicon density-weighted algorithms for analyzing dissimilarity and dynamic alterations of RAPD polymorphisms of *Cordyceps sinensis*. Beijing Da Xue Xue Bao Yi Xue Ban, 46, 618–628. DOI: 10.3969/j.issn.1671-167X.2014.04.027

[60] Yao, Y.-S., Zhou, Y.-J., Gao, L., Lu, J.-H., Wu, Z.-M., Zhu, J.-S., 2011. Dynamic alterations of the differential fungal expressions of *Ophiocordyceps sinensis* and its mutant genotypes in stroma and caterpillar during maturation of natural *Cordyceps sinensis*. J Fungal Res, 9, 37–49,53.

[61] Yao, Y.-S., Zhu, J.-S., 2016. Indiscriminate use of the Latin name for natural *Cordyceps sinensis* and *Ophiocordyceps sinensis* fungi. Chin. J. Chin. Mater. Med, 41(7), 1316–1366.

[62] Zhang, S., Zhang, Y.-J., 2015.. Molecular evolution of three protein-coding genes in the Chinese caterpillar fungus *Ophiocordyceps sinensis*. Microbiology China, 42(8), 1549–1560. DOI: 10.13344/j.microbiol.china.150279

[63] Zhang, S., Zhang, Y.-J., Liu, X.-Z., Zhang, H., Liu, D.-S., 2013a. On the reliability of DNA sequences of *Ophiocordyceps sinensis* in public databases. J Ind Microbiol Biotechnol, 40, 365–378. DOI 10.1007/s10295-012-1228-4

[64] Zhang, S., Zhang, Y.-J., Shrestha, B., Xu, J.-P., Wang, C.-S., Liu, X.-Z., 2013b. *Ophiocordyceps sinensis* and *Cordyceps militaris*: research advances, issues and perspectives. Mycosystema, 32, 577–597. DOI: 10.13346/j.mycosystema.2013.04.001

[65] Zhang, S.-W., Cem, K., Liu, Y., Zhou, X.-W., Wang, C.-S., 2018. Metatranscriptomics analysis of the fruiting caterpillar fungus collected from the Qinghai-Tibetan plateau. Scientia Sinica Vitae, 48(5), 562. doi: 10.1360/N052017-00253

[66] Zhang, Y.-J., Li, E.-W., Wang, C.-S., Li, Y.-L., Liu, X.-Z., 2012. *Ophiocordyceps sinensis*, the flagship
fungus of China: terminology, life strategy and ecology. Mycology, 3, 2–10. DOI:
10.1080/21501203.2011.654354

[67] Zhang, Y.-J., Sun, B.-D., Zhang, S., Wang, M., Liu, X.-Z., Gong, W.-F., 2010. Mycobiotal investigation
of natural *Ophiocordyceps sinensis* based on culture-dependent investigation. Mycosystema, 29,
518–527.

[68] Zhang, Y.-J., Xu, L., Zhang, S., Liu, X.-Z., An, Z., Wang, M., Guo, Y., 2009. Genetic diversity of
Ophiocordyceps sinensis, a medicinal fungus endemic to the Tibetan Plateau: implications for its
evolution and conservation. BMC Evolut. Biol, 9, 290. http://www.biomedcentral.com/1471-2148/9/290

[69] Zhang, Y.-J., Zhang, S., Li, Y.-L., Ma, S.-L., Wang, C.-S., Xiang, M.-C., Liu, X.-Z., An, Z.-Q., Xu, J.-P.,
2014. Phylogeography and evolution of a fungal–insect association on the Tibetan Plateau. Mol Ecol, 23,
5337–5355. doi: 10.1111/mec.12940

[70] Zhao, J., Wang, N., Chen, Y.-Q., Li, T.-H., Qu, L.-H., 1999. Molecular identification for the asexual
stage of *Cordyceps sinensis*. Acta Scientiarum Naturalium Universitatis Sunyatseni, 38, 122–123.

[71] Zhu, J.-S., Gao, L., Li, X.-H., Yao, Y.-S., Zhao, J.-Q., 2010. Maturational alteration of oppositely
orientated rDNA and differential proliferation of GC- and AT-biased genotypes of *Ophiocordyceps*
sinensis and *Paecilomyces hepiali* in natural *Cordyceps sinensis*. Am J Biomed Sci, 2, 217–238. doi:
10.5099/aj100300217

[72] Zhu, J.-S., Guo, Y.-L., Yao, Y.-S., Zhou, Y.-J., Lu, J.-H., Qi, Y., Liu, X.-Z., Wu, Z.-M., Chen, W.,
Zhang, L., Yin. W.-T., Zheng, T.-Y., Zhang, L.-J., 2007. Maturation of *Cordyceps sinensis* associates
with co-existence of *Hirsutella sinensis* and *Paecilomyces hepiali* DNA and dynamic changes in fungal
competitive proliferation predominance and chemical profiles. J Fungal Res, 5(4), 214–224. DOI:
10.3969/j.issn.1672-3538.2007.04.009

[73] Zhu, J.-S., Halpern, G.M., Jones, K., 1998a. The scientific rediscovery of a precious ancient Chinese
herbal regimen: *Cordyceps sinensis*: Part I. J Altern Complem Med, 4(3), 289–303. DOI:
10.1089/acm.1998.4.3-289

[74] Zhu, J.-S., Halpern, G.M., Jones, K., 1998b. The scientific rediscovery of an ancient Chinese herbal
medicine: *Cordyceps sinensis*: Part II. J Altern Complem Med, 4(4), 429–457. DOI:
10.1089/acm.1998.4.429

[75] Zhu, J.-S., Li, Y.-L., 2017. A Precious Transitional Chinese Medicine, *Cordyceps sinensis*: Multiple
Heterogeneous *Ophiocordyceps sinensis* in the Insect-Fungi Complex. Lambert Academic Publishing,
Germany.

[76] Zhu, J.-S., Li, Y.-L., Yao, Y.-S., Xie, W.-D., 2018. Molecular evidence not supporting the use of
Hirsutella sinensis strain EFCC 7287 as the taxonomic standard for multiple *Ophiocordyceps sinensis*
fungi. Int J Mol Biol, 3, 114–115. DOI: 10.15406/ijmboa.2018.03.00062

[77] Zhu, J.-S., Li, Y.-L., Yao, Y.-S., Xie, W.-D., 2019. The multiple genotypes of *Ophiocordyceps sinensis*
and the ITS pseudogene hypothesis. Mol Phylogenetics Evol, 139, 106322.

https://doi.org/10.1016/j.ympev.2018.10.034

[78] Zhu, J.-S., Wu, J.-Y., 2015. Genetic heterogeneity of natural *Cordyceps sinensis* with co-existence of
multiple fungi. Chin J Cell Biol, 37(2), 284–298. DOI: 10.11844/cjcb.2015.02.0284

[79] Zhu, J.-S., Zhao, J.-G., Gao, L., Li, X.-H., Yao, Y.-S., Zhao, J.-Q., 2012. Dynamically altered
expressions of at least 6 *Ophiocordyceps sinensis* mutants in the stroma of *Cordyceps sinensis*. J Fungal
Res, 10, 100–112.

1841