Multiple GC-biased repetitive ITS copies in the *Hirsutella sinensis* genome are not generated by RIP mutagenesis involving transition point mutations

Xiu-Zhang Li[a], Yu-Ling Li[a], Ya-Nan Wang[b], Jia-Shi Zhu[a,c,*]

[a] State Key Laboratory of Plateau Ecology and Agriculture, Qinghai Academy of Animal and Veterinary Sciences, Qinghai University, Xining, Qinghai 810016, China

[b] State Key Laboratory of Breeding Base of Dao-di Herbs, National Resource Center for Chinese Materia Medica, China Academy of Chinese Medical Sciences, Beijing 100700, China

[c] Institute of Biopharmaceutical and Health Engineering, Shenzhen International Graduate School, Tsinghua University, Shenzhen 518055, China

[*] Corresponding author:

E-mail address: zhujosh@163.com (J-S Zhu)

Short title:

H. sinensis genomic ITS repeats are not generated by RIP mutagenesis

Abstract

Li et al. (2013, 2020b) hypothesized that AT-biased genotypes of *Ophiocordyceps sinensis* are generated through "repeat-induced point mutation (RIP)" and exist as "ITS pseudogenes in a single genome" of GC-biased Genotype #1 *Hirsutella sinensis*. This study revealed multiple GC-biased repetitive ITS copies in the *H. sinensis* genome, including multiple insertion/deletion and transversion alleles, which might not be generated through RIP mutagenesis that theoretically causes C-to-T and G-to-A transitions. The repetitive ITS copies are genetically and phylogenetically distinct from the AT-biased genotypes of *O. sinensis*, which possess multiple scattered transition alleles. The sequences of GC- and AT-biased Genotypes #2–17 were absent in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000 and NGJJ00000000 of *H. sinensis* strains but instead in the genomes of independent *O. sinensis* fungi. The GC- and AT-biased genotypes of *O. sinensis* were found to differentially occur in different combinations in different compartments of natural *Cordyceps sinensis*. Their abundances were dynamically altered in an asynchronous, disproportional manner during *C. sinensis* maturation. Metatranscriptomic analysis of natural *C. sinensis* revealed transcriptional silencing of the 5.8S genes of all *C. sinensis*-colonizing fungi, including *H. sinensis*. The problematic design of the 5.8S gene transcription assay and data analysis reported by Li et al. (2013) have provided controversial and incomplete evidence to support the notion that the 5.8S genes of AT-biased genotypes of *O. sinensis* are permanently nonfunctional "ITS pseudogenes". These results demonstrate the genomic independence of 17 genotypes belonging to independent *O. sinensis* fungi. In parallel with GC-biased genotypes, AT-biased *O. sinensis* genotypes might have been generated during the long history of evolution from a common ancestor through RIP or other mutagenesis mechanisms in response to the extremely harsh environment of the Qinghai-Tibet Plateau.

KEY WORDS: Multiple genotypes of *Ophiocordyceps sinensis*; genomic independence; repeat-induced point mutations (RIP); insertion/deletion, transversion and transition point mutations; 5.8S gene transcript; transcriptional silencing.

I. Introduction

Natural *Cordyceps sinensis* is one of most valued therapeutic agents in traditional Chinese medicine and has a rich history of clinical use for health maintenance, disease amelioration, postdisease/surgery recovery and antiaging therapy (Zhu et al. 1998a, 1998b; Li et al. 2001; Tan et al. 2001). The Chinese Pharmacopeia defines natural *C. sinensis* as an insect-fungi complex, which includes the remains of Hepialidae moth larva and >90 fungal species from >37 genera, with differential co-occurrence of some of 17 genotypes of *Ophiocordyceps sinensis*, *i.e.*, natural *C. sinensis* ≠ *O. sinensis* fungus (Engh 1999; Kinjo and Zang 2001; Chen et al. 2004, 2011; Stensrud et al. 2005, 2007; Wei et al. 2006, 2016; Yang et al. 2008; Xiao et al. 2009; Zhang et al. 2009, 2010; Zhu et al. 2010, 2012, 2018, 2019; Gao et al. 2011, 2012; Yao et al. 2011; Mao et al. 2013; Xia et al. 2015; Zhu and Wu 2015; Li et al. 2016b, 2020a, 2022a, 2022c, 2022d, 2022e; Zhu and Li 2017). However, the Latin name *C. sinensis* has been indiscriminately used for both the wild insect-fungi complex and the teleomorph/holomorph of the *C. sinensis* fungus since the 1840s (Yao & Zhu 2016; Li et al. 2022a). The fungus was renamed *Ophiocordyceps sinensis* using *Hirsutella sinensis* Strain EFCC 7287 (GC-biased Genotype #1) as the nomenclature reference (Sung et al. 2007), while the name of the wild and cultivated insect-fungi complex has remained unchanged. Since then, especially after the improper implementation of the IMA's One Fungus=One name nomenclature rule in *O. sinensis* research (Hawksworth et al. 2007; Zhang et al. 2013; Li et al. 2022a), the name *O. sinensis* has been used to refer to multiple teleomorphic and anamorphic fungi that have distinct genomes and has been indiscriminately applied to the natural insect-fungi complex (Zhu and Wu 2015; Yao & Zhu 2016; Li et al. 2016b, 2022a, 2022e; Zhu and Li 2017). The confusion surrounding Latin name use continues and has even spread from the scientific community to the public media and mass markets, causing a significant decrease in the wholesale and retail prices of natural *C. sinensis* and related fermentation products. Because a consensus Latin name for the wild insect-fungi product has not been identified by taxonomists and multiple genotypes of *O. sinensis* with undetermined taxonomic positions are referred to by the same Latin name (Zhang et al. 2012; Ren et al. 2013; Yao & Zhu 2016; Li et al. 2022a, 2022d, 2022e), we temporarily refer to the multiple genotype fungi as *O. sinensis*, including Genotype #1 *H. sinensis*, and we continue to refer to the wild and cultivated insect-fungi complexes as *C. sinensis*; however, this practice only partially ameliorates the academic confusion arising from the indiscriminate use of Latin names and will likely be replaced by the exclusive use of distinct Latin names in the future.

All 17 genotypes of *O. sinensis* are indiscriminately referred to by the same Latin name (Zhu et al. 2010, 2019; Yao & Zhu 2016; Li et al. 2020a, 2022a, 2022c, 2022d, 2022e). Kinjo and Zang (2001) were the first to report 2 groups and 3 clades of mutant *O. sinensis* sequences, which were mistakenly believed to represent intraspecific variants based on the observation of the same *H. sinensis*-like morphological and growth characteristics. Mao et al. (2013) confirmed that the fungi in Groups B–C (Genotypes #4–5) shared the same *H. sinensis*-like morphological and growth characteristics. Stensrud et al. (2007) used a Bayesian phylogenetic algorithm and confirmed the existence of 3 clades of mutant sequences: GC-biased Group A and AT-biased Groups B–C of *O. sinensis*, in addition to 2 other clades (Groups D–E) believed to be unrelated to *O. sinensis*. All the mutants, including the sequences in Groups D–E, were annotated as *C. sinensis* or *O. sinensis* under GenBank taxid 72228. Stensrud et al. (2007) reported that the variation in the 5.8S gene sequences of *O. sinensis* in Groups A–C "far exceeds what is normally observed in fungi … even at higher taxonomic levels (genera and family)" that the mutant genotypes with "profound AT bias … share a common ancestor" and originate from "accelerated nrDNA evolution" (nrDNA: nuclear ribosomal DNA). In addition to the use of "universal" primers, Xiao et al. (2009) designed pairs of genotype-specific primers and used a nested PCR technique to identify the differential coexistence of multiple *O. sinensis* mutants in the stroma and caterpillar body of natural *C. sinensis* in different maturational stages; these authors believed that the genetically variable sequences likely belonged to independent *O. sinensis* fungi.

Figure 3 of (Bushley et al. 2013) demonstrates that *C. sinensis* hyphae and ascospores are heterokaryons containing multiple mono-, bi-, and trinucleated cells. Li et al. (2013) reported the identification of heterogeneous ITS sequences of both GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis* in 8 out of 15 clones following 25 days of liquid incubation of *C. sinensis* mono-ascospores. They reported the identification of the 5.8S gene cDNA of Genotype #1 but not Genotype #5 in a cDNA library of Strain 1220, one of the 8 heterogeneous clones. By ignoring the simultaneous identification of homogenous Genotype #1 in 7 other clones derived from the same mono-ascospores, these authors concluded that "Different types of ITS sequences were thus proved to coexist in a single genome" and proposed the hypothesis of "ITS pseudogene … in a single genome", which was extended to all AT-biased ITS genotypes. Considering the uncertain definition of "ITS pseudogene … in a single genome" based on controversial evidence, Li et al. (2017) extended this hypothesis to "rDNA pseudogenes", and Li et al. (2020b) further hypothesized that all AT-biased *O. sinensis* genotypes "could have emerged either before or after" a new GC-biased "ITS haplotype

was generated" through "repeat-induced point mutations (RIP)" and co-occur as repetitive ITS copies

in "a single genome" of GC-biased Genotype #1 *H. sinensis*.

This paper reanalyzes the relevant genomic and transcriptomic sequences in the GenBank

database. We discuss the genetic and phylogenetic characteristics of multiple repetitive ITS copies in

the genome of Genotype #1 of *O. sinensis*, the genomic residency of AT-biased genotypes in the

genome of Genotype #1 *H. sinensis*, and whether sufficient evidence exists to reach a conclusion that

the 5.8S genes of AT-biased genotypes are permanently nonfunctional pseudogenes and those of GC-

biased genotypes are functional counterparts.

II. Materials and Methods

II-1. Reagents

Common laboratory reagents such as agarose, electrophoresis reagents, nylon membranes, and

blocking solution, etc., were purchased from Beijing Bioland Technology Company. Endonuclease

*Ava*I, *Dra*I, and *Eco*RI are products of New England BioLabs, United States. Mercuric chloride (0.1%)

for surface sterilization of freshly collected *C. sinensis* specimens was a gift from the Institute of

Microbiology, Chinese Academy of Sciences. The Universal DNA Purification kit was a product of

TIANGEN BIOTECH Company, China. The DNeasy Plant Mini Kit was a product of Qiagen

Company, Germany. The Gel Extraction Kit was a product of Omega Bio-Tek, United States. The

Taq PCR reagent kit and Vector NTI Advance 9 software were purchased from Invitrogen, United

States. All primers were synthesized by Invitrogen Beijing Lab. or Beijing Bomaide Technology Co.

The DIG High Prime DNA Labeling and Detection Kit was purchased from Roche Diagnostics,

Indiana, USA.

II-2. Information on fungal species in natural *C. sinensis*

134   Information regarding intrinsic fungal species in natural *C. sinensis* was summarized by Jiang

135 and Yao (2013), and subsequent discoveries of additional species among natural *C. sinensis*

136 specimens were reviewed by Li et al. (2016b, 2022a, 2022e). The sequences of 17 genotypes of *O.*

137 *sinensis* have been obtained and uploaded to GenBank by many molecular mycology groups since

138 1999. A BLAST search of the GenBank database revealed 668 ITS sequences of *O. sinensis*, which

were uploaded. These genotypic sequences were analyzed genetically and phylogenetically by Stensrud et al. (2007), Zhu et al. (2010, 2012), Zhang et al. (2013), Zhu & Wu (2015), Li et al. (2016b, 2022a, 2022d. 2022e), and Zhu & Li (2017).

II-3. Genetic, genomic and transcriptomic sequences from GenBank

Five genome assemblies, ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000 and NGJJ00000000, for *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195 and CC1406-203, respectively, were uploaded to GenBank (Hu et al. 2013; Shu et al. 2020; Li et al. 2016a; Jin et al. 2020; Liu et al. 2020) and used for the analysis of repetitive ITS copies and multiple GC- and AT-biased genotypes.

mRNA transcriptome assembly GCQL00000000, for *H. sinensis* Strain L0106, was uploaded to GenBank by Liu et al. (2015). Two metatranscriptome studies of natural *C. sinensis* have been reported. Xiang et al. (2014) uploaded the GAGW00000000 metatranscriptome assembly to GenBank, which was obtained from natural specimens (unknown maturational stages) collected from Kangding County of Sichuan Province, China. Xia et al. (2017) uploaded the unassembled SRR5428527 Shotgun Read Archive (SRA) to GenBank for mature *C. sinensis* specimens collected from Deqin County of Yunnan Province, China, but the assembled metatranscriptome sequences were uploaded to www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas, which is currently inaccessible, but the file has been downloaded for analysis.

II-4. Genetic and phylogenetic analyses

The ITS sequences of multiple genotypes of *O. sinensis* and multiple genomic repetitive ITS copies as well as metatranscriptomic/transcriptomic sequences were analyzed using the GenBank BLAST algorithms (https://blast.ncbi.nlm.nih.gov/blast/). A Bayesian phylogenetic tree of the ITS sequences was inferred using MrBayes v3.2.7 software (Markov chain Monte Carlo [MCMC] algorithm) based on 3×10^4 samples, with a sampling frequency of 10^3 iterations after discarding the first 25% of samples from a total of 4×10^6 iterations (Huelsenbeck and Ronquist 2001; Li et al. 2022a, 2022d); this phylogenetic analysis was performed at Nanjing Genepioneer Biotechnologies Co.

II-5. Collection and on-site processing of fresh *C. sinensis*

Fresh *C. sinensis* specimens were purchased in a local market in Kangding County of Sichuan Province, China (3800–4600 m' altitude; latitude 30°049'N, longitude 101°959'E) (Zhu et al. 2010; Yao et al. 2014). Immature *C. sinensis* specimens were characterized by a plump caterpillar body and very short stroma (1.0–2.0 cm). Maturing *C. sinensis* specimens were characterized by a plump caterpillar body and stroma of 2.5–5.0 cm in length, without the formation of an expanded fertile portion close to the stromal tip. Mature *C. sinensis* specimens were characterized by a plump caterpillar body and long stroma (>5.0 cm) and by the formation of an expanded fertile portion close to the stromal tip, which was densely covered with numerous ascocarps (Zhu et al. 2010; Li et al. 2022c). Governmental permission was not required for *C. sinensis* purchases in local markets, and the collections of *C. sinensis* specimens from sales by local farmers fall under the governmental regulations for traditional Chinese herbal products.

Freshly collected *C. sinensis* specimens were washed thoroughly in running water with gentle brushing and surface sterilized in 0.1% mercuric chloride for 10 min, followed by washing 3 times with sterile water (Zhu et al. 2010; Li et al. 2022c). The thoroughly cleaned and sterilized *C. sinensis* specimens were immediately frozen in liquid nitrogen and kept frozen during transportation and storage prior to further processing.

II-6. Extraction of genomic DNA and preparation of *C. sinensis* DNA

Caterpillar bodies and stromata of sterilized, frozen *C. sinensis* were separately ground into powder in liquid nitrogen. Genomic DNA was individually extracted from the powder samples using a DNeasy Plant Mini Kit.

II-7. PCR amplification and amplicon sequencing

Genomic DNA of immature, maturing and mature stromata of natural *C. sinensis* was used as the template for PCR amplification. *H. sinensis*-specific primers were designed (a forward primer *Hsprp1*: 5' ATTATCGAGTCACCACTCCCAAACCCCC 3'; and a reverse primer *Hsprp3*: 5'

AGGTTCTCAGCGAGCTA 3') based on alignment analysis of the ITS sequences (AB067721,

AB067740, and AB067744 representing Genotypes #1 and #4–5 of *O. sinensis*, respectively).

A touch-down PCR protocol was used, in which the annealing temperature was initially set at

70 °C and was then decreased stepwise with a decrement of 0.3 °C every cycle, for a total of 36 cycles

(Zhu et al. 2007, 2010). PCR amplicons were subjected to agarose gel electrophoresis, and DNA

bands were recovered, purified using a gel extraction kit, and sequenced.

II-8. Restriction digestion assay and genotypic density analysis

The target PCR moieties (440 bp) based on the design of the primers (*Hsprp1* and *Hsprp3*) from

sequences of GC- and AT-genotypes of *O. sinensis* were subjected to overnight digestion with

endonuclease *Eco*RI, which is specific to the GC-biased Genotype #1 sequence (*Eco*RI-sensitive; *cf*.

Figure 1 below). AT-biased genotypes are *Eco*RI-resistant due to a single-base mutation that results

in the loss of the *Eco*RI enzymatic site. The *Eco*RI digestion products were subjected to agarose gel

electrophoresis, and the digestion fragments were recovered from the gel and sequenced. DNA

sequences were analyzed with Vector NTI Advance 9.

The density of PCR amplicons and the *Eco*RI digestion fragments in agarose gels were analyzed

and quantified using a gel imaging system with Quantity One software (Universal Hood II Molecular

Imager Gel Doc XR System, Bio-Rad, CA, USA). To monitor the alterations in *O. sinensis* genotypes

during *C. sinensis* maturation, the density of each DNA band in different agarose gels was normalized

to the density of the target 440-bp moiety that served as an internal control.

II-9. Preparation of fungus-specific ITS1 probe and 18S internal control probe

An *H. sinensis*-specific probe (158 bp) was designed based on the ITS1 sequence of GC-biased

Genotype #1 *H. sinensis* and generated through PCR using a forward *Hsprp1* primer and a reverse

Hsprp2 primer (5' ATTTGCTTGCTTCTTGACTGAGAGATGCC 3').

A nonspecific 18S internal control probe was designed based on alignments of GenBank 18S

rDNA sequences (DQ838795, AB187268, AB067735, AB067736 and AB080090) of several *C.*

sinensis-associated fungi. A highly homologous region of 18S rDNA, located over 700 bp upstream

of the ITS1 regions of *C. sinensis*-associated fungi, was selected. The 18S control probe was generated through PCR using a forward *Icprp1* primer (5' ACCGTCGTAGTCTTAACCAT 3') and a reverse *Icprp2* primer (5' ATCGGCTTGAGCCGATAGTC 3').

Both amplicons were confirmed by sequencing, and the probes were labeled using the DIG High Prime DNA Labeling and Detection Kit.

II-10. Southern blotting analysis

The stromata and caterpillar bodies of immature and mature *C. sinensis* were used for Southern blotting. Genomic DNA was extracted from the samples and prepared by endonuclease digestion using *Ava*I, *Dra*I, and *Eco*RI. The digested genomic DNA (5 µg of each sample) was subjected to agarose gel electrophoresis and transferred onto a nylon membrane through capillary transfer, followed by baking the membrane at 58 °C for 4 hours for fixation, blocking the binding sites on the membrane in blocking solution, and hybridizing with the labeled *H. sinensis*-specific probe at 40 °C overnight. The hybridized transblotted membrane was viewed by colorimetric photography using a gel imaging system (Bio-Rad) and then stripped and reprobed with the labeled 18S internal control probe.

III. Results and Discussion

III-1. GC-biased characters of multiple repetitive ITS copies in the *H. sinensis* genome

Generally, fungal genomes contain multiple repetitive copies of nrDNA ITS1-5.8S-ITS2 sequences. However, a single copy of the ITS1-5.8S-ITS2 sequences was identified in genome assemblies ANOV01021709, LKHE01000582, and LWBQ01000008 of *H. sinensis* Strains Co18, 1229, and ZJB12195, respectively (Hu et al. 2013; Li et al. 2016a; Jin et al. 2020), probably because the ITS repeats were discarded during the assembly of the genome shotgun sequences. Multiple repetitive ITS copies were identified in 2 additional genome assemblies, JAAVMX000000000 and NGJJ00000000, for *H. sinensis* Strains IOZ07 and CC1406-203, respectively (Liu et al. 2020; Shu et al. 2020) (Table 1), where 17 ITS1-5.8S-ITS2 copies were found within JAAVMX000000000, and 7

251   full-length ITS1-5.8S-ITS2 copies and 2 partial ITS segments were found within NGJJ00000000.

Table 1. GC contents of the repetitive ITS1-5.8S-ITS2 sequences from the genome assemblies of 2 H.
sinensis strains and percent similarities of the repetitive ITS sequences to the sequences of GC-biased
Genotype #1 (AB067721) and AT-biased Genotypes #4–6 and #15–17 of O. sinensis.

Genome segment	Sequence range & direction	GC content	Percent similarity of the repetitive sequence	
			vs. Genotype #1 AB067721	*vs.* AT-biased Genotypes #4–6 and #15–17
JAAVMX010000002	18688917→18689407	64.8%	100%	86.7% – 89.9%
	18702095→18702586	64.5%	97.4%	82.8% – 88.0%
JAAVMX010000008	13823→14313	64.8%	100%	85.5% – 89.9%
	1199→1687	64.8%	99.2%	85.2% – 89.3%
JAAVMX010000017	9147←9637	64.8%	100%	85.5% – 89.9%
	21791←22281	64.8%	100%	85.5% – 89.9%
	34435←34925	64.8%	100%	85.5% – 89.9%
	47079←47569	64.8%	100%	85.5% – 89.9%
JAAVMX010000018	13381→13871	64.8%	100%	85.5% – 89.9%
	26076→26566	64.8%	100%	85.5% – 89.9%
	38771→39261	64.8%	100%	85.5% – 89.9%
	51467→51958	64.6%	98.8%	85.5% – 88.8%
	700→1186	64.5%	97.0%	82.6% – 87.6%
JAAVMX010000019	19404→19894	64.8%	100%	85.5% – 89.9%
	32048→32538	64.8%	100%	85.5% – 89.9%
	6233→6733	65.3%	94.5%	81.3% – 85.5%
	44729→45251	63.3%	90.8%	80.1% – 84.6%
NGJJ01000573	13133←13623	64.8%	99.4%	85.2% – 89.6%
	14624←15113	64.7%	99.4%	84.8% – 89.6%
	1←260 (ITS1 and partial 5.8S)	56.5%	98.5%	83.5% – 88.3%
NGJJ01000582	10703←11192	64.7%	99.8%	85.5% – 89.8%
	24068←24190 (only partial ITS2)	73.2%	98.1%	87.7% – 89.1%
NGJJ01000796	1514→2004	64.8%	100%	85.5% – 89.9%
NGJJ01000798	1177→1665	64.6%	99.6%	84.8% – 89.6%
NGJJ01000799	3295→3785	64.8%	100%	85.5% – 89.9%
	16666→17156	64.8%	100%	85.5% – 89.9%

257   Most of the repetitive ITS copies were highly homologous (>97%) to reference sequence

258   AB067721 of GC-biased Genotype #1 *H. sinensis*. Two copies (6233→6733 and 44729→45251)

259   within JAAVMX010000019 showed similarities of 94.5% and 90.8%, respectively, to AB067721

260   (Table 1). Two other copies, 18702095→18702586 within JAAVMX010000002 and 700→1186

within JAAVMX010000018, showed similarities of 97.4% and 97.0%, respectively (Table 1). All of the repetitive ITS copies listed in Table 1 showed 80.1%–89.9% similarity to the sequences of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* (Tables 1 and 2).

Table 2. *Percent similarities between nrDNA ITS1-5.8S-ITS2 sequences of 5 genome assemblies of Genotype #1 H. sinensis and 17 O. sinensis genotype sequences. (Modified from Table 1 of Li et al. 2022d).*

| GT | GenBank Accession # | % similarity between the *H. sinensis* genomic ITS segments and 17 *O. sinensis* genotypes | | | | |
		ANOV01021709 838→1504	LKHE01000582 2074→2740	LWBQ01000008 991739→992405	JAAVMX010000017 9029←9695	NGJJ01000799 3237→3903
#1	AB067721	99.7%	100%	99.6%	100%	100%
#2	MG770309	94.4%	94.7%	94.7%	94.7%	94.7%
#3	HM595984	95.9%	96.1%	95.6%	95.3%	95.3%
#7	AJ488254	94.7%	95.0%	95.3%	95.0%	95.0%
#8	GU246286	89.5%	89.5%	89.2%	89.5%	89.5%
#9	GU246288	95.0%	95.2%	94.6%	95.2%	95.2%
#10	GU246287	83.0%	83.2%	83.0%	83.2%	83.2%
#11	JQ695935	78.9%	79.1%	78.5%	79.1%	79.1%
#12	GU246296	94.3%	94.5%	93.9%	94.5%	94.5%
#13	KT339190	87.5%	87.5%	87.2%	87.5%	87.5%
#14	KT339178	89.3%	89.6%	89.1%	89.6%	89.6%
#4	AB067744	89.1%	89.4%	89.3%	89.4%	89.4%
#5	AB067740	86.3%	86.7%	86.2%	86.7%	86.7%
#6	KJ720572	85.5%	85.5%	85.5%	85.5%	85.5%
#15	KT232017	89.6%	89.9%	89.4%	89.9%	89.9%
#16	KT232019	87.4%	87.7%	87.6%	87.7%	87.7%
#17	KT232010	87.7%	88.1%	87.5%	88.1%	88.1%

Note: "GT" represents genotype. GC-biased genotypes are listed in blue in the upper panel. AT-biased genotypes are listed in red in the lower panel.

These variable and slightly less variable GC-biased repetitive ITS copies contain predominantly insertion/deletion mutant alleles, with much less predominant transversion mutant alleles and no or a few transition mutant alleles (Table 3). The variable (94.5% and 90.8% similarity) ITS repeats, 6233→6733 and 44729→45251 within JAAVMX010000019, contained 22 and 32 insertion/deletion alleles and 4 and 11 transversion alleles (Table 3; in blue in Figure 1); they contained only 1 and 5 transition mutant alleles, having allelic ratios of insertion/deletion/transversion *vs*. transition of 26:1 and 43:5, respectively. The slightly less variable (97.0–97.4% similarity) ITS repeats 18702095→18702586 within JAAVMX010000002 and 700→1186 within JAAVMX010000018

contained no transition alleles and predominantly included 13 and 15 insertion, deletion and transversion alleles, with allelic ratios of insertion/deletion/transversion *vs.* transition of 13:0 and 15:0 (Table 3; in green in Figure 1). Other repetitive ITS copies with 98.1%–99.8% similarity to AB067721 listed in Table 1 contained only a few insertion/deletion alleles.

Table 3. *Types of mutations in the variable and slightly less variable genomic repetitive ITS sequences compared with GC-biased Genotype #1 (AB067721).*

	Sequence range & direction	% similarity	Number of mutant alleles			Ins./Del./Transv. *vs.* Transit.
			Ins./Del.	Transv.	Transit.	
JAAVMX010000002	18702095→18702586	97.4%	13	0	0	13 : 0
JAAVMX010000018	700→1186	97.0%	10	5	0	15 : 0
JAAVMX010000019	6233→6733	94.5%	22	4	1	26 : 1
	44729→45251	90.8%	32	11	5	43 : 5

Note: "Ins./Del." indicates insertion/deletion point mutations. "Transv." denotes Transversion point mutations. "Transit." refers to transition point mutations.

All the repetitive ITS copies within genome assemblies JAAVMX000000000 and NGJJ00000000 listed in Table 1, for *H. sinensis* Strains IOZ07 and CC1406-203, and the single genomic ITS copies ANOV01021709, LKHE01000582, and LWBQ01000008 listed in Table 2, for Strains Co18, 1229, and ZJB12195, respectively (Hu et al. 2013; Jin et al. 2020; Li et al. 2016a; Liu et al. 2020; Shu et al. 2020), are GC-biased with GC contents of 64.7±2.3% (Table 1), similar to the GC content (64.8%) of the AB067721 Genotype #1 sequence. Regardless of the sequence similarity to the reference sequence AB067721 of Genotype #1, the repetitive ITS sequences contained predominantly insertion and deletion alleles and less frequent transversion alleles, while no or only a few transition alleles were found. These genetic features are distinct from AT-biased Genotypes #4–6 and #15–17 of *O. sinensis*, which included multiple scattered transition alleles (Figure 1) and contained 46.9%–55.2% AT bases and 44.8%–53.1% GC bases.

```
GT#1  AB067721         59     TCGAGTCACCACTCCC AAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCCGGCG CCCCAGGGCCCG 136
GT#3  FJ654148          1     CT----TG-------- ---A----C-T------------------------------------ ----------- 83
JAAVMX010000019    19404     ---------------- -------------------------------------------------- ------------ 19481
JAAVMX010000019    32048     ---------------- -------------------------------------------------- ------------ 32125
JAAVMX010000002 18702095     ---------------- -----------------------------------------T-------- ------------ 18702172
JAAVMX010000018      700     --------- ------ -----------------------T---------------------T 776
JAAVMX010000019     6233     ----------------C- -------------------------------------------- ------------ 6311
JAAVMX010000019    44729     ---------------- -------------------------------------------------- ------------ 44806
GT#4  AB067744         55     ------T-------T ----------T--T----------------------- ---T----- 132
GT#5  AB067740         59     ------T--T----TT ------ATA--T--T-T--------TA-----------TA--- ---T-A------ 136
GT#6  KJ720572          11     ------A---T-T-------------TA-----------TA--- --TT-A------ 70
GT#15 KT232017         37     ------T--------T --------T--T----------------------- ---T------ 114
GT#16 KT232019         37     ------T--T----T ------T-T-T--------TA-----------TA--- ---T-A------ 114
GT#17 KT232010         37     ------T--T----T ------T-T-T--------TA-----------TA--- ---T-A------ 114

GT#1  AB067721        137     GACCAGG GCGCCC GCCGGAGGACCCCCAGAC CCTCCTGTCGCA GTGGCATCT CTCAGTCAAGAA GCAAGCAAAT 210
GT#3  FJ654148         84     ------- ------ ------------------ T---------- -----G--- ------------ ---------- 157
JAAVMX010000019    19482     ------- ------ ------------------ ------------ --------- ------------ ---------- 19555
JAAVMX010000019    32126     ------- ------ ------------------ ------------ --------- ------------ ---------- 32199
JAAVMX010000002 18702173     ------- ------A---------------- ------------A--------- ------------ ---------- 18702247
JAAVMX010000018      777     ------- ------ ------------------ ------------ --------- ---C-------- ---------- 850
JAAVMX010000019     6312     -------- ------ ------T----------C ------------ ---------C----------G---------- 6388
JAAVMX010000019    44807     -------C- ------ ------------------ ------------ --------- ------------ ---------- 44881
GT#4  AB067744        133     ---T--- ------ ----------TT---- ------A---T- --A------ --T---T----- 206
GT#5  AB067740        137     ---T-AA ------ ----------TT---- ------A-TAT- --A-T---- --T---T----- 213
GT#6  KJ720572         71     ---T--- ------ ----------TT---- ------A---T- --A-T---- --T---T----- 182
GT#15 KT232017        115     ---T--- ------ ----------TT---- ------A---- --A------ --T---T----- 188
GT#16 KT232019        115     ---T-AA ------ ----------TT---- ------A---- --A-T---- --T---T----- 188
GT#17 KT232010        115     ---T-AA ------ ----------TT---- ------A-TAT- --A-T---- --T---T----- 188

GT#1  AB067721        211     GAATCAAAACTTTCAACAACGGATCTCTTGGTTCTGGCATC GATG AAGAACGCAGCGAAATGCGATAAGTAATGTGAA 288
GT#3  FJ654148        158     ----------------------------------------- ---- -------------------------------- 235
JAAVMX010000019    19556     ----------------------------------------- ---- -------------------------------- 19633
JAAVMX010000019    32200     ----------------------------------------- ---- -------------------------------- 32277
JAAVMX010000002 18702248     -- ---------------------------------C--- ---- -------------------------------- 18702325
JAAVMX010000018      851     ------------------------------- ---A------ ---- -------------------- --------- 928
JAAVMX010000019     6389     ----------- --CAC---------------------- ---- -------------------- --------- ------- 6463
JAAVMX010000019    44882     ----------------------------------------- ---- -------------------------------- 44959
GT#4  AB067744        207     A---T--------T--T-------------A---T-A----T A-A ------------------A-------------A-- 284
GT#5  AB067740        214     A---T-------T-------------A---A----T A-A --A------------------A---------A-A-- 288
GT#6  KJ720572        183     A---T-------T-------------A---A---- ---A -------T--T-----A-------A-A-- 222
GT#15 KT232017        189     ----T-------T--T-------------A---T-A----T A-A ------------------A-------------A-- 266
GT#16 KT232019        189     ----T-------T-------------A---A---- ---A ------------------A---------A-A-- 266
GT#17 KT232010        189     A---T-------T-------------A---A---- --A --A------------------A---------A-A-- 266

GT#1  AB067721        289     TTGCAGAATTCAGTGAACCA TCGAATCTTT GAACG CACATTG CGCCCGCCAGCACTCTGGCGGGCATGC CTGTC 362
GT#3  FJ654148        236     -------------------- ---------- ----- ------- -------------------------- ----- 309
JAAVMX010000019    19634     -------------------- ---------- ----- ------- -------------------------- ----- 19707
JAAVMX010000019    32278     -------------------- ---------- ----- ------- -------------------------- ----- 32351
JAAVMX010000002 18702326     --------------------- ------T---- ----- ------- -------------------------- ----- 18702398
JAAVMX010000018      929     -------------------- ----- C---------- -----C-------C-- --------------- ------- ----- 1000
JAAVMX010000019     6464     -----------------C--------- ----- ----C----------------------------- ----- 6539
JAAVMX010000019    44960     -------------------- ---------- ----- ------- --------------------------CT----- 45035
GT#4  AB067744        285     ---T------T--A--T- ---------- A---- T-T-C-A -------T--------A---------- ----- 358
GT#5  AB067740        289     --A------T---A---T- ---------- A---- --T---A -------T--------A---------- --A-- 362
GT#6  KJ720572        223     --AT-------T---A---- ---------- A---- T-T---A -------T--------A---------- ----- 296
GT#15 KT232017        267     ---T------T--A--T- ---------- A---- T-T---A -------T--------A---------- ----- 340
GT#16 KT232019        267     --A-------T---A--T- ---------- A---- T-T---A -------T--------A---------- ----- 340
GT#17 KT232010        267     --A-------T--A--T- ---------- A---- --T---- -------T--------A---------- --A-- 340

GT#1  AB067721        363     CGAGCGTCATCTCAACCCTCGAG CCCCCCGCCTCGCGG CGGCGGGGCCCGGCCTT GGGGGTCACGGCC CCGCGCC 437
GT#3  FJ654148        310     ----------------------- --------------- ----------------- ------------- ------- 384
JAAVMX010000019    19708     ----------------------- --------------- ----------------- ------------- ------- 19781
JAAVMX010000019    32352     ----------------------- --------------- ----------------- ------------- ------- 32425
JAAVMX010000002 18702399     ----------------------- --------------- ----------------- ------------- ------- 18702472
JAAVMX010000018     1001     ----------------------- --------------- ----------------- ---G--------- ---- -- 1074
JAAVMX010000019     6540     ----------------------- --------------- ----------------- ---- ------------C------- 6613
JAAVMX010000019    45036     -----------------------C---------A----TA--CG--C--------- ------TC-----AA------ 45119
GT#4  AB067744        359     -------T-C--T------TA-- --------------- ----------------- A----------- --A---- 432
GT#5  AB067740        363     -------T--T------------ --------------- ----------T------ A----T------ ------- 436
GT#6  KJ720572        297     -------T-C--T---- 313
GT#15 KT232017        341     -------T--T------------ --------------- ----------T------ A----T------ ------- 414
GT#16 KT232019        341     -------T-C--T----TA-- --------------- ----------T------ A----T------ --A---- 414
GT#17 KT232010        341     -------T---T---------- --------------- ----------T------ A----T------ ------- 414

GT#1  AB067721        438     G  CCCCCTAAACGC AG     TGGCGACC  CCGCCGCGGCTCC CCTGCGCAGTAGCT  CG  CTGAGAACCT 499
GT#3  FJ654148        385     -  ---------T-- --     -------   -------------  --------------- --  -C---G---C 446
JAAVMX010000019    19782     -  ------------ --     -------   -------------  --------------- --  ---------- 19844
JAAVMX010000019    32426     -  ------------ --     -------   -------------  --------------- --  ---------- 32488
JAAVMX010000002 18702473     -  ------------ --     -------   -------------  --------------- --  ---------- 18702534
JAAVMX010000018     1075     -  ------------ --     -------   -------------  --------------- --  ---------- 1136
JAAVMX010000019     6614     -  ---------G--G --     --------GC-----  -------------  -----GCT-------- 6680
JAAVMX010000019    45120     -TCC-T---GT----- --CCTAAGTG------- ----GA------A---T-A------CA-- ---------- 45189
GT#4  AB067744        433     -  ------------ --     -A-T----  -------------  ---A----A----- TA  --A------- 495
GT#5  AB067740        437     -  ----T------T --     -A------  -------------  ---A--TG------ -A  -C------- 529
GT#6  KJ720572        313
GT#15 KT232017        415     -  ----------T --     -A------  -------------  ---A--T-A----- -A  -C------- 477
GT#16 KT232019        415     -  ----------- --     -A------  -------------  ---A--A------ TA  --A------- 477
GT#17 KT232010        415     -  ----------T --     -A------  -------------  ---A--T-A----- -A  -C------- 477

GT#1  AB067721        500     CGCACCGGGAGCGCGGA     GGCGG TCACGCC  TGG    AA ACCACC     ACACCCTCCA             549
GT#3  FJ654148        447     --------A-------T     ----- ------- ---    -- ------     ---   ---             492
JAAVMX010000019    19845     -----------------     ----- ------- -   --   -- ------     ----------             19894
JAAVMX010000019    32489     -----------------     ----- ------- -   --   -- ------     ----------             32537
JAAVMX010000002 18702535     -----------------     ----- ------GTGT-  -- ------     ----------             18702586
JAAVMX010000018     1137     -----------------     ----- ------- -   --   -- ------     ----------             1186
JAAVMX010000019     6681     C----------------     ----- ------ -    -- ------ACA----------             6733
JAAVMX010000019    45190     ----------------CCCAT-----C--------  ---GATG--C-----ACC--------             45251
GT#4  AB067744        496     -----TA----------     ----- ------- --A    -- ----T    -T------             543
GT#5  AB067740        530     -------A--------     ----- ------- --A    -- ----T    --G--             544
GT#6  KJ720572        313
GT#15 KT232017        478     -------A---------     ----- ------- --A    -- ----T    --G--             522
GT#16 KT232019        478     -----TA----------     ----- ------- --A    -- ----T    -T------             525
GT#17 KT232010        478     -------A---------     ----- ------- --A    -- ----T    --G--             522
```

299   ***Figure 1. ITS sequence alignment of AB067721, repetitive ITS copies within genome JAAVMX000000000***
300   ***of H. sinensis Strain IOZ07, and AT-biased genotypes of O. sinensis.***
301   *The ITS sequences contain complete or partial ITS1-5.8S-ITS2 nrDNA segments. "GT" denotes the O. sinensis*
302   *genotype. The underlined sequence in black represents the 5.8S gene of GC-biased Genotype #1 H. sinensis.*
303   *AB067721 is the ITS sequence of GC-biased Genotype #1 H. sinensis. Genome assemblies*
304   *JAAVMX010000002, JAAVMX010000018 and JAAVMX010000019 were obtained from H. sinensis Strain*
305   *IOZ07 (Shu et al. 2020). One copy each within JAAVMX010000002 and JAAVMX010000018 indicated in*
306   ***green*** *shows 97.4% or 97.0% similarity to AB067721. JAAVMX010000019 contains 4 repetitive ITS copies,*
307   *including 2 copies in **black** (19404→19894 and 32048→32537), which are 100% identical to AB067721, and*
308   *2 other copies in **blue** (6233→6733 and 44729→45251), with 94.5% and 90.8% similarities to AB067721.*
309   *Sequences in **red** (AB067744, AB067740, KJ720572, KT232017, KT232019, and KT232010) represent AT-*
310   *biased Genotypes #4–6 and #15–17 of O. sinensis, respectively. The underlined "GAATTC" sequences in*
311   ***purple*** *represent the EcoRI endonuclease cleavage sites in the sequences of GC-biased Genotype #1 and GC-*
312   *biased genome assembly JAAVMX000000000. EcoRI endonuclease cleavage sites are absent in AT-biased*
313   *ITS sequences due to a single-base C-to-T (cytosine→threonine) transition. Hyphens indicate identical bases,*
314   *and spaces denote unmatched sequence gaps.*

315   Selker and his colleague (1985, 1990, 1997, 2002) reported frequent C-to-T (cytosine→thymine)

316   transition mutations and concomitant epigenetic methylation of nearly all remaining cytosine residues

317   in fungal 5S RNA genes associated with transcriptional silencing. Gladyshev (2017) summarized that

318   RIP (1) "occurs specifically in haploid parental nuclei that continue to divide by mitosis in preparation

319   for karyogamy and ensuing meiosis;" (2) "detects duplications of chromosomal DNA above a certain

320   length threshold (~0.4 kbp);" (3) "mutates cytosines on both strands of each DNA duplex in a pairwise

321   fashion (e.g., preferentially affecting none or both repeated sequences);" (4) "occasionally spreads

322   from duplicated sequences into neighboring nonrepetitive regions;" and (5) "remains unaffected in

323   *mei-2* crosses that exhibit a strong defect in meiotic pairing of homologous chromosomes."

324   The variable repetitive ITS copies characterized by multiple scattered insertion, deletion and

325   transversion point mutations (Table 3; Figure 1) might not have been generated through RIP

326   mutagenesis, which theoretically causes C-to-T (cytosine→threonine) and G-to-A (guanine→adenine)

327   transition mutations. Thus, the hypothesis of Li et al. (2020b) that RIP mutagenesis generated mutant

328   AT-biased genotypes of *O. sinensis* within the single genome of GC-biased Genotype #1 *H. sinensis*

329   lacks supporting evidence.

330

III-2. Genomic independence of multiple genotypes of *O. sinensis*

331

332   Answering the question of whether the AT-biased sequences of *O. sinensis* are components of a

333   single genome of Genotype #1 *H. sinensis* will provide key evidence for the "ITS pseudogene"

hypothesis. However, the sequences of mutant Genotypes #2–17 of *O. sinensis* were not present in genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000 and NGJJ00000000 of *H. sinensis* strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively, as shown in Tables 1–2 and Figure 1 (Hu et al. 2013; Zhu & Wu 2015; Li et al. 2016a, 2016b, 2020a, 2022a, 2022d, 2022e; Zhu & Li 2017; Zhu et al. 2018, 2019; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). The GC-biased sequences of repetitive ITS copies in the genomes of *H. sinensis* strains are 80.1%–89.9% similar to the sequences of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* (Table 1). The sequences of Genotypes #2–17 (both GC- and AT-biases) of *O. sinensis* are absent in the genome of Genotype #1 *H. sinensis,* as shown in Table 2, but instead belong to the genomes of independent *O. sinensis* fungi (Stensrud et al. 2008; Xiao et al. 2009; Zhu and Wu 2015; Li et al. 2016b; 2022d, 2022e; Zhu and Li 2017). The nonresidency measure of AT-biased *O. sinensis* genotypes does not support the notion that the AT-biased sequences are "ITS pseudogene" components of "a single genome" of GC-biased Genotype #1 *H. sinensis*.

The sequences of AT-biased Genotypes #4–6 and #15–17 of *O. sinensis* shown in red in Figure 1 contain numerous scattered transition alleles. They were generated through long-term accelerated nrDNA evolution from a common ancestor during a long history of evolution, regardless of whether this occurred *via* RIP or other mutagenic mechanisms. However, the multiple scattered transition point mutations in the AT-biased genotypes are genetically and phylogenetically distinct from the predominant insertion, deletion and transversion point mutations, as observed in the sequences of repetitive ITS copies within the genome of GC-biased Genotype #1 *H. sinensis* (Figures 1–2 and Tables 1–3). The notion that RIP mutagenesis induces the formation of AT-biased repetitive ITS copies in the single genome of Genotype #1 *H. sinensis* lacks genomic evidence. Even if it is proven that AT-biased genotypes were indeed induced by RIP mutagenesis, there has been no proof that they are genomic components of the GC-biased Genotype #1 genome that emerged after a new GC-biased Genotype #1 "ITS haplotype was generated" as "ITS pseudogenes … in a single genome", as assumed by Li et al. (2013, 2020b).

GC-biased genotypes, including the GC-biased genomic repetitive ITS copies, might have evolved in parallel with AT-biased genotypes from an ancient common ancestor during long-term evolution, as assumed by Stensrud et al. (2007). Zhang et al. (2009) detected GC-biased Genotype #3 in natural *C. sinensis* specimens collected from the Nyingchi District of Tibet and hypothesized that "Nyingchi might have acted as **a center of origin** of *O. sinensis* and *O. sinensis* might have been

transmitted from Nyingchi to other locations". Accordingly, GC-biased Genotype #3 might represent a modern immediate ancestor of Genotype #1 *H. sinensis* that has been identified in other production locations during long-term evolution. The authors described an evolution path from GC-biased Genotype #3 to Genotype #1 and transmission of *O. sinensis* from the "center of origin … spread eastward and westward first among southern regions on the Tibetan Plateau" and then "spread from south to north … at a later time". However, this evolutionary path and transmission of *O. sinensis* are not evolutionarily related to AT-biased genotypes.

The genomic repetitive ITS copies, regardless of their similarities to Genotype #1 AB067721, were phylogenetically clustered into the GC-biased clades in blue alongside the Bayesian tree in Figure 2. Apparently, they were phylogenetically distant from the AT-biased clades containing Genotypes #4–6 and #15–17 of *O. sinensis* shown in red alongside in the tree. Stensrud et al. (2007) concluded that the variation in the 5.8S gene sequences of AT-biased genotypes "far exceeds what is normally observed in fungi … even at higher taxonomic levels (genera and family)". Xiao et al. (2009) concluded that the sequences of AT-biased genotypes likely belong to independent *O. sinensis* fungi.

Figure 2 includes the 4 genomic repetitive ITS copies with low or relatively low similarities shown in Table 3, which are individually scattered in the GC-biased clades. The 2 low-similarity genomic repetitive ITS copies, 6233→6733 (94.5%) and 44729→45251 (90.8%), within genome assembly JAAVMX010000019 (*cf.* Tables 1 and 3), showed greater phylogenetic (horizontal lines) distances from the GC-biased clades. These results indicate that genetic variations in the repetitive ITS sequences shown in blue in Figure 1 had an impact on Bayesian phylogenetic clustering (Figure 2).

Figure 2. A Bayesian majority rule consensus phylogenetic tree.
Twenty-eight ITS segments of the genome assemblies (ANOV01021709, LKHE01000582, LWBQ01000008,
JAAVMX010000002, JAAVMX010000008, JAAVMX0100000017, JAAVMX0100000018, JAAVMX010000019,
NGJJ01000573, NGJJ01000582, NGJJ01000796, NGJJ01000798, and NGJJ01000799) of H. sinensis strains
(Co18, 1229, ZJB12195, IOZ07, and CC1406-203, respectively) and 25 ITS sequences of GC-biased
Genotypes #1–3 and #7–14 (in blue alongside the tree) and AT-biased Genotypes #4–6 and #15–17 of O.
sinensis (in red alongside the tree) were analyzed phylogenetically using MrBayes v3.2.7 software (cf. Section
II-4). Genome assemblies JAAVMX000000000 and NGJJ00000000 contain multiple repetitive ITS copies (cf.
Table 1). The percent similarities of the genomic ITS sequences with the representative Genotype #1 sequence
(AB067721) are shown in green alongside the tree.

Li et al. (2020b) hypothesized that (1) AT-biased *O. sinensis* genotypes "could have emerged either before or after" a new GC-biased "ITS haplotype was generated" and that (2) RIP mutagenesis is "responsible for nrDNA pseudogene formation (*i.e.*, AT-biased genotypes)". These 2 assumptions support the "ITS pseudogene" hypothesis that AT-biased sequences coexist "in a single genome" of GC-biased Genotype #1, as repeatedly stated by Li et al. (2013; 2017, 2019, 2020b). However, the genomic residency measure results obtained from different perspectives confirmed that the sequences of AT-biased genotypes are not genomic components of GC-biased Genotype #1 but instead belong to independent genomes of *O. sinensis* (further discussion below). AT-biased genotypes possessing numerous scattered transition alleles might have independently evolved through RIP or other mutagenic and evolutionary mechanisms, in parallel with GC-biased genotypes, during a long history of evolution from the common genetic ancestor and coexist in different combinations in different compartments of natural *C. sinensis* as independent *O. sinensis* fungal species, even "at higher taxonomic levels (genera and family)" (Stensrud et al. 2007; Xiao et al. 2009).

III-3. Inconsistency of the genetically divergent Genotype #1 *H. sinensis* strains

Li et al. (2013) reported the identification of heterogeneous ITS sequences of both GC-biased Genotype #1 and AT-biased Genotype #5 in 8 of 15 clones after 25 days of inoculation with natural *C. sinensis* mono-ascospores and hypothesized that Genotypes #1 and #5 coexisted "in a single genome". This speculation is irreconcilable with the simultaneous identification of 7 other clones derived from the same mono-ascospores that contained only homogenous sequences of Genotype #1. It is even more inexplicable why Li et al. (2013, 2017, 2019, 2020b) repeatedly called all AT-biased genotypes divergent pseudogenes, which coexisted "in single individuals of *O. sinensis*" or "in a single genome" of GC-biased Genotype #1 (the functional counterpart), while the researchers did not

identify AT-biased Genotypes #4, #6, and #15–17 in all 15 clones obtained from the *C. sinensis* mono-ascospores.

Bushley et al. (2013), who were collaborators of Li et al. (2013), reported that *C. sinensis* ascospores are multicellular heterokaryons with mono-, bi-, and trinucleated cellular structures. Zhang and Zhang (2015) wondered whether the binucleated cells (also trinucleated cells) of *C. sinensis* hyphae and ascospores contain homogenous or heterogeneous genetic material, meaning that these polykaryotic cells may contain heterogeneous sets of chromosomes and genomes. Furthermore, various mono-nucleated hyphal and ascosporic cells of *C. sinensis* might also contain heterogeneous genetic material.

By aggregating the information from molecular and genomic studies (Hu et al. 2013; Li et al. 2013, 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020), different genomic DNA samples of "pure" *H. sinensis* strains were examined and can be grouped as follows.

(1) Group 1 of *H. sinensis* strains, consisting of pure, homokaryotic anamorphic *H. sinensis* strains (1229, CC1406-203, Co18, IOZ07, and ZJB12195) (Hu et al. 2013; Li et al. 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). Total genomic DNA was isolated from these strains and subjected to genome-wide sequencing. The genomes contained no AT-biased genotype sequences.

(2) Group 2 of *H. sinensis* strains, consisting of 7 clones (Strains 1207, 1218, 1219, 1221, 1222, 1225 and 1229) among 15 clones derived from 25 days of incubation of natural *C. sinensis* mono-ascospores (Li et al. 2013). Total genomic DNA was isolated from these strains and shown to contain the homogenous ITS sequence of Genotype #1 *H. sinensis* but included no AT-biased sequences of *O. sinensis* genotypes.

(3) Group 3 of "*H. sinensis*" strains, including 8 other clones (Strains 1206, 1208, 1209, 1214, 1220, 1224, 1227 and 1228) obtained from the incubation of *C. sinensis* mono-ascospores for 25 days (Li et al. 2013). Total genomic DNA was isolated from these strains and showed genetic heterogeneity, with the coexistence of GC-biased Genotype #1 and AT-biased Genotype #5 of *O. sinensis*.

The *H. sinensis* strains of Groups 1 and 2 are similar, possessing a homogenous GC-biased Genotype #1 genome, although genome assemblies JAAVMX000000000 and NGJJ00000000 of *H. sinensis* Strains IOZ07 and CC1406-203 contained multiple GC-biased repetitive ITS copies (*cf.* Tables 1 and 3 and Figures 1–2) (Hu et al. 2013; Li et al. 2013, 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020). Although a single copy of ITS sequences was identified in genome assemblies

ANOV00000000, LKHE00000000, and LWBQ00000000 of *H. sinensis* Strains Co18, 1229, and

ZJB12195, respectively (Hu et al. 2013; Li et al. 2016a; Jin et al. 2020), the genomes of these strains

may theoretically contain more repetitive ITS copies that could have been discarded during the

assembly of genome shotgun sequences. Conceivably, reassembling the genome shotgun reads may

very likely identify additional repetitive ITS copies in the genomes of Group 1 Strains Co18, 1229,

and ZJB12195, among which Strain 1229 is also a Group-2 strain derived from mono-ascospores by

Li et al. (2013, 2016a). Similarly, the genome sequencing of Group-2 strains may very likely identify

multiple repetitive ITS copies.

Although the anamorphic *H. sinensis* strains in Groups 2 and 3 were derived after 25 days of

liquid culture of the same *C. sinensis* mono-ascospores (Li et al. 2013), they are genetically different

and may likely have multicellular heterokaryotic mycelial structures, similar to the mycelia derived

from microcycle conidiation of the *H. sinensis* ascospores initially cultured on the solid medium plate

for 30 days and subsequently in liquid medium for 10–53 days, as shown in Figures 2–3 of (Xiao et

al. 2011). Li et al. (2013) reported that the Group 3 strains included cocultured GC-biased Genotype

#1 and AT-biased Genotype #5 of *O. sinensis* fungi, which were genomically independent and

coexisted in heterokaryotic mycelial cells, indicating either diploid/polyploid mycelial cells

containing heterogeneous sets of chromosomes/genomes or impure mycelia, each containing a

different chromosome/genome. However, the authors did not identify any haploid clones containing

only AT-biased Genotype #5, indicating that this AT-biased *O. sinensis* fungus may not exhibit

independent culturability under the *in vitro* experimental settings, although other studies have

reported the identification of single AT-biased Genotype #4 or #5 of *O. sinensis*, without coculture

with GC-biased Genotype #1, in cultures of natural and cultivated *C. sinensis* under different

experimental settings (Mao et al. 2013; Wei et al. 2016). Unfortunately, Li et al. (2013, 2017, 2019,

2020b) neglected the differences in genetic material between Group 2 and 3 strains, which were

presumably derived from the same or different cells of multicellular heterokaryotic ascospores

(Bushley et al. 2013; Zhang and Zhang 2015; Li et al. 2020a, 2022d, 2022e).

In addition to the genome assembly LKHE00000000 derived from the Group-2 *H. sinensis* Strain

1229 by Li et al. (2016a), the genome sequencing of one of the heterogeneous Group-3 strains may

provide critical genomic evidence to validate the notion that AT-biased Genotype #5 of *O. sinensis*

coexists as an "ITS pseudogene … in a single genome" of GC-biased Genotype #1. Unfortunately,

the authors elected not to do so since 2012 after obtaining the Group-3 strains, or perhaps failed in

such an endeavor due to the technical difficulties in shotgun sequence assembly because of the heterogeneous sequences from more than one set of genomes. Consequently, they elected willingly or unwillingly to pass on the opportunity to experimentally confirm their "ITS pseudogene" hypothesis that AT-biased Genotype #5 and GC-biased Genotype #1 coexist "in a single genome". Studies from other groups that are reviewed in this paper have confirmed that the sequences of all AT-biased genotypes belong to the genomes of independent fungi (Stensrud et al. 2007; Xiao et al. 2009; Zhu et al. 2010; Yao et al. 2011; Li et al. 2016, 2020a, 2022a, 2022d, 2022e) and that heterokaryotic ascospores and hyphae with multiple mono-, bi-, and tri-nucleated cells contain heterogeneous sets of chromosomes and genomes (Bushley et al. 2013; Zhang and Zhang 2015).

Li et al. (2013) did not report the identification of ITS sequences of AT-biased Genotypes #4, #6, and #15–17 of *O. sinensis* in genomic DNA preparations that were extracted from the 15 ascosporic strains in Groups 2 and 3. The expansion of the "ITS pseudogene" hypothesis from only Genotype #5 to all AT-biased genotypes was apparently an overinterpretation with no supporting evidence. As shown in Tables 1–3 and Figures 1–2, genomic and phylogenetic analyses did not support the notion that the "diverged ITS paralogs" (*i.e.*, AT-biased Genotypes #4–6 and #15–17) coexist "in a single genome" of *O. sinensis*. Li et al. (2022d) demonstrated that the sequences of Genotypes #4 and #15 of AT-biased Cluster-B (*cf.* Figure 2) were absent in the genomic DNA pool extracted from natural *C. sinensis* ascospores. Instead, sequences of these genotypes belong to independent genomes of *O. sinensis* fungi, regardless of whether the AT-biased genotypes might have been generated *via* RIP or other mutagenic mechanisms during the long evolutionary history of *O. sinensis*.

III-4. Differential co-occurrence and dynamic alteration of multiple *O. sinensis* genotypes in the compartments of natural *C. sinensis* during maturation

A BLAST search of the GenBank database revealed 668 ITS sequences of *O. sinensis* (GenBank taxid: 72228), 228 of which belong to AT-biased Genotypes #4–6 and #15–17 (Li et al. 2016b, 2022d, 2022e; Zhu et al. 2018, 2019; Zhu & Li 2017). After Kinjo and Zang (2001) reported for the first time the 3 phylogenetic clades of mutant *O. sinensis* sequences, Stensrud et al. (2007) and Zhang et al. (2013) analyzed the phylogeny of 71 and 397 ITS sequences, respectively, and confirmed the 3 phylogenetic clades, *i.e.*, Groups A–C (GC-biased Genotype #1 and AT-biased Genotypes 4–5), which consisted of correct *O. sinensis* sequences. Additional GC- and AT-biased *O. sinensis* genotypes have been successively discovered (*cf.* Table 2) (Yang et al. 2008; Xiao et al. 2009; Zhang

et al. 2009; Zhu et al. 2010, 2012; Chen et al. 2011; Zhu and Wu 2015; Li et al. 2013, 2016, 2022a, 2022d,2022e; Zhu and Li 2017). Molecular mycology studies confirmed the genetic heterogeneity of natural *C. sinensis* with differential co-occurrence of several GC- and AT-biased genotypes of *O. sinensis* in different combinations in the different compartments of natural *C. sinensis,* and the biomasses/abundances of the genotypes were altered dynamically during *C. sinensis* maturation (Xiao et al. 2009; Zhu et al. 2010, 2012, 2018, 2019; Gao et al. 2011, 2012; Yao et al. 2011; Mao et al. 2013; Li et al. 2016b, 2016c, 2022a, 2022c, 2022d, 2022e).

Many studies have used pairs of "universal" primers (*e.g.*, *ITS1–ITS5*) for PCR and identified one of the GC- and AT-biased genotypes of *O. sinensis* in natural *C. sinensis* specimens collected from different production areas or cultivated *C. sinensis* (Engh 1999; Zhao et al. 1999; Kinjo and Zang 2001; Liu et al. 2001; Stensrud et al. 2005; Wei et al. 2006, 2016; Chen et al. 2011; Mao et al. 2013). Wei et al. (2016) reported an industrial artificial cultivation project and identified a single teleomorphic AT-biased Genotype #4 of *O. sinensis* in cultivated *C. sinensis* and a single teleomorphic GC-biased Genotype #1 in natural *C. sinensis* as the teleomorph control, indicating the existence of at least 2 *O. sinensis* teleomorphs. This study revealed a species contradiction between the anamorphic inoculants (reportedly 3 *H. sinensis* strains) and teleomorphic AT-biased Genotype #4 in cultivated *C. sinensis*.

Other studies optimized and combined the use of a variety of molecular techniques and identified naturally cooccurring GC- and AT-biased genotypes of *O. sinensis* in different combinations in the caterpillar body, stroma, stromal fertile portion with numerous ascocarps, and ascospores of natural *C. sinensis* (Yang et al. 2008; Xiao et al. 2009; Zhu et al. 2010, 2012, 2018, 2019; Gao et al. 2011, 2012; Yao et al. 2011; Zhu and Wu 2015; Li et al. 2016b, 2020a, 2022a, 2022c, 2022d, 2022e; Zhu and Li 2017). Xiao et al. (2009) used genotype-specific primers for the first time and a nested PCR approach to amplify mutant sequences of *O. sinensis* from the stroma and caterpillar body of immature and mature *C. sinensis* specimens collected from geographically remote production areas. They reported the consistent occurrence of GC-biased Genotype #1 and inconsistent occurrence of AT-biased Genotypes #4–5 of *O. sinensis* and then concluded that the AT-biased genotypes likely belonged to independent fungal species. Two of the authors also identified the first sequence (EU555436) of AT-biased Genotype #6 and uploaded it to GenBank. The use of genotype-specific primers combined with touchdown PCR, restriction fragment length polymorphism (RFLP), cloning-based amplicon sequencing, SNP mass spectrometry genotyping techniques, etc., in other studies

greatly improved the detection specificity and efficiency and facilitated the identification of differential coexistence and dynamic alterations of GC- and AT-biased genotypes of *O. sinensis* in the caterpillar body, stroma, stromal fertile portion with numerous ascocarps, and ascospores of natural *C. sinensis* during maturation (Zhu et al. 2010, 2012, 2019; Chen et al. 2011; Gao et al. 2011, 2012; Yao et al. 2011; Zhu and Wu 2015; Li et al. 2016b, 2020a, 2022a, 2022c, 2022d, 2022e; Zhu and Li 2017). To date, 17 genotypes of *O. sinensis* have been identified, which are genetically and phylogenetically distinct (Table 2 and Figures 1–2).

In contrast to the inconsistent cooccurrence of Genotypes #1 and #4–5 (Xiao et al. 2009), Zhu et al. (2010) used *H. sinensis*-specific primers, *Hsprp1*/*Hsprp3*, and identified the consistent coexistence of GC-biased Genotypes #1 and #2 in the stromata of immature, maturing, and mature *C. sinensis* specimens collected in Kangding County of Sichuan Province, China. Notably, the ITS sequences of the GC-biased genotypes contained an *Eco*RI endonuclease cleavage site (G▾AATTC) from nucleotides 294–299 of AB067721 (a representative sequence of Genotype #1 *H. sinensis*), which is underlined in purple in Figure 1 and was lost in all AT-biased genotypes due to a single base C-to-T (cytosine→thymine) transition mutation (GAATTT). Sequencing the target amplicons (440 bp, based on the design of primers) obtained ITS sequences of Genotype #1 *H. sinensis* as shown in Figure 10 of (Zhu et al. 2010). A faster migrating amplicon (320 bp) on the agarose gel showed Genotype #2 ITS sequences (MG770309).

Due to the possibility that the predominant 440-bp amplicons might not be homogeneous based on the various similarities of primers *Hsprp1*/*Hsprp3* to the ITS sequences of various *O. sinensis* genotypes (although only a single sequence was obtained from the target amplicons), the 440-bp amplicons obtained from immature, maturing and mature stromata were subjected to overnight *Eco*RI digestion. Two digested fragments (240 and 200 bp) were generated with extremely low densities in the immature stroma, representing GC-biased Genotype #1, which was the minor species in the immature *C. sinensis* stroma; the densities of the digested fragments were increased in the maturing and mature stromata, as shown in Figure 13 of (Zhu et al. 2010). Further quantitative analysis demonstrated different maturational alterations of the co-occurring Genotypes #1 and #2 in an asynchronous and disproportional manner in the *C. sinensis* stromata during maturation (Figure 3, modified from Figure 6 of Zhu et al. 2012). The dynamic but asynchronous alteration of the abundance of Genotypes #1 and #2 during *C. sinensis* maturation indicates genomic independence of these 2 *O. sinensis* genotypes but does not support the idea that "Different types of ITS sequences

575 were thus proved to coexist in a single genome" (Li et al. 2013).

Figure 3. Dynamic alterations in the abundance of amplicons of Genotypes #1 and #2 of O. sinensis in the
stromata of C. sinensis during maturation *(modified from Figure 6 of* Zhu et al. 2012*).*
Immature C. sinensis specimens had very short stromata of 1.5 cm on average. Maturing C. sinensis specimens
had stromata of 4.0 cm on average, without the formation of an expanded fertile portion close to the stromal
tip. Mature C. sinensis specimens had long stromata of 7.0 cm on average and showed the formation of an
expanded stromal fertile portion (SFP in red), which was densely covered with numerous ascocarps.

To examine dynamic alterations of the natural biomasses of GC- and AT-biased genotypes of *O.*

sinensis without PCR manipulation, which may cause nonlinear, exponential amplification of DNA

sequences, the Southern blotting technique was applied to examine genomic DNA after enzymolysis

with endonuclease *Eco*RI (Zhu et al. 2010). This non-PCR approach identified a single *Eco*RI-

sensitive GC-biased DNA moiety in genomic DNA isolated from a pure *H. sinensis* strain (a gift from

a senior mycological taxonomist, Prof. Ying-Lan Guo) (*cf.* Figure 4B of Zhu et al. 2010) and a doublet

(both *Eco*RI-sensitive Genotype #1 and *Eco*RI-resistant AT-biased genotypes) in genomic DNA

isolated from natural *C. sinensis* (Figure 4; reuse of Figure 6 of Zhu et al. 2010 with permission). The

lack of *Eco*RI-resistant AT-biased genotypes in pure *H. sinensis* shown in Figure 4B of (Zhu et al.

2010) indicates that the sequences of AT-biased genotypes of *O. sinensis* are absent in the genome of

GC-biased Genotype #1 *H. sinensis*. The upper panel of Figure 4 shows that Southern blotting without

PCR amplification demonstrated the following:

Figure 4. Southern blot of C. sinensis nrDNA in the stroma and caterpillar body of natural C. sinensis during maturation. *[Reproduced with permission from AJBMS (www.nwpii.com/ajbms) Am J Biomed Sci 2010; 2(3): 217-238] [Zhu et al. 2010].*
*Caterpillar refers to caterpillar body. Genomic DNA templates were isolated from the stromata or caterpillar bodies of immature (Im) or mature (M) C. sinensis specimens collected from Sichuan Province in China and prepared using the restriction enzymes EcoRI, DraI and AvaI. **The upper panel** was probed with an H. sinensis-specific probe. **The lower panel** was probed with a nonspecific 18S internal control probe.*

(1)   Comparison of *O. sinensis* genotypes in the stromata: the left 2 lanes of the upper panel of Figure 4 show that the slower-migrating moieties (*Eco*RI-resistant AT-biased genotypes) predominate naturally without PCR amplification in the stromata of immature and mature *C. sinensis*. The faster-migrating moieties (*Eco*RI-sensitive GC-biased *H. sinensis*) are always minor moieties in the *C. sinensis* stromata.

(2)   Comparison of *O. sinensis* genotypes in the caterpillar bodies: although the second lane from the right in the upper panel of Figure 4 seems to show a single *Eco*RI-sensitive moiety in the immature caterpillar body but a doublet in the mature caterpillar body shown in the far right lane, PCR experiments identified *Eco*RI-resistant AT-biased genotypes in both samples. These results suggest that the natural quantity of *Eco*RI-resistant AT-biased genotypes in the immature caterpillar body was below the detection limit of Southern blotting, which is much less sensitive than PCR-based techniques. The quantity of AT-biased genotypes was dramatically and disproportionally increased from a micro-quantity in the immature caterpillar body to a level comparable to that of *Eco*RI-sensitive GC-biased *H. sinensis* in the mature caterpillar body.

(3)   Comparison of *O. sinensis* genotypes in immature *C. sinensis*: the first and third lanes from the left in Figure 4 show that the abundances and density ratios of *Eco*RI-resistant AT-biased genotypes and *Eco*RI-sensitive GC-biased *H. sinensis* differed significantly in the stroma and caterpillar body of immature *C. sinensis*, which were sampled from the same specimens.

(4) <u>Comparison of *O. sinensis* genotypes in mature *C. sinensis*</u>: the first and third lanes from the right in Figure 4 show that the abundances and density ratio of *Eco*RI-resistant and *Eco*RI-sensitive genotypes differed in the stroma and caterpillar body of mature *C. sinensis*, which were sampled from the same specimens.

The disproportional distribution of GC- and AT-biased *O. sinensis* genotypes in the stromata and caterpillar bodies of natural *C. sinensis* and their asynchronous alterations of the natural biomasses during *C. sinensis* maturation indicate that the sequences of AT-biased *O. sinensis* genotypes do not reside in the genome of Genotype #1 *H. sinensis*. Instead, the genomically independent AT-biased *O. sinensis* genotypes are interindividually accompanied by GC-biased Genotype #1 *H. sinensis* in the stroma and caterpillar body of natural *C. sinensis*.

If "Different types of ITS sequences were thus proved to coexist in a single genome" as assumed by Li et al. (2013), the distribution of GC- and AT-biased *O. sinensis* genotypes should be proportional in the *C. sinensis* stroma and caterpillar body and any biomass alterations of the co-occurring genotypes should be synchronous during *C. sinensis* maturation. Southern blotting, as well as RFLP after pretreatment with *Eco*RI enzymolysis and MassARRAY SNP mass spectrometry genotyping assays, clearly demonstrated asynchronous and disproportional alterations of the GC- and AT-biased *O. sinensis* genotypes (Zhu et al. 2010, 2012; Gao et al. 2011, 2012; Yao et al. 2011) and confirmed the genomic independence of the *O. sinensis* genotypes, which belong to independent *O. sinensis* fungi (Stensrud et al. 2007; Xiao et al. 2009).

Zhu et al. (2010) and Gao et al. (2011, 2012) demonstrated that Genotype #4 of AT-biased Cluster-B of *O. sinensis* predominated in the immature stroma of natural *C. sinensis,* which was in its asexual growth stage; however, its abundance dramatically declined in the mature stroma*,* which was in its sexual reproductive stage. The maturational decline of Genotype #4 in the *C. sinensis* stroma was accompanied by a significant reciprocal elevation in the abundance of Genotype #5 of AT-biased Cluster-A of *O. sinensis*, indicating that the multiple AT-biased genotypes belong to independent *O. sinensis* fungi.

Li et al. (2022c, 2022d) designed several pairs of genotype-specific primers combined with cloning-based amplicon sequencing and identified Genotypes #5–6 and #16 of AT-biased Cluster-A in the ascospores of natural *C. sinensis*, co-occurring with GC-biased Genotype #1. They identified the sequences of Genotypes #4 and #15 of AT-biased Cluster-B in the stromal fertile portion densely covered with numerous ascocarps but not in ascospores obtained from the same *C. sinensis* specimens,

consistent with the findings reported by Li et al. (2013) that AT-biased Genotype #4 was not present in 15 cultures of *C. sinensis* mono-ascospores. Li et al. (2022d) also identified Genotype #13 in semiejected ascospores and Genotype #14 in fully ejected ascospores collected from the same pieces of natural *C. sinensis*. Li et al. (2022c) revealed a high abundance of Genotypes #6 and #15 in the preejection stromal fertile portion of natural *C. sinensis,* but the abundance dramatically declined in the stromal fertile portion after ascospore ejection, while Genotype #5 continued to show high abundance. All the scientific evidence supports the genomic independence hypothesis for multiple *O. sinensis* genotypic fungi initially proposed by Xiao et al. (2009).

After the hypothesis of "ITS pseudogenes … in a single genome" was published and subjected to academic discussion, Li et al. (2013) had sufficient time to validate their hypothesis by conducting genome sequencing on one of the Group 3 strains (*cf.* Section III-3); however, scientific papers addressing this have yet to appear. Instead, Li et al. (2013, 2017, 2019, 2020c) stated repeatedly that "Different types of ITS sequences were thus proved to coexist in a single genome", "from a single genome, and not from different genomes", and "The coexistence of diverged ITS paralogs in a single genome". These scientists either did not conduct genome sequencing of one of the Group 3 strains or may have failed the experiments due to difficulties in assembly of shotgun reads to experimentally verify their hypothesis that Genotypes #1 and #5 coexist in "a single genome", although this genome sequencing strategy may not be able to confirm the coexistence of AT-biased Genotypes #4, #6 and #15–17 as the genomic components of "a single genome" of GC-biased Genotype #1 *H. sinensis*.

Bushley et al. (2013) and Hu et al. (2013) hypothesized self-fertilization as the reproductive mode of Genotype #1 *H. sinensis* under homothallism or pseudohomothallism based on the identification of mating-type genes of both *MAT1-1* and *MAT1-2* idiomorphs in 2 *H. sinensis* strains. Li et al. (2020b) stated accordingly that "*O. sinensis* is one of the only known homothallic species in *Ophiocordycipitaceae* … the homothallic life-style may have increased the likelihood and frequency of sex in *O. sinensis*, leading to increased RIP in this species." However, Zhang and Zhang (2015) revealed differential occurrence of the mating-type genes of *MAT1-1* and *MAT1-2* idiomorphs in numerous *H. sinensis* strains and alternatively hypothesized facultative hybridization as the mode of *O. sinensis* reproduction.

Li et al. (2022b, 2022e) reported significant intraspecific genetic variations among *H. sinensis* strains and differential expression of *H. sinensis* genes. Li et al. (2021) further reviewed the genetic information of 236 *H. sinensis* strains and confirmed the differential occurrence of mating-type genes

of the *MAT1-1* and *MAT1-2* idiomorphs: 22 strains contained only MAT1-1-1 genes, 65 possessed

only MAT1-2-1 genes, and 149 had both genes, consistent with the findings reported by Zhang and

Zhang (2015). Li et al. (2021) identified differential but reciprocal transcription of mating-type genes

in two *H. sinensis* strains. The MAT1-1-1 transcript was absent, but the MAT1-2-1 transcript was

present in transcriptome assembly GCQL00000000 of *H. sinensis* Strain L0106 (Liu et al. 2015; Li

et al. 2021). On the other hand, the MAT1-1-1 gene was shown to be transcribed with normal splicing

of both intron I and II in *H. sinensis* Strain 1229, but the MAT1-2-1 transcript included an unspliced

intron I, which contains 3 stop codons (Bushley et al. 2013; Li et al. 2021). Bushley et al. (2013)

stated that the total "RNA was treated with DNase I … until no DNA contamination could be detected"

prior to reverse transcription and cDNA library construction, and "the products (*i.e.*, cDNA) were

derived from mRNA and did not result from contamination of genomic DNA. In contrast, comparison

of DNA and cDNA sequences of MAT1-2-1 revealed only one spliced intron of 55 bp (Fig S2)." Thus,

the special phenomenon of MAT1-2-1 transcription reported by Bushley et al. (2013) indicates the

production of a largely truncated and dysfunctional MAT1-2-1 protein lacking the majority of the

protein, encoded by exons II and III. Thus, the transcriptional silencing of one of the mating-type

genes and the MAT1-2-1 transcript with an unspliced intron I constitute *O. sinensis* reproduction

controls at the transcriptional level and the coupled transcriptional-translational level.

Accordingly, Li et al. (2021) concluded self-sterility of GC-biased Genotype #1 *H. sinensis*,

suggesting a requirement of sexual partners for reproduction. These partners could be either

intraspecific species, such as *H. sinensis* Strains L0106 and 1229, for physiological heterothallism

regardless of whether *H. sinensis* is monoecious or dioecious, or heterospecific fungal species that

may undergo hybridization if they are able to overcome interspecific reproductive barriers (Zhang

and Zhang 2015; Li et al. 2021, 2022d, 2022e). Because the taxonomic positions of Genotypes #2–

17 of *O. sinensis* have not been determined, it is debatable whether the coexisting genotypes represent

intraspecific species involved in physiological heterothallism or heterospecific species involved in

fungal hybridization. Thus, the notion of "the homothallic life-style in *O. sinensis* … leading to

increased RIP", put forth by Li et al. (2020b), may be groundless because "the homothallic life-style

in *O. sinensis*" is scientifically incorrect and "increased RIP" that theoretically induces C-to-T and

G-to-A transition mutations was not confirmed based on the analysis of the multiple repetitive ITS

copies in the genomes of Genotype #1 *H. sinensis* strains, as analyzed in Section III-1.

III-5. Transcriptional silencing of 5.8S genes

In addition to the identification of both Genotypes #1 and #5 ITS sequences from genomic DNA isolated from 8 of 15 cultures of multicellular heterokaryotic mono-ascospores, Li et al. (2013) examined the 5.8S gene transcripts of these clones. They reported the identification of 5.8S cDNA for Genotype #1, but not for Genotype #5, in a cDNA library constructed from total ascosporic RNA extracted from Clone 1220, one of the 8 genetically heterogeneous clones. However, the authors overinterpreted that the 5.8S genes of all AT-biased genotypes were permanently dysfunctional pseudogenic components coexisting in "a single genome" with a functional copy of GC-biased Genotype #1, although they did not detect the ITS sequences of AT-biased Genotypes #4, #6 and #15–17 in the genomic DNA isolated from any of the ascosporic clones. The logical reasoning underlying the dysfunctional pseudogene hypothesis was explained by Li et al. (2019) as follows: "If AT-biased ITS sequences are not pseudogenes, 5.8S cDNA should have been detected because rRNAs are critical and essential for protein synthesis in all living organisms", which touches on 2 scientific issues. First, are the 5.8S genes of all genotypes of *O. sinensis* actively transcribed in *C. sinensis* ascospores (discussed in this section)? Second, is it correct to conclude that "5.8S cDNA should have been detected" by the techniques designed and used by Li et al. (2013) (discussed in Section III-6)?

Xiang et al. (2014) reported a metatranscriptome study of natural *C. sinensis* specimens collected from Kangding County of Sichuan Province, China. The GAGW00000000 metatranscriptome sequences do not contain any 5.8S gene transcripts of *C. sinensis*-associated fungi (including Genotype #1 and all other genotypes of *O. sinensis*), indicating transcriptional silencing of 5.8S genes of all cocolonized fungi in natural settings.

Xia et al. (2017) reported another metatranscriptome study of mature *C. sinensis* specimens collected from Deqin County of Yunnan Province, China. The assembled metatranscriptome sequences were not uploaded to GenBank but instead to another depository server www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas. Similarly, this assembled metatranscriptome contains no 5.8S gene transcripts. However, the authors uploaded the unassembled shotgun reads (SRR5428527) to GenBank. A BLAST search of the SRR5428527 Shotgun Read Archive (SRA) using the MegaBLAST algorithm identified 1894 metatranscriptome shotgun reads (100 bp each), which were 100% homologous to the 5.8S gene sequences (156 bp) of GC-biased Genotype #1 *H. sinensis*.

It is well established that fungal genomes contain multiple copies of rDNA ITS1-5.8S-ITS2

sequences (Tables 1 and 3; Figures 1–2). The 1894 unassembled metatranscriptome shotgun reads

(SRR5428527) of natural *C. sinensis* may be derived from highly homologous nonrDNA sequences

of slowly evolved genes within fungal metagenomes, while there are no rDNA 18S and 28S gene

segments of *H. sinensis* homologous to the 5.8S gene. Figure 5 shows the alignment of the 5.8S gene

(the query sequence; segment 218–373 of Genotype #1 *H. sinensis* AB067721) and overlap of the

first 75 of the 1894 metatranscriptome shotgun reads of mature natural *C. sinensis*. The short length

(100 bp) of SRA shotgun reads resulted in overlap between the 1894 fragmented sequence reads and

transcripts of nonrDNA genes (further discussion below).

Figure 5. Alignment of the query sequence in green (the 5.8S gene of Genotype #1 H. sinensis AB067721) and the 75 of 1894 overlapping (pink bars) shotgun reads (GenBank SRR5428527) of mature C. sinensis.

Various degrees of similarity were found when the 1894 unassembled metatranscriptome shotgun reads were aligned with the 5.8S gene sequences of the 17 genotypes of *O. sinensis*. Table 4 shows 6 examples of shotgun reads of SRX2718557 aligned with the 5.8S sequences of *O. sinensis* genotypes, *Paecilomyces hepiali*, an AB067719-type Group-E fungus, and *Tolypocladium sinensis* (Group D) (Stensrud et al. 2007). Obviously, the high homology between the shotgun reads and the various 5.8S gene sequences may not conclusively determine their fungal sources in natural *O. sinensis*, which contains >90 species spanning >37 fungal genera (Zhang et al. 2010, 2018; Xiang et al. 2014; Xia et al. 2015, 2017). Because no 5.8S transcript was identified from the assembled metatranscriptome sequences in the same study, the shotgun reads might not be derived from the 5.8S genes of multiple cocolonizing fungi, including the multiple genotypes of *O. sinensis*. Liu et al. (2015) reported the mRNA transcriptome assembly GCQL00000000 derived from *H. sinensis* Strain L0106, which contained no transcript of rRNA or tRNA. Continuous *in vitro* liquid fermentation performed in this study induced a nonlinear reduction in the total number of transcriptomic unigenes, first from 25,511 (after 3 days of fermentation) to 25,214 (after 6 days of fermentation) and then more sharply to 16,245 (after 9 days of fermentation), indicating the nonnatural switching on or off of many genes. The average length of transcripts of Strain L0106 increased nonlinearly, with a gradual decrease in the GC content from 681 nt (GC=58.2%) (3 days) to 682 nt (GC=57.9%) (6 days) and then a dramatic increase to 994 nt (GC=57.0%) (9 days). The differential transcriptional activation and silencing of *H. sinensis* genes were caused by continuous *in vitro* liquid fermentation of *H. sinensis* Strain L0106. Presumably, the prolonged 25-day liquid incubation period adopted by Li et al. (2013) might have a greater impact on the unbalanced transcription of long and short fungal genes with various GC content biases.

Apparently, Li et al. (2013, 2017, 2019, 2020b) neglected the evidence of transcriptional silencing of 5.8S genes demonstrated in metatranscriptome studies in natural settings and that of dynamic and unbalanced transcriptional activation and silencing of numerous functional genes during the continuous *in vitro* fermentation or liquid incubation of Genotype #1 *H. sinensis*. Because of the conservative evolution of 5.8S genes and many other nonrDNA genes and the incomplete study design and controversial findings reported by Li et al. (2013), it needs to be verified (probably using other sophisticated transcriptional expression methods) whether the identified cDNA was truly derived from the 5.8S gene of Genotype #1 *H. sinensis*, or from mutant genotypes of *O. sinensis* or

other colonized fungi, or even from nonrRNA genes (further discussion below) before confidently

and reasonably declaring that the 5.8S genes of the multiple AT-biased genotypes of *O. sinensis* are

truly permanently nonfunctional pseudogenes under the "ITS pseudogene … in a single genome"

hypothesis.

Table 4. *Percent similarity between the unassembled metatranscriptome shotgun reads (SRA:*

SRR5428527) of natural C. sinensis and the 5.8S genes of multiple genotypes of O. sinensis and

Paecilomyces hepiali, an AB067719-type fungus (Group-E) and Tolypocladium sinensis (Group D).

Geno-type	GenBank Accession #	% similarity between 5.8S genes and *C. sinensis* shotgun sequences					
		SRR5428527. 17546327.1	SRR5428527. 17554054.1	SRR5428527. 17565575.2	SRR5428527. 17679995.2	SRR5428527. 17911498.2	SRR5428527. 17929569.2
#1	AB067721	**100%**	**100%**	**100%**	**100%**	**100%**	**100%**
#2	MG770309	96%	96%	**100%**	96%	**98%**	96%
#3	HM595984	**100%**	**99%**	**99%**	**100%**	**99%**	**100%**
#7	AJ488254	**98%**	**98%**	**99%**	**98%**	**98%**	**98%**
#8	GU246286	94%	96%	95%	94%	**97%**	94%
#9	GU246288	**99%**	**99%**	**99%**	**100%**	**99%**	**100%**
#10	GU246287	96%	94%	93%	95%	93%	**97%**
#11	JQ695935	**100%**	**100%**	**100%**	**100%**	**100%**	**100%**
#12	GU246296	**99%**	**99%**	**99%**	**99%**	**99%**	**99%**
#13	KT339190	**98%**	93%	93%	**98%**	93%	**98%**
#14	KT339178	**98%**	93%	93%	**98%**	93%	**98%**
#4	AB067744	86%	87%	87%	83%	87%	83%
#5	AB067740	85%	87%	87%	84%	87%	84%
#6	KJ720572	85%	85%	87%	84%	86%	84%
#15	KT232017	85%	88%	88%	84%	88%	84%
#16	KT232019	86%	86%	87%	85%	87%	85%
#17	KT232010	87%	88%	88%	87%	88%	87%
P.h.	EF555097	**99%**	**98%**	**97%**	**99%**	**98%**	**99%**
	AB067719	**98%**	93%	93%	**98%**	93%	**98%**
T.s.	AB067720	**98%**	**97%**	**96%**	**98%**	**97%**	**98%**

Note: GC-biased genotypes are listed in blue, *and AT-biased genotypes are listed in* red. *GenBank*

sequence ID SRA: SRR5428527 fragments (17546327.1, 17554054.1, 17565575.2, 17679995.2,

17911498.2, and 17929569.2) are the unassembled transcriptome shotgun sequences obtained from

the mature C. sinensis specimen collected from Deqin County of Yunnan Province by Xia et al. *(2017).*

P.h. refers to Paecilomyces hepiali; AB067719 refers to an AB067719-type Group-E fungus; T.s.

refers to Group-D Tolypocladium sinensis (Stensrud et al. *2007). Highly homologous percent*

similarities (97%–100%) are bolded in **green**.

III-6. Identification and analysis of 5.8S cDNA using 5.8S-F/R primers

In addition to the discussion above of the transcriptional silencing of 5.8S genes in natural settings and that of the significant impact of continuous *in vitro* liquid-phase culture on unbalanced gene transcription, the second issue related to the logic given by Li et al. (2019) is whether the methods designed by Li et al. (2013) were scientifically capable of identifying 5.8S gene cDNAs of multiple *O. sinensis* genotypes (Li et al. 2020a).

For instance, Li et al. (2019) referred to "The 5.8S-F/R primers … located in the most conserved region", which may reduce the primers' specificity for 5.8S cDNA because of the conservative evolution of the 5.8S gene and many other genes. Figure 6 shows that primer 5.8S-F is 95% (20/21 nt) similar and 100% (21/21 nt) identical to the 5.8S transcript sequence of AT-biased Genotypes #4–5, respectively. Primer 5.8S-R is 90% (18/20 nt, with 2 flanking residues downstream in the ITS2 region) and 95% (19/20 nt) similar to the 5.8S transcript sequence of AT-biased Genotypes #4–5, respectively. In contrast to the 95% (20/21 nt) and 85% (17/20 nt) similarities of primers 5.8S-F/R to the 5.8S transcript sequence of GC-biased Genotype #1, it is apparent that Li et al. (2013) designed the 5.8S-F/R primers based on the AT-biased 5.8S gene sequences, making it easier to amplify the 5.8S cDNA sequence of Genotype #5 than Genotype #1. The primers 5.8S-F/R showed variable similarity, ranging from 61.9%–100% and 31.8%–95.5%, respectively, among all 17 genotypes of *O. sinensis*, preferentially amplifying the most easily amplified cDNA sequence and consequently causing false-negative results for other 5.8S cDNA sequences.

Section III-5 and Table 4 show 100% homology between the 1894 unassembled metatranscriptome shotgun reads (100 bp) and the 5.8S gene sequence (156 bp) of GC-biased Genotype #1 *H. sinensis*, probably including a large number of nonrRNA transcripts among the 1894 shotgun reads. A BLAST search of the GenBank database showed that the 5.8S-F/R primers were 100% homologous to mRNA transcriptome assemblies GCQL01001191, GCQL01006777, GCQL01007818, and GCQL01013443 of Genotype #1 Strain L0106, which contains no rRNA transcripts and assembled metatranscriptome assemblies GAGW01003691, GAGW01007670, and GAGW01007671 of natural *C. sinensis* (Li et al. 2020a). Alignment analysis showed that metatranscriptome GAGW01003691 (2984 bp) was 99.8% homologous to a segment of mRNA transcriptome GCQL01006777 of *H. sinensis* Strain L0106. Metatranscriptomes GAGW01007670 (234 bp) and GAGW01007671 (1620 bp) are 100% identical to segments of GCQL01007818 and

831 GCQL01001191, respectively. Because mRNA transcriptome assembly GCQL00000000 contains no

832 rRNA transcripts, the extremely high homology between the metatranscriptome and transcriptome

833 sequences indicates that the 5.8S-F/R primers designed by Li et al. (2013) may nonspecifically

834 amplify nonrRNA components in the cDNA library derived from Group 3 clones (*cf*. Section III-3).

```
GT1 AB067721 1     TAGAGGAAGTAAAAGTCGTAACAAGGTCTCCGTTGGTGAACCAGCGGAGGGATCATTATC  60
GT4 AB067744 1     ------------------------------A--A---T----------T------       56
GT5 AB067740 1     --A---------------T-----------A--A---T----------T------       60

GT1 AB067721 61    GAGTCACCACTCCCAAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCC  120
GT4 AB067744 57    -----------C-------------A--C-------------------------------  116
GT5 AB067740 61    ----T--T----TT---------ATA--T--T-T-----------TA-----------T  120

GT1 AB067721 121   GGCGCCCCAGGGCCCGGACCAGGGCGCCCGCCGGAGGACCCCCAGACCCTCCTGTCGCAG  180
GT4 AB067744 117   -------T----------T-------------------TT----------A---T--    176
GT5 AB067740 121   A------T-A---------T-AA------------------TT---------A-TAT--  180

GT1 AB067721 181   TGGCATCTCTCAGTCAAGAAGCAAGCAAATGAATCAAAACTTTCAACAACGGATCTCTTG  240
GT4 AB067744 177   -A--------T---T--------------A---T-A-ACTTTTAATAACGGATCTCTTA  236
GT5 AB067740 181   -A-T------T---T--------------A---T---ACTTTTAACAACGGATCTCTTA  240
Primer 5.8S-F                                          ACTTTTAACAACGGATCTCTT

GT1 AB067721 241   GTTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCA  300
GT4 AB067744 237   ---T-A----TA--A---------------A-------------A-----T------T-  296
GT5 AB067740 241   -----A----TA--A--A------------A--------A-A---A-------T-      300

GT1 AB067721 301   GTGAACCATCGAATCTTTGAACGCACATTGCGCCCGCCAGCACTCTGGCGGGCATGCCTG  360
GT4 AB067744 297   --A---T-----------A----T-T-C-A-------T--------A------ATGCCTG  356
GT5 AB067740 301   --A---T-----------A------T---A-------T--------A------ATGCCTA  360
Primer 5.8S-R                                                         ATGCTTA

GT1 AB067721 361   TCCGAGCGTCATCTCAACCCTCGAGCCCCCCGCCTCGCGGCGGCGGGGCCCGGCCTTGGG  420
GT4 AB067744 357   TCCGAGCGTTACCTT------TA--------------------------------A--  416
GT5 AB067740 361   TCCGAGCGTTATCTT---------------------------------T-------A--  420
Primer 5.8S-R      TCCGAGCGTTATCTT

GT1 AB067721 421   GGTCACGGCCCCGCGCCGCCCCCTAAACGCAGTGGCGACCCCGCCGCGGCTCCCCTGCGC  480
GT4 AB067744 417   -----------A------------------A-T------------------A---      476
GT5 AB067740 421   ---T-----------------T------T---A--------------------A--T  480

GT1 AB067721 481   AGTAGCTCGCTGAGAACCTCGCACCGGGAGCGCGGAGGCGGTCACGCCGTGAAACCACCA  540
GT4 AB067744 477   -A-----TA--A------------TA-----------------------A------T-  536
GT5 AB067740 481   G-------A-C--------------A-----------------------A------T-  540

GT1 AB067721 541   CACCCTCCAGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACTTAAGCATATCAATAA  600
GT4 AB067744 537   T------TG---A---------T----------------A------------------  596
GT5 AB067740 541   -G--T-TT----A---------T----------------A------------T-----  600

GT1 AB067721 601   GCGGAGGAAAAGAAACCAACAGGGATTGCCCCAGTAACGGCGAGTGAAGCGGCAACAGCT  660
GT4 AB067744 597   --------                                                     604
GT5 AB067740 601   ----------------T--T---------TT------------A------T--T----  660
```

Figure 6. ITS sequence alignment of AB067721 of GC-biased Genotype #1, AB067744 and AB067740 of

AT-biased Genotypes #4–5 of O. sinensis and the 5.8S-F/R primers.

"GT1", "GT4" and "GT5" represent Genotypes #1 and #4–5 of O. sinensis, respectively. The sequences in

blue are 18S and 28S genes (5' and 3' end regions, respectively). Those in pink represent sequences of 5.8S

genes. Those in black are ITS1 between the 18S and 5.8S genes and ITS2 between the 5.8S and 28S genes. The

sequences underlined in green are for primers 5.8S-F/R, while primer 5.8S-R was converted to the sense chain

sequence. Hyphens indicate identical bases.

Li et al. (2013) did not disclose the sequences of 5.8S cDNA that they identified. Instead, they

reported that "Of these clones, 28 showed identical sequences (104 bp excluding the primers) to the
most frequent sequence in group A (JQ900148). The remaining 14 sequences … had single
substitutions at various positions", where group A = Genotype #1 *H. sinensis*. GenBank annotates the
location of the 5.8S gene of JQ900148 as nucleotides 192→346, covering 155 nt (not the multiple of
three), which is not in accord with the open reading frame of the gene.

Alternatively, the 5.8S genes are annotated with nucleotide positions of 218→373 in AB067721
of Genotype #1 and AB067740 of Genotype #5, and 214→369 in AB067744 of Genotype #4,
covering 156 nt and 52 codons in the 3 cases (Zhu & Li 2017; Li et al. 2020a, 2022e). Excluding the
5.8S-F/R primer regions, the remaining sequence of the 5.8S gene consists of 114 bp, with 10
additional nucleotides (mismatches) beyond the 104 bp of all cDNA clones reported by Li et al. (2013).
Adding "single substitutions at various positions" in 14 other clones reported by Li et al. (2013), the
42 clones presumably contained either 10 or 11 mismatches (8.8–9.6%) in the 114-bp segment of the
5.8S gene cDNA, corresponding to 90.4%–91.2% sequence similarity to the 5.8S gene sequence of
Genotype #1 of *O. sinensis*. Thus, attributing such high dissimilarity rates to "mismatches during
reverse transcription or PCR amplification" may not provide a convincing explanation, giving rise to
the question of whether the identified "5.8S cDNA" with a large number of mismatches was truly
derived from the 5.8S gene of *O. sinensis* Genotype #1, as assumed by Li et al. (2013, 2019, 2020b).
Considering the intensive epigenetic methylation of cytosine residues in a 5S RNA gene that is
associated with the transcriptional silencing of GC-biased genes, as reported by Selker and his
colleague (1985, 1990, 1997, 2002) and Gladyshev (2017), it may become legitimate to doubt
whether the 5.8S cDNA sequences that were identified by Li et al. (2013) truly belong to GC-biased
Genotype #1; this genotype contains more cytosine residues and is more susceptible to epigenetic
methylation attack and subsequent translational silencing than AT-biased Genotype #5. Accordingly,
it is reasonable to doubt the "ITS pseudogene" hypothesis, which postulates that the 5.8S genes of
AT-biased genotypes are permanently dysfunctional, rather than GC-biased Genotype #1.

Prior to declaring that the 5.8S genes of all AT-biased *O. sinensis* genotypes are nonfunctional
pseudogenes co-occurring with a functional 5.8S gene in the genome of GC-biased Genotype #1,
several other concerns should be addressed.

(1) Li et al. (2013) did not detect the ITS sequences of Genotypes #2–4 and #6–17 of *O. sinensis* in
the genomic DNA pool of ascosporic cultures, indicating that inferring that the 5.8S genes of
these genotypes are dysfunctional "ITS pseudogene" components of the genome of Genotype #1

H. sinensis is an apparent overinterpretation.

(2) The sequences of Genotypes #2–17 are absent in the genome assemblies ANOV00000000,

JAAVMX000000000, LKHE00000000, LWBQ00000000 and NGJJ00000000 of *H. sinensis*

Strains Co18, IOZ07, 1229, ZJB12195 and CC1406-2031229, respectively (Hu et al. 2013; Li et

al. 2016a; Jin et al. 2020; Liu et al. 2020; Shu et al. 2020), indicating genomic independence of

the genotypes of *O. sinensis*.

(3) Selker and his colleague (1985, 1990, 1997, 2002) and Gladyshev (2017) have repeatedly

reported epigenetic methylation of nearly all cytosine residues in 5.8S RNA genes and

subsequent transcription silencing of GC-biased genes, while the 5.8S gene of GC-biased

Genotype #1 contains more cytosine residues and is more susceptible to epigenetic methylation

attack and subsequent translational silencing than 5.8S genes of AT-biased genotypes.

(4) The culture-dependent mycological approach used by Li et al. (2013) might have missed some

fungal species that are nonculturable or difficult to culture under the *in vitro* experimental settings

of the study. The culture-dependent strategy might have a significant impact on the transcription

of many fungal genes, with many genes being switched on or off nonnaturally, as evidenced by

the nonlinear reduction in the total number of transcriptomic unigenes and increases in the

average length but decreases in the GC content of transcripts during 3–9 days of liquid

fermentation of *H. sinensis* Strain L0106 (Liu et al. 2015). A much greater impact on differential

transcriptional activation and silencing of many genes may be expected after the prolonged 25-

day liquid incubation period adopted by Li et al. (2013).

(5) Three distinct types of secondary steric conformations of 5.8S rRNA were predicted for Groups

A–C (*i.e.*, Genotypes #1 and #4–5) of *O. sinensis* by Li et al. (2013). The possibility of producing

circular RNA through back splicing-exon circularization (Dong et al. 2014a) should be

considered during study design because these distinct steric structures may have a considerable

impact on reverse transcription PCR and cDNA sequencing. In addition, the ITS1-5.8S-ITS2

sequences of Genotypes #2 and #6 may adopt different types of steric conformations (Zhu & Wu

2015; Li et al. 2016b, 2022e; Zhu & Li 2017); thus, the design of other genotype-specific primers

and the combined use of other molecular techniques should be considered.

(6) Wei et al. (2016) reported the identification of a single teleomorph of AT-biased Genotype #4 of

O. sinensis in cultivated *C. sinensis* and a single teleomorph of GC-biased Genotype #1 in natural

C. sinensis (*cf*. Figure 6 of Wei et al. 2016). If AT-biased Genotype #4 represents a nonfunctional

ITS pseudogene, as believed by Li et al. (2013, 2017, 2019, 2020b), the single *O. sinensis*

teleomorph in cultivated *C. sinensis* might have disturbed teleomorphic functions, leading to

reproductive sterility and an abnormal lifecycle of cultivated *C. sinensis*.

(7) Li et al. (2022c, 2022e) reported the identification of teleomorphic Genotypes #4 and #15 of *O.*

sinensis of AT-biased Cluster-B in the stromal fertile portion densely covered with numerous

ascocarps but not in ascospores collected from the same pieces of natural *C. sinensis*. AT-biased

Genotypes #6 and #15 were found at high abundance in the stromal fertile portion prior to

ascospore ejection, and their abundance drastically declined after ascospore ejection, while

teleomorphic Genotype #5 maintained its high abundance in the stromal fertile portion before

and after ascospore ejection (Li et al. 2022a, 2022e).

(8) The 5.8S genes of multiple *O. sinensis* genotypes may be transcriptionally activated or silenced

in different developmental and maturational stages of natural *C. sinensis*. Significant changes in

the proteomic expression profiles of the stroma and caterpillar body of natural *C. sinensis* during

maturation constitute evidence of such dynamic alterations in epigenetic, transcriptional,

translational and posttranslational modifications of numerous genes (Dong et al. 2014b).

Apparently, the techniques used by Li et al. (2013) provided insufficient and nonconclusive

evidence to support their "ITS pseudogenes" hypothesis. The whole package of gene expression in

O. sinensis fungi and natural *C. sinensis* needs to be further explored before the 5.8S genes of AT-

biased genotypes of *O. sinensis* can be defined as permanently nonfunctional pseudogenes.

VI. Conclusion

This paper analyzes the data related to the multiple repetitive ITS copies in the genome of

Genotype #1 *H. sinensis* and multiple genotypes of *O. sinensis* fungi and discusses the "ITS

pseudogene … in a single genome" hypothesis for all AT-biased *O. sinensis* genotypes (Li et al. 2013,

2017, 2019, 2020b). The repetitive ITS copies in the genome of Genotype #1 contain multiple

scattered insertion, deletion and transversion point mutations and may not be generated through RIP

mutagenesis because RIP theoretically causes C-to-T and G-to-A transitions. The GC-biased

repetitive ITS copies are genetically and phylogenetically distinct from AT-biased *O. sinensis*

genotypes that contain multiple scattered transition alleles. The AT-biased genotypes are genomically

independent from the genome of Genotype #1 and might have been generated from a common ancestor through RIP or other mutagenic and evolutionary mechanisms in the long history of evolution and become independent *O. sinensis* fungi with independent genomes coexisting with GC-biased genotypes in natural *C. sinensis*. Li et al. (2013) provided incomplete and controversial information to support their "ITS pseudogene … in a single genome" hypothesis, which needs to be further validated.

V. Acknowledgment:

This research was supported by a grant from the Science and Technology Department of Qinghai Province, China, grant number 2021-SF-A4 "Study on key technologies of conservation of natural resource and industrial upgrading of *Cordyceps sinensis*", the major science and technology projects in Qinghai Province.

References

[1] Bushley, K.E., Li, Y., Wang, W.-J., Wan,g X.-L., Jiao, L., Spatafora, J.W., Yao, Y.-J., 2013. Isolation of the MAT1-1 mating type idiomorph and evidence for selfing in the Chinese medicinal fungus *Ophiocordyceps sinensis*. Fungal Biol. 117, 599–610. (PMID: 22181845 DOI: 10.1016/j.funbio.2013.06.001)

[2] Chen C-S, Hseu R-S, Huang C-T. 2011. Quality control of *Cordyceps sinensis* teleomorph, anamorph, and Its products. Chapter 12, *in* (Shoyama, Y., Ed.) *Quality Control of Herbal Medicines and Related Areas*. InTech, Rijeka, Croatia. (www.intechopen.com)

[3] Chen, Y.-Q., Hu, B., Xu, F., Zhang, W., Zhou, H., Qu, L.-H., 2004. Genetic variation of *Cordyceps sinensis*, a fruit-body-producing entomopathogenic species from different geographical regions in China. FEMS Microbiol. Lett. 230, 153–158. (PMID: 14734179 DOI: 10.1016/S0378-1097(03)00889-9)

[4] Dong, R., Che,n L.-L., Yang, L., 2014a. Circular RNA studies in the post-genomic era. Chin. J. Cell Biol. 36 (11), 1455–1459. (DOI: 10.11844/cjcb.2014.11.9001)

[5] Dong, Y.-Z., Zhang, L.-J., Wu, Z.-M., Gao, L., Yao, Y.-S., Tan, N.-Z., Wu, J.-Y., Ni, L.-Q., Zhu, J.-S., 2014b. Altered proteomic polymorphism in the caterpillar body and stroma of natural *Cordyceps sinensis* during maturation. PLoS ONE. 9 (10), e109083. (PMID: 25310818 doi:10.1371/journal.pone.0109083)

[6] Engh, I.B., 1999. Molecular phylogeny of the *Cordyceps-Tolypocladium* complex. Candidate scientific thesis, Department of Biology, University of Oslo, Oslo. 1999.

[7] Gao, L., Li, X.-H., Zhao, J.-Q., Lu, J.-H., Zhu, J.-S., 2011. Detection of multiple *Ophiocordyceps sinensis*
mutants in premature stroma of *Cordyceps sinensis* by MassARRAY SNP MALDI-TOF mass spectrum
genotyping. Beijing Da Xue Xue Bao. 43 (2), 259–266. (PMID: 21503123 doi: 10.3969/j.issn.1671-
167X.2011.02.019)

[8] Gao, L., Li, X.-H., Zhao, J.-Q., Lu, J.-H., Zhao, J.-G., Zhu, J.-S., 2012. Maturation of *Cordyceps sinensis*
associates with alterations of fungal expressions of multiple *Ophiocordyceps sinensis* mutants with
transition and transversion point mutations in stroma of *Cordyceps sinensis*. Beijing Da Xue Xue Bao. 44
(3), 454–463. (PMID: 22692321 doi: 10.3969/j.issn.1671-167X.2012.03.025)

[9] Gladyshev E. 2017. Repeat-Induced Point Mutation (RIP) and Other Genome Defense Mechanisms in
Fungi. Microbiol. Spectr. 5(4): . (PMID: 28721856 doi: 10.1128/microbiolspec.FUNK-0042-2017)

[10] Hawksworth DL, Crous PW, Redhead SA, Reynolds DR, Samson RA, Seifert KA, Taylor JW, Wingfield
MJ, Abaci O, Aime C, Asan A, Bai F-Y, de Beer ZW, Begerow D, Berikten D, Boekhout T, K Buchanan
P, Burgess T, Buzina W, Cai L, Cannon PF, Crane JL, Damm U, Daniel H-M, van Diepeningen AD,
Druzhinina I, Dyer PS, Eberhardt U, Fell JW, Frisvad JC, Geiser DM, Geml J, Glienke C, Gräfenhan T,
Groenewald JZ, Groenewald M, de Gruyter J, Guého-Kellermann E, Guo L-D, Hibbett DS, Hong S-B,
de Hoog GS, Houbraken J, Huhndorf SM, Hyde K D, Ismail A, Johnston PR, Kadaifciler DG, Kirk PM,
Kõljalg U, Kurtzman CP, Lagneau P-E, Lévesque CA, Liu X, Lombard L, Meyer W, Miller A, Minter
DW, Najafzadeh MJ, Norvell L, Ozerskaya SM, Oziç R, Pennycook SR, Peterson SW, Pettersson OV,
Quaedvlieg W, Robert VA, Ruibal C, Schnürer J, Schroers H-J, Shivas R, Slippers B, Spierenburg H,
Takashima M, Taşkın E, Thines M, Thrane U, Uztan AH,van Raak M, Varga J, Vasco A, Verkley G,
Videira SIR, de Vries RP, Weir BS, Yilmaz N, Yurkov A, Zhang N. The Amsterdam declaration on
fungal nomenclature. IMA Fungus 2011; 2: 105−112 (PMID: 22679594 DOI:
10.5598/imafungus.2011.02.01.14).

[11] Hu, X., Zhang, Y., Xiao, G., Zheng, P., Xia, Y., Zhang, X., St Leger, R.J., Liu, X., Wang, C., 2013. Genome
survey uncovers the secrets of sex and lifestyle in caterpillar fungus. Chin. Sci. Bull. 58, 2846–2854. (doi:
10.1007/s11434-013-5929-5)

[12] Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogeny. Bioinformat. 17:
754−755. (PMID: 11524383 DOI: 10.1093/bioinformatics/17.8.754)

[13] Jin L-Q, Xu Z-W, Zhang B, Yi M, Weng C-Y, Lin S, Wu H, Qin X-T, Xu F, Teng Y, Yuan S-J, Liu Z-Q,
Zheng Y-G. 2020. Genome sequencing and analysis of fungus *Hirsutella sinensis* isolated from
Ophiocordyceps sinensis. AMB Expr. 10:105. (PMID: 32494871 DOI: 10.1186/S13568-020-01039-x)

[14] Kinjo, N., Zang, M., 2001. Morphological and phylogenetic studies on *Cordyceps sinensis* distributed in
southwestern China. Mycoscience. 42, 567–574. (https://doi.org/10.1007/BF02460956)

[15] Li C-L, Tan N-Z, Berger JL, Ferguson SB, Zhang Y, Prolla TA, Zhu J-S. 2011. The Combined use of

ageLOC whole-gene expression profiling technology and mouse lifespan test in anti-aging herbal product

study. Proc. 2011 New TCM Prod. Innovat. Indust. Devel. Summit, Hangzhou, China, 2011, pp. 443–448.

(https://xueshu.baidu.com/usercenter/paper/show?paperid=08341c17fa58c8f85584b92572b90f75&site=xueshu_se)

[16] Li X-Z, Li Y-L, Zhu J-S. *Differential expression of mating-type genes in Hirsutella sinensis and natural*

Cordyceps sinensis. Proc. 2021 Intl. Conf. Info. Technol. Biomed. Engineering, 2021, pp. 318−333.

(https://conferences.computer.org/icitbepub/pdfs/ICITBE2021-6JNtJiThlp10s9N5Sodvuy/009900a318/009900a318.pdf)

[17] Li X-Z, Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. 2020a. Further discussion with Li et al. (2013, 2019)

regarding the "ITS pseudogene hypothesis" for *Ophiocordyceps sinensis*. Mol. Phylogenet. Evol. 146:

106728. (PMID: 31904509 DOI: 10.1016/j.ympev.2019.106728)

[18] Li X-Z, Li Y-L, Yao Y-S, Xie W-D, Zhu J-S, 2022a. Molecular identification of *Ophiocordyceps sinensis*

genotypes and the indiscriminate use of the Latin name for the multiple genotypes and the natural insect-

fungi complex. Am J Biomed Sci. 14(3): 115–135. (doi:10.5099/aj220300115)

[19] Li X-Z, Li Y-L, Yao Y-S, Zhu J-S. 2022b. Differential expressions of the *Hirsutella sinensis* genes and

intraspecific genetic variations among *H. sinensis* strains. Biochem Mol Biol J. 8(2): 58. (DOI:

10.36648/2471-8084.22.8.58)

[20] Li, Y., Hsiang, T., Yang, R.-H., Hu, X.-D., Wang, K., Wang, W.-J., Wang, X.-L., Jiao, L., Yao, Y.-J., 2016a.

Comparison of different sequencing and assembly strategies for a repeat-rich fungal genome,

Ophiocordyceps   *sinensis*.   J.   Microbiol.   Methods.   128,   1–6.

(http://dx.doi.org/10.1016/j.mimet.2016.06.025)

[21] Li, Y., Jiang, L., Wang, K., Wu, H.-J., Yang, R.-H., Yan, Y.-J., Bushley, K.E., Hawksworth, D.L., Wu, Z.-

J., Yao, Y.-J., 2020b. RIP mutated ITS genes in populations of *Ophiocordyceps sinensis* and their

implications for molecular systematics. IMA Fungus 11, 18. (PMID: 27343682 DOI:

10.1016/j.mimet.2016.06.025)

[22] Li, Y., Jiao, L., Yao, Y.-J., 2013. Non-concerted ITS evolution in fungi, as revealed from the important

medicinal   fungus   *Ophiocordyceps*   *sinensis*.   Mol.   Phylogenetics   Evol.   68,   373–379.

(http://dx.doi.org/10.1016/j.ympev.2013.04.010)

[23] Li, Y., Wang, Y.-H., Wang, K., Yang, R.-H., Jiao, L., Yao, Y.-J., 2019. Response to "The multiple

genotypes of *Ophiocordyceps sinensis* and the ITS pseudogene hypothesis". Mol. Phylogenetics Evol.

139, 106522. (PMID: 23618625 DOI: 10.1016/j.ympev.2013.04.010)

[24] Li, Y., Yang, R.-H., Jiang, L., Hu, X.-D., Wu, Z.-J., Yao, Y.-J., 2017. rRNA pseudogenes in filamentous

ascomycetes as revealed by genome data. G3 Genes|Genomes|Genetics. 7, 2695–2703. (PMID: 28637809

doi: 10.1534/g3.117.044016)

[25] Li Y-L, Gao L, Yao Y-S, Li X-Z, Wu Z-M, Lou Z-Q, Xie W-D, Wu J-Y, Zhu J-S. 2022c. Altered GC- and

AT-biased genotypes of *Ophiocordyceps sinensis* in the stromal fertile portions and ascospores of natural

Cordyceps sinensis. PLOS ONE (submitted). BIORXIV Preprint 2022-11-12.

(https://www.biorxiv.org/content/10.1101/2022.10.07.511361v1.full.pdf)

[26] Li Y-L, Li X-Z, Yao Y-S, Zhu J-S. 2022d. Differential cooccurrence of multiple genotypes of

Ophiocordyceps sinensis in the stromata, stromal fertile portion (ascocarps) and ascospores of natural

Cordyceps sinensis. PLOS ONE (in revision). BIORXIV Preprint 2022-6-20.

(https://www.biorxiv.org/content/10.1101/2022.06.20.496766v2.full.pdf)

[27] Li Y-L, Li X-Z, Zhu J-S. 2022e. *Cordyceps sinensis* Fungal Heterokaryons. Chemistry Industry Press Co.,

Beijing, China. (ISBN: 978-7-122-40189-2   http://www.cip.com.cn/Book/Index/48939)

[28] Li, Y.-L., Yao, Y.-S., Xie, W.-D., Zhu, J.-S., 2016b. The molecular heterogeneity of natural *Cordyceps*

sinensis with multiple *Ophiocordyceps sinensis* fungi challenges the anamorph-teleomorph connection

hypotheses. Am. J. Biomed. Sci. 8, 123–159. (doi: 10.5099/aj100300217)

[29] Li, Y.-L., Yao, Y.-S., Zhang, Z.-H., Xu, H.-F., Liu, X., Ma, S.-L., Wu, Z.-M., Zhu, J.-S., 2016c. Synergy

of fungal complexes isolated from the intestines of *Hepialus lagii* larvae in increasing infection potency.

J. Fungal Res. 14, 96–112. (DOI: 10.13341/j.jfr.2014.1067)

[30] Liu J, Guo L-N, Li Z-W, Zhou Z, Li Z, Li Q, Bo X-C, Wang S-Q, Wang J-L, Ma S-C, Zheng J, Yang Y.

2020. Genomic analyses reveal evolutionary and geologic context for the plateau fungus *Ophiocordyceps*

sinensis. Clin. Med. 15:107–119. (PMID: 33042212 DOI: 10.1186/s13020-020-00365-3)

[31] Liu, Z.-Q., Lin, S., Baker, P., Wu, L.-F., Wang, X.-R., Wu, H., Xu, F., Wang, H.-Y., Brathwaite, M., Zheng,

Y.-G., 2015. Transcriptome sequencing and analysis of the entomopathogenic fungus *Hirsutella sinensis*

isolated from *Ophiocordyceps sinensis*. BMC Genome. 16, 106. (PMID: 25765329 DOI 10.1186/s12864-

015-1269-y)

[32] Liu Z-Y, Yao Y-J, Liang Z-Q, Liu A-Y, Pegler DN, Chase MW. 2001. Molecular evidence for the

anamorph-teleomorph connection in *Cordyceps sinensis*. Mycology Research 105: 827−832.

(https://doi.org/10.1017/S095375620100377X)

[33] Mao, X.-M., Zhao, S.-M., Cao, L., Yan, X., Han, R.-X., 2013. The morphology observation of

Ophiocordyceps sinensis from different origins. J. Environ. Entomol. 35, 343–353. (doi:

10.3969/j.issn.1674-0858.2013.03.11)

[34] Ren Y, Wan D-G, Lu X-M, Guo J-L. 2013. The study of scientific name discussion for TCM *Cordyceps*.

Lishizhen Medicine and Materia Medica Research 24: 2211−2212 (Doi: 10.3969/j.issn.1008-

0805.2013.09.075).

[35] Selker EU. 1990. Premeiotic instability of repeated sequences in Neurospora crassa. Annu Rev Genet.

24:579–613. (PMID: 2150906 DOI: 10.1146/annurev.ge.24.120190.003051)

[36] Selker EU. 1997. Epigenetic phenomena in filamentous fungi: useful paradigms or repeat-induced

confusion? Trends Genet. 13:296–301. (PMID: 9260514 DOI: 10.1016/S0168-9525(97)01201-8)

[37] Selker EU. 2002. Repeat-induced gene silencing in fungi. Adv Genet. 46:439–450. (PMID: 11931234

DOI: 10.1016/S0065-2660(02)46016-6)

[38] Selker EU, Stevens JN. 1985. DNA methylation at asymmetric sites is associated with numerous

transition mutations. Proc. Natl. Acad. Sci. USA. 82:8114–8118. (PMID: 2415981 DOI:

10.1073/pnas.82.23.8114)

[39] Shu R-H, Zhang J-H, Meng Q, Zhang H, Zhou G-L, Li M-M, Wu P-P, Zhao Y-N, Chen C, Qin Q-L. 2020.

A new high-quality draft genome assembly of the Chinese cordyceps *Ophiocordyceps sinensis*. Genome

Biol. Evol. 12(7): 1074−1079. (PMID: 32579174 DOI: 10.1093/gbe/evaa112)

[40] Stensrud, Ø., Hywel-Jones, N,L,, Schumacher, T., 2005. Towards a phylogenetic classification of

Cordyceps: ITS nrDNA sequence data confirm divergent lineages and paraphyly. Mycol Res. 109 (1), 41–

56. (PMID: 15736862 DOI: 10.1017/S095375620400139X)

[41] Stensrud, Ø., Schumacher, T., Shalchian-Tabrizi, K., Svegården, I.B., Kauserud, H., 2007. Accelerated

nrDNA evolution and profound AT bias in the medicinal fungus *Cordyceps sinensis*. Mycol. Res. 111,

409–415. (PMID: 17509850 doi:10.1016/j.mycres.2007.01.015)

[42] Sung, G.-H., Hywel-Jones, N,L,, Sung, J.-M., Luangsa-ard, J.J., Shrestha, B., Spatafora, J.W., 2007.

Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. Stud. Mycol. 57, 5–59. (PMID:

18490993 DOI: 10.3114/sim.2007.57.01)

[43] Tan N-Z, Berger JL, Zhang Y, Prolla TA, Weindruch R, Zhao C-S, Bartlett M, Zhu J-S. 2011. The lifespan-

prolonging effect of *Cordyceps sinensis* Cs-4 in normal mice and its molecular mechanisms. FASEB J.

25(1):599.1. (http://www.fasebj.org/cgi/content/meeting_abstract/25/1_MeetingAbstracts/599.1)

[44] Wei, J.-C., Wei, X.-L., Zheng, W., Guo, W., Liu, R., 2016. Species identification and component detection

of *Ophiocordyceps sinensis* cultivated by modern industry. Mycosystema. 35, 404–410. (doi:

10.13346/j.mycosystema.160045)

[45] Wei X-L, Yin X-C, Guo Y-L, Shen N-Y, Wei, J.-C. 2006. Analyses of molecular systematics on *Cordyceps*

sinensis and its related taxa. Mycosystema. 25(2): 192–202.

[46] Xia, E.-H., Yang, D.-R., Jiang, J.-J., Zhang, Q.-J., Liu, Y., Liu, Y.-L., Zhang, Y., Zhang, H.-B., Shi, C.,

Tong, Y., Kim, C., Chen, H., Peng, Y.-Q., Yu, Y., Zhang, W., Eichler, E.-E., Gao, L.-Z., 2017. The

caterpillar fungus, *Ophiocordyceps sinensis*, genome provides insights into highland adaptation of fungal

pathogenicity. Sci Rep, 7, 1806. (PMID: 28496210 doi: 10.1038/s41598-017-01869-z)

(www.plantkingdomgdb.com/Ophiocordyceps_sinensis/data/cds/Ophiocordyceps_sinensis_CDS.fas for

the cDNA file)

[47] Xia F, Liu Y, Shen G-R, Guo L-X, Zhou X-W. 2015. Investigation and analysis of microbiological

communities in natural *Ophiocordyceps sinensis*. Can J Microbiol, 61: 104−111. (PMID: 25578897

http://dx.doi.org/10.1139/cjm-2014-0610)

[48] Xiang, L., Li, Y., Zhu, Y.-J., Luo, H.->., Li, C.-F., Xu, X.-L., Sun, C., Song, J.-Y., Shi, L.-C., He, L., Sun, W., Chen, S.-L., 2014. Transcriptome analysis of the *Ophiocordyceps sinensis* fruiting body reveals putative genes involved in fruiting body development and cordycepin biosynthesis. Genomics 103, 154–159. (PMID: 24440419   http://dx.doi.org/10.1016/j.ygeno.2014.01.002)

[49] Xiao, Y-Y., Chen, C., Dong, J.-f., Li, C.-R., Fan, M.-Z. 2011. Morphological observation of ascospores of *Ophiocordyceps sinensis* and its anamorph in growth process. J Anhui Agricult Univ. 38(4): 587–591. (http://www.cnki.net/kcms/detail/34.1162.S.20110624.1628.030.html)

[50] Xiao, W., Yang, J.-L., Zhu, P., Cheng, K., He, H., Zhu, H., Wang, Q., 2009. Non-support of species complex hypothesis of *Cordyceps sinensis* by targeted rDNA-ITS sequence analysis. Mycosystema. 28, 724–730. (DOI: 10.13346/j.mycosystema.2009.05.028)

[51] Yang, J.-L., Xiao, W., He, H., Zhu, H., Wang, S., Cheng, K., Zhu, P., 2008. Molecular phylogenetic analysis of *Paecilomyces hepiali* and *Cordyceps sinensis*. Acta Pharm. Sin. 43, 421–426. (PMID: 18664207)

[52] Yao, Y.-S., Zhou, Y.-J., Gao, L., Lu, J.-H., Wu, Z.-M., Zhu, J.-S., 2011. Dynamic alterations of the differential fungal expressions of *Ophiocordyceps sinensis* and its mutant genotypes in stroma and caterpillar during maturation of natural *Cordyceps sinensis*. J. Fungal Res. 9, 37–49,53. (PMID: 22692321 Doi: 10.3969/j.issn.1672-3538.2011.01.007)

[53] Yao, Y.-S., Zhu, J.-S., 2016. Indiscriminative use of the Latin name for natural *Cordyceps sinensis* insect-fungi complex and multiple *Ophiocordyceps sinensis* fungi. Chin. J. Chin. Mater. Med. 41 (7), 1316–1366. (PMID: 28879757 DOI: 10.4268/cjcmm20160733)

[54] Zhang S, Zhang Y-J. 2015. Molecular evolution of three protein-coding genes in the Chinese caterpillar fungus *Ophiocordyceps sinensis*. Microbiology China 42(8): 1549−1560. (DOI: 10.13344/j.microbiol.china.150279)

[55] Zhang S, Zhang Y-J, Shrestha B, Xu J-P, Wang C-S, Liu X-Z. *Ophiocordyceps sinensis* and *Cordyceps militaris*: research advances, issues and perspectives. Mycosystema 2013; 32: 577−597 (DOI: 10.3969/j.issn.1673-5110.2014.18.061).

[56] Zhang S-W, Cem K, Liu Y, Zhou X-W, Wang C-S. Metatranscriptomics analysis of the fruiting caterpillar fungus collected from the Qinghai-Tibetan plateau. Scientia Sinica Vitae 2018; 48(5): 562. (doi: 10.1360/N052017-00253)

[57] Zhang Y-J, Li E-W, Wang C-S, Li Y-L, Liu X-Z. *Ophiocordyceps sinensis*, the flagship fungus of China: terminology, life strategy and ecology. Mycology 2012; 3: 2−10. (DOI: 10.1080/21501203.2011.654354)

[58] Zhang, Y.-J., Sun, B.-D., Zhang, S., Wang, M., Liu, X.-Z., Gong, W.-F., 2010. Mycobiotal investigation of natural *Ophiocordyceps sinensis* based on culture-dependent investigation. Mycosystema 29, 518−527.

[59] Zhang, Y.-J., Xu, L., Zhang, S., Liu, X.-Z., An, Z., Wang, M., Guo, Y., 2009. Genetic diversity of

Ophiocordyceps sinensis, a medicinal fungus endemic to the Tibetan Plateau: implications for its
evolution and conservation. BMC Evolut. Biol. 9, 290. (PMID: 20003548 DOI: 10.1186/1471-2148-9-
290)

[60] Zhao J, Wang N, Chen Y-Q, Li T-H, Qu L-H. 1999. Molecular identification for the asexual stage of
Cordyceps sinensis. Acta Scientiarum Naturalium Universitatis Sunyatseni 38: 122−123.

[61] Zhu, J.-S., Gao, L., Li, X.-H., Yao, Y.-S., Zhao, J.-Q., 2010. Maturational alteration of oppositely
orientated rDNA and differential proliferation of GC- and AT-biased genotypes of *Ophiocordyceps*
sinensis and *Paecilomyces hepiali* in natural *Cordyceps sinensis*. Am. J. Biomed. Sci. 2, 217–238. (doi:
10.5099/aj100300217)

[62] Zhu J-S, Halpern GM, Jones K. 1998a. The scientific rediscovery of a precious ancient Chinese herbal
regimen: *Cordyceps sinensis*: Part I. J. Altern. Complem. Med. 4(3): 289–303. (PMID: 9764768 DOI:
10.1089/acm.1998.4.3-289)

[63] Zhu J-S, Halpern GM, Jones K. 1998b. The scientific rediscovery of an ancient Chinese herbal medicine:
Cordyceps sinensis: Part II. J. Altern. Complem. Med. 4(4): 429–457. (PMID: 9884180 DOI:
10.1089/acm.1998.4.429)

[64] Zhu, J.-S., Li, Y.-L., 2017. A Precious Transitional Chinese Medicine, *Cordyceps sinensis*: Multiple
Heterogeneous *Ophiocordyceps sinensis* in the Insect-Fungi Complex. Lambert Academic Publishing,
Germany. (ISBN-10: 9783330046689, ISBN-13: 978-3330046689)

[65] Zhu, J.-S., Li, Y.-L., Yao, Y.-S., Xie, W.-D., 2018. Molecular evidence not supporting the use of *Hirsutella*
sinensis strain EFCC 7287 as the taxonomic standard for multiple *Ophiocordyceps sinensis* fungi. Int. J.
Mol. Biol. 3, 114–115. (DOI: 10.15406/ijmboa.2018.03.00062)

[66] Zhu, J.-S., Li, Y.-L., Yao, Y.-S., Xie, W.-D., 2019. The multiple genotypes of *Ophiocordyceps sinensis*
and the ITS pseudogene hypothesis. Mol. Phylogenetics Evol. 139, 106322. (PMID: 30394296 DOI:
10.1016/j.ympev.2018.10.034)

[67] Zhu, J.-S., Wu, J.-Y., 2015. Genetic heterogeneity of natural *Cordyceps sinensis* with co-existence of
multiple fungi. Chin. J. Cell Biol. 37 (2), 284–298. (DOI: 10.11844/cjcb.2015.02.0284)

[68] Zhu, J.-S., Zhao, J.-G., Gao, L., Li, X.-H., Yao, Y.-S., Zhao, J.-Q., 2012. Dynamically altered expressions
of at least 6 *Ophiocordyceps sinensis* mutants in the stroma of *Cordyceps sinensis*. J. Fungal Res. 10, 100–
112. (DOI: 10.13341/j.jfr.2012.02.007)

1165

Multiple GC-biased repetitive ITS copies in the *Hirsutella sinensis* genome are not generated by RIP mutagenesis involving transition point mutations

Xiu-Zhang Li[a], Yu-Ling Li[a], Ya-Nan Wang[b], Jia-Shi Zhu[a,c,*]

Table 1. *GC contents of the repetitive ITS1-5.8S-ITS2 sequences from the genome assemblies of 2 H. sinensis strains and percent similarities of the repetitive ITS sequences to the sequences of GC-biased Genotype #1 (AB067721) and AT-biased Genotypes #4–6 and #15–17 of O. sinensis.*

Genome segment	Sequence range & direction	GC content	Percent similarity of the repetitive sequence	
			vs. Genotype #1 AB067721	*vs.* AT-biased Genotypes #4–6 and #15–17
JAAVMX010000002	18688917→18689407	64.8%	100%	86.7% – 89.9%
	18702095→18702586	64.5%	97.4%	82.8% – 88.0%
JAAVMX010000008	13823→14313	64.8%	100%	85.5% – 89.9%
	1199→1687	64.8%	99.2%	85.2% – 89.3%
JAAVMX010000017	9147←9637	64.8%	100%	85.5% – 89.9%
	21791←22281	64.8%	100%	85.5% – 89.9%
	34435←34925	64.8%	100%	85.5% – 89.9%
	47079←47569	64.8%	100%	85.5% – 89.9%
JAAVMX010000018	13381→13871	64.8%	100%	85.5% – 89.9%
	26076→26566	64.8%	100%	85.5% – 89.9%
	38771→39261	64.8%	100%	85.5% – 89.9%
	51467→51958	64.6%	98.8%	85.5% – 88.8%
	700→1186	64.5%	97.0%	82.6% – 87.6%
JAAVMX010000019	19404→19894	64.8%	100%	85.5% – 89.9%
	32048→32538	64.8%	100%	85.5% – 89.9%
	6233→6733	65.3%	94.5%	81.3% – 85.5%
	44729→45251	63.3%	90.8%	80.1% – 84.6%
NGJJ01000573	13133←13623	64.8%	99.4%	85.2% – 89.6%
	14624←15113	64.7%	99.4%	84.8% – 89.6%
	1←260 (ITS1 and partial 5.8S)	56.5%	98.5%	83.5% – 88.3%
NGJJ01000582	10703←11192	64.7%	99.8%	85.5% – 89.8%
	24068←24190 (only partial ITS2)	73.2%	98.1%	87.7% – 89.1%
NGJJ01000796	1514→2004	64.8%	100%	85.5% – 89.9%
NGJJ01000798	1177→1665	64.6%	99.6%	84.8% – 89.6%
NGJJ01000799	3295→3785	64.8%	100%	85.5% – 89.9%
	16666→17156	64.8%	100%	85.5% – 89.9%

Table 2. *Percent similarities between nrDNA ITS1-5.8S-ITS2 sequences of 5 genome assemblies of Genotype #1 H. sinensis and 17 O. sinensis genotype sequences. (Modified from Table 1 of Li et al. 2022d).*

GT	GenBank Accession #	% similarity between the *H. sinensis* genomic ITS segments and 17 *O. sinensis* genotypes				
		ANOV01021709 838→1504	LKHE01000582 2074→2740	LWBQ01000008 991739→992405	JAAVMX010000017 9029←9695	NGJJ01000799 3237→3903
#1	AB067721	99.7%	100%	99.6%	100%	100%
#2	MG770309	94.4%	94.7%	94.7%	94.7%	94.7%
#3	HM595984	95.9%	96.1%	95.6%	95.3%	95.3%
#7	AJ488254	94.7%	95.0%	95.3%	95.0%	95.0%
#8	GU246286	89.5%	89.5%	89.2%	89.5%	89.5%
#9	GU246288	95.0%	95.2%	94.6%	95.2%	95.2%
#10	GU246287	83.0%	83.2%	83.0%	83.2%	83.2%
#11	JQ695935	78.9%	79.1%	78.5%	79.1%	79.1%
#12	GU246296	94.3%	94.5%	93.9%	94.5%	94.5%
#13	KT339190	87.5%	87.5%	87.2%	87.5%	87.5%
#14	KT339178	89.3%	89.6%	89.1%	89.6%	89.6%
#4	AB067744	89.1%	89.4%	89.3%	89.4%	89.4%
#5	AB067740	86.3%	86.7%	86.2%	86.7%	86.7%
#6	KJ720572	85.5%	85.5%	85.5%	85.5%	85.5%
#15	KT232017	89.6%	89.9%	89.4%	89.9%	89.9%
#16	KT232019	87.4%	87.7%	87.6%	87.7%	87.7%
#17	KT232010	87.7%	88.1%	87.5%	88.1%	88.1%

Note: "GT" represents genotype. GC-biased genotypes are listed in blue in the upper panel. AT-biased genotypes are listed in red in the lower panel.

Table 3. *Types of mutations in the variable and slightly less variable genomic repetitive ITS sequences compared with GC-biased Genotype #1 (AB067721).*

	Sequence range & direction	% similarity	Number of mutant alleles			Ins./Del./Transv. *vs*. Transit.
			Ins./Del.	Transv.	Transit.	
JAAVMX010000002	18702095→18702586	97.4%	13	0	0	13 : 0
JAAVMX010000018	700→1186	97.0%	10	5	0	15 : 0
JAAVMX010000019	6233→6733	94.5%	22	4	1	26 : 1
	44729→45251	90.8%	32	11	5	43 : 5

Note: "Ins./Del." indicates insertion/deletion point mutations. "Transv." denotes Transversion point mutations. "Transit." refers to transition point mutations.

```
GT#1  AB067721  59        TCGAGTCACCACTCCC AAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCCGGCG CCCCAGGGCCCG 136
GT#3  FJ654148  1         CT----TG-------- ---A----C-T-------------------------------------- ------------ 83
JAAVMX010000019 19404     ---------------- -------------------------------------------------- ------------ 19481
JAAVMX010000019 32048     ---------------- -------------------------------------------------- ------------ 32125
JAAVMX010000002 18702095  ---------------- --------------------------------------T----------- ------------ 18702172
JAAVMX010000018 700       --------- ------ --------------------------------T----------------T 776
JAAVMX010000019 6233      ----------------C -------------------------------------------- ------------ 6311
JAAVMX010000019 44729     ---------------- -------------------------------------------------- ------------ 44806
GT#4  AB067744  55        ------T-------T --------------T-T--------------------------- --T----A----- 132
GT#5  AB067740  59        ------T--T----TT -----ATA--T--T-T------------TA------------TA---- --T--A------ 136
GT#6  KJ720572  11              -------A---T-T-----------TA-----------TA--- --TT-A------ 70
GT#15 KT232017  37        ------T-------T --------------T-T--------------------------- --T------- 114
GT#16 KT232019  37        ------T--T----T --------------T-T-T----------TA------------TA--- --T-A------ 114
GT#17 KT232010  37        ------T--T----T --------------T-T-T----------TA------------TA--- --T-A------ 114

GT#1  AB067721  137       GACCAGG GCGCCC GCCGGAGGACCCCCAGAC CCTCCTGTCGCA GTGGCATCT CTCAGTCAAGAA GCAAGCAAAT 210
GT#3  FJ654148  84        ------- ------ ------------------ T---------- -----G--- ------------ ---------- 157
JAAVMX010000019 19482     ------- ------ ------------------ ------------ --------- ------------ ---------- 19555
JAAVMX010000019 32126     ------- ------ ------------------ ------------ --------- ------------ ---------- 32199
JAAVMX010000002 18702173  ------- ------ ----A------------- ------------ --------A -----------  ---------- 18702247
JAAVMX010000018 777       ------- ------ ------------------ ------------ --------- ---C-------- ---------- 850
JAAVMX010000019 6312      ------- ------ ------T----------C ------------ --------C- --------G----- ---------- 6388
JAAVMX010000019 44807     -------C ------ ------------------ ------------ --------- ------------ ---------- 44881
GT#4  AB067744  133       ---T--- ------ ----------TT---- ------A---T- --A------ --T---T---- ---------- 206
GT#5  AB067740  137       ---T-AA ------ ----------TT---- ------A-TAT- --A-T---- --T---T---- ---------- 213
GT#6  KJ720572  71        ---T--- ------ ----------TT---- ------A--T- --A-T---- --T---T---- ---------- 182
GT#15 KT232017  115       ---T--- ------ ----------TT---- ------A---A- --A------ --T---T---- ---------- 188
GT#16 KT232019  115       ---T-AA ------ ----------TT---- ------A---- --A-T---- --T---T---- ---------- 188
GT#17 KT232010  115       ---T-AA ------ ----------TT---- ------A-TAT- --A-T---- --T---T---- ---------- 188

GT#1  AB067721  211       GAATCAAAACTTTCAACAACGGATCTCTTGGTTCTGGCATC GATG AAGAACGCAGCGAAATGCGATAAGTAATGTGAA 288
GT#3  FJ654148  158       ----------------------------------------- ---- -------------------------------- 235
JAAVMX010000019 19556     ----------------------------------------- ---- -------------------------------- 19633
JAAVMX010000019 32200     ----------------------------------------- ---- -------------------------------- 32277
JAAVMX010000002 18702248  -------------------------------------C--- ---- -------------------------------- 18702325
JAAVMX010000018 851       ------------------------------------ ---A------------------------- ---------- 928
JAAVMX010000019 6389      ------------------ --CAC------------------- ---- --------------- ---------- ------ 6463
JAAVMX010000019 44882     ----------------------------------------- ---- -------------------------------- 44959
GT#4  AB067744  207       A---T--------T--T--------A---T-A---T A--A --------------------A-----------A-- 284
GT#5  AB067740  214       A---T--------T-------A----A---T A--A --A------------A---------A-A-- 288
GT#6  KJ720572  183       A---T--------T-----------A---A--- ---A----T--T-----A--------A-A-- 222
GT#15 KT232017  189       ----T--------T--T--------A---T-A---T A--A --------------------A-----------A-- 266
GT#16 KT232019  189       ----T--------T-----------A----A---- --A----------------A---------A-A-- 266
GT#17 KT232010  189       A---T--------T-----------A----A---- ---A----A-------------A---------A-A-- 266

GT#1  AB067721  289       TTGCAGAATTCAGTGAACCA TCGAATCTTT GAACG CACATTG CGCCCGCCAGCACTCTGGCGGGCATGC CTGTC 362
GT#3  FJ654148  236       -------------------- ---------- ----- ------- --------------------------- ----- 309
JAAVMX010000019 19634     -------------------- ---------- ----- ------- --------------------------- ----- 19707
JAAVMX010000019 32278     -------------------- ---------- ----- ------- --------------------------- ----- 32351
JAAVMX010000002 18702326  -------------------- -------T-- ----- ------- - --------------------------- ----- 18702398
JAAVMX010000018 929       -------------------- C--------- -----C------C---------- ------- ----- 1000
JAAVMX010000019 6464      -------------------C--------- ----- ----C------------------ ----- 6539
JAAVMX010000019 44960     -------------------- ---------- ----- ------- ----------------------CT----- 45035
GT#4  AB067744  285       ---T------T--A--T- --------- A---- T-T-C-A -------T-------A---------- ----- 358
GT#5  AB067740  289       --A------T--A--T- --------- A---- --T---A ------T---------A--------- --A-- 362
GT#6  KJ720572  223       --AT------T--A---- --------- A---- T-T---A ------T---------A--------- ----- 296
GT#15 KT232017  267       ---T------T--A--T- --------- A---- T-T---A -------T-------A---------- ----- 340
GT#16 KT232019  267       --A-------T--A--T- --------- A---- T-T---A -------T-------A---------- ----- 340
GT#17 KT232010  267       --A-------T--A--T- --------- A---- --T---- -------T-------A---------- --A-- 340

GT#1  AB067721  363       CGAGCGTCATCTCAACCCTCGAG CCCCCCGCCTCGCGG CGGCGGGGCCCGGCCTT GGGGGTCACGGCC CCGCGCC 437
GT#3  FJ654148  310       ----------------------- --------------- ----------------- ------------- ------- 384
JAAVMX010000019 19708     ----------------------- --------------- ----------------- ------------- ------- 19781
JAAVMX010000019 32352     ----------------------- --------------- ----------------- ------------- ------- 32425
JAAVMX010000002 18702399  ----------------------- ----- ------ ----------------- ------------- ------- 18702472
JAAVMX010000018 1001      ----------------------- --------------- ------------------G------------ ---- -- 1074
JAAVMX010000019 6540      ----------------------- --------------- ----- ------------- --------C------- 6613
JAAVMX010000019 45036     ----------------------C--------- A----TA--CG--C--------- ------TC-----AA------ 45119
GT#4  AB067744  359       -------T-C--T-----TA-- --------------- ----------------- A----------- --A---- 432
GT#5  AB067740  363       -------T----T--------- --------------- -------------T----- A----T------ ------- 436
GT#6  KJ720572  297       -------T-C--T---- 313
GT#15 KT232017  341       -------T----T--------- --------------- -------------T----- A----T------ ------- 414
GT#16 KT232019  341       -------T-C--T-----TA-- --------------- ----------------- A----------- --A---- 414
GT#17 KT232010  341       -------T----T--------- --------------- -------------T----- A----T------ ------- 414

GT#1  AB067721  438       G  CCCCCTAAACGC AG      TGGCGACC  CCGCCGCGGCTCC  CCTGCGCAGTAGCT  CG  CTGAGAACCT 499
GT#3  FJ654148  385       -  ---------T-- --      --------  -------------  ---------------  --  -C---G---C 446
JAAVMX010000019 19782     -  ------------ --      --------  -------------  ---------------  --  ---------- 19844
JAAVMX010000019 32426     -  ------------ --      --------  -------------  ---------------  --  ---------- 32488
JAAVMX010000002 18702473  -  ------------ --      --------  -------------  ---------------  --  ---------- 18702534
JAAVMX010000018 1075      -  ------------ --      --------  -------------  ---------------  --  ---------- 1136
JAAVMX010000019 6614      -  ---------G--G --      --------GC-------  --------------  --GCT--------- 6680
JAAVMX010000019 45120     -TCC--T---GT----- --CCTAAGTG------- ----GA------A----T-A-------CA-- ---------- 45189
GT#4  AB067744  433       -  ------------ --      -A-T----  -------------  ---A----A-----  TA  --A------- 495
GT#5  AB067740  437       -  ----T------T --      -A------  -------------  ---A--TG------  -A  -C-------- 529
GT#6  KJ720572  313
GT#15 KT232017  415       -  ----------T --      -A------  -------------  ---A--T-A-----  -A  -C------- 477
GT#16 KT232019  415       -  ----------T --      -A------  -------------  ---A--A------  TA  --A------- 477
GT#17 KT232010  415       -  ----------T --      -A------  -------------  ---A--T-A-----  -A  -C------- 477

GT#1  AB067721  500       CGCACCGGGAGCGCGGA      GGCGG TCACGCC  TGG     AA ACCACC   ACACCCTCCA              549
GT#3  FJ654148  447       -------------A-------T  ----- ------- ---     -- ------   ---    ---            492
JAAVMX010000019 19845     -----------------      ----- ------- - ---     -- ------   ----------            19894
JAAVMX010000019 32489     -----------------      ----- ------- - ---     -- ------   ----------            32537
JAAVMX010000002 18702535  -----------------      ----- ------GTGT-  --  ------   ----------            18702586
JAAVMX010000018 1137      -----------------      ----- ------- - ---     -- ------   ----------            1186
JAAVMX010000019 6681      C----------------      ----- ------- - -----   -- ------ACA----------            6733
JAAVMX010000019 45190     ----------------CCCAT-----C--------- ---GATG--C------ACC---------            45251
GT#4  AB067744  496       -----TA----------      ----- ------- --A     -- ----T   -T------            543
GT#5  AB067740  530       ------A--------      ----- ------- --A     -- ----T   --G--            544
GT#6  KJ720572  313
GT#15 KT232017  478       ------A----------      ----- ------- --A     -- ----T   --G--            522
GT#16 KT232019  478       -----TA----------      ----- ------- --A     -- ----T   -T------            525
GT#17 KT232010  478       ------A----------      ----- ------- --A     -- ----T   --G--            522
```

Figure 1. ITS sequence alignment of AB067721, repetitive ITS copies within genome JAAVMX000000000
of H. sinensis Strain IOZ07, and AT-biased genotypes of O. sinensis.

The ITS sequences contain complete or partial ITS1-5.8S-ITS2 nrDNA segments. "GT" denotes the O. sinensis
genotype. The underlined sequence in black represents the 5.8S gene of GC-biased Genotype #1 H. sinensis.
AB067721 is the ITS sequence of GC-biased Genotype #1 H. sinensis. Genome assemblies
JAAVMX010000002, JAAVMX010000018 and JAAVMX010000019 were obtained from H. sinensis Strain
IOZ07 (Shu et al. 2020). One copy each within JAAVMX010000002 and JAAVMX010000018 indicated in
green shows 97.4% or 97.0% similarity to AB067721. JAAVMX010000019 contains 4 repetitive ITS copies,
including 2 copies in black (19404→19894 and 32048→32537), which are 100% identical to AB067721, and
2 other copies in blue (6233→6733 and 44729→45251), with 94.5% and 90.8% similarities to AB067721.
Sequences in red (AB067744, AB067740, KJ720572, KT232017, KT232019, and KT232010) represent AT-
biased Genotypes #4–6 and #15–17 of O. sinensis, respectively. The underlined "GAATTC" sequences in
purple represent the EcoRI endonuclease cleavage sites in the sequences of GC-biased Genotype #1 and GC-
biased genome assembly JAAVMX000000000. EcoRI endonuclease cleavage sites are absent in AT-biased
ITS sequences due to a single-base C-to-T (cytosine→threonine) transition. Hyphens indicate identical bases,
and spaces denote unmatched sequence gaps.

Figure 2. A Bayesian majority rule consensus phylogenetic tree.

Twenty-eight ITS segments of the genome assemblies (ANOV01021709, LKHE01000582, LWBQ01000008,

JAAVMX010000002, JAAVMX010000008, JAAVMX0100000017, JAAVMX0100000018, JAAVMX010000019,

NGJJ01000573, NGJJ01000582, NGJJ01000796, NGJJ01000798, and NGJJ01000799) of H. sinensis strains

(Co18, 1229, ZJB12195, IOZ07, and CC1406-203, respectively) and 25 ITS sequences of GC-biased

Genotypes #1–3 and #7–14 (in blue alongside the tree) and AT-biased Genotypes #4–6 and #15–17 of O.

sinensis (in red alongside the tree) were analyzed phylogenetically using MrBayes v3.2.7 software (cf. Section

II-4). Genome assemblies JAAVMX000000000 and NGJJ00000000 contain multiple repetitive ITS copies (cf.

Table 1). The percent similarities of the genomic ITS sequences with the representative Genotype #1 sequence

(AB067721) are shown in green alongside the tree.

Figure 3. Dynamic alterations in the abundance of amplicons of Genotypes #1 and #2 of O. sinensis in the

stromata of C. sinensis during maturation (modified from Figure 6 of Zhu et al. 2012).

Immature C. sinensis specimens had very short stromata of 1.5 cm on average. Maturing C. sinensis specimens

had stromata of 4.0 cm on average, without the formation of an expanded fertile portion close to the stromal

tip. Mature C. sinensis specimens had long stromata of 7.0 cm on average and showed the formation of an

expanded stromal fertile portion (SFP in red), which was densely covered with numerous ascocarps.

Figure 4. Southern blot of C. sinensis nrDNA in the stroma and caterpillar body of natural C. sinensis

during maturation. [Reproduced with permission from AJBMS (www.nwpii.com/ajbms) Am J Biomed Sci 2010;

2(3): 217-238] [Zhu et al. 2010].

Caterpillar refers to caterpillar body. Genomic DNA templates were isolated from the stromata or caterpillar

bodies of immature (Im) or mature (M) C. sinensis specimens collected from Sichuan Province in China and

*prepared using the restriction enzymes EcoRI, DraI and AvaI. **The upper panel** was probed with an H. sinensis-*

*specific probe. **The lower panel** was probed with a nonspecific 18S internal control probe.*

Figure 5. Alignment of the query sequence in green (the 5.8S gene of Genotype #1 H. sinensis AB067721)
and the 75 of 1894 overlapping (pink bars) shotgun reads (GenBank SRR5428527) of mature C. sinensis.

Table 4. *Percent similarity between the unassembled metatranscriptome shotgun reads (SRA: SRR5428527) of natural C. sinensis and the 5.8S genes of multiple genotypes of O. sinensis and Paecilomyces hepiali, an AB067719-type fungus (Group-E) and Tolypocladium sinensis (Group D).*

Geno-type	GenBank Accession #	% similarity between 5.8S genes and *C. sinensis* shotgun sequences					
		SRR5428527. 17546327.1	SRR5428527. 17554054.1	SRR5428527. 17565575.2	SRR5428527. 17679995.2	SRR5428527. 17911498.2	SRR5428527. 17929569.2
#1	AB067721	**100%**	**100%**	**100%**	**100%**	**100%**	**100%**
#2	MG770309	96%	96%	**100%**	96%	**98%**	96%
#3	HM595984	**100%**	**99%**	**99%**	**100%**	**99%**	**100%**
#7	AJ488254	**98%**	**98%**	**99%**	**98%**	**98%**	**98%**
#8	GU246286	94%	96%	95%	94%	**97%**	94%
#9	GU246288	**99%**	**99%**	**99%**	**100%**	**99%**	**100%**
#10	GU246287	96%	94%	93%	95%	93%	**97%**
#11	JQ695935	**100%**	**100%**	**100%**	**100%**	**100%**	**100%**
#12	GU246296	**99%**	**99%**	**99%**	**99%**	**99%**	**99%**
#13	KT339190	**98%**	93%	93%	**98%**	93%	**98%**
#14	KT339178	**98%**	93%	93%	**98%**	93%	**98%**
#4	AB067744	86%	87%	87%	83%	87%	83%
#5	AB067740	85%	87%	87%	84%	87%	84%
#6	KJ720572	85%	85%	87%	84%	86%	84%
#15	KT232017	85%	88%	88%	84%	88%	84%
#16	KT232019	86%	86%	87%	85%	87%	85%
#17	KT232010	87%	88%	88%	87%	88%	87%
P.h.	EF555097	**99%**	**98%**	**97%**	**99%**	**98%**	**99%**
	AB067719	**98%**	93%	93%	**98%**	93%	**98%**
T.s.	AB067720	**98%**	**97%**	**96%**	**98%**	**97%**	**98%**

Note: GC-biased genotypes are listed in blue, and AT-biased genotypes are listed in red. GenBank sequence ID SRA: SRR5428527 fragments (17546327.1, 17554054.1, 17565575.2, 17679995.2, 17911498.2, and 17929569.2) are the unassembled transcriptome shotgun sequences obtained from the mature C. sinensis specimen collected from Deqin County of Yunnan Province by Xia et al. *(2017). P.h. refers to Paecilomyces hepiali; AB067719 refers to an AB067719-type Group-E fungus; T.s. refers to Group-D Tolypocladium sinensis (Stensrud* et al. *2007). Highly homologous percent similarities (97%–100%) are bolded in green.*

```
GT1 AB067721 1     TAGAGGAAGTAAAAGTCGTAACAAGGTCTCCGTTGGTGAACCAGCGGAGGGATCATTATC   60
GT4 AB067744 1     ----------------------------A--A---T-----------T------   56
GT5 AB067740 1     --A---------------T----------A--A---T-----------T------   60

GT1 AB067721 61    GAGTCACCACTCCCAAACCCCCTGCGAACACCACAGCAGTTGCCTCGGCGGGACCGCCCC   120
GT4 AB067744 57    -----------C--------------A--C----------------------------   116
GT5 AB067740 61    ----T--T----TT---------ATA--T--T-T-----------TA------------T   120

GT1 AB067721 121   GGCGCCCCAGGGCCCGGACCAGGGCGCCCGCCGGAGGACCCCCAGACCCTCCTGTCGCAG   180
GT4 AB067744 117   -------T-----------T--------------------TT----------A---T--   176
GT5 AB067740 121   A------T-A---------T-AA-----------------TT----------A-TAT--   180

GT1 AB067721 181   TGGCATCTCTCAGTCAAGAAGCAAGCAAATGAATCAAAACTTTCAACAACGGATCTCTTG   240
GT4 AB067744 177   -A---------T---T---------------A---T-A-ACTTTTAATAACGGATCTCTTA   236
GT5 AB067740 181   -A-T------T---T---------------A---T---ACTTTTAACAACGGATCTCTTA   240
Primer 5.8S-F                                         ACTTTTAACAACGGATCTCTT

GT1 AB067721 241   GTTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTCA   300
GT4 AB067744 237   ---T-A----TA--A---------------A-------------A-----T------T-   296
GT5 AB067740 241   -----A----TA--A--A---------------A---------A-A----A-------T-   300

GT1 AB067721 301   GTGAACCATCGAATCTTTGAACGCACATTGCGCCCGCCAGCACTCTGGCGGGCATGCCTG   360
GT4 AB067744 297   --A---T-----------A----T-T-C-A-------T--------A------ATGCCTG   356
GT5 AB067740 301   --A---T-----------A------T---A-------T--------A------ATGCCTA   360
Primer 5.8S-R                                                    ATGCTTA

GT1 AB067721 361   TCCGAGCGTCATCTCAACCCTCGAGCCCCCCGCCTCGCGGCGGCGGGGCCCGGCCTTGGG   420
GT4 AB067744 357   TCCGAGCGTTACCTT------TA--------------------------------A--   416
GT5 AB067740 361   TCCGAGCGTTATCTT---------------------------------T-------A--   420
Primer 5.8S-R      TCCGAGCGTTATCTT

GT1 AB067721 421   GGTCACGGCCCCGCGCCGCCCCCTAAACGCAGTGGCGACCCCGCCGCGGCTCCCCTGCGC   480
GT4 AB067744 417   -----------A------------------A-T-------------------A---   476
GT5 AB067740 421   ---T--------------T------T---A------------------------A--T   480

GT1 AB067721 481   AGTAGCTCGCTGAGAACCTCGCACCGGGAGCGCGGAGGCGGTCACGCCGTGAAACCACCA   540
GT4 AB067744 477   -A-----TA--A-----------TA-----------------------A-------T-   536
GT5 AB067740 481   G-------A-C--------------A--------------------A-------T-   540

GT1 AB067721 541   CACCCTCCAGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACTTAAGCATATCAATAA   600
GT4 AB067744 537   T------TG---A---------T----------------A----------------   596
GT5 AB067740 541   -G--T-TT----A---------T----------------A------------T-----   600

GT1 AB067721 601   GCGGAGGAAAAGAAACCAACAGGGATTGCCCCAGTAACGGCGAGTGAAGCGGCAACAGCT   660
GT4 AB067744 597   --------   604
GT5 AB067740 601   ---------------T--T---------TT-----------A------T--T----   660
```

Figure 6. ITS sequence alignment of AB067721 of GC-biased Genotype #1, AB067744 and AB067740 of

AT-biased Genotypes #4–5 of O. sinensis and the 5.8S-F/R primers.

"GT1", "GT4" and "GT5" represent Genotypes #1 and #4–5 of O. sinensis, respectively. The sequences in

blue are 18S and 28S genes (5' and 3' end regions, respectively). Those in pink represent sequences of 5.8S

genes. Those in black are ITS1 between the 18S and 5.8S genes and ITS2 between the 5.8S and 28S genes. The

sequences underlined in green are for primers 5.8S-F/R, while primer 5.8S-R was converted to the sense chain

sequence. Hyphens indicate identical bases.