Network analysis of chromophore binding site in LOV domain

Rishab Panda [1], Pritam K. Panda [2,3], Janarthanan Krishnamoorthy [4], Rajiv K. Kar [5*]

[1] Department of Chemical Sciences, Indian Institute of Science Education and Research Mohali, Punjab, India.
[2] Condensed Matter Theory Group, Materials Theory Division, Department of Physics and Astronomy, Uppsala University, Box 516, SE-751 20 Uppsala, Sweden.
[3] Division of Immunology and Chronic Disease, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden.
[4] School of Biomedical Engineering, Jimma institute of Technology, Jimma University, Jimma 378, Ethiopia.
[5] Jyoti and Bhupat Mehta School of Health Sciences and Technology, Indian Institute of Technology Guwahati, Guwahati, Assam – 781039, India.

* Correspondence email: rajivkar@iitg.ac.in

Keywords

Abstract

26
Photoreceptor proteins are versatile toolbox for developing biosensors for optogenetic
applications. These molecular tools get activated upon illumination of blue light, which in
turn offers a non-invasive method for gaining high spatiotemporal resolution and precise
control of cellular signal transduction. The Light-Oxygen-Voltage (LOV) domain family of
proteins is a well-recognized system for constructing optogenetic devices. Translation of
these proteins into efficient cellular sensors is possible by tuning their photochemistry
lifetime. However, the bottleneck is the need for more understanding of the relationship
between the protein environment and photocycle kinetics. Significantly, the effect of the local
environment also modulates the electronic structure of chromophore, which perturbs the
electrostatic and hydrophobic interaction within the binding site. This work highlights the
critical factors hidden in the protein network linking with their experimental photocycle
kinetics. It also presents an opportunity to quantitatively examine the alternation in
chromophore equilibrium geometry and identify details which have substantial implications
in designing synthetic constructs with desirable photocycle efficiency.

41
42

43 **Graphical Abstract**

44
45

46
47 Graphical representation of LOV photoreceptor, translated into residual community network
48 to understand the role of protein network onto the photophysical properties.
49

Introduction

Optogenetics represents a new paradigm that can control biological responses through light. In the past decade, optogenetic tools have rapidly evolved through understanding of the fundamental principles governing the biomolecular systems and technological advancement. These non-invasive tools are promising for diagnosis and treating several diseases afflicting humankind. The advanced application of the optogenetics-based biomedical tool include treatment of diseases such as diabetes, neurological conditions, heart diseases, and cancer.[1] They have also been used for gene-editing and epigenome regulation that have significant applications in molecular and cellular biology.[2] The area of optogenetics could target either neuronal cells that function in neuronal signalling activation or suppression; or non-neuronal cells that affect the signalling cascade for precise monitoring and control of biological functioning. One such example of non-neuronal optogenetic tool development utilizes families of photoactivable proteins - the LOV (light, oxygen, and voltage) family of proteins. LOV domain proteins provide a novel platform owing to their small molecular size and quick response time.[3] These tools take advantage of the fluorescent properties of LOV proteins to develop fluorescence imaging tools and possible implementation in super-resolution microscopy.[4] The LOV-based fluorescent reporter systems have shown increased brightness, improved stability and functionality in low-oxygen and anaerobic conditions, compared to the green fluorescent proteins.[5] In all cases, the core LOV domain signals the effector elements through the highly variable C-ter and N-ter extensions.[6]

The LOV domain is the photosensory module that responds to UVA or blue light[7] and are widespread among several species, including bacteria, fungi, archaea, protists, and plant species. They are a subclass member of the superfamily – PAS domain (Period-ARNT-Singleminded), which comprises α-helical segment and five-stranded antiparallel β-sheets.[8,9] Structurally, the active site of LOV contains a flavin chromophore (Flavin mononucleotide, Flavin adenine dinucleotide, or Riboflavin) and is characterised by the presence of a conserved motif GXNCRFLQ.[8]

The photochemistry of the LOV domain is triggered when the chromophore absorbs light and attains a singlet excited state (S_1), followed by intersystem crossing to a triplet state (T_1). The process induces a proton transfer to the N5 atom, changing the hybridization of the C4a atom to sp^3, and generates a tilt in the planarity of the isoalloxazine ring. This photocycle mediated by a radical-pair mechanism involve the thiol group of a conserved cysteine and the C4a atom of flavin, enabling electron transfer and subsequent formation of a covalent adduct. The ground state (also referred to as the resting or dark state) is characterized by flavin in its oxidized form, absorbing blue light at 450 nm (2.75 eV). The light-adapted signalling state containing the C4a adduct absorbs at wavelength 390 nm (3.18 eV). While the kinetics of the transition, $S_1 \rightarrow T_1$, occurs in the nanosecond time regime, the conversion of radical pair to adduct falls within microseconds timescale. Notably, the overall photocycle attains thermal reversibility in the dark state,[10] the timescale of this process is a scientific puzzle to be solved as it ranges between a few seconds to several hours in LOV variants.[11] It also raises opportunities to tune the limetime of the photocycle of LOV domain and extend its scope for developing an efficient optogenetic tool.[12] Studies on LOV proteins and their variants have

shown that mutation in the chromophore binding site can affect the photocycle duration. In
Avena sativa phototropin 1 (AsLOV2), mutations like Val to Thr reduces the time constant to
a few seconds, Asn to Leu extends to hours, whereas, Asn to Ala has no effect.[13–15] Likewise,
in fungal circadian clock photoreceptor Vivid (VVD), Ile to Val double mutants, Met to Ile
double mutant, and Met to Leu double mutants reduces, increases, or have no effect on the
time decay constant, respectively.[16] The relationship between mutation and changes in the
time constant links to the change in electrostatic environment,[17] hydrogen bonding residues[18]
and steric effect[19] on flavin affecting the adduct's stability.[15] Mutations in the active site can
perturb the solvent accessibility near the chromophore and facilitate proton transfer involving
the photocycle time constants.[20,21]

Several experimental studies based on LOV domain protein structures shed light on the effect
of specific amino-acid point mutation, their side-chain orientation, and the potential of being
H-bond donors /acceptors in the close vicinity of the chromophore. However, an existing gap
in the current pool of knowledge is a comparative structural analysis that could explain the
molecular origin of the mutational effect on the LOV photocycle. Here, we have considered
several LOV domain proteins from diverse sources to construct a phylogenetic tree and
analyzed their clustering pattern. Structural models of protein based on the cluster's distance
are energy minimized and suitable cavities are hydrated with water molecules for structural
analysis. Subsequent analyses are based on normal mode simulations to create community
networks based on the amino acid arrangement. We also employed graph-based algorithms to
understand the H-bonding pattern in LOV proteins and the chromophore binding site
arrangement and function.

Materials and Methods

Phylogenetic analysis
Given the significant number of LOV protein sequences available in the databases, our
foremost approach was to construct a distance-based phylogenetic tree. The objective is to
determine their closest relatives among the eukaryotic and prokaryotic origins. However, the
phylogeny relationship is based on the LOV domain's changes in the amino acid sequences.
The protein sequences were searched using the NCBI server's protein Basic Local Alignment
and Search Tool.[22] The initial seed proteins were collected from the literature.[23–26,] and the
search was performed for non-redundant sequences using a threshold E-value of 0.05.
Notably, the pattern of amino acid and conserved Cys, which forms an adduct with flavin in
the light state, is cross-checked for all sequences.

Multiple sequence alignment was performed using the MUSCLE algorithm (See supporting
information).[27] Construction of distance trees uses aligned sequences. The MEGA11 program
was used to construct the phylogenetic tree employing the maximum likelihood method.[28]
Jones-Taylor-Thornton's substitution model was adopted for amino acids , whereas, the
heuristic nearest-neighbour interchange method was used for building the tree inference . To
test the phylogeny, we employed the Boostrap method with 1000 replicas. The unrooted

distance tree was finally constructed and visualized using the interactive Tree of Life (iTOL)
server.[29]
139
Structural Modelling
The constructed phylogenetic tree retrieved representative three-dimensional structures from
individual clusters from the protein data bank (PDB). In total, 18 such structures constitute
the dataset for comparative analysis. The corresponding photocycle kinetics were collected
for literature. The PDB format of each structure was cleaned to remove any solvent or ion
other than protein, a chromophore (FAD, FMN, or Riboflavin), and water molecules. In the
case of dimeric and multimeric structures, only one chain is processed for further analysis.
Missing residues in the structures are refined using the SWISS-MODEL server.[30] The
structural coordinate in 5DJT contains a mutation in the active site Cys50Ala, which was
back-mutated to retain Cys in the present work.[31] Note that the resulting network becomes
crowded while considering protein residues outside the LOV domain. The resulting network
in such a case indicates several nodes that either have one edge or no edge within the path
connected to the chromophore. Hence, we only considered the residues present in the PAS
domain of LOV protein to refine the network.
154
The presence of water molecules within the LOV chromophore binding site is crucial for its
photochemistry.[20,32] It has been reported to stabilize the internal charges in protein and
participate in hydrogen bonding interaction. To account for the hydration of proteins, we used
the Dowser++ program for hydrating the biomacromolecule.[33] The prediction of water sites at
the cavity (at the protein surface and folded cavity) is based on an internal probe of 1 Å for
finding the cavity and a semi-empirical estimate of binding energy more significant than 10
kcal.mol^{-1}.
162
Hybrid QM/MM calculation
The three-dimensional coordinates of protein structures are input for the hybrid quantum
mechanics/ molecular mechanics (QM/MM) calculations. The QM region comprises flavin's
isoalloxazine ring and is treated with the TD-DFT method using the B3LYP/6-31G* level of
theory. The remaining system constitutes the MM part and is treated with Amber force fields
ff14SB.[34] Dispersion correction is applied using Grimme's D3/B-J damping variant.[35] Other
parameters include resolution of identity (RI) for Coulomb integrals (J) and COSX for
exchange integrals, denoted as RIJCOSX.[36] In these simulations, the backbone atoms of
protein are fixed to their crystallographic position, while side-chain atoms and water
molecules present around 8 Å of the isoalloxazine ring are relaxed. All QM/MM-based
geometry optimization is performed with Orca program[37] (quantum chemistry package)
interfaced with the DL_POLY module (for MM) of Chemshell.[38]
175
Network Construction
Protein networks are constructed based on the QM/MM optimized geometries. The
comparative analysis was based on three different network types, viz., (i) hydrogen-bond
interaction network, (ii) hydrophobic interaction network; and (iii) residue interaction-based
dynamic correlation network.

The residual interaction data obtained from the protein's normal mode analysis (NMA)
displays the residue-by-residue cross-correlations matrix. The preliminary graph consists of
amino acids as nodes and the distance between them as edges. It is followed by a coarse-
grained approach that uses the Girvan-Newman clustering method. The final network thus
converts the all-residue network into communities of highly intra-connected but loosely
interconnected nodes. For simplification of networks, they are represented using centrality
measures like degree centrality and betweenness centrality to quantify the relevance of each
node in the graph. 'Degree centrality' denotes the number of edge incidents to a given node.
In contrast, 'betweenness centrality refers to the shortest path between two nodes, so the
number of edges that pass through is minimized. Notably, the nodes with high betweenness
may have considerable influence within a network as they control the information passing
between other nodes. Removal of such nodes will disrupt communications as they lie on the
most significant number of paths taken by messages transmitted through the network.
Interested readers may refer to the study by Deb et al., where similar methods are used to
study residue interaction and protein contact networks of LOV proteins.[39] The networks in
this study are created using Bio3D working through RStudio IDE, R version 4.2.0.[40]
Customization and visualization of networks are performed using the igraph program.
Description of protocol for network construction and the codes are shared in the Supporting
Information.
200
201 The details of hydrogen bonding and hydrophobic interaction are further collected in the form
202 of graph-based networks. The participating residues/water molecules represent the nodes, and
203 the edges link. The connections between nodes are based on Dijkstra's algorithm. Bridge
204 program, a plugin with Pymol, was used for this analysis.[41] Specific interaction of amino acid
205 with chromophore atoms is analysed using LiBiSCo.[42]
206
207 **Results and Discussion**
208
Sequence-guided phylogenetic reconstruction of LOV proteins. Understanding
phylogenetic connections are crucial for a wide range of biological investigations. Major
evolutionary transitions rely on an accurate phylogenetic tree, such as molecular adaptation
detection, morphological character evolution, and recreating demographic shifts in separated
species. Given a large number of LOV protein sequences available in the annotated form, it is
crucial to establish a phylogenetic tree based on variation in the protein sequences. Multiple
sequence alignment was performed using the Muscle algorithm. The bootstrapped tree was
obtained with 1000 replicates as an evolutionary model. The constructed unrooted
phylogenetic tree is shown in Fig. 1, where the nodes and branches of proteins with similar
sequences are clustered together and coloured distinctly. LOV belonging to the same
kingdom are often found to have diverse functioning. For instance, LOV protein belong to the
same kingdom – Viridiplantae has been classified in to three separate clusters. to the first
cluster belong to the plant phototropins, the second cluster contains LOV proteins from
zeitlupe (ZTL) and flavin-binding kelch repeat F box protein (FKF1) and the third cluster
includes neochromes, aureochromes, and protein twin-LOV.
224

Another such example is the fungi, which got distributed across two clusters. One class of
protein belongs to fungal G-protein receptors, while the other belongs to white collar 1
(WC1) and VVD photoreceptors. Notably, the LOV module is widely dispersed among the
proteobacteria belonging to alpha, gamma, and actinobacteria classes. The differences in the
sequence and its arrangement in the chromophore binding site are responsible for variability
in photoinduced molecular reactivity. Thus, understanding the structural positioning of the
amino acid side chains will provide insights on the factors governing the on/off kinetics.
Available structural models of the experimentally resolved protein were retrieved from the
Protein Data Bank. In total, 18 structural models were considered for network analysis. The
corresponding cluster group of these LOV proteins was marked in the phylogenetic tree (Fig.
1). More details on the organismal source, along with their rate kinetics and experimental
conditions, are provided in the Table 1. For further analysis, the proteins have been
categorized into fast, intermediate, and slow kinetics based on their time constants.

238
239
240 Figure 1: Phylogenetic tree depicting the diversification of the LOV domain proteins.

Table 1: Classification of the proteins based on their time constant values.

PDB code	Organism	Time constant (and relevant details)	Ref
1N9O[a]	*Chlamydomonas reinhardtii*	200/168 sec	43
1G28[a]	*Adiantum capillus-veneris*	90 sec	44
5DJT[a]	*Avena sativa; Staphylococcus aureus*	~50 sec	
4EEP[a]	*Arabidopsis thaliana*	68 sec[d]	45
4KUK[a]	*Dinoroseobacter shibae*	9.6 sec (20 °C)	46
7A6P[a]	*Pseudomonas putida*	210 sec (20 °C)	
3UE6[a]	*Vaucheria frigida*	480 sec	
3P7N[a]	*Erythrobacter litoralis*	25.5 sec	
4HJ3[b]	*Cereibacter sphaeroides*	~39 min	
2PR5[b]	*Bacillus subtilis*	~65 min (23 °C, pH 7)	
6HMJ[b]	*Nakamurella multipartita*	~37 min (22 °C)	52
4R38[b]	*Erythrobacter litoralis*	55 min	53
4WUJ[b]	*Trichoderma reesei*	25 min	54
5DKL[b]	*Phaeodactylum tricornutum*	~25 min	
6I22[b]	*Ochromonas danica*	112 mins[d]	
5SVU[c]	*Arabidopsis thaliana*	~6.6 hrs	
3T50[c]	*Brucella melitensis*	9.5 hrs (20^0 C, 6.5 pH)[d] 5.8 hrs (35^0 C, 7 pH)[d]	58
2PD7[c]	*Neurospora crassa*	2.7 hrs (25 °C)[d]	59

Notes: [a] fast kinetics (<500 sec); [b] intermediate kinetics (500 to 7000 sec "or" up to 112 min); [c] slow kinetics (>7000 sec "or">2 hrs); [d] referred to as half-life in original reports.

Classification of data set based on the time constant. The structural models constituting the dataset in this study are classified based on the adduct decay (time constant) kinetics. In particular, three regimes of photocycle kinetics are used for classification, viz. fast (<500 sec), intermediate (500 to 7000 sec), and slow (>7000 sec) (detailed in Table 1). The protein networks constructed in our work are based on normal mode analysis, aiming to understand the chain connectivity related to the folding pattern of proteins. Processing the coordinate data from PDB with coarse-graining helps decompose constituent amino acid residues' mutual and intrinsic motion. Here the centrality of individual residues represents the topological importance of the residue-residue cross-correlation network. The residues act as nodes and the edge which interconnect two nodes are weighted in proportion to the degree of correlated motion. The betweenness centrality in the network was calculated by finding the shortest path between pairs and scoring each edge (n) of the paths with the inverse value of the number of shortest paths (m); betweenness score = n/m. Note that the residues present at the C-ter and N-ter of model proteins did not show a direct perturbative effect on the chromophore binding site but indirectly through allosteric pathways. Since the total number of residues in each protein structure is not constant, it is challenging to identify a particular trend from the resulted protein network. For structural modelling, the terminal residues were removed to bring the protein networks within a range of betweenness centrality. Thus, the

shortest paths for all pairs were optimized according to the classification of the dataset.
ranging between 14 to 17 for fast cycling, 20 to 24 for intermediate cycling, and 30 to 33 for
slow cycling proteins. Details of individual protein models are provided in Table 2. A
graphical representation of the classification based on betweenness centrality is shown in Fig.
2. It also includes the total number of residues in each protein of the structural dataset. The
coordinates of modelled structures of the LOV domain used for constructing networks are
available in the supporting information.

Table 2: Details of LOV structural models used for network construction.

Protein Name	PDB code	N-ter residue	C-ter residue	Total residue numbers [a]	Water molecules added [b]	Centrality [c]
Phot-LOV1 domain	1N9O	4His	109Ser	106	25	17
Flavin-binding domain, LOV2, from the chimeric phytochrome/phototropin photoreceptor PHY3	1G28	1Lys	104Met	104	9	17
LOV2 Domain of Avena Sativa	5DJT	13Asn	121Asp	109	15	16
LOV2 domain of phototropin 2	4EEP	6Ile	115Val	110	23	15
A superfast recovering full-length LOV protein from the marine phototrophic bacterium	4KUK	20Asp	139Leu	120	17	16
Short LOV protein PpSB2-LOV	7A6P	31Val	146Glu	116	16	16
Blue-light photoreceptor Aureochrome1 LOV	3UE6	46Gln	154Ala	109	12	14
Light-activated transcription factor El222	3P7N	12Ser	123Glu	112	22	17
LOV protein	4HJ3	14Asp	127Ser	114	15	23
Dimeric LOV Photosensor YtvA	2PR5	10Val	117Tyr	108	25	20
RNA-binding Light-Oxygen-Voltage Receptor	6HMJ	243His	350Val	108	19	21
EL346 blue-light-activated histidine kinase	4R38	28Ile	129Gln	102	9	20
Fungal LOV Domain Photoreceptor	4WUJ	37Ser	146Glu	110	27	24
Blue light photoreceptor Aureochrome 1a LOV	5DKL	17Gln	128Val	112	10	20
LOV domain	6I22	19Val	123Tyr	105	19	20
LOV domain of ZEITLUPE	5SVU	14Thr	133Ile	120	24	33
LOV-HK sensory protein	3T50	2Ala	116Val	115	15	33
Fungal Blue-Light Photoreceptor Vivid	2PD7	27Val	148Cys	122	19	30

297 **Notes:** [a] Total number of amino acids present in the structural model used for normal mode
298 analysis. [b] Number of water molecules added in the 8 Å of isoalloxazine ring, using the
299 Dowser program. [c] Betweenness Centrality.

300
301

302
303
304
305 **Figure 2:** Dataset classification LOV domain based on adduct decay kinetics and
306 betweenness centrality of protein networks.
307
Interaction analysis of the chromophore within the active site. Interaction details help
quantify the factors responsible for the stabilization of chromophore. Fig. 3 shows the
interaction details of LOV proteins exhibiting fast kinetics. The ring I in the isoalloxazine
was involved in polar contacts with the electronegative N and O atoms. On the contrary, ring
III engages in hydrophobic interaction, attributed to the presence of steric bulky methyl
groups. The presence of H-bonding of the C2=O group in the active site accelerates the
photocycle kinetics.[60] It also stabilizes the isoalloxazine ring within the active site, as
revealed through vibrational frequency measurements.[61] Except for 7A6P, 5DJT, and 4KUK,
the rest of the fast kinetic class of proteins showed water molecules to be in close contact
with the carbonyl moiety adding to its polarity.[62] (Fig. 3F, G, and H). Within the active site,
the polar contact is also facilitated through Asn, as observed for all fast kinetics proteins. In
4EEP the Asn residue is making polar contact with C4=O, which however could re-orient
during the dynamic conformational changes (Fig. 3E). The dynamic flipping of Leu and Gln
residues stabilizes isoalloxazine from both sides.[63] Interestingly, another Asn residue is
involved in making H-bond with N3 and/or C4=O group. In the case of IG28, 3UE6, 4EEP,
and 5DJT, we observe an interaction between a Gln residue and the N5 of ring II (Fig. 3A, B,
E, and G). Previous reports have identified this residue crucial for tuning the LOV
photochemistry inferred from mutation studies.[59,64,65] In the absence of Gln residue in 3P7N,
1N9O, and 7A6P, the H-bonding within the active site are compensated by water molecule(s)
(Fig. 3C, D, and F). The contour map of hydrophilic interaction also shows substantial
contributions from Arg and other Gln residues. However, their interactions are not associated
with the isoalloxazine ring but with the phosphate chain and are crucial for chromophore

stabilization. Arg residue forms a conserved salt bridge with the phosphate group and reduces
the dynamicity of chromophore.[63] Note that this interaction has also been predicted to
stabilize adducts, by affecting the decay kinetics of the adduct.[66] Overall, we see the
interactions of Gln/Asn residues and water molecules are predominantly electrostatic in
nature and interacts with ring I as they contains polar atoms. Whereas, the hydrophobic Val,
Leu, and Ile residues interacted adjacent to ring III containing two methyl groups. The Ala
residues found in a few structural models, are known to facilitate faster kinetics.[62] Some of
the aromatic-aromatic contacts (π-π stacking type) mediated through Phe and Tyr residues
are also observed.
339

343 **Figure 3:** Interaction analysis of fast-cycling LOV proteins. The contour map shows the
344 relative weight of polar contacts (polar interaction). Residue-wise interaction is shown for
345 each protein model with chromophore in stick model and residues coloured according to the
346 nature of hydrophobic and polar contacts. The colour codes used to describe interaction are
347 shown.
348

In the case of intermediate cycling proteins (Fig. 4A-G), one Asn residue was involved in
making H-bonding with C2=O. Asn residue stabilizes the chromophore interaction by
forming polar contact with N3 and/or C4=O. Exceptions to this trend are 6I22 and 4WUJ
(Fig. 4D and E). The role of Gln residue in stabilizing the isoalloxazine ring through polar
contacts is evident except for 6HMJ (Fig. 4G). The role of water molecules in relaying the H-
bond to the isoalloxazine ring at N5 position is a well-established hypothesis.[67–69] A water
molecule close to N5 is in line to this for the intermediate cycling proteins, except 4R38 and
4WUJ (Fig. 4C and E). Notably, the extent of polar contacts of the ring-I is relatively less

than that of the fast kinetic group. Additional water-mediated contacts strongly involve
hydroxy groups in the extended phosphate chain. Arg residue forms salt-bridges with
phosphate and Gln and/or Asn residues are involved with H-bond with the hydroxy groups,
rendering the chromophore more stable. The hydrophobic contacts involving the π-alkyl and
π-π stacking type interactions are similar to that of the fast kinetic group. Interestingly, the
Cys and Met in this group prefer π-Sulfur type interaction, instead of induced-dipole
interactions.
364
The number of structural models in the slow kinetic groups is comparatively less, making it
difficult to identify a reliable trend. Nevertheless, the water molecules mediating stabilization
through H-bond with hydroxyl groups of extended chains exists only in 2PD7 and 3T50 (Fig.
4H and I). In 5SVU (Fig. 4J), one additional water molecule enabled H-bond with N3 of ring
I and N5 of ring II. The role of Asn in forming conventional H-bonds is consistently observed
in this group, and Gln is found only in the case of 3T50. The hydrophobic interactions were
predominantly mediated by Leu and Ile residues through π-alkyl type contacts with ring III.
The hydrophobic interaction involving a larger side chain or conversely, the absence of
shorter side chain like Val, could be a contributing factor for longer time scale in decay.[62]

374
375
376

Figure 4: Interaction analysis of intermediate cycling (A-G) and slow cycling (H-I) LOV proteins. The contour map shows the relative weight of polar contacts (polar interaction). Residue-wise interaction is shown for each protein model with chromophore in stick model and residues coloured according to the nature of hydrophobic and polar contacts. The colour codes used to describe interaction are the same as in Fig. 3.

Quantifying interacting with chromophore atoms. This section quantitatively illustrates the residue-specific interaction with atoms of the chromophore. Fig. 5 shows the average distance between active-site residues with specific atoms of the isoalloxazine ring of flavin. The H-bonds are mainly formed by three polar atoms, in which nitrogen at N3 and carbonyl groups' oxygen at O2 and O4 contribute significantly. Considering the three kinetic groups, the average distance between N3, O2, and O4 with the corresponding residue follows a similar trend. The distance is in order O4 > O2 > O4 (Fig. 5A, C, and D). However, the contributions with water molecules are not include included in this analysis. We also performed distance analysis for hydrophobic interactions analysis. The pattern of average interaction distance obtained shows differences crucial for their photocycle kinetics. The N5 atom found the minimum distance in all three groups, indicating that strong hydrophobic interaction is incident at the chromophore's reaction centre. For the fast kinetic group (Figure 5B), relatively distant interaction is for C6 and C9a. On the contrary, the maximum interaction distance for the intermediate kinetic group is for C9a and C9, whereas for slow, it is C9 and C7M (Fig. 5D and F). This indicates that the dominant factor determining the photocycle kinetics is the hydrophobic and steric restraint induced due to non-polar residues in the active site.

401
402
403 **Figure 5:** Average interaction distance between chromophore atoms and active site residues
404 are shown for fast (A, B); intermediate (C, D); and slow kinetic group (E, F). H-bond
405 distance on left panel and interaction distance on right panel denotes polar and hydrophobic
406 interaction, respectively.
407
Dynamic correlation network. The dynamic correlation network was constructed using the
information from normal mode analysis. The residue-based network was converted into
community clusters and weighted according to their significance within the network.
Calculation of these undirected network graphs considers the betweenness centrality for
partitioning them into clusters.[40] The networks are optimized for a range of betweenness
centrality, corresponding to their classification according to kinetics. It helps in avoiding
unambiguity and rationalizes the networks for comparison. Tight packing of chromophore
within the LOV binding site is facilitated by β-sheets surrounded by Eα and Fα helices. Their
cross-talk between community have direct effects on the stability of the adduct. It is
evidenced through a network of H-bonds and electrostatic interaction, facilitated mainly by
residues present in Eα, Hβ and hydrophobic by Gβ. While residues in Iβ contribute through
both polar and non-polar contacts.[9,46] The protein correlation network for the fast cycling
group are in Fig. 6, where the community clusters are shown with distinct colours. The
corresponding protein segments are rendered with similar colours to denote the network
nodes.
423

The graph pattern from 1G28 and 3UE6 was devoid of interlinked branches, except for two
short bifurcations (Fig. 6A, B). The number of nodes present in these networks are 10 and 8,
respectively. The β-sheets (Gβ and Hβ) are intensely engaged in community interaction in all
structural models of this group, with an exception of 3UE6. Likewise, Iβ also show strong
network interaction except for 3UE6 and 1G28. The loop regions between β-sheets,
connecting Bβ-Cα, and Eα-Fα were found to be central and are linked to two or more nodes
in the dynamic correlation network. Conformational changes in these regions alter photocycle
kinetics as (i) they are solvent-exposed on one side with buried hydrophobic part; (ii) they are
reported to show temperature dependency.[8,44] The correlation network of 1N9O and 7A6P are
alike, containing 12 and 10 nodes, respectively (Fig. 6D, F). The phototropin 1N9O and
bacterial protein 7A6P can accommodate more water molecules within the chromophore
binding site.[44,47] The triad in the network which links Fα with the loop between Hβ-Iβ and
connecting the β-sheets can facilitate increased solvent accessibility. This promote faster
deprotonation (flavin N5 position) and faster kinetics of 200 and 210 sec (Table 1). The
number of nodes present in the correlation network of 3P7N are 9, whereas, 7A6P and 5DJT
contain 10 nodes. Notably, the number of nodes are high in 4KUK and 4EEP containing 14
and 13 nodes, respectively.
441
442

446 **Figure 6:** Dynamic correlation network of fast-cycling LOV proteins. The spheres represent
447 coarse-grained networks based on dynamically coupled communities, and the linking is based
448 on calculated betweenness. Similar colours represent individual community clusters on the
449 correlation network and protein's secondary structures.
450

The hydrophobic region in β-sheets is crucial for LOV proteins having intermediate kinetics.
In the case of 2PR5, Hβ and Iβ denote the central node connected to several residual
communities (Fig. 7A).[70] The congregation formed in 4R38 through β-sheets and Fα appears
dispersed but are connected to the more residual community. They are crucial in governing

collective normal modes. Though the time constants for 2PR5 and 4R38 are comparable
(3880 and 3300 sec), the key difference could be the presence of chromophore and riboflavin,
respectively. Moreover, the faster adduct decay in 2PR5 (Fig. 7C) are due to factors like - the
presence of Ala in Fα instead of conserved basic residues lacks a polar contact with an
extended chain; the presence of residues Met in re-face; and Phe near isoalloxazine ring.[51,53]
The Gβ-Hβ-Iβ region is linking Aβ-Bβ in one closed loop and to Fα in another loop, within
the network of 4HJ3 (Fig. 7B), restricting the solvent accessibility towards chromophore site.
Ile and Leu residues, positioned re-faced to isoalloxazine ring, govern the solvent
perturbation and the kinetics.[50] The total number of nodes in the network of 2PR5 and 4HJ3
are 11, while it is 9 in 4R38. The Iβ sheet in 4WUJ loosely packs the chromophore site,
indicating the possibility of solvent exposure towards the active site (Fig. 7D). This
hypothesis is based on collected evidence from force-field-based simulations.[69] The
corresponding residual community is linked to Aβ and Bβ, and is collectively crucial for
governing the decay kinetics.[24] The network pattern found in 5DKL and 6I22 are alike, where
the part of Fα with Gβ, Hβ, and Iβ appear as central nodes (Fig. 7E and G), playing a crucial
role in chromophore stability. The simulation-based machine-learning analysis confirmed the
dynamic difference between these regions and the protein's remaining part.[71] Notably, the
hydrophobic residues in Iβ region are essential for cross-talk between other residual
communities of protein.[72,73] The number of nodes revealed in the dynamic correlation
network for 4WUJ, 5DKL, 6HMJ, and 6I22 are 10. Notably, the involvement of Aβ, Bβ, and
Cα remains uniform in all protein models. They are known to participate in solvent mediated
polar contact on the outside and facilitating a hydrophobic environment towards (inside to)
the chromophore. However, the role of Fα in cross-talk with β-sheets is more significant in
the intermediate kinetics group.
479

480
481

Figure 7: Dynamic correlation network of intermediate (A-G) and fast (H-J) cycling LOV proteins. The spheres represent a coarse-grained network based on dynamically coupled communities, and the linking is based on calculated betweenness. Similar colours describe individual community clusters on correlation networks and protein's secondary structures.

All three protein models of slow kinetics, viz. 2PD7, 3T50 and 5SVU have 12 nodes in the complex dynamic correlation network. The role of Bβ is crucial in connecting with other residual communities and maintaining close packing of the chromophore in all three models. In the case of 2PD7, perturbation in Bβ-loop-Cα is critical in maintaining the polar contacts by Gln and Asp (Fig. 7H).[59,74] Met and Arg's role in stabilizing the sheet-helix packing has also been demonstrated through molecular dynamics simulation.[75] Notably, in 3T50, the sheet-loop-helix region at N-ter remains central to other adjacent loops of the residual community (Fig. 7I). The structural compactness of these structural regions remains the same after adduct formation, as determined through NMR chemical shifts.[58] However, in the case of 5SVU, this region is shared by more than one node (Fig. 7J). Overall, the Bβ-loop-Cα region appears crucial for slow kinetic LOV protein; but more structural models are required to understand its significance.

Correlation between bond length and rate kinetics

The Hybrid QM/MM method helps evaluate the protein environment's effect on the chromophore, especially on the bonded and non-bonded interactions.[76] The structural models of the LOV domain are segmented into three groups based on their rate kinetics and subjected to hybrid QM/MM simulation to identify the changes in equilibrium bond length of chromophore. The QM region selection was uniform in all models, which constitutes the isoalloxazine ring. The protein environment was treated as point charges for which the parameters were obtained from the MM force fields. We analyzed the standard deviation in bond lengths over the structural models for each bond. The numerical data obtained from simulations are summarized in Table S1 of Supplementary information, and the graphical representation is shown in Fig. 8. Note that the residues proximal to the chromophore directly impact its geometry. The corresponding sequence logos for each dataset are shown in Fig. 8A by comparing active site sequences for all proteins and for respective groups in Fig. 8B-D.

515
516
517
Figure 8: Alternation in the equilibrium geometry of chromophore due to rearrangement of
polar and hydrophobic interaction within the active site. The standard deviation in bond
lengths was obtained based on hybrid QM/MM calculation. Colour codes used to denote
alternation include green (>0.001 Å), yellow (>0.006 Å), and red (>0.01 Å). The sequence
logo represents the multiple sequence alignment of active site residues within 6 Å of the
chromophore. The height of individual residue at a particular position corresponds to its
relative frequency. The figure panels correspond to all structural models (A), fast kinetic (B),
average kinetic (C), and slow kinetics (D).

The collective analysis of protein models (Fig. 8A) depicts that the least affected bond
lengths (>0.001 Å) are CH$_3$-C8, C8-C9, C9-C9a, and C5a-N5, which are linked to
dimethylbenzimidazole moiety (ring III) of the isoalloxazine ring. On the contrary, the
regions where bond length vary significantly (>0.01 Å) are prevalent across the three rings.
Two bonds with the polar groups, viz. C2-N3 and C4a-C10a are part of ring I, and the
remaining three C5a-C6, C7-C8, and C9a-N10, belonging to hydrophobic groups are
distributed across ring III. Among the bonds constituting an electronegative element (N or O
atom) in the rings I and II show intermediary bond length deviation (>0.006 Å) with C4-C4a
being an exception. However, this pattern differed significantly when the individual groups

were analysed based on their experimental rate kinetics. Substantial differences were found in
the bond length of the isoalloxazine ring for the three kinetics groups.
538
The kinetics and the protein environment at the binding site correlated well with the changes
in the bond lengths of the chromophore (Fig. 8B-D). A significant structural changes were
observed for the chromophores belonging to fast kinetics. Compared to the collective
analysis, the deviation in bond lengths is now altered from intermediate to high deviation
(std. dev.), as reflected by the change in contour colour from yellow to red (Fig. 8A and B).
The bonds concerning the diazabutadiene system (N1=C10a-C4a=N5) and fused rings II and
III (C9a=C5a). The highest perturbation of ring I, in which two more bonds, viz. N1-C2 and
C4-C4a show alteration from intermediate to least (green) and high (red), respectively.
Overall, the isoalloxazine ring shows substantial alterations, implicating that steric constraints
by the active site residues are less onto the chromophore binding site in this group. On the
contrary, the remaining two groups (Fig. 8C and D) reflect the minimum perturbation on the
chromophore. The bonds linked with a ring I show deviation in the group of proteins showing
intermediate kinetics. The bond involving polar elements offers intermediate changes, which
includes N1-C2, N3-C4, C4=O, and C4a-N5, and remains consistent with cumulative
analysis. Likewise, C2-N3 shows significant alteration (>0.01 Å) and is the same as that of
the fast kinetics group. For the slow kinetics group, the isoalloxazine ring has the most
negligible perturbation in bond length as shown by the cumulative analysis. Exceptions
include C8-CH$_3$ and C8-C9 of ring III, which show high and intermediate deviation. Overall,
the bonds involving electron-rich N5 are altered >0.01 Å in the case of fast, >0.006 Å in
intermediate, and >0.001 Å in slow kinetics.
559
Sequence logo analysis with specific chromophore atoms. We extended our analysis for
chromophore's atom-wise interaction with active site residues (Fig. 9). The occurrence of
amino acids interacting with particular atoms of the chromophore is shown as a sequence
logo, where the height of the letter is proportional to its frequency. The average distance
between N3, O2, and O4 for H-bond interactions (ring I) and C8methyl, C7methyl, C6, C5a,
C9, and C9a (ring III) are shown in Fig. 5. The list also includes hydrophobic interaction
mediates through N5 (ring III). Particularly for H-bonding, Asn and Glu are key players,
which is also reflected in Fig. 8. However, their differential impact has been clearly
understood from the QM/MM simulations.
569

Figure 9: Sequence logo analysis for amino acids interacting with individual atoms of chromophore. The analysis from three kinetics group are shown for (A) fast, (B) intermediate, and (C) slow. N3, O2, and O4 analysis represent the H-bonding interaction while the remaining atoms describes the hydrophobic interaction.

The C8M region of isoalloxazine's ring III is highly occupied with several amino acids. The interaction occupancy is similar in fast and intermediate, whereas it is strongly dominated by Ile and is occupied with fewer amino acids. On the contrary, the interacting amino acids near C7M are significant for slow kinetics proteins compared to the other two groups. For the C6 atom, strong hydrophobic restraints are present in proteins with fast kinetics and are minimum for the intermediate. The adjacent atom C5a shows the involvement of Cys residue, which, however, is occupied with other residues only in the case of the slow group. The hydrophobic interaction near the reaction centre N5 was found to be equivalent for the fast and intermediate groups, with a slight difference in the frequency of amino acids. On the other side (near the ribityl chain), the C9 atom interacts only with Ile in the fast group. At the same time, positional restraints are reflective in the other two groups with different amino acids, such as Cys, Leu, Ile, and Met. The C9a atom is also occupied with Cys interaction, especially in the intermediate group. Notably, this atom is occupied with another hydrophobic residue in the fast and slow groups. Thus, the variable hydrophobic interactions near the chromophore are critical for alternating equilibrium bond length. Later they translate it into photocycle kinetics, which provides crucial information for designing synthetic LOV constructs.

Conclusion

The translational utility of the LOV domain as sensors is limited due to a need for sufficient information on the role of the local protein environment in the photocycle kinetics. Several mutational studies in the past provide valuable information concerning the factors that perturb the Flavin-Cys adduct decay kinetics. Typically, the presence of a polar group for mediating

electrostatic interaction, hydrophobic side chain residues in the proximity of chromophore, and water molecule accessibility are influencing features. However, instead of focusing on individual residues, analysis of residual clusters around chromophore could provide a strong lead for constructing synthetic proteins.

In this work, we revisited the local protein environment crucial for the photosensory behaviour in LOV proteins. The analysis constitutes individual residue-wise interaction and a residual community network with the chromophore binding site. Herein, our approach is to find patterns of protein networks correlating with the experimental photocycle kinetics. Notably, we limited our structural analysis to the dark-state conformational models of LOV protein. One reason is their wide occurrence in monomeric conformation in the dark state. In contrast, the light-activate state in certain classes of proteins attains a dimeric conformation linking with their effector protein domain.

Our structural analysis emphasises the role of hydrogen bonding near ring I and the hydrophobic environment surrounding ring III, which cooperates with the chemical nature of the isoalloxazine ring. The role of Asn, Leu, Gln, and water molecules are rendered common among the LOV protein having fact photocycle kinetics. For other kinetic groups, the role of Val, Leu, and Ile in governing hydrophobic interaction and Arg in mediating salt-bridge with phosphate backbone remains common. However, the absence of shorter sidechain residue is responsible for a longer decay time constant. The dynamic correlation network was elucidated based on the normal mode analysis of proteins. Network analysis highlights that the link between the α-helices and β-sheets is critical for restricting water accessibility near N5 of Flavin and affecting the kinetics. Similar cross-talk between the Fα helix with the Hβ-Iβ loop and adjacent β-sheets is crucial in restricting water accessibility near N5 of Flavin and affect the kinetics. In other proteins, similar cross-talk between Fα helix with Gβ-Hβ and Hβ-Iβ region is crucial. region is crucial in other proteins. Notably, the role of Aβ, Bβ, and Cα are non-significant in fast and intermediate proteins.

On the contrary, tight packing of the Bβ-Cα region is responsible for slow kinetics. The local environment's perturbative effect on Flavin's equilibrium geometry correlates well with the experimental time constant. Based on quantum mechanics, we found a significant difference in the bond length change of chromophore. Overall, the analytical approach present in this work is worthwhile for designing synthetic LOV proteins. However, future studies comprising the larger structural dataset can render a more accurate correlation between protein networks and photocycle kinetics.

Acknowledgements

This research was supported by Department of Science and Technology (DST) – Science and Engineering Research Board (SERB), Government of India (SRG/2022/000858) and INUP, MeitY (5(1)/2021-NANO) to RKK. The author also thanks RnDOPs, IIT Guwahati for a research grant (Project reference no. 2022-040113000301).

Declaration of interests
The authors declare that there is no conflict of interest.

Supporting Information
All supplementary data are available in PDF.

References:

1. Ye, H. & Fussenegger, M. Optogenetic medicine: Synthetic therapeutic solutions precision-guided by light. *Cold Spring Harb Perspect Biol* **11**, (2019).
2. Lan, T.-H., He, L., Huang, Y. & Zhou, Y. Optogenetics for transcriptional programming and genetic engineering. *Trends in Genetics* (2022) doi:10.1016/J.TIG.2022.05.014.
3. McCue, A. C. & Kuhlman, B. Design and engineering of light-sensitive protein switches. *Curr Opin Struct Biol* **74**, 102377 (2022).
4. Seifert, S. & Brakmann, S. LOV Domains in the Design of Photoresponsive Enzymes. *ACS Chemical Biology* vol. 13 1914–1920 Preprint at https://doi.org/10.1021/acschembio.8b00159 (2018).
5. Buckley, A. M., Petersen, J., Roe, A. J., Douce, G. R. & Christie, J. M. LOV-based reporters for fluorescence imaging. *Current Opinion in Chemical Biology* vol. 27 Preprint at https://doi.org/10.1016/j.cbpa.2015.05.011 (2015).
6. Manoilov, K. Y., Verkhusha, V. v. & Shcherbakova, D. M. A guide to the optogenetic regulation of endogenous molecules. *Nature Methods* vol. 18 1027–1037 Preprint at https://doi.org/10.1038/s41592-021-01240-1 (2021).
7. Huala, E. *et al.* Arabidopsis NPH1: A protein kinase with a putative redox-sensing domain. *Science (1979)* **278**, (1997).
8. Harper, S. M., Neil, L. C. & Gardner, K. H. Structural basis of a phototropin light switch. *Science (1979)* **301**, (2003).
9. Möglich, A., Ayers, R. A. & Moffat, K. Structure and Signaling Mechanism of Per-ARNT-Sim Domains. *Structure* vol. 17 Preprint at https://doi.org/10.1016/j.str.2009.08.011 (2009).
10. Crosson, S. & Moffat, K. Structure of a flavin-binding plant photoreceptor domain: Insights into light-mediated signal transduction. *Proc Natl Acad Sci U S A* **98**, (2001).
11. Pudasaini, A., El-Arab, K. K. & Zoltowski, B. D. LOV-based optogenetic devices: Light-driven modules to impart photoregulated control of cellular signaling. *Frontiers in Molecular Biosciences* vol. 2 Preprint at https://doi.org/10.3389/fmolb.2015.00018 (2015).
12. Zoltowski, B. D., Vaccaro, B. & Crane, B. R. Mechanism-based tuning of a LOV domain photoreceptor. *Nat Chem Biol* **5**, 827–834 (2009).
13. Zoltowski, B. D., Nash, A. I. & Gardner, K. H. Variations in protein-flavin hydrogen bonding in a light, oxygen, voltage domain produce non-Arrhenius kinetics of adduct decay. *Biochemistry* **50**, (2011).

14. Kawano, F., Aono, Y., Suzuki, H. & Sato, M. Fluorescence imaging-based high-
throughput screening of fast- and slow-cycling LOV proteins. *PLoS One* **8**, (2013).
15. Zayner, J. P. & Sosnick, T. R. Factors that control the chemistry of the LOV domain
photocycle. *PLoS One* **9**, (2014).
16. Zoltowski, B. D. & Crane, B. R. Light activation of the LOV protein vivid generates a
rapidly exchanging dimer. *Biochemistry* **47**, (2008).
17. Swartz, T. E. *et al.* The Photocycle of a Flavin-binding Domain of the Blue Light
Photoreceptor Phototropin. *Journal of Biological Chemistry* **276**, (2001).
18. Raffelberg, S., Mansurova, M., Gärtner, W. & Losi, A. Modulation of the photocycle
of a LOV domain photoreceptor by the hydrogen-bonding network. *J Am Chem Soc*
133, (2011).
19. Yamamoto, A., Iwata, T., Tokutomi, S. & Kandori, H. Role of Phe1010 in light-
induced structural changes of the neo1-LOV2 domain of Adiantum. *Biochemistry* **47**,
(2008).
20. Chan, R. H. & Bogomolni, R. A. Structural water cluster as a possible proton acceptor
in the adduct decay reaction of oat phototropin 1 LOV2 domain. *Journal of Physical
Chemistry B* **116**, (2012).
21. Freddolino, P. L., Gardner, K. H. & Schulten, K. Signaling mechanisms of LOV
domains: New insights from molecular dynamics studies. *Photochemical and
Photobiological Sciences* **12**, 1158–1170 (2013).
22. Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res* **36**, 5–9
(2008).
23. Losi, A. & Gärtner, W. The evolution of flavin-binding photoreceptors: An ancient
chromophore serving trendy blue-light sensors. *Annual Review of Plant Biology* vol.
63 Preprint at https://doi.org/10.1146/annurev-arplant-042811-105538 (2012).
24. Losi, A. & Gärtner, W. Solving Blue Light Riddles: New Lessons from Flavin-binding
LOV Photoreceptors. *Photochemistry and Photobiology* vol. 93 Preprint at
https://doi.org/10.1111/php.12674 (2017).
25. Mandalari, C., Losi, A. & Gärtner, W. Distance-tree analysis, distribution and co-
presence of bilin- and flavin-binding prokaryotic photoreceptors for visible light.
Photochemical and Photobiological Sciences **12**, (2013).
26. Losi, A., Mandalari, C. & Gärtner, W. The Evolution and Functional Role of Flavin-
based Prokaryotic Photoreceptors. in *Photochemistry and Photobiology* vol. 91 (2015).
27. Edgar, R. C. MUSCLE: A multiple sequence alignment method with reduced time and
space complexity. *BMC Bioinformatics* **5**, (2004).
28. Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics
Analysis Version 11. *Mol Biol Evol* **38**, (2021).
29. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v5: An online tool for
phylogenetic tree display and annotation. *Nucleic Acids Res* **49**, (2021).
30. Bienert, S. *et al.* The SWISS-MODEL Repository-new features and functionality.
Nucleic Acids Res **45**, (2017).
31. Wang, H. *et al.* LOVTRAP: An optogenetic system for photoinduced protein
dissociation. *Nat Methods* **13**, (2016).
32. Song, S. H. *et al.* Modulating LOV domain photodynamics with a residue alteration
outside the chromophore binding site. *Biochemistry* **50**, (2011).
33. Morozenko, A. & Stuchebrukhov, A. A. Dowser++, a new method of hydrating
protein structures. *Proteins: Structure, Function and Bioinformatics* **84**, (2016).
34. Maier, J. A. *et al.* ff14SB: Improving the Accuracy of Protein Side Chain and
Backbone Parameters from ff99SB. *J Chem Theory Comput* **11**, 3696–3713 (2015).

35. Grimme, S., Ehrlich, S. & Goerigk, L. Effect of the Damping Function in Dispersion
Corrected Density Functional Theory. *J Comput Chem* **32**, 1456–1465 (2011).
36. Kossmann, S. & Neese, F. Efficient structure optimization with second-order many-
body perturbation theory: The RIJCOSX-MP2 method. *J Chem Theory Comput* **6**,
2325–2338 (2010).
37. Neese, F. Software update: the ORCA program system, version 4.0. *Wiley Interdiscip*
Rev Comput Mol Sci **8**, 4–9 (2018).
38. Metz, S., Kästner, J., Sokol, A. A., Keal, T. W. & Sherwood, P. ChemShell-a modular
software package for QM/MM simulations. *Wiley Interdiscip Rev Comput Mol Sci* **4**,
101–110 (2014).
39. Deb, A., Grewal, R. K., Roy, S. & Mitra, D. Residue interaction dynamics in
Vaucheria aureochrome1 light-oxygen-voltage: Bridging theory and experiments.
Proteins: Structure, Function and Bioinformatics **88**, (2020).
40. Grant, B. J., Skjærven, L. & Yao, X. Q. The Bio3D packages for structural
bioinformatics. *Protein Science* **30**, (2021).
41. Siemers, M. *et al.* Bridge: A Graph-Based Algorithm to Analyze Dynamic H-Bond
Networks in Membrane Proteins. *J Chem Theory Comput* **15**, (2019).
42. Hassan, S., Töpel, M. & Aronsson, H. Ligand Binding Site Comparison — LiBiSCo
— a web-based tool for analyzing interactions between proteins and ligands to explore
amino acid specificity within active sites. *Proteins: Structure, Function and*
Bioinformatics **89**, (2021).
43. Kottke, T., Heberle, J., Hehn, D., Dick, B. & Hegemann, P. Phot-LOV1□: Photocycle
of a Blue-Light Receptor Domain from the Green Alga Chlamydomonas reinhardtii.
Biophys J **84**, 1192–1201 (2003).
44. Iwata, T., Nozaki, D., Tokutomi, S. & Kandori, H. Comparative investigation of the
LOV1 and LOV2 domains in adiantum phytochrome3. *Biochemistry* **44**, (2005).
45. Oide, M. *et al.* Blue light– excited LOV1 and LOV2 domains cooperatively regulate
the kinase activity of full-length phototropin2 from arabidopsis. *Journal of Biological*
Chemistry **293**, (2018).
46. Endres, S. *et al.* Structure and function of a short LOV protein from the marine
phototrophic bacterium Dinoroseobacter shibae. *BMC Microbiol* **15**, (2015).
47. Arinkin, V. *et al.* Structural determinants underlying the adduct lifetime in the LOV
proteins of Pseudomonas putida. *FEBS Journal* **288**, (2021).
48. Mitra, D., Yang, X. & Moffat, K. Crystal structures of aureochrome1 LOV suggest
new design strategies for optogenetics. *Structure* **20**, (2012).
49. Nash, A. I. *et al.* Structural basis of photosensitivity in a bacterial light-oxygen-
voltage/ helix-turn-helix (LOV-HTH) DNA-binding protein. *Proc Natl Acad Sci U S A*
108, (2011).
50. Conrad, K. S., Bilwes, A. M. & Crane, B. R. Light-induced subunit dissociation by a
light-oxygen-voltage domain photoreceptor from Rhodobacter sphaeroides.
Biochemistry **52**, (2013).
51. Möglich, A. & Moffat, K. Structural Basis for Light-dependent Signaling in the
Dimeric LOV Domain of the Photosensor YtvA. *J Mol Biol* **373**, (2007).
52. Weber, A. M. *et al.* A blue light receptor that mediates RNA binding and translational
regulation. *Nat Chem Biol* **15**, (2019).
53. Rivera-Cancel, G., Ko, W. H., Tomchick, D. R., Correa, F. & Gardner, K. H. Full-
length structure of a monomeric histidine kinase reveals basis for sensory regulation.
Proc Natl Acad Sci U S A **111**, (2014).

54.   Lokhandwala, J. *et al.* Structural biochemistry of a fungal LOV domain photoreceptor
reveals an evolutionarily conserved pathway integrating light and oxidative stress.
Structure **23**, (2015).

55.   Heintz, U. & Schlichting, I. Blue light-induced LOV domain dimerization enhances
the affinity of aureochrome 1a for its target DNA sequence. *Elife* **5**, (2016).

56.   Kalvaitis, M. E., Johnson, L. A., Mart, R. J., Rizkallah, P. & Allemann, R. K. A
Noncanonical Chromophore Reveals Structural Rearrangements of the Light-Oxygen-
Voltage Domain upon Photoactivation. *Biochemistry* **58**, (2019).

57.   Pudasaini, A. *et al.* Kinetics of the LOV domain of ZEITLUPE determine its circadian
function in arabidopsis. *Elife* **6**, (2017).

58.   Rinaldi, J. *et al.* The β-scaffold of the LOV domain of the brucella light-activated
histidine kinase is a key element for signal transduction. *J Mol Biol* **420**, (2012).

59.   Zoltowski, B. D. *et al.* Conformational switching in the fungal light sensor vivid.
Science (1979) **316**, (2007).

60.   Zoltowski, B. D., Nash, A. I. & Gardner, K. H. Variations in protein-flavin hydrogen
bonding in a light, oxygen, voltage domain produce non-Arrhenius kinetics of adduct
decay. *Biochemistry* **50**, 8771–8779 (2011).

61.   Ataka, K., Hegemann, P. & Heberle, J. Vibrational spectroscopy of an algal Phot-
LOV1 domain probes the molecular changes associated with blue-light reception.
Biophys J **84**, (2003).

62.   Zayner, J. P. & Sosnick, T. R. Factors that control the chemistry of the LOV domain
photocycle. *PLoS One* **9**, (2014).

63.   Christie, J. M. *et al.* Structural tuning of the fluorescent protein iLOV for improved
photostability. *Journal of Biological Chemistry* **287**, (2012).

64.   Nash, A. I., Ko, W. H., Harper, S. M. & Gardner, K. H. A conserved glutamine plays a
central role in LOV domain signal transmission and Its duration. *Biochemistry* **47**,
(2008).

65.   Nozaki, D. *et al.* Role of Gln1029 in the photoactivation processes of the LOV2
domain in Adiantum phytochrome3. *Biochemistry* **43**, (2004).

66.   Circolone, F. *et al.* Structural basis for the slow dark recovery of a full-length LOV
protein from Pseudomonas putida. *J Mol Biol* **417**, (2012).

67.   Yee, E. F. *et al.* Signal transduction in light-oxygen-voltage receptors lacking the
adduct-forming cysteine residue. *Nat Commun* **6**, (2015).

68.   Pennacchietti, F. *et al.* The dark recovery rate in the photocycle of the bacterial
photoreceptor YtvA is affected by the cellular environment and by hydration. *PLoS*
One **9**, (2014).

69.   Lokhandwala, J. *et al.* A native threonine coordinates ordered water to tune light-
oxygen-voltage (LOV) domain photocycle kinetics and osmotic stress signaling in
trichoderma reesei ENVOY. *Journal of Biological Chemistry* **291**, (2016).

70.   Freddolino, P. L., Dittrich, M. & Schulten, K. Dynamic switching mechanisms in
LOV1 and LOV2 domains of plant phototropins. *Biophys J* **91**, 3630–3639 (2006).

71.   Tian, H., Trozzi, F., Zoltowski, B. D. & Tao, P. Deciphering the Allosteric Process of
the Phaeodactylum tricornutum Aureochrome 1a LOV Domain. *Journal of Physical*
Chemistry B **124**, (2020).

72.   Nakajima, H., Kobayashi, I., Adachi, Y. & Hisatomi, O. Transmission of light signals
from the light-oxygen-voltage core via the hydrophobic region of the β-sheet surface in
aureochrome-1. *Sci Rep* **11**, (2021).

73.   Hepp, S. *et al.* An Optogenetic Tool for Induced Protein Stabilization Based on the
Phaeodactylum tricornutum Aureochrome 1a Light–Oxygen–Voltage Domain. *J Mol*
Biol **432**, (2020).

74.   Ganguly, A., Thiel, W. & Crane, B. R. Glutamine Amide Flip Elicits Long Distance
Allosteric Responses in the LOV Protein Vivid. *J Am Chem Soc* **139**, (2017).
75.   Zhou, H., Zoltowski, B. D. & Tao, P. Revealing hidden conformational space of lov
protein VIVID through rigid residue scan simulations. *Sci Rep* **7**, (2017).
76.   Kollar, J. & Frecer, V. How accurate is the description of ligand–protein interactions
by a hybrid QM/MM approach? *J Mol Model* **24**, (2018).
843