Artificial light at night alters morphology, phenology, and reproductive capacity

in an annual herb

Lu Xiao[1,†], Shuo Wang[1,†], Ayub M. O. Oduor[1,2,3], Zhihui Wang[4], Hongxiang Zhang[1],

Yanjie Liu[1*]

[1]*State Key Laboratory of Black Soils Conservation and Utilization, Northeast Institute*

of Geography and Agroecology, Chinese Academy of Sciences, Changchun, 130102,

China

[2]*Department of Applied Biology, Technical University of Kenya, P.O Box*

52428-00200, Nairobi, Kenya.

[3]*Shandong Key Laboratory of Eco-Environmental Science for Yellow River Delta,*

Shandong University of Aeronautics, Binzhou, Shandong, China

[4]*Heilongjiang Bayi Agricultural University, Daqing, 163319, China*

†*These authors contributed equally to this work.*

Email of corresponding author: liuyanjie@iga.ac.cn

CRediT authorship contribution statement

YL conceived the idea and designed the long-term grassland ALAN manipulation

experiment. SW and YL conceived the idea of the present experiment and designed it.

SW, ZW, and LX performed the experiment. SW and YL analyzed the data. SW and

LX wrote the first draft of the manuscript, with major inputs from YL, and further

inputs from AO, ZW, LX, and HZ.

Funding

We acknowledge the financial support from the International Partnership Program of

Chinese Academy of Sciences (044GJHZ2023023FN) and the National Natural

Science Foundation of China (41971069).

Acknowledgments

We thank the Erguna Forest-Steppe Ecotone Ecosystem Research Station, Institute of Applied Ecology, Chinese Academy of Sciences for providing the experimental field site and practical assistance with the experiment setup. We thank Xue Zhang, Zhengkuan Lu for their help in the experiment setup.

Abstract

The rapid global expansion of artificial light at night (ALAN) has spurred growing interest in its ecological impact on plant life. However, the effects of low-intensity ALAN on plants in their natural habitats remain largely unexplored, particularly concerning the influence of morphological and phenological responses on the overall fitness of these plants. We conducted a field experiment using the annual herbaceous plant *Elsholtzia densa* as a model species to assess the effects of ALAN on plant morphology, reproductive phenology, and reproductive capacity. The results indicate that ALAN increased the specific leaf area and elongated the top inflorescences, but it resulted in a reduction of secondary branches and a decrease in the proportion of individuals with undeveloped top inflorescences. Additionally, ALAN induced a shift in biomass allocation toward the aboveground parts of plants. It also accelerated the onset of budding, blooming, fruiting, and seed maturity by 3.4 to 6.2 days and caused a decrease in the number of fruity inflorescences. This study suggests that ALAN can significantly affect plant morphology, reproductive timing, and potentially the fitness of plants. While ALAN induces potentially adaptive changes in leaf area and biomass allocation, it may also disrupt plant-pollinator interactions and negatively impact plant reproductive capacity.

Keywords: Artificial light at night, Phenological shifts, Light pollution, Plant traits□ Ecological impact

1. Introduction

To adapt to the cyclical changes in environmental light due to Earth's rotation, a wide range of organisms, including algae, plants, and animals, have evolved molecular

circadian clocks (Hölker et al. 2010; Yerushalmi and Green 2009). These clocks have

reduced competition and increased coexistence among sympatric species by enabling

organisms to optimize their physiological processes in synchrony with the diurnal cycle

(Gaston et al. 2013; Krittika and Yadav 2019; Patke et al. 2020; Yerushalmi and Green

2009; Zhao et al. 2021). However, the natural darkness of Earth's night, which has

persisted for hundreds of billions of years, has been rapidly diminished in recent

decades by the widespread use of artificial light at night (ALAN) across vast

geographical areas (Falchi et al. 2016; Gaston 2018; Irwin 2018; Kyba et al. 2017). At a

global scale, satellite data show that artificially lit outdoor areas expanded by 2.2%

annually, with a 1.8% yearly increase in radiance intensity between 2012 and 2016

(Kyba et al. 2017). This rapid global increase of ALAN has led to light pollution, which

has been shown to impact various species critically and potentially threaten ecosystems

(Irwin 2018), making it a defining feature of the Anthropocene epoch. Consequently,

there has been a growing interest in exploring the ecological effects of ALAN within

the field of global change biology and urban ecology, particularly regarding its impacts

on species behavior and ecosystem services (Davies and Smyth 2018; Hölker et al.

2021; Lockett et al. 2022).

A recent meta-analysis revealed that most studies on the impacts of altered

photoperiod caused by ALAN have historically focused on animal behavior,

physiology, and life history (Sanders et al. 2021). However, increasing attention is now

being directed toward the effects of ALAN on plants. Numerous studies on urban trees,

ornamental garden plant species, and crops have demonstrated that ALAN can

significantly affect plant physiological processes, including photosynthesis,

respiration, and stomatal conductance (Haque et al. 2015; Kim et al. 2015; Kwak et al.

2018; Kwak et al. 2017; Segrestin et al. 2021; Singhal et al. 2019). ALAN has also been

found to influence plant morphology, affecting traits such as leaf size, stem elongation,

and biomass allocation (Abonyo and Oduor 2024; Cieraad et al. 2023; Lockett et al.

2022). Moreover, ALAN can significantly impact plant phenology, particularly

flowering time and plant-pollinator interactions (Abonyo and Oduor 2024; Bennie et al.

2018; Ffrench-Constant et al. 2016; Giavi et al. 2021; Liao et al. 2014; Meng et al.

2022). Despite these advances, the ecological consequences of ALAN on native flora in

natural ecosystems remain understudied compared to urban and agricultural contexts.

Therefore, significant knowledge gaps persist regarding the long-term effects of ALAN

on plant community composition, biodiversity, and ecosystem functioning in natural

habitats (Gaston et al. 2015; Secondi et al. 2019). Addressing these gaps should be a

research priority to inform conservation strategies and mitigate ALAN's potential

negative impacts on natural ecosystems.

Given that plants rely on light as a crucial resource and signaling mechanism for

regulating fundamental processes such as photoperiod responses, their growth and

development are likely to be sensitive to ALAN (Bennie et al. 2016; Gaston et al. 2015;

Singhal et al. 2019). Plants perceive light signals via specialised photoreceptors,

including phytochromes, cryptochromes, and phototropins, which play pivotal roles in

regulating growth and development (Quail 2002). The distinct spectral and temporal

characteristics of ALAN may interfere with these photoreceptor-mediated pathways,

disrupting the normal functioning of light-dependent processes (Heinen 2021; Meravi

and Kumar Prajapati 2020). Furthermore, ALAN has the potential to disrupt circadian

rhythms, which are essential for synchronising physiological processes with daily and

seasonal light cycles. This disruption may alter the timing of gene expression and

metabolic processes, leading to misalignment between the endogenous circadian clock

and the external environment (Bordage et al. 2016; Fankhauser and Staiger 2002;

Webb 2003). Such misalignment can have detrimental effects on plant growth,

development, and fitness (Dodd et al. 2015; Pierik and de Wit 2014).

ALAN may also influence hormonal pathways that control plant growth and

reproduction. For instance, Briggs (2006) hypothesized that low-intensity ALAN,

which has a spectral composition different from direct sunlight, might be perceived by

plants as a shaded environment. This perception can trigger shade avoidance responses,

such as stem elongation and reduced branching, which are commonly mediated by

enhanced auxin signaling and gibberellin production (Jiang et al. 2020; Pierik and de

Wit 2014). These hormonal changes can lead to altered biomass allocation patterns,

with plants investing more resources in vertical growth at the expense of leaf and root

development (Hertel et al. 2021; Speißer et al. 2021). Recent studies have indeed

shown that low-intensity ALAN, such as street lighting, can emit photosynthetically

active radiation that affects the growth and resource allocation in vascular plants

(Speißer et al. 2021) and algae (Diamantopoulou et al. 2021). While some plants may

benefit from extended photoperiods under ALAN, others may experience reduced

fitness due to the disruption of circadian rhythms and the metabolic costs associated

with prolonged growth (Cathey and Campbell 1975; Gaston et al. 2013). The potential

effects of ALAN on functional traits, particularly leaf function and resource allocation

in natural plant populations, require further investigation to elucidate the underlying

physiological mechanisms and their ecological consequences.

Understanding the impact of ALAN on plant flowering and other reproductive

processes is critical, as ALAN-induced shifts in phenology may lead to mismatches

between plants and their pollinators, thereby posing direct threats to the reproductive

success of entomophilous plants. Research using satellite data (Zheng et al. 2021) and

phenological observation data (Meng et al. 2022) suggests that ALAN might lead to an

earlier start of the growing season. However, the specific effects of ALAN on flowering

are variable, ranging from negative to neutral to positive (Abonyo and Oduor 2024;

Bennie et al. 2018; Cathey and Campbell 1975), which highlights the complex and

context-dependent nature of ALAN's influence. Such phenological shifts can lead to

pollinator mismatches, reducing pollination success and seed production, ultimately

impacting dependent species (Giavi et al. 2021; Heinen 2021). ALAN may impact

plant-pollinator networks by altering the timing of flowering and the availability of

floral resources, which in turn can affect the abundance and diversity of pollinators

(Knop et al. 2017; MacGregor et al. 2015). Furthermore, ALAN-induced changes in

plant phenology and resource allocation may have cascading effects on plant-herbivore

interactions (Bennie et al. 2015; Grenis and Murphy 2019) and plant-plant interactions,

such as competition (Abonyo and Oduor 2024).These alterations in species

interactions can potentially reshape the structure and composition of plant

communities (Bennie et al. 2016; Bennie et al. 2018; Hölker et al. 2021; Speißer et al.

2021). Given the close link between reproductive traits and fitness (Violle et al.

2007) , ALAN's capacity to alter these traits could directly affect the viability of

natural plant populations. Therefore, to fully understand the ecological consequences

of ALAN on plant reproduction in the wild, research should prioritize field experiments

over laboratory or greenhouse studies, allowing for a more accurate assessment of plant

responses in their natural environments and within the context of their ecological

communities (Secondi et al. 2019).

This study investigated the effects of artificial light at night (ALAN) on the fitness

of the widespread annual herb *Elsholtzia densa* Benth., with a particular focus on

morphological traits, reproductive phenology, and reproductive capacity of the species.

We tested the following predictions: (1) ALAN increases specific leaf area of plant; (2)

ALAN enhances plant height and total biomass; (3) ALAN advances the onset of

budding, blooming, and fruiting as a result of extended photoperiod and disruption of

circadian rhythms; and (4) ALAN increases both the size and number of inflorescences

due to changes in vegetative growth and reproductive investment. These hypotheses

reflect the increasing recognition that ALAN can interfere with light signaling

pathways, alter photoperiod-sensitive processes, and affect internal resource

distribution in plants. By examining these responses in a natural setting, the study aims

to elucidate the mechanisms by which ALAN influences plant performance, thereby

contributing to a more comprehensive understanding of its ecological impacts on plant

fitness and population dynamics.

2. Materials And Methods

2.1. Study sites and species

This study is part of an ongoing long-term ALAN manipulation experiment conducted in a temperate continental monsoon climate grassland in Inner Mongolia, China (50.17°N, 119.39°E, 523 m a.s.l.). This grassland experiences a mean annual precipitation of 375 mm and a mean annual temperature of –3 °C (Liu et al. 2021) . The broader experiment encompasses 40 square plots, each 4 m², arranged in four combinations of two light pollution levels (ALAN vs. no-ALAN) and two animal-exclusion levels (open cage vs. closed cage), with ten replicates for each combination. The present study specifically utilized 20 plots from the larger experiment, employing only the no-animal-exclusion treatment (open cage) and the two light pollution levels.

To ensure homogeneity in environmental conditions across all plots, we selected a relatively uniform area within the grassland, characterized by similar soil texture, moisture levels, and nutrient availability. The soil was initially tilled, and clonal propagules were removed to homogenize species composition. On April 18, 2021, artificial communities were re-established using seeds from 54 common herbaceous species collected in 2020 from various local grassland sites, ensuring a diverse representation of populations and genetic diversity. The seeds were mixed and randomly distributed across all plots, maintaining a consistent species proportion. All species included were capable of independent propagation without human intervention. We chose the widespread annual herb, *E. densa*, as our model organism to assess the impacts of ALAN on plant morphology, reproductive phenology, and reproductive capacity. Its dominance in our experimental plots, where it exhibited a relative cover of 33.78 ± 3.0% in 2021 (Supplementary Data Fig. S1), alongside its broad distribution

from East Afghanistan to Southeast Siberia and China (POWO database;

https://powo.science.kew.org), made it an ideal candidate for studying the effects of

light pollution on common plant species.

2.2. Set up of the artificial-light-at-night experiment

The ALAN treatment was implemented using LED spotlights (10 W, DC 12–24 V, IP

66, cool white 6500 K; Shenzhen Mingkeming Lighting Co., Ltd., China) mounted on

metal frames 2 meters above the center of each plot. These lights emit

photosynthetically active radiation (PAR; Supplementary Data Fig. S2). To minimize

lateral light radiation and prevent interference between plots, each spotlight was

encased in a cubic lampshade ($30 \times 30 \times 30$ cm) made of stainless steel, and plots were

spaced 2-3 meters apart. To replicate the intensity of light pollution typically

experienced at ground level, we covered each lampshade with two layers of white

nylon netting and one layer of black light-shading netting. This setup resulted in a

consistent light intensity of 21.15 ± 0.30 lx, comparable to that found under streetlights

(Bennie et al. 2016; Speißer et al. 2021). This value represents the spatiotemporal

average per plot, based on measurements taken at five randomly selected points on the

ground within each plot. While some variability existed, statistical analyses confirmed

that light intensity remained significantly different between ALAN and No-ALAN

treatments. The lamps for the ALAN treatment were regulated by a photoelectric

switch (XT KG-F, Zhejiang Xintuo New Energy Co., Ltd., China), which automatically

turned the lights on when ambient light was below 30 lux and off when above this

threshold during the growing season. Control plots (no-ALAN) also had metal frames

and lampshades made of stainless steel covered with nylon netting but lacked LED

spotlights. Light intensity measurements in control plots using a SpectraPen LM500

spectroradiometer (Photon Systems Instruments) yielded zero readings, indicating

levels below the device's detection limit (< 0.1 lux).

2.3. Trait measurements

We measured 14 traits related to morphology, reproductive phenology, and

reproductive capacity of *E. densa* from the onset of budding on July 6, 2021, to the first

frost on September 23, 2021 (Table 1). Ten of these traits were measured

non-destructively on 10 randomly selected and marked individuals of *E. densa* within a

1×1 m subplot at the center of each plot. From July 6 to September 23, we monitored

the marked *E. densa* individuals daily to record the onset of budding, blooming,

fruiting, and seed maturity. The onset of budding was defined as the appearance of the

first bud, blooming as the emergence of the first purple inflorescence, fruiting as the

initial swelling of the inflorescence, and seed maturity as the presence of the first

dehiscent inflorescence, following the methodology of Dorji et al. (2013). At the end of

the growing season (September 23), we measured branch number and reproductive

height of each marked individual plant. Branches were classified as primary if arising

directly from the main stem, and as secondary if originating from the primary branches,

like the approach used by Li et al. (2016). To assess the impact of ALAN on

reproductive capacity, we recorded the number of fruiting inflorescences

(inflorescences bearing seeds) per individual and the number of individuals with

undeveloped top inflorescences (remaining at the bud stage until the end of the growing

season). In this study, we used the number of fruiting inflorescences as a proxy for

reproductive capacity in *E. densa* due to the challenges associated with directly

measuring seed production in the field. Premature seed fall makes it difficult to

accurately quantify the total number of seeds produced by individual plants. The

number of fruiting inflorescences, however, is a more reliable and practical measure

that can be assessed in the field without the risk of underestimating reproductive output

due to seed loss. The number of fruiting inflorescences often correlates with seed

production (Kudo and Harder 2005; Rodger et al. 2021), serving as a proxy for

reproductive capacity and offering insights into the impact of ALAN on plant fitness.

This approach is supported by studies showing that fruiting inflorescences can be a

reliable indicator of reproductive capacity, especially when direct seed production is

challenging to measure (Kudo and Harder 2005; Waser and Ollerton 2006).

Additionally, noting a potential difference in top inflorescence length between ALAN

and no-ALAN plots (Fig. S3), we measured this trait and calculated the proportion of

individuals with undeveloped top inflorescences in each plot as: the number of

individuals with undeveloped top inflorescence / 10 marked individuals per plot.

We conducted destructive sampling to measure four additional traits. To assess the

response of specific leaf area to ALAN, we collected ten mature, fully expanded, and

healthy leaves from five unmarked *E. densa* individuals adjacent to the central subplot

within each plot on August 27. Each individual contributed two leaves. We scanned the

entire leaves using a CanoScan LiDE 300 electronic scanner (Canon Inc., Lake

Success, NY, USA) and calculated total leaf area with ImageJ software (version 1.53g).

Subsequently, the leaves were dried at 60 °C for 72 hours and weighed. Specific leaf

area was calculated for each plot as the total leaf area divided by the total dry leaf

biomass of the corresponding plot. On the same day (August 27), we also measured the

aboveground, inflorescence, and total biomass of *E. densa* under both ALAN

treatments. For this, we randomly selected an additional 3-6 individuals adjacent to

each subplot, resulting in 45 individuals for ALAN plots and 52 for no-ALAN plots.

We harvested the belowground parts, inflorescences, and other aboveground parts

separately for each of the 97 individuals. All materials were dried at 60°C for 72 hours

and weighed. From this data, we calculated the total biomass (belowground parts +

inflorescences + other aboveground parts), aboveground biomass fraction (sum of

inflorescences and other aboveground parts divided by total biomass), and

inflorescence biomass fraction (inflorescences divided by total biomass) for each *E.*

densa individual. It is important to note that both leaf function and biomass allocation

measurements were taken during the vigorous growth stage (August 27), when some

inflorescences were blooming but not yet senescing. This timing was chosen to capture

trait values representative of the plant's peak physiological performance. As

demonstrated in previous studies, the timing of trait measurements can substantially

influence observed patterns and their responses to environmental conditions

(Famiglietti et al. 2024). Given the extended reproductive period of *E. densa*

(approximately 50 days), the relatively small ALAN-induced reproductive phenology

shift (3-6 days) likely had only minimal impact on the observed differences in leaf

functionality and biomass allocation. Sampling too early could have missed important

developmental effects of ALAN on plant traits, while sampling too late might have

introduced confounding effects due to senescence-related trait decline.

2.4. Statistical analyses

To investigate the effects of ALAN on various plant traits, we fitted a series of

Bayesian multilevel models using the *brm* function from the *brms* package (Bürkner

2017) in R v.4.0.2 (R Core Team 2020). These models accounted for the hierarchical

structure of the data and the non-independence of repeated measurements. A total of 14

traits were analyzed, with ALAN included as a fixed effect and the specific trait as the

response variable in each model. Prior to selecting the appropriate model for each trait,

we conducted exploratory data analysis to assess the distribution pattern. Histograms

were used to visually inspect the distribution of each variable. The number of fruity

inflorescences, primary branches, and secondary branches were modeled using

negative binomial distributions. The proportion of individuals with undeveloped top

inflorescences was modeled using a binomial distribution. Plant height and total

biomass were modeled with gamma distributions, while the above-ground and

inflorescence biomass fractions were modeled using beta distributions. The remaining

traits, including specific leaf area (square root transformed), budding date, blooming

date, fruiting date, seed maturity date, and length of top inflorescence were modeled

using Gaussian distributions. To account for the non-independence of individuals

within plots, plot identity was included as a random effect in all models, except for

those analyzing specific leaf area and the proportion of individuals with undeveloped

top inflorescences, which were measured at the plot level. All models used the default

priors provided by the *brms* package and were run with four independent chains, each

consisting of 6000 iterations, including 2000 warm-up iterations. Model fit was

assessed using the *pp_check* function from *brms*, which compares model-generated

data with the observed data via visual inspection of density and scatter plots (Gabry et

al. 2019). A model was considered adequate if the simulated data closely approximated

the observed data in terms of shape and spread. To select the optimal model for each

trait, we used the *loo* function from the *brms* package, which implements leave-one-out

cross-validation (LOO-CV) to estimate the expected log predictive density and the

Widely Applicable Information Criterion (WAIC) (Vehtari et al. 2017). The model

with the lowest WAIC was selected as the optimal model for each trait, indicating the

best balance between model fit and complexity. The significance of the ALAN

treatment was assessed based on the credible intervals of the posterior distribution.

Specifically, the ALAN treatment was deemed significant when the 95% credible

interval of the posterior distribution did not include zero, and marginally significant if

the 90% credible interval did not overlap zero.

3. Results

3.1. Effects of ALAN on morphological traits

Two of the six morphological traits of *E. densa* were significantly affected by ALAN

(Table 2). Specifically, ALAN significantly increased the specific leaf area (+9.2%;

Fig. 2a, Table 2) but decreased the number of secondary branches (−44.3%; Fig. 2c,

Table 2) compared to the no-ALAN treatment. Relative to the no-ALAN treatment,

ALAN also tended to increase the aboveground biomass fraction (+0.68%; Fig. 2f,

Table 2). The number of primary branches, height, and total biomass were not

significantly affected by ALAN treatment (Fig. 2b, d, e; Table 2).

3.2. Effects of ALAN on reproductive phenology

ALAN significantly affected the reproductive phenology of *E. densa* (Table 2).

Specifically, ALAN advanced the averaged Julian days for the onset of budding,

blooming, fruiting, and seed maturity by 3.4, 6.2, 3.4, and 4.6 days, respectively (Fig. 3,

Table 2).

3.3. Effects of ALAN on reproductive capacity

ALAN significantly increased the length of the top inflorescences of *E. densa* (+29.2%;

Fig. 4a, Table 2). Additionally, ALAN significantly decreased the proportion of

individuals with undeveloped top inflorescences (−81.8%; Fig. 4b, Table 2). ALAN

caused a marginally significant reduction in the number of fruity inflorescences relative

to the no-ALAN treatment (−25.2%; Fig. 4d, Table 2). The ALAN treatment did not

have a significant effect on the biomass fraction of inflorescence (Table 2).

4. Discussion

Our study contributes to the growing body of literature on the ecological effects of ALAN on plants (Bennie et al. 2016; Gaston et al. 2017) by demonstrating that ALAN significantly influenced the morphology, reproductive phenology, and reproductive capacity of natural populations of the annual herb *E. densa*. Specifically, ALAN increased specific leaf area and aboveground biomass, while decreasing secondary branch numbers, suggesting a shift in resource allocation towards vertical growth and light capture. Additionally, ALAN advanced the timing of key reproductive stages, including budding, blooming, fruiting, and seed maturity, indicating a disruption in the plant's natural phenological cycles. These findings highlight the multifaceted effects of ALAN on plant development and reproduction.

The morphological responses of *E. densa* to ALAN present a complex picture, possibly reflecting a combination of adaptive strategies. The observed increase in specific leaf area under ALAN conditions aligns with previous research demonstrating that high light exposure promotes leaf expansion and reduced leaf thickness to maximize light capture and photosynthetic efficiency (Poorter et al. 2009). This adaptation, coupled with the observed tendency for increased aboveground biomass allocation under ALAN, suggests that *E. densa* may be strategically prioritizing resources towards leaves and stems at the expense of belowground structures. This shift in resource allocation likely optimizes light capture and photosynthetic performance under extended photoperiod caused by ALAN. Similar shifts in biomass allocation in response to ALAN have been documented in other plant species (Abonyo and Oduor 2024). The reduction in secondary branches, a shade-avoidance response, might also prioritize vertical growth and leaf expansion for maximizing light interception (Gratani 2014). However, the lack of significant effects of ALAN on the number of primary

branches, height, and total biomass highlights the importance of considering multiple

morphological traits when assessing the ecological consequences of ALAN. While

these morphological changes could potentially enhance the competitive ability of *E.*

densa by optimizing light capture, they may also make the plant more susceptible to

environmental stressors such as drought or herbivory. This is because a greater

allocation of resources to above-ground structures might reduce the plant's resilience to

such stressors (Gupta et al. 2020). These findings suggest that while ALAN-induced

changes in resource allocation may provide short-term benefits in terms of light capture,

they could have long-term consequences for plant survival and fitness in natural

ecosystems. Future research should investigate whether these morphological and

phenological changes translate into increased fitness and survival of *E. densa* in natural

ecosystems, or if they have unintended consequences for the plant's overall ecological

performance.

The observed advancement in the budding, blooming, fruiting, and seed maturity

of *E. densa* under ALAN condition are consistent with studies showing that ALAN can

cause earlier leaf-out and flowering in plants, extend the growing season, and alter

reproductive timing (Cieraad et al. 2023; Ffrench-Constant et al. 2016; Meng et al.

2022). This acceleration of reproductive phenology is likely due to the extended

photoperiod and increased light availability under ALAN, which modifies the plants'

perception of seasonal cues and triggers earlier initiation of reproductive processes

(Gaston et al. 2017). Our observations suggest that *E. densa* is a long-day plant,

typically initiating bud differentiation in mid-July under ambient light when day length

is approximately 15.8 hours. The light spectrum emitted by our LED spotlights,

containing red and far-red light (650nm-740nm, Fig. S2), can activate phytochromes,

key light-sensing molecules that influence flowering (Bennie et al. 2016). This

suggests that ALAN likely induces *E. densa* to perceive a shorter night length,

triggering earlier flowering. This early flowering likely extends to subsequent

phenological stages, as evidenced by the synchronized advancement across blooming,

fruiting, and seed maturity observed in our study. While ALAN is primarily associated

with photoperiod extension, other mechanisms, such as hormonal regulation, may also

play a role. For instance, gibberellins, which promote cell elongation and stem growth,

may increase under ALAN conditions and contribute to the observed phenological

shifts (Sena et al. 2024). Similarly, abscisic acid, involved in stress responses and

dormancy, may also be affected, potentially altering the timing of leaf senescence

(Lazzarin et al. 2021). Likewise, jasmonates, which regulate defense and reproductive

development, may be disrupted by ALAN, influencing flowering and fruit maturation

(Lazzarin et al. 2021). Therefore, it is plausible that ALAN-induced hormonal changes,

alongside altered photoperiod perception, contribute to the phenological shifts

observed in *E. densa.*

The phenological shifts observed in *E. densa* under ALAN may have important

ecological consequences, particularly by disrupting the timing of interactions with

pollinators, seed dispersers, and other species (Hölker et al. 2010; Knop et al. 2017).

Although the observed shift in flowering time, ranging from 3 to 6 days in our study,

may appear small relative to the total flowering period (*ca*. 65 days), even slight

mismatches can affect ecological interactions. Diurnal pollinators such as bees and

butterflies may miss peak floral resource availability (Giavi et al. 2021), while

nocturnal pollinators like moths and bats may reduce activity under continuous

illumination (Knop et al. 2017). Similarly, earlier seed maturation may become

decoupled from the peak activity of seed dispersers, potentially lowering dispersal

efficiency and recruitment success (Schupp et al. 2010). Given that the ecological

consequences of ALAN are complex and can extend beyond direct effects on

individual plant species (Giavi et al. 2021; Macgregor et al. 2017), there is a need for

more comprehensive studies to fully understand the ecological consequences of

ALAN.

The present study reveals the complex effects of ALAN on plant reproductive

traits, demonstrating both potential benefits and drawbacks (Fig. 4). ALAN

significantly increased the length of top inflorescences (+29.2%) and decreased the

proportion of individuals with undeveloped top inflorescences (-81.8%), suggesting a

stimulatory effect on inflorescence growth and maturation. These responses may be

driven by extended photoperiod and increased light availability, which are known to

enhance vegetative growth and promote the development of reproductive structures

(Abonyo and Oduor 2024; Ffrench-Constant et al. 2016). However, the marginally

significant reduction in the number of fruiting inflorescences under ALAN (-25.2%)

raises the possibility of trade-offs or constraints on reproductive success. The lack of

change in inflorescence biomass fraction suggests that ALAN influences the timing and

developmental efficiency of reproduction rather than altering overall reproductive

investment. Additionally, ALAN may affect plant physiology and resource allocation,

contributing to the observed shifts in reproductive traits (Heinen 2021). To fully

understand the implications of these changes, future research should investigate how

ALAN affects broader ecological interactions, including plant-pollinator dynamics,

herbivory, and competition within plant communities (Heinen 2021; Knop et al. 2017) .

Such studies would provide a more holistic understanding of ALAN's multifaceted

impacts on plant reproductive success.

Changes in the physical characteristics and reproductive timing of *E. densa* due to

ALAN may lead to reduced fitness. While directly measuring seed production in the

field is difficult because of early seed drop, we used the number of fruiting

inflorescences as a proxy for reproductive capacity and fitness. Our results show that

ALAN led to fewer fruiting inflorescences and a lower proportion of plants with

undeveloped top inflorescences. Although our additional study (data not shown) did

not find ALAN exposure in the parent generation to affect seed germination or seedling

survival, the overall decrease in reproductive capacity suggests a potential negative

impact on *E. densa* fitness. However, the long-term consequences for future

generations remain uncertain. Importantly, the significant reduction in inflorescences

of this nectar-rich plant could lead to food shortages for pollinating insects like bees,

potentially disrupting ecosystem services through a bottom-up effect (Jiang et al. 1996)

5. Conclusions

In conclusion, our study contributes to the growing body of literature on the ecological

effects of ALAN on plants by showing that ALAN can significantly affect plant

morphology, reproductive timing, and potentially the fitness of plants. We observed

both potentially adaptive responses (increased leaf area and altered biomass allocation)

and potential negative impacts (reduced reproductive capacity). These findings have

important implications for plant populations and ecosystem functioning in the context

of global change. As light pollution continues to increase worldwide, the consequences

of ALAN on plants may extend beyond individual-level responses and alter population

dynamics, community structure, and ecosystem processes. To fully understand the

complex direct and indirect consequences of ALAN on plants and ecosystems, future

research should include long-term field experiments that track plant demographic

responses to ALAN across multiple growing seasons. Such studies will provide

valuable insights into the potential cumulative effects of ALAN on plant fitness,

population viability, and ecosystem resilience. Additionally, studies examining

species-specific ALAN responses could help identify which plant species or functional

groups are most vulnerable to light pollution. This information could guide

conservation prioritization efforts and inform management strategies to mitigate the

adverse effects of ALAN on sensitive plant communities.

Data availability

Data will be made available on request.

Declaration of competing interest

We declare we have no competing interests.

References

Abonyo, C.R.K., Oduor, A.M.O., 2024. Artificial night☐time lighting and nutrient

enrichment synergistically favour the growth of alien ornamental plant sp

ecies over co☐occurring native plants. J. Ecol. 112(2), 319-337. https://doi.

org/10.1111/1365-2745.14235.

Bennie, J., Davies, T.W., Cruse, D., Gaston, K.J., Swenson, N., 2016. Ecological

effects of artificial light at night on wild plants. J. Ecol. 104(3), 611-620.

https://doi.org/10.1111/1365-2745.12551.

Bennie, J., Davies, T.W., Cruse, D., Inger, R., Gaston, K.J., 2015. Cascading effe

cts of artificial light at night: resource-mediated control of herbivores in a

grassland ecosystem. Phil. Trans. R. Soc. B 370(1667), 20140131. https://d

oi.org/10.1098/rstb.2014.0131.

Bennie, J., Davies, T.W., Cruse, D., Inger, R., Gaston, K.J., Lewis, O., 2018. Arti

ficial light at night causes top☐down and bottom☐up trophic effects on in

vertebrate populations. J. Appl. Ecol. 55(6), 2698-2706. https://doi.org/10.1

111/1365-2664.13240.

Bordage, S., Sullivan, S., Laird, J., Millar, A.J., Nimmo, H.G., 2016. Organ speci

ficity in the plant circadian system is explained by different light inputs to

the shoot and root clocks. New Phytol. 212(1), 136-149. https://doi.org/10.

1111/nph.14024.

Briggs, W.R., 2006. Physiology of plant responses to artificial lighting.

Bürkner, P.-C., 2017. brms: An R Package for Bayesian Multilevel Models Using

Stan. J. Stat. Softw. 80(1), 1-28. https://doi.org/10.18637/jss.v080.i01.

Cathey, H.M., Campbell, L.E., 1975. Effectiveness of five vision-lighting sources

on photo-regulation of 22 species of ornamental plants. J. Am. Soc. Hort.

Sci. 100(1), 65-71.

Cieraad, E., Strange, E., Flink, M., Schrama, M., Spoelstra, K., 2023. Artificial li

ght at night affects plant–herbivore interactions. J. Appl. Ecol. 60(3), 400-

410. https://doi.org/10.1111/1365-2664.14336.

Davies, T.W., Smyth, T., 2018. Why artificial light at night should be a focus for
global change research in the 21st century. Global Change Biol. 24(3), 87
2-882. https://doi.org/10.1111/gcb.13927.

Diamantopoulou, C., Christoforou, E., Dominoni, D.M., Kaiserli, E., Czyzewski, J.
, Mirzai, N., Spatharis, S., 2021. Wavelength-dependent effects of artificial
light at night on phytoplankton growth and community structure. Proc. R.
Soc. B 288(1953), 20210525. https://doi.org/10.1098/rspb.2021.0525.

Dodd, A.N., Belbin, F.E., Frank, A., Webb, A.A., 2015. Interactions between circ
adian clocks and photosynthesis for the temporal and spatial coordination
of metabolism. Front Plant Sci 6, 245. https://doi.org/10.3389/fpls.2015.002
45.

Dorji, T., Totland, O., Moe, S.R., Hopping, K.A., Pan, J., Klein, J.A., 2013. Plant
functional traits mediate reproductive phenology and success in response t
o experimental warming and snow addition in Tibet. Global Change Biol.
19(2), 459-472. https://doi.org/10.1111/gcb.12059.

Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C.C.M., Elvidge, C.D., Baugh, K., P
ortnov, B.A., Rybnikova, N.A., Furgoni, R., 2016. The new world atlas of
artificial night sky brightness. Sci. Adv. 2(6), e1600377. https://doi.org/10.
1126/sciadv.1600377.

Famiglietti, C.A., Worden, M., Anderegg, L.D.L., Konings, A.G., 2024. Impacts o
f climate timescale on the stability of trait-environment relationships. New
Phytol. 241(6), 2423-2434. https://doi.org/10.1111/nph.19416.

Fankhauser, C., Staiger, D., 2002. Photoreceptors in Arabidopsis thaliana: light pe
rception, signal transduction and entrainment of the endogenous clock. Pla
nta 216(1), 1-16. https://doi.org/10.1007/s00425-002-0831-4.

Ffrench-Constant, R.H., Somers-Yeates, R., Bennie, J., Economou, T., Hodgson,
D., Spalding, A., McGregor, P.K., 2016. Light pollution is associated with
earlier tree budburst across the United Kingdom. Proc. R. Soc. B 283(183
3), 20160813. https://doi.org/10.1098/rspb.2016.0813.

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., Gelman, A., 2019. Visualizat
ion in Bayesian Workflow. J. R. Stat. Soc. A 182(2), 389-402. https://doi.
org/10.1111/rssa.12378.

Gaston, K.J., 2018. Lighting up the nighttime. Science 362(6416), 744-746. https:/
/doi.org/10.1126/science.aau8226.

Gaston, K.J., Bennie, J., Davies, T.W., Hopkins, J., 2013. The ecological impacts
of nighttime light pollution: a mechanistic appraisal. Biol. Rev. 88(4), 91
2-927. https://doi.org/10.1111/brv.12036.

Gaston, K.J., Davies, T.W., Nedelec, S.L., Holt, L.A., 2017. Impacts of Artificial
Light at Night on Biological Timings. Annu Rev. Ecol. Evol. Syst. 48(1),
49-68. https://doi.org/10.1146/annurev-ecolsys-110316-022745.

Gaston, K.J., Visser, M.E., Ho¨lker, F., 2015. The biological impacts of artificial
light at night: the research challenge. Phil. Trans. R. Soc. B 370(1667), 20
140133. https://doi.org/10.1098/rstb.2014.0133.

Giavi, S., Fontaine, C., Knop, E., 2021. Impact of artificial light at night on diurn
al plant-pollinator interactions. Nat. Commun. 12(1), 1690. https://doi.org/1
0.1038/s41467-021-22011-8.

Gratani, L., 2014. Plant Phenotypic Plasticity in Response to Environmental Facto
rs. Advances in Botany 2014, 1-17. https://doi.org/10.1155/2014/208747.

Grenis, K., Murphy, S.M., 2019. Direct and indirect effects of light pollution on t
he performance of an herbivorous insect. Insect Sci. 26(4), 770-776. https:/
/doi.org/10.1111/1744-7917.12574.

Gupta, A., Rico-Medina, A., Cano-Delgado, A.I., 2020. The physiology of plant r
esponses to drought. Science 368(6488), 266-269. https://doi.org/10.1126/sc
ience.aaz7614.

Haque, M.S., Kjaer, K.H., Rosenqvist, E., Ottosen, C.O., 2015. Continuous light i
ncreases growth, daily carbon gain, antioxidants, and alters carbohydrate m
etabolism in a cultivated and a wild tomato species. Front. Plant Sci. 6, 52
2. https://doi.org/10.3389/fpls.2015.00522.

Heinen, R., 2021. A spotlight on the phytobiome: Plant-mediated interactions in a
n illuminated world. Basic Appl. Ecol. 57, 146-158. https://doi.org/10.1016
/j.baae.2021.10.007.

Hertel, M.F., Araújo, H.H., Stolf-Moreira, R., Pereira, J.D., Pimenta, J.A., Bianchi
ni, E., Oliveira, H.C., 2021. Different leaf traits provide light-acclimation r
esponses in two neotropical woody species. Theoretical and Experimental
Plant Physiology 33(4), 313-327. https://doi.org/10.1007/s40626-021-00213-
1.

Hölker, F., Bolliger, J., Davies, T.W., Giavi, S., Jechow, A., Kalinkat, G., Longc
ore, T., Spoelstra, K., Tidau, S., Visser, M.E., Knop, E., 2021. 11 Pressing
Research Questions on How Light Pollution Affects Biodiversity. Front.
Ecol. Evol. 9, 767177. https://doi.org/10.3389/fevo.2021.767177.

Hölker, F., Wolter, C., Perkin, E.K., Tockner, K., 2010. Light pollution as a biodi
versity threat. Trends Ecol. Evol. 25(12), 681-682. https://doi.org/10.1016/j.
tree.2010.09.007.

Irwin, A., 2018. The dark side of light: how artificial lighting is harming the natu
ral world. Nature 553(7688), 268-271. https://doi.org/10.1038/d41586-018-0
0665-7.

Jiang, H., Shui, Z., Xu, L., Yang, Y., Li, Y., Yuan, X., Shang, J., Asghar, M.A.,
Wu, X., Yu, L., Liu, C., Yang, W., Sun, X., Du, J., 2020. Gibberellins m
odulate shade-induced soybean hypocotyl elongation downstream of the m
utual promotion of auxin and brassinosteroids. Plant Physiol. Biochem. 15
0, 209-221. https://doi.org/10.1016/j.plaphy.2020.02.042.

Jiang, Y., Deng, Y., Yang, W., Dang, R., 1996. Anatomic studies of the floral ne
ctaries in Elsholtzia densa Benth. Acta Bot. Boreali-Occidential Sinica 16(
3), 239-244.

Kim, Y.J., Yu, D.J., Rho, H., Runkle, E.S., Lee, H.J., Kim, K.S., 2015. Photosynt
hetic changes in Cymbidium orchids grown under different intensities of n

591   ight interruption lighting. Sci. Hortic. 186, 124-128. https://doi.org/10.1016/
592   j.scienta.2015.01.036.

593 Knop, E., Zoller, L., Ryser, R., gerpe, C., Hörler, M., Fontaine, C., 2017. Artifici
594   al light at night as a new threat to pollination. Nature 548(7666), 206-209.
595   https://doi.org/10.1038/nature23288.

596 Krittika, S., Yadav, P., 2019. Circadian clocks: an overview on its adaptive signif
597   icance. Biol. Rhythm Res. 51(7), 1109-1132. https://doi.org/10.1080/092910
598   16.2019.1581480.

599 Kudo, G., Harder, L.D., 2005. Floral and inflorescence effects on variation in poll
600   en removal and seed production among six legume species. Funct. Ecol. 1
601   9(2), 245-254. https://doi.org/10.1111/j.1365-2435.2005.00961.x.

602 Kwak, M., Je, S., Cheng, H., Seo, S., Park, J., Baek, S., Khaine, I., Lee, T., Jang,
603   J., Li, Y., Kim, H., Lee, J., Kim, J., Woo, S., 2018. Night Light-Adaptati
604   on Strategies for Photosynthetic Apparatus in Yellow-Poplar (Liriodendron
605   tulipifera L.) Exposed to Artificial Night Lighting. Forests 9(2), 74. https:
606   //doi.org/10.3390/f9020074.

607 Kwak, M.J., Lee, S.H., Khaine, I., Je, S.M., Lee, T.Y., You, H.N., Lee, H.K., Jan
608   g, J.H., Kim, I., Woo, S.Y., 2017. Stomatal movements depend on interact
609   ions between external night light cue and internal signals activated by rhyt
610   hmic starch turnover and abscisic acid (ABA) levels at dawn and dusk. Ac
611   ta Physiol. Plant. 39(8), 162. https://doi.org/10.1007/s11738-017-2465-y.

612 Kyba, C.C.M., Kuester, T., Miguel, A.S.d., Baugh, K., Jechow, A., Hölker, F., Be
613   nnie, J., Elvidge, C.D., Gaston, K.J., Guanter, L., 2017. Artificially lit surf
614   ace of Earth at night increasing in radiance and extent. Sci. Adv. 3(11), e1
615   701528. https://doi.org/10.1126/sciadv.1701528.

616 Lazzarin, M., Meisenburg, M., Meijer, D., van Ieperen, W., Marcelis, L.F.M., Ka
617   ppers, I.F., van der Krol, A.R., van Loon, J.J.A., Dicke, M., 2021. LEDs
618   Make It Resilient: Effects on Plant Growth and Defense. Trends Plant Sci.
619   26(5), 496-508. https://doi.org/10.1016/j.tplants.2020.11.013.

Li, F., Chen, B., Xu, K., Gao, G., Yan, G., Qiao, J., Li, J., Li, H., Li, L., Xiao,
X., Zhang, T., Nishio, T., Wu, X., 2016. A genome-wide association study
of plant height and primary branch number in rapeseed (Brassica napus).
Plant Sci. 242, 169-177. https://doi.org/10.1016/j.plantsci.2015.05.012.

Liao, Y., Suzuki, K., Yu, W., Zhuang, D., Takai, Y., Ogasawara, R., Shimazu, T.,
Fukui, H., 2014. Night break effect of LED light with different wavelengt
hs on floral bud differentiation of Chrysanthemum morifolium Ramat 'Jim
ba'and 'Iwa' no hakusen. Environ. Control Biol. 52(1), 45-50. https://doi.or
g/10.2525/ecb.52.45.

Liu, H., Wang, R., Lu, X.T., Cai, J., Feng, X., Yang, G., Li, H., Zhang, Y., Han,
X., Jiang, Y., 2021. Effects of nitrogen addition on plant-soil micronutrie
nts vary with nitrogen form and mowing management in a meadow steppe.
Environ. Pollut. 289, 117969. https://doi.org/10.1016/j.envpol.2021.117969.

Lockett, M.T., Rasmussen, R., Arndt, S.K., Hopkins, G.R., Jones, T.M., 2022. Art
ificial light at night promotes bottom-up changes in a woodland food chai
n. Environ. Pollut. 310, 119803. https://doi.org/10.1016/j.envpol.2022.11980
3.

Macgregor, C.J., Evans, D.M., Fox, R., Pocock, M.J., 2017. The dark side of stre
et lighting: impacts on moths and evidence for the disruption of nocturnal
pollen transport. Global Change Biol. 23(2), 697-707. https://doi.org/10.111
1/gcb.13371.

MacGregor, C.J., Pocock, M.J., Fox, R., Evans, D.M., 2015. Pollination by noctur
nal Lepidoptera, and the effects of light pollution: a review. Ecol. Entomol.
40(3), 187-198. https://doi.org/10.1111/een.12174.

Meng, L., Zhou, Y., Román, M.O., Stokes, E.C., Wang, Z., Asrar, G.R., Mao, J.,
Richardson, A.D., Gu, L., Wang, Y., 2022. Artificial light at night: an un
der-appreciated effect on phenology of deciduous woody plants. PNAS nex
us 1(2), pgac046. https://doi.org/10.1093/pnasnexus/pgac046/6569705.

Meravi, N., Kumar Prajapati, S., 2020. Effect street light pollution on the photosy
nthetic efficiency of different plants. Biol. Rhythm Res. 51(1), 67-75. https:
//doi.org/10.1080/09291016.2018.1518206.

Patke, A., Young, M.W., Axelrod, S., 2020. Molecular mechanisms and physiolog
ical importance of circadian rhythms. Nat. Rev. Mol. Cell Biol. 21(2), 67-
84. https://doi.org/10.1038/s41580-019-0179-2.

Pierik, R., de Wit, M., 2014. Shade avoidance: phytochrome signalling and other
aboveground neighbour detection cues. J. Exp. Bot. 65(11), 2815-2824. htt
ps://doi.org/10.1093/jxb/ert389.

Poorter, H., Niinemets, U., Poorter, L., Wright, I.J., Villar, R., 2009. Causes and
consequences of variation in leaf mass per area (LMA): a meta-analysis. N
ew Phytol. 182(3), 565-588. https://doi.org/10.1111/j.1469-8137.2009.02830.
x.

Quail, P.H., 2002. Photosensory perception and signalling in plant cells: new para
digms? Curr. Opin. Cell Biol. 14(2), 180-188. https://doi.org/10.1016/s0955
-0674(02)00309-5.

R Core Team, 2020. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.

Rodger, J.G., Bennett, J.M., Razanajatovo, M., Knight, T.M., van Kleunen, M., A
shman, T.-L., Steets, J.A., Hui, C., Arceo-Gómez, G., Burd, M., 2021. Wi
despread vulnerability of flowering plant seed production to pollinator decl
ines. Sci. Adv. 7(42), eabd3524. https://doi.org/10.1126/sciadv.abd3524.

Sanders, D., Frago, E., Kehoe, R., Patterson, C., Gaston, K.J., 2021. A meta-anal
ysis of biological impacts of artificial light at night. Nat. Ecol. Evol. 5(1),
74-81. https://doi.org/10.1038/s41559-020-01322-x.

Schupp, E.W., Jordano, P., Gomez, J.M., 2010. Seed dispersal effectiveness revisi
ted: a conceptual review. New Phytol. 188(2), 333-353. https://doi.org/10.1
111/j.1469-8137.2010.03402.x.

Secondi, J., Davranche, A., Théry, M., Mondy, N., Lengagne, T., Isaac, N., 2019.

Assessing the effects of artificial light at night on biodiversity across latit

ude – Current knowledge gaps. Global Ecol. Biogeogr. 29(3), 404-419. htt

ps://doi.org/10.1111/geb.13037.

Segrestin, J., Mondy, N., Boisselet, C., Guigard, L., Lengagne, T., Poussineau, S.,

Secondi, J., Puijalon, S., 2021. Effects of artificial light at night on the le

af functional traits of freshwater plants. Freshwat. Biol. 66(12), 2264-2271.

https://doi.org/10.1111/fwb.13830.

Sena, S., Kumari, S., Kumar, V., Husen, A., 2024. Light emitting diode (LED) li

ghts for the improvement of plant performance and production: A compreh

ensive review. Current Res. Biotechnol. 7, 100184. https://doi.org/10.1016/j.

crbiot.2024.100184.

Singhal, R.K., Kumar, M., Bose, B., 2019. Eco-physiological Responses of Artific

ial Night Light Pollution in Plants. Russ. J. Plant Physiol. 66(2), 190-202.

https://doi.org/10.1134/s1021443719020134.

Speißer, B., Liu, Y., van Kleunen, M., Oduor, A., 2021. Biomass responses of wi

dely and less☐widely naturalized alien plants to artificial light at night. J.

Ecol. 109(4), 1819-1827. https://doi.org/10.1111/1365-2745.13607.

Vehtari, A., Gelman, A., Gabry, J., 2017. Practical Bayesian model evaluation usi

ng leave-one-out cross-validation and WAIC. Statistics and Computing 27,

1413-1432. https://doi.org/10.1007/s11222-016-9696-4.

Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., Garnie

r, E., 2007. Let the concept of trait be functional! Oikos 116(5), 882-892.

https://doi.org/10.1111/j.0030-1299.2007.15559.x.

Waser, N.M., Ollerton, J., 2006. Plant-pollinator interactions: from specialization t

o generalization. University of Chicago Press.

Webb, A.A., 2003. The physiology of circadian rhythms in plants. New Phytol. 1

60(2), 281-303. https://doi.org/10.1046/j.1469-8137.2003.00895.x.

Yerushalmi, S., Green, R.M., 2009. Evidence for the adaptive significance of circ
adian rhythms. Ecol. Lett. 12(9), 970-981. https://doi.org/10.1111/j.1461-02
48.2009.01343.x.

Zhao, K., Ma, B., Xu, Y., Stirling, E., Xu, J., 2021. Light exposure mediates circ
adian rhythms of rhizosphere microbial communities. ISME J. 15(9), 2655-
2664. https://doi.org/10.1038/s41396-021-00957-3.

Zheng, Q., Teo, H.C., Koh, L.P., 2021. Artificial Light at Night Advances Spring
Phenology in the United States. Remote Sens. 13(3), 399. https://doi.org/1
0.3390/rs13030399.

Table 1. The 14 traits measured in this study.

Category	Trait	Measuring time	Replication No.
Morphology	*Specific leaf area*	27 August, 2021	20
	Primary branch number	23 September, 2021	200
	Secondary branch number	23 September, 2021	200
	Aboveground biomass fraction	27 August, 2021	97
	Height	23 September, 2021	200
	Total biomass	27 August, 2021	97
Reproductive phenology	*Budding date*	From 6 July to	200
	Blooming date	23 September, 2021	200
	Fruiting date	(before plant budding	200
	Seed maturity date	to the start of frost)	200
Reproduction capacity	*Length of top inflorescence*	23 September, 2021	200
	Proportion of individuals with undeveloped top inflorescence	23 September, 2021	20
	Inflorescence biomass fraction	27 August, 2021	97
	Fruity inflorescence number	23 September, 2021	200

Table 2. Output of Bayesian models testing the effects of artificial light at night (ALAN) on each of the 14 traits observed in this study. Asterisks (*) and daggers (†) indicate terms whose 95% and 90% credible intervals (CIs) do not overlap with zero, respectively.

No.	Trait	brm family	Term	Fixed effect			Random effect
				Estimate	95% CIs	90% CIs	Estimate (95% CIs)
1	*Specific leaf area* (n = 20)	Gaussian	Intercept (No-ALAN) *	14.087	(13.707, 14.473)	(13.767, 14.404)	—
			ALAN *	0.630	(0.067, 1.181)	(0.171, 1.084)	—
2	*Primary branch number* (n = 200)	Negative binomial	Intercept (No-ALAN) *	2.132	(2.009, 2.252)	(2.03, 2.232)	0.14 (0.04, 0.25)
			ALAN	0.060	(-0.110, 0.237)	(-0.08, 0.203)	—
3	*Secondary branch number* (n = 200)	Negative binomial	Intercept (No-ALAN) *	1.192	(0.763, 1.659)	(0.832, 1.575)	0.26 (0.01, 0.78)
			ALAN *	-0.637	(-1.322, -0.006)	(-1.183, -0.111)	—
4	*Aboveground biomass fraction* (n = 97)	Beta	Intercept (No-ALAN) *	2.375	(2.32, 2.44)	(2.326, 2.425)	—
			ALAN †	0.081	(-0.007, 0.170)	(0.007, 0.155)	—
5	*Height* (n = 200)	Gamma	Intercept (No-ALAN) *	0.024	(0.021, 0.027)	(0.021, 0.026)	0.01 (0.00, 0.02)
			ALAN	0.001	(-0.004, 0.004)	(-0.003, 0.003)	—
6	*Total biomass* (n = 97)	Gamma	Intercept (No-ALAN) *	0.330	(0.234, 0.438)	(0.25, 0.418)	0.13 (0.08, 0.21)
			ALAN	0.006	(-0.139, 0.153)	(-0.114, 0.124)	—
7	*Budding date* (n = 200)	Gaussian	Intercept (No-ALAN) *	205.513	(204.369, 206.629)	(204.575, 206.446)	0.49 (0.02, 1.35)
			ALAN *	-3.588	(-4.954, -2.243)	(-4.713, -2.464)	—
8	*Blooming date* (n = 200)	Gaussian	Intercept (No-ALAN) *	229.190	(228.101, 230.243)	(228.313, 230.059)	1.29 (0.39, 2.46)
			ALAN *	-5.494	(-7.673, -3.457)	(-7.294, -3.785)	—
9	*Fruiting date* (n = 200)	Gaussian	Intercept (No-ALAN) *	238.063	(237.033, 239.111)	(237.216, 238.896)	0.96 (0.09, 2.07)
			ALAN *	-3.335	(-5.107, -1.583)	(-4.814, -1.885)	—
10	*Seed maturity date* (n = 200)	Gaussian	Intercept (No-ALAN) *	255.622	(254.520, 256.745)	(254.722, 256.548)	0.81 (0.04, 2.12)
			ALAN *	-4.526	(-6.609, -2.462)	(-6.259, -2.784)	—
11	*Length of top inflorescence* (n = 200)	Gaussian	Intercept (No-ALAN) *	5.373	(4.942, 5.798)	(5.015, 5.728)	0.23 (0.01, 0.60)
			ALAN *	0.836	(0.255, 1.429)	(0.348, 1.329)	—
12	*Proportion of individuals with undeveloped top inflorescence* (n = 20)	Binomial	Intercept (No-ALAN) *	-2.105	(-2.775, -1.509)	(-2.658, -1.598)	—
			ALAN *	-1.920	(-3.786, -0.480)	(-3.421, -0.693)	—
13	*Inflorescence biomass fraction* (n = 97)	Beta	Intercept (No-ALAN) *	-1.809	(-2.015, -1.610)	(-1.977, -1.645)	0.22 (0.04, 0.41)
			ALAN	0.020	(-0.264, 0.299)	(-0.215, 0.254)	—
14	*Fruity inflorescence number* (n = 200)	Negative binomial	Intercept (No-ALAN) *	3.208	(2.966, 3.446)	(3.007, 3.404)	0.32 (0.17, 0.51)
			ALAN †	-0.285	(-0.628, 0.050)	(-0.567, -0.008)	—

Figure captions

Fig. 1. (a) Top view of the entire experimental setup, showcasing the 20 open cages (no animal exclusion) used in this study (marked with borders), both during the day and at night (lower left inset). (b) Close-up photograph of a representative open cage plot.

Fig. 2. Effects of artificial light at night (ALAN) on specific leaf area (a), primary branch number (b), secondary branch number (c), plant height (d), total biomass (e), and aboveground biomass fraction (f). Asterisks (*) indicate that the 95% credible intervals of the estimated effects of ALAN do not include zero, indicating a statistically significant difference. Daggers (†) represent 90% credible intervals that do not include zero, suggesting a marginally significant difference. Details on each trait are provided in Table 2.

Fig. 3. Effects of artificial light at night (ALAN) on the onset of different reproductive phenophases (budding, blooming, fruiting, and seed maturity), expressed as Julian days for the year 2021. Asterisks (*) indicate that the 95% credible intervals of the estimated effects of ALAN do not include zero, indicating a statistically significant difference. Details on each trait are provided in Table 2. Data points in different colors represent measurements from various plots, as detailed in Figure 2.

Fig. 4. Effects of artificial light at night (ALAN) on length of top inflorescence (a), proportion of individuals with undeveloped top inflorescence (b), inflorescence biomass fraction (c), and fruity inflorescence number (d). Asterisks (*) indicate that the 95% credible intervals of the estimated effects of ALAN do not include zero, indicating a statistically significant difference. Daggers (†) represent 90% credible intervals that do not include zero, suggesting a marginally significant difference. Information on each trait can be found in Table 2. Data points in different

742  colors represent measurements from various plots, as detailed in Fig. 2.

743 **Fig. 1**

746 **Fig. 2**

750   **Fig. 3**

753 **Fig. 4**

757