1

Distinct stem-like cell populations facilitate functional regeneration of the *Cladonema* medusa tentacle

[1,2] Sosuke Fujita**,** [3] Mako Takahashi, [3] Gaku Kumano, [2] Erina Kuranaga,
[1] Masayuki Miura, and [1,2,4] Yu-ichiro Nakajima*

1. Graduate School of Pharmaceutical Sciences, The University of Tokyo
2. Graduate School of Life Sciences, Tohoku University
3. Asamushi Research Center for Marine Biology, Graduate School of Life Sciences, Tohoku University
4. Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp
Tel: +81-3-5841-4863

Running title: Distinct stem-like cells enable tentacle regeneration in jellyfish
Main text: 6593 words (Abstract: 190 words)
Figures: 7
Supplementary information (Figures: 11)
Keywords: regeneration, blastema, non-bilaterians, cnidarians, jellyfish, tentacle, i-cell

Abstract

Blastema formation is a crucial process that provides a cellular source for regenerating tissues and organs. While bilaterians have diversified blastema formation methods, its mechanisms in non-bilaterians remain poorly understood. Cnidarian jellyfish, or medusae, represent early-branching metazoans that exhibit complex morphology and possess defined appendage structures highlighted by tentacles with stinging cells (nematocytes). Here we investigate the mechanisms of tentacle regeneration, using the hydrozoan jellyfish *Cladonema pacificum*. We show that proliferative cells accumulate at the tentacle amputation site and form a blastema composed of cells with stem cell morphology. Lineage tracing experiments indicate that repair-specific proliferative cells in the blastema are distinct from resident stem cells. We further demonstrate that resident stem cells control nematogenesis and tentacle elongation during both homeostasis and regeneration as homeostatic stem cells, while repair-specific proliferative cells preferentially differentiate into epithelial cells in the newly formed tentacle, analogous to lineage-restricted stem/progenitor cells observed in salamander limbs. Taken together, our findings reveal a regeneration mechanism that utilizes both resident homeostatic stem cells and repair-specific proliferative cells, which in conjunction efficiently enable functional appendage regeneration, and provide novel insight into the diversification of blastema formation across animal evolution.

Introduction

Regeneration, the phenomenon of re-forming missing body parts, is widespread among metazoans. Common regenerative processes include wound closure immediately after injury; formation of the cellular source, or blastema, that reconstitutes the lost tissue; and regrowth of the tissue that integrates different cellular behaviors such as proliferation and differentiation [1]. Among these regenerative responses, blastema formation is a critical step that can distinguish regenerative and non-regenerative systems. Indeed, most mammalian and avian species do not form blastema upon injury while wound closure normally occurs [2]. Understanding blastema formation mechanisms in highly regenerative animals may therefore help us identify the necessary elements to potentially improve our regenerative abilities.

Blastema can be defined as an undifferentiated cellular mass that contains cells with mitotic capacity and appears after damage such as amputation [3,4]. Accumulating evidence has suggested that methods of blastema formation vary among animals with high regenerative abilities [5]. For example, in planarians, pluripotent stem cells called neoblasts that are distributed throughout the body, are recruited to the injury site to produce blastema [6]. Salamanders can regenerate adult limbs upon amputation via blastema formation, but the underlying cellular mechanisms vary among species: in the axolotl, tissue-specific stem cells contribute to blastema, while in the newt, muscle fibers dedifferentiate into proliferative progenitor cells to behave as blastema [7]. During zebrafish caudal fin regeneration, both osteoblast-derived dedifferentiated cells and resident progenitor cells migrate to the wound site to form blastema [8,9]. These studies support the idea that the supply of blastema has diversified across the animal kingdom, whose members utilize resident stem/progenitor cells and/or repair-specific *de novo* proliferative cells to reconstruct lost body parts. While mechanisms of blastema formation have been extensively studied in the limited number of regenerative animals, little is known about their evolutionary characteristics: which elements are acquired as lineage-specific novelties and which are widely conserved within highly regenerative species? In particular, the current understanding of blastema formation largely relies on bilaterian models, and thus the mechanisms of blastema formation outside of bilaterians remain poorly understood.

Among various regeneration contexts, appendage regeneration is widely
observed in bilaterians (e.g. amphibian limbs and fish fins), and is suitable for
understanding the evolution of regenerative processes [3,10]. The bilaterian
program of blastema formation during appendage regeneration is associated
with repair-specific stem/progenitor cells [7,9,11]. Yet, to elucidate the
evolutionary history of blastema formation programs, their mechanisms must be
studied in early-branching metazoans in addition to bilaterian models.
Cnidarians (corals, sea anemones, hydroids, and jellyfish) are among the
earliest branching metazoans, composed of two major groups Anthozoa and
Medusozoa, forming a diverse phylum that contains over 10,000 species, and
stand at a unique phylogenetic position as the sister group to bilaterians (Fig
1A). While cnidarians display considerably divergent morphologies and life
cycles, represented by the polyp and medusa stages; their common traits are a
diploblastic radially-symmetric body along with tentacles as the well-defined
appendages that bear the stinging cells, nematocytes (cnidocytes) [12,13].
Although regenerative potential may vary among the group, most documented
cnidarian species are capable of regenerating lost tissues and organs, and
some can even regenerate their entire body [14,15].
Among cnidarians, polyp-type animals such as *Hydra*, *Hydractinia,* and
Nematostella have been utilized as models to understand the mechanism of
whole-body regeneration; including patterning, body axis formation, and
mechanical responses [16–18]. After mid-gastric bisection in *Hydra*, cell
proliferation of the stem cells, or i-cells, around the wound site is accelerated by
Wnt3a produced from dying cells to generate blastema [19]. Upon decapitation
of the colony polyp *Hydractinia*, i-cells remotely located in the body column
migrate to the injury site to form blastema [20]. During head regeneration of the
sea anemone *Nematostella*, two adult stem-like cell populations, fast-cycling
cells in the body wall epithelium and slow-cycling cells in the mesenteries,
migrate toward the amputation site to form blastema [21]. These studies
suggest that the recruitment of resident stem cells to the injury site is a
prerequisite for blastema formation after amputation of the body. However, it
remains unclear whether repair-specific proliferative cells constitute the cellular
source of blastema during cnidarian regeneration.
In contrast to the sessile polyp stage, medusae, commonly called
jellyfish, exhibit a more complex body structure that includes multiple types of
muscle such as smooth and striated muscles, developed neural networks, and

distinct organs and appendages, which together enable free swimming and
avoidance behaviors [22,23]. While most sexually-reproducing medusae do not
clonally propagate like polyps, medusae can still regenerate various organs and
appendages (e.g., manubrium, tentacle, umbrella, gonad, eye) after injury and
reconstitute *de novo* structures after their removal [24–30]. A recent report
using the hydrozoan jellyfish *Clytia* has suggested that, upon removal of the
entire manubrium, i-cells and differentiated cells localized in neighboring organs
migrate to the damage site through the canals, contributing to *de novo*
manubrium regeneration [27]. Despite the importance of proliferating blastema
cells in highly regenerative animals, the detailed mechanisms of organ and
appendage regeneration in medusae, particularly the mechanism of blastema
formation and its specific role are largely unknown.
The hydrozoan jellyfish *Cladonema pacificum* is an emerging jellyfish
model that has been utilized to study development and physiology [31–33].
Because *Cladonema* allows for easy lab maintenance with its high spawning
rate, it enables monitoring all life cycle stages and exploring organismal
responses to different stimuli. The genus *Cladonema* is morphologically
characterized by branched tentacles at the medusa stage (Fig 1B) [31,34–36].
The *Cladonema* medusa tentacle is primarily composed of bi-layered epithelial
tissues (epidermis and gastrodermis) that include muscle fiber, also called
epitheliomuscular cells [23]: i-cells, neurons, and nematocytes are located in the
epidermal layer while neurons and gland cells are located in the gastrodermal
layer. Resident stem cells, i-cells, are localized at the basal side of the tentacle
(tentacle bulb), and are thought to give rise to progenitors and differentiated
cells during normal development and homeostasis [24,37]. The continuous
growth and branching potential of the medusa tentacle makes it an attractive
model to understand the mechanisms of appendage morphogenesis, growth,
and regeneration in cnidarians.
In this study, we investigate the cellular mechanism of appendage
regeneration using the *Cladonema* medusa tentacle. Establishing the
Cladonema tentacle as an organ that can efficiently and functionally regenerate
upon amputation, we show that highly proliferative cells accumulate at the injury
site within 24 hours to behave as blastema. Lineage tracing experiments using
nucleoside analogs as well as dye labeling reveal that most blastema cells are
not derived from resident stem cells but rather appear locally after damage. We
further identify the role of blastema cells as the principal cellular source of the

epithelium in the newly formed tentacle, while resident stem cells contribute to
nematogenesis and tissue elongation during both homeostasis and
regeneration. These results suggest the existence of two distinct proliferative
cell populations: blastema as repair-specific proliferative cells and resident stem
cells as homeostatic stem cells, both of which collectively enable functional
tentacle regeneration. In a broader context, our findings highlight the
diversification of blastema formation in non-bilaterian systems, providing an
evolutionary insight into animal regeneration.
185
186

Materials and Methods

Animal cultures and surgical manipulations

Cladonema pacificum (female strain 6W) medusae were used for this research.
The medusae were maintained in plastic containers (V-type container, V-7, V-8,
and V-9; AS ONE) at 22 ℃ in artificial sea water (ASW). ASW was prepared
using SEA LIFE (Marin Tech) dissolved in tap water with chlorine neutralizer
(Coroline off; GEX Co. ltd) (24 p.p.t). The medusae were fed Vietnamese brine
shrimp (A&A Marine LLC).
Before surgical manipulation, the medusae were anesthetized with 7%
$MgCl_2$ in deionized water for 2 min in a petri dish (Falcon, #351008). Surgical
manipulations were performed with micro scissors (Teraoka, NY33408). After
manipulations, the medusae were quickly returned to ASW. When the tentacle
was amputated with the bulb left intact, the medusae were fed 1 day before
manipulation and were starved after amputation. When the tentacle was
amputated along with the bulb, the medusae were fed 1 day before
manipulation and were fed 3 times a week afterward.
Bright field pictures of medusae were taken with a LEICA S8APO
microscope with a Nikon digital camera (D5600) or a Zeiss AXIO Zoom.V16.
206

Immunofluorescence

208 The medusae were anesthetized with 7% $MgCl_2$ in deionized water for 5
209 min and fixed at room temperature (R.T.) for 1 h or overnight at 4 ℃ with 4%
210 paraformaldehyde (PFA) in ASW. After fixation, the samples were washed three
211 times (10 min each) with PBS containing 0.1% Triton X-100 (0.1% PBT). The
212 samples were incubated in primary antibodies in 0.1% PBT overnight at 4 ℃.

The antibodies used were mouse anti-acetylated tubulin (1:500; Sigma-Aldrich, T6793), rabbit anti-FMRFamide (1:1000; Wako, 20091), rabbit anti-Phospho-Histone H3 (Ser10) (1:500; Upstate, 06–570), mouse anti-β-catenin (8E4) (1:100; Sigma-Aldrich, ALX-804-260-C100), and rabbit anti-PKC ζ C20 (1:100; Santa Cruz, sc-216). After primary antibody incubation, the samples were washed three times (10 min each) with 0.1% PBT. The samples were incubated with secondary antibodies (1:500; ALEXA FLUOR 488, 555, 647 Goat anti-mouse IgG, ALEXA FLUOR 488, 555, 647 Goat anti-rabbit IgG; Thermo Fisher Scientific, A28175, A-11008) and Hoechst 33342 (1:500; Thermo Fisher Scientific, H1399) in 0.1% PBT for 1 h in the dark. After three washes (10 min each) in 0.1% PBT, the samples were mounted on slides with 70% glycerol. Nuclear and poly-γ-glutamate were stained with DAPI (1:250; Invitrogen, D1306), and actin fibers were stained with Alexa 546 phalloidin (1:400; Invitrogen, A22283) in 0.1% PBT for 1 h. Confocal images were collected through a Zeiss LSM 880 confocal microscope. Image processing (color display) and quantification were performed using ImageJ/Fiji software.

EdU labeling

The medusae were incubated with 150 µM 5-ethynyl-2'-deoxyuridine (EdU) (EdU kit; Invitrogen, C10337) in ASW for 1 h or 24 h. In chase labeling, the medusae were washed with ASW at least twice (up to three times, depending on amount). In pulse labeling, after EdU treatment, the medusae were anesthetized with 7% $MgCl_2$ in deionized water and fixed 4% PFA in ASW. After fixation, the samples were washed and incubated with a EdU reaction cocktail (1× reaction buffer, $CuSO_4$, Alexa Fluor azide 488 or 647, and 1× reaction buffer additive; all included in EdU kit; Invitrogen, C10337 or C10340) for 30 min in the dark. After the EdU reaction, the samples were washed and incubated with Hoechst 33342 in 0.1% PBT for 30 min.

In combination with antibody staining, EdU reaction was performed after incubation with the secondary antibody. To combine EdU labeling with fluorescent *in situ* hybridization (FISH), the EdU reaction was performed after the incubation with Cy5-tyramide solution.

EdU and BrdU dual labeling

The medusae were incubated with 150 µM EdU in ASW for 1h or 24 h before amputation, depending on the experiments. Just after incorporation of EdU, the

medusae were washed in ASW three times, and tentacles were amputated. At

24 hpa, the medusae were incubated with 2.5 mM BrdU in ASW for 1 h. The

medusae were washed out in ASW three times.

The medusae were anesthetized with 7% $MgCl_2$ for 5 min and fixed with

4% PFA in ASW. The samples were washed three times with 0.1% PBT and

incubated with a EdU reaction cocktail for 30 min in dark. The samples were

washed three times and treated with 2N HCl for 30 min and then washed three

times and incubated in primary antibodies in 0.1% PBT overnight at 4 ℃. The

primary antibody used was mouse anti-BrdU (1:100; BD, #347580). The

samples were washed three times and incubated with secondary antibodies

(1:500; ALEXA FLUOR 555 Goat anti-mouse IgG) and Hoechst 33342 in 0.1%

PBT for 1 h in the dark and washed three times.

262   Fluorescent *in situ* hybridization

The following FISH protocol was created from the published protocol [38]. The

sequences of *Nanos1, Nanos2,* and *Piwi* have been previously described [39],

the presumptive sequence of *Minicollagen1* (*Mcol1*) and *Vasa1* were acquired

from the annotation of an RNA-seq result [33]. Purified total RNA was reverse

transcribed into cDNA by PrimeScript II 1st strand cDNA Synthesis Kit (Takara,

6210A) or SMARTer™ RACE cDNA Amplification Kit (Clontech). A cDNA library

was used as a template for PCR (*Nanos1, Nanos2, Piwi and Mcol1*) and 3'

RACE (*Vasa1*). The primer sets used for PCR cloning are as follows: *Nanos1*:

5′-AAGAGACACAGTCATTATCAAGCGA-3′ (forward) and 5′-

AGCACGTAAAATTGGACACGTCG-3′ (reverse), *Nanos2*: 5′-

ACTTCTCCAAAACCTCATGCCGAG-3′ (forward) and 5′-

GAATGGCGGGCGATTTGACATCC-3′ (reverse), *Piwi*: 5′-

CACACAAGAGTTGGACCGGA-3′ (forward) and 5′-

ACCGGCTTATCGATGCAACA-3′ (reverse), *Vasa1*: 5′-

GCCACCCAAAGAAGACAGACAGACAC-3′ (forward for 3′ RACE) and 5′-

CGAAACGACTTGCTGATTTTCTCGCCAG -3′ (nested forward for 3′

RACE*)*, *Mcol1*: 5′-CTCGTCGGTATTGCCCTCTC-3′ (forward) and 5′-

CCAACCTATCGTGGACGTGT-3′ (reverse). PCR products were sub-cloned

into the TAK101 vector (TOYOBO) and RACE product was sub-cloned into the

pGEM-T Easy vector (Promega). The resulting plasmids were used for RNA

probe synthesis with digoxigenin (DIG) labeling mix (Roche, #11277073910) or

fluorescein (FITC) labeling mix (Roche, 11685619910), and T7 (Roche,

#10881767001) or T3 RNA polymerase (Roche, #11031163001) was used,
according to the insert direction.

Medusae were anesthetized 7% $MgCl_2$ for 5 min and fixed overnight at
4 ℃ with 4% PFA in ASW. Briefly, fixed samples were washed three times with
PBS containing 0.1% Tween-20 (PBST), followed by pre-hybridization in
hybridization buffer (HB buffer: 5 × SSC, 50% formamide, 0.1% Tween-20, 50
µg/ml tRNA, 50 µg/ml heparin) at 55 ℃ for 2 h. The samples were hybridized
with HB Buffer containing the antisense probes (final probe concentration: 0.5–1
ng/µL in HB Buffer) at 55 ℃ overnight. The samples were washed twice each
with wash buffer 1 (5 × SSC, 50% formamide, 0.1% Tween-20), wash buffer 2
(2 × SSC, 50% formamide, 0.1% Tween-20), and 2 × SSC. The samples were
then washed with 0.1% PBST and incubated in 1% blocking buffer (1% blocking
reagent [Roche] in Maleic acid) for 1 h. The samples were incubated with anti-
DIG-POD antibodies (1:500; Roche, #11207733910) in 1% blocking buffer
overnight at 4 ℃. The samples were then washed with Tris-NaCl-Tween Buffer
and incubated with Cy5-tyramide solution (TSA Plus Cyanine 5; AKOYA
Biosciences, NEL745001KT) for 10 min. Finally, the samples were washed with
0.1% PBST and incubated with Hoechst 33342 in 0.1% PBST for 30 min in the
dark and washed three times.

For double FISH, DIG for low expression genes and FITC probes for
high expression genes were used (S3E Fig; *Nanos1*-DIG and *Nanos2*-FITC,
S3F Fig; *Mcol1*-DIG and *Nanos2*-FITC). The samples were then washed with
Tris-NaCl-Tween Buffer after incubation with anti-DIG-POD antibodies and
incubated with Cy3-tyramide solution (TSA Plus Cyanine 3; AKOYA
Biosciences, NEL744001KT) for 10 min. The samples were washed with 0.1%
PBST and incubated with 3% H_2O_2 for 15 min. The samples were washed with
0.1% PBST and incubated with anti-Fluorescein-POD antibodies (1:500; Roche,
#11426346910) in 1% blocking buffer overnight at 4 ℃. The samples were then
washed with Tris-NaCl-Tween Buffer and incubated with Cy5-tyramide solution
for 10 min. Finally, the samples were washed with 0.1% PBST and incubated
with Hoechst 33342 in 0.1% PBST for 30 min in the dark and washed three
times.

DiI injection
Medusae were relaxed in 7% $MgCl_2$ on a petri dish. 10 mM CellTracker CM-DiI
(Invitrogen, C7001) was injected into the epidermal layer of the tentacle bulb or

the center of the tentacle, using a micro injector (Eppendorf, Femtojet 4i). The
quartz capillary (Sutter Instrument, QF100-70-10) was pulled by Laser-Based
Micropipette Puller P2000 (Sutter Instrument). After injection, injected tentacles
were amputated. To identify DiI labeled cells, medusae were incubated with 150
µM EdU for 1 h before DiI injection or antibody staining was performed after DiI
injection using anti-β-catenin. To monitor DiI labeled cells during blastema
formation, at 24 hpa, the injected medusae were incubated with 150 µM EdU in
ASW for 1 h. The medusae were relaxed in 7% MgCl$_2$ and fixed in 4% PFA.
After washing in 0.1% PBT, the medusae were incubated with an EdU reaction
cocktail for 30 min in the dark. After the EdU reaction, the samples were
washed and incubated with Hoechst 33342 in 0.1% PBT.
332
X-ray irradiation
The medusae were placed in a V7 plastic container with 6-8 ml of ASW, the
minimum required for their maintenance. X-ray irradiations were performed
using an X-ray machine (mediXtec, MX-160Labo). Medusae were positioned 19
cm from the X-ray source, and a dose of 30 Gy (19.3 minutes at 160 kV 3 mA),
50 Gy (32.2 minutes at 160 kV 3 mA), or 75 Gy (48.3 minutes at 160 kV 3 mA)
was delivered. After irradiation, the medusae were moved to a plastic container
with fresh ASW.
341
Drug treatment
The medusae were incubated with 10 mM hydroxyurea (HU) (085-06653;
Wako, Osaka, Japan) or 30 µM Mitomycin C (139-18711; Wako, Osaka, Japan)
in ASW (ASW only for control). Drug incubation was sustained over 3 days.
Medusae were fed every other day, and the drug solutions were renewed after
feeding.
348
qPCR
Total RNA was purified from the whole tentacles of three medusae with RNeasy
Mini kits (Qiagen). Lysate was treated with Dnase I (Qiagen, #79254) for 15
min at RT. cDNA was synthesized from 200 ng of total RNA by Prime script RT
master Mix (Takara, RR036A). RT-qPCR was performed using TB Green
Premix Ex TaqII (Tli RnaseH Plus) (Takara, RR820L) and a Quantstudio6 Flex
Real-Time PCR system (Thermo Fisher) using *F-actin capping protein subunit*
beta as an internal control [32]. The qPCR primer sets are as follows:

CpCapZbeta: 5'-AAAGAAAGCTGGAGACGGTTCA-3' (Forward), 5'-
GTAGTGGGCATTTCTTCCGC-3' (Reverse), *Nanos1*: 5'-
TTCGTCAAGTGGCAGTCGTG-3' (Forward), 5-
CAAGCCCTGGTACAAACGGA-3' (Reverse).
361

UV laser exposure
Tentacles were amputated at 1 day before UV exposure. At 24 hpa, the
medusae were incubated with Hoechst 33342 (1:250) for 20 min and washed
three times in ASW. After washing, the relaxed medusae were placed on slides
with 7% $MgCl_2$ and enclosed with cover glass. UV laser was exposed to the
blastema region of the amputated tentacle using the photo bleaching application
(50 cycles at 100% 405 laser power) of the Zeiss LSM 880 confocal
microscope. The bleached region is shown with a white dot square (400 x 400
pixels, W x H) in S11B Fig. After UV exposure, the medusae were moved to
ASW in plastic containers and monitored to examine the effect on regeneration.
372

Quantification and statistical analysis
Distribution of cells
Three areas were defined (the basal, middle, and distal regions) by dividing the
tentacle into three equal lengths in confocal images (Figs 2A-2B, 3C-3D and
S2A-S2B). The number of PH3$^+$, EdU$^+$, or *Nanos1*$^+$ cells were counted manually
in each area using the multipoint tool of ImageJ/Fiji.
379

Counting the number of cells
The PH3$^+$, EdU$^+$, or *Nanos1*$^+$ cell ratio (Figs 2D-2F, 4I-4J and S7B-S7C, S7E-
S7F) was measured by counting the total number of cells as well as the PH3$^+$,
EdU$^+$, or *Nanos1*$^+$ cells using ImageJ/Fiji. Each signal$^+$ cell was counted
manually using the multipoint tool of ImageJ/Fiji. Total cell number quantification
was performed as follows: (1) Binarize the signal of Hoechst staining using the
'Threshold' command. (2) Divide continuously adjacent multiple nuclei using the
'Watershed' command. (3) Measure the number of nuclei using the 'Analyze
Particles' command.
Other cell number counting (Figs 3A-3B, 3E-3F, 4C-4F, 5B-5C, 5F-5G,
6B-6C, 6E-6F and S2C, S2E, S4C-S4E, S5A-S5D, S6B-S6C, S8A-S8B, S10,
S11C-S11D) was performed manually using the multipoint tool of ImageJ/Fiji.
392

Measurement of tentacle length and area, signal intensity

The tentacle length (Fig 5K and 6I) was measured using the segment line tool

of ImageJ/Fiji. The tentacle area (S5C-S5E Fig) was recorded as the ROI with

the polygon selection tool and measured by ImageJ/Fiji. The quantification area

was marked with rectangle tool, and the signal intensity was measured by

ImageJ/Fiji (Fig S2D).

Statistics

Statistical analyses were performed using Excel and Graphpad Prism9. Two

tailed *t* tests were used for comparisons between two groups. Significance is

indicated in the figures as follows: *$P \leq 0.05$, **$P \leq 0.05$, ***$P \leq 0.001$, Not

Significant (N.S.): $P > 0.05$. Bar graphs show mean ± standard error. Dots in bar

graphs and boxplots indicate individual values.

Results

For cnidarians, tentacles are essential and common appendages for capturing

prey and defending against predators. Cnidarian polyps such as *Hydra*,

Hydractinia, and *Nematostella* have tentacles in the head region, near the

hypostome, which can restore functional tentacles during head regeneration

[18,20,40]. Similarly, studies using hydrozoan jellyfish have shown that

medusae tentacles can regenerate upon amputation [24,27]. Although

cnidarians commonly exhibit high regenerative capacity for tentacles, the exact

cellular processes and the limitation of regeneration are poorly understood.

To monitor the process of tentacle regeneration and to explore its

responses to different dissections, we utilized the medusa tentacle of the

hydrozoan jellyfish *Cladonema* (Fig 1). In *Cladonema* tentacles, proliferative

cells including stem cells, or i-cells, are localized on the basal side, called the

tentacle bulb, while nematocytes are distributed as clusters on the distal side,

similar to the jellyfish *Clytia* (Figs 1C, 2Ai, 3Ci, and S1A, S3A-S3D, S4Ai, S4Bi)

[24,37,39]. The existence of localized stem cells at the bulb prompted us to

examine tentacle regeneration as a model for appendage regeneration in non-

bilaterians.

***Cladonema* medusa tentacles functionally regenerate upon amputation**

When the tentacle was amputated with the bulb left intact (Fig 1D and 1E),

wound closure finished by 24 hours post-amputation (hpa) (S1B and S1C Fig),

at which point the injury site became smooth surface (Fig 1Di and 1Dii).

Subsequently, tentacles began to elongate distally, and nematocyte clusters,

the spherical structures stained with the mature nematocyte marker poly-γ-

glutamate, formed at the tip of the regenerating tentacle at 48 hpa (Figs 1Diii,

1Fiii, and S1A). The number of nematocyte clusters increased during tentacle

elongation, and these clusters were localized along the distal side at 72 hpa,

which is when the regenerating tentacle branched, a characteristic feature of

the genus *Cladonema* (Fig 1Div and 1E). Notably, these regeneration

processes occurred without feeding, suggesting that tentacle regeneration is an

immediate response for the organism to re-construct the organ that is required

for capturing food.

To investigate the subtleties of tentacle morphology at cellular

resolution during regeneration, we visualized the muscle fibers, mature

nematocytes, and neurons by staining F-actin, poly-γ-glutamate, and

FMRFamide and acetylated tubulin, respectively (Figs 1F, 1G and S1D, S1E).

During the early stages of regeneration (0-24 hpa), wound closure occurred

progressively via the accumulation of supra-cellular actin fibers (Fig 1Fi and

1Fii). Accordingly, muscle fibers regenerated during tentacle elongation (Fig

1Fiii and 1Fiv). While mature nematocytes were rarely found in the bulb or the

injury site until 24 hpa (Fig 1Fi and 1Fii), nematocytes began to accumulate at

the tip of the regenerating tentacle at 48 hpa and formed clusters at 72 hpa (Fig

1Fiii and 1Fiv). The re-formation of neurons on the distal side occurred at 48

hpa and completed at 72 hpa (Figs 1G and S1D, S1E). These results suggest

that the regenerating tentacle is fully functional after 48 hpa. To test this

possibility, we examined whether the regenerating tentacle is capable of

feeding. While the regenerating tentacle was not functional until 24 hpa, more

than half of the regenerating tentacles captured prey at 48 hpa (55%), and this

ratio reached 100% at 72 hpa (Fig 1H and 1I). These results indicate that

Cladonema medusae can fully regenerate a functional tentacle within 2-3 days

of amputation.

While the *Cladonema* medusa tentacle can almost always regenerate

with the bulb left intact, it is unclear whether the tentacle can still regenerate

even after the bulb has been removed (Fig 1Ji). We thus ablated the entire

tentacle, including the bulb, and monitored the regeneration process. After

amputation, the ring canal and the radial canal adhered, and the wound closure
completed at 1 day post-amputation (dpa) (Fig 1Jii). The injured area began to
bulge as early as 5 dpa, and the transparent wound site turned orange, which
reflects the color of its food, *Artemia* (Fig 1Jiii). The *de novo* tentacle bulb
reformed within 1-2 weeks with a nematocyte cluster at the tip, and the small
tentacle gradually elongated (Fig 1Jiv and 1K). Because the rate of successful
regeneration varied dramatically while rearing medusae under the same
conditions and feeding frequency, tentacle bulb regeneration likely depends on
the health condition of the individual medusa (S1F Fig).
We also examined the regenerative potential of isolated tentacles.
While the whole body cannot be regrown from isolated tentacles in *Hydra* or
Nematostella [21,41], polyps in the jellyfish *Aurelia aurita* can regenerate from
isolated tentacles [42]. To examine the regeneration capacity of the isolated
tentacle in *Cladonema*, we cultured *Cladonema* medusa tentacles either with or
without the bulb left intact. Although wound closure successfully occurred
regardless of the existence of the bulb, tissues and organs located to the
proximal side of the tentacle including the umbrella, the radial canal, and the
bulb, did not regenerate, and the isolated tentacle gradually degenerated (S1G
and S1H Fig).
These observations together show that the *Cladonema* medusa tentacle can
functionally regenerate after amputation, but that its organ repair capacity is
restricted to the organism that recognizes the missing tentacle, likely through
the canals, as previously shown in the *Clytia* medusa [27]. For our remaining
studies, we focused our attention on the process of tentacle regeneration with
the bulb intact, which provides a highly reproducible and tractable system.
489
Blastema is formed in the early phase of tentacle regeneration
During organ regeneration, blastema forms at the regenerating tip after wound
closure and contributes to reconstruction of the new tissue as a source of cell
proliferation and differentiation [1,2]. In order to investigate the timing of
blastema formation during *Cladonema* tentacle regeneration, we performed
pulse labeling of EdU, a S-phase marker (Fig 2A) [43]. To understand the
spatial pattern of cell proliferation, we defined three areas in the regenerating
tentacle (the basal, middle, and distal regions) and quantified the distribution of
proliferating cells. Consistent with the localization of stem cells (Figs 3Ci and
S4Ai, S4Bi)[39], cell proliferation in the intact tentacle was mainly detected on

the basal side (Fig 2Ai and 2B). By contrast, cell proliferation in the regenerating tentacle was frequently observed on the distal side near the wound area at 24 hpa (Fig 2Aii and 2B). At 72 hpa, unlike at 24 and 48 hpa, cell proliferation was detected in the middle and on basal side, not on the distal side (Fig 2Aiv and 2B). To investigate the exact timing of the cell proliferation increase at the injury site, we examined the localization of EdU$^+$ cells every four hours immediately after amputation to 24 hpa (S2C Fig). We found that the number of proliferating cells significantly increased at 20 hpa and peaked at 24 hpa (S2D-S2E Fig). We further confirmed the pattern of proliferating cells using the M-phase marker, anti-Phospho-Histone 3 (PH3) antibody. While PH3$^+$ cells were distributed across the basal side before amputation, in the regenerating tentacle, PH3$^+$ cells accumulated on the distal side at 24 hpa, and gradually localized on the basal side starting at 48 hpa (Figs 2D and S2A, S2B), which is consistent with the results of EdU$^+$ cell distribution (Fig 2A and 2B). These results also reflect the fact that differentiated nematocytes accumulate on the distal side at 48 hpa and 72 hpa (Fig 1D and 1F). Together, these observations suggest that cell proliferation is accelerated near the injury site at 24 hpa.

Blastema cells are undifferentiated cell populations that possess a high mitotic capacity during regeneration [44]. We thus investigated proliferative activity in detail during the early phase of tentacle regeneration (Fig 2D-2F). In the regenerating tip, the number of PH3$^+$ and EdU$^+$ cells was about 4 and 20 times greater, respectively, than the corresponding area (amputation site) in the intact tentacle (Fig 2D-2F). Moreover, these PH3$^+$ and EdU$^+$ cell numbers at the regenerating tip were significantly greater than those in the bulb of the intact tentacle where resident stem cells are localized (Fig 2D-2F). Notably, nearly half of the cells in the regenerating tip are EdU$^+$ cells (Fig 2Dii and 2F). Altogether, these results indicate that proliferative activity in the regenerating tip is much higher than that of cells not in the regenerating tip, which provides evidence that blastema cells appear during tentacle regeneration.

To determine whether blastema formation is specific to the location of amputation, we examined the localization of proliferating cells after dissecting tentacles in different ways. When the distal side of the tentacle is amputated, ectopic cell proliferation was observed near the injury site at 24 hpa (Fig 2G). Similarly, when the tentacle was amputated with the bulb, cell proliferation occurred around the injury site where the canals merged at 24 hpa (Fig 2H).

These results suggest that blastema is formed around the injury site regardless
of the location of amputation.
Based on these observations, we hypothesized that blastema is a
prerequisite for proper regeneration of the tentacle. To test this possibility, we
examined whether blastema formation occurs in the isolated tentacle, which
fails to regenerate basal tissue (S1G-S1H Fig). In the isolated tentacle, EdU+
cells were not detected at the injury site at 24 hpa (Fig 2I). These results
together suggest that blastema formation is associated solely with the
regenerating tentacle.
544
Stem cells compose blastema
What cell types constitute blastema? In hydrozoans, i-cells behave as pluri-
/multipotent stem cells that can differentiate into progenitors and differentiated
lineages [45]. In order to confirm the cell types that form blastema, we utilized
the combination of β-catenin antibody staining and nuclear staining, which can
distinguish i-cells, nematocytes, and epithelial cells (Fig 3A). In both *Hydractinia*
and *Cladonema*, i-cells were identified by cytoplasmic signals of β-catenin,
together with the large nucleus-to-cytoplasm ratio and prominent nucleoli
[39,46]. While nematocytes exhibit crescent-shaped nuclei, epithelial cells show
a polygonal shape delineated by the membrane-localized β-catenin (Fig 3Aiii).
Through the quantification of cell types, we found that around 80% of blastema
cells were i-cells, 10% were epithelial cells, and 10% were nematocytes (Fig 3A
and 3B), indicating that most cells in blastema are morphologically i-cells.
To molecularly characterize these blastema cells, we examined the
expression of the stem cell markers, *Nanos1*, *Piwi*, and *Vasa1* by fluorescent *in*
situ hybridization (FISH). Note that, while *Cladonema* possess two *Nanos*
genes, *Nanos1* and *Nanos2, Nanos2+* cells co-express a nematoblast marker
Mcol1, suggesting that *Nanos2* is a nematoblast marker rather than a stem cell
marker in *Cladonema* medusa (S3E and S3F Fig), similar to *Hydractinia* [47]. In
the intact tentacle, *Nanos1+*, *Piwi+* or *Vasa1+* cells, are primarily distributed in
the bulb and branching site (Figs 3Ci and S3A-S3D, S4Ai, S4Bi) [38,39]. By
contrast, at 24 hpa *Nanos1+* cells accumulated around the injury site where
blastema formed (Fig 3Cii), and *Piwi+* and *Vasa1+* cells were also localized in
the blastema region (S4Aii and S4Bii Fig). While *Nanos1+* cells accumulated to
the new branching bud after 48 hpa (Fig 3Ciii, white arrow) [38,39], *Nanos1*
expression was no longer detected at the regenerating tip and was restricted to

the middle and basal side of the tentacle (Fig 3Ciii). We further quantified the
distribution of *Nanos1*[+] cells and found a gradual transition of their distribution
from the distal to the basal side of the regenerating tentacle (Fig 3D). These
results are consistent with the distribution pattern of proliferative cells (Figs 2A-
2C and S2A-S2B), further supporting the idea that proliferative cells in blastema
exhibit characteristics of stem cells.
577
Most blastema cells are not derived from resident stem cells localized at
the bulb
Previous reports on cnidarian regeneration have suggested that undifferentiated
blastema cells are supplied by the migration of resident stem/progenitor cells in
response to injury [20,21,27]. To test whether resident stem cells, or originally-
localized i-cells, in the bulb migrate to the injury site to form blastema, we
performed EdU-chase experiments during *Cladonema* tentacle regeneration.
We leveraged the fact that short-time incubation of EdU labels proliferating i-
cells in the intact tentacle, and, after incubating, amputated and visualized the
regenerating tentacle at different time points (S5A Fig). Until 24 hpa, most EdU[+]
cells remained localized to the bulb while a small number of EdU[+] cells were
distributed near blastema (S5Ai-vi and S5B Fig). Following 48 h and 72 h
chasing, EdU[+] cells continued to localize at the basal side, including the bulb,
rather than in the regenerating tip, although the population of EdU[+] cells
propagated in the distal direction, likely through cell proliferation and migration
(S5Avii and S5Aviii Fig). Considering the minimal contribution of EdU[+] i-cells to
blastema, these observations suggest that resident stem cells localized at the
bulb may not be sufficient to form blastema.
We further conducted an alternative tracing experiment that specifically
labels cells in the bulb region by DiI microinjection followed by amputation and
cell tracking, which allows monitoring the labeled cells *in vivo* during tentacle
regeneration (S6 Fig). To confirm the cell-type of DiI[+] cells in the bulb, we first
combined EdU staining or β-catenin antibody staining with DiI injection (S6A-
S6C Fig) to show that more than 75% of EdU[+] cells or i-cells (β-catenin
cytoplasm signal) were DiI[+] in the bulb (S6A-S6C Fig), indicating that the
majority of resident i-cells are labeled by DiI injection. After a 24 h chase, DiI[+]
cells were still localized near the bulb, not merged with EdU[+] cells in blastema
(S6D and S6Ei-ii Fig). In contrast, when DiI was injected into the central area
near the amputation site, DiI[+] cells overlapped with EdU[+] cells, participating in

blastema formation (S6D and S6Eiii-iv Fig). These results, taken together with
the EdU chase results, suggest that resident stem cells do not migrate to the
blastema region.
The above experiments support the presence of two populations of
proliferative cells during regeneration: one population are the "resident
homeostatic stem cells (RHSCs)," composed of i-cells localized at the bulb, and
the other are "repair-specific proliferative cells (RSPCs)" that appear as
blastema upon amputation. In order to test this hypothesis, we performed
chasing experiments by labeling the two proliferative cell populations with
distinct nucleoside analogs, EdU and BrdU (Fig 4A). This approach has been
utilized in different animal systems as a way to track proliferative cell
populations and their lineages [48–50]. We labeled RHSCs in the bulb by EdU
before amputation and RSPCs in the blastema by BrdU at 24 hpa (Fig 4A). At
the time of blastema formation at 24 hpa, EdU$^+$ RHSCs were still localized to
the bulb, while BrdU$^+$ RSPCs accumulated at the blastema (Fig 4Bi).
Importantly, only 13.6% of BrdU$^+$ RSPCs were EdU$^+$ in blastema, supporting the
notion that the migration of RHSCs may not be sufficient to form blastema (Fig
4Bi, 4Cii, and 4D). Indeed, following 24-48 h, BrdU$^+$ RSPCs were distributed on
the distal side of the regenerating tentacle, EdU$^+$ RHSCs remained localized on
the basal side, although their distribution expanded slightly (Fig 4Bii and 4Biii).
Consistent with these observations, when DiI was injected to the center area
near the injury site, these DiI$^+$ cells initially contributed to RSPCs and were
gradually distributed to distal side in the regenerating tentacle (Figs 4Bii, 4Biii
and S6H-S6I). These results suggest that RSPCs at the blastema participate in
the formation of the new structure while RHSCs expand locally, supporting the
existence of two distinct proliferative cell populations during tentacle
regeneration (Fig 4G).
To determine the characteristics of these distinct proliferative cell
populations, we examined their respective proliferative capacities. For this
purpose, we quantified the increase in cell numbers of the two populations
during tentacle regeneration. At 24 hpa, RHSCs increased approximately two-
fold, but no dramatical increase was observed in the following 48 h (Fig 4E; 0
hpa: n=206±66.6, 24 hpa: n=441.3±100.6, 48 hpa: n=493.3±123.2, 72 hpa:
n=593.7±250.3). Notably, the number of these RHSCs also increased during 24
h without amputation (S5C and S5D Fig), suggesting that the increase of
RHSCs is not an injury-specific response. In contrast, the number of RSPCs in

the blastema tripled at 48 h, which led to an approximate 4-fold increase at 72

hpa (Fig 4F; 24 hpa: n=275.1±64.2, 48 hpa: n=929.3±151.9, 72 hpa:

n=1125±225.1). These results indicate a pronounced difference in proliferative

activity between the two populations, further implicating their distinct responses

to injury.

If RSPCs do not derive from RHSCs in the bulb, blastema should form

when such stem cells are removed. To test this possibility, we examined

blastema formation after RHSCs are eliminated by X-ray irradiation. We first

determined a condition that would eliminate resident stem cells after exposure

to different doses of X-ray irradiation (30, 50, 75 Gy) by monitoring expression

of the stem cell marker *Nanos1*. FISH staining of *Nanos1* showed that *Nanos1*$^+$

cells decreased in an X-ray dose dependent manor, and most *Nanos1*$^+$ cells

were removed at 75 Gy at 3 days post-irradiation (dpi) (S7A-S7C Fig). qPCR

analysis further confirmed a significant reduction of *Nanos1* mRNA levels at 3

dpi (S7G Fig). We thus amputated the tentacle at 3 dpi and investigated

blastema formation by EdU pulse labeling (Fig 4H). At 24 hpa, blastema formed

successfully regardless of X-ray irradiation, although the rate of EdU$^+$ cells in

the blastema decreased slightly (Fig 4I and 4J). In order to support the

conclusion derived from X-ray irradiation, we also performed pharmacological

treatments using hydroxyurea (HU) or Mitomycin C, both of which are used to

eliminate i-cells in other hydrozoan species [20,51,52]. The results of FISH and

qPCR confirmed that most *Nanos1*$^+$ cells are removed by HU or Mitomycin C

treatment for 3 days (Fig S7D-S7G). We then examined blastema formation

after HU or Mitomycin C treatment and found that blastema formed at 24 hpa,

consistent with the results of X-ray irradiation (Fig 4I and 4J). Combined, these

results demonstrate that distinct proliferative cell populations appear during

tentacle regeneration and that most blastema cells are not derived from

RHSCs.

RHSCs contribute to nematogenesis and tentacle elongation

In the tentacles of hydrozoan medusae such as *Clytia* and *Cladonema*,

proliferative i-cells in the bulb are proposed to migrate from the basal to the

distal side while differentiating into nematocytes, which is known as the belt

conveyor model [24,37,53]. Based on this proposed model, we hypothesized

that RHSCs in the bulb contribute to nematogenesis during both homeostasis

and regeneration. In order to test this hypothesis, we performed EdU and BrdU

double chase experiments with some modifications (Fig 5A). We found that
long-incubation with EdU for 24 h allows for the labeling of RHSCs in the bulb
as well as nematocyte progenitors labeled by *Mcol1* (Figs 5Bi and S8). After
EdU labeling for 24 h, followed by amputation, most EdU$^+$ cells remained in the
bulb, and 26.9% of the BrdU$^+$ RSPCs were EdU$^+$ in the blastema, indicating that
the majority of RSPCs are of different lineage than RHSCs (Fig 5Bii, 5Biv, and
5C). This result is consistent with the results of the chasing experiment using
short duration EdU labeling (Fig 4A-4D), further supporting the conclusion that
the migration of RHSCs in the bulb is not sufficient to form blastema. At 72 hpa,
while EdU$^+$ cells were still distributed in the basal region, including the bulb,
some EdU$^+$ cells were located in the distal nematocyte clusters (Fig 5Bv: yellow
arrow), suggesting that one of the lineages of RHSCs are nematocytes during
regeneration. We also performed DiI labeling of the bulb to show that these DiI$^+$
cells were distributed in nematocyte clusters in the regenerated tentacle (S6F
and S6G Fig: white arrow).
To further confirm the above observations, we identified cell types
located in the distal side of the tentacle. We found that atypical protein kinase C
(aPKC) antibody can be utilized to detect cell types with nuclear staining, which
is a method similar to using β-catenin antibody (Figs 3A, 5D and S9) [39,46].
The combination of EdU chasing and cell type identification revealed that nearly
half of the EdU$^+$ cells are nematocytes in the distal side of the tentacle
(nematocytes: 45.7%, epithelial cells: 24.2%, i-cells: 30.1%; Fig 5E-5G). These
results together indicate that RHSCs in the bulb differentiate into nematocytes
during regeneration. Intriguingly, we also noticed that EdU-labeled RHSCs
merged with FMRFamide$^+$ neurons (S10A and S10B Fig), suggesting that
RHSCs can differentiate into neurons during tentacle regeneration.
Given that the RHSCs mainly differentiate into both nematocytes and
epithelial cells during regeneration (Fig 5F and 5G), it is expected that these
cells play a role in supplying new nematocytes as well as in tentacle elongation.
We first examined the timing of nematocyte cluster formation as a readout of
nematogenesis when RHSCs were eliminated by X-ray or drug treatments
using HU or Mitomycin C (Fig 5H-5J) and found that it was delayed, particularly
at the early stage of regeneration after elimination of RHSCs (Fig 5I and 5J),
indicating that RHSCs migrate from the basal to the distal side while
differentiating into nematocytes during regeneration.

Next, we measured tentacle length during regeneration with RHSCs
cells removed by X-ray irradiation or drug treatments (Fig 5H and 5K). When
tentacles were amputated at 3 dpi by which time RHSCs had been eliminated
by X-ray, tentacle elongation was severely inhibited compared to the non-
irradiated controls (Fig 5K, regenerated tentacle length at 4 dpa, Control: 1.66
mm, X-ray: 0.540 mm). It is possible that X-ray irradiation affects the source of
RSPCs to some extent, and indeed when RHSCs were removed with HU or
Mitomycin C treatment, tentacle elongation was mildly but still significantly
inhibited (Fig 5K, regenerated tentacle length at 4 dpa, HU: 1.14 mm, Mitomycin
C: 1.19 mm). Given that EdU$^+$ blastema cells appear after X-ray irradiation as
well as drug treatments (Fig 4I and 4J), RHSCs contribute to tentacle elongation
independently of blastema cells. We speculate that RHSCs, and particularly
their decedents, participate in tissue elongation likely through pushing out from
the basal side as progenitors and differentiating into epithelial cells.
728
RSPCs contribute to reconstruction of the new tentacle
In order to investigate the role of RSPCs during regeneration, we performed
EdU chase labeling of the blastema with cell type identification by aPKC
antibody staining (Fig 6A-6C). While 94.4% of the RSPCs exhibited i-cell
morphology at 24 hpa (Fig 6Bi, 6Biii and 6C, consistent with the results of β-
catenin antibody staining in Fig 3A and 3B), 59.2% of EdU$^+$ cells were epithelial
cells in the distal portion of the regenerating tentacle at 72 hpa, (Fig 6Bii, 6Biv
and 6C). Strikingly, in contrast to RHSCs, EdU$^+$ cells derived from the blastema
did not become nematocytes (Fig 5E-5G and 6A-6C). These results suggest
that RSPCs preferentially differentiate into epithelial cells in the regenerating
tentacle. Of note, we found that a small portion of EdU-labeled RSPCs merged
with FMRFamide$^+$ neurons, although these populations are significantly smaller
than the RHSCs (S10C-S10E Fig). Combined, these findings indicate that
RSPCs are multipotent, or at least bipotent, stem cells that preferentially
generate epithelial cells during regeneration.
To identify the role of RSPCs, we examined the impact of locally
inhibiting cell proliferation on tentacle regeneration (Figs 6D-6I and S11). UVA
radiation (320 to 400 nm) induces various forms of cellular damage via the
production of reactive oxygen species or the impairment of proteasomal
function, which blocks cell cycle progression at the G$_2$/M phase in both
mammalian cultured cells and fly epidermal progenitors [54–57]. In addition,

Hoechst and BrdU are known photosensitizers [58,59], which can facilitate the
process of damage induction. To verify the effects of UVA on RSPCs, we
performed UV laser illumination around the blastema region at 25 hpa and
counted the number of EdU labeled cells (Fig 6D-6F). Following around 24-48
h, the number of EdU$^+$ cells decreased significantly in the UV exposed tentacle
compared to the control, indicating that UV laser exposure inhibits cell
proliferation of RSPCs (Fig 6F). In contrast, we confirmed that cell proliferation
and the timing of nematocyte cluster formation by RHSCs were similar to the
non-UV exposure controls, suggesting a minimal impact of UV laser exposure
on RHSCs (S11 Fig). In order to examine the contribution of the proliferation of
RSPCs toward tentacle regeneration, we monitored the length of regenerating
tentacles after UV laser exposure (Fig 6G). Upon UV laser illumination, tentacle
elongation during regeneration was significantly inhibited (Fig 6H and 6I). These
results, together with the distribution of their descendants on the distal side of
the regenerating tentacle (Fig 4Biii and 5Biii), indicate that proliferation of
RSPCs contribute to the reconstruction of new tentacles after amputation.
766
767

Discussion
In this study, we uncovered the cellular mechanism underlying blastema
formation during *Cladonema* tentacle regeneration. We found that, upon
amputation of the tentacle, RSPCs accumulate as blastema, most of which is
not derived from RHSCs. While RHSCs contribute to nematogenesis and tissue
elongation during both homeostasis and regeneration, RSPCs differentiate into
the epithelial cells of the newly formed tentacle. Based on these results, we
propose that distinct proliferative cell populations facilitate rapid functional
regeneration of the tentacle, which enables critical behaviors of jellyfish
organismal physiology such as feeding and swimming (Fig 7A).
778
779

Heterogeneity of stem-like cells
In hydrozoans, i-cells behave as pluripotent or multipotent stem cells that can
differentiate into several progenitors and differentiated lineages such as
nematocytes, neurons, gland cells, and gametes. Although i-cells exhibit
common features such as their interstitial localization, cellular morphology, and
expression of the conserved stem cell markers (*Nanos, Piwi, Vasa, PL10*); their

potency and heterogeneity likely differ across species [45]. Indeed, *Hydra* i-cells
differentiate into most cell types except epithelial cells, while *Hydractinia* i-cells
behave as pluripotent cells that can differentiate into all cell types including
somatic and germline cells [16,60,61].
In the intact *Cladonema* jellyfish, *Nanos1*$^+$, *Piwi*$^+$, and *Vasa1*$^+$ cells are
primarily distributed at the bulb (Figs 3C and S4A, S4B), and about 70% of
RHSCs in the bulb are *Nanos1*$^+$ cells (Fig 3E and 3F). From lineage-tracing
experiments, these RHSCs differentiate into epithelia as well as nematocytes
(Fig 5F and 5G), suggesting their role as multipotent stem cells. These
observations are consistent with the jellyfish *Clytia*, in which i-cells in the bulb
proliferate and differentiate into nematocytes [37,53], indicating that i-cells in the
tentacle bulb are commonly involved in organ homeostasis including
nematogenesis in hydrozoan jellyfish.
In contrast to RHSCs in the bulb, only 17-36% of RSPCs at blastema
express *Nanos1*$^+$, *Piwi*$^+$, and *Vasa1*$^+$ (Figs 3E-3F and S4C-S4E)*,* although more
than 80% of blastema cells show i-cell morphology (Fig 3A-3B and 6B-6C).
These results imply that the stem-like population of the blastema is likely
heterogenous, and differs from that of resident i-cells in the bulb. Indeed,
RSPCs predominantly differentiate into epithelial cells rather than nematocytes
(Fig 6A-6C), suggesting their role as lineage-restricted stem/progenitor cells
while maintaining the potential to become neurons in the regenerating tentacle
(S10 Fig). Such stem-like cells also accumulate at the newly-formed branch
starting at 48-72 hpa (Fig 4Bii-iii), consistent with the prediction by the previous
report proposing that i-cell population in the branching bud differs from RHSCs
in the bulb [39]. While previous studies using other cnidarians have described
active cell proliferation upon injuries, these proliferating cells are
pluri/multipotent stem cells migrating to the injury site [20,21,27,62], not lineage-
restricted stem/progenitor cells like RSPCs. Altogether, our findings support the
heterogeneity of stem-like cells in *Cladonema* jellyfish (Fig 7A), which is similar
to the subpopulations of *Hydra* i-cells [52,63].
816
817
How are RSPCs supplied in the blastema?
Our work is the first to demonstrate that RSPCs, which exhibit preferential
differentiation into epithelial cells, appear upon injury in cnidarians, but how this
cell population is supplied remains unknown. Lineage-tracing experiments,

together with the elimination of RHSCs show that blastema formation is mostly
independent of RHSCs in the bulb (Figs 4 and S6). When the distal side of the
tentacle, where few proliferating cells exist, is amputated, local cell proliferation
is still observed around the injury site (Fig 2G). These observations raise two
possibilities: mitotic re-entry of quiescent/slow-cycling stem cells or
dedifferentiation from differentiated cells into stem-like cells.

Previous reports have showed that some cnidarian species possess
quiescent or slow-cycling stem cells. For instance, while *Hydra* has three types
of adult stem cells—i-cells, epidermal stem cells, and gastrodermis epithelial
stem cells—the subpopulation of each stem cell is slow-cycling [52]. These
slow-cycling cells are resistant to hydroxyurea treatment due to G2 phase
arrest, and re-enter the cell cycle during head regeneration. *Nematostella* also
has slow-cycling/quiescent cells that are predominantly present in the
mesenteries and are resistant to X-ray irradiation, and these cells' cell cycle is
activated upon injury, when they migrate toward the wound site [21]. Given that
a small number of i-cells exist in the *Cladonema* intact tentacle, outside the
bulb, and branching bud (Figs 3C and S3A, S3C, S3D, S4A, S4B)[39]; such a
minor population of i-cells may behave as slow-cycling/quiescent cells and re-
enter the cell cycle in response to injury.

By contrast, although cnidarian dedifferentiation potential is still debated,
some species appear to exhibit context-dependent cellular plasticity [64]. For
instance, classical experiments using the hydrozoan jellyfish *Podocoryna*
carnea have indicated their dedifferentiation potential *in vitro*: when isolated
striated muscles are cultured after collagenase treatment, the muscle cells
dedifferentiate into stem-like cells and re-differentiate into multiple cell types
[65]. Additionally, in green *Hydra*, the isolated gastrodermis, which is i-cell free,
can reconstitute a complete polyp including the epidermis [66]. In this case,
gastrodermal gland cells appear to dedifferentiate into i-cells [67], while
gastrodermal digestive cells transdifferentiate into epidermal epitheliomuscular
cells [68]. Moreover, a recent report using *Hydractinia* has shown that, when the
hypostome is isolated from the body, cellular reprogramming triggered by cell
senescence occurs in somatic cells, which convert into pluripotent *Piwi1*+ i-cells
[69]. Although these reports support the possibility that dedifferentiation
produces pluri/multipotent stem cells after isolation from the body, it is also
possible that RSPCs are derived from their descendant cell types. During newt
limb regeneration, cellular lineage is restricted such that muscle progenitors

dedifferentiated from resident muscle fibers produce only muscles [7,70]. Given

that repair-specific proliferative cells in the *Cladonema* tentacle mainly

differentiate into epithelial cells (Fig 6A-6C), epithelial cells may dedifferentiate

into progenitors that give rise to epithelial cells in the newly formed tentacle.

Evolutionary conservation and diversification of blastema formation

Although the molecular and cellular mechanisms of animal regeneration have

been addressed in detail using representative species such as planaria and

salamanders, questions such as the evolutionary history of regeneration and

the homology of regeneration mechanisms across species remain largely

unknown. Repair-specific stem cells provided by quiescent stem cells and/or

dedifferentiation are common blastema components in bilaterian appendage

regeneration. As in vertebrates where lineage-restricted repair-specific

stem/progenitor cells participate in appendage regeneration [7,9,11], the

protostome ecdysozoa *Parhyale hawaiensis* utilize re-activation of muscle

satellite cells during thoracic leg regeneration [71]. By contrast, during

echinoderm starfish arm regeneration, dedifferentiated dermal cells form

blastema-like structures [72]. Similarly, during insect cricket leg regeneration,

the source of blastemal cells is derived from injury-induced dedifferentiation

[73,74]. These reports suggest that repair-specific production of blastema cells

is relatively conserved throughout bilaterians (Fig 7B).

In non-bilaterian metazoans, resident pluripotent stem cells are thought

to contribute to both homeostasis and whole-body regeneration. During the

regeneration of the demosponge in Porifera, one of the most primitive animals,

archeocytes known as homeostatic stem cells contribute to an undifferentiated

cell mass beneath the wound as a blastema-like structure [75]. While the

regenerating ctenophore cydippids do not apparently form blastema-like

structure, slow-cycling stem-like cells are recruited to the injury site [76]. In

cnidarians, resident stem cells, which are involved in homeostasis, are recruited

to the injury site after amputation of the body and removal of the organs [19–

21,27]. By contrast, as shown in this study, the main cellular source of

blastemal cells is not RHSCs but rather RSPCs during *Cladonema* tentacle

regeneration (Fig 4). Our findings suggest that, in terms of the regeneration

processes and blastemal cellular source, the regeneration mechanism of the

Cladonema tentacle is more similar to that of bilaterian species (Fig 7B).

One important remaining question is whether bilaterians' and cnidarians'
most recent common ancestor possessed a system to supply blastema through
repair-specific stem/proliferative cells. Although sponges and ctenophores have
been proposed to possess dedifferentiation or transdifferentiation potentials that
can provide a cellular source of regenerated bodies [64], it is currently unknown
whether they are repair-specific cell populations or not. In this study, we
demonstrate that RSPCs contribute to the newly generated epithelium during
Cladonema tentacle regeneration. Intriguingly, recent genomic studies have
revealed that jellyfish-specific genes are not conserved among bilaterians or
even other cnidarians including corals and sea anemones, and the anthozoan
Nematostella is the most 'bilaterian-like' cnidarian sequenced [77–79]. These
findings raise the possibility that medusozoans had evolved the medusa stage
after branching from their common ancestor and that bilaterians and cnidarian
jellyfish likely independently acquired a similar mechanism of blastema
formation upon amputation of appendages. It is thus tempting to speculate that
blastema formation by RSPCs is a common feature acquired for complex organ
and appendage regeneration during animal evolution.
910
911

References

1.   McCusker C, Bryant SV, Gardiner DM. The axolotl limb blastema: cellular
and molecular mechanisms driving blastema formation and limb regeneration in
tetrapods: The Axolotl Limb Blastema. Regeneration. 2015;2: 54–71.
doi:10.1002/reg2.32
2.   Seifert AW, Muneoka K. The blastema and epimorphic regeneration in
mammals. Developmental Biology. 2018;433: 190–199.
doi:10.1016/j.ydbio.2017.08.007
3.   Aztekin C. Appendage regeneration is context dependent at the cellular level.
Open Biol. 2021;11: 210126. doi:10.1098/rsob.210126
4.   Bely AE, Nyberg KG. Evolution of animal regeneration: re-emergence of a
field. Trends in Ecology & Evolution. 2010;25: 161–170.
doi:10.1016/j.tree.2009.08.005
5.   Tanaka EM, Reddien PW. The Cellular Basis for Animal Regeneration.
Developmental Cell. 2011;21: 172–185. doi:10.1016/j.devcel.2011.06.016

6.   Reddien PW, Alvarado AS. FUNDAMENTALS OF PLANARIAN

REGENERATION. Annu Rev Cell Dev Biol. 2004;20: 725–757.

doi:10.1146/annurev.cellbio.20.010403.095114

7.   Sandoval-Guzmán T, Wang H, Khattak S, Schuez M, Roensch K, Nacu E, et

al. Fundamental Differences in Dedifferentiation and Stem Cell Recruitment during

Skeletal Muscle Regeneration in Two Salamander Species. Cell Stem Cell. 2014;14:

174–187. doi:10.1016/j.stem.2013.11.007

8.   Ando K, Shibata E, Hans S, Brand M, Kawakami A. Osteoblast Production by

Reserved Progenitor Cells in Zebrafish Bone Regeneration and Maintenance.

Developmental Cell. 2017;43: 643-650.e3. doi:10.1016/j.devcel.2017.10.015

9.   Knopf F, Hammond C, Chekuru A, Kurth T, Hans S, Weber CW, et al. Bone

Regenerates via Dedifferentiation of Osteoblasts in the Zebrafish Fin. Developmental

Cell. 2011;20: 713–724. doi:10.1016/j.devcel.2011.04.014

10.   Daponte V, Tylzanowski P, Forlino A. Appendage Regeneration in

Vertebrates: What Makes This Possible? Cells. 2021;10: 242.

doi:10.3390/cells10020242

11.   Lehoczky JA, Robert B, Tabin CJ. Mouse digit tip regeneration is mediated by

fate-restricted progenitor cells. Proc Natl Acad Sci USA. 2011;108: 20609–20614.

doi:10.1073/pnas.1118017108

12.   Fritz AE, Ikmi A, Seidel C, Paulson A, Gibson MC. Mechanisms of tentacle

morphogenesis in the sea anemone *Nematostella vectensis*. Development. 2013;140:

2212–2223. doi:10.1242/dev.088260

13.   Technau U, Steele RE. Evolutionary crossroads in developmental biology:

Cnidaria. Development. 2011;138: 1447–1458. doi:10.1242/dev.048959

14.   Fujita S, Kuranaga E, Nakajima Y. Regeneration Potential of Jellyfish: Cellular

Mechanisms and Molecular Insights. Genes. 2021;12: 758. doi:10.3390/genes12050758

15.   Holstein TW, Hobmayer E, Technau U. Cnidarians: An evolutionarily

conserved model system for regeneration? Dev Dyn. 2003;226: 257–267.

doi:10.1002/dvdy.10227

16.   Gahan JM, Bradshaw B, Flici H, Frank U. The interstitial stem cells in

Hydractinia and their role in regeneration. Current Opinion in Genetics & Development.

2016;40: 65–73. doi:10.1016/j.gde.2016.06.006

17.   Röttinger E. Nematostella vectensis, an Emerging Model for Deciphering the

Molecular and Cellular Mechanisms Underlying Whole-Body Regeneration. Cells.

2021;10: 2692. doi:10.3390/cells10102692

18. Vogg MC, Galliot B, Tsiairis CD. Model systems for regeneration: Hydra.
Development. 2019;146. doi:10.1242/dev.177212

19. Chera S, Ghila L, Dobretz K, Wenger Y, Bauer C, Buzgariu W, et al.
Apoptotic Cells Provide an Unexpected Source of Wnt3 Signaling to Drive Hydra Head
Regeneration. Developmental Cell. 2009;17: 279–289.
doi:10.1016/j.devcel.2009.07.014

20. Bradshaw B, Thompson K, Frank U. Distinct mechanisms underlie oral vs
aboral regeneration in the cnidarian Hydractinia echinata. eLife. 2015;4: e05506.
doi:10.7554/eLife.05506

21. Amiel AR, Foucher K, Ferreira S, Röttinger E. Synergic coordination of stem
cells is required to induce a regenerative response in anthozoan cnidarians. bioRxiv;
2019. p. 2019.12.31.891804. doi:10.1101/2019.12.31.891804

22. Katsuki T, Greenspan RJ. Jellyfish nervous systems. Current Biology. 2013;23:
R592–R594. doi:10.1016/j.cub.2013.03.057

23. Leclère L, Röttinger E. Diversity of Cnidarian Muscles: Function, Anatomy,
Development and Regeneration. Front Cell Dev Biol. 2017;4.
doi:10.3389/fcell.2016.00157

24. Fujita S, Kuranaga E, Nakajima Y. Cell proliferation controls body size
growth, tentacle morphogenesis, and regeneration in hydrozoan jellyfish *Cladonema*
pacificum. PeerJ. 2019;7: e7579. doi:10.7717/peerj.7579

25. Gamero-Mora E, Halbauer R, Bartsch V, Stampar SN, Morandini AC.
Regenerative Capacity of the Upside-down Jellyfish Cassiopea xamachana. Zool Stud.
2019;58. doi:10.6620/ZS.2019.58-37

26. Ostendarp M, Plewka J, Flathmann J, Tilstra A, El-Khaled YC, Wild C.
Complete and rapid regeneration of fragments from the upside-down jellyfish
Cassiopea. Front Mar Sci. 2022;9: 949233. doi:10.3389/fmars.2022.949233

27. Sinigaglia C, Peron S, Eichelbrenner J, Chevalier S, Steger J, Barreau C, et al.
Pattern regulation in a regenerating jellyfish. eLife. 2020;9: e54868.
doi:10.7554/eLife.54868

28. Stamatis S-A, Worsaae K, Garm A. Regeneration of the Rhopalium and the
Rhopalial Nervous System in the Box Jellyfish *Tripedalia cystophora*. The Biological
Bulletin. 2018;234: 22–36. doi:10.1086/697071

29. Stierwald M, Yanze N, Bamert RP, Kammermeier L, Schmid V. The Sine
oculis/Six class family of homeobox genes in jellyfish with and without eyes:
development and eye regeneration. Developmental Biology. 2004;274: 70–81.
doi:10.1016/j.ydbio.2004.06.018

30.   Weber C. Structure, histochemistry, ontogenetic development, and regeneration
of the ocellus ofCladonema radiatum dujardin (cnidaria, hydrozoa, anthomedusae). J
Morphol. 1981;167: 313–331. doi:10.1002/jmor.1051670306

31.   Fujiki A, Hou S, Nakamoto A, Kumano G. Branching pattern and
morphogenesis of medusa tentacles in the jellyfish Cladonema pacificum (Hydrozoa,
Cnidaria). Zoological Lett. 2019;5: 12. doi:10.1186/s40851-019-0124-4

32.   Masuda-Ozawa T, Fujita S, Nakamura R, Watanabe H, Kuranaga E, Nakajima
Y. siRNA-mediated gene knockdown via electroporation in hydrozoan jellyfish
embryos. Sci Rep. 2022;12: 16049. doi:10.1038/s41598-022-20476-1

33.   Takeda N, Kon Y, Artigas GQ, Lapébie P, Barreau C, Koizumi O, et al.
Identification of jellyfish neuropeptides that act directly as oocyte maturation-inducing
hormones. Development. 2018;145. doi:10.1242/dev.156786

34.   Rees JT. The polyp and immature medusa stages of *Cladonema californicum* ,
Hyman, 1947 (Anthomedusae: Cladonemidae) with biological notes and a discussion of
the taxonomy of the genus *Cladonema*. Journal of Natural History. 1979;13: 295–302.
doi:10.1080/00222937900770231

35.   Suga H, Schmid V, Gehring WJ. Evolution and Functional Diversity of
Jellyfish Opsins. Current Biology. 2008;18: 51–55. doi:10.1016/j.cub.2007.11.059

36.   Zhou K, Gu Y, Wang L, Chen J. Discovery of Cladonema multiramosum sp.
nov. (Cnidaria: Hydrozoa: Cladonematidae) using DNA barcoding and life cycle
analyses. Acta Oceanol Sin. 2022;41: 44–52. doi:10.1007/s13131-021-1900-5

37.   Denker E, Manuel M, Leclère L, Le Guyader H, Rabet N. Ordered progression
of nematogenesis from stem cells through differentiation stages in the tentacle bulb of
Clytia hemisphaerica (Hydrozoa, Cnidaria). Developmental Biology. 2008;315: 99–113.
doi:10.1016/j.ydbio.2007.12.023

38.   Fujita S, Kuranaga E, Miura M, Nakajima Y-I. Fluorescent In Situ
Hybridization and 5-Ethynyl-2'-Deoxyuridine Labeling for Stem-like Cells in the
Hydrozoan Jellyfish Cladonema pacificum. J Vis Exp. 2022. doi:10.3791/64285

39.   Hou S, Zhu J, Shibata S, Nakamoto A, Kumano G. Repetitive accumulation of
interstitial cells generates the branched structure of *Cladonema* medusa tentacles.
Development. 2021;148: dev199544. doi:10.1242/dev.199544

40.   Passamaneck YJ, Martindale MQ. Cell proliferation is necessary for the
regeneration of oral structures in the anthozoan cnidarian Nematostella vectensis. BMC
Dev Biol. 2012;12: 34. doi:10.1186/1471-213X-12-34

41.   Galliot B, Buzgariu W, Schenkelaars Q, Wenger Y. Non-developmental

dimensions of adult regeneration in Hydra. Int J Dev Biol. 2018;62: 373–381.

doi:10.1387/ijdb.180111bg

42.   Lesh-Laurie GE, Hujer A, Suchy P. Polyp regeneration from isolated tentacles

of Aurelia scyphistomae: a role for gating mechanisms and cell division. Hydrobiologia.

1991;216: 91–97. doi:10.1007/BF00026448

43.   Salic A, Mitchison TJ. A chemical method for fast and sensitive detection of

DNA synthesis *in vivo*. Proc Natl Acad Sci USA. 2008;105: 2415–2420.

doi:10.1073/pnas.0712168105

44.   Tasaki J, Shibata N, Sakurai T, Agata K, Umesono Y. Role of c-Jun N-terminal

kinase activation in blastema formation during planarian regeneration: JNK activation in

planarian regeneration. Development, Growth & Differentiation. 2011;53: 389–400.

doi:10.1111/j.1440-169X.2011.01254.x

45.   Gold DA, Jacobs DK. Stem cell dynamics in Cnidaria: are there unifying

principles? Dev Genes Evol. 2013;223: 53–66. doi:10.1007/s00427-012-0429-1

46.   Plickert G, Frank U, Müller WA. Hydractinia, a pioneering model for stem cell

biology and reprogramming somatic cells to pluripotency. Int J Dev Biol. 2012;56:

519–534. doi:10.1387/ijdb.123502gp

47.   Kanska J, Frank U. New roles for Nanos in neural cell fate determination

revealed by studies in a cnidarian. Development. 2013;140: e1608–e1608.

doi:10.1242/dev.101246

48.   Harris L, Zalucki O, Piper M. BrdU/EdU dual labeling to determine the cell-

cycle dynamics of defined cellular subpopulations. J Mol Hist. 2018;49: 229–234.

doi:10.1007/s10735-018-9761-8

49.   Ribeiro RP, Egger B, Ponz-Segrelles G, Aguado MT. Cellular proliferation

dynamics during regeneration in Syllis malaquini (Syllidae, Annelida). Front Zool.

2021;18: 27. doi:10.1186/s12983-021-00396-y

50.   Vega CJ, Peterson DA. Stem cell proliferative history in tissue revealed by

temporal halogenated thymidine analog discrimination. Nat Methods. 2005;2: 167–169.

doi:10.1038/nmeth741

51.   Bode HR, Scott Smith G. Regulation of interstitial cell differentiation inHydra

attenuata: II. Correlation of the axial position of the interstitial cell with nematocyte

differentiation. Wilhelm Roux' Archiv. 1977;181: 203–213. doi:10.1007/BF00848421

52.   Govindasamy N, Murthy S, Ghanekar Y. Slow-cycling stem cells in hydra

contribute to head regeneration. Biology Open. 2014;3: 1236–1244.

doi:10.1242/bio.201410512

53.   Condamine T, Jager M, Leclère L, Blugeon C, Lemoine S, Copley RR, et al.
Molecular characterisation of a cellular conveyor belt in Clytia medusae.
Developmental Biology. 2019;456: 212–225. doi:10.1016/j.ydbio.2019.09.001

54.   Catalgol B, Ziaja I, Breusing N, Jung T, Höhn A, Alpertunga B, et al. The
Proteasome Is an Integral Part of Solar Ultraviolet A Radiation-induced Gene
Expression. Journal of Biological Chemistry. 2009;284: 30076–30086.
doi:10.1074/jbc.M109.044503

55.   Girard P-M, Pozzebon M, Delacôte F, Douki T, Smirnova V, Sage E.
Inhibition of S-phase progression triggered by UVA-induced ROS does not require a
functional DNA damage checkpoint response in mammalian cells. DNA Repair.
2008;7: 1500–1516. doi:10.1016/j.dnarep.2008.05.004

56.   Latonen L, Laiho M. Cellular UV damage responses—Functions of tumor
suppressor p53. Biochimica et Biophysica Acta (BBA) - Reviews on Cancer.
2005;1755: 71–89. doi:10.1016/j.bbcan.2005.04.003

57.   Nakajima Y, Kuranaga E, Sugimura K, Miyawaki A, Miura M.
Nonautonomous Apoptosis Is Triggered by Local Cell Cycle Progression during
Epithelial Replacement in Drosophila. Mol Cell Biol. 2011;31: 2499–2512.
doi:10.1128/MCB.01046-10

58.   Herweijer H, And WS, Visser JWM. High-speed photodamage cell selection
using bromodeoxyuridine/Hoechst 33342 photosensitized cell killing. Cytometry.
1988;9: 143–149. doi:10.1002/cyto.990090208

59.   Singh S, Dwarakanath B, Mathew T. DNA ligand Hoechst-33342 enhances UV
induced cytotoxicity in human glioma cell lines. Journal of Photochemistry and
Photobiology B: Biology. 2004;77: 45–54. doi:10.1016/S1011-1344(04)00122-8

60.   Varley Á, Horkan HR, McMahon ET, Krasovec G, Frank U. Pluripotent, germ
cell competent adult stem cells underlie cnidarian plant-like life history. Developmental
Biology; 2022 Nov. doi:10.1101/2022.11.09.515637

61.   Watanabe H, Hoang VT, Mättner R, Holstein TW. Immortality and the base of
multicellular life: Lessons from cnidarian stem cells. Seminars in Cell & Developmental
Biology. 2009;20: 1114–1125. doi:10.1016/j.semcdb.2009.09.008

62.   Chera S, Ghila L, Wenger Y, Galliot B. Injury-induced activation of the
MAPK/CREB pathway triggers apoptosis-induced compensatory proliferation in hydra
head regeneration: MAPK-dependent injury-induced apoptosis in Hydra. Development,
Growth & Differentiation. 2011;53: 186–201. doi:10.1111/j.1440-169X.2011.01250.x

63.   David CN. Interstitial stem cells in Hydra: multipotency and decision-making.
Int J Dev Biol. 2012;56: 489–497. doi:10.1387/ijdb.113476cd

64.   Nakanishi N, Jacobs DK. The Early Evolution of Cellular Reprogramming in
Animals. 1st ed. In: Bishop CD, Hall BK, editors. Deferring Development. 1st ed. CRC
Press; 2019. pp. 67–86. doi:10.1201/9780429445446-4
65.   Schmid V, Alder H. Isolated, mononucleated, striated muscle can undergo
pluripotent transdifferentiation and form a complex regenerate. Cell. 1984;38: 801–809.
doi:10.1016/0092-8674(84)90275-7
66.   Haynes J, Burnett AL. Dedifferentiation and Redifferentiation of Cells in
Hydra viridis. Science. 1963;142: 1481–1483. doi:10.1126/science.142.3598.1481
67.   Burnett AL, Davis LE, Ruffing FE. A histological and ultrastructural study of
germinal differentiation of interstitial cells arising from gland cells inHydra viridis. J
Morphol. 1966;120: 1–8. doi:10.1002/jmor.1051200102
68.   Davis LE, Burnett AL, Haynes JF, Mumaw VR. A histological and
ultrastructural study of dedifferentiation and redifferentiation of digestive and gland
cells in Hydra viridis. Developmental Biology. 1966;14: 307–329. doi:10.1016/0012-
1606(66)90018-2
69.   Salinas-Saavedra M, Febrimarsa, Krasovec G, Horkan HR, Baxevanis AD,
Frank U. Senescence-induced cellular reprogramming drives cnidarian whole-body
regeneration. Cell Reports. 2023;42: 112687. doi:10.1016/j.celrep.2023.112687
70.   Tanaka HV, Ng NCY, Yang Yu Z, Casco-Robles MM, Maruo F, Tsonis PA, et
al. A developmentally regulated switch from stem cells to dedifferentiation for limb
muscle regeneration in newts. Nat Commun. 2016;7: 11069. doi:10.1038/ncomms11069
71.   Konstantinides N, Averof M. A Common Cellular Basis for Muscle
Regeneration in Arthropods and Vertebrates. Science. 2014;343: 788–791.
doi:10.1126/science.1243529
72.   Fan T, Fan X, Du Y, Sun W, Zhang S, Li J. Patterns and cellular mechanisms
of arm regeneration in adult starfish Asterias rollestoni bell. J Ocean Univ China.
2011;10: 255–262. doi:10.1007/s11802-011-1837-y
73.   Bando T, Mito T, Hamada Y, Ishimaru Y, Noji S, Ohuchi H. Molecular
mechanisms of limb regeneration: insights from regenerating legs of the cricket Gryllus
bimaculatus. Int J Dev Biol. 2018;62: 559–569. doi:10.1387/ijdb.180048ho
74.   Truby PR. Blastema formation and cell division during cockroach limb
regeneration. Development. 1983;75: 14.
75.   Borisenko IE, Adamska M, Tokina DB, Ereskovsky AV. Transdifferentiation
is a driving force of regeneration in *Halisarca dujardini* (Demospongiae, Porifera).
PeerJ. 2015;3: e1211. doi:10.7717/peerj.1211

76.   Ramon-Mateu J, Ellison ST, Angelini TE, Martindale MQ. Regeneration in the
ctenophore Mnemiopsis leidyi occurs in the absence of a blastema, requires cell
division, and is temporally separable from wound healing. BMC Biol. 2019;17: 80.
doi:10.1186/s12915-019-0695-8
77.   Khalturin K, Shinzato C, Khalturina M, Hamada M, Fujie M, Koyanagi R, et
al. Medusozoan genomes inform the evolution of the jellyfish body plan. Nat Ecol Evol.
2019;3: 811–822. doi:10.1038/s41559-019-0853-y
78.   Putnam NH, Srivastava M, Hellsten U, Dirks B, Chapman J, Salamov A, et al.
Sea Anemone Genome Reveals Ancestral Eumetazoan Gene Repertoire and Genomic
Organization. Science. 2007;317: 86–94. doi:10.1126/science.1139158
79.   Steele RE, David CN, Technau U. A genomic view of 500 million years of
cnidarian evolution. Trends in Genetics. 2011;27: 7–13. doi:10.1016/j.tig.2010.10.002

1151
1152

Acknowledgement

We thank R. Deguchi (Miyagi Univ. Education, Japan) for sharing *Cladonema pacificum*. We thank D. Umetsu, H. Nagai and T. Akiyama for helpful discussion. We thank I. Nagai, A. Sasaki, and H. Nakatani for technical assistance and animal maintenance. We thank Y. Nashimoto for technical support. This work was supported by JSPS/MEXT KAKENHI (grant numbers JP21H05767 to G.K., JP21H05255 to E.K., JP21H04774, JP21K19206 to M.M., and JP17H06332, JP22H02762, JP23H04696 to Y.N.), JST grant (JPMJCR1852 to E.K. and JPMJSP2114 to S.F.), AMED-Aging (JP21gm5010001 to M.M.), and AMED-PRIME (JP22gm6110025 to Y.N.), and NIBB Collaborative Research Program (23NIBB202 to Y.N.).

Author contributions

S.F. conceptualized and designed the project, performed experiments, analyzed data, prepared figures, and wrote the manuscript. M.T. performed experiments, and analyzed data. G.K., E.K., and M.M. supervised the project. Y.N. conceptualized, supervised and designed the project, prepared figures, and wrote the manuscript. All authors approved the final manuscript.

Competing interests

1175 The authors declare no competing interests.

1176

1177

1178 **Figure legends**

1179

Fig 1. Regeneration processes and potentials of the *Cladonema* medusa
tentacle. (A) The phylogenetic tree of Bilateria and Cnidaria, composed of
Anthozoa and Medusozoa. (B) Medusa of *Cladonema pacificum*. U: umbrella,
M: manubrium, Ra: radial canal, Ri: ring canal, T: tentacle. (C) Tentacle of
Cladonema medusa. O: ocellus, TB: tentacle bulb, Br: branch, NC: nematocyte
cluster. (D) Tentacle regeneration processes after amputation while retaining
the bulb. hpa: hour post-amputation. Yellow arrows indicate nematocyte
clusters. n = 234/236 (tentacles). (E) Scheme of the tentacle regeneration that
retains the bulb. (F) Distribution of mature nematocytes and localization of
muscle fibers during tentacle regeneration. DAPI for poly-γ-glutamate (green)
and nuclei (blue), and Phalloidin for F-actin (red). Yellow arrows indicate
nematocyte clusters. (G) Neural morphology in the regenerating tentacle
stained with the anti-FMRFamide antibody. Yellow arrows indicate neural fibers.
(H) The regenerating tentacle is fully functional at 72 hpa. Image of the tentacle
capturing prey, brine shrimp (White arrow). (I) The rate of functional tentacles
across the regeneration time course. Intact: n = 40 (tentacles), 0-72 hpa: n = 36.
(J) Tentacle regeneration process after removing the bulb from canals. Yellow
arrow in (iv) indicates a nematocyte cluster. Dpa: day post-amputation. n =
117/219 (tentacles). Ra: radial canal, Ri: ring canal. (K) Scheme of tentacle
regeneration without bulb. Scale bars: (B-D, J) 1 mm, (F, Gi) 100 μm, (Gii) 50
μm.

1201

Fig 2. Blastema formation at early phase of tentacle regeneration.
(A) Distribution of S-phase cells in intact and regenerating tentacle stained with
EdU 1 h pulse labeling at intact as well as 24, 48, and 72 hpa, as shown in the
scheme (A'). The tentacles were divided into three parts defined as basal (grey
arrows), middle (pink arrows), and distal (red arrows). (B) Quantification of EdU$^+$
cells' distribution in regenerating tentacle based on the defined areas in (A).
Intact: n = 19 (tentacles), 24 hpa: n = 10, 48 hpa: n = 9, 72 hpa: n = 8. (C)
Scheme of proliferative cells distribution during tentacle regeneration. (D)
Distribution of S-phase and M-phase cells stained with EdU 1 h pulse labeling

and anti-Phospho-Histone 3 (PH3) antibody. EdU (green) and PH3 (magenta). Yellow dot squares show each quantification area (74.72^2 μm^2) in (E, F). (E, F) The proportion of PH3$^+$ and EdU$^+$ cells per all cells in each area. The regions of bulb (a) and amputation (amp) site were defined in intact tentacles while those of bulb (b) and blastema were defined in the regenerating tentacle at 24 hpa. Each area: n = 14. (G-I) Distribution of proliferative cells at 24 hpa: (G) After the distal side of tentacle is amputated, (H) after removing the bulb from canals, and (I) in isolated tentacles at 24 hpa. EdU 1 h pulse labeling at 24 hpa. Unpaired two-tailed *t*-test. *p < 0.05, **p < 0.005, ***p < 0.001. Scale bars: (A, G-I) 100 μm, (D) 50 μm.

Fig 3. Stem cells and progenitor cells in blastema.
(A) Identification of cell types in blastema stained with anti-β-catenin staining. White arrow: epithelial cell, white arrow head: i-cell, yellow arrow head: nematocyte. Yellow dot square (ii) shows quantification area (35.10^2 μm^2) in (B). Magnified images for each cell type (iii). (B) Cell types in blastema at 24 hpa by anti-β-catenin staining. n = 3 (areas). (C) FISH images of *Nanos1* gene expression from the proximal, abaxial side of the tentacle. Distribution of stem (*Nanos1*$^+$) cells in intact and regenerating tentacle. The tentacles were divided into three parts defined as basal (grey arrows), middle (pink arrows), and distal (red arrows). Yellow arrow indicates main tentacle, and white arrow indicates new branch. (D) Quantification of *Nanos1*$^+$ cells distribution in tentacles based on the defined areas in (C). Intact: n = 5, 24 hpa: n = 4 (tentacles), 48 hpa: n = 4, 72 hpa: n = 5. (E) Images of the intact and regenerating tentacle co-labeled with *Nanos1* FISH and EdU 1 h pulse labeling. EdU (green) and *Nanos1* (magenta). White squares show the quantification area (74.72^2 μm^2) in (3F and S4E). Yellow arrowheads indicate EdU$^+$ and *Nanos1*$^+$ cells. (F) The ratio of *Nanos1*$^+$ cells / EdU$^+$ cells in intact and regenerating tentacles. The averages of *Nanos1*$^+$ cells / EdU$^+$ cells, intact bulb: 69.7% (n = 18), blastema: 30.3% (n = 22). Unpaired two-tailed *t*-test. *p < 0.05, **p < 0.005, ***p < 0.001. Scale bars: (Ai, C, Ei and Eiii) 50 μm, (Eii and Eiv) 25 μm, (Aii) 10 μm.

Fig 4. Distinct proliferative cell populations during tentacle regeneration.
(A) Scheme of lineage-tracing experiments using EdU and BrdU in (B-D). Resident homeostatic stem cells (RHSCs) are labeled by EdU before amputation; repair-specific proliferative cells (RSPCs) are labeled by BrdU at 24

hpa. (B) Distribution of EdU$^+$ and BrdU$^+$ cells during tentacle regeneration. EdU
(green) and BrdU (red). (C) The magnified images of bulb and blastema in (Bi).
White arrowheads indicate EdU$^+$ and BrdU$^+$ cells. (D) Quantification of
EdU$^+$/BrdU$^+$ cells in (C). Each quantification area size is 74.72^2 μm^2. The ratio
of EdU$^+$/BrdU$^+$ cells. bulb: n = 10, blastema: n = 22. (E) The number of RHSCs
during regeneration. Intact: n = 20 (tentacles), 24 hpa: n = 19, 48 hpa: n = 20,
72 hpa: n = 16. (F) The number of RSPCs during regeneration. 24 hpa: n = 12,
48 hpa: n = 12, 72 hpa: n = 6. For (E) and (F), quantification area is the whole
regenerating tentacle. (G) Scheme of two proliferative cell populations during
regeneration. (H) Scheme of stem cell elimination by X-ray irradiation (75 Gy) or
drug treatments (HU 10 mM and Mitomycin C 30 μM) followed by labeling
blastema with EdU (I-J). Tentacle amputation at 3 days post-irradiation/after
treatment of drugs. (I) Distribution of proliferative cells after X-ray irradiation or
drug treatments. EdU 1 h pulse labeling at 24 hpa. Yellow dot squares show
each quantification area (74.72^2 μm^2) in (J). (J) The ratio of EdU$^+$ cells in
blastema. Control: n = 12, X-ray: n = 13, HU: n = 12, Mitomycin C: n = 20.
Unpaired two-tailed *t*-test. *$p < 0.05$, **$p < 0.005$, ***$p < 0.001$. Scale bars: (B, I)
100 μm, (C) 20 μm.
1265
Fig 5. Contribution of resident homeostatic stem cells to nematogenesis
and tentacle elongation.
(A) Scheme of EdU long-time (24 hours) and BrdU double chasing experiment
in (B-C). (B) The distributions of EdU$^+$ and BrdU$^+$ cells during tentacle
regeneration. EdU (green) and BrdU (red). Yellow arrows indicate accumulation
of EdU$^+$ cells at nematocyte clusters in distal side at 72 hpa. White arrowheads
indicate EdU$^+$/BrdU$^+$ cells. Yellow dot squares show magnified views (iv, v). (C)
Quantification of EdU$^+$/BrdU$^+$ cells (%) in the blastema region (Biv). The
quantification area is 74.72^2 μm^2. n = 29. (D) Cell type identification by anti-
aPKC staining. i-cells, nematocytes, and epithelial cells, similar to the anti-β-
catenin staining (see Fig 3A). (E) Scheme of experiments (F-G). RHSCs are
labeled by EdU 24 hours incorporation before amputation and anti-aPKC
staining at 72 hpa. (F) Cell type identification of EdU labeled cells in distal side
at 72 hpa by anti-aPKC staining. EdU (green) and aPKC (magenta). White
arrowheads show nematocytes. (G) Quantification of the cell types of EdU$^+$ cells
on the distal side of a regenerating tentacle at 72 hpa. Quantification area of
regenerating tentacle is 74.72^2 μm^2. n = 10. (H) Experimental scheme for

 tentacle regeneration after X-ray irradiation (75 Gy) or drug treatments (HU 10

mM and Mitomycin C 30 µM) in (I-K). Tentacle amputation at 3 days post-

irradiation or after 3 days treatment of drugs. (I) Representative images of

regenerating tentacles at 3 dpa. Yellow arrowheads indicate nematocyte

clusters. (J) The timing of nematocyte cluster formation in regenerating

tentacles. Control: n = 48, HU: n = 51, Mitomycin C: n = 51, X-ray: n = 60. (K)

Length of regenerating tentacle after irradiation and drug treatments. Control: n

= 48, HU: n = 51, Mitomycin C: n = 51, X-ray: n = 60. Unpaired two-tailed t-test.

*p < 0.05, **p < 0.005, ***p < 0.001. Scale bars: (I) 1 mm, (Bi-iii) 100 µm, (Biv

and Bv) 50 µm, (F) 25 µm.

1293

1294 Fig 6. Contribution of repair-specific proliferative cells to tentacle

1295 elongation.

(A) Experimental scheme for combining the identification of cell types by anti-

aPKC staining and EdU chasing in (B-C). (B) Identification of cell types of EdU$^+$

cells in blastema at 24 hpa and the distal side of the regenerating tentacle at 72

hpa by anti-aPKC staining. EdU (green) and aPKC (magenta). White arrow:

epithelial cell, white arrow head: i-cell. Magnified views (iii, iv). (C) Quantification

of cell types in EdU$^+$ cells at 24 and 72 hpa. Quantification area of regenerating

tentacle is 35.06^2 µm^2. 24 hpa: n = 9, 72 hpa: n = 12. (D) Experimental scheme

depicting blastema monitoring after UV exposure in (E-F). EdU 1 h

incorporation at 24 hpa and UV exposure around the blastema region. (E)

Distribution of EdU labeled cells during regeneration, Ctl (no UV) vs UV. (F) The

number of EdU$^+$ cells from 24 hpa to 72 hpa, Ctl (no UV) vs UV. 24 hpa: n = 11

(tentacles), 48 hpa Ctl: n = 18, 48 hpa UV: n = 20, 72 hpa Ctl: n = 10, 72 hpa

UV: n = 10. (G) Experimental scheme for measuring of regenerating tentacle

after UV exposure to blastema at 24 hpa (H & I). (H) Representative images of

regenerating tentacles at 96 hpa. (I) Length of the regenerating tentacle after

UV exposure on blastema. Ctl: n = 12 (tentacles), UV: n = 12. Unpaired two-

tailed t-test. *p < 0.05, **p < 0.005, ***p < 0.001. Scale bars: (H) 1 mm, (E) 100

µm, (Bi and Bii) 50 µm, (Biii and Biv) 10 µm.

1314

1315 Fig 7. Summary of *Cladonema* tentacle regeneration and evolutionary

1316 history of appendage regeneration.

(A) When a *Cladonema* medusa tentacle is amputated, wound closure is

completed by 24 hpa. After wound healing, repair-specific proliferative cells

(RSPCs) appear and actively proliferate at the wound tip. Ectopic cell
proliferation on the distal side is gradually suppressed, and the cell
differentiation phase begins. While resident homeostatic stem cells (RHSCs) in
the bulb supply nematocytes and epithelial cells, RSPCs mainly differentiate
into epithelial cells. (B) Blastema formation mechanism during appendage
regeneration across animal evolution. When an appendage is amputated,
RSPCs (stem/progenitor cells) contribute to blastema formation in both
regenerative bilaterians (e.g. salamanders) and cnidarian jellyfish.
1327
Supplementary Figures
Fig S1. Wound healing, neuron regeneration and the rate of regeneration.
(A) Representative images of nematocyte clusters (NC) in the medusa tentacle.
Yellow arrows indicate nematocyte clusters that include mature nematocytes
(Poly-γ-glutamate$^+$). (B) The process of wound healing from 4 hpa to 24 hpa
with Phalloidin and anti-α-Tubulin antibody. Phalloidin for F-actin (red) and α-
Tubulin (green). (C) The extent of wound healing during tentacle regeneration.
Full open: actin fibers not attached (Bi), Partially close: disorganized actin fibers
(Bii and Biii), Completed close: actin fibers are fully attached (Biv). 4 hpa: n = 15
(tentacles), 8 hpa: n = 17, 16 hpa: n = 15, 24 hpa: n = 14. (D) Neural
morphology in intact tentacle and the regenerating tentacle stained with the
anti-FMRFamide antibody. Yellow allows indicate cell bodies of FMRFamide
neurons. FMRFamide (magenta). (i, ii) Intact whole tentacle, (iii, iv) basal side of
intact tentacle, (v, vi) regenerating tentacle at 24 hpa. (E) Neural morphology in
the regenerating tentacle stained with the anti-acetylated-Tubulin antibody.
White arrows indicate neural fibers. acetylated-Tubulin (green). (F) The rate of
the tentacle regeneration after removing the bulb from canals. Difference of the
regeneration rate between each experiment. (G) Images of the isolated tentacle
without bulb at 1 dpa and 7 dpa. (H) Images of the isolated whole tentacle 1 dpa
and 7 dpa. Scale bars: (A, Di, Diii, Dv and Ei, Eiii) 100 μm, (Div, Dvi and Eii, Eiv)
50 μm, (G, H) 1 mm.
1349
1350 **Fig S2. The distribution of mitotic cells during regeneration.**
1351 (A) Distribution of mitotic cells detected by anti-PH3 in regenerating tentacle. (B)
1352 Quantification of PH3$^+$ cells' distribution in regenerating tentacle based on the
1353 defined areas in (A). Intact: n = 11 (tentacles), 24 hpa: n = 10, 48 hpa: n = 9, 72
1354 hpa: n = 8. (C) Distribution of S-phase cells in regenerating tentacle stained with

EdU 1 h pulse labeling. EdU 1h pulse labeling was performed at 0, 4, 8, 12, 16, 20, and 24 hpa as shown in the scheme. White dot squares show each quantification area (150^2 μm^2) in (D, E). (D) EdU relative intensity in blastema. 0 hpa: n = 6, 4 hpa: n = 7, 8 hpa: n = 10, 12 hpa: n = 11, 16 hpa: n = 10, 20 hpa: n = 26, 24 hpa: n = 12. (E) The number of EdU$^+$ cells in blastema. 0 hpa: n = 6, 4 hpa: n = 7, 8 hpa: n = 10, 12 hpa: n = 11, 16 hpa: n = 10, 20 hpa: n = 26, 24 hpa: n = 12. Scale bars: (A, C) 100 μm.

Fig S3. The distribution of stem cells in intact tentacle.
(A-D) Expression of *Nanos1, Nanos2, Piwi,* and *Vasa1* in intact tentacles by FISH. Yellow arrows indicate the branching site. (E) Expression of *Nanos1* and *Nanos2* in intact tentacle by double FISH. White arrowheads indicate *Nanos1$^+$,* and white arrows indicate *Nanos2$^+$*. (F) Expression of *Nanos2* and *Mcol1* in intact tentacle by double FISH. Yellow arrows indicate representative co-expression of *Nanos2* and *Mcol1*. Scale bars: (A-D) 100 μm, (Ei, Fi) 50 μm, (Eii, Fii) 25 μm.

Fig S4. The distribution of stem-like cells in regenerating tentacle.
(A, B) Distribution of cells with stem marker genes (*Piwi$^+$* or *Vasa1$^+$*) in intact tentacle and regenerating tentacle by FISH. Yellow arrow indicates the branching site. (C, D) Images of the regenerating tentacle co-labeled with *Piwi* or *Vasa1* FISH and EdU 1 h pulse labeling at 24 hpa. EdU (green) and *Piwi* or *Vasa1* (magenta). White squares show each quantification area (74.72^2 μm^2) in (E). Yellow arrowheads indicate EdU$^+$/*Piwi$^+$* or EdU$^+$/*Vasa1$^+$* cell. (E) The rate of cells positive for stem cell marker genes in blastema. *Nanos1*: n = 22, *Piwi*: n = 12, *Vasa1*: n = 11. Scale bars: (Aiv, Biv) 100 μm, (Ai-iii, Bi-iii, Ci and Di) 50 μm, (Cii and Dii) 25 μm.

Fig S5. Distribution and increase of resident homeostatic stem cells (RHSCs) during tentacle regeneration.
(A) Transition of labeled RHSCs during regeneration by chasing EdU$^+$ cells. EdU 1 h labeling before amputation and chasing at 0, 4, 8, 12, 16, 24, 48, 72 hpa. White dot squares show each quantification area (150^2 μm^2) in (B). (B) The number of EdU$^+$ cells around blastema during regeneration. 0 hpa: n = 6, 4 hpa: n = 6, 8 hpa: n = 13, 12 hpa: n = 12, 16 hpa: n = 8, 24 hpa: n = 10. (C) The comparison of proliferative cell number in intact vs no amputation vs 24 hpa.

EdU 1 h pulse labeling before amputation and chasing with amputation or without amputation (no amp). (D) The number of EdU$^+$ cells labeled before amputation. Counted area is the whole regenerating tentacle at 24 hpa. Detailed information is in (E). Intact: n = 6 (tentacles), 24 h (no amp): n = 6, 24 hpa: n = 6. (E) Area size used for quantification. Note that the area of quantification at 24 hpa is similar. Unpaired two-tailed t-test. *$p < 0.05$, **$p < 0.005$, ***$p < 0.001$. Scale bars: (A, C) 100 μm.

Fig S6. DiI injection revealed that resident stem cells do not migrate to blastema during regeneration.

(A) Experimental scheme of DiI labeled cell identification in the bulb with EdU or anti-β-catenin staining in (B). (B) Co-staining of DiI and EdU or anti-β-catenin in intact tentacle. White dot squares show each quantification area (74.72^2 μm^2) in (C). (C) Rate of DiI labeled cells per proliferative cells or i-cells in bulb. Quantification of DiI$^+$/EdU$^+$ cells or DiI$^+$/i-cells (β-catenin cytoplasmic signal$^+$ cells). DiI$^+$/EdU$^+$ cells: n = 5 (areas), DiI$^+$/i-cells: n = 5. (D) Experimental scheme of chasing DiI-labeled cells in (E). EdU 1h pulse labeling at 24 hpa. (E) (i and ii) Little migration of DiI labeled cells in bulb from the moment of amputation to 24 hpa. n = 9/9. (iii and iv) DiI labeled cells in center area that merged with EdU (yellow arrow). n = 10/12. (F) Experimental scheme for chasing DiI-labeled cells in the bulb in (G). (G) Distribution change of DiI labeled cells in the bulb during tentacle regeneration of an animal. White arrow indicates migration of labeled cells to nematocyte cluster at distal side. n = 6/10. (H) Experimental scheme for chasing DiI-labeled cells in the central area of tentacle in (I). (I) Distribution change of DiI-labeled cells in central area during tentacle regeneration of an animal. White arrow indicates DiI-labeled cells in the newly regenerated tentacle. n = 17/19. Scale bars: (G, I) 250 μm, (Bi, Biii, E) 100 μm, (Bii, Biv) 50 μm.

Fig S7. *Nanos1*$^+$ cells in the bulb decrease after X-ray irradiation or drug treatments.

(A) Experimental scheme of X-ray irradiation (30, 50, 75 Gy). FISH and qPCR at 3 days post-irradiation (3 dpi). (B) Expression of *Nanos1* in intact tentacle at 3 dpi by FISH. (C) Quantification of *Nanos1*$^+$ cells number in tentacle. Quantification area is the entire tentacle in confocal images. Control: n = 9, 30 Gy: n = 5, 50 Gy: n = 5, 75 Gy: n = 17. (D) Experimental scheme of drug

treatments (HU 10 mM or Mitomycin C 30 μM). FISH and qPCR after 3 days of
treatment. (E) Expression of *Nanos1* in intact tentacle after 3 days drug
treatments by FISH. (F) Quantification of *Nanos1*$^+$ cells in tentacle.
Quantification area is the entire tentacle in confocal images. Control: n = 9, 75
Gy: n = 17, HU: n = 10, Mitomycin C: n = 8. (G) Relative expression of *Nanos1*
after irradiation and drug treatments by qPCR. Unpaired two-tailed *t*-test. *$p <$
0.05, **$p <$ 0.005, ***$p <$ 0.001. Scale bars: (B, E) 50 μm.
1434
Fig S8. EdU long incorporation labels nematocyte progenitors.
(A) Nematocyte progenitor cells in the bulb with dual staining of *Mcol1* FISH and
EdU staining. EdU 1 h or 24 h pulse labeling. The expression level of *Mcol1* is
extremely high such that the *Mcol1* signal invades the wavelength of the EdU
signal. White arrows indicate EdU$^+$ cells, and yellow arrowheads indicate
Mcol1$^+$/EdU$^+$ cells. (B) The rate of *Mcol1*/EdU$^+$ cells in intact tentacle.
Quantification area is the entire tentacle in single section of confocal images. 1
h: n = 7 (tentacles), 24 h: n = 7. Unpaired two-tailed *t*-test. **$p <$ 0.005. Scale
bars: (A) 50 μm.
1444
1445 **Fig S9. Similar pattern between β-catenin and aPKC antibody staining.**
1446 Identification of cell types in blastema stained by β-catenin and aPKC antibody
1447 staining. Scale bar: 50 μm.
1448
Fig S10. Neurons are newly-generated from two stem-like cell populations
during regeneration.
(A) Experimental scheme of chasing RHSCs in (B). (B) Neuron derived from
RHSCs by co-staining with EdU and anti-FMRFamide antibody at 72 hpa.
Yellow arrowhead indicates EdU$^+$ & FMRFamide$^+$ cell. (C) Experimental scheme
of chasing RSPCs in (D). (D) Neuron derived from RSPCs by co-staining wtih
EdU and anti-FMRFamide antibody at 72 hpa. Yellow arrowhead indicates
EdU$^+$ & FMRFamide$^+$ cell. (E) (i) The number of neurons in each chasing
experiment. Quantification area is the entire tentacle in confocal images (B ii
and D ii). RSPCs: n = 19 (images), RHSCs: n = 18. (ii) Rate of
EdU$^+$/FMRFamide$^+$ cell in (Ei). Unpaired two-tailed *t*-test. *$p <$ 0.05. Scale bars:
(Bi, Di) 100 μm, (Bii, Dii) 50 μm.
1461
1462 **Fig S11. Blastema exposure to UV has no significant effect on RHSCs.**

(A) Experimental scheme depicting the combination of EdU chasing and UV exposure in (B-D). (B) Bleaching of Hoechst signal only in UV exposure area at 24 hpa. White dot square is UV exposure area. (C) Distribution of EdU labeled cells from intact to 72 hpa, Ctl (no UV) vs UV. (D) The number of EdU$^+$ cells during regeneration, Ctl (no UV) vs UV. 24 hpa: n = 11 (tentacles), 48 hpa Ctl: n = 18, 48 hpa UV: n = 20, 72 hpa Ctl: n = 10, 72 hpa UV: n = 10. (E) Experimental scheme depicting nematogenesis monitoring after UV exposure in (F-G). (F) Representative images of Ctl vs UV at 72 hpa. Yellow arrowheads show nematocyte clusters. (G) Timing of nematocyte cluster formation during tentacle regeneration, Ctl vs UV. Each tentacle: n = 12. Unpaired two-tailed t-test. ***$p < 0.001$. Scale bars: (F) 500 μm, (B, C) 100 μm.

Figure 1

Figure 2

Figure 3

Figure 4

A Experiment scheme (Fig 4B-4D)

Chasing

−1   0   24   48   72   hpa

EdU 1 h   Amp   BrdU 1 h

Labeling RHSCs in bulb   Labeling RSPCs in blastema

B DNA EdU BrdU   24 hpa   48 hpa   72 hpa

(i)   (ii)   (iii)

C (i)   C (ii)

EdU   (i')   (ii')   (iii')

BrdU   (i'')   (ii'')   (iii'')

C DNA EdU BrdU

Bulb (i)

Blastema (ii)

D EdU+ cells / BrdU+ cells (%)

Bulb   Blastema

E RHSCs — # of cells labeled before amputation

Intact   24 hpa   48 hpa   72 hpa

***   n.s.   *   n.s.

F RSPCs — # of cells labeled at 24 hpa

24 hpa   48 hpa   72 hpa

***   n.s.

G 24   48   72   hpa

● Resident homeostatic stem cells (RHSCs)
● Repair-specific proliferative cells (RSPCs)

H Experiment scheme (Fig 4I-4J)

X-ray

Drugs treatment (for 3 day)

−72   −48   −24   0   24   hpa

Amputation   EdU 1 h

Stem cell elimination   Blastema formation?

I DNA EdU

No irradiation (i)   X-ray (ii)   HU (iii)   Mitomycin C (iv)

EdU   (i')   (ii')   (iii')   (iv')

J % EdU+ cells

Control   X-ray   HU   MitomycinC

n.s.   **   ***

Figure 5

A Experiment scheme (Fig 5B-5C)

B Intact / 24 hpa / 72 hpa — DNA EdU BrdU

C EdU+ cells /BrdU+ cells (%)

D DNA aPKC — Nematocytes / Epithelial cells / i-cells

E Experiment scheme (Fig 5F-5G)

F 72 hpa — DNA EdU aPKC

G % cell types — Nematocytes 45.7, Epithelial cells 24.2, i-cells 30.1

H Experiment scheme (Fig 5I-5K)

I 3 dpa — No irradiation / X-ray / HU / Mitomycin C

J % tentacle with nematocytes vs dpa — Control, X-ray, HU, Mitomycin C

K Regenerated Tentacle length (mm) vs dpa — Control, X-ray, HU, Mitomycin C

Figure 6

A Experiment scheme (Fig 6B-6C)

B DNA EdU aPKC | 24 hpa | 72 hpa

C % cell types — Nematocytes, Epithelial cells, i-cells; 24 hpa (94.4, 4.7, 0.9), 72 hpa (40.8, 59.2)

D Experiment scheme (Fig 6E-6F)

E Ctl — DNA EdU / EdU: 24 hpa, 48 hpa, 72 hpa; UV — DNA EdU / EdU: 48 hpa, 72 hpa

F # of EdU+ cells — 24 hpa, 48 hpa Ctl, 48 hpa UV, 72 hpa Ctl, 72 hpa UV

G Experiment scheme (Fig 6H-6I)

H 96 hpa — Ctl, UV

I Regenerated Tentacle length (mm) vs hpa — Ctl, UV

Figure 7

A

B Appendage regeneration evolution model