1

Distinct stem-like cell populations facilitate functional regeneration of

the *Cladonema* medusa tentacle

[1] Sosuke Fujita**,** [1] Erina Kuranaga, [2] Masayuki Miura, and [2] Yu-ichiro Nakajima*****

1. Graduate School of Life Sciences, Tohoku University

2. Graduate School of Pharmaceutical Sciences, The University of Tokyo

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp

Tel: +81-3-5841-4863

Running title: Distinct stem-like cells enable tentacle regeneration in jellyfish

Main text: 6352 words (Abstract: 186 words)

Figures: 6   **Supplementary information** (Figures: 8)

Keywords: regeneration, blastema, non-bilaterians, cnidarians, jellyfish, tentacle, i-cell

Abstract

Blastema formation is a crucial process that provides a cellular source for regenerating tissues and organs. While bilaterians have diversified blastema formation methods, its mechanisms in non-bilaterians remain poorly understood. Cnidarian jellyfish, or medusae, represent early-branching metazoans that exhibit complex morphology and possess defined appendage structures highlighted by tentacles with stinging cells (nematocytes). Here we investigate the mechanisms of tentacle regeneration, using the hydrozoan jellyfish *Cladonema pacificum*. We show that proliferative cells accumulate at the tentacle amputation site and form a blastema composed of cells with stem-like cell morphology. Lineage tracing experiments indicate that repair-specific proliferative cells in the blastema are distinct from resident stem-like cells. We further demonstrate that resident stem-like cells control nematogenesis and tentacle elongation during both homeostasis and regeneration, while repair-specific proliferative cells preferentially differentiate into epithelial cells in the newly formed tentacle, analogous to lineage-restricted stem/progenitor cells observed in salamander limbs. Taken together, our findings reveal a regeneration mechanism that utilizes both resident stem-like cells and repair-specific proliferative cells, which in conjunction efficiently enable functional appendage regeneration, and provide novel insight into the diversification of blastema formation across animal evolution.

Introduction

Regeneration, the phenomenon of re-forming missing body parts, is widespread among metazoans. Common regeneration processes include wound closure immediately after injury; formation of the cellular source, or blastema, that reconstitutes the lost tissue; and regrowth of the tissue that integrates different cellular behaviors such as proliferation and differentiation (McCusker et al., 2015). Among these regenerative responses, blastema formation is a critical step that can distinguish regenerative and non-regenerative systems. Indeed, most mammalian and avian species do not form blastema upon injury while wound closure normally occurs (Seifert and Muneoka, 2018). Understanding blastema formation mechanisms in highly regenerative animals may thus help us identify the necessary elements to potentially improve our regenerative abilities.

Blastema is defined as undifferentiated cells with mitotic capacity that appear after damage such as amputation or irradiation (Aztekin, 2021; Bely and Nyberg, 2010). Accumulating evidence has suggested that methods of blastema formation vary among animals with high regenerative abilities (Tanaka and Reddien, 2011). For example, in planarians, pluripotent stem cells called neoblasts that are distributed throughout the body, are recruited to the injury site to produce blastema (Reddien and Alvarado, 2004). Salamanders can regenerate adult limbs upon amputation via blastema formation, but the underlying cellular mechanisms vary among species: in the axolotl, tissue-specific stem cells contribute to blastema, while in the newt, muscle fibers dedifferentiate into proliferative progenitor cells to behave as blastema (Sandoval-Guzmán et al., 2014). During zebrafish caudal fin regeneration, both osteoblast-derived dedifferentiated cells and resident progenitor cells migrate to the wound site to form blastema (Ando et al., 2017; Knopf et al., 2011). These studies support the idea that the supply of blastema has diversified across the animal kingdom, whose members utilize resident stem/progenitor cells and/or repair-specific *de novo* proliferative cells to reconstruct lost body parts. While mechanisms of blastema formation have been intensively studied in the limited number of regenerative animals, little is known about their evolutionary characteristics: which elements are acquired as lineage-specific novelties and which are widely conserved within highly regenerative species? In particular, the current

understanding of blastema formation largely relies on bilaterian models, and thus the

mechanisms of blastema formation outside of bilaterians remain poorly understood.

Among various regeneration contexts, appendage regeneration is widely

observed in bilaterians (e.g. amphibian limbs and fish fins), and is suitable for

understanding the evolution of regenerative processes (Aztekin, 2021; Daponte et al.,

2021). The bilaterian program of blastema formation during appendage regeneration is

associated with repair-specific stem/progenitor cells (Knopf et al., 2011; Lehoczky et al.,

2011; Sandoval-Guzmán et al., 2014). Yet, to elucidate the evolutionary history of

blastema formation programs, their mechanisms must be studied in early-branching

metazoans in addition to bilaterian models.

Cnidarians (corals, sea anemones, hydroids, and jellyfish) are among the earliest

branching metazoans, composed of two major groups Anthozoa and Medusozoa,

forming a diverse phylum that contains over 10,000 species, and stand at a unique

phylogenetic position as the sister group to bilaterians (Figure 1A). While cnidarians

display considerably divergent morphologies and life cycles, represented by the polyp

and medusa stages; their common traits are a diploblastic radially-symmetric body

along with tentacles as the well-defined appendages that bear the stinging cells,

nematocytes (cnidocytes) (Fritz et al., 2013; Technau and Steele, 2011). Although

regenerative potential may vary among the group, most documented cnidarian species

are capable of regenerating lost tissues and organs, and some can even regenerate

their entire body (Fujita et al., 2021; Holstein et al., 2003).

Among cnidarians, polyp-type animals such as *Hydra*, *Hydractinia,* and

Nematostella have been utilized as models to understand the mechanism of whole-body

regeneration; including patterning, body axis formation, and mechanical responses

(Gahan et al*.,* 2016; Röttinger, 2021; Vogg et al., 2019). After mid-gastric bisection in

Hydra, cell proliferation of the stem cells, or i-cells, around the wound site is accelerated

by Wnt3a produced from dying cells to generate blastema (Chera et al., 2009). Upon

decapitation of the colony polyp *Hydractinia*, i-cells remotely located in the body column

migrate to the injury site to form blastema (Bradshaw et al., 2015). During head

regeneration of the sea anemone *Nematostella*, two adult stem-like cell populations,

fast-cycling cells in the body wall epithelium and slow-cycling cells in the mesenteries,

migrate toward the amputation site to form blastema (Amiel et al., 2019). These studies

suggest that the recruitment of resident stem cells to the injury site is prerequisite for

blastema formation after amputation of the body. However, it remains unclear whether

repair-specific proliferative cells constitute the cellular source of blastema during

cnidarian regeneration.

In contrast to the sessile polyp stage, medusae, commonly called jellyfish,

exhibit a more complex body structure that includes multiple types of muscle such as

smooth and striated muscles, developed neural networks, and distinct organs and

appendages, which together enable free swimming and avoidance behaviors (Katsuki

and Greenspan, 2013; Leclère and Röttinger, 2017). While most sexually-reproducing

medusae do not clonally propagate like polyps, medusae can still regenerate various

organs and appendages (e.g., manubrium, tentacle, umbrella, gonad, eye) after injury

and reconstitute *de novo* structures after their removal (Fujita et al., 2019; Gamero-

Mora et al., 2019; Ostendarp et al., 2022; Sinigaglia et al., 2020; Stamatis et al., 2018;

Stierwald et al., 2004; Weber, 1981). A recent report using the hydrozoan jellyfish *Clytia*

has suggested that, upon removal of the entire manubrium, i-cells and differentiated

cells localized in neighboring organs migrate to the damage site through the canals,

contributing to *de novo* manubrium regeneration (Sinigaglia et al., 2020). Despite the

importance of proliferating blastema cells in highly regenerative animals, the detailed

mechanisms of organ and appendage regeneration in medusae, particularly the

mechanism of blastema formation and its specific role are largely unknown.

The hydrozoan jellyfish *Cladonema pacificum* is an emerging jellyfish model that

has been utilized to study development and physiology (Fujiki, 2019; Masuda-Ozawa et

al., 2022; Takeda et al., 2018). Because *Cladonema* allows for easy lab maintenance

with its high spawning rate, it enables monitoring all life cycle stages and exploring

organismal responses to different stimuli. The genus *Cladonema* is morphologically

characterized by branched tentacles at the medusa stage (Figure 1B) (Fujiki, 2019;

Rees, 1979; Suga et al., 2008; Zhou et al., 2022). The *Cladonema* medusa tentacle is

primarily composed of bi-layered epithelial tissues (ectoderm and endoderm): i-cells,

neurons, and nematocytes are located in the ectodermal layer while neurons and gland

cells are located in the endodermal layer. Resident stem-like cells, i-cells, are localized

at the basal side of the tentacle (tentacle bulb), and are thought to give rise to

progenitors and differentiated cells during normal development and homeostasis

(Denker et al., 2008; Fujita et al., 2019). The continuous growth and branching potential
of the medusa tentacle makes it an attractive model to understand the mechanisms of
appendage morphogenesis, growth, and regeneration in cnidarians.

In this study, we investigate the cellular mechanism of appendage regeneration
using the *Cladonema* medusa tentacle. Establishing the *Cladonema* tentacle as an
organ that can efficiently and functionally regenerate upon amputation, we show that
highly proliferative cells accumulate at the injury site within 24 hours to behave as
blastema. Lineage tracing experiments using nucleoside analogs as well as dye labeling
reveal that blastema cells are not derived from resident i-cells but rather appear locally
after damage. We further identify the role of blastema cells as the principal cellular
source of the epithelium in the newly formed tentacle, while resident i-cells contribute to
nematogenesis and tissue elongation during both homeostasis and regeneration. These
results suggest the existence of two distinct proliferative cell populations: blastema as
repair-specific proliferative cells and resident i-cells as homeostatic proliferative cells,
both of which collectively enable functional tentacle regeneration. In a broader context,
our findings highlight the diversification of blastema formation in non-bilaterian systems,
providing an evolutionary insight into animal regeneration.

172
173

Materials and Methods

Animal cultures and surgical manipulations
Cladonema pacificum (female strain 6W) medusae were used for this research. The
medusae were maintained in plastic containers (V-type container, V-7, V-8, and V-9; AS
ONE) at 22 ℃ in artificial sea water (ASW). ASW was prepared using SEA LIFE (Marin
Tech) dissolved in tap water with chlorine neutralizer (Coroline off; GEX Co. ltd) (24
p.p.t). The medusae were fed Vietnamese brine shrimp (A&A Marine LLC).

Before surgical manipulation, the medusae were anesthetized with 7% $MgCl_2$ in
deionized water for 2 min in a petri dish (Falcon, #351008). Surgical manipulations were
performed with micro scissors (Teraoka, NY33408). After manipulations, the medusae
were quickly returned to ASW. When the tentacle was amputated with the bulb left
intact, the medusae were fed 1 day before manipulation and were starved after

amputation. When the tentacle was amputated along with the bulb, the medusae were

fed 1 day before manipulation and were fed 3 times a week afterward.

Bright field pictures of medusae were taken with a LEICA S8APO microscope

with a Nikon digital camera (D5600).

Immunofluorescence

The medusae were anesthetized with 7% $MgCl_2$ in deionized water for 5 min

and fixed at room temperature (R.T.) for 1 h or overnight at 4 ℃ with 4%

paraformaldehyde (PFA) in ASW. After fixation, the samples were washed three times

(10 min each) with PBS containing 0.1% Triton X-100 (0.1% PBT). The samples were

incubated in primary antibodies in 0.1% PBT overnight at 4 ℃. The antibodies used

were rabbit anti-Phospho-Histone H3 (Ser10) (1:500; Upstate, 06–570), mouse anti-β-

catenin (8E4) (1:100; Sigma-Aldrich, ALX-804-260-C100), and mouse anti-PKC ζ C20

(1:100; Santa Cruz, sc-216). After primary antibody incubation, the samples were

washed three times (10 min each) with 0.1% PBT. The samples were incubated with

secondary antibodies (1:500; ALEXA FLUOR 488, 555, 647 Goat anti-mouse IgG,

ALEXA FLUOR 488, 555, 647 Goat anti-rabbit IgG; Thermo Fisher Scientific, A28175,

A-11008) and Hoechst 33342 (1:500; Thermo Fisher Scientific, H1399) in 0.1% PBT for

1 h in the dark. After three washes (10 min each) in 0.1% PBT, the samples were

mounted on slides with 70% glycerol. Nuclear and poly-γ-glutamate were stained with

DAPI (1:250; Invitrogen, D1306), and actin fibers were stained with Alexa 546 phalloidin

(1:400; Invitrogen, A22283) in 0.1% PBT for 1 h. Confocal images were collected

through a Zeiss LSM 880 confocal microscope. Image processing (color display) and

quantification were performed using ImageJ/Fiji software.

EdU labeling

The medusae were incubated with 20 or 150 µM 5-ethynyl-2'-deoxyuridine (EdU) (EdU

kit; Invitrogen, C10337) in ASW for 1h or 24 h. In chase labeling, the medusae were

washed with ASW at least twice (up to three times, depending on amount). In pulse

labeling, after EdU treatment, the medusae were anesthetized with 7% $MgCl_2$ in

deionized water and fixed 4% PFA in ASW. After fixation, the samples were washed

and incubated with a EdU reaction cocktail (1× reaction buffer, $CuSO_4$, Alexa Fluor

azide 488 or 647, and 1× reaction buffer additive; all included in EdU kit; Invitrogen,

C10337 or C10340) for 30 min in the dark. After the EdU reaction, the samples were

washed and incubated with Hoechst 33342 in 0.1% PBT for 30 min.

222   In combination with antibody staining, EdU reaction was performed after

incubation with the secondary antibody. To combine EdU labeling with fluorescent *in*

situ hybridization (FISH), the EdU reaction was performed after the incubation with Cy5-

tyramide solution. In combination with DAPI poly-γ-glutamate staining, the EdU reaction

was performed before incubation with DAPI.

EdU and BrdU dual labeling

The medusae were incubated with 150 µM EdU in ASW for 1h or 24 h before

amputation, depending on the experiments. Just after incorporation of EdU, the

medusae were washed in ASW three times, and tentacles were amputated. At 24 hpa,

the medusae were incubated with 2.5 mM BrdU in ASW for 1h. The medusae were

washed out in ASW three times.

234   The medusae were anesthetized with 7% $MgCl_2$ for 5 min and fixed with 4%

PFA in ASW. The samples were washed three times with 0.1% PBT and incubated with

a EdU reaction cocktail for 30min in dark. The samples were washed three times and

treated with 2N HCl for 30min and then washed three times and incubated in primary

antibodies in 0.1% PBT overnight at 4 ℃. The primary antibody used was mouse anti-

BrdU (1:100; BD, #347580). The samples were washed three times and incubated with

secondary antibodies (1:500; ALEXA FLUOR 555 Goat anti-mouse IgG) and Hoechst

33342 in 0.1% PBT for 1 h in the dark and washed three times.

Fluorescent *in situ* hybridization

The following FISH protocol was created from the published protocol (Fujita et al.,

2022). The sequence of *Nanos1* and *Piwi* was previously described (Hou et al., 2021),

the presumptive sequence of *Minicollagen1* (*Mcol1*) was acquired from Dr. Kumano,

and *Nematocyte Outer Wall Antigen* (*Nowa*) was acquired from the annotation of our

RNA-seq results. Purified total RNA was reverse transcribed into cDNA by PrimeScript

II 1st strand cDNA Synthesis Kit (Takara, 6210A). The target gene fragments were

amplified from a cDNA library. The primer sets used for PCR cloning are as follows:

Mcol1: 5′-CTCGTCGGTATTGCCCTCTC-3′ (forward) and 5′-
CCAACCTATCGTGGACGTGT-3′ (reverse), *Nowa*: 5′-GGTCGGTCGAAAGCCACTAA-
3′ (forward) and 5′-CCAGGAGGTGAAAGCGAGTT-3′ (reverse). PCR products were
sub-cloned into the TAK101 vector (TOYOBO). The resulting plasmids were used for
RNA probe synthesis with digoxigenin (DIG) labeling mix (Roche, #11277073910), and
T7 (Roche, #10881767001) or T3 RNA polymerase (Roche, #11031163001) was used,
according to the insert direction.

Medusae were anesthetized 7% $MgCl_2$ for 5 min and fixed overnight at 4 ℃ with
4% PFA in ASW. Briefly, fixed samples were washed three times with PBS containing
0.1% Tween-20 (PBST), followed by pre-hybridization in hybridization buffer (HB buffer:
5 × SSC, 50% formamide, 0.1% Tween-20, 50 μg/ml tRNA, 50 μg/ml heparin) at 55 ℃
for 2 h. The samples were hybridized with HB Buffer containing the antisense probes
(final probe concentration: 0.5–1 ng/μL in HB Buffer) at 55 ℃ overnight. The samples
were washed twice each with wash buffer 1 (5 × SSC, 50% formamide, 0.1% Tween-
20), wash buffer 2 (2 × SSC, 50% formamide, 0.1% Tween-20), and 2 × SSC. The
samples were then washed with 0.1% PBST and incubated in 1% blocking buffer (1%
blocking reagent [Roche] in Maleic acid) for 1 h. The samples were incubated with anti-
DIG-POD antibodies (1:500; Roche, #11207733910) in 1% blocking buffer overnight at
4 ℃. The samples were then washed with Tris-NaCl-Tween Buffer and incubated with
Cy5-tyramide solution (TSA Plus Cyanine 5; AKOYA Biosciences, NEL745001KT) for
10 min. Finally, the samples were washed with 0.1% PBST and incubated with Hoechst
33342 in 0.1% PBST for 30 min in the dark and washed three times.
273

274 **DiI injection**

Medusae were relaxed in 7% $MgCl_2$ on a petri dish. 10 mM CellTracker CM-DiI
(Invitrogen, C7001) was injected into the ectodermal layer of the tentacle bulb, using a
micro injector (Eppendorf, Femtojet 4i). The quartz capillary (Sutter Instrument, QF100-
70-10) was pulled by Laser-Based Micropipette Puller P2000 (Sutter Instrument). After
injection, injected tentacles were amputated. At 24 hpa, the injected medusae were
incubated with 150 μM EdU in ASW for 1h. The medusae were relaxed in 7% $MgCl_2$ and
fixed in 4% PFA. After washing in 0.1% PBT, the medusae were incubated with an EdU

reaction cocktail for 30 min in the dark. After the EdU reaction, the samples were

washed and incubated with Hoechst 33342 in 0.1% PBT.

X-ray irradiation

The medusae were placed in a V7 plastic container with 6-8 ml of ASW, the minimum

required for their maintenance. X-ray irradiations were performed using an X-ray

machine (mediXtec, MX-160Labo). Medusae were positioned 19 cm from the X-ray

source, and a dose of 30 Gy (19.3 minutes at 160 kV 3 mA) was delivered. After

irradiation, the medusae were moved to a plastic container with fresh ASW.

qPCR

Total RNA was purified from the whole bodies of three medusae with RNeasy Mini kits

(Qiagen). Lysate was treated with DNase I (Qiagen, #79254) for 15 min at RT. cDNA

was synthesized from 200 ng of total RNA by Prime script RT master Mix (Takara,

RR036A). RT-qPCR was performed using TB Green Premix Ex TaqII (Tli RNaseH Plus)

(Takara, RR820L) and a Quantstudio6 Flex Real-Time PCR system (Thermo Fisher)

using *F-actin capping protein subunit beta* as an internal control (Masuda-Ozawa et al.,

2022). The qPCR primer sets are as follows: *CpCapZbeta*: 5'-

AAAGAAAGCTGGAGACGGTTCA-3' (Forward), 5'-GTAGTGGGCATTTCTTCCGC-3'

(Reverse), *Nanos1*: 5'-TTCGTCAAGTGGCAGTCGTG-3' (Forward), 5-

CAAGCCCTGGTACAAACGGA-3' (Reverse).

UV laser exposure

Tentacles were amputated at 1 day before UV exposure. At 24 hpa, the medusae were

incubated with Hoechst 33342 (1:250) for 20 min and washed three times in ASW. After

washing, the relaxed medusae were placed on slides with 7% $MgCl_2$ and enclosed with

cover glass. UV laser was exposed to the blastema region of the amputated tentacle

using the photo bleaching application (50 cycles at 100% 405 laser power) of the Zeiss

LSM 880 confocal microscope. The bleached region is shown with a white dot square

(400 x 400 pixels, W x H) in Figure S8B. After UV exposure, the medusae were moved

to ASW in plastic containers and monitored to examine the effect on regeneration.

Results

For cnidarians, tentacles are essential and common appendages for capturing prey and defending against predators. Cnidarian polyps such as *Hydra*, *Hydractinia*, and *Nematostella* have tentacles in the head region, near the hypostome, which can restore functional tentacles during head regeneration (Bradshaw et al., 2015; Passamaneck and Martindale, 2012; Vogg et al., 2019). Similarly, studies using hydrozoan jellyfish have shown that medusae tentacles can regenerate upon amputation (Fujita et al., 2019; Sinigaglia et al., 2020). Although cnidarians commonly exhibit high regenerative capacity for tentacles, the exact cellular processes and the limitation of regeneration are poorly understood.

To monitor the process of tentacle regeneration and to explore its responses to different dissections, we utilized the medusa tentacle of the hydrozoan jellyfish *Cladonema* (Figure 1). In *Cladonema* tentacles, proliferative cells including stem-like cells, or i-cells, are localized on the basal side, called the tentacle bulb, while nematocytes are distributed as clusters on the distal side, similar to the jellyfish *Clytia* (Figures 1C, 2A, 3C and S3A-S3C) (Denker et al., 2008; Fujita et al., 2019; Hou et al., 2021). The existence of localized stem-like cells at the bulb prompted us to examine tentacle regeneration as a model for appendage regeneration in non-bilaterians.

Cladonema medusa tentacles functionally regenerate upon amputation

When the tentacle was amputated with the bulb left intact (Figures 1D and 1E), wound closure finished by 24 hours post-amputation (hpa) (Figures S1A and S1B), at which point the injury site became smooth surface (Figures 1D i and 1D ii). Subsequently, tentacles began to elongate distally, and nematocyte clusters formed at the tip of the regenerating tentacle at 48 hpa (Figure 1D iii). The number of nematocyte clusters increased during tentacle elongation, and these clusters were localized along the distal side at 72 hpa, which is when the regenerating tentacle branched, a characteristic feature of the genus *Cladonema* (Figures 1D iv and 1E). Notably, these regeneration processes occurred without feeding, suggesting that tentacle regeneration is an immediate response for the organism to re-construct the organ that is required for capturing food.

To investigate the subtleties of tentacle morphology at cellular resolution during
regeneration, we visualized the muscle fibers, mature nematocytes, and neurons by
staining F-actin, poly-γ-glutamate, and FMRFamide and acetylated tubulin, respectively
(Figures 1F, 1G and S1C, S1D). During the early stages of regeneration (0-24 hpa),
wound closure occurred progressively via the accumulation of supra-cellular actin fibers
(Figures 1F i and 1F ii). Accordingly, muscle fibers regenerated during tentacle
elongation (Figures 1F iii and 1F iv). While mature nematocytes were rarely found in the
bulb or the injury site until 24 hpa (Figures 1F i and 1F ii), nematocytes began to
accumulate at the tip of the regenerating tentacle at 48 hpa and formed clusters at 72
hpa (Figures 1F iii and 1F iv). The re-formation of neurons on the distal side occurred at
48 hpa and completed at 72 hpa (Figures 1G and S1C, S1D). These results suggest
that the regenerating tentacle is fully functional after 48 hpa. To test this possibility, we
examined whether the regenerating tentacle is capable of feeding. While the
regenerating tentacle was not functional until 24 hpa, more than half of the regenerating
tentacles captured prey at 48 hpa (55%), and this ratio reached 100% at 72 hpa
(Figures 1H and 1I). These results indicate that *Cladonema* medusae can fully
regenerate a functional tentacle within 2-3 days of amputation.
While the *Cladonema* medusa tentacle can almost always regenerate with the
bulb left intact, it is unclear whether the tentacle can still regenerate even after the bulb
has been removed (Figure 1J i). We thus ablated the entire tentacle, including the bulb,
and monitored the regeneration process. After amputation, the ring canal and the radial
canal adhered, and the wound closure completed at 1 day post-amputation (dpa)
(Figure 1J ii). The injured area began to bulge as early as 5 dpa, and the transparent
wound site turned orange, which reflects the color of its food, *Artemia* (Figure 1J iii). The
de novo tentacle bulb reformed within 1-2 weeks with a nematocyte cluster at the tip,
and the small tentacle gradually elongated (Figures 1J iv and 1K). Because the rate of
successful regeneration varied dramatically while rearing medusae under the same
conditions and feeding frequency, tentacle bulb regeneration likely depends on the
health condition of the individual medusa (Figure S1E).
We also examined the regenerative potential of isolated tentacles. While the
whole body cannot be regrown from isolated tentacles in *Hydra* or *Nematostella* (Amiel
et al., 2019; Galliot et al., 2018), polyps in the jellyfish *Aurelia aurita* can regenerate

from isolated tentacles (Lesh-Laurie et al., 1991). To examine the regeneration capacity
of the isolated tentacle in *Cladonema*, we cultured *Cladonema* medusa tentacles either
with or without the bulb left intact. Although wound closure successfully occurred
regardless of the existence of the bulb, tissues and organs located to the proximal side
of the tentacle including the umbrella, the radial canal, and the bulb, did not regenerate,
and the isolated tentacle gradually degenerated (Figures S1F and S1G).
These observations together show that the *Cladonema* medusa tentacle can
functionally regenerate after amputation, but that its organ repair capacity is restricted to
the organism that recognizes the missing tentacle, likely through the canals, as
previously shown in the *Clytia* medusa (Sinigaglia et al., 2020). For our remaining
studies, we focused our attention on the process of tentacle regeneration with the bulb
intact, which provides a highly reproducible and tractable system.
390

Blastema is formed in the early phase of tentacle regeneration
During organ regeneration, blastema forms at the regenerating tip after wound closure
and contributes to reconstruction of the new tissue as a source of cell proliferation and
differentiation (McCusker et al., 2015; Seifert and Muneoka, 2018). In order to
investigate the timing of blastema formation during *Cladonema* tentacle regeneration,
we performed pulse labeling of EdU, a S-phase marker (Figure 2A) (Salic and
Mitchison, 2008). To understand the spatial pattern of cell proliferation, we defined
basal, middle, and distal regions in the regenerating tentacle and quantified the
distribution of proliferating cells. Consistent with the localization of stem-like cells
(Figure 3C i), cell proliferation in the intact tentacle was mainly detected on the basal
side (Figures 2A i and 2B). By contrast, cell proliferation in the regenerating tentacle
was frequently observed on the distal side near the wound area at 24 hpa (Figures 2A ii
and B). At 72 hpa, unlike at 24 and 48 hpa, cell proliferation was detected in the middle
and on basal side, not on the distal side (Figures 2A iv and 2B). We further confirmed
the pattern of proliferating cells using the M-phase marker, anti-Phospho-Histone 3
(PH3) antibody. While PH3$^+$ cells were distributed across the basal side before
amputation, in the regenerating tentacle, PH3$^+$ cells accumulated on the distal side at
24 hpa, and gradually localized on the basal side starting at 48 hpa (Figures 2D and
S2), which is consistent with the results of EdU$^+$ cell distribution (Figures 2A and 2B).

These results also reflect the fact that differentiated nematocytes accumulate on the

distal side at 48 hpa and 72 hpa (Figures 1D and 1F). Together, these observations

suggest that cell proliferation is accelerated near the injury site at 24 hpa after

amputation.

Blastema cells are undifferentiated cell populations that possess a high mitotic

capacity during regeneration (Tasaki et al., 2011). We thus investigated proliferative

activity in detail during the early phase of tentacle regeneration (Figures 2D-2F). In the

regenerating tip, the number of PH3[+] and EdU[+] cells was about 4 and 20 times greater,

respectively, than the corresponding area in the intact tentacle (Figures 2D-2F).

Moreover, these PH3[+] and EdU[+] cell numbers at the regenerating tip were significantly

greater than those in the bulb of the intact tentacle where stem-like cells are localized

(Figures 2D-2F). Notably, nearly half of the cells in the regenerating tip are EdU[+] cells

(Figures 2D ii and 2F). Altogether, these results indicate that proliferative activity in the

regenerating tip is much higher than that of cells not in the regenerating tip, which

provides evidence that blastema cells appear during tentacle regeneration.

To determine whether blastema formation is specific to the location of

amputation, we examined the localization of proliferating cells after dissecting tentacles

in different ways. When the distal side of the tentacle is amputated, ectopic cell

proliferation was observed near the injury site at 24 hpa (Figure 2G). Similarly, when the

tentacle was amputated with the bulb, cell proliferation occurred around the injury site

where the canals merged at 24 hpa (Figure 2H). These results suggest that blastema is

formed around the injury site regardless of the location of amputation.

Based on these observations, we hypothesized that blastema is prerequisite for

proper regeneration of the tentacle. To test this possibility, we examined whether

blastema formation occurs in the isolated tentacle, which fails to regenerate basal tissue

(Figure S1F). In the isolated tentacle, EdU[+] cells were not detected at the injury site at

24 hpa (Figure 2I). These results together suggest that blastema formation is

associated solely with the regenerating tentacle.

Stem-like cells compose blastema

What cell types constitute blastema? In hydrozoans, i-cells behave as pluri-/multipotent

stem cells that can differentiate into progenitors and differentiated lineages (Gold and

Jacobs, 2013). In order to confirm the cell types that form blastema, we utilized the

combination of β-catenin antibody staining and nuclear staining, which can distinguish i-

cells, nematocytes, and epithelial cells (Figure 3A). In both *Hydractinia* and *Cladonema*,

i-cells were identified by cytoplasmic signals of β-catenin, together with the large

nucleus-to-cytoplasm ratio and prominent nucleoli (Hou et al., 2021; Plickert et al.,

2012). While nematocytes exhibit crescent-shaped nuclei, epithelial cells show a

polygonal shape delineated by the membrane-localized β-catenin (Figure 3A iii).

Through the quantification of cell types, we found that around 80% of blastema cells

were i-cells, 10% were epithelial cells, and 10% were nematocytes (Figures 3A and 3B),

indicating that most cells in blastema are i-cells.

452   To molecularly characterize these blastema cells, we examined the expression

of the stem cell markers *Nanos1* and *Piwi* by fluorescent *in situ* hybridization (FISH). In

the intact tentacle, *Nanos1*[+] cells are primarily distributed in the bulb of the intact

tentacle (Fujita et al., 2022; Hou et al., 2021). By contrast, at 24 hpa *Nanos1*[+] cells

accumulated around the injury site where blastema formed (Figure 3C ii), and *Piwi*[+]

cells were also localized in the blastema region (Figure S3C). While *Nanos1*[+] cells

accumulated to the new branching bud after 48 hpa (Figure 3C iii, Fujita et al., 2022;

Hou et al., 2021), *Nanos1* expression was no longer detected at the regenerating tip

and was restricted to the middle and basal side of the tentacle (Figure 3C iii). We further

quantified the distribution of *Nanos1*[+] cells and found a gradual transition of their

distribution from the distal to the basal side of the regenerating tentacle (Figure 3D).

These results are consistent with the distribution pattern of proliferative cells (Figures

2A-2C and S2), further supporting the idea that proliferative cells in blastema are stem-

like cells.

Blastema cells are not derived from stem-like cells localized at the bulb

Previous reports on cnidarian regeneration have suggested that undifferentiated

blastema cells are supplied by the migration of resident stem/progenitor cells in

response to injury (Amiel et al., 2019; Bradshaw et al., 2015; Sinigaglia et al., 2020). To

test whether resident stem cells, or originally-localized i-cells, in the bulb migrate to the

injury site to form blastema, we performed EdU-chase experiments during *Cladonema*

tentacle regeneration. We leveraged the fact that short-time incubation of EdU labels

proliferating i-cells in the intact tentacle, and, after incubating, amputated and visualized

the tentacle at different time points (Figure S4A i). At 24 hpa, EdU$^+$ cells remained fairly

distributed across the tentacle (Figure S4A ii). Following 48 h and 72 h chasing, EdU$^+$

cells continued to localize at the basal side, including the bulb, rather than in the

regenerating tip, although the population of EdU$^+$ cells propagated in the distal direction

likely through cell proliferation and migration (Figures S4A iii and S4A iv). We further

conducted an alternative tracing experiment that specifically labels cells in the bulb

region by DiI microinjection followed by amputation and cell tracking (Figures S4B and

S4C). After a 24 h chase, DiI$^+$ cells were still localized near the bulb (Figure S4C ii),

which, taken together with the EdU chase results, suggests that proliferating stem-like

cells do not migrate to the blastema region.

The above experiments support the presence of two populations of proliferative

cells during regeneration: one population are the "homeostatic proliferative cells,"

composed of stem-like cells localized at the bulb, and the other are "repair-specific

proliferative cells" that appear *de novo* as blastema upon amputation. In order to test

this hypothesis, we performed chasing experiments by labeling the two proliferative cell

populations with distinct nucleoside analogs, EdU and BrdU (Figure 4A). This approach

has been utilized in different animal systems as a way to track proliferative cell

populations and their lineages (Harris et al., 2018; Ribeiro et al., 2021; Vega and

Peterson, 2005). We labeled homeostatic proliferative cells in the bulb by EdU before

amputation and repair-specific proliferative cells in the blastema by BrdU at 24 hpa

(Figure 4A). At the time of blastema formation at 24 hpa, EdU$^+$ homeostatic proliferative

cells were still localized to the bulb, and only 12.6% merged with BrdU$^+$ cells, which

accumulated at the blastema (Figure 4B i). Following 24-48 h, BrdU$^+$ cells were

distributed on the distal side of the regenerating tentacle while EdU$^+$ cells remained

localized on the basal side, although their distribution expanded slightly (Figures 4B ii

and 4B iii). These results suggest that repair-specific proliferative cells at the blastema

participate in the formation of the new structure while homeostatic proliferative cells

expand locally, supporting the existence of two distinct proliferative cell populations

during tentacle regeneration (Figure 4E).

To determine the characteristics of these distinct proliferative cell populations,

we examined their respective proliferative capacities. For this purpose, we quantified

the increase in cell numbers of the two populations during tentacle regeneration. At 24

hpa, homeostatic proliferative cells increased approximately two-fold, but no significant

difference was observed in the following 48 h (Figure 4C; 0 hpa: n=161.2±13.5, 24 hpa:

n=392.2±32.1, 48 hpa: n=426.8±28.5, 72 hpa: n=385.7±29.1). Notably, the number of

these proliferative cells also increased during 24 h without amputation (Figures S4D and

S4E), suggesting that the increase of homeostatic proliferative cells is not an injury-

specific response. In contrast, the number of repair-specific proliferative cells in the

blastema tripled at 48 h, which led to an approximate 4-fold increase at 72 hpa (Figure

4D; 24 hpa: n=275.1±19.4, 48 hpa: n=929.3±45.8, 72 hpa: n=1125±100.7). These

results indicate a pronounced difference in proliferative activity between the two

populations, further implicating their distinct responses to injury.

If repair-specific proliferative cells do not derive from the resident stem-like cells

in the bulb, blastema should form when such stem-like cells are removed. To test this

possibility, we examined blastema formation after homeostatic proliferative cells are

eliminated by X-ray irradiation. We first confirmed the elimination of stem-like cells after

X-ray irradiation (30 Gy) by monitoring expression of the stem cell marker *Nanos1*.

qPCR analysis after X-ray irradiation revealed a gradual reduction of *Nanos1* mRNA

levels, particularly after 3 days post-irradiation (dpi) (Figure S5A). FISH staining of

Nanos1 also showed that its signal in the bulb decreased significantly at 3 dpi (Figures

S5B and S5C). We thus amputated the tentacle at 3 dpi and investigated the blastema

formation by EdU pulse labeling (Figure 4F). At 24 hpa, blastema successfully formed

regardless of X-ray irradiation (Figure 4G). Of note, the rate of EdU$^+$ cells in the

blastema was equivalent between tentacles with X-ray irradiation and controls (Figure

4H). Combined, these results demonstrate that distinct proliferative cell populations

appear during tentacle regeneration and that blastema cells are not derived from

resident stem-like cells.

Homeostatic proliferative cells contribute to nematogenesis and tentacle

elongation

In the tentacles of hydrozoan medusae such as *Clytia* and *Cladonema*, proliferative i-

cells in the bulb are proposed to migrate from the basal to the distal side while

differentiating into nematocytes, which is known as the belt conveyor model

(Condamine et al., 2019; Denker et al., 2008; Fujita et al., 2019). Based on this

proposed model, we hypothesized that homeostatic proliferative cells in the bulb

contribute to nematogenesis during both homeostasis and regeneration. In order to test

this hypothesis, we performed EdU and BrdU double chase experiments with some

modifications. We found that long-incubation with EdU for 24 h allows for the labeling of

most proliferative cells in the bulb as well as nematocyte progenitors labeled by *Mcol*

and *Nowa* (Figures 5B i and S6A-S6C). After EdU labeling for 24 h, followed by

amputation, most EdU$^+$ cells remained in the bulb and did not merge with repair-specific

proliferative cells labeled with BrdU in the blastema (Figure 5B ii), which is consistent

with the results of the chasing experiment using EdU pulse labeling (Figure 4B). At 72

hpa, while EdU$^+$ cells were still distributed in the basal region, including the bulb, some

EdU$^+$ cells were located in the distal nematocyte clusters (Figure 5B iii: yellow arrow),

suggesting that one of the lineages of homeostatic proliferative cells are nematocytes

during regeneration. To further confirm the above observation, we identified cell types

located in the distal side of the tentacle. We found that atypical protein kinase C (aPKC)

antibody can be utilized to detect cell types with nuclear staining, which is a method

similar to using β-catenin antibody (Figures 3A, 5C ii, and S7) (Hou et al., 2021; Plickert

et al., 2012). The combination of EdU chasing and cell type identification revealed that

nearly half of the EdU$^+$ cells are nematocytes in the distal side of the tentacle

(nematocytes: 45.7%, epithelial cells: 24.2%, i-cells: 30.1%; Figures 5C and 5D). These

results together indicate that homeostatic proliferative cells in the bulb differentiate into

nematocytes during regeneration.

Given that the homeostatic proliferative cells differentiate into both nematocytes

and epithelial cells during regeneration (Figures 5C and 5D), it is expected that these

cells play a role in supplying new nematocytes as well as in tentacle elongation. We first

examined the number of nematocytes when the homeostatic proliferative cells were

eliminated by X-ray (Figure 5E). At 72 hpa, the number of nematocyte clusters and

particularly stenoteles, large rugby ball-shaped nematocytes, were significantly reduced

in the X-ray irradiated medusae tentacles (nematocyte clusters, control: 11.7, X-ray:

6.52; stenoteles, control: 47.8, X-ray: 26.8) (Figures 5F-5H), indicating that homeostatic

proliferative cells migrate from the basal to the distal side while differentiating into

nematocytes during regeneration.

Next, we measured tentacle length during regeneration with homeostatic

proliferative cells removed by X-ray irradiation (Figure 5I). When tentacles were

amputated at 3 dpi and 5 dpi, by which time homeostatic proliferative cells had been

eliminated, tentacle elongation was significantly inhibited compared to the non-irradiated

controls (Figure 5J, regenerated tentacle length at 96 hpa, Control: 1.70mm, 1 dpi:

1.44mm, 3 dpi: 0.928mm, 5 dpi: 1.04mm). Although we cannot exclude the possibility

that X-ray irradiation affects the source of repair-specific proliferative cells to some

extent, we confirmed that EdU$^+$ blastema cells appear after X-ray irradiation (Figures 4G

and 4H), suggesting that homeostatic proliferative cells contribute to tentacle elongation

independently of blastema cells. We speculate that homeostatic proliferative cells, and

particularly their decedents, participate in tissue elongation likely through pushing out

from the basal side as progenitors and differentiating into epithelial cells.

Repair-specific proliferative cells contribute to reconstruction of the new tentacle

In order to investigate the role of repair-specific proliferative cells during regeneration,

we performed EdU pulse labeling of the blastema with cell type identification by aPKC

antibody staining (Figure 5K). While 94.4% of the injury-responsive proliferative cells

exhibited i-cell morphology at 24 hpa (Figures 5L i and 5M, consistent with the results of

β-catenin antibody staining in Figures 3A and 3B), 59.2% of EdU$^+$ cells were epithelial

cells in the distal portion of the regenerating tentacle at 72 hpa, (Figures 5L ii and 5M).

Strikingly, in contrast to homeostatic proliferative cells, EdU$^+$ cells derived from the

blastema did not become nematocytes (Figures 5A-5D and 5K-5M). These results

suggest that repair-specific proliferative cells preferentially differentiate into epithelial

cells in the regenerating tentacle.

To identify the role of repair-specific proliferative cells, we examined the impact

of locally inhibiting cell proliferation on tentacle regeneration (Figure 5N). UVA radiation

(320 to 400 nm) induces various forms of cellular damage via the production of reactive

oxygen species or the impairment of proteasomal function, which blocks cell cycle

progression at the G_2/M phase in both mammalian cultured cells and fly epidermal

progenitors (Catalgol et al., 2009; Girard et al., 2008; Latonen and Laiho, 2005;

Nakajima et al., 2011). In addition, Hoechst and BrdU are known photosensitizers

(Herweijer et al., 1988; Singh et al., 2004), which can facilitate the process of damage

induction. To verify the effects of UVA on the repair-specific proliferative cells, we

performed UV laser illumination around the blastema region at 24 hpa and counted the

number of EdU labeled cells (Figure S8). Following 24 h, the number of EdU$^+$ cells

decreased significantly in the UV exposed tentacle compared to the control, indicating

that UV laser exposure inhibits cell proliferation (Figure S8). In order to examine the

contribution of the proliferation of repair-specific proliferative cells toward tentacle

regeneration, we monitored the length of regenerating tentacles after UV laser exposure

(Figure 5N). Upon UV laser illumination, tentacle elongation during regeneration was

significantly inhibited (Figures 5N and 5O). These results, together with the distribution

of their descendants on the distal side of the regenerating tentacle (Figures 4B and 5B),

indicate that proliferation of repair-specific proliferative cells contribute to the

reconstruction of new tentacles after amputation.

616   Discussion

In this study, we uncovered the cellular mechanism underlying blastema formation

during *Cladonema* tentacle regeneration. We found that, upon amputation of the

tentacle, repair-specific proliferative cells accumulate as blastema, which is not derived

from resident i-cells. While resident i-cells contribute to nematogenesis and tissue

elongation during both homeostasis and regeneration, repair-specific proliferative cells

differentiate into the epithelial cells of the newly formed tentacle. Based on these

results, we propose that distinct proliferative cell populations facilitate rapid functional

regeneration of the tentacle, which enables critical behaviors of jellyfish organismal

physiology such as feeding and swimming (Figure 6A).

627   Heterogeneity of i-cells

In hydrozoans, i-cells behave as pluripotent stem cells that can differentiate into several

progenitors and differentiated lineages such as nematocytes, neurons, gland cells, and

gametes. Although i-cells exhibit common features such as their interstitial localization,

cellular morphology, and expression of the conserved stem cell markers (*Nanos, Piwi,*

Vasa, PL10); their potency and heterogeneity likely differ across species (Gold and

Jacobs, 2013). Indeed, *Hydra* i-cells differentiate into most cell types except epithelial

634 cells, while *Hydractinia* i-cells behave as pluripotent cells that can differentiate into all

635 cell types including somatic and germline cells (Watanabe et al., 2009; Gahan et al.,

636 2016; Varley et al., 2022).

637   In the intact *Cladonema* jellyfish, *Nanos1*$^+$ and *Piwi*$^+$ cells are primarily distributed

638 at the bulb (Figures 3C and S3), and about 70% of homeostatic proliferative cells in the

639 bulb are *Nanos1*$^+$ i-cells (Figures 3E and 3F). From lineage-tracing experiments, these

640 homeostatic proliferative cells differentiate into epithelia as well as nematocytes

641 (Figures 5C and 5D), suggesting their role as multipotent stem cells. These

642 observations are consistent with the jellyfish *Clytia*, in which i-cells in the bulb proliferate

643 and differentiate into nematocytes (Condamine et al., 2019; Denker et al., 2008),

644 indicating that i-cells in the tentacle bulb are commonly involved in organ homeostasis

645 including nematogenesis in hydrozoan jellyfish.

646   In contrast to i-cells in the bulb, only about 30% of repair-specific proliferative

647 cells at blastema express *Nanos1*$^+$ and *Piwi*$^+$, although more than 80% of blastema cells

648 show i-cell morphology (Figures 3A, B and 5L, 5M). These results imply that the i-cell

649 population of the blastema differs from that of resident i-cells in the bulb. Indeed, repair-

650 specific proliferative cells predominantly differentiate into epithelial cells rather than

651 nematocytes (Figures 5K-5M), suggesting their role as unipotent stem/progenitor cells.

652 Such repair-specific proliferative cells also accumulate at the newly-formed branch

653 starting at 48 hpa (Figure 4B), consistent with the prediction by the previous report

654 proposing that i-cell population in the branching bud differs from resident i-cells in the

655 bulb (Hou et al., 2021). Altogether, our findings support the heterogeneity of i-cells in

656 *Cladonema* jellyfish (Figure 6A), which is similar to the subpopulations of *Hydra* i-cells

657 (David, 2012; Govindasamy et al., 2014).

659 **How are repair-specific proliferative cells supplied in the blastema?**

660 Our work is the first to demonstrate that repair-specific proliferative cells appear upon

661 injury in cnidarians, but how this cell population is supplied remains unknown. Lineage-

662 tracing experiments show that blastema formation is independent of resident i-cells in

663 the bulb (Figures 4A-4E). When the distal side of the tentacle, where few proliferating

664 cells exist, is amputated, local cell proliferation is still observed around the injury site

(Figure 2G). These observations raise two possibilities: mitotic re-entry of

quiescent/slow-cycling i-cells or dedifferentiation from differentiated cells into i-cells.

Previous reports have showed that some cnidarian species possess quiescent or

slow-cycling stem cells. For instance, while *Hydra* has three types of adult stem cells—i-

cells, ectodermal stem cells, and endoderm epithelial stem cells—the subpopulation of

each stem cell is slow-cycling (Govindasamy et al., 2014). These slow-cycling cells are

resistant to hydroxyurea treatment due to G2 phase arrest, and re-enter the cell cycle

during head regeneration. *Nematostella* also has slow-cycling/quiescent cells that are

predominantly present in the mesenteries and are resistant to X-ray irradiation, and

these cells' cell cycle is activated upon injury, when they migrate toward the wound site

(Amiel et al., 2019). Given that a small number of i-cells exist in the *Cladonema* intact

tentacle, outside the bulb, and branching bud (Figure S3A) (Hou et al., 2021); such a

minor population of i-cells may behave as slow-cycling/quiescent cells and re-enter the

cell cycle in response to injury.

By contrast, although cnidarian dedifferentiation potential is still debated, some

species appear to exhibit context-dependent cellular plasticity (Nakanishi and Jacobs,

2019). For instance, classical experiments using the hydrozoan jellyfish *Podocoryna*

carnea have indicated their dedifferentiation potential *in vitro*: when isolated striated

muscles are cultured after collagenase treatment, the muscle cells dedifferentiate into

stem-like cells and re-differentiate into multiple cell types (Schmid and Alder, 1984).

Additionally, in green *Hydra*, the isolated gastrodermis, which is i-cell free, can

reconstitute a complete polyp including the ectoderm (Haynes and Burnett, 1963). In

this case, endodermal gland cells appear to dedifferentiate into i-cells (Burnett et al.,

1966), while endodermal digestive cells transdifferentiate into epidermal

epitheliomuscular cells (Davis et al., 1966). Moreover, a recent report using *Hydractinia*

has shown that, when the hypostome is isolated from the body, cellular reprogramming

triggered by cell senescence occurs in somatic cells, which convert into pluripotent

Piwi1[+] i-cells (Salinas-Saavedra et al., 2022). Although these reports support the

possibility that dedifferentiation produces pluri/multipotent stem cells after isolation from

the body, it is also possible that repair-specific proliferative cells are derived from their

descendant cell types. During newt limb regeneration, cellular lineage is restricted such

that muscle progenitors dedifferentiated from resident muscle fibers produce only

muscles (Sandoval-Guzmán et al., 2014; Tanaka et al., 2016). Given that repair-specific

proliferative cells in the *Cladonema* tentacle differentiate into only epithelial cells

(Figures 5K-5M), epithelial cells may dedifferentiate into unipotent progenitors that give

rise to only epithelial cells in the newly formed tentacle.

Evolutionary conservation and diversification of blastema formation

Although the molecular and cellular mechanisms of animal regeneration have been

addressed in detail using representative species such as planaria and salamanders,

questions such as the evolutionary history of regeneration and the homology of

regeneration mechanisms across species remain largely unknown. Repair-specific stem

cells provided by quiescent stem cells and/or dedifferentiation are common blastema

components in bilaterian appendage regeneration. As in vertebrates where lineage-

restricted repair-specific stem/progenitor cells participate in appendage regeneration

(Ando et al., 2017; Knopf et al., 2011; Lehoczky et al., 2011; Sandoval-Guzmán et al.,

2014), the protostome ecdysozoa *Parhyale hawaiensis* utilize re-activation of muscle

satellite cells during thoracic leg regeneration (Konstantinides and Averof, 2014). By

contrast, during echinoderm starfish arm regeneration, dedifferentiated dermal cells

form blastema-like structures (Fan et al., 2011). Similarly, during insect cricket leg

regeneration, the source of blastemal cells is derived from injury-induced

dedifferentiation (Bando et al., 2018; Truby, 1983). These reports suggest that repair-

specific production of blastema cells is relatively conserved throughout bilaterians

(Figure 6B).

In non-bilaterian metazoans, resident pluripotent stem cells are thought to

contribute to both homeostasis and whole-body regeneration. During the regeneration

of the demosponge in Porifera, one of the most primitive animals, archeocytes known

as homeostatic stem cells contribute to an undifferentiated cell mass beneath the

wound as a blastema-like structure (Borisenko et al., 2015). While the regenerating

ctenophore cydippids do not apparently form blastema-like structure, slow-cycling stem-

like cells are recruited to the injury site (Ramon-Mateu et al., 2019). In cnidarians,

resident stem cells, which are involved in homeostasis, are recruited to the injury site

after amputation of the body and removal of the organs (Amiel et al., 2019; Bradshaw et

al., 2015; Chera et al., 2009; Sinigaglia et al., 2020). By contrast, as shown in this study,

the source of blastemal cells are not resident i-cells but rather repair-specific

proliferative cells during *Cladonema* tentacle regeneration (Figure 4). Our findings

suggest that, in terms of the regeneration processes and blastemal cellular source, the

regeneration mechanism of the *Cladonema* tentacle is more homologous to that of

bilaterian species (Figure 6B).

One important remaining question is whether bilaterians' and cnidarians' most

recent common ancestor possessed a system to supply blastema through repair-

specific proliferative cells. Although sponges and ctenophores have been proposed to

possess dedifferentiation or transdifferentiation potentials that can provide a cellular

source of regenerated bodies (Nakanishi and Jacobs, 2019), it is currently unknown

whether they are repair-specific cell populations or not. In this study, we demonstrate

that repair-specific proliferative cells contribute to the newly generated epithelium during

Cladonema tentacle regeneration. Intriguingly, recent genomic studies have revealed

that jellyfish-specific genes are not conserved among bilaterians or even other

cnidarians including corals and sea anemones, and the anthozoan *Nematostella* is the

most 'bilaterian-like' cnidarian sequenced (Khalturin et al., 2019; Putnam et al., 2007;

Steele et al., 2011). These findings raise the possibility that medusozoans had evolved

the medusa stage after branching from their common ancestor and that bilaterians and

cnidarian jellyfish likely independently acquired a similar mechanism of blastema

formation upon amputation of appendages. It is thus tempting to speculate that

blastema formation by repair-specific proliferative cells is a common feature acquired for

complex organ and appendage regeneration during animal evolution.

References

Amiel AR, Foucher K, Ferreira S, Röttinger E. 2019. Synergic coordination of stem cells is

required to induce a regenerative response in anthozoan cnidarians.

doi:10.1101/2019.12.31.891804

Ando K, Shibata E, Hans S, Brand M, Kawakami A. 2017. Osteoblast Production by Reserved

Progenitor Cells in Zebrafish Bone Regeneration and Maintenance. *Developmental Cell* **43**:643-

650.e3. doi:10.1016/j.devcel.2017.10.015

Aztekin C. 2021. Appendage regeneration is context dependent at the cellular level. *Open Biol*
11:210126. doi:10.1098/rsob.210126

Bando T, Mito T, Hamada Y, Ishimaru Y, Noji S, Ohuchi H. 2018. Molecular mechanisms of
limb regeneration: insights from regenerating legs of the cricket Gryllus bimaculatus. *Int J Dev*
Biol **62**:559–569. doi:10.1387/ijdb.180048ho

Bely AE, Nyberg KG. 2010. Evolution of animal regeneration: re-emergence of a field. *Trends*
in Ecology & Evolution **25**:161–170. doi:10.1016/j.tree.2009.08.005

Borisenko IE, Adamska M, Tokina DB, Ereskovsky AV. 2015. Transdifferentiation is a driving
force of regeneration in *Halisarca dujardini* (Demospongiae, Porifera). *PeerJ* **3**:e1211.
doi:10.7717/peerj.1211

Bradshaw B, Thompson K, Frank U. 2015. Distinct mechanisms underlie oral vs aboral
regeneration in the cnidarian Hydractinia echinata. *eLife* **4**:e05506. doi:10.7554/eLife.05506

Burnett AL, Davis LE, Ruffing FE. 1966. A histological and ultrastructural study of germinal
differentiation of interstitial cells arising from gland cells inHydra viridis. *J Morphol* **120**:1–8.
doi:10.1002/jmor.1051200102

Catalgol B, Ziaja I, Breusing N, Jung T, Höhn A, Alpertunga B, Schroeder P, Chondrogianni N,
Gonos ES, Petropoulos I, Friguet B, Klotz L-O, Krutmann J, Grune T. 2009. The Proteasome Is
an Integral Part of Solar Ultraviolet A Radiation-induced Gene Expression. *Journal of Biological*
Chemistry **284**:30076–30086. doi:10.1074/jbc.M109.044503

Chera S, Ghila L, Dobretz K, Wenger Y, Bauer C, Buzgariu W, Martinou J-C, Galliot B. 2009.
Apoptotic Cells Provide an Unexpected Source of Wnt3 Signaling to Drive Hydra Head
Regeneration. *Developmental Cell* **17**:279–289. doi:10.1016/j.devcel.2009.07.014

Condamine T, Jager M, Leclère L, Blugeon C, Lemoine S, Copley RR, Manuel M. 2019.

Molecular characterisation of a cellular conveyor belt in Clytia medusae. *Developmental Biology*

456:212–225. doi:10.1016/j.ydbio.2019.09.001

Daponte V, Tylzanowski P, Forlino A. 2021. Appendage Regeneration in Vertebrates: What

Makes This Possible? *Cells* **10**:242. doi:10.3390/cells10020242

David CN. 2012. Interstitial stem cells in Hydra: multipotency and decision-making. *Int J Dev*

Biol **56**:489–497. doi:10.1387/ijdb.113476cd

Davis LE, Burnett AL, Haynes JF, Mumaw VR. 1966. A histological and ultrastructural study of

dedifferentiation and redifferentiation of digestive and gland cells in Hydra viridis.

Developmental Biology **14**:307–329. doi:10.1016/0012-1606(66)90018-2

Denker E, Manuel M, Leclère L, Le Guyader H, Rabet N. 2008. Ordered progression of

nematogenesis from stem cells through differentiation stages in the tentacle bulb of Clytia

hemisphaerica (Hydrozoa, Cnidaria). *Developmental Biology* **315**:99–113.

doi:10.1016/j.ydbio.2007.12.023

Fan T, Fan X, Du Y, Sun W, Zhang S, Li J. 2011. Patterns and cellular mechanisms of arm

regeneration in adult starfish Asterias rollestoni bell. *J Ocean Univ China* **10**:255–262.

doi:10.1007/s11802-011-1837-y

Fritz AE, Ikmi A, Seidel C, Paulson A, Gibson MC. 2013. Mechanisms of tentacle

morphogenesis in the sea anemone *Nematostella vectensis*. *Development* **140**:2212–2223.

doi:10.1242/dev.088260

Fujiki A. 2019. Branching pattern and morphogenesis of medusa tentacles in the jellyfish

Cladonema pacificum (Hydrozoa, Cnidaria). *Zoological Letters* **5**:13. doi:10.1186/s40851-019-

0124-4

Fujita S, Kuranaga E, Miura M, Nakajima Y-I. 2022. Fluorescent In Situ Hybridization and 5-
Ethynyl-2'-Deoxyuridine Labeling for Stem-like Cells in the Hydrozoan Jellyfish Cladonema
pacificum. *J Vis Exp*. doi:10.3791/64285

Fujita S, Kuranaga E, Nakajima Y. 2021. Regeneration Potential of Jellyfish: Cellular
Mechanisms and Molecular Insights. *Genes* **12**:758. doi:10.3390/genes12050758

Fujita S, Kuranaga E, Nakajima Y. 2019. Cell proliferation controls body size growth, tentacle
morphogenesis, and regeneration in hydrozoan jellyfish *Cladonema pacificum*. *PeerJ* **7**:e7579.
doi:10.7717/peerj.7579

Gahan JM, Bradshaw B, Flici H, Frank U. 2016. The interstitial stem cells in Hydractinia and
their role in regeneration. *Current Opinion in Genetics & Development* **40**:65–73.
doi:10.1016/j.gde.2016.06.006

Galliot B, Buzgariu W, Schenkelaars Q, Wenger Y. 2018. Non-developmental dimensions of
adult regeneration in Hydra. *Int J Dev Biol* **62**:373–381. doi:10.1387/ijdb.180111bg

Gamero-Mora E, Halbauer R, Bartsch V, Stampar SN, Morandini AC. 2019. Regenerative
Capacity of the Upside-down Jellyfish Cassiopea xamachana. *Zool Stud* **58**.
doi:10.6620/ZS.2019.58-37

Girard P-M, Pozzebon M, Delacôte F, Douki T, Smirnova V, Sage E. 2008. Inhibition of S-phase
progression triggered by UVA-induced ROS does not require a functional DNA damage
checkpoint response in mammalian cells. *DNA Repair* **7**:1500–1516.
doi:10.1016/j.dnarep.2008.05.004

Gold DA, Jacobs DK. 2013. Stem cell dynamics in Cnidaria: are there unifying principles? *Dev
Genes Evol* **223**:53–66. doi:10.1007/s00427-012-0429-1

Govindasamy N, Murthy S, Ghanekar Y. 2014. Slow-cycling stem cells in hydra contribute to
head regeneration. *Biology Open* **3**:1236–1244. doi:10.1242/bio.201410512

Harris L, Zalucki O, Piper M. 2018. BrdU/EdU dual labeling to determine the cell-cycle

dynamics of defined cellular subpopulations. *J Mol Hist* **49**:229–234. doi:10.1007/s10735-018-

9761-8

Haynes J, Burnett AL. 1963. Dedifferentiation and Redifferentiation of Cells in Hydra viridis.

Science **142**:1481–1483. doi:10.1126/science.142.3598.1481

Herweijer H, And WS, Visser JWM. 1988. High-speed photodamage cell selection using

bromodeoxyuridine/Hoechst 33342 photosensitized cell killing. *Cytometry* **9**:143–149.

doi:10.1002/cyto.990090208

Holstein TW, Hobmayer E, Technau U. 2003. Cnidarians: An evolutionarily conserved model

system for regeneration? *Dev Dyn* **226**:257–267. doi:10.1002/dvdy.10227

Hou S, Zhu J, Shibata S, Nakamoto A, Kumano G. 2021. Repetitive accumulation of interstitial

cells generates the branched structure of *Cladonema* medusa tentacles. *Development*

148:dev199544. doi:10.1242/dev.199544

Katsuki T, Greenspan RJ. 2013. Jellyfish nervous systems. *Current Biology* **23**:R592–R594.

doi:10.1016/j.cub.2013.03.057

Khalturin K, Shinzato C, Khalturina M, Hamada M, Fujie M, Koyanagi R, Kanda M, Goto H,

Anton-Erxleben F, Toyokawa M, Toshino S, Satoh N. 2019. Medusozoan genomes inform the

evolution of the jellyfish body plan. *Nat Ecol Evol* **3**:811–822. doi:10.1038/s41559-019-0853-y

Knopf F, Hammond C, Chekuru A, Kurth T, Hans S, Weber CW, Mahatma G, Fisher S, Brand

M, Schulte-Merker S, Weidinger G. 2011. Bone Regenerates via Dedifferentiation of Osteoblasts

in the Zebrafish Fin. *Developmental Cell* **20**:713–724. doi:10.1016/j.devcel.2011.04.014

Konstantinides N, Averof M. 2014. A Common Cellular Basis for Muscle Regeneration in

Arthropods and Vertebrates. *Science* **343**:788–791. doi:10.1126/science.1243529

Latonen L, Laiho M. 2005. Cellular UV damage responses—Functions of tumor suppressor p53.

Biochimica et Biophysica Acta (BBA) - Reviews on Cancer **1755**:71–89.

doi:10.1016/j.bbcan.2005.04.003

Leclère L, Röttinger E. 2017. Diversity of Cnidarian Muscles: Function, Anatomy, Development

and Regeneration. *Front Cell Dev Biol* **4**. doi:10.3389/fcell.2016.00157

Lehoczky JA, Robert B, Tabin CJ. 2011. Mouse digit tip regeneration is mediated by fate-

restricted progenitor cells. *Proc Natl Acad Sci USA* **108**:20609–20614.

doi:10.1073/pnas.1118017108

Lesh-Laurie GE, Hujer A, Suchy P. 1991. Polyp regeneration from isolated tentacles of Aurelia

scyphistomae: a role for gating mechanisms and cell division. *Hydrobiologia* **216**:91–97.

doi:10.1007/BF00026448

Masuda-Ozawa T, Fujita S, Nakamura R, Watanabe H, Kuranaga E, Nakajima Y. 2022. siRNA-

mediated gene knockdown via electroporation in hydrozoan jellyfish embryos. *Sci Rep*

12:16049. doi:10.1038/s41598-022-20476-1

McCusker C, Bryant SV, Gardiner DM. 2015. The axolotl limb blastema: cellular and molecular

mechanisms driving blastema formation and limb regeneration in tetrapods: The Axolotl Limb

Blastema. *Regeneration* **2**:54–71. doi:10.1002/reg2.32

Nakajima Y, Kuranaga E, Sugimura K, Miyawaki A, Miura M. 2011. Nonautonomous Apoptosis

Is Triggered by Local Cell Cycle Progression during Epithelial Replacement in Drosophila. *Mol*

Cell Biol **31**:2499–2512. doi:10.1128/MCB.01046-10

Nakanishi N, Jacobs DK. 2019. The Early Evolution of Cellular Reprogramming in Animals In:

Bishop CD, Hall BK, editors. Deferring Development. CRC Press. pp. 67–86.

doi:10.1201/9780429445446-4

Ostendarp M, Plewka J, Flathmann J, Tilstra A, El-Khaled YC, Wild C. 2022. Complete and

rapid regeneration of fragments from the upside-down jellyfish Cassiopea. *Front Mar Sci*

9:949233. doi:10.3389/fmars.2022.949233

Passamaneck YJ, Martindale MQ. 2012. Cell proliferation is necessary for the regeneration of

oral structures in the anthozoan cnidarian Nematostella vectensis. *BMC Dev Biol* **12**:34.

doi:10.1186/1471-213X-12-34

Plickert G, Frank U, Müller WA. 2012. Hydractinia, a pioneering model for stem cell biology

and reprogramming somatic cells to pluripotency. *Int J Dev Biol* **56**:519–534.

doi:10.1387/ijdb.123502gp

Putnam NH, Srivastava M, Hellsten U, Dirks B, Chapman J, Salamov A, Terry A, Shapiro H,

Lindquist E, Kapitonov VV, Jurka J, Genikhovich G, Grigoriev IV, Lucas SM, Steele RE,

Finnerty JR, Technau U, Martindale MQ, Rokhsar DS. 2007. Sea Anemone Genome Reveals

Ancestral Eumetazoan Gene Repertoire and Genomic Organization. *Science* **317**:86–94.

doi:10.1126/science.1139158

Ramon-Mateu J, Ellison ST, Angelini TE, Martindale MQ. 2019. Regeneration in the ctenophore

Mnemiopsis leidyi occurs in the absence of a blastema, requires cell division, and is temporally

separable from wound healing. *BMC Biol* **17**:80. doi:10.1186/s12915-019-0695-8

Reddien PW, Alvarado AS. 2004. FUNDAMENTALS OF PLANARIAN REGENERATION.

Annu Rev Cell Dev Biol **20**:725–757. doi:10.1146/annurev.cellbio.20.010403.095114

Rees JT. 1979. The polyp and immature medusa stages of *Cladonema californicum* , Hyman,

1947 (Anthomedusae: Cladonemidae) with biological notes and a discussion of the taxonomy of

the genus *Cladonema*. *Journal of Natural History* **13**:295–302.

doi:10.1080/00222937900770231

Ribeiro RP, Egger B, Ponz-Segrelles G, Aguado MT. 2021. Cellular proliferation dynamics

during regeneration in Syllis malaquini (Syllidae, Annelida). *Front Zool* **18**:27.

doi:10.1186/s12983-021-00396-y

Röttinger E. 2021. Nematostella vectensis, an Emerging Model for Deciphering the Molecular

and Cellular Mechanisms Underlying Whole-Body Regeneration. *Cells* **10**:2692.

doi:10.3390/cells10102692

Salic A, Mitchison TJ. 2008. A chemical method for fast and sensitive detection of DNA

synthesis *in vivo*. *Proc Natl Acad Sci USA* **105**:2415–2420. doi:10.1073/pnas.0712168105

Salinas-Saavedra M, Febrimarsa, Krasovec G, Horkan HR, Baxevanis AD, Frank U. 2022.

Senescence-induced cellular reprogramming drives cnidarian whole-body regeneration.

doi:10.1101/2022.09.02.506310

Sandoval-Guzmán T, Wang H, Khattak S, Schuez M, Roensch K, Nacu E, Tazaki A, Joven A,

Tanaka EM, Simon A. 2014. Fundamental Differences in Dedifferentiation and Stem Cell

Recruitment during Skeletal Muscle Regeneration in Two Salamander Species. *Cell Stem Cell*

14:174–187. doi:10.1016/j.stem.2013.11.007

Schmid V, Alder H. 1984. Isolated, mononucleated, striated muscle can undergo pluripotent

transdifferentiation and form a complex regenerate. *Cell* **38**:801–809. doi:10.1016/0092-

8674(84)90275-7

Seifert AW, Muneoka K. 2018. The blastema and epimorphic regeneration in mammals.

Developmental Biology **433**:190–199. doi:10.1016/j.ydbio.2017.08.007

Singh S, Dwarakanath B, Mathew T. 2004. DNA ligand Hoechst-33342 enhances UV induced

cytotoxicity in human glioma cell lines. *Journal of Photochemistry and Photobiology B: Biology*

77:45–54. doi:10.1016/S1011-1344(04)00122-8

Sinigaglia C, Peron S, Eichelbrenner J, Chevalier S, Steger J, Barreau C, Houliston E, Leclère L.
2020. Pattern regulation in a regenerating jellyfish. *eLife* **9**:e54868. doi:10.7554/eLife.54868

Stamatis S-A, Worsaae K, Garm A. 2018. Regeneration of the Rhopalium and the Rhopalial
Nervous System in the Box Jellyfish *Tripedalia cystophora*. *The Biological Bulletin* **234**:22–36.
doi:10.1086/697071

Steele RE, David CN, Technau U. 2011. A genomic view of 500 million years of cnidarian
evolution. *Trends in Genetics* **27**:7–13. doi:10.1016/j.tig.2010.10.002

Stierwald M, Yanze N, Bamert RP, Kammermeier L, Schmid V. 2004. The Sine oculis/Six class
family of homeobox genes in jellyfish with and without eyes: development and eye regeneration.
Developmental Biology **274**:70–81. doi:10.1016/j.ydbio.2004.06.018

Suga H, Schmid V, Gehring WJ. 2008. Evolution and Functional Diversity of Jellyfish Opsins.
Current Biology **18**:51–55. doi:10.1016/j.cub.2007.11.059

Takeda N, Kon Y, Artigas GQ, Lapébie P, Barreau C, Koizumi O, Kishimoto T, Tachibana K,
Houliston E, Deguchi R. 2018. Identification of jellyfish neuropeptides that act directly as oocyte
maturation-inducing hormones. *Development* **145**. doi:10.1242/dev.156786

Tanaka EM, Reddien PW. 2011. The Cellular Basis for Animal Regeneration. *Developmental*
Cell **21**:172–185. doi:10.1016/j.devcel.2011.06.016

Tanaka HV, Ng NCY, Yang Yu Z, Casco-Robles MM, Maruo F, Tsonis PA, Chiba C. 2016. A
developmentally regulated switch from stem cells to dedifferentiation for limb muscle
regeneration in newts. *Nat Commun* **7**:11069. doi:10.1038/ncomms11069

Tasaki J, Shibata N, Sakurai T, Agata K, Umesono Y. 2011. Role of c-Jun N-terminal kinase
activation in blastema formation during planarian regeneration: JNK activation in planarian
regeneration. *Development, Growth & Differentiation* **53**:389–400. doi:10.1111/j.1440-
169X.2011.01254.x

Technau U, Steele RE. 2011. Evolutionary crossroads in developmental biology: Cnidaria.

Development **138**:1447–1458. doi:10.1242/dev.048959

Truby PR. 1983. Blastema formation and cell division during cockroach limb regeneration.

Development **75**:14.

Varley Á, Horkan HR, McMahon ET, Krasovec G, Frank U. 2022. Pluripotent, germ cell

competent adult stem cells underlie cnidarian plant-like life history (preprint). Developmental

Biology. doi:10.1101/2022.11.09.515637

Vega CJ, Peterson DA. 2005. Stem cell proliferative history in tissue revealed by temporal

halogenated thymidine analog discrimination. *Nat Methods* **2**:167–169. doi:10.1038/nmeth741

Vogg MC, Galliot B, Tsiairis CD. 2019. Model systems for regeneration: Hydra. *Development*

146. doi:10.1242/dev.177212

Weber C. 1981. Structure, histochemistry, ontogenetic development, and regeneration of the

ocellus ofCladonema radiatum dujardin (cnidaria, hydrozoa, anthomedusae). *J Morphol*

167:313–331. doi:10.1002/jmor.1051670306

Zhou K, Gu Y, Wang L, Chen J. 2022. Discovery of Cladonema multiramosum sp. nov.

(Cnidaria: Hydrozoa: Cladonematidae) using DNA barcoding and life cycle analyses. *Acta*

Oceanol Sin **41**:44–52. doi:10.1007/s13131-021-1900-5

Acknowledgement

We thank R. Deguchi (Miyagi Univ. Education, Japan) for sharing *Cladonema*

pacificum. We thank D. Umetsu and T. Akiyama for helpful discussion. We thank I.

Nagai, A. Sasaki, and H. Nakatani for technical assistance. This work was supported by

JSPS/MEXT KAKENHI (grant numbers JP22J01430 to H.N., JP21H05255 to E.K.,

JP21H04774, JP21K19206 to M.M., and JP17H06332, JP19K22550, JP22H02762 to

Y.N.), JST grant (JPMJCR1852 to E.K. and JPMJSP2114 to S.F.), AMED-Aging

(JP21gm5010001 to M.M.), and AMED-PRIME (JP22gm6110025 to Y.N.).

Author contributions

974   S.F. conceptualized and designed the project, performed experiments, analyzed data,

975   prepared figures, and wrote the manuscript. E.K. supervised the project. M.M.

976   supervised the project. Y.N. conceptualized, supervised and designed the project,

977   prepared figures, and wrote the manuscript. All authors approved the final manuscript.

Competing interests

981   The authors declare no competing interests.

Figure legends

Figure 1. Regeneration processes and potentials of the *Cladonema* medusa

tentacle. (A) The phylogenetic tree of Bilateria and Cnidaria, composed of Anthozoa

and Medusozoa. (B) Medusa of *Cladonema pacificum*. U: umbrella, M: manubrium, Ra:

radial canal, Ri: ring canal, T: tentacle. (C) Tentacle of *Cladonema* medusa. O: ocellus,

TB: tentacle bulb, Br: branch, NC: nematocyte cluster. (D) Tentacle regeneration

processes after amputation while retaining the bulb. hpa: hour post-amputation. Yellow

arrows indicate nematocyte clusters. n = 234/236 (tentacles). (E) Scheme of the

tentacle regeneration that retains the bulb. (F) Distribution of mature nematocytes and

localization of muscle fibers during tentacle regeneration. DAPI for poly-γ-glutamate

(green) and nuclei (blue), and Phalloidin for F-actin (red). Yellow arrows indicate

nematocyte clusters. (G) Neural morphology in the regenerating tentacle stained with

the anti-FMRFamide antibody. White arrows indicate neural fibers. (H) The regenerating

tentacle is fully functional at 72 hpa. Image of the tentacle capturing prey, brine shrimp

(Arrow). (I) The rate of functional tentacles across the regeneration time course. Intact:

n = 40 (tentacles), 0-72 hpa: n = 36. (J) Tentacle regeneration process after removing

the bulb from canals. Dpa: day post-amputation. n = 117/219 (tentacles). Ra: radial

canal, Ri: ring canal. (K) Scheme of tentacle regeneration without bulb. Scale bars: (A-

C, I) 1 mm, (E, F i) 100 μm, (F ii) 50 μm.

1004

Figure 2. Blastema formation at early phase of tentacle regeneration.

(A) Distribution of S-phase cells in intact and regenerating tentacle stained with EdU 1 h

pulse labeling. (B) Quantification of EdU$^+$ cells' distribution in regenerating tentacle. 3

areas defined as basal (grey arrows), middle (pink arrows), distal (red arrows) in (A).

Intact: n = 19 (tentacles), 24 hpa: n = 10, 48 hpa: n = 9, 72 hpa: n = 8. (C) Scheme of

proliferative cells distribution during tentacle regeneration. (D) Distribution of S-phase

and M-phase cells stained with EdU 1 h pulse labeling and anti-Phospho-Histone 3

(PH3) antibody. EdU (green) and PH3 (magenta). Yellow dot squares show each

quantification area (74.72^2 μm^2) in (E, F). (E, F) The proportion of PH3$^+$ and EdU$^+$ cells

per all cells in each area. Each area: n = 14. (G-I) Distribution of proliferative cells at 24

hpa: (G) After the distal side of tentacle is amputated, (H) after removing the bulb from

canals, and (I) in isolated tentacles at 24 hpa. EdU 1 h pulse labeling. Unpaired two-

tailed *t*-test. *$p < 0.05$, **$p < 0.005$, ***$p < 0.001$. Scale bars: (A, G-I) 100 μm, (D) 50 μm.

1018

Figure 3. Stem cells and progenitor cells in blastema.

(A) Identification of cell types in blastema stained with anti-β-catenin staining. White

arrow: epithelial cell, white arrow head: i-cell, yellow arrow head: nematocyte. Yellow

dot squares show each quantification area (35.10^2 μm^2) in (B). (B) Cell types in

blastema at 24 hpa by anti-β-catenin staining. n = 3 (areas). (C) FISH images of

Nanos1 gene expression from the proximal, abaxial side of the tentacle. Distribution of

stem (*Nanos1$^+$*) cells in intact and regenerating tentacle. Yellow arrow indicates main

tentacle, and white arrow indicates new branch. (D) Quantification of *Nanos1+* cells

distribution in regenerating tentacle. 3 areas defined as basal (grey arrows), middle

(pink arrows), distal (red arrows) in (C). Intact: n = 5, 24 hpa: n = 4 (tentacles), 48 hpa: n

= 4, 72 hpa: n = 5. (E) Images of the intact and regenerating tentacle co-labeled with

Nanos1 FISH and EdU 1 h pulse labeling. EdU (green) and *Nanos1* (magenta). White

arrowheads indicate EdU$^+$ and *Nanos1$^+$* cells. (F) The rate of *Nanos1+* cells / EdU+

cells in intact and regenerating tentacles. Quantification area of intact tentacle bulb and

blastema is 74.72^2 µm^2. Intact: n = 18, 24 hpa: n = 22. Unpaired two-tailed *t*-test. *$p <$

0.05, **$p <$ 0.005, ***$p <$ 0.001. Scale bars: (A i, C and E) 50 µm, (A ii) 10 µm.

1035

Figure 4. Distinct proliferative cell populations during tentacle regeneration.

(A) Scheme of lineage-tracing experiments using EdU and BrdU. (B) Distribution of

EdU$^+$ and BrdU$^+$ cells during tentacle regeneration. EdU (green) and BrdU (red). (C)

The number of homeostatic proliferative cells during regeneration. Intact: n = 12

(tentacles), 24 hpa: n = 10, 48 hpa: n = 12, 72 hpa: n = 9. (D) The number of repair-

specific proliferative cells during regeneration. 24 hpa: n = 12, 48 hpa: n = 12, 72 hpa: n

= 6. (E) Scheme of two proliferative cell populations during regeneration. (F) Scheme of

stem cell elimination by 30 Gy X-ray irradiation. Day post-irradiation (dpi). (G)

Distribution of proliferative cells after X-ray irradiation. EdU 1 h pulse labeling. Yellow

dot squares show each quantification area (74.72^2 µm^2) in (H). (H) The rate of EdU$^+$

cells in blastema. No irradiation: n = 14, X-ray irradiation: n = 7. Unpaired two-tailed *t*-

test. *$p <$ 0.05, **$p <$ 0.005, ***$p <$ 0.001. Scale bars: (B, G) 100 µm.

1048

Figure 5. Distinct function of two proliferative cell populations.

(A) Scheme of EdU long-time and BrdU double chasing experiment. (B) The

distributions of EdU$^+$ and BrdU$^+$ cells during tentacle regeneration. EdU (green) and

BrdU (red). Yellow arrows indicate accumulation of EdU$^+$ cells at nematocyte clusters in

distal side at 72 hpa. (C) Cell type identification of EdU labeled cells in distal side at 72

hpa by anti-aPKC staining. EdU (green) and aPKC (magenta). Yellow arrow heads

show nematocytes. (D) Cell types of EdU$^+$ cells on the distal side of a regenerating

tentacle at 72 hpa. Quantification area of regenerating tentacle is 74.72^2 µm^2. n = 10.

(E) Experimental scheme for nematogenesis after 30 Gy X-ray irradiation. (F)

Distribution of mature nematocytes at 72 hpa after irradiation stained with DAPI. DAPI

for poly-γ-glutamate (green) and and nuclei (blue). Yellow arrows indicate nematocyte

clusters. (G) The number of stenoteles (large mature nematocytes) in the regenerating

tentacle at 72 hpa. Ctl: n = 36 (tentacles), X-ray: n = 42. (H) The number of nematocyte

clusters in the regenerating tentacle at 72 hpa. Ctl: n = 36, X-ray: n = 42. (I)

Experimental scheme for tentacle elongation after 30 Gy X-ray irradiation. (J) Length of

regenerating tentacle after irradiation. Ctl: n = 58 (tentacles), 1 dpi: n = 58, 3 dpi: n = 58,

5 dpi: n = 58. (K) Experimental scheme for combining the identification of cell types and
EdU chasing. (L) Identification of cell types of EdU$^+$ cells in blastema and the distal side
of the regenerating tentacle stained with anti-aPKC staining. EdU (green) and aPKC
(magenta). White arrow: epithelial cell, white arrow head: i-cell, yellow arrow head:
nematocyte. (M) Quantification of cell types in EdU$^+$ cells at 24 and 72 hpa.
Quantification area of regenerating tentacle is 35.06^2 µm^2. 24 hpa: n = 9, 72 hpa: n =
12. (N) Experimental scheme for measuring of regenerating tentacle after damage to
the blastema from UV exposure. (O) Length of the regenerating tentacle after UV
exposure on blastema. Ctl: n = 12 (tentacles), UV: n = 12. Unpaired two-tailed *t*-test. *p
< 0.05, **p < 0.005, ***p < 0.001. Scale bars: (B, F) 100 µm, (C, L) 50 µm.
1075

Figure 6. Summary of *Cladonema* tentacle regeneration and evolutionary history
of appendage regeneration.
(A) When a *Cladonema* medusa tentacle is amputated, wound closure is completed by
24 hpa. After wound healing, repair-specific proliferative cells appear and actively
proliferate at the wound tip. Ectopic cell proliferation on the distal side is gradually
suppressed, and the cell differentiation phase begins. While homeostatic proliferative
cells in the bulb supply nematocytes and epithelial cells, repair-specific proliferative cells
only differentiate into epithelial cells. (B) Blastema formation mechanism during
appendage regeneration across animal evolution. When an appendage is amputated,
repair-specific proliferative cells (stem/progenitor cells) contribute to blastema formation
in both regenerative bilaterians (e.g. salamanders) and cnidarian jellyfish.
1087

Figure S1. Wound healing, neuron regeneration and the rate of regeneration.
(A) The process of wound healing from 4 hpa to 24 hpa with Phalloidin and anti-α-
Tubulin antibody. Phalloidin for F-actin (red) and α-Tubulin (green). (B) The extent of
wound healing during tentacle regeneration. 4 hpa: n = 15 (tentacles), 8 hpa: n = 17, 16
hpa: n = 15, 24 hpa: n = 14. (C) Neural morphology in intact tentacle and the
regenerating tentacle stained with the anti-FMRFamide antibody. Yellow allows indicate
cell bodies of FMRFamide neurons. FMRFamide (magenta). (i, ii) Intact whole tentacle,
(iii, iv) basal side of intact tentacle, (v, vi) regenerating tentacle at 24 hpa. (D) Neural
morphology in the regenerating tentacle stained with the anti-acetylated-Tubulin

antibody. White arrows indicate neural fibers. acetylated-Tubulin (green). (E) The rate of
the tentacle regeneration after removing the bulb from canals. (F) Images of the isolated
tentacle without bulb at 1 dpa and 7 dpa. (G) Images of the isolated whole tentacle 1
dpa and 7 dpa. Scale bars: (A, C i-iii, v and D i, iii) 100 μm, (C iv, vi and D ii, iv) 50 μm,
(F, G) 1 mm.

1102

1103 **Figure S2. The distribution of mitotic cells during regeneration.**
1104 (A) Distribution of mitotic cells detected by anti-PH3 in regenerating tentacle. (B)
1105 Quantification of PH3$^+$ cells distribution in regenerating tentacle. Intact: n = 11
1106 (tentacles), 24 hpa: n = 10, 48 hpa: n = 9, 72 hpa: n = 8.

1107

Figure S3. The distribution of stem cells in intact tentacle and regenerating
tentacle.
(A) Expression pattern of *Nanos1* in intact tentacle by FISH. (B) Expression pattern of
Piwi in intact tentacle by FISH. (C) Distribution of stem (*Piwi$^+$*) cells in intact and
regenerating tentacle. White arrowhead indicates *Piwi$^+$* cells. (D) Images of the proximal
abaxial side of regenerating tentacle co-labeled with *Piwi* expression and EdU. EdU 1 h
pulse labeling. White arrowhead indicates *Piwi$^+$* and EdU$^+$ cells. (E) The rate of *Piwi$^+$*
cells / EdU$^+$ cells around blastema in regenerating tentacles. n = 5 (tentacles). Scale
bars: (A and B) 100 μm, (C and D) 50 μm.

1117

Figure S4. Distribution and increase of homeostatic proliferative cells during
tentacle regeneration.
(A) Transition of labeled cells during regeneration by chasing EdU signals. EdU 1 h
pulse labeling before amputation. (B) Experiment scheme of chasing the cells in bulb
labeled by DiI. (C) No migration of DiI labeled cells in bulb from amputation to 24 hpa.
EdU 1h pulse labeling at 24 hpa. (D) The comparison of proliferative cell number in
intact vs no amputation vs 24 hpa. EdU 1 h pulse labeling before amputation. (E)
Quantification of cells labeled before amputation. Intact: n = 6 (tentacles), 24 h: n = 6,
24 hpa: n = 6. Unpaired two-tailed *t*-test. *$p < 0.05$, **$p < 0.005$, ***$p < 0.001$. Scale
bars: (A, C and D) 100 μm.

1128

Figure S5. _Nanos1_⁺ cells in bulb decreased after X-ray irradiation.

(A) Relative expression of _Nanos1_ after irradiation by qPCR. (B) Expression of _Nanos1_ in bulb of intact tentacle after irradiation by FISH. (C) Quantification of _Nanos1_⁺ cells number in bulb. Control: n = 6 (tentacles), 3 dpi: n = 6. Unpaired two-tailed _t_-test. *$p <$ 0.05, **$p <$ 0.005, ***$p <$ 0.001. Scale bars: (B) 100 μm.

Figure S6. EdU long incorporation also labeled nematocyte progenitors.

(A) Nematocyte progenitor cells in bulb of intact tentacle with dual staining of _Nowa_ FISH and EdU staining. EdU 1 h or 24 h pulse labeling. (B) The rate of _Nowa_⁺ cells / EdU⁺ cells in intact tentacle. 1 h: n = 20 (tentacles), 24 h: n = 22. (C) Nematocyte progenitor cells in bulb with dual staining of _Mcol1_ FISH and EdU staining. EdU 1 h or 24 h pulse labeling. Unpaired two-tailed _t_-test. *$p <$ 0.05, **$p <$ 0.005, ***$p <$ 0.001. Scale bars: (A, C) 50 μm.

Figure S7. Similar pattern between β-catenin and aPKC antibody staining

Identification of cell types in blastema stained by β-catenin and aPKC antibody staining. Scale bar: 50 μm.

Figure S8. Injury-responsive cells decreased after UV exposure on blastema.

(A) Experimental scheme depicting the combination of EdU chasing and UV exposure. EdU 1 h pulse labeling at 24 hpa. (B) Bleaching of Hoechst signal only in UV exposure area at 24 hpa. White dot square is UV exposure area. (C) Distribution of EdU labeled cells 1 day after UV exposure. (D) The number of EdU⁺ cells in UV exposed tentacle at 48 hpa. Ctl: n = 7 (tentacles), UV: n = 8. Unpaired two-tailed _t_-test. *$p <$ 0.05, **$p <$ 0.005, ***$p <$ 0.001. Scale bars: (B, C) 100 μm.

Figure 1

Figure 2

Figure 3

Figure 4

A

Chasing

-1   0   24   48   72   hpa

EdU 1 h — Labeling homeostatic proliferative cells in bulb

Amputation

BrdU 1 h — Labeling repair-specific proliferative cells in blastema

B

DNA EdU BrdU

24 hpa   48 hpa   72 hpa

(i)   (ii)   (iii)

100 um

EdU

(i')   (ii')   (iii')

BrdU

(i'')   (ii'')   (iii'')

C Homeostatic proliferative cells

\# of cells labeled before amputation

n.s.   n.s.   ***

Intact   24 hpa   48 hpa   72 hpa

D Repair-specific proliferative cells

\# of cells labeled at 24 hpa

n.s.   ***

24 hpa   48 hpa   72 hpa

E

24   48   72   hpa

● Homeostatic proliferative cells
● Repair-specific proliferative cells

F

0 dpi   3 dpi   24 hpa

X-ray   Amputation   EdU 1 h

Stem cell elimination   Blastema formation

G DNA EdU

No irradiation   X-ray

(i)   (ii)

100 um

H

% EdU+ cells in blastema

n.s.

No irradiation   X-ray

Figure 5

Figure 6

A

Amputation 24 48 72 hpa

Wound healing · Blastema formation · Regenerative outgrowth

Cell proliferation

Differentiation

Homeostatic proliferative cells contribute to nematogenesis and tentacle elongation

Nematocytes

Epithelial cells

Repair-specific proliferative cells contribute to mass enlargement

Epithelial cells

● Homeostatic proliferative cells
● Repair-specific proliferative cells

B Appendage regeneration evolution model

Cnidarian Jellyfish

Injury

Blastema formation

Resident stem cell

Dedifferentiation?

Quiescent stem cell re-activation?

Repair-specific stem cell

Differentiated cell

Bilateria
e.g. salamander limbs

Injury

Blastema formation

Quiescent stem cell re-activation

Dedifferentiation

Repair-specific stem cell

Differentiated cell