Resurrection of plant disease resistance proteins via helper NLR bioengineering

Mauricio P. Contreras[1], Hsuan Pai[1], Muniyandi Selvaraj[1], AmirAli Toghani[1], David M. Lawson[2],

Yasin Tumtas[3], Cian Duggan[3], Enoch Lok Him Yuen[3], Clare E. M. Stevenson[2], Adeline Harant[1],

Chih-Hang Wu[4], Tolga O. Bozkurt[3], Sophien Kamoun[1,*] and Lida Derevnina[1,†,*]

1: The Sainsbury Laboratory, University of East Anglia, Norwich, United Kingdom.

2: Department of Biochemistry and Metabolism, John Innes Centre, Norwich, United Kingdom.

3: Department of Life Sciences, Imperial College, London, United Kingdom.

4: Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan.

†Current address: Crop Science Centre, Department of Plant Sciences, University of Cambridge, Cambridge, United Kingdom.

*Correspondence to: sophien.kamoun@tsl.ac.uk (SK); ld645@cam.ac.uk (LD)

One sentence summary: A helper NLR is mutated to evade inhibition by a parasite effector.

Abstract

Parasites counteract host immunity by suppressing helper NLR proteins that function as central nodes in immune receptor networks. Understanding the mechanisms of immunosuppression can lead to strategies for bioengineering disease resistance. Here, we show that a cyst nematode virulence effector binds and inhibits oligomerization of the helper NLR protein NRC2 by physically preventing intramolecular rearrangements required for activation. A single amino acid polymorphism at the binding interface between NRC2 and the inhibitor is sufficient for this helper NLR to evade immune suppression, thereby restoring the activity of multiple disease resistance genes. This points to a novel strategy for resurrecting disease resistance in crop genomes.

Introduction

The nucleotide binding and leucine-rich repeat (NLR) class of intracellular immune receptors are an important component of innate immunity in plants and animals. They mediate intracellular recognition of pathogens and subsequently initiate an array of immune responses in order to counteract infection (*1, 2*). NLRs can be activated by pathogen-secreted virulence proteins, termed effectors, which pathogens deliver into host cells to modulate host physiology (*2*). A hallmark of plant and animal NLR activation is their oligomerization into higher order immune complexes termed resistosomes or inflammasomes, respectively (*3-9*). These complexes initiate immune signaling via diverse mechanisms, often leading to a form of programmed cell death, termed hypersensitive response (HR) in plants or pyroptosis in animals (*10, 11*). Recent studies have reported NLR-like proteins mediating antiviral immunity and programmed cell death in prokaryotes via mechanisms analogous to those found in eukaryotic NLRs, suggesting that this is a conserved defense mechanism across all three domains of life (*12*). Remarkably, pathogen effectors can act both as triggers and suppressors of NLR-mediated immunity (*13*). In some cases, adapted pathogens deploy effectors which directly or indirectly interfere with NLR signaling to suppress immune activation (*12, 14-17*). However, the exact biochemical mechanisms by which pathogen effectors compromise NLR-mediated immunity to promote disease remain largely unknown. Moreover, whereas multiple strategies to bioengineer novel effector recognition specificities in NLRs have been proposed in recent years (*18*), approaches to mitigate the impact of effector-mediated immune suppression of NLRs are lacking.

NLRs belong to the signal ATPases with numerous domains (STAND) superfamily. They typically exhibit a tripartite domain architecture consisting of an N-terminal signaling domain, a central nucleotide binding domain and C-terminal superstructure forming repeats (*2*). The central domain, termed NB-ARC (nucleotide binding adaptor shared by APAF-1, plant R proteins and CED-4) in plant NLRs, is a hallmark of this protein family and plays a key role as a molecular switch, mediating conformational changes required for activation. NB-ARC domains consist of a nucleotide binding domain (NB), a helical domain (HD1), and a winged-helix domain (WHD) (*2, 19*). Diverse NLR activation and signaling strategies are found in nature. In some cases one NLR protein, termed a singleton, can mediate both elicitor perception and subsequent immune signaling (*20*). However, some NLRs can function as receptor pairs or in higher order configurations termed immune receptor networks (*13, 21*). In these cases, one NLR acts as a pathogen sensor, requiring a second helper NLR to initiate immune signaling. Such is the case in the solanaceous NRC immune receptor network,

which is comprised of multiple sensor NLRs that require an array of downstream helper NLRs termed

NRCs (NLRs required for cell death) to successfully initiate immune signaling. Upon perception of

their cognate effectors, sensors in this network activate oligomerization of their downstream NRC

helpers into a putative NRC resistosome, without stably forming part of the mature complex. This

has been termed the activation-and release model (*4*). The NRC network can encompass up to half of

the NLRome in some solanaceous plant species and plays a key role in mediating immunity against a

variety of plant pathogens including oomycetes, bacteria, viruses, nematodes and insects (*15, 21*).

Plant and metazoan parasites have evolved effectors that interfere with host NLR signaling to promote

disease. Parasite effectors can suppress NLR-mediated immunity indirectly by interfering with host

proteins that act downstream of NLR signaling (*15, 17, 22, 23*), or directly by interacting with NLRs

to inhibit their functions (*15, 16, 24*). One such example is the potato cyst nematode (*Globodera*

rostochiensis) effector, SPRYSEC15 (SS15), which can suppress signaling mediated by *Nicotiana*

benthamiana helper NLRs NRC2 and NRC3 and tomato (*Solanum lycopersicum*) helper NLR NRC1, by

binding to their central NB-ARC domains (*15*) (**Fig. 1A, Fig. S1A**). In this study, we reasoned that

mapping the binding interface between SS15 and its target helper NLRs would enable us to

bioengineer NLR variants that evade pathogen suppression, therefore resurrecting the immune

signaling activity of upstream sensors in the NRC network.

Results

First, we investigated how SS15 binding to NRC2 prevents immune signaling., notably whether SS15

prevents oligomerization of NRC following sensor NLR activation. To test this hypothesis, we

transiently co-expressed NRC2 or NRC4 with their upstream sensor Rx and the effector SS15 in leaves

of *nrc2/3/4* CRISPR KO *N. benthamiana* plants and leveraged previously established BN-PAGE-based

readouts for NRC resistosome formation (*4*). For biochemical analyses, we used previously described

NRC2 and NRC4 variants with mutations in their N-terminal MADA motifs (NRC2^{EEE} and NRC4^{AAA},

respectively) which abolish cell death induction without compromising receptor activation,

oligomerization, or localization (*4, 25, 26*). We activated the Rx-NRC system by co-expressing PVX

CP-GFP or free GFP as an inactive control. In the absence of SS15, both NRC2 and NRC4

oligomerize upon effector-triggered activation mediated by their upstream sensor Rx. However, when

SS15 is present, Rx/CP-activated NRC2 is unable to oligomerize and appears as a band of ~240 kDa,

which co-migrates with SS15. Inactive NRC2 co-expressed with SS15 also migrates as a band of ~240

kDa, which is slower-migrating relative to inactive NRC2 in the absence of SS15, indicative of *in vivo*

NRC2-SS15 complex formation (**Fig. 1B, Fig. S1**). We also observed that SS15 co-expression not

only blocks NRC2 oligomerization but also prevents the previously reported shift of NRC2 from

cytoplasm to PM as well as the formation of NRC2 PM-associated puncta upon Rx/CP activation

(**Fig. S2**) (*4*). In contrast, NRC4 oligomerization is not affected in the presence of SS15, which is in

line with previous findings that NRC4 immune signaling is not suppressed by SS15 (*15*). We conclude

that SS15 can suppress immune signaling by acting as a direct proteinaceous inhibitor of NRC2, but

not NRC4, by directly binding to its NB-ARC domain to block the formation of a signal-competent

oligomeric resistosome.

Given that NRC4 retains the capacity to oligomerize in the presence of SS15, we leveraged this

differential SS15 sensitivity between NRC2 and NRC4 to identify the domain that determines SS15

association and inhibition. We generated a series of NRC2-NRC4 chimeric proteins (**Fig. 2A-2B, Fig.**

S3), which we subsequently assayed for SS15 association via *in planta* co-immunoprecipitation. We

identified one chimeric variant of NRC4, carrying the HD1-1 region of NRC2 (NRC4[2HD1-1]), which

gains association to SS15 (**Fig. 2C**). Unlike NRC4, NRC4[2HD1-1] is susceptible to inhibition by SS15

and is unable to oligomerize and trigger cell death in the presence of SS15 (**Fig. 2D-2E, Fig. S3, Fig.**

S4). We conclude that the HD1-1 region is important for association to SS15 and for the effector to

directly inhibit NRC oligomerization and programmed cell death.

To further define the interface between SS15 and NRC proteins, we crystallized SS15 in complex with

the NB-ARC domain of SlNRC1, a tomato NRC that is inhibited by the nematode effector. We

subsequently solved the structure using X-ray diffraction data collected to 4.5 Å resolution (**Fig. 3A**,

Fig. S5, **Table S3**), which allowed us to determine that SS15 binds to a loop in the HD1-1 region of

NRCs which connects the NB domain to the HD1 and WHD domains. This provides orthogonal

evidence that the SS15-NRC interactions are mediated by this region as shown with the chimera

experiments (**Fig. 2**). This loop was previously shown to act as a "hinge", allowing the NB domain to

rotate relative to the HD1 and WHD domains following activation (**Fig. S5**, **Movie S1**) (*27*). We

propose that SS15 prevents conformational changes that are critical for NLR activation by binding

and immobilizing the NB-HD1 hinge.

Given that SS15 suppresses cell death induction mediated by SlNRC1, NRC2 and NRC3 but not
NRC4 or other well characterized NLRs such as ZAR1 (**Fig. S1**) (*15*), we leveraged the high degree
of conservation that is characteristic of plant NB-ARC domains to narrow down residues within the
binding interface that underpin this interaction. We shortlisted residues within the HD1-1 region that
are similar in SlNRC1, NRC2 and NRC3 but different in NRC4 or AtZAR1 (**Fig. 3B**). Combining
information from the co-crystal structure and the alignments allowed us to select 13 candidate residues
to test by mutagenesis in NRC2 (**Fig. 3B**). We mutated each of these residues to the corresponding
amino acid found in NRC4 and screened these NRC2 variants for susceptibility to SS15 inhibition in
cell death assays. This revealed two variants, NRC2^{E316P} and NRC2^{D317K} which triggered cell death
when activated by Rx/CP and were no longer inhibited by SS15 (**Fig. 3C, Fig. S6, Fig. S7**). We also
tested all 13 single amino acid mutants for association with SS15 by *in planta* co-immunoprecipitation
and found that NRC2^{E316P} and NRC2^{D317K} exhibited reduced association with SS15 relative to NRC2
(**Fig. 3D, Fig. S6**), which is in line with the observation that SS15 is not able to suppress these variants
(**Fig. 3C**). We conclude that the E316 and D317 residues are critical for SS15-mediated inhibition of
NRC2 and that mutating these residues to their equivalent amino acid in NRC4 allows Rx/CP-
activated NRC2 to evade SS15 association and inhibition.

We next tested whether these two SS15-evading variants of NRC2 could restore functionality of
NRC2-dependent sensor NLRs that are suppressed by the parasite effector. We tested this by
performing complementation assays with NRC2^{E316P} and NRC2^{D317K} in *nrc2/3/4* CRISPR KO *N.*
benthamiana plants. We activated the NRC2 variants with a panel of agronomically important sensor
NLRs mediating resistance to diverse pathogens, including the potato cyst nematode R gene Gpa2 (an
allele of Rx), as well as other well characterized oomycete and bacterial resistance proteins.
Remarkably, NRC2^{D317K} evaded SS15 inhibition with all tested sensor NLRs restoring their capacity
to activate immune signaling (**Fig 4A**, **Fig. S8, Fig. S9**). In contrast, NRC2^{E316P} could evade SS15
suppression when activated by Rx, but not when activated by other sensors. We therefore selected
NRC2^{D317K} for follow-up biochemical studies, using BN-PAGE-based assays. Unlike NRC2, activated
NRC2^{D317K} oligomerized even in the presence of SS15 and did not form an *in vivo* complex with the
inhibitor (**Fig. 4B**). We conclude that NRC2^{D317K} can fully evade SS15-mediated immune suppression,
retaining the capacity to oligomerize and mediate cell death when activated by multiple agronomically
important sensor NLRs.

Conclusion

Our study reveals how a parasite effector has evolved as an inhibitor of a helper NLR by directly binding its NB-ARC domain to prevent resistosome formation and immune signaling. By binding and immobilizing a critical hinge loop in the HD1 region of the NB-ARC, SS15 restricts movement of the NB domain relative to the HD1 and WHD domains, preventing immune receptor activation. Remarkably, while SS15 can bind and inhibit NRC2, it cannot bind or inhibit the NRC2 paralog NRC4. We exploited NRC4's resilience to SS15 inhibition by making chimeric NRC2-NRC4 variants which, together with structural information, helped us identify the inhibitor binding interface. Mutational studies of this interface allowed us to generate a single amino acid variant of NRC2 (NRC2^{D317K}) that evades SS15 inhibition without compromising receptor signaling capacity. This NRC2^{D317K} variant can now support signaling by any NRC2-dependent sensor even in the presence of SS15.

The existence of plant parasite secreted NLR inhibitors suggests that suppressed resistance genes may occur in crop genomes. Leveraging the approach detailed in this study, it may be possible to resurrect cryptic or defeated resistance proteins to enhance disease resistance. Moreover, considering that multiple sensors can signal through the same downstream helper, applying this approach to helper NLRs holds potential to simultaneously resurrect multiple upstream sensor NLRs. Importantly, the single amino acid NRC2 variants we identified could in theory be generated in-locus using gene editing technologies in agronomically important crop species, making deployment of this technology viable in countries where transgenic approaches are not feasible. Our work describes a novel approach to achieve robust immunity by engineering NLRs that avoid parasite suppression. This could in theory be applied to other plant, metazoan or even prokaryotic NLR immune receptors that are directly targeted by parasite effectors (*12, 15, 16, 24*). Combined with recent advances in NLR engineering to achieve novel pathogen recognition specificities (*18, 28*), this technology holds the potential to unlock a new era in disease resistance breeding.

Materials and Methods

Plant growth conditions

Wild-type and *nrc2/3/4* CRISPR mutant *Nicotiana benthamiana* lines were grown in a controlled environment growth chamber with a temperature range of 22 to 25 ºC, humidity of 45% to 65% and a 16/8-hour light/dark cycle.

Plasmid construction

We used the Golden Gate Modular Cloning (MoClo) kit (*29*) and the MoClo plants part kit (*30*) for cloning. All vectors used were generated with these kits unless otherwise stated. Cloning design and sequence analysis were done using Geneious Prime (v2021.2.2; https://www.geneious.com). Plasmid construction is described in more detail in **Table S1**.

Cell death assays by agroinfiltration

Proteins of interest were transiently expressed in *N. benthamiana* according to previously described methods (*31*). Briefly, leaves from 4–5-week-old plants were infiltrated with suspensions of *Agrobacterium tumefaciens* GV3101 pM90 strains transformed with expression vectors coding for different proteins indicated. Final OD_{600} of all *A. tumefaciens* suspensions were adjusted in infiltration buffer (10 mM MES, 10 mM $MgCl_2$ and 150 μM acetosyringone (pH 5.6)). Final OD_{600} used for each construct is described in **Table S2**.

Extraction of total proteins for BN-PAGE and SDS-PAGE assays

Four to five-week-old *N. benthamiana* plants were agroinfiltrated as described above with constructs of interest and leaf tissue was collected 3 days post agroinfiltration for NRC2 and 2 days post agroinfiltration for NRC4. Final OD_{600} used for each construct is described in **Table S2**. BN-PAGE was performed using the Bis-Tris Native PAGE system (Invitrogen) according to the manufacturer's instructions, as described previously (*31*). Leaf tissue was ground using a Geno/Grinder tissue homogenizer. For NRC2, GTMN extraction buffer was used (10% glycerol, 50 mM Tris-HCl (pH 7.5), 5 mM $MgCl_2$ and 50 mM NaCl) supplemented with 10 mM DTT, 1x protease inhibitor cocktail (SIGMA) and 0.2% Nonidet P-40 Substitute (SIGMA). For NRC4, GHMN buffer (10% glycerol, 50 mM HEPES (pH 7.4), 5 mM $MgCl_2$ and 50 mM NaCl) buffer supplemented with 10mM DTT, 1x protease inhibitor cocktail (SIGMA) and 1% Digitonin (SIGMA) was used for extraction. Samples were incubated in extraction buffer on ice for 10 minutes with short vortex mixing every 2 minutes.

216 Following incubation, samples were centrifuged at 5,000 x*g* for 15 minutes and the supernatant was
217 used for BN-PAGE and SDS-PAGE assays.
218

BN-PAGE assays
For BN-PAGE, samples extracted as detailed above were diluted as per the manufacturer's
instructions by adding NativePAGE 5% G-250 sample additive, 4x Sample Buffer and water. After
dilution, samples were loaded and run on Native PAGE 3%-12% Bis-Tris gels alongside either
NativeMark unstained protein standard (Invitrogen) or SERVA Native Marker (SERVA). The
proteins were then transferred to polyvinylidene difluoride membranes using NuPAGE Transfer
Buffer using a Trans-Blot Turbo Transfer System (Bio-Rad) as per the manufacturer's instructions.
Proteins were fixed to the membranes by incubating with 8% acetic acid for 15 minutes, washed with
water and left to dry. Membranes were subsequently re-activated with methanol to correctly visualize
the unstained native protein marker. Membranes were immunoblotted as described below.
229

SDS-PAGE assays
For SDS-PAGE, samples were diluted in SDS loading dye and denatured at 72 °C for 10 minutes.
Denatured samples were spun down at 5,000 x*g* for 3 minutes and supernatant was run on 4%-20%
Bio-Rad 4%-20% Mini-PROTEAN TGX gels alongside a PageRuler Plus prestained protein ladder
(Thermo Scientific). The proteins were then transferred to polyvinylidene difluoride membranes using
Trans-Blot Turbo Transfer Buffer using a Trans-Blot Turbo Transfer System (Bio-Rad) as per the
manufacturer's instructions. Membranes were immunoblotted as described below.
237

Immunoblotting and detection of BN-PAGE and SDS-PAGE assays
Blotted membranes were blocked with 5% milk in Tris-buffered saline plus 0.01% Tween 20 (TBS-T)
for an hour at room temperature and subsequently incubated with desired antibodies at 4 °C overnight.
Antibodies used were anti-GFP (B-2) HRP (Santa Cruz Biotechnology), anti-HA (3F10) HRP
(Roche), anti-Myc (9E10) HRP (Roche), and anti-V5 (V2260) HRP (Roche), all used in a 1:5000
dilution in 5% milk in TBS-T. To visualize proteins, we used Pierce ECL Western (32106, Thermo
Fisher Scientific), supplementing with up to 50% SuperSignal West Femto Maximum Sensitivity
Substrate (34095, Thermo Fishes Scientific) when necessary. Membrane imaging was carried out with
an ImageQuant LAS 4000 or an ImageQuant 800 luminescent imager (GE Healthcare Life Sciences,
Piscataway, NJ). Rubisco loading control was stained using Ponceau S (Sigma).

Co-Immunoprecipitation assays

CoIP assays were performed as described previously (*32*). Total soluble protein was extracted as described above from leaves of *N. benthamiana* 3 days post agro-infiltration using GTEN buffer (10% Glycerol, 25 mM Tris-HCl (pH 7.5), 1mM EDTA, 150 mM NaCl) supplemented with 2% (w/v) polyvinylpolypyrrolidone, 10 mM DTT and 1x protease inhibitor cocktail (SIGMA), 0.3% IGEPAL (SIGMA). OD_{600} used can be found in **Table S2**. Protein extracts were filtered using Minisart 0.45 µm filter (Sartorius Stedim Biotech, Goettingen, Germany). Part of the extract was set aside prior to immunoprecipitation and used for SDS-PAGE as described above. These were the inputs. 1.4 ml of the remaining filtered total protein extract was mixed with 30 µl of anti-c-Myc agarose beads (A7470, SIGMA) and incubated end over end for 90 minutes at 4 °C. Beads were washed 5 times with immunoprecipitation wash buffer (GTEN extraction buffer with 0.3% v/v IGEPAL (SIGMA)) and resuspended in 60 µl of SDS loading dye. Proteins were eluted from beads by heating for 10 minutes at 72 °C. Immunoprecipitated samples were used for SDS-PAGE and immunoblotted as described above and compared to the inputs.

Confocal microscopy

Three to four-week-old plants were agroinfiltrated as described above with constructs of interest. Final OD_{600} used for each construct is described in **Table S2**. Leaf tissue was prepared for imaging by sectioning of desired area surrounding an infiltration spot using a cork borer size 4, and were mounted, live, in wells containing dH2O made in Carolina Observation Gel to enable diffusion of gasses. The abaxial of the leaf tissue was imaged using a Leica SP8 with 40x water immersion objective. Laser excitations for fluorescent proteins were used as described previously (*33*), namely 488 nm (Argon) for GFP, 561/594 nm (Diode) for RFP and 405 nm (Diode) for BFP.

Membrane enrichment assays

Membrane enrichment was carried out by slightly modifying a previously described protocol (*34*). In brief, leaf material was ground to fine powder using liquid nitrogen and 2x volume of extraction buffer was added. Extraction buffer consisted of 0.81M sucrose, 5% (v/v) glycerol, 10 mM EDTA, 10 mM EGTA, 5mM KCl, and 100 mM Tris-HCl (pH 7.5) supplemented with 5 mM DTT, 1% Sigma Plant Protease Inhibitor Cocktail, 1 mM PMSF and 0.5% PVPP. After addition of the buffer, the samples were vortexed for a minute and the cell debris was cleared out by two subsequent centrifugation steps at 1000 xg for 5 min. The supernatant was diluted 1:1 using distilled water and an aliquot of the

supernatant was separated as the total fraction (T). The remaining supernatant (200-300 μL) was
further centrifuged at 21,000 xg for 90 min at 4°C. This centrifugation yielded the supernatant (soluble
fraction, S) and membrane enriched pellet (membrane fraction, M). After separating the soluble
fraction, the pellet was resuspended in diluted extraction buffer (without PVPP). All the fractions were
diluted with SDS loading dye, and proteins were denatured by incubating at 50°C for 15 min. Western
blotting was performed as previously described following SDS-PAGE. Endogenous plasma
membrane ATPase was detected using anti-H+ATPase (AS07 260) antibody (Agrisera) as a marker to
show the success of membrane enrichment.

Heterologous protein production and purification from _E. coli_

Heterologous production and purification of SS15 was performed as previously described (*32*).
Recombinant SS15 protein (lacking signal peptide) was expressed by cloning in pOPIN-S3C plasmid,
with an N-terminal tandem 6xHis-SUMO followed by a 3C protease cleavage site. pOPIN-S3C:SS15
was transformed into _E. coli_ SHuffle cells. 8L of these cells were grown at 30 ºC in autoinduction
media (*35*) to an OD600 of 0.6 to 0.8 followed by overnight incubation at 18 ºC and harvested by
centrifugation. Pelleted cells were resuspended in 50 mM Tris HCl (pH 8), 500 mM NaCl, 50 mM
Glycine, 5% (vol/vol) glycerol and 20 mM imidazole (buffer A) supplemented with cOmplete EDTA-
free protease inhibitor tablets (Roche) and lysed by sonication. The clarified cell lysate was applied to
a Ni^{2+}-NTA column connected to an AKTA pure system. 6xHis-SUMO-SS15 was step eluted with
elution buffer (buffer A with 500 mM imidazole) and directly injected onto a Superdex 200 26/600
gel filtration column pre-equilibrated with buffer B (20 mM HEPES (pH 7.5), 150 mM NaCl). The
fractions containing 6xHis-SUMO-SS15 were pooled, and the N-terminal 6xHis-SUMO tag was
cleaved by addition of 3C protease (10 μg/mg of fusion protein), incubating overnight at 4ºC. Cleaved
SS15 was further purified using a Ni^{2+}-NTA column, this time collecting the flow through to separate
the cleaved tag from the SS15 protein. Un-tagged SS15 was further purified by another round of gel
filtration as described above. The concentration of protein was judged by absorbance at 280 nm (using
a calculated molar extinction coefficient of 35920 M^{-1cm-1} for SS15).

Heterologous production and purification of SlNRC1[NB-ARC] was performed as previously described
(*36*). Recombinant SlNRC1[NB-ARC] protein was also expressed cloning in pOPIN-S3C plasmid as
described above. pOPIN-S3C:SlNRC1[NB-ARC] was transformed into _E. coli_ Lemo21 (DE3) cells. 8L of
these cells were grown at 37 ºC in autoinduction media (*35*) to an OD600 of 0.6 to 0.8 followed by

overnight incubation at 18 ℃ and harvested by centrifugation. Pelleted cells were resuspended in 50

mM Tris HCl (pH 8), 500 mM NaCl, 50 mM Glycine, 5% (vol/vol) glycerol and 20 mM imidazole

(buffer A) supplemented with cOmplete EDTA-free protease inhibitor tablets (Roche) and lysed by

sonication. The clarified cell lysate was applied to a Ni^{2+}-NTA column connected to an AKTA pure

system. 6xHis-SUMO-SlNRC1^{NB-ARC} was step eluted with elution buffer (buffer A with 500 mM

imidazole) and directly injected onto a Superdex 200 26/600 gel filtration column pre-equilibrated

with buffer B (20 mM HEPES (pH 7.5), 150 mM NaCl). The fractions containing 6xHis-SUMO-

SlNRC1^{NB-ARC} were pooled, and the N-terminal 6xHis-SUMO tag was cleaved by addition of 3C

protease (10 µg/mg of fusion protein), incubating overnight at 4℃. Cleaved SlNRC1^{NB-ARC} was further

purified using a Ni^{2+}-NTA column, this time collecting the flow through to separate the cleaved tag

from the SlNRC1^{NB-ARC} protein. Un-tagged SlNRC1^{NB-ARC} was further purified by another round of gel

filtration as described above. The concentration of protein was judged by absorbance at 280 nm (using

a calculated molar extinction coefficient of 63370 M^{-1cm-1} for SlNRC1^{NB-ARC}).

To obtain SlNRC1^{NB-ARC} in complex with SS15, both proteins were incubated in a 1:1 molar ratio

overnight at 4 ℃ and subjected to gel filtration on a Superdex 200 26/600 gel filtration column as

described above. The fractions containing SlNRC1^{NB-ARC} in complex with SS15 were pooled,

concentrated to 10-15 mg/ml and subsequently used for crystallisation screens.

Crystallisation, data collection, and structure solution

Crystallisation screens were performed at 18°C using the sitting-drop vapour diffusion technique.

Drops composed of 0.3 µL of protein solution and 0.3 µL of reservoir solution were set up in MRC

96-well crystallisation plates (Molecular Dimensions), which were dispensed using an Oryx Nano or

an Oryx8 robot (Douglas Instruments). Crystal growth was monitored using a Minstrel Desktop

Crystal Imaging System (Rikagu). Suitable crystals grew after 72 hrs in a Morpheus screen

crystallisation condition containing 0.1M MES buffer (pH 6.5), 10% (w/v) PEG8000 and 20% (v/v)

ethylene glycol (Molecular Dimensions) and were harvested by flash-cooling in liquid nitrogen using

LithoLoops (Molecular Dimensions). X-ray diffraction data were collected at the Diamond Light

Source (Didcot, UK) on beamline I03 using an Eiger2 XE 16M pixel array detector (Dectris) with

crystals maintained at 100 K by a Cryojet cryocooler (Oxford Instruments).

X-ray data were integrated, and scaled using XDS (*37*), as implemented through the XIA2 (*38*)

pipeline, and then merged using AIMLESS (*39*), via the CCP4i2 graphical user interface (*40*). The

NRC1^{NB-ARC}-SS15 complex crystallised in space group $P6_1$ with cell parameters $a=b=$128.6, $c=$170.7 Å,

and the best crystal yielded diffraction data to 4.5 Å resolution (see **Table S3** for a summary of data

collection and refinement statistics). Given the small size of the dataset, we assigned 10% of the data

(883 unique reflections) for the R_{free} calculation, to give a more statistically meaningful metric. The

crystal structure of NRC1^{NB-ARC} alone was already available (PDB 6S2P), but there was no

experimentally determined structure for SS15. Thus, we made use of AlphaFold2-multimer (*41*), as

implemented through Colabfold

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb)

(*42*) to generate structural predictions for the complex. There was very good sequence coverage for

both proteins and the five independent models of the individual components were closely similar. The

pLDDT scores were generally good (e.g. averages of 82 and 75 for NRC1^{NB-ARC} and SS15 models,

respectively, from the rank 1 predictions). However, the relative placement of the two components

varied across the five models and the corresponding PAE scores indicated very low confidence in

these predictions. A comparison of the five NRC1^{NB-ARC} models with the known crystal structure

showed a good agreement (e.g. superposition of the rank 1 model gave an rmsd of 1.77 Å). Given that

the AlphaFold2 (AF2) model provided starting coordinates for several loops missing from the crystal

structure, we decided to use this model in molecular replacement. Templates for both components

were prepared using the "Process Predicted Models" CCP4i2 task, which removed low confidence

regions (based on pLDDT) and converted the pLDDT scores in the B-factor field of the PDB

coordinate files to pseudo B-factors. Analysis of the likely composition of the asymmetric unit (ASU)

suggested that it contained two copies of each component, giving an estimated solvent content of

~67%. PHASER (*43*) was able to place the four chains within the ASU, although the second SS15

domain required manual repositioning with respect to its neighbouring NRC1 domain in order to

avoid a number of clashes and improve the fit to the density. This was achieved using COOT (*44*)

and guided by the arrangement of the other NRC1-SS15 complex (**Fig. S4**). The structure was then

subjected to jelly body refinement in REFMAC5 (*45*) using ProSMART restraints (*46*) generated from

the AF2 models, giving R_{work} and R_{free} values of 0.357 and 0.401, respectively, to 4.5 Å resolution.

Now it was possible to generate more complete models for the components by superposing the

original unprocessed AF2 models and trimming these with reference to the improved electron density.

Furthermore, a significant region of positive difference density was present at the cores of both NRC1

domains, which corresponded to the ADP in the crystal structure; thus we incorporated ADP into the

model. Due to the low resolution of the dataset, only very limited rebuilding was possible in COOT,
where Geman-McClure and Ramachandran restraints were used to maintain good stereochemical
parameters. After several cycles of restrained refinement in REFMAC5 and editing in COOT, a
reasonable model was obtained with R_{work} and R_{free} values of 0.258 and 0.298, respectively. However,
there remained a region of positive difference density near the N-termini of both SS15 domains that
we could not adequately explain. At this point we re-ran the AlphaFold2-multimer predictions, but
this time with one copy of the complex taken from the crystal structure as a reference template.
Although the new predictions did not produce complexes that were consistent with the X-ray data,
and the models for the individual components did not appear to be noticeably improved based on
pLDDT scores, we used them as starting points to rebuild the X-ray structure.

Significantly, for several models, the N-terminal region of SS15 adopted conformations that partially
accounted for the region of positive difference density electron density, and this could be improved
by careful rebuilding and refinement. This "AlphaFold recycling" procedure was repeated a further
two times before finalising the structure, which included residues 153-494 for SlNRC1 (numbered
with respect to full length protein) and residues 18-223 for SS15, where residues 33-43 in both copies
of the latter formed α-helices that occupied the regions of positive difference density observed earlier.
For the last refinement in REFMAC5, the following options were used: ProSMART restraints
generated from the latest AF2 models, overall B-factor refinement with TLS restraints (where each
protein chain was treated as a separate domain), and non-crystallographic symmetry restraints. The
final model gave R_{work} and R_{free} values of 0.237 and 0.275, respectively, to 4.5 Å resolution (see **Table
S3** for a summary of refinement statistics). All structural figures were prepared using ChimeraX (*47*)
and PyMOL (*48*).

 Figures:

 Figure 1

Fig. 1: SS15 directly inhibits NRC2, but not NRC4 oligomerization.

(**A**) Schematic representation of the NRC immune receptor network, consisting of multiple sensor

NLRs (in blue) and their downstream helper NLRs, NRC2 and NRC4 (in purple and green,

respectively). Effector-triggered activation of a sensor leads to downstream helper oligomerization

and resistosome formation. The *G. rostochiensis* effector SS15 (in yellow) can directly bind to the NB-

ARC domain of NRC2 but not NRC4, thereby inhibiting signaling by directly binding to the NB-ARC

domain of this helper NLR. (**B**) BN-PAGE assays with inactive and activated Rx together with NRC2

or NRC4, in the absence or presence of SS15. C-terminally V5-tagged Rx and C-terminally 4xMyc-

tagged NRC2^{EEE} or NRC4^{AAA} were co-expressed with either free GFP or C-terminally GFP-tagged

PVX CP. These effector-sensor-helper combinations were co-infiltrated together with a 6xHA-

mCherry fusion protein or with N-terminally 6xHA-tagged SS15. Total protein extracts were run in

parallel on native and denaturing PAGE assays and immunoblotted with the appropriate antisera

labelled below. Approximate molecular weights (kDa) of the proteins are shown on the left.

Corresponding SDS-PAGE assays can be found in **Fig. S1**. The experiment was repeated three times

with similar results.

Figure 2

Fig. 2: The HD1-1 region of NRC NB-ARC domains determines sensitivity to SS15.

(A) Schematic representation of NRC domain architecture, highlighting regions within the NB-ARC domain swapped between NRC2-NRC4 chimeric proteins, namely NRC4[2HD1-1] and NRC4[2HD1-2]. Association (+) or lack thereof (-) between these NLR immune receptors and SS15 determined by *in planta* co-immunoprecipitation is detailed on the right. (**B**) Close-up view of amino acid sequence alignment between AtZAR1, NRC2 and NRC4 focused on the HD1 region of NB-ARC domain. Predicted secondary structure is shown above the alignment. Well-characterized motifs within this region, such as RNBS-C and GLPL are underlined below the alignment. (**C**) Co-Immunoprecipitation (Co-IP) assays between SS15 and chimeric NRC2-NRC4 variants. C-terminally 4xMyc-tagged NRC proteins were transiently co-expressed with N-terminally 4xHA-tagged SS15. IPs were performed with agarose beads conjugated to Myc antibodies (Myc IP). Total protein extracts were immunoblotted

with the appropriate antisera labelled on the left. Approximate molecular weights (kDa) of the proteins

are shown on the right. Rubisco loading control was carried out using Ponceau stain (PS). The

experiment was repeated three times with similar results. (**D**) Photo of representative leaves from *N.*

benthamiana nrc2/3/4 KO plants showing HR after co-expression of Rx and PVX CP with NRC2,

NRC4 and the two NRC2-NRC4 chimeras, NRC4^{2HD1-1} and NRC4^{2HD1-2}. These effector-sensor-helper

combinations were co-expressed with a free mCherry-6xHA fusion protein (EV) or with N-terminally

4xHA-tagged SS15. The experiment consisted of 3 biological replicates with at least 6 technical

replicates each. A quantitative analysis of the HR phenotypes can be found in **Fig. S4**. (**E**) BN-PAGE

assay with inactive and activated Rx together with NRC4 or an NRC2-NRC4 chimeric protein in the

absence or presence of SS15. C-terminally V5-tagged Rx and C-terminally 4xMyc-tagged NRC4^{AAA} or

NRC4$^{AAA-2HD1-1}$ were co-expressed with either free GFP or C-terminally GFP-tagged PVX CP. These

effector-sensor-helper combinations were co-infiltrated together with a mCherry-6xHA fusion protein

or with N-terminally 4xHA-tagged SS15. Total protein extracts were run in parallel on native and

denaturing PAGE assays and immunoblotted with the appropriate antisera labelled on the left.

Corresponding SDS-PAGE blots can be found in **Fig. S3**. Approximate molecular weights (kDa) of

the proteins are shown on the left. The experiment was repeated three times with similar results.

Figure 3

Fig. 3: SS15-NRC binding interface enables bioengineering NRC2 variants that evade suppression. (A) Crystal structure of the SS15-SlNRC1[NB-ARC] complex. The NB, HD1 and WHD domains of SlNRC1[NB-ARC] are shown in cyan, pale blue and magenta, respectively; SS15 is in dark blue. The inset displays a close-up of the interface between SS15 and the HD1 domain of SlNRC1[NB-ARC] with the residues corresponding to those selected for mutagenesis in NRC2 highlighted in stick representation. See **Table S3** for a summary of data collection and processing statistics. (**B**) Alignment of HD1-1 region of AtZAR1, SlNRC1, NRC2, NRC3, and NRC4. Candidate residues (highlighted in red) were shortlisted based on the interface identified in the co-crystal structure of SS15 and the

SlNRC1 NB-ARC domain, as well as being conserved in SlNRC1, NRC2 and NRC3 but not NRC4

and AtZAR1. 13 NRC2 variants were generated by mutating individual candidate positions to the

corresponding amino acid in NRC4 (detailed underneath the alignment). (**C**) Photo of representative

leaves from *N. benthamiana nrc2/3/4* KO plants showing HR after co-expression of Rx and PVX CP

with NRC2, or different NRC2 variants generated. These effector-sensor-helper combinations were

co-expressed with a free mCherry-6xHA fusion protein (EV) or with N-terminally 4xHA-tagged SS15.

The experiment consisted of 3 biological replicates. A quantitative analysis of HR phenotypes can be

found in **Fig. S7**. (**D**) Co-Immunoprecipitation (Co-IP) assays between SS15 and NRC2 variants. C-

terminally 4xMyc-tagged NRC2 variants were transiently co-expressed with N-terminally 4xHA-

tagged SS15. IPs were performed with agarose beads conjugated to Myc antibodies (Myc IP). While

only data for E316P, D317K and W320F NRC2 variants is shown, additional variants were also tested

and can be found in **Fig S6**. Total protein extracts were immunoblotted with appropriate antisera

labelled on the left. Approximate molecular weights (kDa) of the proteins are shown on the right.

Rubisco loading control was carried out using Ponceau stain (PS). The experiment was repeated three

times with similar results.

Figure 4

Fig. 4: NRC2^{D317K} helper restores immune signaling of multiple disease resistance genes in

the presence of the effector SS15. (A) Photo of representative leaves from *N. benthamiana nrc2/3/4*

KO plants showing HR after co-expression of Rx/CP, Pto/AVRPto or Gpa2/RBP1 together with

NRC2, NRC2^{E316P} or NRC2^{D317K} in the absence or presence of SS15. The experiment consisted of 3

biological replicates. A quantitative analysis of HR phenotypes can be found in **Fig. S8**. (**B**) BN-

PAGE assays with inactive and activated Rx together with NRC2 or NRC2^{D317K}, in the absence or

presence of SS15. C-terminally V5-tagged Rx and C-terminally 4xMyc-tagged NRC2^{EEE} or NRC2^{EEE-}

D317K were co-expressed with either free GFP or C-terminally GFP-tagged PVX CP. These effector-

sensor-helper combinations were co-infiltrated together with a 6xHA-mCherry fusion protein or with

N-terminally 4xHA-tagged SS15. Total protein extracts were run in parallel on native and denaturing

PAGE assays and immunoblotted with the appropriate antisera labelled below. Approximate

molecular weights (kDa) of the proteins are shown on the left. Corresponding SDS-PAGE blots can

be found in **Fig. S9**. The experiment was repeated three times with similar results.

Supplemental Data

Figure S1

Fig. S1: SS15 suppresses cell death mediated by SlNRC1, NRC2 and NRC3 but not NRC4 or NbZAR1.

(**A**) Photo of representative leaves from *N. benthamiana nrc2/3/4* KO plants showing HR after co-expression of various autoactive NLR variants with a free mCherry-6xHA fusion protein (EV) or with N-terminally 4xHA-tagged SS15. Images are representative of (**B**) SDS-PAGE accompanying BN-PAGE shown in **Fig. 1B**. Total protein extracts were immunoblotted with the appropriate antisera labelled on the left. Approximate molecular weights (kDa) of the proteins are shown on the right. Rubisco loading control was carried out using Ponceau stain (PS). The experiment was repeated three times with similar results.

Figure S2

Fig. S2: SS15 inhibits plasma membrane-association of activated NRC2.

(**A-D**) C-terminally GFP-tagged NRC2^{EEE} and C-terminally RFP-tagged Rx were co-expressed with

an EV-4xMyc construct or a CP-4xMyc construct in leaves of *nrc2/3/4* KO *N. benthamiana.*

Representative single-plane confocal micrographs show the localization of both components of the

inactive and active Rx-NRC2 system. Scale bars represent 10 μm. (**A**) NRC2^{EEE}-GFP and Rx-RFP co-

localize in the cytoplasm. (**B**) As reported previously, Rx/CP activated NRC2^{EEE} forms plasma

membrane-associated puncta while Rx remains in the cytoplasm. (**C**) Co-expression with SS15 does

not alter the localization of inactive NRC2^{EEE}-GFP or Rx-RFP. (**D**) Upon co-expression CP-4xMyc,

Rx-RFP and NRC2^{EEE}-GFP with SS15, the punctate localization for NRC2^{EEE}-GFP is no longer

observed. (**E**) Membrane enrichment assays are consistent with microscopy. As reported previously,

inactive NRC2^{EEE}-GFP is mostly present in the soluble fraction (S) whereas activated NRC2^{EEE}-GFP

exhibits equal distribution across soluble and membrane (M) fractions. Upon co-expression with SS15,

NRC2^{EEE}-GFP distribution remains in the soluble fraction regardless of the presence or absence of

PVX CP. The experiment was repeated twice with similar results.

Figure S3

Fig. S3: SS15 inhibits NRC2 by interacting with the HD1-1 region of the NB-ARC domain.

(A) Schematic representation of all NRC2-NRC4 chimeric proteins generated. Association with SS15

(+) or lack thereof (-) is indicated on the right. (**B**) Co-Immunoprecipitation (Co-IP) assays between

SS15 and chimeric NRC2-NRC4 variants. C-terminally 4xMyc-tagged NRC proteins were transiently

co-expressed with N-terminally 4xHA-tagged SS15. IPs were performed with agarose beads

conjugated to Myc antibodies (Myc IP). Total protein extracts were immunoblotted with the

appropriate antisera labelled on the left. Approximate molecular weights (kDa) of the proteins are

shown on the right. Rubisco loading control was carried out using Ponceau stain (PS). The experiment

was repeated three times with similar results. (**C**) SDS-PAGE accompanying BN-PAGE shown in

Fig. 2E. Total protein extracts were immunoblotted with the appropriate antisera labelled on the left.

Approximate molecular weights (kDa) of the proteins are shown on the right. Rubisco loading control

was carried out using Ponceau stain (PS). The experiment was repeated three times with similar results.

Figure S4

Fig. S4: NRC4^{HD1-1} chimera is susceptible to inhibition by SS15.

HR scores accompanying **Fig. 2D**. In all cases, Rx/CP was used to activate the system. HR was scored

based on a modified 0–7 scale (*49*) between 5–7 days post-infiltration. HR scores are presented as dot

plots, where the size of each dot is proportional to the number of samples with the same score (Count).

Results are based on 3 biological replicates. Statistical tests were implemented using the besthr R

library (*50*). We performed bootstrap resampling tests using a lower significance cut-off of 0.025 and

an upper cut-off of 0.975. Mean ranks of test samples falling outside of these cut-offs in the control

samples bootstrap population were considered significant. Significant differences between the

conditions are indicated with an asterisk (*).

Figure S5

Fig. S5: Crystal structure of SS15 in complex with SlNRC1[NB-ARC].

Electron density map showing the relative orientation and arrangement of SS15 (orange) and SlNRC1[NB-ARC] (violet) within an asymmetric unit. 2Fo-Fc map countered at 1σ **(B)** Two possible interfaces between SS15 and SlNRC1[NB-ARC] revealed from the crystal packing. Both interfaces (Interface 1 and Interface 2) are outlined (Left). Modelling of both potential binding interfaces for

SS15 complex with full length SlNRC1 (magenta) reveals a steric clash between the CC-domain of

SlNRC1 and SS15, making interface 2 unlikely to be biologically relevant in the full-length context

(Right). **(C)** Close up view of interaction between SS15-SlNRC1^{NB-ARC} interaction interface relative to

the ATP-binding site within the NB-ARC domain of SlNRC1. The pyrophosphate moiety of ADP is

oriented facing opposite the SS15 binding interface (shown as ball and sticks), suggesting that SS15 is

unlikely to displace bound nucleotide or prevent ATP hydrolysis. **(D)** Structure of SS15-SlNRC1^{NB-ARC}

(yellow, PDB 8BV0) is superimposed over the NB-ARC domain of AtZAR1 in its inactive (green,

PDB 6J5W), intermediate (cyan, PDB 6J5V), and active resistosome (magenta, 6J5T) conformations.

Visualizing these three states reveals the trajectory of the NB domain as it moves relative to the HD1

and WHD domains while changing from inactive to activated states. The binding of SS15 at the critical

hinge region between the NB and HD1-WHD domains likely immobilizes this loop, preventing these

critical intramolecular rearrangements and therefore preventing NLR activation. See **Movie S1**.

Figure S6

A

B

Fig. S6: Out of 13 NRC2 variants tested, only E316P and D317K mutations abolish SS15 association and HR suppression. (A) Photo of representative leaves from *N. benthamiana nrc2/3/4* KO plants showing HR after co-expression of Rx and PVX CP with NRC2, or the different NRC2 variants generated. These effector-sensor-helper combinations were co-expressed with a free mCherry-6xHA fusion protein (EV) or with N-terminally 4xHA-tagged SS15. (**B**) Co-Immunoprecipitation (Co-IP) assays between SS15 and NRC2 variants. C-terminally 4xMyc-tagged

577   NRC2 variants were transiently co-expressed with N-terminally 4xHA-tagged SS15. IPs were
578   performed with agarose beads conjugated to Myc antibodies (Myc IP). Total protein extracts were
579   immunoblotted with appropriate antisera labelled on the left. Approximate molecular weights (kDa)
580   of the proteins are shown on the right. Rubisco loading control was carried out using Ponceau stain
581   (PS). The experiment was repeated three times with similar results.
582

Figure S7

Fig. S7: NRC2^{E316P} and NRC2^{D317K} abolish SS15-mediated suppression of Rx.

HR scores accompanying **Fig. 3C**. In all cases, Rx/CP was used to activate the system. HR was scored based on a modified 0–7 scale (*49*) between 5–7 days post-infiltration. HR scores are presented as dot plots, where the size of each dot is proportional to the number of samples with the same score (Count). Results are based on 3 biological replicates. Statistical tests were implemented using the besthr R library (*50*). We performed bootstrap resampling tests using a lower significance cut-off of 0.025 and an upper cut-off of 0.975. Mean ranks of test samples falling outside of these cut-offs in the control samples bootstrap population were considered significant. Significant differences between the conditions are indicated with an asterisk (*).

Figure S8

Fig. S8: NRC2^{D317K} abolishes SS15-mediated suppression of Rx, Gpa2 and Prf.

HR scores accompanying **Fig. 4A**. NRCs were activated using Rx/CP (**A**), Pto/AVRPto (**B**) or

Gpa2/RBP1 (**C**). HR was scored based on a modified 0–7 scale (*49*) between 5–7 days post-

infiltration. HR scores are presented as dot plots, where the size of each dot is proportional to the

number of samples with the same score (count). Results are based on 3 biological replicates. Statistical

tests were implemented using the besthr R library (*50*). We performed bootstrap resampling tests using

a lower significance cut-off of 0.025 and an upper cut-off of 0.975. Mean ranks of test samples falling

outside of these cut-offs in the control samples bootstrap population were considered significant.

Significant differences between the conditions are indicated with an asterisk (*).

Figure S9

A

B

Fig. S9: NRC2^{D317K} abolishes SS15-mediated suppression of all NRC2-dependent sensors tested and restores NRC2 resistosome formation.

Photo of representative leaves from *N. benthamiana nrc2/3/4* KO plants showing HR after co-expression of NRC2, or different NRC2 variants generated with various sensor/effector pairs. These effector-sensor-helper combinations were co-expressed with a free mCherry-6xHA fusion protein (EV) or with N-terminally 4xHA-tagged SS15. (**B**) SDS-PAGE accompanying BN-PAGE shown in **Fig. 4B**. Total protein extracts were immunoblotted with the appropriate antisera labelled on the left. Approximate molecular weights (kDa) of the proteins are shown on the right. Rubisco loading control was carried out using Ponceau stain (PS). The experiment was repeated three times with similar results.

Supplementary Tables

620 **Table S1: List of constructs used in this study.** (As separate file)

621 **Table S2: List of OD$_{600}$ used for agroinfiltration experiments.** (As separate file)

622 **Table S3: Summary of X-ray data and model parameters for NRC1-SS15.**

Data collection	
Diamond Light Source beamline	I03
Wavelength (Å)	0.9763
Detector	Eiger2 XE 16M
Resolution range (Å)	51.34 – 4.50 (5.03 – 4.50)
Space Group	$P6_1$
Cell parameters (Å)	$a = b = 128.6$, $c = 170.7$
Total no. of measured intensities	77102 (13865)
Unique reflections	8981 (2263)
Multiplicity	8.6 (6.1)
Mean $I/\sigma(I)$	7.2 (1.5)
Completeness (%)	94.0 (84.0)
R_{merge}[a]	0.084 (1.513)
R_{meas}[b]	0.033 (0.612)
$CC_{1/2}$[c]	0.998 (0.588)
Refinement	
Resolution range (Å)	51.34 – 4.50 (4.62 – 4.50)
Reflections: working/free[d]	8075/883
R_{work}/ R_{free}[e]	0.237/0.275
MolProbity score/Clashscore[f]	1.58/5.41
Ramachandran plot: favoured/allowed/disallowed[f] (%)	95.9/4.1/0.0
R.m.s. bond distance deviation (Å)	0.003
R.m.s. bond angle deviation (°)	0.79
NRC1 – chains/no. protein residues/ranges	A,C/343/153-494
SS15 – chains/no. protein residues/ranges	B,D/206/18-223
No. ADP molecules/RSCC[g]	2/0.72,0.84

PDB accession code	8BV0

Values in parentheses are for the outer resolution shell.

[a] $R_{merge} = \sum_{hkl} \sum_I |I_i(hkl) - \langle I(hkl) \rangle| / \sum_{hkl} \sum_i I_i(hkl)$.

[b] $R_{meas} = \sum_{hkl} [N/(N-1)]^{1/2} \times \sum_I |I_i(hkl) - \langle I(hkl) \rangle| / \sum_{hkl} \sum_i I_i(hkl)$, where $I_i(hkl)$ is the ith observation of reflection hkl, $\langle I(hkl) \rangle$ is the weighted average intensity for all observations I of reflection hkl and N is the number of observations of reflection hkl.

[c] $CC_{1/2}$ is the correlation coefficient between symmetry equivalent intensities from random halves of the dataset.

[d] The data set was split into "working" and "free" sets consisting of 90 and 10% of the data respectively. The free set was not used for refinement.

[e] The R-factors R_{work} and R_{free} are calculated as follows: $R = \sum(|F_{obs} - F_{calc}|)/\sum|F_{obs}|$, where F_{obs} and F_{calc} are the observed and calculated structure factor amplitudes, respectively.

[f] As calculated using MolProbity (51).

[g] Real Space Correlation Coefficient as calculated by the PDB validation server.

Supplementary Movies

Movie S1: SS15 immobilizes hinge region between NB and HD1-WHD domains of NRCs to prevent NLR activation.

Acknowledgements

We are very thankful to Jiorgos Kourelis, Clemence Marchal, Daniel Lüdke and Hee-Kyung Ahn (The Sainsbury Laboratory, Norwich, UK) for invaluable scientific discussions, ideas, and feedback. We thank Julia Mundy for assistance with crystallography experiments. We thank the Diamond Light Source, UK (beamline i03 under proposal MX18565) for access to X-ray data collection facilities. M.P.C. thanks Emiliano "Dibu" Martínez and Lionel Messi for inspiration. We thank all members of the TSL Support Services for their invaluable assistance.

Funding:

The Kamoun Lab is funded primarily from the Gatsby Charitable Foundation, Biotechnology and Biological Sciences Research Council (BBSRC, UK, BB/WW002221/1, BB/V002937/1, BBS/E/J/000PR9795 and BBS/E/J/000PR9796) and the European Research Council (BLASTOFF).

Author Contributions:

Conceptualization: M.P.C, C-H.W., S.K., L.D.

Methodology: M.P.C., H.P., M.S., D.M.L., C.E.M.S., L.D.

Validation: M.P.C., H.P., M.S., Y.T., C.D., E.L.H.Y., A.H.

Formal Analysis: M.P.C., H.P., M.S., A.T., D.M.L., Y.T., C.D., C.E.M.S.

Investigation: M.P.C., H.P., M.S., Y.T., C.D., E.L.H.Y., A.H., L.D.

Data Curation: M.P.C., H.P., M.S., A.T., D.M.L., Y.T., C.D., E.L.H.Y., C.E.M.S., A.H.

Visualization: M.P.C., H.P., M.S., A.T., Y.T., C.D., E.L.H.Y., L.D.

Writing – Original Draft: M.P.C., S.K., L.D.

Writing – Review & Editing: M.P.C., D.M.L., S.K., L.D.

Supervision: M.P.C., T.O.B., S.K., L.D.

Project Administration: M.P.C., T.O.B., S.K., L.D.

Funding Acquisition: T.O.B., S.K., L.D.

Competing interests:

C.D., T.O.B and S.K. receive funding from industry on NLR biology. M.P.C., S.K. and L.D. have filed patents on NLR biology.

Data and materials availability:

All relevant data are within the paper, in the Supplementary Materials and Source Data files. Data

relevant to the structure presented in **Fig. 3** can be found in the Protein Data Bank, PDB 8BV0.

References

1.   J. D. Jones, R. E. Vance, J. L. Dangl, Intracellular innate immune surveillance devices in
plants and animals. *Science* **354**, aaf6395 (2016).
2.   Z. Duxbury, C.-h. Wu, P. Ding, A comparative overview of the intracellular guardians of
plants and animals: NLRs in innate immunity and beyond. *Annual review of plant biology*
72, 155-184 (2021).
3.   J. Wang *et al.*, Reconstitution and structure of a plant NLR resistosome conferring
immunity. *Science* **364**, eaav5870 (2019).
4.   M. P. Contreras *et al.*, Sensor NLR immune proteins activate oligomerization of their NRC
helper. *The EMBO Journal, in press* (2022).
5.   H.-K. Ahn *et al.*, Effector-dependent activation and oligomerization of NRC helper NLRs by
Rpi-amr3 and Rpi-amr1. *The EMBO Journal, in press* (2022).
6.   A. Förderer *et al.*, A wheat resistosome defines common principles of immune receptor
channels. *bioRxiv*, 2022.2003.2023.485489 (2022).
7.   R. Martin *et al.*, Structure of the activated ROQ1 resistosome directly recognizing the
pathogen effector XopQ. *Science* **370**, eabd9993 (2020).
8.   I. V. Hochheiser *et al.*, Structure of the NLRP3 decamer bound to the cytokine release
inhibitor CRID3. *Nature* **604**, 184-189 (2022).
9.   Z. Hu *et al.*, Structural and biochemical basis for induced self-propagation of NLRC4.
Science **350**, 399-404 (2015).
10.   Z. Duxbury *et al.*, Induced proximity of a TIR signaling domain on a plant-mammalian NLR
chimera activates defense in plants. *Proceedings of the National Academy of Sciences* **117**,
18832-18839 (2020).
11.   B. P. M. Ngou, P. Ding, J. D. Jones, Thirty years of resistance: Zig-zag through the plant
immune system. *The Plant Cell* **34**, 1447-1478 (2022).
12.   L. A. Gao *et al.*, Prokaryotic innate immunity through pattern recognition of conserved
viral proteins. *Science* **377**, eabm4096 (2022).
13.   C.-H. Wu, L. Derevnina, S. Kamoun, Receptor networks underpin plant immunity. *Science*
360, 1300-1301 (2018).
14.   L. D. Cunha *et al.*, Inhibition of inflammasome activation by Coxiella burnetii type IV
secretion system effector IcaA. *Nature communications* **6**, 1-13 (2015).
15.   L. Derevnina *et al.*, Plant pathogens convergently evolved to counteract redundant nodes
of an NLR immune receptor network. *PLoS biology* **19**, e3001136 (2021).
16.   H. Yen, N. Sugimoto, T. Tobe, Enteropathogenic Escherichia coli uses NleA to inhibit NLRP3
inflammasome activation. *PLoS pathogens* **11**, e1005121 (2015).
17.   Q. Chai *et al.*, A bacterial phospholipid phosphatase inhibits host pyroptosis by hijacking
ubiquitin. *Science* **378**, eabq0132 (2022).

18.   C. Marchal, H. Pai, S. Kamoun, J. Kourelis, Emerging principles in the design of
716   bioengineered made-to-order plant immune receptors. *Current Opinion in Plant Biology*,
717   102311 (2022).

718   19.   J. Kourelis, T. Sakai, H. Adachi, S. Kamoun, RefPlantNLR is a comprehensive collection of
719   experimentally validated plant disease resistance proteins from the NLR family. *PLoS*
720   *Biology* **19**, e3001124 (2021).

721   20.   H. Adachi, L. Derevnina, S. Kamoun, NLR singletons, pairs, and networks: evolution,
722   assembly, and regulation of the intracellular immunoreceptor circuitry of plants. *Current*
723   *opinion in plant biology* **50**, 121-131 (2019).

724   21.   C.-H. Wu *et al.*, NLR network mediates immunity to diverse plant pathogens. *Proceedings*
725   *of the National Academy of Sciences* **114**, 8113-8118 (2017).

726   22.   C. N. LaRock, B. T. Cookson, The Yersinia virulence effector YopM binds caspase-1 to arrest
727   inflammasome assembly and processing. *Cell host & microbe* **12**, 799-805 (2012).

728   23.   H. Chen *et al.*, The bacterial T6SS effector EvpP prevents NLRP3 inflammasome activation
729   by inhibiting the Ca2+-dependent MAPK-Jnk pathway. *Cell host & microbe* **21**, 47-58
730   (2017).

731   24.   H. S. Karki, S. Abdullah, Y. Chen, D. A. Halterman, Natural genetic diversity in the potato
732   resistance gene RB confers suppression avoidance from Phytophthora effector IPI-O4.
733   *Molecular Plant-Microbe Interactions* **34**, 1048-1056 (2021).

734   25.   H. Adachi *et al.*, An N-terminal motif in NLR immune receptors is functionally conserved
735   across distantly related plant species. *Elife* **8**, e49956 (2019).

736   26.   J. Kourelis *et al.*, The helper NLR immune protein NRC3 mediates the hypersensitive cell
737   death caused by the cell-surface receptor Cf-4. *bioRxiv*, (2021).

738   27.   J. Wang *et al.*, Ligand-triggered allosteric ADP release primes a plant NLR complex. *Science*
739   **364**, eaav5868 (2019).

740   28.   J. Kourelis, C. Marchal, S. Kamoun, NLR immune receptor-nanobody fusions confer plant
741   disease resistance. *bioRxiv*, (2021).

742   29.   E. Weber, C. Engler, R. Gruetzner, S. Werner, S. Marillonnet, A modular cloning system for
743   standardized assembly of multigene constructs. *PloS one* **6**, e16765 (2011).

744   30.   C. Engler *et al.*, A golden gate modular cloning toolbox for plants. *ACS synthetic biology* **3**,
745   839-843 (2014).

746   31.   M. P. Contreras *et al.*, Sensor NLR immune proteins activate oligomerization of their NRC
747   helper. *BioRxiV*, (2022).

748   32.   L. Derevnina *et al.*, Plant pathogens convergently evolved to counteract redundant nodes
749   of an NLR immune receptor network. *PLoS biology* **19**, e3001136 (2021).

750   33.   C. Duggan *et al.*, Dynamic localization of a helper NLR at the plant–pathogen interface
751   underpins pathogen recognition. *Proceedings of the National Academy of Sciences* **118**,
752   (2021).

753   34.   L. Abas, C. Luschnig, Maximum yields of microsomal-type membranes from small amounts
754   of plant material without requiring ultracentrifugation. *Analytical biochemistry* **401**, 217-
755   227 (2010).

756   35.   F. W. Studier, Protein production by auto-induction in high-density shaking cultures.
757   *Protein expression and purification* **41**, 207-234 (2005).

758   36.   J. F. Steele, R. K. Hughes, M. J. Banfield, Structural and biochemical studies of an NB-ARC
759   domain from a plant NLR immune receptor. *PLoS One* **14**, e0221226 (2019).

760   37.   W. Kabsch, xds. *Acta Crystallographica Section D: Biological Crystallography* **66**, 125-132
761   (2010).

762   38.   G. Winter, xia2: an expert system for macromolecular crystallography data reduction.
763   *Journal of applied crystallography* **43**, 186-190 (2010).

764   39.   P. R. Evans, G. N. Murshudov, How good are my data and what is the resolution? *Acta*
765   *Crystallographica Section D: Biological Crystallography* **69**, 1204-1214 (2013).

766   40.   M. Winn, An overview of the CCP4 project in protein crystallography: an example of a
767   collaborative project. *Journal of synchrotron radiation* **10**, 23-25 (2003).

768   41.   R. Evans *et al.*, Protein complex prediction with AlphaFold-Multimer. *bioRxiv*,
769   2021.2010.2004.463034 (2022).

770   42.   M. Mirdita *et al.*, ColabFold: making protein folding accessible to all. *Nature Methods*, 1-
771   4 (2022).

772   43.   A. J. McCoy *et al.*, Phaser crystallographic software. *Journal of applied crystallography* **40**,
773   658-674 (2007).

774   44.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
775   *Crystallographica Section D: Biological Crystallography* **66**, 486-501 (2010).

776   45.   G. N. Murshudov *et al.*, REFMAC5 for the refinement of macromolecular crystal
777   structures. *Acta Crystallographica Section D: Biological Crystallography* **67**, 355-367
778   (2011).

779   46.   R. A. Nicholls, M. Fischer, S. McNicholas, G. N. Murshudov, Conformation-independent
780   structural comparison of macromolecules with ProSMART. *Acta Crystallographica Section*
781   *D: Biological Crystallography* **70**, 2487-2499 (2014).

782   47.   E. F. Pettersen *et al.*, UCSF ChimeraX: Structure visualization for researchers, educators,
783   and developers. *Protein Science* **30**, 70-82 (2021).

784   48.   The PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC.

785   49.   M. E. Segretin *et al.*, Single amino acid mutations in the potato immune receptor R3a
786   expand response to Phytophthora effectors. *Molecular Plant-Microbe Interactions* **27**,
787   624-637 (2014).

788   50.   D. MacLean, Besthr. *Zenodo*, (2019).

789   51.   I. W. Davis *et al.*, MolProbity: all-atom contacts and structure validation for proteins and
790   nucleic acids. *Nucleic acids research* **35**, W375-W383 (2007).

791