The Structural Layers of the Porcine Iris Exhibit Inherently Different Biomechanical Properties

Royston K. Y. Tan[1,2,3], Satish K. Panda[1,4], Fabian A. Braeu[1,2,3,5], Arumugam R. Muralidharan[2,3], Monisha E. Nongpiur[2,3], Anita S. Y. Chan[2,3], Tin Aung[2,3,6], Raymond P. Najjar[2,3,6,7], Michaël J.A. Girard[1,2,3,8]

1. Ophthalmic Engineering & Innovation Laboratory, Singapore Eye Research Institute, Singapore National Eye Centre, Singapore
2. Singapore Eye Research Institute, Singapore National Eye Centre, Singapore
3. Duke-NUS Medical School, Singapore
4. Department of Mechanical Engineering, Indian Institute of Technology, Bhubaneswar, India
5. Critical Analytics for Manufacturing Personalized-Medicine, Singapore-MIT Alliance for Research and Technology, Singapore
6. Department of Ophthalmology, Yong Loo Lin School of Medicine, National University of Singapore and National University Health System, Singapore
7. Centre for Innovation & Precision Eye Health, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
8. Institute of Molecular and Clinical Ophthalmology, Basel, Switzerland

Keywords: Permeability, stress relaxation, angle closure, iris stroma, dilator muscle

Word Count: 4429 (Manuscript text)
250 (Abstract text)

Tables: 1

Figures: 4

Corresponding Author: Michaël J.A. Girard
Ophthalmic Engineering & Innovation Laboratory
Singapore Eye Research Institute
The Academia
20 College Road, Discovery Tower Level 6
Singapore 169856
mgirard@ophthalmic.engineering

The authors have no proprietary or commercial interest in any materials discussed in this article.

Abstract

Purpose: To isolate the structural components of the *ex vivo* porcine iris tissue and to

determine their biomechanical properties.

Methods: The porcine stroma and dilator tissues were separated, and their dimensions were

assessed using optical coherence tomography (OCT). The stroma underwent flow test ($n =$

32) to evaluate for permeability using Darcy's Law (ΔP = 2000 Pa, A = 0.0391 mm^2), and both

tissues underwent stress relaxation experiments (ε = 0.5 with initial ramp of $\delta\varepsilon$ = 0.1) to

evaluate for their viscoelastic behaviours (n = 28). Viscoelasticity was characterised by the

parameters β (half width of the Gaussian distribution), τ_m (mean relaxation time constant), E_0

(instantaneous modulus) and E_∞ (equilibrium modulus).

Results: For the stroma, the hydraulic permeability was 9.49 ± 3.05 × 10^{-6} mm^2/Pa·s, and the

viscoelastic parameters were β = 2.50 ± 1.40, and τ_m = 7.43 ± 4.96 s, with the two moduli

calculated to be E_0 = 14.14 ± 6.44 kPa and E_∞ = 6.08 ± 2.74 kPa. For the dilator tissue, the

viscoelastic parameters were β = 2.06 ± 1.33 and τ_m = 1.28 ± 1.27 s, with the two moduli

calculated to be E_0 = 9.16 ± 3.03 kPa and E_∞ = 5.54 ± 1.98 kPa.

Conclusion: We have established a new protocol to evaluate the biomechanical properties

of the structural layers of the iris. Overall, the stroma was permeable and exhibited smaller

moduli than those of the dilator muscle. An improved characterisation of iris biomechanics

may form the basis to further our understanding of angle closure glaucoma.

Introduction

68   Research in angle closure glaucoma explores a myriad of possible risk factors and

69 characteristics that are associated with the disease, ranging from genetic studies[1-5] to

70 anatomical measurements[6-10] and computational studies[11-15]. Being able to predict whether a

71 patient will be afflicted with angle closure is paramount, so that preventive measures can be

72 undertaken[16, 17]. Similarly, clinical examinations can identify patients with a higher risk of

73 developing angle closure so that laser iridotomy can be done to widen the drainage angle,

74 lowering the chance of glaucoma development in many instances.

75   Whilst clinical examinations are crucial to aid early angle closure detection[18, 19], the

76 main pathology of angle closure disease is apposition of the peripheral iris to the trabecular

77 meshwork impeding aqueous outflow. It is believed that dynamic movement of the iris,

78 primarily the way the iris dilates, plays a role in angle closure pathogenesis[20-22]. Modern

79 technology can also play its part to study the iris using computational simulations in the form

80 of biomechanical analyses. An example would be to input ocular parameters from a patient's

81 optical coherence tomography (OCT) scan to evaluate how hypothetical changes in iris

82 stiffness could influence angle closure development. These biomechanical models could

83 evaluate the physics and predict the behaviour of the eye in any prescribed scenarios that

84 would otherwise be difficult to analyse *in vivo*.

85   For example, Wang et al.[23] explored aqueous humour dynamics to investigate its

86 influence on the iris and pupillary block in angle closure. This was done using a physical model

87 of the anterior chamber and tracer particles together with computational simulations. Huang

88 et al.[24] and Heys et al.[25] modelled the iris-lens interactions to determine the influence of

89 different morphological parameters (i.e. iris thickness, pupil size, anterior chamber depth, lens

90 diameter, etc.) affecting pupillary block that would otherwise be impossible to determine

91 experimentally. Models could also be invaluable to interpret clinical data; from dynamic OCT

92 images, one could extract parameters such as iris stiffness and permeability[11, 12] in addition to

more traditional morphologic parameters in order to better describe the structural and

biomechanical phenotype of angle closure.

However, current computational models[26-29] of the iris are still limited in describing the

physiological behaviour *in vivo*, particularly outflow of aqueous during pupil constriction and

dilation. The inability to model volume changes of the iris severely limits research capabilities

in angle closure, since the primary mechanism is pupillary block by the iris tissue. Our group

had also previously attempted to describe the iris as a biphasic tissue; with a permeable

stroma to allow for fluid exchange, and a non-linear hyperelastic model to describe both the

solid phases of the stroma and the iris muscles[11, 30, 31] However, to date, no attempts were

made to isolate these tissues experimentally to describe their individual biomechanical

behaviours, partly due to the difficulty in separating them *ex vivo*.

The aim of this study was to characterise the biomechanics of the main structural

components of the iris (stroma and dilator muscle) experimentally. Specifically, we aimed to

determine (1) the permeability of the iris stroma using a refined experimental set-up, and (2)

the viscoelastic properties of both the stroma and dilator muscle through stress relaxation

testing.

Methods

For this study, we isolated two of the primary load bearing components of the iris: the

stroma and the dilator muscle. The iris stroma is composed of a sponge-like meshwork[32],

porous in the middle to allow fluid movements within the anterior volume. It is also composed

of fibroblasts and melanocytes that form a cellular matrix, along with vasculature, neural

elements and a connective tissue extracellular matrix[33]. We isolated the stroma tissue to

determine its fluid permeability, as well as the viscoelastic properties of the solid matrix.

On the other hand, the dilator muscle is made of smooth muscle cells, arranged in

bundles separated by tissue septae. These interdigital systems of muscle fibres are innervated

from autonomic nerve fibres within these tissue septae, communicating through other muscle

fibres through large gap junctions. The surrounding cytoplasm contains a uniform distribution

of myofilaments that contribute to the mechanical strength of the muscle. As the dilator muscle

is impermeable due to the presence of tissue septae and gap junctions[33-35], only its viscoelastic

properties would be investigated. As the iris posterior pigmented epithelium is a component of

the blood aqueous barrier, it was removed to study the stromal permeability.

Tissue separation

Sixty freshly enucleated porcine eyes (were purchased from a local porcine abattoir

(approximately aged 6 months, Singapore Food Industries Pte Ltd., Singapore). The fresh

porcine eyes were refrigerated after excision, chilled in an ice box during the transport to the

laboratory. Experiments were performed at room temperature of 24°C and completed within

24 hours post-mortem.

The procedure to separate the stroma and dilator tissues was similar to our previous

study, which had been validated through histological analysis[31]. We made further

improvements to the technique by implementing the following refinements in the procedure.

First, for each eye, the entire iris was isolated from the porcine globe, and the iris pigmented

epithelium gently scraped off. Next, a discontinuous cut was made to the iris. Using a mixture

of tying forceps (2-500-2E, Duckworth & Kent, Hertfordshire, UK), jewellers forceps (2-900E,

Duckworth & Kent, Hertfordshire, UK) and Chihara conjunctival forceps (2-500-4E, Duckworth

& Kent, Hertfordshire, UK), the dilator muscle was gently lifted and peeled away from the

stroma inward radially up to the sphincter, where the piece of tissue was cut off using a pair

of vannas scissors (1-111, Duckworth & Kent, Hertfordshire, UK). When isolating the stroma,

the dilator was removed in pieces to retain as much stroma tissue as possible, whereas when

isolating the dilator muscle, care was taken to ensure no holes were present in the isolated

sample, even though small amounts of stroma tissue may still be present. This method

ensured proper tissue isolation to preserve the structural integrity of each sample and little to

no impact on the measurement of biomechanical properties. After tissue isolation, the samples

were stored in phosphate buffer saline (PBS) without calcium or magnesium to limit any

potential post-mortem muscle activity[36], diluted from 10× stock solution (17-517Q, Lonza

Pharma & Biotech, Maryland, USA).

Stroma permeability study

149   Isolated stroma samples (n = 32) underwent a flow experiment to assess their

150   permeability. The principle of the permeability experiment was similar to that described in our

151   previous study, whereby homogenisation of the Navier-Stokes equation[37, 38] will yield Darcy's

152   law[39, 40] :

$$Q = \frac{\kappa A \Delta P}{\mu \Delta x} \ (\text{mm}^3/\text{s}) \tag{1}$$

where Q is the flow rate (mm³/s), κ is the permeability (mm²), A is the surface area of flow

(mm²), ΔP is the pressure drop across the tissue in the direction of flow (Pa), μ is the dynamic

viscosity of the fluid (Pa·s) and Δx is the thickness of the tissue (mm). Our measurement

parameter is the hydraulic permeability, used in similar biphasic biological tissue

experimentation such as the meniscus[41]. It is defined by the permeability of the tissue divided

by the viscosity of the fluid (here: phosphate buffered saline or PBS):

$$\text{Hydraulic Permeability} = \frac{\kappa}{\mu} = \frac{Q \Delta x}{A \Delta P} \ (\text{mm}^2/[\text{Pa} \cdot \text{s}]) \tag{2}$$

Using Darcy's law, it was possible to obtain the hydraulic permeability of the iris stroma

by setting a specific pressure difference and measuring the resulting flow rate. We designed

an experimental setup consisting of a PBS column (30 mL syringe reservoir), itself connected

by a tube (internal diameter (ID) = 1.00 mm) to a custom 3D-printed holder where the stroma

tissue was positioned (**Figure 1**). The cross-sectional area for flow was measured using

optical coherence tomography (OCT) and estimated as an ellipse (**Figure 1B and 2A**). The

syringe reservoir was sufficiently large to prevent any noticeable drops in fluid level (< 1 mm)

from direct flow or evaporation. The vertical height between the stroma tissue and the fluid

level was approximately 204 mm to provide a pressure difference of approximately 2,000 Pa,

or 15 mmHg. Outflow from the holder was through the large tube (ID = 1.00 mm), followed by

a much smaller tube (ID = 0.397 mm) on the horizontal plane for the measurement of flow rate

(**Figure 1C**).

The tubes of different inner diameters were selected for the following reasons: 1) The

larger tube (ID = 1.00 mm) was sufficiently large that the tube resistance was negligible; 2)

The smaller tube allowed for accurate measurement of flow rate, which was measured by the

time taken for 1.237 mm^3 of fluid to flow through the stroma tissue (10 mm of the tube). A

smaller tube also created outflow resistance (positive pressure), which was taken into account

during the calculations. The hydraulic permeability values were reported as the mean ±

standard deviation, with outliers excluded beyond 3 standard deviations.

To ensure accurate and consistent results, we calibrated the experimental setup at the

start of each experiment by measuring the flow rate in the absence of tissue resistance. The

measured flow rate closely matched theoretical calculations using Poiseuille's equation, with

a margin of error of less than 5%. Before testing each sample, we thoroughly flushed and

inspected the entire system for bubbles and leaks to ensure the reliability of our results.

Stress relaxation study

Iris stroma ($n = 28$) and dilator muscles ($n = 28$) underwent stress relaxation tests (i.e.,

stress response over time over a constant strain) using a uniaxial testing machine (5542A,

Instron Corp, Norwood, MA, USA). The tested length (in the direction of the uniaxial stretch)

was 1.5 mm for all stroma and dilator muscle samples, with the cross-sectional areas

evaluated using OCT scans performed for each sample (**Figure 2C**) prior to the stress

relaxation tests. As the iris tissues typically exhibit extremely large strains over several

seconds (up to the second order of magnitude in percentage), we decided to perform the

stress relaxation experiment at 50% strain. The samples were extended with a strain rate of

10% per second for 5.0 seconds and held at constant strain (ε_0) for the next 300 seconds

(**Figure 3**). All samples were clamped using sandpaper to prevent slipping (**Figure 3C**).

Preconditioning was performed for 25 cycles from 0% to a maximum of 25% strain at a rate of

1% per second. Overall, samples were found to exhibit little to no hysteresis.

To quantify the relaxation behaviour of the iris tissues, we used the linear viscoelastic

theory initially described by Fung[42] and the same mathematical model as in our previous study

by Girard et al., with the stress relaxation response[43]:

$$\sigma(t) = \frac{\sigma_0}{1+c}\left(1 + \frac{c}{\beta\sqrt{\pi}}\int_{-\infty}^{+\infty}\exp\left(-\frac{z^2}{\beta^2} - \frac{t}{\tau_m 10^z}\right)dz\right) \tag{3}$$

where c is the relaxation ratio, β is the half width of the Gaussian distribution[44], $z = \log(\tau/\tau_m)$,

τ is the relaxation time[45] and τ_m is the mean relaxation time constant. σ_0 is defined as the

peak stress and was directly obtained from the experimental data.

Using MATLAB (Version 2020b, Mathworks, Inc., Natick, MA, USA), we reduced noise

by signal averaging and performed curve fitting of the experimental data to **Equation 3** using

the differential evolution[46] (DE) algorithm developed by Girard et al.[43] to estimate the three

model parameters: c, β and τ_m. For the DE algorithm, we utilised the following parameter

settings: a population size (NP) of 15, a mutation factor (F) of 0.8, a crossover probability (CR)

of 0.9. The range of the variables were defined (e.g., 0 < c, β, τ_m < 10) within the DE algorithm,

which represented values within the physiological range[43]. The unique global minimum

determined exhibited an excellent fit, with a cost function (defined by square root of the sum

of the squares) of less than 10^{-3} (**Figure 4**). The two elastic moduli, E_0 and E_∞ was determined

by:

$$E_0 = \frac{\sigma_0}{\varepsilon_0} \tag{4}$$

$$E_\infty = \frac{E_0}{1+c} \tag{5}$$

212   *OCT tissue measurements*

213   Images obtained from this study were performed using OCT (RTVue-100, Optovue Inc, Fremont, CA, USA). The stroma and dilator tissue samples were placed on a glass slide with minimal liquid to reduce influence from liquid refraction. The acquisition mode was the 8-line radial scan (each cross-sectional scan was 22.5° apart). The images were processed using Photoshop (v2022, Adobe Inc, CA, USA) and MATLAB. Thickness and cross-sectional area were evaluated using the scale from the OCT machine provided in the images (**Figure 2**).

Stroma permeability study: Three random images were selected per sample to determine the tissue thickness after the experiment was completed for a better reflection of the tissue state during the experiment. An area of 45 × 100 pixels was selected for each image and manually converted to binary black and white using photoshop, with the average of the 3 images computed for the tissue thickness (**Figure 3A**).

Stress relaxation study: Here, tissue thickness was measured prior the experiment. It was important to position the tissue uniformly so that the horizontal width of the tissue would be accurately reflected and measured by the cross-sectional images. The entire cross-section was manually converted to binary black and white using photoshop, with the width and cross-section computed for analysis (**Figure 3B**).

Statistical Analysis

The Wilcoxon rank sum test (two-tailed) was employed to compare the five material parameters (c, β, τ_m, E_0 and E_∞) between the dilator muscle and stroma groups ($n = 28$), to evaluate whether the tissues were inherently different.

Results

Stroma permeability study

The 3D tissue holder area was evaluated using OCT and determined to be 0.0391 mm^2. The average thickness of the stroma tissue samples was 0.152 ± 0.0259 mm. The PBS flow through the stroma samples was found to be slow and laminar, with a Reynold's number

< 1, thus satisfying the Darcy's law approximation. Specifically, flow through the experimental

setup was 0.00500 ± 0.00191 mm^3/s, exerting minimal outlet pressure (7.65 ± 2.93 Pa). The

hydraulic permeability of the porcine iris stroma was then calculated as 9.49 ± 3.05 × 10^{-6}

mm^2/Pa·s, with a total of 31 samples evaluated, with 1 outlier excluded from the analysis.

Stress relaxation study

Dilator muscle tissue. The average width, thickness and cross-sectional area of the

dilator tissue samples were 3.78 ± 0.74 mm, 0.192 ± 0.028 mm and 0.733 ± 0.183 mm^2

respectively. The biomechanical properties of the dilator muscle, as determined by the

differential evolution algorithm, were c = 1.41 ± 0.60, β = 2.50 ± 1.40, and τ_m = 7.43 ± 4.96 s,

with the two moduli calculated to be E_0 = 14.14 ± 6.44 kPa and E_∞ = 6.08 ± 2.74 kPa.

Stroma tissue. The average width, thickness and cross-sectional area of the stroma

tissue samples were 3.71 ± 0.70 mm, 0.220 ± 0.04 mm and 0.830 ± 0.263 mm^2 respectively.

The biomechanical properties of the stroma, as determined by the differential evolution

algorithm, were c = 0.707 ± 0.38, β = 2.06 ± 1.33, and τ_m = 1.28 ± 1.27 s, with the two moduli

calculated to be E_0 = 9.16 ± 3.03 kPa and E_∞ = 5.54 ± 1.98 kPa.

Overall, the dilator muscle tissue exhibited greater E_0 (p < 0.001) compared to the

stroma tissue but did not show significant difference in E_∞ (p = 0.64). The dilator muscle tissue

also showed greater c and τ_m (p < 0.001) but no significant difference in β (p = 0.20).

Discussion

In this study, we identified the stroma and dilator muscles as two structural tissues

influencing iris biomechanical behaviour. To the best of our knowledge, the biomechanics of

such tissue layers (when isolated) had not been studied, and we proposed herein a

methodology for doing so. Overall, we found that (1) The iris stroma was highly permeable (as

opposed to the dilator); (2) the instantaneous modulus (but not the equilibrium modulus) of the

dilator muscle was significantly higher than that of the stroma; and that (3) the stroma exhibited

a significantly shorter mean relaxation time constant as compared to the dilator muscle. Our

work may ultimately help us identify an 'healthy' range for iris biomechanical properties in

human, any deviations from which may be associated with the development of angle closure.

The dilator muscle is stiffer than the stroma tissue in a healthy iris

In this study, we found that the instantaneous modulus of the dilator muscle was

significantly higher than that of the stroma tissue – but this was not true for the equilibrium

modulus. Such a difference could potentially be explained by the different fibres that constitute

both tissues, including stiffer smooth muscle fibres for the dilator tissue, but a sparse collagen

fibre meshwork permeable to aqueous humour for the stroma, thus allowing for greater

compliance. This softer stroma layer is essential for iris accommodation, since the contraction

forces (for dilation) originate from a much thinner muscle layer. Recently, it was reported that

glaucomatous eyes, including eyes with primary open angle glaucoma and primary angle

closure glaucoma, generally display a reduction in the pupillary light response amplitudes[47-49].

Although such alterations in the pupillary light response can be attributed to ganglion cell loss

or dysfunction in the disease, we hypothesise that decreased muscle innervation and changes

in biomechanical behaviour such as increased stroma stiffness, are additional underlying

biological changes that may contribute to pupillometric alterations in response to light for

patients with glaucoma.

The iris accommodates movement rapidly to reach biomechanical equilibrium

In this study, we found that the mean relaxation time constants for both the dilator

muscle (7.43 ± 4.96 s) and stroma (1.28 ± 1.27 s) were relatively small. These features may

reflect the highly dynamic physiological properties of the iris that is constantly changing in

shape in response to the accommodation reflex or the lighting environment. This is in contrast

to other ocular tissues such as the corneo-scleral shell that exhibits considerably smaller

changes in tissue stretch. The viscoelastic behaviour of stiffer, load bearing structural tissues

(in terms of intraocular pressure) such as the sclera exhibits high moduli and greater mean

relaxation time constant, whereas the small amount of muscle fibres orchestrating fine and

rapid pupil dilator changes result in much smaller moduli (E_∞ = 3,300 kPa for rabbit sclera vs.

5.54 kPa for iris stroma) and short mean relaxation time constants (τ_m = > 10s for rabbit sclera

vs. < 10s for iris tissues)[43].

The results also show a shorter mean relaxation time constant for the stroma tissue,

with it seemingly behaving close to a linear elastic material (**Figure 4**). This could be due to

its porous nature and composition; most of the stroma is composed short collagen fibrils

meshwork with fibroblasts and melanosomes[33, 34]. The stroma matrix deforms rapidly in both

extension and compression as aqueous humour gets quickly displaced through its pores,

reaching structural equilibrium within seconds. Conversely, the dilator muscle takes longer to

reach biomechanical equilibrium, the simple bundles of smooth muscle cells[33] containing actin

and myosin filaments passively extend irreversibly, until the filaments are activated to overlap

and contract during pupil constriction.

Changes in iris tissue properties can affect iris curvature and induce angle closure

In angle closure, the behaviour and shape of the iris in the mydriatic state affects the

chamber angle. Patients with angle closure glaucoma were found to have anterior convex-to-

convex (in dark and light) iris configuration[50], possibly signalling a biomechanical mismatch

between the iris stroma and muscle tissues. There are several potential mechanisms that can

result in an anteriorly convex iris. Firstly, if the iris stroma undergoes stiffening, the tissue may

bow upward to counteract compression by the dilator muscle. Secondly, changes in the

composition of the iris stroma could lead to an increased matrix-to-aqueous ratio, reducing

compressibility and resulting in anterior bowing. Thirdly, a decrease in stromal permeability

could impede the outflow of aqueous humour from the iris[32], leading to volume retention[51] and

increased iris thickness[52], ultimately causing convexity. This change would likely be attributed

to morphological changes at the anterior boundary layer, where the density of the tissue is the

primary limiting factor restricting aqueous fluid migration. Finally, a mismatch in biomechanical

behaviour between the stroma and surrounding muscles, such as an increase in stroma mean

relaxation time constant, could affect how the iris behaves in unison. In the case of pupil

dilation, an anteriorly convex iris could result initially before the stresses are evenly distributed,

only becoming less convex over a longer period of time.

319   The lack of iris features such as crypts and furrows[53] had been linked to the

development of angle closure. The presence of crypts, which are diamond shape lacunae

causing discontinuities on the anterior boundary layer, could facilitate aqueous humour

movement across the stroma tissue and prevent the iris from bulging and bowing during pupil

dilation. The presence of lower number of crypts had been found to be proportional to the

incidence of angle closure in patients[54]. Furrows, which are circumferential creases where the

iris folds at the periphery during pupil dilation, could also be associated with the prevalence of

angle closure. With longer furrow lengths, it was found that the irides were more likely to have

greater volume and less aqueous loss during pupil dilation[55]. These iris features provide not

only greater insight into their associations with angle closure, but also possible reasons for the

deviations from mean iris behaviour that could be quantified biomechanically.

Our work allows us to better describe the passive extensive behaviour of the ex vivo

porcine iris

332   This study represented the iris as a structural bilayer; the anterior stroma as a solid-

liquid biphasic tissue with solid viscoelastic properties and fluid permeability, and the posterior

dilator muscle as a solid with viscoelastic properties. We believe this captures the unique

behaviour of the iris, which is unlike many other tissues in the body. The iris experiences large

deformations created by the posteriorly positioned muscles, coupled with interaction with the

surrounding fluid medium from the partially porous and anteriorly positioned stroma. By

isolating the individual primary load bearing layers, it was possible to accurately determine

their biomechanical behaviour. Previous research into the iris took different approaches:

Whitcomb et al. performed nano-indentation of the porcine iris anterior and posterior segments

to determine overall iris stiffness[28], while Narayanaswamy et al. used atomic force

microscopy[56]. There are some anatomical and biomechanical points to consider for these

approaches. First, for the anterior indentation, the small level of indentation in a porous stroma

could effectively be flexing fibres between the gaps of the tissue, undermining the true tissue

stiffness. Second, when the antagonistic muscles contract, the iris is either in a compressive

or extensive state in the radial direction, orthogonal to the direction of indentation. The

assumption of an isotropic tissue may not be biomechanically accurate to represent the

radially oriented dilator muscles, and it may be critical to test the tissues according to their

preferred fibre orientations as we had performed herein.

Whilst the method for measuring biomechanical properties is only possible *ex vivo*, we

believe that capturing a better representation of the iris behaviour is invaluable for clinical

research. First, we could utilise this technique to investigate the tissue biomechanical

differences between normal and angle closure iridectomy samples. Next, through evaluation

of miosis and mydriasis OCT data, we believe that iris biomechanical differences in angle

closure can be validated *in vivo* using computational methods such as inverse finite element[11]

and AI-driven approaches to access tissue biomechanics. From a clinical perspective, our

goal is to identify differences in biomechanical data between normal and disease subjects by

evaluating complex OCT data, and the understanding of differences can be combined with AI

to improve disease prognosis.

Limitations

In this study, several limitations warrant further discussion. First, a more complete

biomechanical breakdown of the iris tissue properties should also include the sphincter muscle.

The omission of the sphincter muscle was due to the difficulty in isolating this circular tissue.

However, it is theoretically possible to equate the passive biomechanical behaviour of both

the dilator and sphincter muscle since they are antagonistic tissues. The differences in speeds

of pupil constriction and dilation suggest possible small tissue differences, perhaps due to

different nerve innervations, but microscopic studies have shown that these muscles are

typical smooth muscles[33], and hence are likely to exhibit similar biomechanical properties.

Moreover, a better approach would involve incorporating the biphasic poroviscoelastic (BPVE)

constitutive model, which accounts for both viscoelasticity and permeability. In our study, we

focused on evaluating the solid linear viscoelastic theory for modelling the iris dilator muscle

and the iris stroma. Additionally, we separately assessed the hydraulic permeability of the iris

stroma due to our previous study's[31] findings suggesting that the high permeability had little

impact on tissue behaviour (Appendix A). This corroborates the results investigated by Greiner

et al.[57], who found that the fluid nominal stresses were negligible for permeability values above

10^{-6} mm^2/Pa·s. However, it is advisable for future investigations to consider employing the

solid linear viscoelastic theory for the muscle tissue and the BPVE theory for the stroma tissue,

especially when considering other loading scenarios.

Second, a limitation of experimental *ex vivo* biomechanical determination of tissues is

the inability to determine active muscle properties. Changes in pupil consists of at least three

tissue changes; pupil constriction requires active sphincter muscle contraction, dilator muscle

extension (relaxation) and stroma extension, whilst pupil dilation requires active dilator muscle

contraction, sphincter muscle extension (relaxation) and stroma compression. Whilst it would

be a logistical challenge to determine active muscle forces by keeping the tissues alive, such

forces might be able to be estimated with *in vivo* imaging such as OCT.

Third, the stroma undergoes both compressive and tensile stresses with pupillary

changes, however, on our study we only performed tensile experiments. In addition,

permeability could be both affected by tensile and compressive stress, as is true for other soft

tissues such as cartilage[58, 59]. Novel mechanical test equipment might need to be designed to

assess such properties.

Fourth, we had loaded the tissue at a rate of 10% per second for 5 seconds. Our

original tissue length measured approximately 4 – 4.5 mm, trimmed to 1.5 mm. Hence our

loading rate corresponds to a constriction rate of approximately 0.4 – 0.45 mm/s. *In vivo*

constriction rates[60, 61] range can be as slow as 0.5 mm/s and increase up to 7.5 mm/s

depending on light intensity, with the maximum rate achieved at 550 cd/m^2. We modelled our

loading rate based on physiological dynamic accommodation constriction rates[62], which also

corresponded to the lower limits of dark-to-light constriction rates. It would be important for

future studies to investigate the upper limits of constriction rates for a more comprehensive

analysis of the iris' biomechanical behaviour.

Conclusion

Determining the biomechanical properties of the individual iris layers may be

invaluable in uncovering how and why the iris deforms in the anterior chamber in normal and

eyes with angle closure. This porcine study established a protocol to be replicated on human

iris tissues, by isolating the individual components for analysis. In learning more about iris

biomechanics, we hope to establish a better understanding of the pathogenesis of angle

closure glaucoma.

Acknowledgments

The authors thank the donors of the National Glaucoma Research, a program of the

BrightFocus Foundation, for support of this research (G2021010S [MJAG]); the NMRC-LCG

grant 'TAckling & Reducing Glaucoma Blindness with Emerging Technologies (TARGET)',

award ID: MOH-OFLCG21jun-0003 [MJAG]; and the "Retinal Analytics through Machine

learning aiding Physics (RAMP)" project that is supported by the National Research

Foundation (NRF2019-THE002-0006 [MJAG]).

References

1.   Wang J, Yusufu M, Khor CC, Aung T, Wang N. The genetics of angle closure glaucoma. *Experimental eye research* 2019;189:107835.

2.   Nongpiur ME, Khor CC, Jia H, et al. ABCC5, a gene that influences the anterior chamber depth, is associated with primary angle closure glaucoma. *PLoS genetics* 2014;10:e1004089.

3.   Vithana EN, Khor C-C, Qiao C, et al. Genome-wide association analyses identify three new susceptibility loci for primary angle closure glaucoma. *Nature genetics* 2012;44:1142-1146.

4.   Suri F, Yazdani S, Chapi M, et al. COL18A1 is a candidate eye iridocorneal angle-closure gene in humans. *Human molecular genetics* 2018;27:3772-3786.

5.   Ahram D, Alward W, Kuehn M. The genetic mechanisms of primary angle closure glaucoma. *Eye* 2015;29:1251-1259.

6.   Congdon NG, Friedman DS. Angle-closure glaucoma: impact, etiology, diagnosis, and treatment. *Current opinion in ophthalmology* 2003;14:70-73.

7.   Cheung CY, Zheng C, Ho C-L, et al. Novel anterior-chamber angle measurements by high-definition optical coherence tomography using the Schwalbe line as the landmark. *British Journal of Ophthalmology* 2011;95:955-959.

8.   Wang B, Narayanaswamy A, Amerasinghe N, et al. Increased iris thickness and association with primary angle closure glaucoma. *British Journal of Ophthalmology* 2011;95:46-50.

9.   Sihota R, Lakshmaiah N, Agarwal H, Pandey R, Titiyal J. Ocular parameters in the subgroups of angle closure glaucoma. *Clinical & experimental ophthalmology* 2000;28:253-258.

10.   Zhou M, Wang W, Ding X, et al. Choroidal thickness in fellow eyes of patients with acute primary angle-closure measured by enhanced depth imaging spectral-domain optical coherence tomography. *Investigative Ophthalmology & Visual Science* 2013;54:1971-1978.

11.   Panda SK, Tan RK, Tun TA, et al. Changes in Iris Stiffness and Permeability in Primary Angle Closure Glaucoma. *Investigative Ophthalmology & Visual Science* 2021;62:29-29.

12.   Pant AD, Gogte P, Pathak-Ray V, Dorairaj SK, Amini R. Increased iris stiffness in patients with a history of angle-closure glaucoma: an image-based inverse modeling analysis. *Investigative ophthalmology & visual science* 2018;59:4134-4142.

13.   Thomas VS, Salinas SD, Pant AD, Dorairaj SK, Amini R. Biomechanical assessment of the Iris in relation to angle-closure glaucoma: a multi-scale computational approach. *International Symposium on Computer Methods in Biomechanics and Biomedical Engineering*: Springer; 2019:470-482.

14.   Safa BN, Bahrani Fard MR, Ethier CR. In vivo biomechanical assessment of iridial deformations and muscle contractions in human eyes. *Journal of the Royal Society Interface* 2022;19:20220108.

15.   Safa BN, Wong CA, Ha J, Ethier CR. Glaucoma and biomechanics. *Current Opinion in Ophthalmology* 2022;33:80-90.

16.   Ang LP, Aung T, Chew PT. Acute primary angle closure in an Asian population: long-term outcome of the fellow eye after prophylactic laser peripheral iridotomy. *Ophthalmology* 2000;107:2092-2096.

17.   Playfair TJ, Watson PG. Management of acute primary angle-closure glaucoma:
a long-term follow-up of the results of peripheral iridectomy used as an initial
procedure. *British Journal of Ophthalmology* 1979;63:17-22.
18.   Tarongoy P, Ho CL, Walton DS. Angle-closure glaucoma: the role of the lens
in the pathogenesis, prevention, and treatment. *Survey of ophthalmology*
2009;54:211-225.
19.   Wright C, Tawfik MA, Waisbourd M, Katz LJ. Primary angle‐closure glaucoma:
an update. *Acta ophthalmologica* 2016;94:217-225.
20.   He M, Lu Y, Liu X, Ye T, Foster PJ. Histologic changes of the iris in the
development of angle closure in Chinese eyes. *Journal of glaucoma* 2008;17:386-392.
21.   Hao L, Hu Y, Xu Y, et al. Dynamic analysis of iris changes and a deep learning
system for automated angle-closure classification based on AS-OCT videos. *Eye and
Vision* 2022;9:1-10.
22.   Quigley HA, Silver DM, Friedman DS, et al. Iris cross-sectional area decreases
with pupil dilation and its dynamic behavior is a risk factor in angle closure. *Journal of
glaucoma* 2009;18:173-179.
23.   Wang W, Qian X, Song H, Zhang M, Liu Z. Fluid and structure coupling analysis
of the interaction between aqueous humor and iris. *Biomedical Engineering Online*
2016;15:569-586.
24.   Huang EC, Barocas VH. Active iris mechanics and pupillary block: steady-state
analysis and comparison with anatomical risk factors. *Annals of biomedical
engineering* 2004;32:1276-1285.
25.   Heys JJ, Barocas VH, Taravella MJ. Modeling passive mechanical interaction
between aqueous humor and iris. *J Biomech Eng* 2001;123:540-547.
26.   Zhang K, Qian X, Mei X, Liu Z. An inverse method to determine the mechanical
properties of the iris in vivo. *Biomedical engineering online* 2014;13:1.
27.   Heys J, Barocas V. Mechanical characterization of the bovine iris. *Journal of
biomechanics* 1999;32:999-1003.
28.   Whitcomb JE, Amini R, Simha NK, Barocas VH. Anterior–posterior asymmetry
in iris mechanics measured by indentation. *Experimental eye research* 2011;93:475-
481.
29.   Whitcomb JE, Barnett VA, Olsen TW, Barocas VH. Ex vivo porcine iris stiffening
due to drug stimulation. *Experimental eye research* 2009;89:456-461.
30.   Tan RK, Wang X, Perera SA, Girard MJ. Numerical stress analysis of the iris
tissue induced by pupil expansion: Comparison of commercial devices. *PloS one*
2018;13:e0194141.
31.   Tan RKY, Wang X, Chan ASY, et al. Permeability of the porcine iris stroma.
Exp Eye Res 2019;181:190-196.
32.   Quigley HA. The iris is a sponge: a cause of angle closure. *Ophthalmology*
2010;117:1-2.
33.   Freddo TF. Ultrastructure of the iris. *Microscopy research and technique*
1996;33:369-389.
34.   Tousimis A, FINE BS. Ultrastructure of the iris: the intercellular stromal
components. *AMA Archives of Ophthalmology* 1959;62:974-976.
35.   Tonjum A, Pedersen O. The permeability of the human ciliary and iridial
epithelium to horseradish peroxidase. An in vitro study. *Acta ophthalmologica*
1977;55:781-788.
36.   Greaser M, Cassens R, Briskey E, Hoekstra W. Post‐mortem changes in
subcellular fractions from normal and pale, soft, exudative porcine muscle. 1. Calcium

accumulation and adenosine triphosphatase activities. *Journal of Food Science* 1969;34:120-124.

37. Chorin AJ. Numerical solution of the Navier-Stokes equations. *Mathematics of computation* 1968;22:745-762.

38. Temam R. *Navier-Stokes equations: theory and numerical analysis*: American Mathematical Soc.; 2001.

39. Keller JB. Darcy's law for flow in porous media and the two-space method. *Nonlinear partial differential equations in engineering and applied sciences* 1980;31:429-443.

40. Whitaker S. Flow in porous media I: A theoretical derivation of Darcy's law. *Transport in porous media* 1986;1:3-25.

41. Périé D, Korda D, Iatridis JC. Confined compression experiments on bovine nucleus pulposus and annulus fibrosus: sensitivity of the experiment in the determination of compressive modulus and hydraulic permeability. *Journal of biomechanics* 2005;38:2164-2171.

42. Fung Y-c. *Biomechanics: mechanical properties of living tissues*: Springer Science & Business Media; 2013.

43. Girard M, Suh J-KF, Hart RT, Burgoyne CF, Downs JC. Effects of storage time on the mechanical properties of rabbit peripapillary sclera after enucleation. *Current eye research* 2007;32:465-470.

44. Olberding JE, Suh JF. A dual optimization method for the material parameter identification of a biphasic poroviscoelastic hydrogel: potential application to hypercompliant soft tissues. *Journal of biomechanics* 2006;39:2468-2475.

45. Nowick AS, Berry BS. Lognormal distribution function for describing anelastic and other relaxation processes I. theory and numerical computations. *IBM Journal of Research and Development* 1961;5:297-311.

46. Storn R. On the usage of differential evolution for function optimization. *Proceedings of north american fuzzy information processing*: Ieee; 1996:519-523.

47. Najjar RP, Rukmini A, Finkelstein MT, et al. Handheld chromatic pupillometry can accurately and rapidly reveal functional loss in glaucoma. *British Journal of Ophthalmology* 2021.

48. Najjar RP, Sharma S, Atalay E, et al. Pupillary responses to full-field chromatic stimuli are reduced in patients with early-stage primary open-angle glaucoma. *Ophthalmology* 2018;125:1362-1371.

49. Rukmini AV, Milea D, Baskaran M, et al. Pupillary responses to high-irradiance blue light correlate with glaucoma severity. *Ophthalmology* 2015;122:1777-1785.

50. Cheung CY-l, Liu S, Weinreb RN, et al. Dynamic analysis of iris configuration with anterior segment optical coherence tomography. *Investigative ophthalmology & visual science* 2010;51:4040-4046.

51. Aptel F, Denis P. Optical coherence tomography quantitative analysis of iris volume changes after pharmacologic mydriasis. *Ophthalmology* 2010;117:3-10.

52. Mak H, Xu G, Leung CK-S. Imaging the iris with swept-source optical coherence tomography: relationship between iris volume and primary angle closure. *Ophthalmology* 2013;120:2517-2524.

53. Pan C-W, Qian Y-X, Li J, Zhong H. Iris surface features and anterior chamber depth in Chinese adolescents. *BMC ophthalmology* 2020;20:1-6.

54. Chua J, Thakku SG, Tun TA, et al. Iris crypts influence dynamic changes of iris volume. *Ophthalmology* 2016;123:2077-2084.

55. Chua J, Thakku SG, Pham TH, et al. Automated detection of iris furrows and their influence on dynamic iris volume change. *Scientific reports* 2017;7:1-8.

56.   Narayanaswamy A, Nai MH, Nongpiur ME, et al. Young's Modulus
Determination of Normal and Glaucomatous Human Iris. *Investigative ophthalmology*
& visual science 2019;60:2690-2695.
57.   Greiner A, Reiter N, Paulsen F, et al. Poro-viscoelastic effects during
biomechanical testing of human brain tissue. *Frontiers in Mechanical Engineering*
2021;7:708350.
58.   Chen X, Zhou Y, Wang L, Santare MH, Wan LQ, Lu XL. Determining tension–
compression nonlinear mechanical properties of articular cartilage from indentation
testing. *Annals of biomedical engineering* 2016;44:1148-1158.
59.   Lai W, Mow VC, Roth V. Effects of nonlinear strain-dependent permeability and
rate of compression on the stress behavior of articular cartilage. 1981.
60.   Ellis C. The pupillary light reflex in normal subjects. *British Journal of*
Ophthalmology 1981;65:754-759.
61.   Ciuffreda KJ, Joshi NR, Truong JQ. Understanding the effects of mild traumatic
brain injury on the pupillary light reflex. *Concussion* 2017;2:CNC36.
62.   Hirota M, Morimoto T, Miyoshi T, Fujikado T. Simultaneous Measurement of
Objective and Subjective Accommodation in Response to Step Stimulation.
Investigative Ophthalmology & Visual Science 2020;61:38-38.

Figure 1. Hydraulic permeability measurement set-up. **A.** Fluid of approximately 2,000 Pa (15

mmHg) flows through a 3D printed holder. **B.** The tissue is placed between over the channel

(brown dotted circle) and clamped between 2 identical 3D printed channels. The area for flow

is approximately 0.0391 mm^2, suitable for samples that are obtained from iridotomy. **C.** Flow

rate through a 0.397 mm diameter tube is measured and recorded.

Figure 2. OCT images were processed to evaluate tissue parameters. The white arrows on

the left depict the direction of the B-scans on the right. **A.** The channel area of the custom 3D-

printed holder for the permeability experiment was estimated as an ellipse using

measurements from OCT. **B.** For the permeability experiment, an area of 45 × 100 pixels was

selected for each image and manually converted to binary black and white, **C.** whereas the

width and area were computed for the stress relaxation experiment.

Figure 3. Stress relaxation experiment was conducted on the iris tissues by **A.** Applying linear

strain rate of 10% per second for 5 seconds and holding constant strain as the tissue relaxes.

B. The same protocol was performed on both the isolated dilator (left) and stroma (right)

tissues. **C.** To prevent slippage, fine sandpaper was used at the clamps of the Instron machine.

Figure 4. Stress relaxation behaviour of the porcine dilator tissue (left) and stroma tissue

(right). The blue line represents the experimental data, and the orange line represents the

fitted data.

Table 1. Viscoelastic parameters of the iris stroma and dilator muscle tissues (mean

value ± standard deviation).

	c	β	τ_m (s)	E_0 (kPa)	E_∞ (kPa)
Dilator muscle	1.41 ± 0.60	2.50 ± 1.40	7.43 ± 4.96	14.14 ± 6.44	6.08 ± 2.74
Stroma tissue	0.707 ± 0.38	2.06 ± 1.33	1.28 ± 1.27	9.16 ± 3.03	5.54 ± 1.98

606