Modelling Spontaneous Firing Activity of the Motor Cortex in a Spiking Neural Network with Random and Local Connectivity

Lysea Haggie [1] ✉, **Thor Besier**[1], **Angus McMorland**[1,2]

[1]Auckland Bioengineering Institute, University of Auckland, Auckland, New Zealand;

[2]Department of Exercise Sciences, University of Auckland, Auckland, New Zealand

✉ **For correspondence:**
lmun373@aucklanduni.ac.nz
(Lysea Haggie)

Present address: Auckland Bioengineering Institute, Level 6, 70 Symonds Street University of Auckland, Auckland 1010, New Zealand

Code availability: Code and data available on GitHub.

Funding: L.H. was supported by funding from Callaghan Innovation. The funders had no role in the template design or decision to publish.

Competing interests: The authors declare no competing interests.

Abstract

Computational models of cortical activity can provide insight into the mechanisms of higher-order processing in the human brain including planning, perception and the control of movement. Activity in the cortex is ongoing even in the absence of sensory input or discernable movements and is thought to be linked to the topology of the underlying cortical circuitry (Ringach, 2009). However, the connectivity and its functional role in the generation of spatio-temporal firing patterns and cortical computations are still vastly unknown. Movement of the body is a key function of the brain, with the motor cortex the main cortical area implicated in the generation of movement. We built a spiking neural network model of the motor cortex which incorporates a laminar structure and circuitry based on a previous cortical model by Potjans and Diesmann (2014). A local connectivity scheme was implemented to introduce more physiological plausbility to the cortex model, and the effect on the rates, distributions and irregularity of neuronal firing, was compared to the original random connectivity method and experimental data. Local connectivity increased the distribution of and overall rate of neuronal firing. It also resulted in the irregularity of firing being more similar to those observed in experimental measurements, and a reduction in the variability in power spectrum measures. The larger variability in dynamical behaviour of the local connectivity model suggests that the topological structure of the connections in neuronal population plays a significant role in firing patterns during spontaneous activity. This model aims to take steps towards replicating the macroscopic network of the motor cortex, replicating realistic firing in order to shed light on information coding in the cortex. Large scale computational models such as this one can capture how structure and function relate to observable neuronal firing behaviour, and investigates the underlying computational mechanisms of the brain.

Introduction

The motor cortex is critical for movement. However, neurons in the motor cortex show patterns of ongoing activity even during resting state periods with no discernible movement (Schieber, 2011). This baseline spontaneous activity, which is observed in awake and conscious experimental subjects, is differentiable from task-induced states in recordings of neuron spiking behaviour (Dabrowska et al., 2021; Velliste et al., 2014; Churchland et al., 2012; Sauerbrei et al., 2020). Spontaneous activity of the brain at rest has spatially structured patterns of activity and is considered to

play an important role in neural encoding for brain processes (Ringach, 2009; Deco, Ponce-Alvarez, et al., 2013). In the past, activity in the cortex was thought to be driven by external sensory input, but it recently has been suggested that spontaneous activity arises from the dynamics of the neuronal circuitry and inputs modulate and modify the dynamics of the network rather than drive it (Destexhe, 2011; Chen and Gong, 2019). Thus, there is an interest in understanding the ongoing behaviour of neural activity, how it reflects the cortical circuitry and shapes the variable functions of the cortex (Tan, 2015; Destexhe, 2011; Baker et al., 1999).

Spontaneous activity in the motor cortex, similar to other cortical areas, is characterised by irregular firing of individual neurons, with low average firing rates (< 10 Hz) but a wide frequency range (0 - 100 Hz) (Dabrowska et al., 2021; Tomov et al., 2014; Hahn et al., 2010; Lacey et al., 2014; Borges et al., 2020). Firing rates in cortical networks measured in vivo are similarly reported to have highly positively skewed, long-tail distributions (Roxin et al., 2011). Neuron populations in the cortex also display oscillatory firing activity covering a broad frequency spectrum ranging from less than one hertz to hundreds of hertz (Buzsaki and Draguhn, 2004; Tomov et al., 2014). In the motor cortex, beta oscillations (13–30 Hz) are consistently observed in electroencephalography (EEG) measurements during resting states or prior to movement, and disappear during movement (Espenhahn et al., 2019; Schmidt et al., 2019).

The origin of the variability in the firing of cortical neurons is unknown and could be due to morphology of dendrites resulting in non-linear integration of inputs, or the properties of the network and synaptic coupling (Softky and Koch, 1993; Renart et al., 2010; Ostojic, 2014; Khanna and Carmena, 2017). The variability in neuronal activity is also believed to play a significant role in information encoding in the cortex (Habenschuss, Jonke, and Maass, 2013; Hawkins, Ahmad, and Cui, 2017; Shamir, 2014). Recently, the generation of neocortical activity has been addressed through computational models of somatosensory cortical circuits based on anatomical and physiological data, which generate realistic spontaneous dynamics (Markram et al., 2015; Potjans and Diesmann, 2014). However, specific model structure or features have not been clearly linked to cortical firing patterns.

The cortex is thought to have a consistent laminar organisation and patterns of connectivity across different regions (Douglas and Martin, 2004; Costa and Martin, 2010; Hawkins, Ahmad, and Cui, 2017). A canonical circuit which generally defines recurrent connections in each layer, layer 4 and 6 as input layers, and layers 2/3 and 5 as output layers, was based originally on Hubel and Wiesel (1962)'s work on the cat visual cortex and a similar pattern observed in other cortical areas in primate studies including the auditory cortex, somatosensory cortex and motor cortex (Douglas and Martin, 2004; Ghosh and Porter, 1988; Huntley and Jones, 1991; Costa and Martin, 2010). Binzegger, Douglas, and Martin (2004) quantified connectivity of the cortical circuit through in vivo intracellular recordings and morphometry of cell types and laminar distribution. This and other photostimulation and optogenetic studies of the mouse motor cortex have shown a dominant connection of layer 2/3 neurons to layer 5 neurons (Weiler et al., 2008; Hooks, Hires, et al., 2011; Hooks, Mao, et al., 2013).

Experiments involving in vivo extracellular injections of neuronal tracers in the cortex observe 'patchy' projections, in which synapses are highly localised (Huntley and Jones, 1991) (also see Voges, Schüz, et al. (2010) for review). Recent photostimulation experiments and digital image reconstruction suggest a distance-based connectivity model with strong connectivity within 0.2 mm of the soma and connectivity decreasing as a function of distance (Bender, Rangel, and Feldman, 2003; Shepherd et al., 2005). Diffusion tensor imaging also shows small-world, patchy connections in the cortex and significantly greater local connectivity than "random" connections (Gong et al., 2009). Highly localised connections are also thought to be optimal in regards to minimising wiring but maximising connectivity between nodes in a network. Local connections with some long range connections may be an efficient method of information transfer in the brain (Voges, Schüz, et al., 2010). Structural variations in network topology have a significant impact on the spatio-temporal activity of the cortex, with changes in connectivity patterns resulting in different dynamics (Senk

 et al., 2022).

Large scale neural network models can be used in exploring the question of how network struc-
ture influences the generation of cortical activity. Voges, Schüz, et al. (2010) have done significant
work in modelling the horizontal connectivity in the neocortex and showed more complex activity
patterns arise in structured, spatially clustered synapses. There recently has also been the devel-
opment of larger-scale cortical models which contain tens of thousands of spiking neuron models
(Esser, Hill, and Tononi, 2005; Potjans and Diesmann, 2014). Previous models of the motor cortex
have been limited in replicating physiological detail and mainly been focused on generating move-
ment dynamics or responses to stimulation (Farokhniaee and Lowery, 2021; Moody and Zipser,
1998; Esser, Hill, and Tononi, 2005; Manola et al., 2007). To the authors' knowledge, the role of net-
work connectivity in generation of the spontaneous cortical activity in physiologically-based spiking
neural network models, however, has not been explored.

Computational models can aid in the understanding of the electrophysiological mechanisms
which underlie the generation of spontaneous activity. In this study, a large scale spiking neural
network containing over 38,000 neurons and 150 million synapses, based on previous modelling
work by Potjans and Diesmann (2014), was implemented in the Python-based neural network simu-
lator, Brian2 (Stimberg, Brette, and Goodman, 2019), to replicate the spontaneous firing behaviour
of the motor cortex and explore the effect of local connectivity on the network behaviour. The aim
of this work was to provide insights into resting state motor cortex dynamics using a spiking neural
network model and investigate connectivity as a potential source of variability and efficient infor-
mation transmission in spontaneous firing behaviour.

Methods

This model was based on previous work by Potjans and Diesmann (2014) and implemented in
Brian2 with reference to source code from Shimoura et al. (2018). The original cortical circuit model
was adapted to represent a 1 mm^2 surface area of the motor cortex. The model follows a laminar
structure, grouping the neurons into four layers, 2/3, 4, 5, and 6. Each layer is further divided
into excitatory and inhibitory cell groups. Layer 1 is ignored due to its low density of neuronal cell
bodies (Muralidhar, Wang, and Markram, 2014). Connection weights in the circuit are shown in
figure 1 below.

Figure 1. Circuit diagram of model showing connections between the populations of excitatory (E) and
inhibitory (I) neuron groups. Groups are organised according to their layer. The colour of the lines indicates
the source group. Thickness is indicative of relative number of connections and weighting of synapses.

Individual neuron dynamics were governed by leaky-integrate-and-fire equations simulated us-
ing the linear state updater with a time step of 0.1 ms. Equation 1 below describes the membrane
potential (V) of each neuron, where τ_m is the time constant, C_m is the membrane capacitance, V_r

is the reset value for the membrane potential following a spike and I_{syn} is the total input current
described by equation 2. Action potentials were fired whenever $V(t)$ became more positive than
the threshold (θ). On the firing of a presynaptic neuron, the synaptic current of the postsynaptic
neuron (I_{syn}^{post}) was changed by the value determined by the conductance (g) multiplied by the weight
(w) after a delay (d) which accounts for the finite time interval of an action potential propagation in
a presynaptic neuron (equation 3). The delay between the spiking activity of a presynaptic neuron
and postsynaptic neuron was drawn from a normal distribution with a mean of 1.4 ms for exci-
tatory neurons and 0.8 ms for inhibitory neurons; the standard deviation of delay times was half
of the mean value. Following a spike being fired, the membrane voltage was reset to -65 mV. The
refractory period of the neuron was 2 ms meaning another spike could not be fired within that
period of time following a previous spike. As in the original Potjans and Diesmann (2014) model,
the conductance value in the layer 4E to layer 2/3E connection was doubled compared to the other
excitatory connections. Parameter descriptions and values are given in table 1.

$$\frac{dV(t)}{dt} = \frac{-(V(t) - V_r)}{\tau_m} + \frac{I_{syn}(t)}{C_m} \tag{1}$$

$$\frac{dI_{syn}}{dt} = -\frac{I_{syn}}{\tau_{syn}} \tag{2}$$

$$I_{syn}^{post}(t + d) = I_{syn}^{post}(t + d) + g \cdot w \tag{3}$$

$$\text{when presynaptic neuron V(t)} \geq \theta$$

Table 1. Parameters for neuron model. Taken from Potjans & Diesman, 2014.

Symbol	Name	Value
C_m	Membrane Capacitance	250 pF
θ	Threshold	-50 mV
τ_{ref}	Refractory Time	2 ms
τ_m	Membrane Time Constant	10 ms
τ_{syn}	Synaptic Time Constant	0.5 ms
V_r	Reset Value	-65 mV
g_e	Excitatory Conductance	1
g_i	Inhibitory Conductance	-4
w	Synaptic Weight	87.8 pA
d_e	Mean Excitatory Delay	1.5 ms
d_i	Mean Inhibitory Delay	0.8 ms

Table 2. Neuron groups in the motor cortex model, the proportions of neurons in each group (kept constant in sensitivity analysis), and absolute number of neurons in each neuron group in the final model.

Group	2/3E	2/3I	4E	4I	5E	5I	6E	6I
Proportion	0.268	0.076	0.063	0.014	0.284	0.071	0.187	0.038
Absolute	10332	2916	2412	540	10944	2736	7200	1476

Neurons were distributed over eight populations, according to the numbers in table 2. Neuron
numbers were adjusted from the original Potjans and Diesmann (2014) model, switching the pro-
portion of layer 4 and 5 cells based on the motor cortex having a dominant layer 5 connectivity and
sparser cell distributions than other layers of the cortex (ie. the somatosensory and visual cortices
on which the original model was based) (Young, Collins, and Kaas, 2013). Previously, the motor
cortex has thought to lack a distinct layer 4 but recent studies support a functional, albeit small,

layer 4 in the motor cortex with similar connections to layer 4 in the somatosensory cortex, notably
as an input pathway from the thalamus (Yamawaki et al., 2014; Barbas and Garcia-Cabezas, 2015).
The superficial layers of 2/3 and 4 contain approximately 40% of the neurons, layer 5 has 35% of
the neurons and layer 6 has 25% of the neurons, which is similar to reported physiological experi-
mental data (Amunts et al., 1995). A sensitivity analysis was carried out to investigate the effect of
the total number of neurons in the network, to select a minimal number of neurons at which the
rate and irregularity of firing activity converges.

Potjans and Diesmann (2014) original connectivity map integrated the findings of multiple anat-
omical and electrophysiological studies to estimate the probabilities of connection and number of
synapses formed between each neuron group. The combining of these data from different species
and areas, including rat visual and somatosensory areas and cat visual striate cortex, builds on the
theoretical framework which suggests an equivalency across cortical areas. Comparative studies
of primary motor cortex also show functional conservation across mammals (Bakken et al., 2021).
The connectivity profile was based on a modified version of Peter's rule which proposes that the
number of synapses is dependent on the number of neurons collocated in the presynaptic and
postsynaptic layers and a probability value of connection between layers, derived from experimen-
tal data (Peters and Payne, 1993; Braitenberg and Schuz, 1998; Binzegger, Douglas, and Martin,
2004; Rees, Moradi, and Ascoli, 2017).

Based on Peter's rule and probabilities derived from anatomical and physiological studies, the
number of connections (K) was calculated for neuron groups using equation 4 (Equation 3 in Shi-
moura et al. (2018)) where C_a is the connection probability defined in table 3, and N_{pre} and N_{post} are
the sizes of the presynaptic and postsynaptic populations, respectively. The C_a values in the con-
nectivity matrix (Table 3) describing the probabilities of connections between groups were adapted
from the original Potjans and Diesmann (2014) study in this motor cortex model based on changes
to the number of neurons in each layer, while maintaining the average relative number of connec-
tions in each neuron target group as the original model. The resulting connectivity was similar to
experimental investigations of layer specific wiring in the motor cortex (Weiler et al., 2008), notably
the dominant layer 2/3 to L5 connection pathway (Figure 2).

$$K = \frac{log(1 - C_a)}{log(1 - 1/(N_{pre} N_{post}))} \tag{4}$$

Table 3. Connectivity probability values (C_a) used in equation (4) to determine the number of synapses (K) between each group.

Target Group	Source Group							
	2/3E	2/3I	4E	4I	5E	5I	6E	6I
2/3E	0.192	0.3095	0.3356	0.5802	0.0143	0	0.0159	0
2/3I	0.252	0.2553	0.2558	0.4183	0.034	0	0.008	0
4E	0.016	0.012	0.3725	0.7704	0.0031	0	0.0879	0
4I	0.1334	0.006	0.5266	0.8295	0.0013	0	0.2007	0
5E	0.1902	0.1202	0.3785	0.0592	0.0377	0.1662	0.0396	0
5I	0.1071	0.0533	0.2129	0.0201	0.027	0.1374	0.0179	0
6E	0.0318	0.014	0.1754	0.1597	0.0257	0.0078	0.0784	0.399
6I	0.0708	0.002	0.0269	0.0101	0.0125	0.0031	0.1276	0.267

The original Potjans and Diesmann (2014) cortical model implemented a random connectivity
scheme. This was reimplemented, adapted to model the motor cortex, as described above, and
a new local connectivity scheme was developed. In the random and local connectivity schemes,
the total number of synapses in both models was kept constant for fair comparison. Neurons
were given spatial parameters (X, Y, and Z coordinates) to be distributed over a 1 mm^2 surface and
a depth range based on the group's layer. The likelihood of a neuron connecting with another

Figure 2. Connectivity matrix of neuron populations in the model

neuron was determined by a Gaussian distribution over distance, where x and y are a neuron's spatial location and the $radius$ parameter defines the extent of the spread (Equation 5). This connectivity is compatible with the synapse definition of the original random model described earlier but the added spatial description of neurons allowed constraining the synapses to a localised connectivity. The spatial connectivity description was also adapted from previous cortical modelling work (Lumer, Edelman, and Tononi, 1997; Hill and Tononi, 2004; Esser, Hill, and Tononi, 2005). Intralaminar connectivity had a greater radius of connectivity than interlaminar connectivity and inhibitory connections had a lower connectivity radius than excitatory connections as supported by physiological data (Bender, Rangel, and Feldman, 2003; Boucsein et al., 2011; Fino, Packer, and Yuste, 2013; Pelt and Ooyen, 2013). The radius value for different connection types is given in table 4. Multiple connections could be established between two neurons in both the random and local connectivity schemes and self connections in the local connectivity model were not allowed. Figure 3 shows the postsynaptic connections of a single neuron (red) on the random connectivity model and the connectivity of a single neuron in the distance based connectivity model.

$$Probability = exp\left(-\frac{(x_{pre} - x_{post})^2 + (y_{pre} - y_{post})^2}{2 * radius^2}\right) \tag{5}$$

Table 4. Connection radius for various connection types

Connection Type	Radius (μm)
Horizontal Intralaminar Excitatory Connections (Layer 2/3, 4 & 5)	300
Horizontal Intralaminar Excitatory Connections (Layer 6)	225
Vertical Interlaminar Excitatory Connections	50
Inhibitory Connections	175

External input was applied to each neuron group as Poisson spike trains at a frequency of 8 Hz. The number of these inputs for each neuron was constant at 2000 inputs for excitatory neuron groups and 1850 inputs for inhibitory neuron groups. This followed the layer-independent input protocol from the original Potjans and Diesmann (2014) model. A sensitivity analysis on the effect of the input on the resulting firing rates was also carried out by manipulating the number of inputs and frequency.

To characterise the activity of the network, spike trains and firing rates of neurons and neu-

Figure 3. Connectivity of a single excitatory neuron on layer 2/3 (a) random connectivity (b) local connectivity. Model represents $1\,mm^2$ of cortical surface. Units along all axes are in mm.

ron groups were monitored. A spike was defined by the neuron reaching the threshold value of -50 mV. Firing rates were calculated by counting the total number of spikes in each time step of the simulation (0.1 ms). Simulations were run for 500 ms and the first 50 ms was ignored to allow the model to settle. The interspike intervals (ISIs) for each neuron were calculated as the time between spikes. Irregularity was characterised by the coefficient of variation (CV) of the ISI distribution. CVs were calculated as the ratio of the standard deviation to the mean ISIs (Equation 6) for each individual neuron in the population, as defined previously (Shimoura et al., 2018; Potjans and Diesmann, 2014). Firing rates and CVs reported are the mean across neurons in the neuron group, and over time. Notably, firing rates and CVs closely matched values reported by the original model when the settling period was included, consistent with the analyses conducted by Shimoura et al. (2018), but due to the inconsistency in the spiking behaviour in the first 50 ms, this settling period was excluded for the analyses carried out in this study.

$$CV = \frac{\sigma_{ISI}}{\mu_{ISI}} \tag{6}$$

Results

The motor cortex receives connections from multiple brain regions such as the thalamus, premotor cortex and somatosensory cortex; this complexity makes selection of an appropriate surrogate input challenging. The effect of the input number and frequency to the model was explored while keeping the ratio of excitatory and inhibitory input and relative input between the different neuron groups constant (ie. 2000 for excitatory neurons, 1850 for inhibitory neurons). A single input frequency was used across the model though physiologically there may be variation in the number of inputs and range of input frequencies. Networks with local connectivity show greater sensitivity to the frequency and number of inputs, with a greater range of resultant average firing rates. Figure 4 shows a heat map of the effect of input number and frequency on the average firing rates of neurons in the model, note the change in scale in each colour map.

An analysis of the effect of the number of neurons in the model shows that consistent frequencies and CVs are reached in a random network containing 35,000 or more neurons. Neuron counts by isotropic fractionator and flow fractionator methods of the motor cortex in old and new world primate species lie in the estimated range of 50,000–90,000 neurons per mm² (Young, Collins, and Kaas, 2013; Collins et al., 2016). Figure 5 shows the firing rates and CVs in the network over a range of 0–80,000 neurons for both random and local connectivity, with standard deviation measures over ten simulations. The firing rates and CVs were more stable in the local connectivity model at lower number of neurons. In both models, layer 6E neuron CVs were not able to be calculated or were highly variable, probably due to very low firing rates in the neuron group, and so are excluded from the graphs below.

Figure 4. Average firing rates in model with changes to input number and frequency for (a) random connectivity and (b) local connectivity. Inhibitory neuron inputs were kept proportional at 92.5% of Excitatory inputs. Scales represent average firing rate across all neurons. Note the different range in scales of the colour maps.

Figure 5. Numerical sensitivity analysis for firing rates of neuron groups in (a) random connectivity and (b) local connectivity, and the coefficient of variation in neuron groups with (c) random connectivity model of the motor cortex and (d) local connectivity. The proportions of each neuron group and relative number of synapses were kept the same.

The firing rates of the model lie in an appropriate range of baseline physiological recordings of
a number of studies, with lower firing rates in layer 2/3 and layer 6 of the model than layer 5 (Cro-
chet and Petersen, 2009). Beloozerova, Sirota, and Swadlow (2003) reported on microelectrode
measurements of the motor cortex in awake rabbits, with mean discharge rates of 0.6 Hz in cortic-
ocortical neurons, prevalent in layer 2/3, 5.7 Hz in layer 5 neurons and 0.4 Hz in layer 6 neurons.
Schiemann et al. (2015) published similar values from whole cell patch clamp recordings in mice.
Dabrowska et al. (2021) recorded firing rates in two awake monkeys and firing rates in a resting
state were 6.3 Hz and 7.81 Hz and CV measures of 0.83 and 0.79 which is lies closer to the irregular-
ity of a random poisson distribution (CV = 1). This was recorded using a 100 electrode Utah Array in
the hand motor area of macaque monkeys, at a depth of 1.5 mm which would correspond to deep
layer 3 or superficial layer 5 (Ninomiya et al., 2019). Lacey et al. (2014) reported a similar mean
value of 7.1 Hz firing in layer 5 motor cortex cells at rest, from micro-electrode recordings in rats.
CVs of spiking activity in cortical neurons have been consistently reported in previous literature as
being close to 1 (Noda and Adey, 1970; Holt et al., 1996).

Table 5. Mean Firing Rates (FR) and Standard Deviation (Std) of Neuron Groups in Random and Local Connectivity schemes.

Neuron Group	Random Connectivity				Local Connectivity			
	Mean FR (Hz)	Std FR	Mean CV	Std CV	Mean FR (Hz)	Std FR	Mean CV	Std CV
2/3E	1.86	8.44	0.51	0.27	3.24	6.98	0.42	0.21
2/3I	4.81	11.78	0.56	0.26	6.57	8.81	0.57	0.25
4E	3.99	7.44	0.48	0.24	2.55	5.25	0.48	0.23
4I	5.51	12.97	0.51	0.24	7.42	14.71	0.75	0.28
5E	6.90	15.16	0.58	0.24	10.51	10.87	0.79	0.27
5I	8.13	9.22	0.51	0.22	9.90	10.49	0.67	0.26
6E	0.008	0.034	-	-	0.125	0.59	0.55	0.27
6I	6.42	9.10	0.51	0.22	8.66	10.96	0.67	0.25

Based on these experimentally reported values, the firing rates of the neural network model
with random connectivity were closer to the physiological range, though with slightly higher firing
in layer 2/3 and lower in 6. The local connectivity model showed an increase in mean firing rates
over all the neuron populations and a wider range in mean CV. The mean CV values in the local con-
nectivity model more closely matched experimental recordings in the middle-deep layers, notably
in 5E which is the dominant neuron group. Firing frequencies were also observed to be higher
in inhibitory neuron groups. Table 5 reports the mean values and standard distributions of firing
rates and irregularity (CV) in the random connectivity model and local connectivity model.
The activity of neurons in the model showed a different pattern of activity in the local connectiv-
ity model compared to the random connectivity. Figure 6 shows a raster plot of 10% of the neurons
in the model and a plot of the frequencies across the neuron groups. Firing rates of the neuron
groups generally remained < 100 Hz but the random connectivity model showed a greater ten-
dency for sudden high frequency firing activity across the network. This synchrony of firing activity
could be neuronal avalanches, bursts of firing, which are a noted feature of brain dynamics and
indicator of scale-free critical dynamics in complex systems (Arviv, Goldstein, and Shriki, 2019).
The range of firing rates in recorded neurons also varies with long-tailed distributions of firing
rates (Roxin et al., 2011). The range of firing is reported to be 0–40 Hz in measurements of awake
mice and cats (Armstrong and Drewt, 1984; Estebanez et al., 2017), while recordings in monkeys
ranged from 0–100 Hz (Dabrowska et al., 2021). The firing rates of the model also showed long
tailed distributions (Figure 7), with longer tailed distributions, closer to physiological data, in the
local connectivity model.
A power spectrum analysis shows a peak between 10–30 Hz which lies within the beta oscillation

Figure 6. Raster plots of the model spiking behaviour with (a) random connectivity and (b) local connectivity. Population rates of model with (c) random connectivity and (d) local connectivity.

Figure 7. Firing rates box and whiskers for (a) Random connectivity and (b) Local Connectivity. Irregularity (CV) box and whiskers plot for (c) Random Connectivity and (d) Local Connectivity.

range (13–30 Hz). Experimentally, beta waves are observed by EEG, MEG or ECoG, though the origin and role of this rhythm is still unclear (Jensen et al., 2005; Khanna and Carmena, 2017; Kilavik et al., 2013; Barone and Rossiter, 2021). This model suggests that oscillatory rhythms could be emergent properties of the cortical network since the input to this model is non-oscillatory. The standard deviation, taken over ten simulations, was slightly reduced in the local connectivity model suggesting a potential regularisation of the power spectrum and beta peak (Figure 8).

Figure 8. Power spectrum of random (left) and local (right) connectivity with observed beta 'peaks' between $10 - 30$Hz.

Discussion

This study builds on previous cortical modelling work, and investigates the effects of a physiologically-realistic, spatially defined local connectivity scheme on neuron activity, while focusing on replicating the spontaneous firing in the motor cortex. We were able to replicate spontaneous firing behaviour in regards to neuron firing rates, irregularity, and power spectrum peaks, as recorded by previous experimental data. Local and random connectivity was compared in the model to shed light on the topological network properties that influence population-based firing behaviour. Local connectivity showed qualitatively different temporal patterns of firing and more realistic irregularity (CV). It also resulted in a wider tailed distribution of firing and a narrower standard deviation in the power spectrum.

Spontaneous activity in the cortex has been characterised by low firing rates and asynchronous activity. The model with random connectivity replicated previously measured values with lower firing rates in layers 2/3 and 6 and higher firing rates in layer 5. The increase in firing rates in the local connectivity model may be due to more coordinated inputs within a local cluster resulting in a greater likelihood to reach threshold. This could be broadly explained by the theory of neurons as coincidence detectors, which states that information is encoded and propagated by the timing of action potentials (Softky and Koch, 1993; Shadlen and Newsome, 1994; Chen, Zhang, et al., 2013; deCharms, 1998). Though local connectivity increased the overall firing of the network, similar proportions of firing rate values between layers were maintained, which suggests additional tuning of input or overall connectivity density might resolve the differences in absolute firing rate and be able to reduce firing to match experimental results.

The local connectivity model better replicates experimentally derived CV values in the deeper layers of 5 and 6, with similar CV values to the random connectivity model in the superficial layers 2/3 and 4. This means that the topological structure could play a key role in the variability of firing activity. The local connectivity model also shows a slightly lower standard deviation in the power spectrum. The narrower range in the power spectrum suggests a more reliable occurrence of beta wave frequencies that are consistently observed in EEG recordings. The more realistic, local connectivity model could therefore be a topological pattern in the cortex which contributes to the occurrence of the beta wave.

The model showed more sensitivity and instability in response to input activity in the local connectivity scheme. Within the range of changes in the input, the local connectivity model covered a substantially larger range of frequencies in the firing behaviour of the network. The greater sensitivity to input could be a key factor in the highly variable activity in the cortex which might be necessary for phase transitions or changes in state, a notion described by criticality (Beggs and Timme, 2012; Arviv, Goldstein, and Shriki, 2019; Chialvo, 2010; Deco and Jirsa, 2012). This critical state is hypothesised to be functionally beneficial for efficient information transmission in the cortex (Shew and Plenz, 2013). In the motor cortex, being in the critical state may play a role in the generation of a wide range of voluntary muscle movements.

The random connectivity model has been used in previous studies of cortical dynamics (van Vreeswijk and Sompolinsky, 1996; Brunel, 2000). Random networks are less clustered in their connections, have longer range connections, and are considered to be inefficient with regards to information transfer (Westlake and Nagarajan, 2011). Mehring et al. (2003) implemented a locally connected random network (LCRN), with distance-based connectivity in a large scale neural network model containing 100,000 neurons. Kumar, Rotter, and Aertsen (2008) also used a LCRN to investigate the propagation of synchronous and asynchronous activity. Typically, these models have two populations, excitatory and inhibitory neuron groups with recurrent intra- and intergroup connectivity. Previous investigations of connectivity have been limited to 2D networks which have not aimed to replicate physiological behaviour (Yger et al., 2011; Voges and Perrinet, 2012). However, to further investigate the spatio-temporal activity of the cortex, particularly with longer range connections, across a cortical network a larger spatial surface area is still needed and this will require increased computing power.

Whether the connection of neurons in the cortex are specific or random has not yet been fully resolved, though recent studies have suggested that neuron morphology and patterns of recorded activity support a notion of specific connections (Callaway, 2002; Udvary et al., 2022). Axons do not simply connect to neurons based on spatially overlapped locations but selectively target specific neuron types or groups, for example in layer specific connectivity patterns (Callaway, 2002; White, 2002). There also appears to be both local and long-range connections in the cortex which is suggested to be efficient in regards to wiring cost and information transfer (Voges, Schüz, et al., 2010; Voges and Perrinet, 2012). A combination of short and long-range connections were included within the local connectivity model with horizontal, intralaminar connections spanning the surface area of the model while vertical, interlaminar connections were narrower in radius.

Over 70 years ago, Mountcastle (1957), through experimental recordings of neurons in the cat somatosensory cortex, suggested an organisational unit of a 'vertical group of cells extending through all cortical layers' known as a cortical column. Columns are groups of neurons, in the range of 0.5 mm in diameter, tuned to a specific stimulation or attribute in the range of 0.5 mm in diameter, with the arrangement of columns patterned like a 'mosaic' (Horton and Adams, 2005). However, the function of this columnar organisation remains unclear and there is still no agreed upon function or definition (Haueis, 2016; Schwalger, Deger, and Gerstner, 2017; Horton and Adams, 2005). In the motor cortex, recordings of neurons show directional tuning during upper limb movements with activity (Georgopoulos, Schwartz, and Kettner, 1986). Multi-unit recordings also support functional clusters which exhibit directional tuning, with widths of the tuning field reported in the range of 50-250 μm in diameter Amirikian and Georgopoulos, 2003; Georgopoulos and Stefanis, 2007; Hatsopoulos, 2010. Individual neurons can make connections outside of columns, with horizontal connections spanning up to 1.5 mm (Boucsein et al., 2011). This model only represented 1 mm^2 of cortical surface area: a future development of this model could, however, be an expansion in area representation with and without an added constraint to model cortical columns. The topographic structure of the motor cortex and its relation to muscle of movement parameters is still undefined and with this model, we can begin to explore these intricate connections in the motor cortex.

The effect of neuron types also may also be an interesting inclusion in any future investigation of cortical dynamics. Spontaneous activity may physiologically reflect a wider range of neuron

types or synaptic dynamics, such as bursting or faster and slower transients, than the simplified excitatory and inhibitory neurons captured in this model (Tan, 2015; Zeldenrust, Wadman, and Englitz, 2018; Masquelier and Deco, 2013; Zhao, Li, and Gu, 2020; Tomov et al., 2014). Firing activity could also be affected by differing input mechanisms. Stevens and Zador (1998), showed in vivo that independent synaptic current injections cannot account for variability in cortical neurons, but inputs arriving synchronously can lead to variability. Though our model only incorporated a simplified model of neuron dynamics and input, we showed that more realistic patterns of spontaneous activity can be replicated in a cortical circuit with local connectivity.

Our approach increases the biological plausibility of previous cortical modelling work and contrasts with previous models of the motor cortex which have used continuous-value recurrent neural networks (Sussillo et al., 2016; Michaels et al., 2019). We recognise that we have not considered other properties such as neuron types, dendritic processing and synaptic plasticity that are also likely to play a role in firing dynamics. Biological learning paradigms such as reinforcement learning with spike-timing dependent plasticity (STDP) have previously been incorporated in spiking neuronal models of motor control, though not specifically replicating motor cortex activity (Spüler, Nagel, and Rosenstiel, 2015). Concurrently, experimental data will also be vital for continued model development and validation, a synergy of computational and experimental techniques will be required to fully elucidate the connectivity of cortical circuits and how it contributes to the generation of dynamical activity.

In summary, the implementation of a local connectivity scheme in a spiking neural network model has shown that the topology of the network plays a substantial role in the resulting cortical dynamics. Our results support theories of structured cortical circuitry and local, patchy connectivity in the generation of spontaneous activity patterns in the motor cortex. Random networks appear to be more regular or synchronous in their behaviour, while the local connectivity model showed more realistic irregularity, particularly in the large pyramidal, output neurons of layer 5. Our model builds on previous work incorporating local connectivity in a more complex laminar structure and cortical circuit, and reproduces firing patterns comparable to those measured in vivo in the motor cortex. The output from our model indicates that the pattern of activity changes substantially, increasing in the range and irregularity in firing but with more stability in the power spectrum, as a result of the implementation of a more physiologically-based local network topology.

Conclusion

In developing this model, we have aimed to build on previous modelling work and keep parameters as physiologically realistic as possible to replicate the spontaneous firing activity in the motor cortex. To our knowledge, this is the first physiologically based spiking neural-network model to explore the effects of 3D spatially realistic connectivity on spontaneous neuron firing activity in the motor cortex. The pattern of connectivity is shown to play a significant role in the generation of irregular firing dynamics and long-tailed firing distributions of cortical neurons which could have an impact on criticality and information transmission. With the implementation of local connectivity in a structured cortical circuit, this model links neuroscientific theories of structure to functional network dynamics.

Acknowledgments

The author thanks Gonzalo Maso Talou and Gurleen Singh for their valuable discussions and Mark Sagar for initial project funding and administration. This preprint was formatted using the LaPreprint template (https://github.com/roaldarbol/lapreprint) by Mikkel Roald-Arbøl .

Author contributions

Conceptualisation: A.M., L.H.; Methodology: A.M., L.H.; Software: L.H.; Writing - original draft: L.H.; Review & editing: A.M., T.B., L.H.; Supervision: A.M. T.B.; Administration & Funding acquisition:

399 A.M., T.B.

References

401 Amirikian, B. and Georgopoulos, A. P. (2003). "Modular organization of directionally tuned cells in
402 the motor cortex: Is there a short-range order?" In: DOI: 10.1073pnas.2037719100.
403 Amunts, K. et al. (1995). "Postnatal development of the human primary motor cortex: a quantitative
404 cytoarchitectonic analysis". In: *Anat Embryol.* 192, pp. 557–571. DOI: 10.1007/BF00187186.
405 Armstrong, D. M. and Drewt, T. (1984). "Discharges of Pyramidal Tract and Other Motor Cortical
406 Neurones During Locomotion in the Cat". In: *J. Physiol* 346, pp. 471–495. DOI: 10.1113/jphysiol.
407 1984.sp015036.
408 Arviv, O., Goldstein, A., and Shriki, O. (2019). "Neuronal avalanches and time-frequency representa-
409 tions in stimulus-evoked activity". In: *Scientific Reports* 9.1. ISSN: 20452322. DOI: 10.1038/s41598-
410 019-49788-5.
411 Baker, S. et al. (1999). "The role of synchrony and oscillations in the motor output". In: *Exp Brain Res*
412 128, pp. 109–117. DOI: 10.1007/s002210050825.
413 Bakken, T. E. et al. (2021). "Comparative cellular analysis of motor cortex in human, marmoset and
414 mouse". In: *Nature* 598.7879, pp. 111–119. ISSN: 14764687. DOI: 10.1038/s41586-021-03465-8.
415 Barbas, H. and Garcia-Cabezas, M. A. (2015). "Motor cortex layer 4: less is more". In: *Trends in Neu-
416 rosciences* 38.5, pp. 259–261. DOI: https://doi.org/10.1016/j.tins.2015.03.005.
417 Barone, J. and Rossiter, H. E. (2021). "Understanding the Role of Sensorimotor Beta Oscillations".
418 In: *Frontiers in Systems Neuroscience* 15. DOI: 10.3389/fnsys.2021.655886.
419 Beggs, J. M. and Timme, N. (2012). "Being critical of criticality in the brain". In: *Frontiers in Physiology*
420 3. DOI: 10.3389/fphys.2012.00163.
421 Beloozerova, I. N., Sirota, M. G., and Swadlow, H. A. (2003). "Activity of Different Classes of Neurons
422 of the Motor Cortex during Locomotion". In: DOI: 10.1523/JNEUROSCI.23-03-01087.2003.
423 Bender, K. J., Rangel, J., and Feldman, D. E. (2003). "Development of Columnar Topography in the
424 Excitatory Layer 4 to Layer 2/3 Projection in Rat Barrel Cortex". In: *Journal of Neuroscience* 23(25).
425 DOI: 10.1523/JNEUROSCI.23-25-08759.2003.
426 Binzegger, T., Douglas, R. J., and Martin, K. A. (2004). "A quantitative map of the circuit of cat primary
427 visual cortex". In: *Journal of Neuroscience* 24.39, pp. 8441–8453. DOI: 10.1523/JNEUROSCI.1400-
428 04.2004.
429 Borges, F. S. et al. (2020). "Self-sustained activity of low firing rate in balanced networks". In: *Physica*
430 *A: Statistical Mechanics and its Applications* 537. DOI: 10.1016/j.physa.2019.122671.
431 Boucsein, C. et al. (2011). "Beyond the cortical column: Abundance and physiology of horizontal
432 connections imply a strong role for inputs from the surround". In: *Frontiers in Neuroscience*. DOI:
433 10.3389/fnins.2011.00032.
434 Braitenberg, V. and Schuz, A. (1998). *Peters' Rule and White's Exceptions*. Berlin, Heidelberg: Springer
435 Berlin Heidelberg, pp. 99–101. ISBN: 978-3-662-03733-1.
436 Brunel, N. (2000). "Dynamics of Sparsely Connected Networks of Excitatory and Inhibitory Spiking
437 Neurons". In: *Journal of Computational Neuroscience* 8, pp. 183–208. DOI: 10.1023/a:1008925309027.
438 Buzsaki, G. and Draguhn, A. (2004). "Neuronal Oscillations in Cortical Networks". In: *Science* 304,
439 pp. 1926–1930. DOI: 10.1126/science.1099745.
440 Callaway, E. (2002). "Cell type specificity of local cortical connections". In: *Journal of Neurocytology*
441 31, pp. 231–237. DOI: 10.1023/A:1024165824469.
442 Chen, G. and Gong, P. (2019). "Computing by modulating spontaneous cortical activity patterns as
443 a mechanism of active visual processing". In: *Nature Communications* 10.1. ISSN: 20411723. DOI:
444 10.1038/s41467-019-12918-8.
445 Chen, Y., Zhang, H., et al. (2013). "The Role of Coincidence-Detector Neurons in the Reliability and
446 Precision of Subthreshold Signal Detection in Noise". In: *PLoS ONE* 8 (2). ISSN: 19326203. DOI:
447 10.1371/journal.pone.0056822.

Chialvo, D. R. (2010). "Emergent complex neural dynamics". In: *Nature Physics* 6 (10), pp. 744–750. ISSN: 17452481. DOI: 10.1038/nphys1803.

Churchland, M. M. et al. (2012). "Neural population dynamics during reaching". In: *Nature* 487.7405, pp. 51–56. ISSN: 00280836. DOI: 10.1038/nature11129.

Collins, C. E. et al. (2016). "Cortical cell and neuron density estimates in one chimpanzee hemisphere". In: *Proceedings of the National Academy of Sciences of the United States of America* 113.3, pp. 740–745. ISSN: 10916490. DOI: 10.1073/pnas.1524208113.

Costa, N. M. da and Martin, K. A. (2010). "Whose cortical column would that be?" In: *Frontiers in Neuroanatomy*. ISSN: 16625129. DOI: 10.3389/fnana.2010.00016.

Crochet, S. and Petersen, C. C. (2009). "Cortical Dynamics by Layers". In: *Neuron* 64 (3), pp. 298–300. ISSN: 08966273. DOI: 10.1016/j.neuron.2009.10.024.

Dabrowska, P. A. et al. (2021). "On the complexity of resting state spiking activity in monkey motor cortex". In: *Cereb Cortex Commun* 2.3. ISSN: 26928205. DOI: 10.1101/2020.05.28.121095.

deCharms, R. C. (1998). "Information coding in the cortex by independent or coordinated populations". In: *Proc. Natl. Acad. Sci.* 95 (26), pp. 15166–15168. DOI: 10.1073/pnas.95.26.15166.

Deco, G. and Jirsa, V. K. (2012). "Ongoing cortical activity at rest: Criticality, multistability, and ghost attractors". In: *Journal of Neuroscience* 32 (10), pp. 3366–3375. ISSN: 02706474. DOI: 10.1523/JNEUROSCI.2523-11.2012.

Deco, G., Ponce-Alvarez, A., et al. (2013). "Resting-state functional connectivity emerges from structurally and dynamically shaped slow linear fluctuations". In: *Journal of Neuroscience* 33.27, pp. 11239–11252. ISSN: 02706474. DOI: 10.1523/JNEUROSCI.1091-13.2013.

Destexhe, A. (2011). "Intracellular and computational evidence for a dominant role of internal network activity in cortical computations". In: 21.5, pp. 717–725. ISSN: 09594388. DOI: 10.1016/j.conb.2011.06.002.

Douglas, R. J. and Martin, K. A. C. (2004). "Neuronal circuits of the neocortex". In: 27, pp. 419–415. ISSN: 0147006X. DOI: 10.1146/annurev.neuro.27.070203.144152.

Espenhahn, S. et al. (2019). "Cortical beta oscillations are associated with motor performance following visuomotor learning". In: *NeuroImage* 195, pp. 340–353. ISSN: 10959572. DOI: 10.1016/j.neuroimage.2019.03.079.

Esser, S. K., Hill, S. L., and Tononi, G. (2005). "Modeling the Effects of Transcranial Magnetic Stimulation on Cortical Circuits". In: *Journal of Neurophysiology* 94, pp. 622–639. ISSN: 0022-3077. DOI: 10.1152/jn.01230.2004.

Estebanez, L. et al. (2017). "Parvalbumin-Expressing GABAergic Neurons in Primary Motor Cortex Signal Reaching". In: *Cell Reports* 20.2, pp. 308–318. DOI: 10.1016/j.celrep.2017.06.044.

Farokhniaee, A. and Lowery, M. M. (2021). "Cortical network effects of subthalamic deep brain stimulation in a thalamo-cortical microcircuit model". In: *Journal of Neural Engineering* 18.5. ISSN: 17412552. DOI: 10.1088/1741-2552/abee50.

Fino, E., Packer, A. M., and Yuste, R. (2013). "The logic of inhibitory connectivity in the neocortex". In: 19.3, pp. 228–237. ISSN: 10738584. DOI: 10.1177/1073858412456743.

Georgopoulos, A. P., Schwartz, A. B., and Kettner, R. E. (1986). "Neuronal Population Coding of Movement Direction". In: *Science* 233 (4771), pp. 1416–1419. DOI: 10.1126/science.3749885.

Georgopoulos, A. P. and Stefanis, C. N. (2007). "Local shaping of function in the motor cortex: Motor contrast, directional tuning". In: *Brain Research Reviews* 55 (2 SPEC. ISS.), pp. 383–389. ISSN: 01650173. DOI: 10.1016/j.brainresrev.2007.05.001.

Ghosh, S. and Porter, R. (1988). "Morphology of pyramidal neurones in monkey motor cortex and the synaptic actions of their intracortical axon collaterals." In: *The Journal of Physiology* 400.1, pp. 593–615. ISSN: 14697793. DOI: 10.1113/jphysiol.1988.sp017138.

Gong, G. et al. (2009). "Mapping anatomical connectivity patterns of human cerebral cortex using in vivo diffusion tensor imaging tractography". In: *Cerebral Cortex* 19.3, pp. 524–536. ISSN: 10473211. DOI: 10.1093/cercor/bhn102.

Habenschuss, S., Jonke, Z., and Maass, W. (2013). "Stochastic Computations in Cortical Microcircuit Models". In: *PLoS Comput Biol* 9.11, p. 1003311. DOI: 10.1371/journal.pcbi.1003311.

Hahn, G. et al. (2010). "Neuronal Avalanches in Spontaneous Activity In Vivo". In: *J Neurophysiol* 104, pp. 3312–3322. DOI: 10.1152/jn.00953.2009.

Hatsopoulos, N. G. (2010). "Columnar organization in the motor cortex". In: *Cortex* 46 (2), pp. 270–271. ISSN: 00109452. DOI: 10.1016/j.cortex.2008.07.005.

Haueis, P. (2016). "The life of the cortical column: opening the domain of functional architecture of the cortex (1955–1981)". In: *History and Philosophy of the Life Sciences* 38.3. ISSN: 17426316. DOI: 10.1007/s40656-016-0103-4.

Hawkins, J., Ahmad, S., and Cui, Y. (2017). "A theory of how columns in the neocortex enable learning the structure of the world". In: *Frontiers in Neural Circuits* 11.81, pp. 1–18. ISSN: 16625110. DOI: 10.3389/fncir.2017.00081.

Hill, S. and Tononi, G. (2004). "Modeling Sleep and Wakefulness in the Thalamocortical System". In: *Journal of Neurophysiology* 93.3, pp. 1671–1698. ISSN: 0022-3077. DOI: 10.1152/jn.00915.2004.

Holt, G. R. et al. (1996). "Comparison of Discharge Variability In Vitro and In Vivo in Cat Visual Cortex Neurons". In: *Journal of Neurophysiology* 75 (5), pp. 1806–14. DOI: 10.1152/jn.1996.75.5.1806.

Hooks, B. M., Hires, S. A., et al. (2011). "Laminar analysis of excitatory local circuits in vibrissal motor and sensory cortical areas". In: *PLoS Biology* 9.1. ISSN: 15449173. DOI: 10.1371/journal.pbio.1000572.

Hooks, B. M., Mao, T., et al. (2013). "Organization of cortical and thalamic input to pyramidal neurons in mouse motor cortex". In: *Journal of Neuroscience* 33.2, pp. 748–760. ISSN: 02706474. DOI: 10.1523/JNEUROSCI.4338-12.2013.

Horton, J. G. and Adams, D. L. (2005). "The cortical column: A structure without a function". In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 360 (1456), pp. 837–862. ISSN: 09628436. DOI: 10.1098/rstb.2005.1623.

Hubel, D. H. and Wiesel, A. T. N. (1962). "Receptive Fields, Binocular Interaction and Functional Architecture in the Cat Visual Cortex". In: *J. Physiol* 160 (1), pp. 106–54. DOI: 10.1113/jphysiol.1962.sp006837.

Huntley, G. W. and Jones, E. G. (1991). "Relationship of Intrinsic Connections to Forelimb Movement Representations in Monkey Motor Cortex: A Correlative Anatomic and Physiological Study". In: *Journal of Neurophysiology* 66 (2), pp. 390–413. DOI: 10.1152/jn.1991.66.2.390.

Jensen, O. et al. (2005). "On the human sensorimotor-cortex beta rhythm: Sources and modeling". In: *NeuroImage* 26.2, pp. 347–355. ISSN: 10538119. DOI: 10.1016/j.neuroimage.2005.02.008.

Khanna, P. and Carmena, J. M. (2017). "Beta band oscillations in motor cortex reflect neural population signals that delay movement onset". In: *Elife*. DOI: 10.7554/eLife.24573.001.

Kilavik, B. E. et al. (2013). "The ups and downs of beta oscillations in sensorimotor cortex". In: *Experimental Neurology* 245, pp. 15–26. ISSN: 00144886. DOI: 10.1016/j.expneurol.2012.09.014.

Kumar, A., Rotter, S., and Aertsen, A. (2008). "Conditions for propagating synchronous spiking and asynchronous firing rates in a cortical network model". In: *Journal of Neuroscience* 28.20, pp. 5268–5280. ISSN: 02706474. DOI: 10.1523/JNEUROSCI.2542-07.2008.

Lacey, M. G. et al. (2014). "Spike firing and IPSPs in layer V pyramidal neurons during beta oscillations in rat primary motor cortex (M1) in vitro". In: *PLoS ONE* 9 (1). ISSN: 19326203. DOI: 10.1371/journal.pone.0085109.

Lumer, E., Edelman, G., and Tononi, G. (1997). "Neural Dynamics in a Model of the Thalamocortical System". In: *Cerebral Cortex* 7, pp. 207–227. DOI: 10.1093/cercor/7.3.207.

Manola, L. et al. (2007). "Anodal vs cathodal stimulation of motor cortex: A modeling study". In: *Clinical Neurophysiology* 118.2, pp. 464–474. ISSN: 13882457. DOI: 10.1016/j.clinph.2006.09.012.

Markram, H. et al. (2015). "Reconstruction and Simulation of Neocortical Microcircuitry". In: *Cell* 163.2, pp. 456–492. ISSN: 10974172. DOI: 10.1016/j.cell.2015.09.029.

Masquelier, T. and Deco, G. (2013). "Network Bursting Dynamics in Excitatory Cortical Neuron Cultures Results from the Combination of Different Adaptive Mechanism". In: *PLoS ONE* 8 (10). ISSN: 19326203. DOI: 10.1371/journal.pone.0075824.

Mehring, C. et al. (2003). "Activity dynamics and propagation of synchronous spiking in locally connected random networks". In: *Biological Cybernetics* 88.5, pp. 395–408. ISSN: 03401200. DOI: 10.1007/s00422-002-0384-4.

Michaels, J. A. et al. (2019). "A neural network model of flexible grasp movement generation". In: *bioRxiv*. DOI: 10.1101/742189.

Moody, S. L. and Zipser, D. (1998). "A model of reaching dynamics in primary motor cortex". In: *Journal of Cognitive Neuroscience* 10 (1), pp. 35–45. DOI: 10.1162/089892998563761.

Mountcastle, V. B. (1957). "Modality and Topographic Properties of Single Neurons of Cat's Somatic Sensory Cortex". In: *Journal of Physiology* 20.4, pp. 408–434. DOI: 10.1152/jn.1957.20.4.408.

Muralidhar, S., Wang, Y., and Markram, H. (2014). "Synaptic and cellular organization of layer 1 of the developing rat somatosensory cortex". In: *Frontiers in Neuroanatomy* 7. ISSN: 16625129. DOI: 10.3389/fnana.2013.00052.

Ninomiya, T. et al. (2019). "Layer specificity of inputs from supplementary motor area and dorsal premotor cortex to primary motor cortex in macaque monkeys". In: *Scientific Reports* 9 (1). ISSN: 20452322. DOI: 10.1038/s41598-019-54220-z.

Noda, H. and Adey, W. R. (1970). "Firing Variability in Cat Association Cortex During Sleep and Wakefulness". In: *Brain Research* 18 (3), pp. 513–526. DOI: 10.1016/0006-8993(70)90134-4.

Ostojic, S. (2014). "Two types of asynchronous activity in networks of excitatory and inhibitory spiking neurons". In: *Nature Neuroscience* 17, pp. 594–600. DOI: 10.1038/nn.3658.

Pelt, J. van and Ooyen, A. van (2013). "Estimating neuronal connectivity from axonal and dendritic density fields". In: *Frontiers in Computational Neuroscience* NOV. ISSN: 16625188. DOI: 10.3389/fncom.2013.00160.

Peters, A. and Payne, B. R. (1993). "Numerical Relationships between Geniculocortical Afferents and Pyramidal Cell Modules in Cat Primary Visual Cortex". In: *Cerebral Cortex* 3 (1), pp. 69–78. DOI: 10.1093/cercor/3.1.69.

Potjans, T. C. and Diesmann, M. (2014). "The cell-type specific cortical microcircuit: Relating structure and activity in a full-scale spiking network model". In: *Cerebral Cortex* 24.3, pp. 785–806. ISSN: 10473211. DOI: 10.1093/cercor/bhs358.

Rees, C. L., Moradi, K., and Ascoli, G. A. (2017). "Weighing the Evidence in Peters' Rule: Does Neuronal Morphology Predict Connectivity?" In: 40.2, pp. 63–71. ISSN: 1878108X. DOI: 10.1016/j.tins.2016.11.007.

Renart, A. et al. (2010). "The Asynchronous State in Cortical Circuits". In: *Science* 327, pp. 587–592. DOI: 10.1162/neco.2009.06-08-806.

Ringach, D. L. (2009). "Spontaneous and driven cortical activity: implications for computation". In: *Current Opinion in Neurobiology* 19 (4), pp. 439–444. ISSN: 09594388. DOI: 10.1016/j.conb.2009.07.005.

Roxin, A. et al. (2011). "On the distribution of firing rates in networks of cortical neurons". In: *Journal of Neuroscience* 31.45, pp. 16217–16226. ISSN: 02706474. DOI: 10.1523/JNEUROSCI.1677-11.2011.

Sauerbrei, B. A. et al. (2020). "Cortical pattern generation during dexterous movement is input-driven". In: *Nature* 577.7790, pp. 386–391. ISSN: 14764687. DOI: 10.1038/s41586-019-1869-9.

Schieber, M. H. (2011). "Dissociating motor cortex from the motor". In: *Journal of Physiology* 589.23, pp. 5613–5624. ISSN: 00223751. DOI: 10.1113/jphysiol.2011.215814.

Schiemann, J. et al. (2015). "Cellular Mechanisms Underlying Behavioral State-Dependent Bidirectional Modulation of Motor Cortex Output". In: *Cell Reports* 11.8, pp. 1319–1330. ISSN: 22111247. DOI: 10.1016/j.celrep.2015.04.042.

Schmidt, R. et al. (2019). "Beta oscillations in working memory, executive control of movement and thought, and sensorimotor function". In: 39.42, pp. 8231–8238. ISSN: 15292401. DOI: 10.1523/JNEUROSCI.1163-19.2019.

Schwalger, T., Deger, M., and Gerstner, W. (2017). "Towards a theory of cortical columns: From spiking neurons to interacting neural populations of finite size". In: *PLoS Computational Biology* 13.4. ISSN: 15537358. DOI: 10.1371/journal.pcbi.1005507. arXiv: 1611.00294.

Senk, J. et al. (2022). "Connectivity Concepts in Neuronal Network Modeling". In: *PLOS Computational Biology*. DOI: 10.1371/journal.pcbi.1010086.

Shadlen, M. N. and Newsome, W. T. (1994). "Noise, neural codes and cortical organization". In: *Current Opinion in Neurobiology* 4, pp. 569–579. DOI: 10.1016/0959-4388(94)90059-0.

Shamir, M. (2014). "Emerging principles of population coding: In search for the neural code". In: *Current Opinion in Neurobiology* 25, pp. 140–148. ISSN: 09594388. DOI: 10.1016/j.conb.2014.01.002.

Shepherd, G. M. et al. (2005). "Geometric and functional organization of cortical circuits". In: *Nature Neuroscience* 8.6, pp. 782–790. ISSN: 10976256. DOI: 10.1038/nn1447.

Shew, W. L. and Plenz, D. (2013). "The functional benefits of criticality in the cortex". In: *Neuroscientist* 19 (1), pp. 88–100. ISSN: 10738584. DOI: 10.1177/1073858412445487.

Shimoura, R. O. et al. (2018). "Reimplementation of the Potjans-Diesmann cortical microcircuit model: From NEST to Brian". In: *bioRxiv*. ISSN: 2692-8205. DOI: 10.1101/248401.

Softky, W. R. and Koch, C. (1993). "The Highly Irregular Firing of Cortical Cells Is Inconsistent with Temporal Integration of Random EPSPs". In: *The Journal of Neuroscience* 13 (l), p. 334350. DOI: 10.1523/JNEUROSCI.13-01-00334.1993.

Spüler, M., Nagel, S., and Rosenstiel, W. (2015). "A spiking neuronal model learning a motor control task by reinforcement learning and structural synaptic plasticity". In: *Proceedings of the International Joint Conference on Neural Networks*. Vol. 2015-September. Institute of Electrical and Electronics Engineers Inc. ISBN: 9781479919604. DOI: 10.1109/IJCNN.2015.7280521.

Stevens, C. F. and Zador, A. M. (1998). "Input synchrony and the irregular firing of cortical neurons". In: *Nature Neuroscience* 1, pp. 210–217. DOI: 10.1038/659.

Stimberg, M., Brette, R., and Goodman, D. (2019). "Brian 2, an intuitive and efficient neural simulator". In: *eLIFE* 8, pp. 1–41. DOI: 10.7554/eLife.47314.001.

Sussillo, D. et al. (2016). "A Neural Network that Finds a Naturalistic Solution for the Production of Muscle Activity". In: *Nature Neuroscience* 18 (7), pp. 1025–1033. DOI: 10.1038/nn.4042.

Tan, A. Y. (2015). "Spatial diversity of spontaneous activity in the cortex". In: *Frontiers in Neural Circuits* 9.September. ISSN: 16625110. DOI: 10.3389/fncir.2015.00048.

Tomov, P. et al. (2014). "Sustained oscillations, irregular firing, and chaotic dynamics in hierarchical modular networks with mixtures of electrophysiological cell types". In: *Frontiers in Computational Neuroscience* 8.SEP. ISSN: 16625188. DOI: 10.3389/fncom.2014.00103.

Udvary, D. et al. (2022). "The impact of neuron morphology on cortical network architecture". In: *Cell Reports* 39.2. ISSN: 22111247. DOI: 10.1016/j.celrep.2022.110677.

van Vreeswijk, C. and Sompolinsky, H. (1996). "Chaos in Neuronal Networks with Balanced Excitatory and Inhibitory Activity". In: *Science* 274 (5293), pp. 1724–1726. DOI: 10.1126/science.274.5293.1724.

Velliste, M. et al. (2014). "Motor cortical correlates of arm resting in the context of a reaching task and implications for prosthetic control". In: *Journal of Neuroscience* 34.17, pp. 6011–6022. ISSN: 15292401. DOI: 10.1523/JNEUROSCI.3520-13.2014.

Voges, N. and Perrinet, L. (2012). "Complex dynamics in recurrent cortical networks based on spatially realistic connectivities". In: *Frontiers in Computational Neuroscience* 6.JUL. ISSN: 16625188. DOI: 10.3389/fncom.2012.00041.

Voges, N., Schüz, A., et al. (2010). *A modeler's view on the spatial structure of intrinsic horizontal connectivity in the neocortex*. DOI: 10.1016/j.pneurobio.2010.05.001.

Weiler, N. et al. (2008). "Top-down laminar organization of the excitatory network in motor cortex". In: *Nature Neuroscience* 11.3, pp. 360–366. ISSN: 10976256. DOI: 10.1038/nn2049.

Westlake, K. P. and Nagarajan, S. S. (2011). "Functional connectivity in relation to motor performance and recovery after stroke". In: *Frontiers in Systems Neuroscience* 5, p. 8. DOI: 10.3389/fnsys.2011.00008.

White, E. L. (2002). "Specificity of cortical synaptic connectivity: Emphasis on perspectives gained from quantitative electron microscopy". In: *Journal of Neurocytology* 31 (3-5), pp. 195–202. DOI: 10.1023/a:1024109606722.

Yamawaki, N. et al. (2014). "A genuine layer 4 in motor cortex with prototypical synaptic circuit connectivity". In: *eLife* 3, e05422. ISSN: 2050084X. DOI: 10.7554/eLife.05422.

Yger, P. et al. (2011). "Topologically invariant macroscopic statistics in balanced networks of conductance-based integrate-and-fire neurons". In: *Journal of Computational Neuroscience* 31.2, pp. 229–245. ISSN: 15736873. DOI: 10.1007/s10827-010-0310-z.

Young, N., Collins, C., and Kaas, J. (2013). "Cell and neuron densities in the primary motor cortex of primates". In: *Frontiers in Neural Circuits* 7.30, pp. 1–11. ISSN: 16625110. DOI: 10.3389/fncir.2013.00030.

Zeldenrust, F., Wadman, W. J., and Englitz, B. (2018). "Neural coding with bursts—Current state and future perspectives". In: *Frontiers in Computational Neuroscience* 12. ISSN: 16625188. DOI: 10.3389/fncom.2018.00048.

Zhao, Z., Li, L., and Gu, H. (2020). "Different dynamical behaviors induced by slow excitatory feedback for type II and III excitabilities". In: *Scientific Reports* 10 (1). ISSN: 20452322. DOI: 10.1038/s41598-020-60627-w.