Sfrp2 is a multifunctional regulator of rodent color patterns

Matthew R. Johnson[1*], Sha Li[1*], Christian F. Guerrero-Juarez[2,3,4,5,6], Pearson Miller[7], Benjamin

J. Brack[1], Sarah A. Mereby[1], Charles Feigin[1], Jenna Gaska[1], Qing Nie[3,4,5,6], Jaime A. Rivera-

Perez[8], Alexander Ploss[1], Stanislav Y. Shvartsman[1,7], and Ricardo Mallarino[1+]

[1]Department of Molecular Biology, Princeton University; Princeton, New Jersey, 08540,

USA.

[2]Carle Illinois College of Medicine, University of Illinois at Urbana-Champaign, Urbana, IL

61801 USA.

[3]Department of Developmental and Cell Biology, University of California, Irvine, Irvine, CA

92697, USA.

[4]Department of Mathematics, University of California, Irvine, Irvine, CA, 92697, USA

[5]Center for Complex Biological Systems, University of California, Irvine, Irvine, CA 92697,

USA

[6]NSF-Simons Center for Multiscale Cell Fate Research, University of California, Irvine,

Irvine, CA, 92697, USA

[7]Center for Computational Biology, Developmental Dynamics, Flatiron Institute, Simons

Foundation; New York, New York, 10010, USA.

[8]University of Massachusetts Chan Medical School, Worcester, MA 01655, USA

*Equal contributions

+Corresponding author. Email: rmallarino@princeton.edu

Summary

Animal color patterns are strikingly diverse and can serve as a useful model for understanding

how tissues acquire positional information. Here, we study the coat of the African striped mouse

(*Rhabdomys pumilio*) to uncover mechanisms regulating the formation of stripe patterns. By

combining transcriptomic profiling, mathematical modeling, mouse transgenics, and *in vivo* gene

editing in striped mice, we show that the Wnt modulator, *Sfrp2,* plays two distinct roles in stripe

patterning. During embryogenesis, it regulates patterns of hair placode formation, producing the

embryonic prepattern that foreshadows pigment stripes while, in postnatal stages, it modulates

differences in hair color. This dual effect is achieved through spatiotemporal shifts in expression

and opposing effects on Wnt signaling within the same tissue. Thus, by uncovering a

multifunctional regulator of stripe formation, our work provides insights into the mechanisms by

which spatial patterns are established in developing embryos and the molecular basis of

phenotypic novelty.

Introduction

During embryogenesis, skin develops via reciprocal interactions between the epithelial cells that give rise to the epidermis and the underlying mesodermal cells that originate the dermis (Fuchs, 2007). This core developmental program is locally modified within a single organism, leading to marked spatial differences in thickness, pigmentation, and types of cutaneous appendages (Fuchs and Raghavan, 2002; Johansson and Headon, 2014). Such extraordinary variation evidences the rich complexity of the skin's underlying gene regulatory code and makes this tissue a fascinating model for understanding both the mechanisms that generate spatially restricted morphological patterns and the molecular basis of phenotypic diversity.

The skin of many vertebrate species displays characteristic periodic pigment patterns, such as spots and stripes, which play key roles in mediating intra- and inter-specific communication (Caro, 2005; Caro and Mallarino, 2020; Mills and Patterson, 2009). Theoretical simulations involving Turing reaction-diffusion systems, in which activators and inhibitors interact to establish organized spatial patterns, can closely recapitulate a wide variety of periodic color patterns seen in nature (Kondo, 2017; Kondo and Miura, 2010; Turing, 1990). Notably, however, Turing-generated patterns are variable and sensitive to stochastic perturbations (Vittadello et al., 2021). That is, Turing mechanisms are insufficient, by themselves, to explain the formation of periodic color patterns seen in many species, which are characterized by a high degree of spatial organization and reproducibility across individuals (Haupaix and Manceau, 2020; Kaelin et al., 2012, 2021; Mallarino et al., 2016; Patterson and Parichy, 2019). Therefore, a long-standing challenge has been to uncover the molecular, cellular, and developmental events by which periodic color patterns are specified and organized.

The African striped mouse (*Rhabdomys pumilio*), a rodent that exhibits a naturally evolved coat

pattern of dark and light parallel stripes (Fig. 1A, B), is a useful model in which to explore

molecular mechanisms underlying periodic color patterns because this mouse can be maintained

and reared in the lab, allowing for controlled experiments and development of molecular tools

(Mallarino et al., 2018). Here, we use a variety of multidisciplinary approaches to uncover the

developmental mechanisms controlling coat pattern formation in striped mice.

Results

Striped mice display regional variation in embryonic placode formation

Striped mice exhibit differences in hair length between striped skin regions that are apparent at

early postnatal stages (Fig. 1B) (Mallarino et al., 2016). We hypothesized that these differences

may reflect stripe-specific alterations in the developmental timing of hair follicle placodes

(hereafter referred to as placodes). In laboratory mice (*Mus musculus*), placodes are visible at

embryonic day 15.5 (E15.5) and are evenly distributed throughout the dorsal skin (Fig. 1C)

(Hardy, 1992; Millar, 2002). Remarkably, this stereotyped spatial pattern differs considerably in

striped mice. At the equivalent embryonic stage (*i.e.*, the emergence of visible placodes, ~E16.5

in striped mice), whole mount *in situ* hybridizations for early markers of placode formation (e.g.,

Dkk4, Ctnnb1, Wnt10b, Wif1, Eda, and *Dkk1*) (Andl et al., 2002) revealed that striped mouse

dorsal placodes develop in a stripe-like, spatially restricted manner, whereby they are present in

some regions along the dorsal skin but absent in others (Fig. 1C and Fig. S1). Approximately two

days later, at E18.5, placodes eventually become visible in regions previously devoid of them, as

evidenced by thickening of the epidermis and the appearance of molecular markers (Fig. S2).

Notably, stripe-like dorsal areas where placodes fail to form initially correspond to regions that

Figure 1. Development of hair placodes and pigment patterns in the African striped mouse. (A) An adult striped mouse in the field. (B) Postnatal Day 3 (P3) striped mouse pup displaying the characteristic striped pattern and corresponding differences in hair length across dorsal regions. (C) Side and dorsal views of Embryonic Day 15.5 (E15.5) laboratory mouse and E16.5 striped mouse embryos displaying spatial patterns of hair placode formation, as visualized by whole-mount *in situ* hybridization for *Dkk4*. (D) Light-microscopy images of striped mouse flat-mount skins from developmental and postnatal stages (E16.5 to P4) showing how patterns of hair placode development foreshadow pigmentation stripes. Photo credit in (A): Trevor Hardaker

94   will constitute the eventual dark pigment stripes, explaining why hair from dark stripes is shorter

95   at birth and suggesting that spatially restricted patterns of embryonic placode formation

96   foreshadow pigmentation stripes (Fig. 1D) (Mallarino et al., 2016).

Transcriptomic profiling and mathematical modeling identify a potential regulator of

placode formation

To identify regulators of the striped mouse placode developmental pattern, we performed an

unbiased bulk-level RNA-sequencing (RNA-seq) screen for genes differentially expressed

between placode rich and placode barren skin regions in E16.5 embryos. We used the boundaries

between these regions as a proxy to mark and isolate four distinct dorsal skin regions: R1 (the

dorsalmost, placode barren region), R2 (the dorsalmost, placode rich region), R3 (the

ventralmost, placode barren region), and R4 (the ventralmost, placode rich region) (Fig. 2A).

After data processing and read mapping (see STAR Methods), we performed a series of pairwise

comparisons (R1 vs. R2; R1 vs. R4; R2 vs. R3; and R3 vs. R4) to identify differentially

expressed genes between placode barren (R1, R3) and placode rich (R2, R4) regions. Although

each pairwise comparison yielded several differentially expressed genes (Fig. 2B, Table S1),

only one gene, *Sfrp2*, was differentially expressed in all four comparisons. Surprisingly,

however, *Sfrp2* was neither up nor downregulated in placode barren (R1, R3) or placode rich

regions (R2, R4). Rather, this gene was always upregulated in the dorsalmost region among the

two being compared (*i.e.,* R1>R2, R1>R4, R2>R3, and R3>R4). Indeed, when we extended our

comparisons to include R1 vs. R3 and R2 vs. R4, we confirmed that *Sfrp2* was differentially

expressed among these regions in a dorsoventral gradient (i.e., R1>R3 and R2>R4) (Fig. S3).

The dorsoventral expression pattern of a potential regulator of placode patterning was, *a priori*,

unexpected. However, previous theoretical work has pointed to upstream morphogen gradients

as a means for establishing stripe-like patterns (Hiscock and Megason, 2015). In mammals,

placode spacing is regulated by a self-organizing Turing reaction-diffusion system in which

Figure 2. Transcriptomics and mathematical modeling of placode development. (A) Schematic of a striped mouse E16.5 embryo and cross section displaying the regions (R1, R2, R3, R4) that were dissected for bulk-level RNA-seq analysis. Placodes are visible in R2 and R4 but not in R1 and R3. (B) UpSet plot summarizing genes differentially expressed (padj < 0.05) between Regions 1-4. Each row of dots represents a single pairwise comparison. Black dots indicate there were differentially expressed genes between a given comparison (number of differentially expressed genes plotted as columns) whereas grey dots indicate there were not. (C-E) Mathematical simulations of hair placode spacing in laboratory mouse and in striped mouse. (C) Equations used to model how a theoretical modulator (blue) could guide the activity of an activator-inhibitor reaction-diffusion system. Four parameters ($\mu, \nu, \rho,$ and K) governing chemical kinetics and representative of distinct mechanisms through which a modulator could influence patterning were tested. Approximately ~40,000 modulator distributions were tested for each parameter. (D) Simulations of a uniform modulator expression pattern (blue line) on variable µ produces a uniform pattern of Turing activity. This pattern is consistent with what is seen in laboratory mice *in vivo* (Fig. 1C). (E) Simulations in which a dorsoventral gradient of a modulator (blue line) applied to the same system in (D), leads to a stripe-like pattern of Turing activity. This recapitulates what is seen in striped mice *in vivo* (Fig 1C).

124   activators and inhibitors interact to establish an organized spatial pattern (Sick et al., 2006). We

125   therefore used mathematical modeling to ask whether a dorsoventral gradient could guide the

126   establishment of parallel stripes in a standard activator-inhibitor reaction-diffusion model. Our

127   model equations have four parameters governing local dynamics that are plausible targets of

128   upstream regulation (Fig. 2C). Each parameter represents a distinct mechanism through which a

129 regulator could influence the formation of parallel stripes, and each one can theoretically be

130 modulated by a spatial gradient. By directly simulating these equations, the specific effect of any

131 gradient on pattern formation can be tested. To this end, we used spectral numerical techniques

132 to perform exhaustive numerical screens to explore the effects of a wide range of plausible

133 regulator profiles, both monotonic and multi-peaked.

135 To specifically investigate how well a given upstream modulator can steer the activator-inhibitor

136 system into producing the parallel stripe pattern observed in striped mice, we performed four

137 computational screens, one for each parameter, in which we simulated pattern formation for a

138 sequence of ~40,000 different modulator distributions (see STAR Methods) to find distributions

139 which replicated a pattern consistent with the observed pattern *in vivo* (Fig. 2D, E). Optimal

140 regulator patterns were selected from a test set (Supplemental Movie 1) that produced final

141 patterns scored as having a high similarity to the stripe pattern of placode formation seen in

142 striped mice (Fig. 2E). Surprisingly, for three of the four potential pathways (μ, ρ, and ν), a

143 simple dorsoventral gradient was the most consistent distribution to robustly produce localized,

144 simulated peaks in a stripe-like manner. It is important to note that, while this is the simplest

145 solution computationally, in nature there are likely to be multiple reinforcing, partially redundant

146 mechanisms to assure robustness. However, our computational analysis strongly implies that a

147 dorsoventral orienting gradient can control establishment of the striped pattern.

149 Taken together, our transcriptomic experiments coupled to our mathematical simulation data

150 suggest that *Sfrp2*, a gene expressed in a dorsoventral gradient at a stage coinciding with the

emergence of embryonic placodes, may control the pattern of placode formation seen in striped

mice.

***Sfrp2* expression correlates negatively with Wnt signaling**

Sfrp2 is a known soluble modulator of Wnt signaling, having dual roles as both an activator and

repressor, depending on cellular microenvironment (Kim et al., 2016; Liang et al., 2019; Lin et

al., 2016; van Loon et al., 2021). To uncover the relationship between *Sfrp2* and Wnt signaling in

striped mouse skin, we dissociated dorsal skin from regions R1, R2, and R3 of E16.5 embryos,

as before, and performed single-cell 3' - RNA-sequencing (scRNA-seq, 10X Genomics). We

combined an equal number of cells from R1-R3 to generate a single pool representative of

striped mouse E16.5 dorsal skin and identified fourteen distinct cell types based on expression of

known molecular markers (Fig. 3A, Table S2) (Gupta et al., 2019; Rezza et al., 2016; Sennett et

al., 2015). *Sfrp2* was specifically expressed in a subset of dermal fibroblasts, as judged by co-

localization with canonical fibroblast markers (Fig. 3B, Table S2). Next, we clustered cells based

on *Sfrp2* expression to define a population of *Sfrp2*-high (i.e., > 2UMIs) and *Sfrp2*-low (i.e., < 2

UMIs) fibroblasts (Fig. 3C). Differential gene expression analysis between *Sfrp2*-high and *Sfrp2*-

low cells yielded numerous genes (P_{adj} <0.05), 482 of which were upregulated and 272 that were

downregulated in *Sfrp2*-high compared to *Sfrp2*-low cells (Table S3). Among the genes

Figure 3. Relationship between *Sfrp2* and Wnt signaling in striped mouse's embryonic skin. (A) Schematic of a striped mouse E16.5 embryo highlighting dorsal skin regions (R1-R3) dissected for scRNA-seq (left) and clustering of skin cell types based on their gene expression profiles (right). (B) Cells in green correspond to dermal fibroblasts, as determined by their high fibroblast score (left, see STAR Methods). *Sfrp2* expressing cells cluster with a subset of dermal fibroblast cells (right). (C) Fibroblasts clustered based on high (> 2 UMI) and low (< 2 UMI) *Sfrp2* expression. (D) Ontology analysis for genes that are downregulated in *Sfrp2* low cells, showing Wnt signaling as one of the main enriched pathways (Enrichr, $P_{adj} < 0.05$). (E) Expression levels for markers of canonical Wnt signaling, *Lef1* and *Ctnnb1*, show upregulation in *Sfrp2* low cells (non-parametric Wilcoxon Rank Sum test). (F-H*) In situ* hybridization for *Sfrp2* coupled with immunofluorescence for the keratin marker KRT14 showing differences in dermal *Sfrp2* expression between stage-matched striped mouse (F) and laboratory mouse (H) embryos. (G-I) Immunofluorescence for CTNNB1 and LEF1 in the developing skin of stage-matched striped mouse (G) and laboratory mouse (I) embryos. Red boxes denote zoomed-in regions of dorsal and lateral skin. (J) Representative GFP and Hematoxylin-Eosin (H&E) stained transverse sections showing, respectively, the distribution of SFRP2 and hair follicles in control and transgenic mice. (K) Quantification of hair follicle number in transgenic mice and controls (N = 4; *** P < 0.001). Dotted lines in (J) delineate the dermis-epidermis boundary. Scale bars: 200µm (zoomed out) and 25µm (zoomed in) in (F and H); 200µm (zoomed out) and 50µm (zoomed in) in (G and I); and 50µm in (J). NT = neural tube.

upregulated in *Sfrp2*-low fibroblasts, Gene Ontology analysis identified regulation of Wnt

signaling as the most significantly enriched pathway (Fig. 3D). Indeed, expression of numerous

Wnt-related genes, including key activators of the pathway (i.e., *Lef1* and *Ctnnb1*), were higher

in *Sfrp2*-low (Fig. 3E, Table S3) compared to *Sfrp2*-high cells, demonstrating that *Sfrp2* and Wnt

activation are negatively correlated in striped mice dermal fibroblasts.

To confirm and extend our previous observation, we examined spatial patterns of *Sfrp2*

expression and Wnt signaling markers. RNA *in situ* hybridizations in cross sections from E16.5

striped mice embryos, coupled to immunohistochemistry (IHC) for an epidermal marker

(KRT14), indicated that expression of *Sfrp2* was strong in mesenchymal cells directly above the

neural tube and decreased laterally, a result consistent with our bulk-level RNA-seq analysis

(Fig. 3F). At this same stage, immunofluorescence (IF) for LEF1 and CTNNB1 showed that

these markers were readily detectable in epidermal cells from lateral and ventral regions of the

embryo but were absent from cells above the neural tube and adjacent areas, where *Sfrp2*

expression was highest (Fig. 3G). This pattern contrasted with what was seen in stage-matched

laboratory mouse embryos (E15.5), which had overall lower levels of dermal *Sfrp2* expression

(Fig. 3H) and uniform levels of LEF1 and CTNNB1 throughout the epidermis, including above

the neural tube (Fig. 3I). Taken together, our scRNA-seq data coupled to RNA *in situ*/IF analysis

indicate that *Sfrp2* and Wnt signaling are negatively correlated and suggest that dermal *Sfrp2*

expression acts to inhibit Wnt signaling in striped mouse embryonic skin.

***Sfrp2* overexpression alters placode formation in laboratory mice**

Wnt signaling is a primary determinant of hair follicle formation and spacing, as perturbation of

members of this pathway leads to alterations in hair follicle number and distribution (Andl et al.,

2002; Gat et al., 1998; Millar, 2002; Tsai et al., 2014). This raises the intriguing hypothesis that

Sfrp2 may be acting to directly regulate striped mouse placode development. If this is the case,

one would predict that increasing expression of *Sfrp2* in embryonic skin with low endogenous

levels of *Sfrp2*, such as that of laboratory mouse (Rezza et al., 2016; Sennett et al., 2015), should

alter placode formation. To test this, we generated *Rosa^{Sfrp2-GFP}* mice, a strain that allowed us to

activate *Sfrp2* transcription in the presence of tissue-specific Cre drivers. To establish whether

dermal expression of *Sfrp2* was sufficient to regulate placode formation, as implied by our

striped mouse data, we crossed *Rosa^{Sfrp2-GFP}* mice with *Dermo-Cre* mice, a mesenchymal driver

of Cre (Šošić et al., 2003; Yu et al., 2003) (Fig. 3J). Indeed, double-transgenic embryos (*Dermo-*

Cre;*Rosa^{Sfrp2-GFP}*) analyzed at E17.5 showed a significant reduction in the number of

placodes/hair follicles, compared to control littermates (N= 4) (Fig. 3J, K). Thus, our transgenic

experiments in laboratory mice demonstrate a functional relationship between *Sfrp2* and

embryonic placode development in which elevated levels of *Sfrp2* in the dermis leads to a

marked reduction in the number of placodes/hair follicles.

The establishment of the *Sfrp2* dorsoventral gradient

In striped mice, *Sfrp2* is expressed in a dorsoventral gradient at E16.5, a stage coinciding with

the emergence of embryonic placodes. To understand how such a pattern is established in space

and time, we characterized the expression dynamics of *Sfrp2* by performing additional scRNA-

seq analysis from the days prior to E16.5 (Fig. 4A). We found that the number of fibroblasts

Figure 4. Spatial and temporal Sfrp2 expression patterns in striped mice and lab mice. (A) Schematic of striped mouse skin dissections from different developmental stages. For E16.5, scRNA-seq was performed on each individual region and data were pooled for an integrated analysis representative of whole striped mouse dorsal skin. (B) Both the percentage of fibroblasts that express *Sfrp2* (left, Spearman's Correlation, R = 0.79) and the expression levels of *Sfrp2* within those fibroblasts (right) increase in the days prior to visible placode formation in striped mouse dorsal skin. (C) Developmental trajectories inferred from striped mouse fibroblasts show regional differences in *Sfrp2* expression in latent time. (D) Schematic of laboratory mouse dissections from different developmental stages used for scRNA-seq. For all stages (E12.5-E15.5), whole dorsal skin was dissected (purple). (E) Both the percentage of fibroblasts expressing *Sfrp2* (left, Spearman's correlation, R = -0.78) and *Sfrp2* expression levels within those fibroblasts (right) decrease in the days prior to visible placode formation. (F) Developmental trajectories inferred from laboratory mouse fibroblasts show no substantial change in *Sfrp2* expression in latent time. Black lines in (B) and (E) indicate mean *Sfrp2* expression levels.

expressing *Sfrp2* was lowest at the earliest time-point sampled, E13.5 (36.03%), and gradually

increased in subsequent days (40.03% at E14.5, 45.34% at E15.5, and 44.23% at E16.5)

(Spearman correlation=0.79) (Fig. 4B). Moreover, average *Sfrp2* expression levels within those

fibroblasts also increased during this period (Fig. 4B). In agreement with these observations, we

observed a marked change in *Sfrp2* expression in a developmental trajectory of dermal

fibroblasts modeled in latent time. Specifically, *Sfrp2* exhibited a prominent spatiotemporal

increase across development toward R1-R3 regions at E16.5, having higher levels of expression

in the region closest to the dorsal midline relative to the more lateral regions (*i.e.*, R1>R2 and

R2>R3) (Fig. 4C).

We next asked whether the sharp increase in *Sfrp2* expression prior to placode formation was a

unique feature of striped mice or, instead, a conserved process in rodent skin development. To

address this, we performed scRNA-seq on laboratory mouse dorsal skin from multiple stages,

including those preceding placode formation as well as at stages when placodes have formed

(E12.5-E15.5) (Fig. 4D). In stark contrast to what was observed in striped mice, we found that

the percentage of *Sfrp2*-expressing fibroblasts was highest early on (48.44% at E12.5) and

decreased in subsequent days (28.44% at E15.5) (Spearman correlation=-0.78). In addition,

average *Sfrp2* expression levels within fibroblasts also decreased from E13.5-E15.5 (Fig. 4E).

Lastly, contrary to what we found in striped mice, *Sfrp2* expression levels in laboratory mice

fibroblasts showed no substantial change in latent time (Fig. 4F).

Together, our results indicate that *Sfrp2* levels in the dorsal mesenchyme of striped mice

gradually increase prior to placodes becoming visible and reach peak numbers at the onset of

placode formation. In line with our mathematical model and transgenic experiments, increased

Sfrp2 levels are sufficient to modulate the underlying developmental program controlling hair

follicle formation. Moreover, our comparative analysis reveals opposing dynamics in laboratory

mice, where *Sfrp2* expression levels decrease as placode formation nears, thereby allowing this

process to proceed uniformly across the dorsal skin.

246 ***Sfrp2* shifts in spatial expression and regulates pigmentation by activating Wnt signaling**

247 Our phenotypic characterization of striped mouse embryonic skin shows that, at E18.5, placodes

248 become visible in regions where they were previously absent (Fig. S2). We reasoned that if *Sfrp2*

249 indeed contributes to establishing patterns of placode formation in stripe mice, as our data

250 suggest, there should be a detectable change in the spatial expression of this gene once placodes

251 emerge throughout the dorsal skin. To test this, we characterized the spatial-specific patterns of

252 *Sfrp2* expression in late embryogenesis (E19.5) and early postnatal stages (P0 and P4) using

253 qPCR. For all samples, we dissected three different skin regions (R1-R3) using either hair-length

254 (E19.5) or visible pigmentation differences (P0 and P4) as a proxy for defining region

255 boundaries and isolating tissue (Mallarino et al., 2016) (Fig. 1D). In agreement with our

256 prediction, we found no regional differences in *Sfrp2* levels at E19.5, demonstrating there had

257 been a shift in the spatial expression pattern of this gene and that the dorsoventral gradient seen

258 at E16.5 was no longer present (Fig. 5A). Unexpectedly however, when we analyzed skin

259 regions from pups (P4), we found that *Sfrp2* had undergone an additional shift in its spatial

260 pattern and was now expressed at significantly lower levels in the light stripes (R2) compared to

261 the flanking dark stripes (R1 and R3) (Fig. 5A). The light-stripe specific decrease in *Sfrp2*

262 expression seen in P4 pups suggested this gene could be playing an additional role in striped

263 mouse skin, regulating hair color during the period in which pigment is synthesized and

264 deposited in the growing hair shaft. Furthermore, previous studies have shown that pigment

265 production/melanogenesis can be controlled by Wnt signals secreted from the dermal papilla

266 (Enshell-Seijffers et al., 2010; Morgan, 2014), a specialized mesenchymal component of the hair

267 follicle that signals to pigment-producing cells, or melanocytes. To explore a potential link

268 between *Sfrp2* and melanogenesis, we first performed fluorescent *in situ* hybridizations on P4

Figure 5. *Sfrp2* promotes melanogenesis by activating Wnt signaling. (A) Quantitative PCR (qPCR) showing *Sfrp2* mRNA fold change levels along different dorsal skin regions in embryonic and postnatal stages. (B) *In situ* hybridization showing specific *Sfrp2* expression in the dermal papilla of a P4 striped mouse hair follicle. (C-D) Functional *in vitro* experiments using lentiviruses. (C) Schematic of the design of *in vitro* functional experiments. Melanocytes were stably transduced with either a control (LV-*GFP*) or an experimental (LV-*Sfrp2^GFP*) lentivirus and sorted using FACS. (D) Fold change differences in expression of Wnt targets and melanogenesis genes in stably transduced control and experimental cells, as determined via qPCR. Statistical significance in panel (A) (**** $P < 0.0001$; *** $P < 0.001$; ** $P < 0.01$; *$P < 0.05$, N = 3 for E16.5, E19.5, and P0, N = 4 for P4) and in panel (D) (** $P < 0.01$; *$P < 0.05$; N = 4) was assessed using an ANOVA test.

striped mouse skin sections and found that *Sfrp2* was specifically expressed in the dermal papilla

(Fig. 5B), a pattern consistent with what is seen in laboratory mouse (Rezza et al., 2016; Sennett

et al., 2015) and with our hypothesis that this gene may regulate melanogenesis. Next, to test the

effect of *Sfrp2* expression on melanocyte behavior, we stably transduced an immortalized mouse

melanocyte cell line (Melan-A) with a lentiviral vector carrying the *Sfrp2* cDNA and used qPCR

to measure the expression of Wnt targets and of genes involved in melanogenesis (Fig. 5C, D).

Compared to melanocytes transduced with a control virus, cells overexpressing *Sfrp2* showed a

significant increase in Wnt target genes, including *Axin2*, *C-myc*, and *Ccnd1* (He et al., 1998; Jho

et al., 2002; Shtutman et al., 1999). Moreover, relative to controls, cells overexpressing *Sfrp2*

showed a marked increase in *Tyr* and *Mitf*, two genes with key roles in promoting melanin

synthesis (Iozumi et al., 1993; Steingrímsson et al., 2004) (Fig 5D). Notably, these results are

consistent with previous findings, in which recombinant SFRP2 was found to activate

Wnt/melanogenesis in human primary melanocytes via paracrine signaling and to promote

darkening of skin explants (Kim et al., 2016).

Taken together, our results indicate that the expression pattern of *Sfrp2* during embryogenesis

and early postnatal stages is dynamic, shifting from a 'high-low', dorsoventral gradient (E16.5)

to a 'high-low-high' pattern (P4) in which regions of low expression (i.e., light stripes) are

flanked by regions where expression is higher (i.e., dark stripes). Moreover, our spatial

expression data and *in vitro* functional experiments indicate that postnatal expression of *Sfrp2*

acts to regulate melanocyte behavior via Wnt activation, thereby revealing an additional

regulatory function for *Sfrp2* during stripe pattern formation.

***Sfrp2 in vivo* knockout in striped mice leads to changes in coat pattern and pigmentation**

Our data implicate *Sfrp2* as a key regulator of striped mouse placode formation and stripe-

specific pigmentation differences. However, to establish a direct causal link between *Sfrp2* and

coat pattern phenotypes, it is critical to perform functional experiments directly in striped mice.

To this end, we adapted a CRISPR-based experimental strategy in which recombinant adeno-

associated viruses (rAAVs) are used to deliver CRISPR-CAS9 reagent into the oviduct of

pregnant females. This *in vivo* gene editing technique is based on the ability of rAAVs to

transduce the zona pellucida on intact pre-implantation embryos (Yoon et al., 2018) (Fig. 6A).

First, to establish the feasibility of using this approach for inducing gene knockouts (KO) in

Figure 6. *In vivo* **genome editing in African striped mice.** (A) Schematic of the CRISPR-based approach used for generating targeted genome deletions in the striped mouse. (B) Postnatal Day 5 (P5) striped mouse littermates showing differences between a Tyrosinase (Tyr) wildtype individual ($Tyr^{+/+}$) and a knockout animal ($Tyr^{-/-}$) (C-D) Sfrp2 regulates stripe pattern width. (C) Representative differences in stripe width seen between wildtype ($Sfrp2^{+/+}$) controls and Sfrp2 knockout ($Sfrp2^{-/-}$) pups. (D) Measurements of stripe width in $Sfrp2^{+/+}$ and $Sfrp2^{-/-}$ pups. Statistical significance in panel (D) (**** $P < 0.0001$; *** $P < 0.001$; ** $P < 0.01$; *$P < 0.05$; $Sfrp2^{+/+}$, N = 10; $Sfrp2^{-/-}$, N = 7) was assessed using an ANOVA test. (E) Representative differences in coat color between $Sfrp2^{+/+}$ and $Sfrp2^{-/-}$ adult mice taken as a single image. (F) Quantification of pigmentation between $Sfrp2^{+/+}$ and $Sfrp2^{-/-}$ adult mice reveal stripe-specific changes in color. Statistical significance in panel (F) (*** $P < 0.001$; *$P < 0.05$; $Sfrp2^{+/+}$, N = 3; $Sfrp2^{-/-}$, N = 3) (G) Model describing the role of Sfrp2 in establishing and implementing the stripe color pattern in striped mice.

striped mice *in vivo*, we injected pregnant females carrying pre-implantation embryos with an

rAAV packaged with Nme2Cas9 and a single-guide RNA (sgRNA) expression cassettes directed

to the coding sequence of *Tyrosinase* (*Tyr*), the rate limiting enzyme for melanin production

(Edraki et al., 2019; Iozumi et al., 1993). Visual inspection of F0 pups coupled to genotyping

revealed this approach resulted in efficient gene editing, enabling us to generate complete KO

animals (Fig. 6B). To our knowledge, this is the first time that *in vivo* gene editing has been

performed in a mammalian species outside of laboratory mouse.

Encouraged by our proof-of-principle results, we next designed multiple sgRNA's against the 5'

coding region of *Sfrp2* (Fig. S4A; STAR Methods). We then tested each sgRNA in striped

mouse immortalized dermal fibroblasts and chose the one that had the highest indel-inducing

efficiency. Next, we injected an rAAV containing Nme2Cas9 and sg*Sfrp2* expression cassettes

into the oviducts of plugged females. From a total of 21 females injected, we obtained 57 pups,

14 of which carried gene edits in *Sfrp2*. After mating F0 founders and wildtype animals

(*Sfrp2$^{+/+}$*) to ensure germline transmission, we crossed F1 heterozygous animals (*Sfrp2$^{+/-}$*) to

generate F2 *Sfrp2* KO (*Sfrp2$^{-/-}$*) mice. We used western blot to confirm the successful elimination

of SFRP2 protein (Fig. S4B). Analysis of coat phenotypes from early postnatal (P3) stages

revealed significant and consistent differences in the coat patterns of F2 *Sfrp2$^{-/-}$* and *Sfrp2$^{+/+}$*

striped mice. Specifically, while the width of the striped area (i.e., the distance between both R3

stripes on either side of the midline) remained the same, we observed significant changes in the

relative width of the stripes, with *Sfrp2$^{-/-}$* mice having a narrower middle stripe and wider dark

stripes than *Sfrp2$^{+/+}$* mice (Fig. 6C, D). Thus, deletion of *Sfrp2* induces significant alterations in

the stripe boundaries within the striped area, effectively altering the stripe pattern. In addition to

the changes in the stripe pattern, analysis of coat coloration in adults, when pigment deposition

has been completed, revealed that the coats of $Sfrp2^{-/-}$ mice were lighter than $Sfrp2^{+/+}$ mice (Fig.

6E). Such difference was particularly evident in regions with light/banded hair (e.g., middle

stripe and light stripes), as determined by quantitative measurements taken from photographs

(Fig. 6F).

The stereotyped coat pattern of striped mice likely results from multiple, reinforcing mechanisms

and, thus, knockout of a single gene is not expected to completely abrogate the pattern (see

Discussion). However, our genome editing experiments in striped mice demonstrate a key role

for $Sfrp2$ in regulating both pigment distribution (i.e., color pattern) and pigment type (i.e.,

overall color) in this species.

Discussion

Over the course of millions of years of evolution, the skin developmental program has been

altered many times to produce a stunning array of phenotypic diversity that is apparent from

comparisons both across species and within the body of a single individual. Here, by taking

advantage of naturally evolved phenotypic variation in striped mouse, a species that we have

now transformed into a genetically tractable laboratory model, we studied the developmental

processes involved in regulating periodic stripe patterns. During mid embryogenesis, striped

mouse placodes emerge in a spatially restricted way, foreshadowing the coat pigment pattern

visible in early postnatal days. Embryonic regions where placodes initially fail to form give rise

to dark colored stripes, implying that a molecular pre-pattern is established either simultaneously

or prior to placode formation. Once this pre-pattern is established, cells in placode barren skin

regions acquire permanent changes that are then relayed to hair follicle signaling centers during

pigment production, or the implementation of the pre-pattern.

In birds and mammals, periodic pigment patterns arise from two sequential processes: the first

occurs early in embryogenesis and specifies the positional boundaries of the future pattern (*i.e.,*

pattern establishment, or prepattern); the second implements this prepattern through genes

mediating pigment production during hair growth and cycling, giving rise to the observed color

differences (*i.e., pattern implementation*) (Haupaix and Manceau, 2020; Kaelin and Barsh,

2013). We identify *Sfrp2* as a regulator of both the establishment of the pre-pattern and its

implementation, a dual role that is made possible by its dynamic expression pattern throughout

embryogenesis and early postnatal stages (Fig. 6G). Prior to placode formation, expression levels

of *Sfrp2* gradually increase in dorsal dermal fibroblasts, as does the number of *Sfrp2* expressing

dermal fibroblasts. When placodes become visible, *Sfrp2* is expressed in a sharp dorsoventral

gradient, with maximum levels in cells above the neural tube. Our simulations coupled to our

transgenic experiments in both laboratory mice and striped mice show that the *Sfrp2* dorsoventral

gradient can act to influence placode formation. Late in embryogenesis, the *Sfrp2* dorsoventral

expression gradient disappears and placodes form throughout the dorsal skin. Then, at early

postnatal stages, *Sfrp2* becomes differentially expressed in the dermal papilla of light and dark

stripe hair follicles, where it acts to regulate melanocyte behavior. These results contrast with

what is seen in felids, where the molecular players regulating pattern establishment are different

from the ones controlling the implementation events that follow (Eizirik et al., 2010; Kaelin et

al., 2012, 2021). The involvement of *Sfrp2* at multiple stages of pattern formation implies a fine-

tuned spatiotemporal regulatory control, likely achieved through a complex *cis*-regulatory

architecture and the action of multiple binding factors (Keller et al., 2020).

Secreted frizzled-related proteins (SFRPs) are known modulators of Wnt signaling that can exert

their effects through a wide variety of mechanisms, including Wnt ligand sequestering, binding

to Frizzled receptors, stabilization of Wnt-Frizzled complexes, and direct activation of Frizzled

(van Loon et al., 2021). As such, SFRPs can either inhibit or activate Wnt signaling, and their

effect is dependent on cell type, developmental stage, and cellular microenvironment(van Loon

et al., 2021). In striped mice, embryonic expression of *Sfrp2* in a subset of dermal fibroblasts

causes a decrease of epidermal CTNNB1 and LEF1, leading to alterations in placode number.

Later in development, dermal papilla-specific expression of *Sfrp2* alters melanocyte behavior,

causing an increase in the expression of Wnt targets and genes involved in melanogenesis. Thus,

our results show that *Sfrp2* can lead to opposing modulation (i.e., activation and inhibition) of

Wnt signaling within the same tissue. Future dissection of the downstream pathways by which

Sfrp2 acts to respectively inhibit and activate Wnt signaling during placode formation and

melanogenesis will reveal key insights into the precise mechanisms by which such opposing

effects are achieved.

It is noteworthy that elimination of *Sfrp2* led to a subtle, albeit consistent difference in both the

pattern and overall coat coloration of striped mice. This result can be explained by different

factors, including the overall effect size of *Sfrp2* as well as possible redundancy and/or

compensation with other molecules within the Wnt pathway and/or other pathways. For example,

with respect to placode formation, our bulk-level RNA-seq data from E16.5 embryos shows that

there are additional Wnt inhibitors expressed at higher levels in R1 than in R4, including known

repressors of placode formation such as *Dkk2* (Song et al., 2018). It is likely that such molecules

act in coordination with *Sfrp2* to specify patterns of placode morphogenesis and are therefore

still playing a substantial role in regulating this trait in *Sfrp2*$^{-/-}$ striped mice. Similarly, our

previous work on pigmentation showed that hair color differences in striped mice are controlled

by the interplay of melanocyte-autonomous (e.g., *Alx3*-mediated suppression of melanocyte

differentiation) and non-autonomous processes (e.g., regulation of pigment-type switching by

paracrine factors secreted from the dermal papilla, including *Asip* and *Edn3*) (Mallarino et al.,

2016; Morgan, 2014). Thus, although *Sfrp2* influences pigmentation, as our experiments show, it

is clear that modulation of hair color in striped mice is achieved through the combinatorial effect

of multiple genes (Mallarino et al., 2016).

Stripe patterns have independently evolved multiple times among rodents, and different species

display marked variation in both number of stripes and their position along the dorsoventral axis

(Johnson et al., 2019). In agreement with our mathematical simulations, the ability of external

gradients to robustly enforce simple parallel stripe patterns is well established as a generic

mechanism in the theory of pattern formation (Hiscock and Megason, 2015). At the same time,

because this mechanism works through selection of just a single spatial wavelength, it does not

account for more complex patterns, which typically feature variations over several length scales

simultaneously. Thus, while the *Sfrp2* dorsoventral gradient can account for the pattern seen in

striped mice, the coat patterns of other rodents, such as those seen in the thirteen-lined ground

squirrel (*Ictidomys tridecemlineatus*) or the zebra mouse (*Lemniscomys barbarus*), are likely

generated through modifications of this mechanism or entirely different molecular processes.

In addition to the pigment stripes found in striped mice and other taxa (Bailleul et al., 2019;

Haupaix et al., 2018; Kuriyama and Hasegawa, 2017; Murakami et al., 2018; Parichy, 2003),

parallel stripe patterns are commonly seen in other tissues during animal development. In a

companion paper, Kingsley et al report that the parallel orientation of the cartilage rings in the

developing chick airway is generated through the interplay of a Turing-like mechanism and an

orienting dorsoventral gradient of SHH. Thus, it is likely that interacting signaling gradients and

Turing systems represent a widespread mechanism for generating stripe-like patterns in multiple

tissues and shaping animals during embryogenesis.

By integrating multidisciplinary approaches, our work reveals insights into the mechanisms by

which spatial patterns are established in developing embryos. In addition, by extending *in vivo*

genome editing approaches to striped mice, a technique previously restricted to traditional

mammalian laboratory models (i.e., laboratory mouse), our work exemplifies the potential of

using emerging model species to uncover the mechanistic basis of naturally evolved phenotypic

traits and will open the door for functional studies in other mammals.

Acknowledgements

We thank members of the Mallarino lab; Princeton LAR (Chris Dmytrow, Kirsten Gerhart,

Grace Barnett, and Jamus McGuire) for help with striped mice husbandry; the LSI Genomics

Core (Wei Wang, Jennifer M. Miller, Jessica Wiggins, and Jean Arley Volmar) for help with

library preparation and sequencing; the Nikon Center of Excellence Confocal Microscopy Core

(Gary Laevsky, Sha Wang); and members of the Rivera-Perez lab (Yeonsoo Yoon and Judith

Gallant) for help with *in vivo* genome editing experiments. We also thank Eric F. Wieschaus and

Ghirish Deshpande for insights and discussion.

Author contributions

MRJ and RM conceived the project and designed experiments. MRJ performed all RNA

sequencing experiments and bulk RNA sequencing analysis. CFG-J led the scRNAseq analysis,

with support from MRJ and QN. SL performed the *in vitro* and *in vivo* genome editing in striped

mice, with help from JAR-P and SAM. MRJ and SL performed all downstream processing and

analysis of genome edited animals. MRJ and RM performed *in situ* hybridizations. PM and SYS

did the mathematical modeling. MRJ, SAM, and BB performed the phenotypic characterization

of striped mouse and laboratory mouse tissues, including immunofluorescence and histology.

MRJ and SAM performed the melanocyte cell culture experiments. CF generated the

"*Rhabdomyzed" Mus* genome and Liftover annotation. JG and AP generated the immortalized

Rhabdomys fibroblasts. MRJ and RM wrote the manuscript with input from all authors.

Funding

This project was supported by an NIH grant to RM (R35GM133758). MRJ was supported by an

NIH fellowship (F32 GM139253). SL was supported by a Presidential Postdoctoral Research

fellowship (Princeton University). BB was supported by an NIH training grant (T32GM007388).

CF was supported by an NIH fellowship (F32 GM139240-01). C.F.G-J. is partially supported by

UC Irvine Chancellor's ADVANCE Postdoctoral Fellowship Program. QN was partially

supported by an NSF grant DMS1763272 and a Simons Foundation grant (594598).

Competing interests

The authors of this publication declare that they have no competing interests.

Data and materials availability

The bulk RNA-seq and scRNA-seq reads are submitted under NCBI BioProject: XYZ.

Methods

Striped mouse husbandry

F10 descendants of wild-derived striped mice (*Rhabdomys pumilio)*, originating from Goegap

Nature Reserve, South Africa, S 29° 41.56′, E 18° 1.60′, were originally obtained from a captive

colony at the University of Zurich (Switzerland) and are maintained at Princeton University. *R.*

pumilio are kept at a 16:8 light-dark cycle and given food *ad libitum*. All experiments performed

were approved by Princeton University's Institutional Animal Care and Use Committee.

Embryo staging

To create a developmental timeseries of striped mouse embryonic development, embryos

spanning mid-gestation were collected and ordered chronologically, based on embryo size and

various morphological features including eye pigmentation, webbing of the forelimb and

hindlimb, and skull morphology (Kaufman, 1992). The stage at which epidermal hair placodes

first become visible in striped mice was considered "stage matched" to *Mus* E15.5, the stage at

which hair placodes first become visible by eye in *Mus*. Indeed, other morphological features in

striped mice at this stage agreed remarkably well with that of E15.5 laboratory mouse embryos.

This stage in striped mice was later confirmed to be E16.5 via a combination of vaginal cytology

and ultrasound imaging. Staging of all striped mouse embryos was confirmed using vaginal

cytology and ultrasound imaging.

Sample preparation for *in situ* hybridization and immunofluorescence

Pregnant females of the appropriate stage were euthanized following approved protocols.

Embryos were harvested, fixed in 4% paraformaldehyde (PFA), dehydrated in increasing

concentrations of Methanol/PBS (25%, 50%, 75%, and 100%), and stored at -20°C.

For whole-mount *in situ* hybridizations, embryos were rehydrated in decreasing concentrations

of Methanol/PBT (75%, 50%, and 25%), and washed with PBS. For section *in situ*

hybridizations and immunofluorescence, embryos were rehydrated as described above and then

incubated in 10% sucrose overnight (O/N) at 4°C, followed by a 30% sucrose O/N incubation at

4°C. Embryos were then incubated in 50% Sucrose, 50% OCT (optimal cutting temperature)

compound for 3 hours at room temperature, embedded in OCT, flash frozen, and cryosectioned

(16um thickness) using a Leica CM3050S Cryostat. Slides were frozen at -80°C until use.

Whole Mount *in situ* hybridizations

mRNA sequences for target genes were obtained either from NCBI (*Mus musculus*) or a denovo

transcriptome (*R. pumilio*) (Mallarino et al., 2016). For *R. pumilio*, we generated the following

anti-sense riboprobes: *Dkk4* (543bp), *Wif1* (563bp), *Bmp4* (619bp), *Wnt10b* (360bp), *Dkk1*

(546bp), and *Ctnnb1* (400bp). For *M. musculus*, we generated a *Dkk4* probe (550bp). All primer

sequences are reported in Table S4. All probes were sequence-confirmed using Sanger

Sequencing. Whole-mount *in situ* hybridizations were then performed using previously described

protocols(Wu and Wang, 2019). Briefly, embryos were post-fixed with 4% PFA in PBS, washed

with PBS, treated with 20ug/mL proteinase in PBT for 45 min, and incubated O/N with

riboprobes at 65°C. The following morning, probes were washed with MABT (Maleic acid,

NaCl, Tween-20) and incubated O/N with secondary anti-DIG antibody (1:2000) diluted in

MABT, 2% Boehringer Blocking Reagent, and 20% heat-treated sheep serum. After washing

several times with MABT, the signal was developed by incubating with NBT/BCIP. Once signal

had developed sufficiently, the reaction was stopped by washing several times with PBT and

fixing in 4% PFA O/N. Embryos were visualized using a SMZ18 stereo microscope (Nikon). At

least 3 embryos per probe per species were analyzed.

Bulk-level RNA sequencing

E16.5 striped mouse embryos (N = 3) were harvested from pregnant females and placed in 1x

PBS on ice. Regions R1 (dorsalmost, placode barren), R2 (dorsalmost, placode rich), R3

(ventralmost, placode barren), and R4 (ventralmost, placode rich) were micro-dissected using

morphological placodes as reference points. Regions from both sides of the midline were

combined into a single tube containing 500ul of RNALater (Invitrogen) and stored at -20°C until

RNA extraction. RNA was extracted using the RNeasy fibrous tissue mini kit (QIAGEN) per the

manufacturer's protocol. RNA-sequencing libraries were prepped using the TruSeq RNA Library

Prep kit v2 (Illumina) and sequenced on a NovaSeq 6000 (2 x 65bp, paired-end). Pairwise

differential expression analyses between the transcriptomes of dorsal skin regions R1-R4 were

performed using DeSeq2 v1.30.1 from BioConductor (https://bioconductor.org/) (Michael I

Love, Wolfgang Huber, Simon Anders, 2014) . Only genes differentially expressed with an

adjusted p-value < 0.05 were considered. For visualization of all comparisons simultaneously,

UpSet plots were generated using Upset R (Conway et al., 2017).

Modeling stripe patterns of placode formation

Model Preliminaries

We modeled the pattern forming process using a variant of the classical Gierer-Meinhardt (GM)

model of the Turing mechanism (Gierer and Meinhardt, 1972). This two-species activator-

inhibitor model is an archetypal model of stripe and spot formation in biological systems, and its

behavior with respect to pre-existing spatial gradients has been extensively studied (Van Gorder,

2021). The use of this model is further justified by its previous success in reproducing follicle

patterning in *M. musculus*(Sick et al., 2006). We use the following form of this model:

562
$$\partial_t u = D_u \partial_{xx} u + \frac{k_2 u^2}{(g + v)\left(1 + \frac{u^2}{k_3}\right)} - k_1 u$$

563
$$\partial_t v = D_v \partial_{xx} v + \frac{k_5 u^2}{\left(1 + \frac{u^2}{k_3}\right)} - k_4 v$$

564   defined on a 1D periodic domain $[-L, L]$. Note that $g \ll 1$ is a small regularizing term added to

565   reduce numerical stiffness and will be neglected in our analysis. It is convenient for our analysis

566   to shift to a non-dimensional version. We rescale our variables as $\overline{x} = \frac{x}{L}, \overline{t} = \frac{D_v t}{L^2}, \overline{u} =$

567    $\frac{k_3 u}{k_2}$ and $\overline{v} = \frac{k_5 v}{k_2}$ for dimensionless variables $\bar{x}, \bar{t}, \bar{u},$ and $\bar{v}$. Under this transformation, after

568   dropping the overbars for convenience, our model takes the form

569
$$\partial_t u = \delta^2 \partial_{xx} u + \frac{\rho\, u^2}{(\gamma + v)\left(1 + \frac{u^2}{K}\right)} - \mu u$$

570
$$\partial_t v = \partial_{xx} v + \frac{\rho\, u^2}{\left(1 + \frac{u^2}{K}\right)} - v v$$

571
$$\text{with K} = \frac{k_5^2 k_3}{k_2^2},\ \rho = \frac{k_2 L^2}{D_v},\ \mu = \frac{k_1 k_2}{k_5},\ v = \frac{k_4 k_2}{k_5},\ \gamma = \frac{k_5\, g}{k_3}\ \text{and}\ \delta^2 = \frac{D_u}{D_v}.$$

In this form, discounting diffusivity δ, there are four possible parameters which could represent

the effect of an upstream regulator such as *Sfrp2*.

Stability and Pattern Formation

For correct choice of parameters, this system of equations exhibits a Turing instability and

subsequent pattern formation. A full bifurcation analysis is outside the scope of this paper, and

similar studies have been performed for several variants of the GM system (Song et al., 2017;

Wei and Winter, 2014). However, we do highlight sufficient conditions for the system to be

unstable to spatial perturbation: in the limit $\gamma \to 0$ there must exist some integer $n > 0$ such that

one of two cases holds:

Case 1:

584
$$\mu > n^2\pi^2(1+\delta^2) + \nu \text{ and } K > \frac{\nu^2(n^2\pi^2(1+\delta^2)+\mu+\nu)}{\mu^2(-n^2\pi^2(1+\delta^2)+\mu-\nu)}$$

585   Case 2:

586
$$K > \frac{(n^2\pi^2\delta^2+\mu)\nu^2(n^2\pi^2+\nu)}{\mu^2(-n^4\pi^4\delta^2-\mu\nu+n^2\pi^2(\mu-\delta^2\nu))}, \nu < n^2\pi^2 \text{ and } \mu > \frac{n^2\pi^2\delta^2(n^2\pi^2+\nu)}{n^2\pi^2-\nu}.$$

Numerical analysis of upstream modulators

Suppose we choose some point in the Turing-unstable region of the parameter space, for

example for the values listed in Table S5. Starting from random initial conditions, a pattern of

peaks and valleys will spontaneously form as the system converges to a steady state. This steady

state is typically not very robust: pick a slightly different initial condition and peaks will form at

different places and possibly in different numbers. To robustly place peaks where we want them

to go, we proposed to take this random pattern forming state and control it by introducing a

spatially varying regulator. We assume that this regulator exists over the entire domain and

serves to modify local chemical kinetics. These kinetics are wholly controlled by the four

parameters $\mu, \nu, \rho,$ and K, so, to systematically examine possible modulations, we explore the

effect of coupling each parameter in turn to a spatially varying, temporally constant field (e.g.,

$\mu \rightarrow \mu(x) = \mu \cdot S(x)$). Modern spectral integration methods, in combination with elementary

parallelization, have made it computationally cheap to simulate equation systems like ours: it is

feasible to aggressively sample a large function space of possible prepatterns. Our basic

methodology, then, is to choose to repeatedly modulate each parameter according to different

choices of $S(x)$, leaving the other parameters fixed, and observing what patterns emerge

consequently. We ask, for each parameter, what choice of $S(x)$ guides the reaction diffusion

equations to the best fit to an empirically derived stripe spacing. This function belongs to the

following family of functions

608
$$S_{C_i}(x) = \sum_{i=1}^{12} C_i \, T_{2i}(x)$$

Here, $T_n(x)$ is the nth Chebyshev polynomial, making $S(x)$ an element of a twelve-dimensional

function space specified by the amplitudes C_i. The cap of $n = 12$ was chosen under the

assumption that the prepattern should not have a substantially finer spatial structure than the

downstream pattern. The characteristic diffusion scale of the above system is $1/\sqrt{\nu} = 0.1$, so we

chose an upper mode with a length scale somewhat smaller than this to ensure we could cover a

large set of biologically plausible $S(x)$.

All simulations were carried out using the Chebfun library for MATLAB (T.A. Driscoll, N.

Hale, L.N. Trefethen, 2014). Our spatial domain was consistently the periodic box *[-1, 1]*, with a

discretization of 128 modes and a time step of $dt = 0.01$. Simulations were integrated to a final

time $t_{end} = 30$. This value was determined numerically to be large relative to the timescale of

relaxation to the steady state, and a sample of results were run to $2t_{end}$ and $4t_{end}$ to verify

convergence. Initial conditions were generated as small random perturbations about the uniform

steady state using Chebfun's randfun function, with spatial scale $\lambda = 0.1$. Upon convergence of

each simulation, we defined a loss function estimating the relative quality of the final steady

state. A weighting function ω was chosen as a measure of stripe location:

625
$$\omega(x) = \tanh\big(1000(|x| - 0.55)\big) - \tanh\big(1000(|x| - 0.75)\big)$$

626
$$+ \tanh\big(1000(|x| - 0.85)\big)$$

627
$$\tanh\big(1000(|x| - 0.85)\big).$$

628   For a given simulation, this produced a 'coloring inside the lines' score via convolution against

629   the final activator concentration:

630
$$L_1 = - \int_{-L}^{L} u_{C_i}(x)\omega(x)dx$$

Here, $u_{C_i}(x)$ is the final steady state observed subject to a modulator specified by amplitudes

C_i. This functional value has the property that it decreases when u has high concentration in the

expected stripe regions and increases when u is present between the desired peaks. Over the

course of our investigation, we found that many potential prepatterns of varying complexity

yielded good fits to the target. To this end, we added in a second component to the loss function

as

637
$$L_2 = \frac{\epsilon_2 \sum_{i=1}^{12} |C_i| i}{\sqrt{\sum_{i=1}^{12} C_i^2}}$$

The weighting parameter ϵ_2 serves to modulate the degree to which higher spatial complexity of

the prepattern is penalized. This value must be high enough so that simple patterns are optimal,

but if too high the loss function will favor simple functions even if they yield subpar stripe

patterns. After trial and error, we found the value of 0.1 was an effective middle ground, reliably

yielding gradients which provided close fits to the target patterns. Finally, we included a third

component of the loss function, L_3, which equals 10^{10} if u takes on negative, uniform, or

exhibits NaN values and is zero otherwise. Collectively, then, we sought to find a coefficient set

C_i which minimizes

646
$$L[C_i] = L_1[C_i] + L_2[C_i] + L_3[C_i] .$$

Taking advantage of the speed and accuracy of spectral methods, we approached this

optimization problem by sampling ~40,000 sets of C_i per kinetic parameter by engaging

MATLAB's implementation of the *patternsearch* algorithm, a widely used derivative-free

optimization procedure. Starting from a set of random points in the function space, we identified

the amplitudes C_i which established a Turing pattern that best matched the experimentally

observed stripe pattern for each parameter discussed above. The gradient shown in Fig. 2E, along

with the corresponding stripe pattern, is the optimized prepattern for the parameter μ.

3' -Single-cell RNA sequencing (scRNA-seq)

Sample preparation and library generation

Embryos from *R. pumilio and M. musculus* females were harvested at the relevant stages of

pregnancy (i.e., E12.5-E15.5 for *M. musculus*; E13.5-E16.5 for *R. pumilio*). For all stages

(except *R. pumilio* E16.5, see below), a single, rectangular piece of dorsal skin, symmetrical

across the midline, was micro-dissected, using the forelimbs and hindlimbs as reference points to

maintain consistency across stages and species. For *R. pumilio* E16.5 embryos, regions R1

(dorsalmost, placode barren), R2 (dorsalmost, placode rich), and R3 (ventralmost, placode

barren) were micro-dissected using morphological placodes as markers and regions from both

sides of the midline were combined. Dorsal skins from 3 embryos were pooled (with the

exception of *M. musculus* E13, N=2) into a single tube containing 1mL 0.25% Trypsin/PBS.

Skins were minced with fine scissors and transferred to an Eppendorf tube containing 10 mL of

0.25% Trypsin/PBS. Tubes were incubated at 37°C in a hybridization oven with gentle rotation

for 15-30 minutes (timing was dependent on embryonic stage) until the tissue was well

dissociated. Primary cell suspensions were passed through a 70um filter and diluted 1:5 with 4%

Bovine calf serum (BCS)/PBS. Cells were centrifuged at 300 x g in a refrigerated centrifuge set

at 4°C for 10 minutes. Supernatant was discarded, the cell pellet was gently washed with 4%

BCS/PBS and resuspended in 100 µl of 0.1% Ultrapure Bovine Serum Albumin (BSA)/PBS

(Invitrogen). Cell number and viability was assessed using trypan exclusion and a TC20

automated cell counter (BioRad). Single cell RNA-sequencing libraries were generated using the

Chromium Single Cell 3' Reagent Kits (v2) (10x Genomics) following the manufacturer's

instructions. Libraries were sequenced on an Illumina NovaSeq 6000 (25 x 125bp, paired-end)

R. pumilio annotation

To generate gene annotations for the striped mouse genome suitable for analysis of bulk RNA-

Seq data, we transferred high-quality NCBI gene annotations from the laboratory mouse

(Mus_musculus.GRCm38.100.chr.gff3) to the *de novo R. pumilio* assembly. To accomplish this,

we used a homology-based lift-over procedure, implemented by the program LiftOff v1.4.1

using default parameters. Raw reads, genomes, and annotations have all been desposited under

BioProject: PRJNA858857 BioSample: SAMN29758252.

Generation of a "Rhabdomyzed" genome

To both facilitate comparisons of transcriptomic data between *R. pumilio* and *M. musculus*, and

to leverage the high quality gene annotations available for the latter, we generated an alternative,

reference-guided assembly for the *R. pumilio* genome against which sequencing data was

analyzed using previously described methods (Feigin et al., 2017). Briefly, genome reads for *R.*

pumilio were aligned against the GRCm38 *M. musculus* genome using bwa-mem2 v2.1.

Alignments were sorted and duplicates were removed with SortSam and MarkDuplicates from

Picard tools (http://broadinstitute.github.io/picard/). Next, read pileups were generated using

bcftools v1.11 (Li, 2011) programs mpileup, ignoring indels to preserve the reference GRCm38

coordinate system. Base calls were then generated using bcftools call's multiallelic caller.

Finally, an initial reference-guided assembly was generated by projecting striped mouse base

calls onto the reference mouse genome with bcftools consensus. This procedure was then

repeated once more using the resulting consensus sequence from the first round as the new

starting reference genome. Metrics using the *'Rhabdomyzed'* genome compared favorably (e.g.,

number of cells identified, median genes identified per cell, and percentage of genes confidently

mapped to transcriptome) to a *R. pumilio* assembly/Liftover annotation and was thus chosen for

downstream analyses.

Processing of raw single-cell RNA-seq data

Raw FASTQ reads from *R. pumilio* samples were mapped with Cell Ranger (version 3.1.0) to the

Rhabdomyzed Mus genome (see above)*:*

(Rhabdomys_to_Mus_bcftools_consensus3.fa/Mus_Musculus.GRCm38.101.gtf)

Raw FASTQ reads from *M. musculus* samples were mapped with Cell Ranger (version 3.1.0) to

the *M. musculus* mm10 reference genome (GRCm38.94.dna/GRCm38.84.gtf).

Doublet/multiplet simulation and low-quality cell filtering

Raw count matrices from *R. pumilio* and *M. musculus* obtained from alignment with the

'Rhabdomyzed' Mus genome *or* mm10 reference genome, respectively, were pre-processed and

doublet/multiplets were simulated using Single-Cell Remover of Doublets (Scrublet) (Wolock et

al., 2019) (version 0.2.1) with default parameters enabled. The doublet/multiplet score threshold

was adjusted manually to intersect the bimodal distribution of the observed transcriptomes on the

probability density histogram as suggested. Putative singlets were kept and used for downstream

query and comparative analyses if and only if they met the following user-defined collective

quality control metrics filtering criteria: a) cells with more than 350 but less than 5,000 genes

were kept; b) cells containing less than 10% total mitochondrial gene content were kept; c) cells

not identified as outliers falling outside a prediction confidence interval defined by a quadratic in

a model of a gene vs. counts plot (P-value $1e^{-3}$) were kept (Fan et al., 2016).

Anchoring, integration, and downstream analysis of single-cell RNA-seq data

We performed anchoring and integration of *R. pumilio* or *M. musculus* data sets using the Seurat

package (Version 3.2.2, R Studio version 3.6.1) (Stuart et al., 2019). In brief, species-specific

Seurat objects were created using individual, raw digitized count matrices as suggested by

developer. Objects were merged and individual gene expression matrices were normalized, and

variable genes/features identified. Data sets were anchored (n=30 dimensions) and integrated

(n=30 dimensions). The integrated object was then scaled. Significant PCs used for clustering

and neighbor-finding were identified using a combination of statistical and heuristic methods.

Neighbors and clusters were identified with dimensions specified by user and visualized using

two-dimensional t-distributed Stochastic Neighbor Embedding (tSNE) embedding. For all

analysis, R1, R2, and R3 from *R. pumilio* E16.5 embryos were randomly down sampled to equal

cell numbers before integration into a single object that was representative of E16.5 *R. pumilio*

dorsal skin.

Cell type annotation

Cell types in *R. pumilio* and *M. musculus* were defined and annotated using a core set of *bona*

fide E14.5 *M. musculus* gene biomarkers (Sennett et al., 2015) and are reported in Table S2.

Aggregate biomarker gene module scores were computed in Seurat. Aggregate biomarker gene

module scores and individual gene expression profiles were Log-normalized and visualized as

feature plots in two-dimensional tSNE embedding.

Quantification of Sfrp2$^+$ fibroblasts

To quantify the number of *Sfrp2$^+$* fibroblasts in *R. pumilio* and *M. musculus*, we randomly sub-

sampled *Sfrp2$^+$* fibroblasts (*Sfrp2*$_{normalized\ counts}$ > 0) in *R. pumilio* or *M. musculus* (n=25

iterations). To mitigate potential differences owed to sample size, we sub-sampled fibroblasts to

the lowest number of total fibroblasts in anyone comparison. *Sfrp2$^+$* fibroblasts in *R. pumilio* or

M. musculus were then normalized to total fibroblasts. Results from each iteration are plotted

with confidence intervals and their correlation was determined with Spearman's correlation.

Results are represented as the average ratio of *Sfrp2*$^+$ fibroblasts to total fibroblasts from 25

different iterations ± STDV.

Gene expression analysis

To quantify gene expression changes in *R. pumilio* fibroblasts that correlate with *Sfrp2*

expression levels, we performed three sub-samplings of fibroblasts based on *Sfrp2* UMI counts:

(1.) *Sfrp2*-high > 1.0; *Sfrp2*-low < 1, (2.) *Sfrp2*-high > 2; *Sfrp2*-low < 2, and (3.) *Sfrp2*-high > 3;

Sfrp2-low < 3. Gene expression analysis was performed in Seurat (Wilcoxon Rank Sum test) to

identify differentially expressed genes (DEGs, adjusted p-value < 0.05). Although good

agreement was observed in all sub-samplings used, only genes identified as differential in all

three sub-sampling were subjected to Gene Ontology analysis using Enrichr and significant GOs

(P_{adj} <0.05) were considered.

Latent time and RNA dynamics analysis

Putative fibroblasts from all developmental stages from *R. pumilio* or *M. musculus* were sub-

clustered and used to compute a two-dimensional Force Layout Embedding with Pegasus

(version 1.3.0). Loom files were created using the Python script *velocyto.py* (version 0.17.17;

Anaconda version 2020.07) for each individual *R. pumilio* and *M. musculus*

library/developmental stage. Loom files were combined using loompy (version 2.0.16; Anaconda

version 3.6-5.0.1; Python version 3.7.2; Spyder version 4.1.5). We computed individual region-

specific terminal states (i.e., R1, R2, or R3 in E15.5) from initial state E12.5 and estimated their

fate probabilities toward them using CellRank (version 0.2.2; Python version 3.7.0) (Lange et al.,

2022). Gene trends were calculated along distinct lineages in the identified terminal states in

latent time (set manually to 1.0).

In situ **hybridization chain reaction (HCR)**

A conserved region in the *Sfrp2* mRNA between *R. pumilio* and *M. musculus* was used to

generate 16 probe binding sequences for *in situ* hybridization chain reaction (HCR). HCR was

performed using the standard protocol for fixed frozen tissue sections available from Molecular

Instruments®. High signal within limb sections of both species was used as a positive control.

Immunofluorescence

Immuno-fluorescence was performed on tissue sections using standard procedures. Briefly,

slides were washed with 1x PBS with 0.1% Tween (PBT) and blocked with 1x PBT/3% Bovine

serum albumin (BSA) for 1 hour. Primary antibodies rabbit anti-LEF1 (Cell signaling #2230S,

1:100), rabbit anti-β-catenin (Sigma # C2206, 1:500), chicken anti-GFP (Novus Biologicals #

NB100-1614, 1:200), or rabbit anti-Krt14 (BioLegend #905301, 1:1000) were diluted in 1x

PBT/3%BSA and slides were incubated at 4°C O/N. The next morning, slides were washed

several times with 1x PBT and secondary antibodies (goat anti-mouse Alexa Fluor 488,

ThermoFisher, #A-21133) or (goat anti-mouse Alexa Fluor 546, ThermoFisher, #A-11001) were

diluted 1:500 in 1x PBT/3%BSA and slides were incubated for 1 hour at room temperature.

Slides were washed several times with 1x PBT, stained with DAPI to visualize nuclei, and

mounted for imaging. Images were taken on a Nikon A1R confocal microscope. At least 3

embryos per condition were analyzed.

Generation of *Rosa^{Sfrp2-GFP}* laboratory mice

Srfp2 cDNA was synthesized (Azenta) and cloned into pR26 CAG/T2AGFP Asc (pR26

CAG/GFP Asc was a gift from Ralf Kuehn, Addgene plasmid # 74285, and modified to contain a

T2A instead of IRES). The plasmid was inserted into the *Rosa26* locus by CRISPR-Cas9-

mediated gene targeting. Cas9 (IDT) was complexed with an sgRNA (Millipore-Sigma) having

the spacer sequence 5'- ACTCCAGTCTTTCTAGAAGA-3'. C57BL/6J zygotes were

microinjected and transferred into pseudopregnant recipients. Founders were screened for the

presence of GFP by PCR using primers GFP(F) and GFP(R)12 founders were further screened

for 5' end targeting using primers ROSA26J and SAR and for 3' end targeting using GFP(F) and

ROSA26L. All 12 were correctly targeted, 10 were chosen to confirm the presence of the cDNA

by Sanger sequencing. Primer sequences are available in Table S4.

Quantification of hair follicle number

To examine the effect of dermal *Sfrp2* over-expression on hair follicle number, *Rosa^{Sfrp2-GFP}*

females were crossed to *Dermo1-Cre* (JAX stock #008712) (Šošić et al., 2003; Yu et al., 2003)

males and embryos were collected at E17.5. Embryos were fixed, embedded, and immunostained

for GFP, as described above. GFP expression was mosaic, but a posterior dorsal region

consistently showed high levels of GFP signal across all GFP (+) embryos. Within this region,

follicle quantification was performed on equally sized areas of the skin after performing

hematoxylin-eosin (H&E) stains to reliably identify hair follicles. In total, 4 double-transgenic

and 4 control embryos were quantified. For each embryo, hair follicle number from 4 skin

sections were counted and averaged with an equal sized region of the embryo. Statistical

significance was assessed using an unpaired t-test. Statistical significance was assigned to p-

values: p<0.05(*); p<0.01(**), p<0.001(***), p<0.0001(****).

Quantitative Real Time PCR (qPCR)

Individual dorsal skin regions R1-R3 were dissected using either hair placodes (E16.5), hair

length (E19.5) or pigmentation (P0, P4) as markers. Skin sections were stored in 500ul of

RNALater (Invitrogen) and kept at -20°C until RNA extraction. RNA was extracted using the

RNeasy fibrous tissue mini kit (QIAGEN) per the manufacturer's protocol. cDNA synthesis was

carried out using qScript cDNA SuperMix (Quantabio) and qPCR for *Sfrp2* was performed using

SYBR Green qPCR Reagent (Quantabio) on a ViiA 7 Real-Time PCR machine (Applied

Biosystems). Primers used for qPCR are listed in Table S4.

Over-expression of *Sfrp2* in melanocytes

The LV-*GFP* plasmid used in this study has been described previously(Beronja et al., 2010). To

generate LV-*SFRP2-GFP*, we amplified the *Sfrp2* coding sequence from an *Sfrp2* (NM_009144)

Mouse Tagged ORF Clone (Origene CAT#: MR204070) using CloneAmp™ HiFi PCR Premix

(Takara 639298) and cloned it (in frame) into the linearized LV-*GFP* plasmid using In-Fusion®

Snap Assembly Master Mix (Takara #638947). High titer viruses were generated as previously

described (Beronja et al., 2010). Prior to transfection, MelanA cells (purchased from the

Wellcome Trust Functional Genomics Cell Bank at St. George's, University of London) were

seeded in RPMI 1640 media supplemented with penicillin (100,000 U/L), streptomycin (100

mg/L), fetal bovine serum (10%), and tetradecanoyl phorbol acetate (TPA) (200nM). At ~80%

confluency, cells were centrifuged for 30min at 37°C and media was aspirated away and

replenished with fresh media lacking antibiotics. LV-*GFP* or LV-*SFRP2-GFP* viral supernatant

was added to each well along with a 1mg/mL polybrene diluted in bovine calf serum. Plates were

gently swirled and incubated at 37°C/5% CO_2 for 30 minutes. Plates were then centrifuged at

1100 x g for 30 minutes at 37°C. Media was removed, cells were rinsed twice with PBS, and

fresh media was added. To enrich for cells with high transfection efficiency, cells with high GFP

expression were sorted by fluorescence activated cell sorting (FACS) on a BD FACSAria II (BD

Biosciences), grown to confluency, sorted one additional time, and grown to ~80% confluency.

Media was removed, cells were incubated in 500ul TriReagent (Zymo) for 5 minutes, flash

frozen in liquid nitrogen, and stored at -80°C. RNA was isolated using a direct-zol RNA

microprep kit (Zymo). cDNA synthesis was carried out using qScript cDNA SuperMix

(Quantabio) and qPCR for *Mitf, Tyr, Axin2, C-myc*, and *Ccnd1* was performed using SYBR

Green qPCR Reagent (Quantabio) on a ViiA 7 Real-Time PCR machine (Applied Biosystems),

as described above. Primers used for qPCR are listed in Table S4.

In vivo* genome editing in *R. pumilio

Generation of immortalized R. pumilio fibroblasts

A skin biopsy was obtained from the flank of an adult male *R. pumilio* dermal fibroblasts were

subsequently isolated following a previously published protocol(Aasen and Izpisúa Belmonte,

2010). In brief, hair, fat, and connective tissue were scraped away from the biopsy before

digesting O/N at 4°C in HBSS without Ca^{2+} and Mg^{2+} (Thermo Fisher) containing dispase at a

final concentration of 500 caseinolytic units/mL (Corning) and an antibiotic/antimycotic solution

final concentrations of 100 μg/mL streptomycin, 100 IU/mL penicillin, and 250 ng/mL

amphotericin B (HyClone). Following digestions, the epidermis was removed and discarded. The

dermis was cut into small pieces, moistened with Dulbecco's modified Eagle medium (Thermo

Fisher; DMEM), and pressed into grooves scored into the wells of a 6-well tissue culture dish.

The dermis was maintained in DMEM containing 10% (vol/vol) fetal bovine serum (Omega

Scientific) and 1% (vol/vol) penicillin/streptomycin (Corning) at 37°C, 5% (vol/vol) CO_2. Media

was changed every 4–5 days and cultures monitored for fibroblast growth. Once sufficient

outgrowth had occurred, the dermis was removed from the plate and the fibroblasts removed by

trypsinization (0.05% trypsin-EDTA; Gibco) for expansion into larger culture dishes.

To generate the immortalized dermal fibroblast cell line, γ-retroviral pseudoparticles containing a

transfer plasmid encoding simian virus 40 (SV40) large T antigen (Hahn et al., 2002) were

produced in HEK293T cells. Cells were cultured on poly-L-lysine−coated 10 cm plates at 37°C,

5% (vol/vol) CO_2 in 10% FBS DMEM. At ~80% confluency, Xtremegene HP DNA transfection

reagent (Roche Applied Science) was utilized per manufacturer's directions to co-transfect the

cells with 4 µg pBABE-neo-SV40 large T, a generous gift from Bob Weinberg (Addgene

plasmid #1780; http://n2t.net/addgene:1780; RRID:Addgene_1780); 4 µg of a plasmid

containing the genes for Moloney murine leukemia virus gag-pol; and 0.57 µg of a plasmid

containing the gene for the G envelope protein of vesicular stomatitis virus. Supernatants were

harvested 24, 48, and 72 hr post-transfection, stored at 4°C, and then pooled before passing

through a 0.45 µm membrane filter (Millipore). Polybrene (Sigma-Aldrich; final concentration, 4

µg/mL) and HEPES (Gibco; final concentration, 2 mM) were added to the filtered supernatants;

aliquots were prepared and at −80°C until needed. Primary dermal fibroblasts were seeded in 6-

well plates for transduction so that cell confluency was 30–40% at the time of transduction. The

cells were "spinoculated" in a centrifuge at 37°C, 931 rcf for 2 h with 2 mL of thawed, undiluted

γ-retroviral pseudoparticles per well. The cells were subsequently kept at 37°C, 5% (vol/vol)

CO_2 and the media replaced with 10% FBS DMEM 6 hours post-spinoculation. The transduced

cells were pooled once they achieved ~80% confluency in the 6-well plate and subsequently

expanded to prepare immortalized cell stocks. Cells were verified as negative for mycoplasma by

testing with the MycoAlert Mycoplasma Detection Assay kit (Lonza) per the manufacturer's

instructions.

Guide-RNA design and testing

We designed 9 CRISPR sgRNAs targeting the *R. pumilio Sfrp2* exon1 using CRISPOR

(Concordet and Haeussler, 2018). We then cloned each guide into the rAAV Nme2*Cas9* plasmid

(Edraki et al., 2019) by replacing sg*Tyr* with sg*Sfrp2* (Yoon et al., 2018). Next, we transfected

each AAV.Nme2*Cas9*.sg*Sfrp2* into immortalized striped mouse immortalized fibroblasts using

lipofectamine3000 (L300008, Thermo Fisher). ~3-4 days post transfection, we extracted

genomic DNA using the Zymo Quick DNA Miniprep Plus Kit (Zymo) and performed targeted

PCR on *Sfrp2* (primers are listed in Table S4). We then performed a T7E1 nuclease assay

(M0302, New England Biolabs) according to the manufacturer's protocol and resolved digested

products on a 1.2% agarose gel containing STBY™ safe DNA gel stain (Invitrogen) to assess

excision efficiency of each designed sgRNA.

rAAV production, in vivo transduction, and genotyping

rAAV6. Nme2*Cas9*.sg*Sfrp2* was produced, concentrated, and purified by the PNI viral core

facility at Princeton University. Female *R. pumilio* between 4 and 6 months of age were chosen

to breed with age matched males. The presence of vaginal mating plugs or sperm was designated

as days post coitum (d.p.c) 0.5. Pregnant females at d.p.c. 0.75 (N = 21) were anesthetized,

administered with analgesics, and aseptically prepared for survival surgery. To deliver rAAV, we

slowly injected 0.5-1 μL rAAV (about 1~3×10^9 genome copy (GC)) mixed with Chicago Sky

Blue dye (0.1%, Fisher Scientific, Cat # AAA1424214) into the oviduct ampulla using a glass

micropipette with tip diameter of ~10-30 μm. The injection technique in *R. pumilio* was

optimized based on previously described procedures in *M. musculus* (Sato, 2005; Yoon et al.,

2018). Operated animals were allowed to carry to term and pups were genotyped as follows: We

collected ear punch tissue from individual animals and extracted genomic DNA using a Zymo

Quick DNA Miniprep Plus Kit (Zymo). Next, we PCR amplified the genomic region containing

the guide targeting site (primers are listed on Table S4) and resolved PCR amplicons using

agarose gel electrophoresis. For F0 striped mice, PCR amplicons were further cloned into a pSC-

amp/kan vector using StrataClone PCR cloning kit (240205, Agilent Technologies). We picked

~20-50 single colonies per sample to determine the presence, composition, and frequency of

targeted mutations via Sanger sequencing.

For knocking out *Tyr* in *R. pumilio*, 29 females were injected, 43 pups were obtained, and 9 of

them carried gene edits in *Tyr*. Of these, we obtained one albino pup of homozygous knockout

(*Tyr*$^{-/-}$) (Fig.6B). 6 F0 founders were mated to wild-type animals to obtain germline transmission.

Of these 6 founders, all of them successfully produced heterozygous animals.

For knocking out *Sfrp2* in *R. pumilio*, 21 females were injected, 57 pups were obtained, and 14

of them carried gene edits in *Sfrp2*. Of these, 11 F0 founders were mated to wild-type animals to

obtain germline transmission. We obtained a total of 4 mutated alleles (2bp deletion, 13bp

deletion, 466bp deletion, and 527 bp deletion); all of which produce a frameshift. 10 out of 11

founders tested successfully produced heterozygous F1 animals. F1 heterozygous striped mice

containing the same mutated alleles were crossed to produce F2 animals. Genotypes of all F2

animals, determined by a combination of PCR, TOPO cloning, and Sanger sequencing, showed a

mendelian ratio distribution (WT: N = 30; Heterozygous: N = 65; Homozygous: N = 27). At

least two independently generated null alleles were tested in phenotypic analyses. Our

designation of $Sfrp2^{-/-}$ mice encompasses animals that either share the same mutated allele or

have a combination of two different mutated alleles.

Western blots

We collected fresh liver tissues of euthanized animals and prepared protein lysates in RIPA (50

mM Tris-HCl pH 8.0, 150 mM NaCl, 0.1% SDS, 0.5% Sodium deoxycholate, 0.1% Triton-X, 1

mM EDTA) supplemented with protease inhibitor cocktail (ab271306, Abcam). Protein

concentration was quantified by Pierce Detergent Compatible Bradford Assay (Thermal

Scientific). For SDS-PAGE, we loaded 20 μg of total protein per sample. Membranes were

probed with primary antibodies at 4℃ O/N, washed with TBST 4x, and incubated in secondary

antibody for 1 hour at room temperature. Primary antibodies: 1:100 *Sfrp2* (ab137560, Abcam,

Cambridge, UK), 1:1000 β-tubulin (mAb #86298, Cell Signaling). Secondary antibodies:

1:20000 IRDye® 800CW Goat anti-Rabbit IgG (926-32211, LI-COR Biosciences), 1:20000

IRDye 680RD Goat anti-Mouse IgG system (926-68070, LI-COR Biosciences). Membranes

were developed using an Odyssey CLx imaging system (LI-COR Biosciences).

Quantification of *Sfrp2^{-/-}* phenotypes

Stripe width: Dorsal skin from equally staged P3 striped mouse *Sfrp2^{-/-}* (N=7) or *Sfrp2^{+/+}* (N=10)

pups was dissected and flat-mount skins were prepped. Images of pinned skins were taken within

a uniform lighting box with an EIOS2000D camera (Canon) at equal magnification. To quantify

stripe width, FIJI (ImageJ) was used to draw line segments spanning each region of interest. For

each image, 10 horizontal lines were measured along the Anterior posterior axis of the skin and

an average width was calculated. For R1, R2, and R3, this was performed on each side of the

midline. To measure width of the "striped area", 10 lines were drawn connecting the most

ventral boundary of R3 on both sides and averaged. "Width (% of striped area)" was calculated

by dividing the average width of each region (i.e., R1, R2, R3, and midline) by the average width

of the stripe area. Statistical significance was assessed using unpaired t-tests. Statistical

significance was assigned to p-values: $p < 0.05$(*); $p < 0.01$(**), $p < 0.001$(***), $p < 0.0001$(****).

Pigmentation: A single image of 6 *R. pumilio* adults (3 *Sfrp2^{+/+}*, 3 *Sfrp2^{-/-}*) was taken within a

uniform lighting box with an EIOS2000D camera (Canon). To ensure image was true to color,

color correction was performed on the image using a Pixel Perfect color checker and Adobe

Lightroom software. To quantify pigment differences across animals, a consistently sized region

of interest (ROI) was drawn within each stripe region and mean gray value within the ROI was

measured using Adobe Photoshop. For R1-R3, mean gray values were measured in 3 subregions

within each stripe region, on each side of the midline (6 ROIs total) and averaged. For the

midline, mean gray values were measured in 5 subregions within the midline and averaged.

Statistical significance was assessed using unpaired t-tests. Statistical significance was assigned

to p-values: $p < 0.05$(*); $p < 0.01$(**), $p < 0.001$(***), $p < 0.0001$(****).

Excel Tables

Table S1: Differentially expressed genes between Region 1-4 of E16.5 striped mice skin.

(Table_S1.xls)

Table S3: Differentially expressed genes between Sfrp2^{high} and Sfrp2^{low} fibroblast populations.

(Table_S3.xls)

Multimedia Files

Supplemental Movie 1: Changes in regulator distribution alter the simulated stripe pattern.

(Supplemental_Movie_1.mp4)

References

Aasen, T., and Izpisúa Belmonte, J.C. (2010). Isolation and cultivation of human keratinocytes
from skin or plucked hair for the generation of induced pluripotent stem cells. Nat. Protoc. *5*,
371–382.
Andl, T., Reddy, S.T., Gaddapara, T., and Millar, S.E. (2002). WNT signals are required for the
initiation of hair follicle development. Dev. Cell *2*, 643–653.
Bailleul, R., Curantz, C., Desmarquet-Trin Dinh, C., Hidalgo, M., Touboul, J., and Manceau, M.
(2019). Symmetry breaking in the embryonic skin triggers directional and sequential plumage
patterning. PLoS Biol. *17*, e3000448.
Beronja, S., Livshits, G., Williams, S., and Fuchs, E. (2010). Rapid functional dissection of
genetic networks via tissue-specific transduction and RNAi in mouse embryos. Nat. Med. *16*,
821–827.
Caro, T. (2005). The Adaptive Significance of Coloration in Mammals. Bioscience *55*, 125–136.
Caro, T., and Mallarino, R. (2020). Coloration in Mammals. Trends Ecol. Evol. *35*, 357–366.
Concordet, J.-P., and Haeussler, M. (2018). CRISPOR: intuitive guide selection for
CRISPR/Cas9 genome editing experiments and screens. Nucleic Acids Res. *46*, W242–W245.
Conway, J.R., Lex, A., and Gehlenborg, N. (2017). UpSetR: an R package for the visualization
of intersecting sets and their properties. Bioinformatics *33*, 2938–2940.
Edraki, A., Mir, A., Ibraheim, R., Gainetdinov, I., Yoon, Y., Song, C.-Q., Cao, Y., Gallant, J.,
Xue, W., Rivera-Pérez, J.A., et al. (2019). A Compact, High-Accuracy Cas9 with a Dinucleotide
PAM for In Vivo Genome Editing. Mol. Cell *73*, 714-726.e4.

Eizirik, E., David, V.A., Buckley-Beason, V., Roelke, M.E., Schäffer, A.A., Hannah, S.S., Narfström, K., O'Brien, S.J., and Menotti-Raymond, M. (2010). Defining and Mapping Mammalian Coat Pattern Genes: Multiple Genomic Regions Implicated in Domestic Cat Stripes and Spots. Genetics *184*, 267–275.

Enshell-Seijffers, D., Lindon, C., Wu, E., Taketo, M.M., and Morgan, B.A. (2010). β-Catenin activity in the dermal papilla of the hair follicle regulates pigment-type switching. Proceedings of the National Academy of Sciences *107*, 21564–21569.

Fan, J., Salathia, N., Liu, R., Kaeser, G.E., Yung, Y.C., Herman, J.L., Kaper, F., Fan, J.-B., Zhang, K., Chun, J., et al. (2016). Characterizing transcriptional heterogeneity through pathway and gene set overdispersion analysis. Nat. Methods *13*, 241–244.

Feigin, C.Y., Newton, A.H., Doronina, L., Schmitz, J., Hipsley, C.A., Mitchell, K.J., Gower, G., Llamas, B., Soubrier, J., Heider, T.N., et al. (2017). Genome of the Tasmanian tiger provides insights into the evolution and demography of an extinct marsupial carnivore. Nature Ecology & Evolution *2*, 182–192.

Fuchs, E. (2007). Scratching the surface of skin development. Nature *445*, 834–842.

Fuchs, E., and Raghavan, S. (2002). Getting under the skin of epidermal morphogenesis. Nat. Rev. Genet. *3*, 199–209.

Gat, U., DasGupta, R., Degenstein, L., and Fuchs, E. (1998). De Novo hair follicle morphogenesis and hair tumors in mice expressing a truncated beta-catenin in skin. Cell *95*, 605–614.

Gierer, A., and Meinhardt, H. (1972). A theory of biological pattern formation. Kybernetik *12*, 30–39.

Gupta, K., Levinsohn, J., Linderman, G., Chen, D., Sun, T.Y., Dong, D., Taketo, M.M., Bosenberg, M., Kluger, Y., Choate, K., et al. (2019). Single-Cell Analysis Reveals a Hair Follicle Dermal Niche Molecular Differentiation Trajectory that Begins Prior to Morphogenesis. Dev. Cell *48*, 17-31.e6.

Hahn, W.C., Dessain, S.K., Brooks, M.W., King, J.E., Elenbaas, B., Sabatini, D.M., DeCaprio, J.A., and Weinberg, R.A. (2002). Enumeration of the simian virus 40 early region elements necessary for human cell transformation. Mol. Cell. Biol. *22*, 2111–2123.

Hardy, M.H. (1992). The secret life of the hair follicle. Trends Genet. *8*, 55–61.

Haupaix, N., and Manceau, M. (2020). The embryonic origin of periodic color patterns. Dev. Biol. *460*, 70–76.

Haupaix, N., Curantz, C., Bailleul, R., Beck, S., Robic, A., and Manceau, M. (2018). The periodic coloration in birds forms through a prepattern of somite origin. Science *361*. https://doi.org/10.1126/science.aar4777.

He, T.C., Sparks, A.B., Rago, C., Hermeking, H., Zawel, L., da Costa, L.T., Morin, P.J., Vogelstein, B., and Kinzler, K.W. (1998). Identification of c-MYC as a target of the APC pathway. Science *281*, 1509–1512.

Hiscock, T.W., and Megason, S.G. (2015). Orientation of Turing-like Patterns by Morphogen Gradients and Tissue Anisotropies. Cell Syst *1*, 408–416.

Iozumi, K., Hoganson, G.E., Pennella, R., Everett, M.A., and Fuller, B.B. (1993). Role of tyrosinase as the determinant of pigmentation in cultured human melanocytes. J. Invest. Dermatol. *100*, 806–811.

Jho, E.-H., Zhang, T., Domon, C., Joo, C.-K., Freund, J.-N., and Costantini, F. (2002). Wnt/beta-catenin/Tcf signaling induces the transcription of Axin2, a negative regulator of the signaling pathway. Mol. Cell. Biol. *22*, 1172–1183.

Johansson, J.A., and Headon, D.J. (2014). Regionalisation of the skin. Semin. Cell Dev. Biol. *25–26*, 3–10.

Johnson, M.R., Barsh, G.S., and Mallarino, R. (2019). Periodic patterns in Rodentia: Development and evolution. Exp. Dermatol. *28*, 509–513.

Kaelin, C.B., and Barsh, G.S. (2013). Genetics of pigmentation in dogs and cats. Annu Rev Anim Biosci *1*, 125–156.

Kaelin, C.B., Xu, X., Hong, L.Z., David, V.A., McGowan, K.A., Schmidt-Küntzel, A., Roelke, M.E., Pino, J., Pontius, J., Cooper, G.M., et al. (2012). Specifying and sustaining pigmentation patterns in domestic and wild cats. Science *337*, 1536–1541.

Kaelin, C.B., McGowan, K.A., and Barsh, G.S. (2021). Developmental genetics of color pattern establishment in cats. Nat. Commun. *12*, 5127.

Kaufman, M.H. (1992). The Atlas of Mouse Development (London, England: Academic Press Limited).

Keller, S.H., Jena, S.G., Yamazaki, Y., and Lim, B. (2020). Regulation of spatiotemporal limits of developmental gene expression via enhancer grammar. Proceedings of the National Academy of Sciences *117*, 15096–15103.

Kim, M., Han, J.H., Kim, J.-H., Park, T.J., and Kang, H.Y. (2016). Secreted Frizzled-Related Protein 2 (sFRP2) Functions as a Melanogenic Stimulator; the Role of sFRP2 in UV-Induced Hyperpigmentary Disorders. J. Invest. Dermatol. *136*, 236–244.

Kondo, S. (2017). An updated kernel-based Turing model for studying the mechanisms of biological pattern formation. J. Theor. Biol. *414*, 120–127.

Kondo, S., and Miura, T. (2010). Reaction-Diffusion Model as a Framework for Understanding Biological Pattern Formation. Science *329*, 1616–1620.

Kuriyama, T., and Hasegawa, M. (2017). Embryonic developmental process governing the conspicuousness of body stripes and blue tail coloration in the lizardPlestiodon latiscutatus. Evol. Dev. *19*, 29–39.

Lange, M., Bergen, V., Klein, M., Setty, M., Reuter, B., Bakhti, M., Lickert, H., Ansari, M., Schniering, J., Schiller, H.B., et al. (2022). CellRank for directed single-cell fate mapping. Nat. Methods *19*, 159–170.

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics *27*, 2987–2993.

Liang, C.-J., Wang, Z.-W., Chang, Y.-W., Lee, K.-C., Lin, W.-H., and Lee, J.-L. (2019). SFRPs Are Biphasic Modulators of Wnt-Signaling-Elicited Cancer Stem Cell Properties beyond Extracellular Control. Cell Rep. *28*, 1511-1525.e5.

Lin, H., Angeli, M., Chung, K.J., Ejimadu, C., Rosa, A.R., and Lee, T. (2016). sFRP2 activates Wnt/β-catenin signaling in cardiac fibroblasts: differential roles in cell growth, energy metabolism, and extracellular matrix remodeling. Am. J. Physiol. Cell Physiol. *311*, C710–C719.

van Loon, K., Huijbers, E.J.M., and Griffioen, A.W. (2021). Secreted frizzled-related protein 2: a key player in noncanonical Wnt signaling and tumor angiogenesis. Cancer Metastasis Rev. *40*, 191–203.

Mallarino, R., Henegar, C., Mirasierra, M., Manceau, M., Schradin, C., Vallejo, M., Beronja, S., Barsh, G.S., and Hoekstra, H.E. (2016). Developmental mechanisms of stripe patterns in rodents. Nature *539*, 518–523.

Mallarino, R., Pillay, N., Hoekstra, H.E., and Schradin, C. (2018). African striped mice. Curr. Biol. *28*, R299–R301.

Michael I Love, Wolfgang Huber, Simon Anders (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. Genome Biology.
Millar, S.E. (2002). Molecular mechanisms regulating hair follicle development. J. Invest.
Dermatol. *118*, 216–225.
Mills, M.G., and Patterson, L.B. (2009). Not just black and white: pigment pattern development
and evolution in vertebrates. Semin. Cell Dev. Biol. *20*, 72–81.
Morgan, B.A. (2014). The dermal papilla: an instructive niche for epithelial stem and progenitor
cells in development and regeneration of the hair follicle. Cold Spring Harb. Perspect. Med. *4*,
a015180.
Murakami, A., Hasegawa, M., and Kuriyama, T. (2018). Developmental mechanisms of
longitudinal stripes in the Japanese four-lined snake. J. Morphol. *279*, 27–36.
Parichy, D.M. (2003). Pigment patterns: fish in stripes and spots. Curr. Biol. *13*, R947-50.
Patterson, L.B., and Parichy, D.M. (2019). Zebrafish Pigment Pattern Formation: Insights into
the Development and Evolution of Adult Form. Annu. Rev. Genet. *53*, 505–530.
Rezza, A., Wang, Z., Sennett, R., Qiao, W., Wang, D., Heitman, N., Mok, K.W., Clavel, C., Yi,
R., Zandstra, P., et al. (2016). Signaling Networks among Stem Cell Precursors, Transit-
Amplifying Progenitors, and their Niche in Developing Hair Follicles. Cell Rep. *14*, 3001–3018.
Sato, M. (2005). Intraoviductal introduction of plasmid DNA and subsequent electroporation for
efficient in vivo gene transfer to murine oviductal epithelium. Mol. Reprod. Dev. *71*, 321–330.
Sennett, R., Wang, Z., Rezza, A., Grisanti, L., Roitershtein, N., Sicchio, C., Mok, K.W.,
Heitman, N.J., Clavel, C., Ma'ayan, A., et al. (2015). An Integrated Transcriptome Atlas of
Embryonic Hair Follicle Progenitors, Their Niche, and the Developing Skin. Dev. Cell *34*, 577–
591.
Shtutman, M., Zhurinsky, J., Simcha, I., Albanese, C., D'Amico, M., Pestell, R., and Ben-Ze'ev,
A. (1999). The cyclin D1 gene is a target of the beta-catenin/LEF-1 pathway. Proc. Natl. Acad.
Sci. U. S. A. *96*, 5522–5527.
Sick, S., Reinker, S., Timmer, J., and Schlake, T. (2006). WNT and DKK determine hair follicle
spacing through a reaction-diffusion mechanism. Science *314*, 1447–1450.
Song, Y., Yang, R., and Sun, G. (2017). Pattern dynamics in a Gierer–Meinhardt model with a
saturating term. Appl. Math. Model. *46*, 476–491.
Song, Y., Boncompagni, A.C., Kim, S.-S., Gochnauer, H.R., Zhang, Y., Loots, G.G., Wu, D., Li,
Y., Xu, M., and Millar, S.E. (2018). Regional Control of Hairless versus Hair-Bearing Skin by
Dkk2. Cell Rep. *25*, 2981-2991.e3.
Šošić, D., Richardson, J.A., Yu, K., Ornitz, D.M., and Olson, E.N. (2003). Twist Regulates
Cytokine Gene Expression through a Negative Feedback Loop that Represses NF-κB Activity.
Cell *112*, 169–180.
Steingrímsson, E., Copeland, N.G., and Jenkins, N.A. (2004). Melanocytes and the
microphthalmia transcription factor network. Annu. Rev. Genet. *38*, 365–411.
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data.
Cell *177*, 1888-1902.e21.
T.A. Driscoll, N. Hale, L.N. Trefethen (2014). Chebfun Guide (Oxford: Pafnuty Publications).
Tsai, S.-Y., Sennett, R., Rezza, A., Clavel, C., Grisanti, L., Zemla, R., Najam, S., and Rendl, M.
(2014). Wnt/β-catenin signaling in dermal condensates is required for hair follicle formation.
Dev. Biol. *385*, 179–188.

Turing, A.M. (1990). The chemical basis of morphogenesis. 1953. Bull. Math. Biol. *52*, 153–
197; discussion 119-52.
Van Gorder, R.A. (2021). Pattern formation from spatially heterogeneous reaction-diffusion
systems. Philos. Trans. A Math. Phys. Eng. Sci. *379*, 20210001.
Vittadello, S.T., Leyshon, T., Schnoerr, D., and Stumpf, M.P.H. (2021). Turing pattern design
principles and their robustness. Philosophical Transactions of the Royal Society A:
Mathematical, Physical and Engineering Sciences *379*, 20200272.
Wei, J., and Winter, M. (2014). The Gierer-Meinhardt System with Saturation. In Mathematical
Aspects of Pattern Formation in Biological Systems, J. Wei, and M. Winter, eds. (London:
Springer London), pp. 249–261.
Wolock, S.L., Lopez, R., and Klein, A.M. (2019). Scrublet: Computational Identification of Cell
Doublets in Single-Cell Transcriptomic Data. Cell Syst *8*, 281-291.e9.
Wu, J., and Wang, X. (2019). Whole-Mount In Situ Hybridization of Mouse Embryos Using
DIG-Labeled RNA Probes. Methods Mol. Biol. *1922*, 151–159.
Yoon, Y., Wang, D., Tai, P.W.L., Riley, J., Gao, G., and Rivera-Pérez, J.A. (2018). Streamlined
ex vivo and in vivo genome editing in mouse embryos using recombinant adeno-associated
viruses. Nat. Commun. *9*, 1–12.
Yu, K., Xu, J., Liu, Z., Sosic, D., Shao, J., Olson, E.N., Towler, D.A., and Ornitz, D.M. (2003).
Conditional inactivation of FGF receptor 2 reveals an essential role for FGF signaling in the
regulation of osteoblast function and bone growth. Development *130*, 3063–3074.