Detection of isoforms and genomic alterations by high-throughput full-length single-cell RNA sequencing in ovarian cancer

Arthur Dondi[1,2,], Ulrike Lischetti[1,3,*], Francis Jacob[3], Franziska Singer[2,4], Nico Borgsmüller[1,2], Ricardo Coelho[3], Tumor Profiler Consortium, Viola Heinzelmann-Schwarz[3], Christian Beisel[1,*], Niko Beerenwinkel[1,2,*]

These authors contributed equally: Arthur Dondi, Ulrike Lischetti
* Corresponding authors

[1] ETH Zurich, Department of Biosystems Science and Engineering, Mattenstrasse 26, 4058 Basel, Switzerland
[2] SIB Swiss Institute of Bioinformatics, Mattenstrasse 26, 4058 Basel, Switzerland
[3] Current affiliation: University Hospital Basel and University of Basel, Ovarian Cancer Research, Department of Biomedicine, Hebelstrasse 20, 4031 Basel, Switzerland
[4] ETH Zurich, NEXUS Personalized Health Technologies, Wagistrasse 18, 8952 Schlieren, Switzerland

Keywords: full-length single-cell RNA sequencing, long-read PacBio sequencing, transcript concatenation, isoforms, mutations, gene fusions, ovarian cancer

Abstract

Understanding the complex background of cancer requires genotype-phenotype information in single-cell resolution. Here, we perform long-read single-cell RNA sequencing (scRNA-seq) on clinical samples from three ovarian cancer patients presenting with omental metastasis and increase the PacBio sequencing depth to 12,000 reads per cell. Our approach captures 152,000 isoforms, of which over 52,000 are novel. Isoform-level analysis accounting for non-coding isoforms reveals 20% overestimation of protein-coding gene expression on average. We also detect cell type-specific isoform and poly-adenylation site usage in tumor and mesothelial cells, and find that mesothelial cells transition into cancer-associated fibroblasts in the metastasis, partly through the TGF-β/miR-29/Collagen axis. Furthermore, we identify gene fusions, including an experimentally validated *IGF2BP2::TESPA1* fusion, which is misclassified as high *TESPA1* expression in matched short-read data, and call mutations confirmed by targeted NGS cancer gene panel results. With these findings, we envision long-read scRNA-seq to become increasingly relevant in oncology and personalized medicine.

Introduction

Cancer is a complex disease characterized by genomic and transcriptomic alterations[1] that drive multiple tumor-promoting capabilities or hallmarks[2]. Among others, these alterations include point mutations, insertions and deletions (indels), and gene fusions on the genomic level, and splice isoforms on the transcriptomic level. Their detection offers great potential for personalized oncology as they can serve as direct therapeutic targets[3,4] or potential neoantigens informing on the immunogenicity of the tumor[5]. Gene fusions arising from large-scale genomic rearrangements, for example, play an oncogenic role in a variety of tumor types[6], and are successfully used as therapeutic targets[7,8]. Like mutations[9] and copy number variations[10], fusion rates can vary widely across cancer types, and gene fusions are thought to be drivers in 16.5% of cancer cases, and even the only driver in more than 1%[11]. Furthermore, out-of-frame gene fusions are more immunogenic than mutations and indels, making them an ideal target for immunotherapies and cancer vaccines[12,13]. On the transcriptomic level, alternative splicing is a major mechanism for the diversification of a cell's transcriptome and proteome[14] and can impact all hallmarks of tumorigenesis. It also presents a fairly novel non-genomic source of potential neoantigens[15]. In breast and ovarian cancer, 68% of samples had at least one isoform with novel exon-exon junction (neojunction) detected in proteomic data[16].

The complexity of cancer further extends to intra-tumor heterogeneity[17] and its intricate interplay with the tumor microenvironment (TME)[18]. Ultimately, to fully decipher functional tumor heterogeneity and its effect on the TME, single-cell resolution providing both phenotype and genotype information is required. Single-cell RNA sequencing (scRNA-seq) is now widely used for the phenotypic dissection of heterogeneous tissues. It can be divided into short-read, high-throughput technologies allowing for gene expression quantification and long-read, low-throughput technologies that cover full-length isoforms[19]. Up to now, short- and long-read methods had to be used in parallel to combine the advantages of each technology. The long-read sequencing field is rapidly expanding[20], with novel scRNA-seq methods being constantly developed and improved on Nanopore[21,22] and PacBio[23–27] platforms. So far, long-read RNA-seq has however only been applied on the bulk level in the field of oncology[25,28,29]. High-quality, high-throughput, long-read scRNA-seq has the potential to provide isoform-level cell type-specific readouts and capture tumor-specific genomic alterations. With near ubiquitous p53 mutations and defective DNA repair pathways causing frequent non-recurrent gene fusions, high-grade serous ovarian cancer (HGSOC) is an ideal candidate to investigate these alterations[10,30,31].

Here, we used high-quality, high-throughput long-read scRNA-seq to capture cell type-specific genomic and transcriptomic alterations in clinical cancer patients. We applied both short-read and long-read scRNA-seq to five samples from three HGSOC patients, comprising 2,571 cells, and generated the PacBio scRNA-seq dataset with the deepest coverage to date. We were

able to identify over 150,000 isoforms, of which a third were novel, as well as novel cell type-
specific isoforms. Isoform-level analysis revealed that, on average, 20% of the protein-coding
gene expression was non-coding, leading to an overestimation of the protein expression. By
combining differential isoform and poly-adenylation site usage analysis between cells from the
metastatic TME and distal omental biopsies, we found evidence that in omental metastases,
mesothelial cells transition into CAFs, partly through the TGF-β/miR-29/Collagen axis.
Additionally, we discovered dysregulations in the insulin-like growth factor (IGF) network in
tumor cells on the genomic and transcriptomic levels. Thereby, we demonstrated that scRNA-
seq can capture genomic alterations accurately, including cancer- and patient-specific
germline and somatic mutations in genes such as *TP53*, as well as gene fusions, including a
novel *IGF2BP2::TESPA1* fusion.

Results

Long-read scRNA-seq creates a catalog of isoforms in ovarian cancer patient-derived tissue samples

We generated short-read and long-read scRNA-seq data from five omentum biopsy samples (**Extended Data Tables 1 and 2**) from three HGSOC patients. Three samples were derived from HGSOC omental metastases and two from matching distal tumor-free omental tissues (**Fig. 1a**). To generate long-reads, we opted for the PacBio platform for its generation of high-fidelity (HiFi) reads through circular consensus sequencing (CCS). To overcome its limitations in sequencing output and optimize for longer library length, we 1) removed template-switch oligo artifacts that can account for up to 50% of reads through biotin enrichment, 2) concatenated multiple cDNA molecules per CCS read, and 3) sequenced on the PacBio Sequel II platform (2-4 SMRT 8M cells per sample, **Methods**). This allowed the generation of a total of 212 Mio HiFi reads in 2,571 cells, which, after demultiplexing, deduplication, and artifact removal, resulted in 30.7 Mio unique molecular identifiers (UMIs), for an average of 12k UMIs per cell. (**Extended Data Table 1**). There was a mean of four cDNA molecules concatenated per sequencing read overall, and cDNA length was similar across samples (**Extended Data Fig. 1a,b**). Artifact removal filtered 51% of the reads, and it included the removal of intra-priming (63%), non-canonical isoforms (36%), and reverse transcriptase switching (1%)[32] (**Extended Data Fig. 1c**). It must be emphasized that those artifact reads emerge from the single-cell library preparation and are also present in short-read data, where they cannot be filtered and are hence accounted for as valid reads.

The long-read dataset revealed 152,546 isoforms, each associated with at least three UMIs. We classified the isoforms according to the SQANTI classification[32] and calculated their proportions (**Methods**, **Fig. 1b**,**c**): full splice match (FSM) - isoforms already in the GENCODE database (32.8%), incomplete splice matches (ISM) - isoforms corresponding to shorter versions of the FSM (35.1%), novel in catalog (NIC) - isoforms presenting combinations of known splice donors and acceptors (15.9%), and novel not in catalog (NNC) - isoforms harboring at least one unknown splice site, or neojunction (14.4%). Novel isoforms (classes NIC and NNC) accounted for 30% of the isoforms, and 11% of the total reads in all samples, while FSM accounted for 33% of the isoforms and 80% of the reads (**Fig. 1c,d**), indicating that high coverage is required for the reliable detection of new, low abundant, transcripts.

To evaluate the structural integrity of all isoforms, we compared their 5' end to the FANTOM5 CAGE database[33] and their 3' end to the PolyASite database[34] (**Fig. 1e**). More than 82% of the NIC and 74% of NNC isoforms could be validated on 3' and 5' ends, similarly to FSM. As expected, fewer ISM isoforms were found to be complete (42%): they are either incompletely sequenced isoforms missing their 5' end (30%) or the result of early 3' termination (55%).

FSM, NIC, and NNC had overall better 3' and 5' validation than the full-length tagged isoforms
in the GENCODE database (**Fig. 1e**). Only the 'Matched Annotation from NCBI and EMBL-
EBI' (MANE[35]) containing curated representative transcripts cross-validated between the
GENCODE and RefSeq database had a better 3' and 5' validation of 95%. A total of 52,884
novel isoforms were complete (NIC+NNC), of which 40,046 were confirmed as valid novel
isoforms by GENCODE (corresponding to 17% of the current GENCODE v36 database), and
3,695 were extended versions of existing isoforms. Isoforms that were not confirmed were
mainly either "partially redundant with existing transcripts", or "overlapping with multiple loci".
Finally, we assessed the biotypes of our newly discovered isoforms, indicative of their
presumed functional categorization. We found that 42% are protein-coding, more than the
36% of protein-coding isoforms found in the GENCODE database (230k entries) (**Fig. 1f,g**).
This demonstrates the ability of concatenated long-read sequencing to generate high-yield,
high-quality data and discover novel isoforms with enhanced annotation.

Long-read sequencing allows for short-read-independent cell type identification

Next, through comparison to short-read data, we assessed the ability of long-read sequencing
to cluster cells and to identify cell types. We generated short- and long-read gene count
matrices and removed non-protein-coding, ribosomal, and mitochondrial genes. After filtering,
we obtained 16.5 Mio unique long-reads associated with 12,757 genes, and 26.3 Mio unique
short-reads associated with 13,122 genes (**Extended Data Table 2**). The short- and long-
read datasets were of similar sequencing depth with a median of 4,930 and 2,750 UMIs per
cell, respectively (average 10,235 and 6,413 UMIs, **Extended Data Fig. 2a**). Long-read data
contained slightly fewer detected genes, and genes detected in both datasets overlapped by
86.4% (**Extended Data Fig. 2b,c**). Paradoxically, the genes detected were overall shorter in
long-reads than short-reads, likely due to the concatenation step (**Extended Data Fig. 2d**).
169
We first identified cell types independently per cell, using cell type marker gene lists (**Methods,**
Extended Data Fig. 3a). We compared short- and long-read data and found that both data
types identified cell types with very similar percentages, namely HGSOC (13,4% in short-read
vs 13.6% in long-read data), mesothelial cells (20.2 vs 20.5%), fibroblasts (10.7 vs 10.7%), T
cells (38.7 vs 38.5%), myeloid cells (14.6 vs 14.9%), B cells (1.4 vs 1.1%), and endothelial
cells (1.1 vs 1.4%) (**Extended Data Fig. 3b**). We then projected short-read gene, long-read
gene, and long-read isoform expression onto 2-dimensional embeddings using UMAP[36] (**Fig.**
2a). We manually clustered cell types based on the embeddings and calculated the Jaccard
distance between clusters. Cell clusters based on short- and long-reads were very similar,
with a Jaccard distance >94% for all cell types except B-cells, where the Jaccard distance was
>75% (**Fig. 2b**). Furthermore, Jaccard similarity analysis between cell type clusters and
attributed cell type labels were analogous between short- and long-read data, with a better

prediction of B cells and endothelial cells for long-reads (**Extended Data Fig. 3c**). Except for
tumor cells, cell types did not show patient-specific clustering (**Extended Data Fig. 3d**).

Long-read sequencing captures germline and somatic mutations and identifies increased neojunctions in tumor cells

Next, we assessed the potential of long-read data for mutation detection, and used somatic
mutations to further validate the cell type annotation. Germline mutations are expected in all
cell types, whereas somatic mutations should be present only in tumor cells. As reference, we
used mutations called from a panel covering 324 genes on patient-matched bulk DNA samples
(**Methods**). We identified germline variants in 48 cells belonging to all cell types from the distal
omentum and tumor sites (**Fig. 2c, Supplementary Table 1**). Somatic mutations were called
in 34 cells, all in the cell cluster annotated as tumor cells (**Fig. 2d**). In 20 of those cells, *TP53*
was found mutated (**Supplementary Table 1**). Thus, high-fidelity, long-read data can be
leveraged for both germline and somatic mutation calling.

Cell type-specific isoform expression analysis (overall and per gene) revealed increased
isoform and transcript expression in HGSOC cells compared to other cell types (**Extended
Data Fig. 4a-c**). This difference does however not translate into mean UMIs per isoform, as
isoforms expressed in cancer cells harbor fewer UMIs than in mesothelial cells (**Extended
Data Fig. 4d**). This means that cancer cells express more low-abundant isoforms suggesting
wider isoform diversity and broader cellular functions, and controls. Isoform class distribution
between cell types revealed a higher fraction of novel isoforms and neojunctions (NNC) in
tumor cells (**Fig. 2e**). Looking into unique isoform expression in the different cell types, we
found that cancer cells contained more than 8% (9,476) of cell type-specific isoforms, and
35% of those contained neojunctions (NNC) (**Methods, Extended Data Fig. 4e**). This was an
increase of 2.3-10.6 times compared to the other cell types. At the cellular level, 0.5% of the
cancer-specific isoforms were also unique to a single cell, which is between 3-6 times the
percentage of unique isoforms in other cell types, and 50% contained neojunctions (**Extended
Data Fig. 4f**). Those rare isoforms were difficult to detect with previous methods, hence their
novelty. Taken together, cancer cells expressed at least twice as many unique isoforms than
other cell types, indicating an increased transcriptomic diversification and supporting previous
findings of cancer-specific neojunction expression in bulk data[16].

Tumor microenvironment shows epithelial-to-mesenchymal transition through TGFβ-driven miR-29 downregulation

In the subsequent analysis, we compared stromal fibroblasts and mesothelial cells derived
from metastasis (TME) and matched tumor-free (distal) omentum. In distal samples,
fibroblasts and mesothelial cells formed distinct clusters in both short- and long-read data (**Fig.
3a left, Extended Data Fig. 5a**). In metastatic samples, however, TME fibroblasts and

mesothelial cells formed a bridge in the UMAP embedding, suggesting that these cells might
undergo a cell state transition. To test this hypothesis, we conducted gene set enrichment
analysis and found that the epithelial-to-mesenchymal transition (EMT) pathway was enriched
in TME compared to distal mesothelial and fibroblast cells (**Fig. 3a, right**). Similarly, collagen
fibril organization and extracellular matrix (ECM) pathways were enriched in TME cells,
indicating a reprogramming of the TME cells during metastasis formation (**Extended Data Fig.**
5b). Additionally, we compared the alternative poly-adenylation (APA) (**Methods**). Among the
2,876 genes tested for APA, the isoforms of 26 genes in TME mesothelial cells exhibited
significant 3'UTR lengthening compared to distal mesothelial cells, while 13 genes showed
shortened 3'UTRs (**Fig. 3b**). Collagen-encoding genes *COL1A2*, *COL3A1*, *COL5A2*, and
COL6A1 were similarly lengthened and up-regulated in TME cells (**Fig. 3c**), with *COL1A2*
having the highest effect size ($P_{corr}=3.42 \times 10^{-67}$, 53% change).
230
As 3'UTR lengthening is usually associated with a decrease in expression due to microRNA
(miRNA) silencing[37], the increased usage of lengthened 3'UTR in TME mesothelial cells
suggests that distal cells may express a distinct set of miRNAs not present in TME cells.
Collagen-encoding genes are known to be regulated by the miR-29 family in fibroblasts[38].
Thus, we used miRDB[39] to predict gene targets for silencing by miR-29a/b/c (**Supplementary**
Table 2). Among the 26 isoforms with lengthened 3'UTR, 9 were predicted targets of miR-
29a, almost all described as EMT actors (collagens[40–43], *KDM6B*[44], *TNFAIP3*[45], *FKBP5*[46] and
RND3[47]). In contrast, none of the 13 shortened 3'UTR isoforms were predicted to be miR-29
targets (**Fig. 3b**). Furthermore, we compared gene expression of TME and distal mesothelial
cells (**Supplementary Table 3**), and genes with lengthened 3'UTR isoforms predicted to be
silenced by miR-29 were significantly overexpressed ($P = 1.35 \times 10^{-3}$) in TME mesothelial cells
compared to the ones not predicted to be silenced by miR-29a (**Fig. 3d**). Mesothelial TME
cells also differentially expressed miR-29 targets which are major ECM genes, such as
collagen gene *COL1A1* (fold-change=9.0, $P_{corr}=1.72 \times 10^{-124}$), *MMP2*[48] (fold-change=4.1,
$P_{corr}=1.51 \times 10^{-30}$), and *LOX*[49] (fold-change=10.6, $P_{corr}=6.33 \times 10^{-51}$), which is also lengthened
($P_{corr}=7.07 \times 10^{-2}$, 77% change). Overall, ECM-related genes known to be targeted by miR-29
were upregulated in TME cells compared to distal cells (**Fig. 3e**), supporting the hypothesis
that the observed EMT was potentially linked with the miR-29 downregulation. When
comparing differentially expressed isoforms in TME mesothelial cells to distal cells, *COL1A1*
was also the gene with the highest change in relative isoform abundance amongst all its
isoforms ($P_{corr}=6.34 \times 10^{-49}$, 86% usage change, **Methods**) (**Fig. 3f**). In TME mesothelial cells,
the *COL1A1* canonical 3' poly-adenylation site was used, whereas distal cells used a
premature poly-adenylation site, leading to the formation of truncated isoforms (**Fig. 3f**). When
incorporating only protein-coding isoforms and removing the truncated isoforms from the
analysis, the gene expression fold-change increased from 9 to 62-folds ($P_{corr}=3.15 \times 10^{-183}$).
This overexpression of canonical *COL1A1* in the TME can be explained by the absence of
miR-29 silencing, as previously described[50].

The miR-29 family is known to be an EMT inhibitor[50]. Its silencing through the TGFβ pathway

correlates with the upregulation of ECM-encoding genes, including multiple collagens, as

reported in the present study. The main TGFβ gene *TGFB1* was found to be enriched in TME

mesothelial cells (fold-change=1.4, P_{corr}=2.32x10^{-2}). Furthermore, in distal mesothelial cells,

38% of *TGFB1* isoforms comprise an alternative 3' exon, leading to aberrant protein

expression (**Extended Data Fig. 5c**), while the canonical protein-coding *TGFB1* isoform

ENST00000221930.6 is overexpressed in TME cells (fold-change=2.3, P_{corr}=6.59x10^{-9}). miR-

29 is also regulated through the expression of non-coding RNAs that act as molecular

sponges, directly binding to miR-29 and, therefore, leading to the overexpression of their

targets. The TGFβ-regulated long non-coding RNA H19, which enhances carboplatin

resistance in HGSOC, has been reported to promote EMT through the *H19*/miR-29b/*COL1A1*

axis[51–53] and was found to be overexpressed in the TME mesothelial cells (fold-change=4.6,

P_{corr}=3.46x10^{-34}). Circular RNAs have also been described as miR-29 sponges, notably

circMYLK, and circKRT7 in HGSOC. circMYLK and circKRT7 originate from *MYLK* and *KRT7*,

respectively, which are both significantly overexpressed in TME mesothelial cells (fold-

change=4.1, P_{corr}=6.97x10^{-96} and fold-change=4.6, P_{corr}=2.18x10^{-13})[54,55]. Similarly, TME

mesothelial cells expressed the endogenous isoform of *GSN* (*cGSN*), while distal cells only

expressed the secreted isoform (pGSN)[56] (**Extended Data Fig. 5d**). *cGSN* has been shown

to be under TGF-β control in breast cancer and to increase EMT marker expression[57]. In

conclusion, our findings strongly support that, in omental metastases, the mesothelial cells

transition into cancer-associated fibroblasts (CAFs), partly through the TGF-β/miR-

29/Collagen axis.

Differential isoform usage in cancer reveals changes in biotypes

After comparing cells from the TME with distal cells, we investigated which isoforms, biotypes,

and poly-adenylation sites were differentially used between cancer and all distal cells. HGSOC

cells expressed isoforms differentially with a change in relative isoform abundance of more

than 20% in 960 genes (15.1%), compared to all distal cells (6,353 genes tested in total, **Fig.**

4a, **Supplementary Table 4, Methods)**. In 36% of those 960 genes, the highest expressed

isoform biotype changed between conditions (**Fig. 4b)**. In 32% of instances, there was a

transition from a protein-coding to a non-protein-coding isoform, and in 17% of cases, cancer

cells expressed a protein-coding isoform while distal cells expressed a non-protein-coding

isoform. Only 39 genes (0.6%) had an isoform switch with a change in relative isoform

abundance of more than 50%. 59% of these switched isoforms demonstrated a biotype

transition (49% of protein-coding to non-protein-coding transition, **Fig. 4c)**, and in 33% of

cases, cancer cells expressed a protein-coding isoform while distal cells expressed a non-

protein-coding isoform. Additionally, in cancer, distal, and TME cells, on average 20-21% of

the expression in protein-coding genes was non-coding (**Extended Data Fig. 6a)**, and 13-

14% of protein-coding genes had more than 50% of non-coding expression. This means that,

on average, using only gene-level information to estimate protein expression (as done in short-read data) will lead to an overestimation of 20%.

The ten genes with the statistically most significant switches were *IGF1*, *TPM2*, *NCALD*, *VAMP8-VAMP5*, *EXOSC7*, *ICAM3*, *CERK*, *OBSL1*, *GSN*, and *RTN1* (**Supplementary Table 4, Methods**). In *IGF1*, cancer cells across all patients predominantly used the second exon of the gene as their transcription start site (secreted isoform, *Class II*), whereas non-cancerous cells primarily used the first exon (endogenous isoform, *Class I*)[58] (**Fig. 4d**). On the contrary, similarly to TME mesothelial cells, cancer cells expressed the endogenous isoform of *GSN* (*cGSN*), while distal cells only expressed the secreted isoform (*pGSN*)[56] (**Extended Data Fig. 5d**). In *RTN1*, distal cells expressed the isoform *RTN1-A* and *RTN1-B*, while Patient 3's cancer cells expressed *RTN1-C*, an isoform known to bind to the anti-apoptotic protein Bcl-xL and reduce its activity[59,60] (**Extended Data Fig. 6b**). In the tropomyosin gene *TPM2*, which is involved in TGF-β-induced EMT, cancer cells differentially expressed exon 6b (isoform *TPM2.3*, expressed in epithelial cells[61]) and the alternative 3'UTR exon 9a (**Extended Data Fig. 6c**). In *VAMP5*, the overexpressed isoform in HGSOC cells was a *VAMP8-VAMP5* read-through gene, i.e., a novel gene formed of two adjacent genes, previously described in human prostate adenocarcinoma[62] (**Extended Data Fig. 6d**). HGSOC cells expressed almost no wild-type (wt) *VAMP5* but had a significantly higher *VAMP8* expression than distal cells (P_{corr}=1.0x10^{-15}), indicating that this read-through gene was under transcriptional control of *VAMP8*. With a short-read 3' capture method, this *VAMP8-VAMP5* expression cannot be distinguished from the wt *VAMP5* expression. For *NCALD* and *OBSL1*, only cancer cells expressed canonical protein-coding isoforms, while other cells expressed short non-coding isoforms (**Extended Data Fig. 6e,f**). For *CERK*, by contrast, Patient 2's HGSOC cells strongly expressed a novel isoform leading to a shortened protein (**Extended Data Fig. 6g**). Finally, in *ICAM3,* cancer cells mainly expressed a short protein-coding isoform, while distal cells (mainly T cells) expressed the canonical isoform. More characterization of those isoforms will be necessary in the future to explore the biological implications linked to their expression (**Extended Data Fig. 6h**).

Although isoforms differentially expressed in cancer cells were similar among patients, there was one significant case of patient-specific expression. For *OAS1*, Patient 3 predominantly expressed isoform *p42*, while Patients 1 and 2 exhibited a balanced distribution of isoforms *p42* and *p46*. The *p42* and *p46* expressions are known to be allele-specific, caused by the rs10774671 SNP, a splice acceptor A/G variation. However, this cannot explain the different expression levels as all patients have both the A and G alleles (**Fig. 4e**). Given that isoform *p42* is more susceptible to Nonsense-Mediated mRNA Decay (NMD) and possesses diminished enzymatic activity[63], the observed differences could potentially indicate a diminished OAS1 activity in Patient 3. Whether *OAS1* has an impact on ovarian cancer is still to be elucidated.

When testing for differential APA between HGSOC and distal cells, we found shortened 3'UTR
in 85 and lengthened 3'UTR in 203 genes (n=4758) (**Fig. 4f**). There was a notable trend toward
lengthening of the 3'UTR in cancer cells (P=5.59x10^{-20}), with *COL1A2* emerging once more
as the most prominent finding (P$_{corr}$=7.48x10^{-47}, 61% change) (**Fig. 3b-c**, **Fig. 4f**). Expression
levels remained consistent between genes featuring either shortened or lengthened 3'UTRs.
Furthermore, neither miRNA profiles nor canonical pathways exhibited an overlap exceeding
20% with either the lengthened or shortened gene sets (**Methods**).

Long-read sequencing captures gene fusions and identifies an *IGF2BP2::TESPA1* fusion that was misidentified in short-read data

To detect fusion transcripts, we aligned long-reads to the reference genome and filtered for
reads split-aligned across multiple genes. We then ranked fusion transcripts with counts
across all cells of more than 10 UMIs (**Supplementary Table 5**). Out of the 34 detected fusion
entries, 21 were genes fused with mitochondrial ribosomal RNA (*mt-rRNA1-2*) and ubiquitous
among all cell types, 11 isoforms were *IGF2BP2*::*TESPA1* fusions specific to Patient 2, one
was a cancer cell-specific *CBLC* (chr8:43.064.215) fusion to a long non-coding RNA (lncRNA)
expressed in Patient 3, and one was a cancer cell-specific fusion of *FNTA* with a lncRNA
expressed in Patient 1. The ubiquitous *mt-rRNA* fusions were likely template-switching
artifacts from the library preparation, as *rRNA* makes up to 80% of RNA in cells[64].
IGF2BP2::TESPA1 was a highly expressed fusion event in Patient 2: 2,174 long-reads
mapped to both *IGF2BP2* (Chr3) and *TESPA1* (Chr12). The gene fusion consisted of 5'
located exons 1-4 of *IGF2BP2,* corresponding to 112 amino acids (aa) and including the RNA
recognition motif 1 (RRM1) and half of the RRM2 domain*,* linked to the terminal *TESPA1* 3'
untranslated region (UTR) exon, encoding 69 aa as in-frame fusion and including no known
domains (**Fig. 5a**). In total, the gene fusion encoded 181 aa, compared to 599 aa of wt
IGF2BP2 and 521 aa of wt *TESPA1* (**Fig. 5b**). 98.9% of fusion reads were found in HGSOC
cells and the fusion was detected in 86.8% of Patient 2's cancer cells, making it a highly cancer
cell- and patient-specific fusion event (**Fig. 5c**). Cancer cells lacking the gene fusion had lower
overall UMI counts, suggesting low coverage as a possible reason for the absence of the gene
fusion (**Fig. 5d**).

We next investigated the footprint of the gene fusion in the short-read data. The *TESPA1* gene
was expressed in T cells, as well as in HGSOC cells, where its expression values were
elevated. High expression was exclusive to Patient 2 HGSOC cells and colocalized with
IGF2BP2 expression (**Fig. 5e,f**). *TESPA1* was the highest differentially expressed gene in
cancer cells compared to non-cancer cells in Patient 2 (P$_{corr}$<1.17x10^{-14}). Next, we re-aligned
Patient 2's short-reads to a custom reference including the *IGF2BP2::TESPA1* transcriptomic
breakpoint as well as wt *TESPA1* and wt *IGF2BP2* junctions (**Extended Data Fig. 7a,**

Methods). Out of the 994 reads mapping to the custom reference, 93% preferentially aligned to *IGF2BP2::TESPA1* (99.8% of those were from HGSOC cells). This means that, when given the option, reads previously aligning to *IGF2BP2* or *TESPA1* are preferentially mapping to the fusion reference, and the reported overexpression of *TESPA1* in short reads is likely an *IGF2BP2::TESPA1* expression. Furthermore, reads covering the *TESPA1* 3' UTR region harbored three heterozygous single nucleotide polymorphisms (hSNPs): chr12:54.950.144 A>T (rs1047039), chr12:54.950.240 G>A (rs1801876), and chr12:54.950.349 C>G (rs2171497). In long-reads, wt *TESPA1* was either triple-mutated or not mutated at all, indicating two different alleles. All fusion long-reads, however, were triple-mutated, indicating a genomic origin and monoallelic expression of the fusion (**Fig. 5g**). In short reads, the three loci were mutated in nearly all reads, supporting the hypothesis that the observed *TESPA1* expression represents almost completely *IGF2BP2::TESPA1* expression and that it has a genomic origin.

Genomic breakpoint validation of the *IGF2BP2::TESPA1* fusion

To validate that the *IGF2BP2::TESPA1* gene fusion is the result of genomic rearrangements, both bulk and single-cell DNA sequencing (scDNA-seq) data from matched omental metastasis was used to query the genomic breakpoint. A putative genomic breakpoint was first found in the RNA data. Two long-read fusions were mapped to intronic regions of *IGF2BP2* and *TESPA1* (**Extended Data Fig. 7b**), pinpointing the location of the breakpoint at position chr3:185,694,020-chr12:54,960,603. Subsequent genotyping PCR on genomic DNA extracted from patient-matched tissue samples using *IGF2BP2::TESPA1*, wt *IGF2BP2*, and wt *TESPA1* primer pairs flanking the genomic breakpoint confirmed the presence of *IGF2BP2::TESPA1* in Patient 2 in 3 out of 4 tested samples (**Fig. 6a, Methods**). In contrast and as expected, the fusion was not found in Patient 1.

To assess whether the fusion was exclusive to cancer cells, we further investigated scDNA-seq data from Patient 2. For the identification of cancer cells, we inferred the scDNA-seq copy number profiles of all cells. We successfully identified two distinct clones within the pool of 162 cells. These clones encompassed a cancer clone designated as "Subclone 0" and a presumably non-cancer clone without copy number alterations labeled as "Subclone 1" (**Fig. 6b**). We next aligned the scDNA-seq data to a custom reference covering the breakpoint (**Methods)** and only found cancer reads mapping to the fusion breakpoint (P=0.032), while a mixture of reads from cancer and non-cancer cells mapped to wt *IGF2BP2* and wt *TESPA1* (P=0.78 and P=1.00, respectively) (**Fig. 6c**). Thus, genotyping PCR of bulk extracted DNA and scDNA-seq data confirmed the genomic fusion breakpoint in the intronic region detected in long-read scRNA-seq data. scDNA-seq also confirmed that the *IGF2BP2::TESPA1* fusion was cancer-cell specific, as suggested by long-read scRNA-seq data.

IGF2BP2 was also overexpressed in cancer cells from Patient 2 compared to other patients

on both RNA and protein levels (**Extended Data Fig. 8a-c**). In Patient 2, there was an elevated

copy number observed within the genomic region encompassing *IGF2BP2* (**Fig. 6b**).

Therefore, the presence of a fusion allele on one allele does not seem to hinder the

transcription of the wt *IGF2BP2* allele. Coherent with the high IGF2BP2 protein levels, *IGF2*

RNA, which is bound by the wt IGF2BP2 protein, is also largely overexpressed in Patient 2

cancer cells compared to other patients (**Extended Data Fig. 8d**). This could indicate that the

fusion happened partly due to accessible chromatin. In the ovarian cancer TCGA RNA

dataset, the expression levels of exons surrounding breakpoints of *IGF2BP2* and *TESPA1* did

not change (**Extended Data Fig. 9a,b**), and the overall expression of the genes did not

correlate in any patient, suggesting that we detected an uncommon, patient-specific fusion

(**Extended Data Fig. 9c**).

Discussion

The detection of genomic alterations like mutations[65,66] and gene fusions[67,68], alongside isoform expression and usage[15] at the single-cell scale offers valuable insights into cancer development, metastasis formation, the role of the tumor microenvironment (TME), potential drug targets, and therapy responses[69]. Here, we applied PacBio HiFi high-throughput long-read RNA-seq on five omental metastases and matching tumor-free samples from chemo-naive HGSOC patients to detect and quantify all of these alterations.

Until now, a combination of single-cell short- and long-read sequencing was necessary to identify cell-specific isoforms: the increased depth of short-read sequencing allowed for cell typing based on gene expression, while long-read sequencing was used for isoform identification[23]. Leveraging multiple strategies to maximize the long-read sequencing output, we achieved a 50-fold increased sequencing depth compared to the first long-read PacBio scRNA-seq study[23], allowing for short read-comparable cell type identification. As a result, future studies employing comparable or higher long-read capacities will no longer depend on parallel short-read sequencing, leading to cost and labor savings.

Taking advantage of the long-read technology, our study provides evidence that cancer cells may induce EMT in TME mesothelial cells. In omental metastases, it has been shown that HGSOC-secreted TGF-β triggers EMT and converts TME cells, including mesothelial cells, into CAFs, which in turn may favor tumor cell adhesion, invasion, and proliferation[70–72]. Our findings provide further evidence of this phenomenon and reveal that the process might partly be controlled through the TGF-β/miR-29/Collagen axis: secreted and endogenous TGF-β downregulates miR-29 expression, thus increasing the expression of its targets, including collagens[73]. Coincidentally, Han and colleagues[48] recently showed that omental CAF-derived exosomes from HGSOC patients contained significantly lower miR-29c than normal fibroblasts. Furthermore, the reduced levels of miR-29 have been demonstrated to play a role in the development of cisplatin resistance by upregulating collagen expression[74].

In addition to TGF-β, HGSOC cells also promote EMT and CAF activation through the secretion of other growth factors such as insulin-like growth factors (*IGF*s)[75]. In our data, HGSOC cells revealed a profoundly modified IGF system among all patients with a drastic switch from endogenous *Class I* to secreted *Class II IGF1* isoform, *IGF2* overexpression, and a highly expressed *IGF2BP2::TESPA1* gene fusion in one patient. Secreted IGF1 (Class II) and IGF2 activate EMT through IGF1R, by triggering an uncontrolled wound healing response[76]. The *IGF* gene family also promotes cancer growth, survival, proliferation, and drug resistance through signaling via *PI3K-AKT* or *MAPK*, and is a known clinical target in ovarian cancer[77].

Additionally, we demonstrated the potential of the technology in terms of coverage and
sequencing accuracy to detect mutations and gene fusions. In particular, in one patient, the
novel fusion *IGF2BP2::TESPA1* was highly overexpressed compared to wt *IGF2BP2* (~10x
more) and *TESPA1* (~150x more). *IGF2BP2* is known to be regulated via 3'UTR miRNA
silencing[78], however, the *IGF2BP2::TESPA1* fusion has the unregulated 3'UTR of *TESPA1*,
which could explain its overexpression. *TESPA1* is normally expressed in T cells[79] and long-
read data confirmed T cell-specific wt *TESPA1* expression. Short read data however
erroneously reported *TESPA1* as the most differentially expressed gene in cancer cells,
resulting from 3' end capture of the fusion transcripts. This highlights that short-read scRNA-
seq data fails to distinguish between gene and fusion expression, potentially leading to wrong
biological conclusions.

In addition to their canonical mRNA isoforms, numerous protein-coding genes express non-
coding RNA isoforms, which are wrongly accounted for as protein-coding on the gene
expression level and can serve as miRNA sponges and competing endogenous RNAs,
especially in EMT and metastasis[80,81]. In our data, we found that 20% of the protein-coding
gene expression was non-coding and, in the genes exhibiting a significant isoform switch
between cancer and healthy cells, nearly half (49%) transitioned from a protein-coding to a
non-coding isoform. Furthermore, 51% of the UMIs were composed of intra-priming and non-
canonical artifacts before filtering, and their detection was only possible through isoform
classification. Overall, our findings highlight the need for an isoform-specific quantification to
accurately assess protein-coding gene expression[82] and narrow the RNA-protein gap[83–86].
Additionally, a better isoform characterization is needed to understand their biological
implications, as we partly demonstrated with *IGF1 Class I/II*, *cGSN, TPM2.3, RTN1-C,* and
OAS1 p42.

Although the achieved sequencing depth allowed for short-read-independent cell typing and
clustering, a further increased depth is needed to capture low-abundance transcripts. For
example, we did not obtain sufficient reads to retrieve and characterize the T cell receptor
repertoire. This is consistent with a long-read scRNA-seq study in blood lymphocytes that
reported a 3.6-fold lower pairing rate for T cell receptors than the higher abundant B cell
receptors from plasmablasts[87]. Using a single-cell long-read targeted sequencing approach,
Byrne and colleagues were able to achieve a T cell receptor alpha and beta pairing rate of
50% in an ovarian cancer patient sample, showing the potential to extract immune repertoire
information from long-read data with increased sequencing depth[88]. Enrichment for low
abundant transcripts for long-read sequencing or depletion of mitochondrial and ribosomal
RNA[89] represent interesting avenues forward.

Taken together, we demonstrate that long-read sequencing provides a more complete picture
of cancer-specific changes. These findings highlight the manifold advantages and new

opportunities that this technology provides to the field of precision oncology, opening the
premise of personalized drug prediction and neoantigen detection for cancer vaccines[90,91].

Materials and Methods

Omentum patient cohort

The use of material for research purposes was approved by the corresponding cantonal ethic commissions (EKNZ: 2017–01900, to V.H.S.), and informed consent was obtained for all human primary material. Tissue samples were immediately collected from the theater and transferred on ice to the Department of Biomedicine of the University Hospital Basel for tissue dissociation.

Sample processing

Fresh omentum and omental HGSOC tumor metastasis biopsy samples were cut into small pieces and dissociated in digestion solution (1 mg/mL collagenase/Dispase [Sigma cat. no. 10269638001], 1 unit/mL DNase I [NEB, cat. no. M0303] and 10% fetal bovine serum (FBS) in DMEM [Sigma, cat. no. D8437-500mL]) for 30 min at 37°C. To focus on the non-adipose cell fraction, adipocytes were separated by centrifugation and the cell pellet was collected. Red blood cell lysis (RBC) was performed using MACS red blood lysis solution (cat. no. 130-094-183). Then, the cell pellet was resuspended into MACS dead cell removal microbeads (cat. no. 130-090-101) and loaded into the AutoMACS separator to remove dead cells. After counting cell numbers, cells were resuspended in PBS with 1% BSA and transferred to the Genomics Facility Basel. The cell suspension was again filtered and cell number and viability were assessed on a Cellometer K2 Image Cytometer (Nexcelom Bioscience, cat. no. Cellometer K2) using ViaStain AOPI Staining Solution (Nexcelom Bioscience, cat. no. CS2-0106-5mL) and PD100 cell counting slides (Nexcelom Bioscience, cat. no. CHT4-PD100-003). For samples with viability below 70% and when cell numbers allowed (>10^5 cells total), apoptotic and dead cells were removed by immunomagnetic cell separation using the Annexin Dead Cell Removal Kit (StemCell Technologies, cat. no. 17899) and EasySep Magnet (StemCell Technologies, cat. no. 18000). If the cell pellet appeared still red, additional RBC lysis was performed. Cells were washed with a resuspension buffer (PBS with 0.04% BSA), spun down, and resuspended in a resuspension buffer. Finally, cells were again counted and their viability was determined. The cell concentration was set according to 10x Genomics protocols (700-1,200 cells/µL).

10x Genomics single-cell capture and short-read sequencing

Cell suspensions were loaded and processed using the 10x Genomics Chromium platform with the 3P v3.1 kit on the 10x Genomics Chromium Single Cell Controller (10x Genomics, cat. no. PN-120263) according to the manufacturer's instructions. 500 or 1,000 cells were targeted per lane. The quality of cDNA traces and GEX libraries were profiled on a 5200 Fragment Analyzer (Agilent Technologies).

Paired-end sequencing was performed on the Illumina NovaSeq platform (100 cycles, 380pm loading concentration with 1% addition of PhiX) at recommended sequencing depth (20,000-50,000 reads/cell).

Long-read library preparation and PacBio sequencing

To increase long-read PacBio sequencing throughput, we followed the strategy of cDNA concatenation of the HIT-scISOseq protocol[24] with the modification of two rounds of biotin-PCR in order to further reduce template-switch oligo (TSO) artifacts from the data. Full protocol details:

cDNA amplification and biotin-enrichment

15 ng of each patient's cDNA library were amplified using the KAPA HiFi HotStart Uracil+ ReadyMix 2x (Kapa Biosystems, cat. no. KK2801) with 0.5 µM final concentration of custom-primers (Integrated DNA Technologies, HPLC purified). Primers contained overhang sequences adapted from Hebelstrup *et al.*[92] with a single deocxyuredine (dU) residue at a 10 nt distance from the 5' terminus enabling USER enzyme digestion and creating single-stranded overhangs. Generated PCR fragments thus contain a single dU residue per DNA strand. The forward primer was specific to the 10x Genomics partial Read 1 sequence and contained a biotin modification allowing for biotin enrichment of amplified full-length cDNA molecules. The reverse primer was specific to the 10x Genomics partial TSO sequence.

Forward Primer: /5Biosg/AGGTCTTAA/ideoxyU/CTACACGACGCCTTCCGATCT

Reverse Primer: ATTAAGACC/ideoxyU/AAGCAGTGGTATCAACGCAGAG

The PCR was run according to the manufacturer's instruction with two cycles at an annealing temperature of 63°C followed by 7 cycles at an annealing temperature of 67°C; annealing time was 30 seconds. Extension was performed at 72°C for 90 seconds. PCR products were purified at 0.6X SPRIselect bead cleanup (Beckman Coulter, cat. no. B23318) according to the manufacturer's instructions and eluted in 22 µL EB buffer (Qiagen, cat. no. 19086). DNA concentrations were measured using the Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific, cat. no. Q32854), which were in the range of 1.5 µg per sample. cDNA traces were additionally evaluated on a 5200 Fragment Analyzer System (Agilent Technologies) using the HS NGS Fragment Kit, 1-6000 bp (Agilent, cat. no. DNF-474-0500). Full-length cDNAs were enriched through capture on 5 µL streptavidin-coated M-280 dynabeads using the Dynabeads™ kilobaseBINDER™ Kit (Invitrogen, cat. no. 60101), thus depleting TSO-TSO artifacts. Washed Dynabeads containing the DNA-complexes were directly resuspended in 20 µL USER reaction buffer containing 10 µL StickTogether DNA Ligase Buffer 2x (NEB, cat. no. B0535S), 1.5 µL USER Enzyme (NEB, cat. no. M5505S) and 8.5 µL Nuclease-free water (Invitrogen, AM9939) and incubated in a thermocycler at 37°C for 20 min and held at 10°C (no annealing). This created a nick at the deoxyuracil site forming palindrome overhangs and releasing the biotin-bound DNA molecules from the beads. Beads were removed by magnetic

separation and the supernatant with the biotin-released cleaved PCR products was subjected
to a 0.6X SPRIselect cleanup step. Approximately 100 ng of purified product per sample were
split into two aliquots and subjected to a second PCR amplification step with 6 cycles using
an annealing temperature of 67°C. Reactions were pooled, purified by 0.6X SPRIselect
cleanup, and quality checked on both Qubit and Fragment Analyzer. Total DNA yield was
between 5-8 µg, which were subjected to a second round of streptavidin-purification using 10
µL of beads.

Transcript ligation

Beads were incubated in 19 µL USER reaction buffer at 37°C for 20 min for USER digestion
and 25°C for 17 min for overhang annealing. Beads were then removed by magnetic
separation and the supernatant was transferred to a new PCR tube. 1 µL of T4 DNA ligase
high-concentration (2,000,000, units/mL, NEB, cat. no. M0202T) was added, mixed and
incubated at 10°C for >24hrs and heat inactivated at 65°C for 10 min. To efficiently deplete
any non-ligated transcripts, 0.38X SPRIselect cleanup was performed, eluted in 20 µL EB
buffer and traces were evaluated on the Fragment Analyzer using the HS Large Fragment kit
(Agilent Technologies, cat. no. DNF-492-0500) at 1:5 dilutions. Ligation products were 8-11kb
long; average yield was 100 ng per sample.

End repair/dA tailing, adapter ligation, and PCR amplification

To enable PCR amplification of the ligated construct, the NEBNext Ultra II DNA Library Prep
Kit for Illumina was followed (NEB, cat. no. E7645S) using total DNA yield as input material.
2.5 µL of 5 µM dT overhang adapter (Roche, cat. no. KK8727) was used for the End Prep
reaction. Adapter-ligated libraries were purified by 0.39X SPRIselect cleanup, eluted in 22 µL
EB buffer and products were evaluated by HS Large Fragment kit. Total yield of around 40 ng
was split in two and PCR amplified using 2X KAPA HiFi Hot-Start ReadyMix (Roche, cat. no.
KK2602) and KAPA Library Amplification Primer Mix (10X concentration, Roche, cat. no.
KK2623), 10 µL library input each with 11 cycles and 9 min extension time. Following a 0.38X
SPRIselect cleanup and elution in 48 µL EB buffer, products were evaluated on a large
fragment gel revealing an average fragment length of libraries of 4.6 kb and an average total
of 1.1 µg DNA. To increase the total yield to 2 µg DNA required for SMRTbell library
preparation of a product with 5 kb amplicon size, the PCR was repeated with three additional
cycles and 5 min extension time. After 0.4X SPRI cleanup and Fragment Analyzer inspection,
the final yield was 2 µg per library.

PacBio SMRTbell library preparation

The SMRTbell Express Template Kit (PacBio, cat. no. 100-938-900) was used following
manufacturer's instructions for DNA damage repair, end repair/dA-tailing and ligation of a
hairpin adapter (double amount used). Final purification of the SMRTbell template was

performed by 0.42X SPRIselect cleanup and elution in 43 µL EB buffer. Exonuclease
treatment was performed by addition of 5 µL of NEBbuffer1 (NEB, cat. no. B7001S) and 1 µL
of each Exonuclease I (NEB, cat. no. M0293S) and Exonuclease III (NEB, cat. no. M0206S)
bringing the total volume to 50 µL per reaction. Enzyme treatment was performed at 37°C for
60 min. After SPRIselect cleanup, products were quantified on a large fragment gel at 1:30
dilution. Final yield was approximately 650 ng per sample, a sufficient amount for long-read
sequencing.

615 PacBio Sequel II sequencing

616 Libraries were sequenced on the PacBio Sequel II platform with the SMRT cell 8M. Omentum
617 metastasis and tumor-free omentum were run on three and two 8M cells, respectively.

618 Genotyping PCR on genomic DNA

Genomic DNA was extracted from homogenized tumor tissue samples (n=8 samples matching
sampling time, Basel Ovarian Biobank) derived from patients using the DNeasy Blood &
Tissue kit (QIAGEN, cat. no. 69504). Isolated DNA underwent QC using Nanodrop and Qubit
measurements. Genotyping PCR on genomic DNA was performed using the MyTaq DNA
Polymerase system (Bioline, cat. no. BIO-21105). Briefly, MyTaq reaction buffer and MyTaq
DNA polymerase were pipetted together with 200 nM primer (Sigma Aldrich) pairs
(gPCR_IGF2BP2-TESPA1_Bp_F 5'-CCT GCT TTG AGG AGG GGA GGG A-3' &
gPCR_IGF2BP2-TESPA1_Bp_R 5'-ACT GAG GAC AAT GCT ACG CAA GA-3';
gPCR_TESPA1_F 5'-CCT GCT TTG AGG AGG GGA GGG A-3' & gPCR_TESPA1_R 5'-TGA
GAA CTG CTG TTC CAG GAG ACA-3'; gPCR_IGF2BP2_F 5'-ACA CTG GAC CCA TGC
TTG AGC T-3' & gPCR_IGF2BP2_R 5'-GCG TGC TAT GAA CAC TCC AGG CC-3'), and 50
ng genomic DNA (gDNA). PCR conditions were 1 cycle at 94°C for 5 min followed by 35 cycles
at 95°C for 20 sec, 58°C for 15 sec, 72°C for 1 min, and finished with 1 cycle at 72°C for 5
min. Amplicons were visualized on a 1.2 % agarose gel together with DNA ladder.
633

634 Immunofluorescence

Formalin-fixed and paraffin-embedded tissue samples were obtained from the Basel Ovarian
Biobank matching with patients 1 and 2 on sampling time and site. Briefly, samples were
deparaffinized and immersed for 10 min in a 10 mM sodium citrate buffer at pH 6.0 (Sigma
Aldrich, cat. no. C9999) at 95°C for antigen retrieval. Samples were permeabilized in 0.25%
(v/v) Triton™ X-100 (Roth, cat. no. 3051.3) in PBS (Sigma Aldrich, cat. no. D8537-500M) for
5 min and blocked in 5% FBS (Sigma Aldrich, cat. no. F7524-500ml), 0.1% Triton™ X-100,
1% BSA (Roth, Fraction V, cat. no. 8076.4) in PBS for 1 hour. The following antibodies were
used for this study: rabbit IGF2BP2 (C-terminal-specific, ThermoFisher Scientific, cat. no.
MA5-42874), EpCAM (Cell Signaling Technologies, cat. no. 5488S,) and goat anti-rabbit Alexa
Fluor® 647 (Cell Signaling Technology, cat. no. 4414). Slides were mounted using ProLong®

Gold Antifade Reagent with DAPI (Cell Signaling Technology, cat. no. 8961). Images were
acquired using a Nikon spinning-disk confocal microscope (Nikon CSU-W1 spinning-disk
confocal microscope) and processed with Fiji. Cell quantification was performed using an in-
house developed QuPath script for cell detection and annotations.

Single-cell DNA-sequencing

650 Cell suspensions were loaded and processed using the 10x Genomics Chromium platform
651 with the single-cell CNV kit on the 10x Genomics Chromium Single Cell Controller (10x
652 Genomics, cat. no. PN-120263) according to the manufacturer's instructions. Paired-end
653 sequencing was performed on the Illumina NovaSeq platform (100 cycles, 380pm loading
654 concentration with 1% addition of PhiX) at recommended sequencing depth.

Data Analysis

Short-read data analysis

Preprocessing

658 Raw reads were mapped to the GRCh38 reference genome using 10x Genomics Cell Ranger
659 3.1.0 to infer read counts per gene per cell. We performed index-hopping removal using a
660 method developed by Griffiths *et al.*[93].

10x Genomics read data processing, normalization, and visualization.

Expression data of each sample was analyzed using the scAmpi workflow[94]. In brief, UMI
counts were quality-controlled and cells and genes were filtered to remove mitochondrial and
ribosomal contaminants. Cells for which over 50% of the reads mapped to mitochondrial genes
and cells with fewer than 400 genes expressed were removed. By default, all non-protein-
coding genes, genes coding for ribosomal and mitochondrial proteins, and genes that were
expressed in less than 20 cells were removed. Doublet detection was performed using
scDblFinder[95]. Subsequently, counts were normalized with sctransform[96], regressing out cell
cycle effects, library size, and sample effects as non-regularized dependent variables. Similar
cells were grouped based on unsupervised clustering using Phenograph[97], and automated
cell type classification was performed independently for each cell[98] using gene lists defining
highly expressed genes in different cell types. Major cell type marker lists were developed in-
house based on unpublished datasets (manuscripts in preparation), including the Tumor
Profiler Study[99], using the Seurat FindMarkers method[100]. Immune subtype marker gene lists
were obtained from Newman *et al.*[101] and enriched with T cell subtypes from Sade-Feldman
et al.[102] The results of the unsupervised clustering and cell typing are visualized in a low-
dimensional representation using Uniform Manifold Approximation and Projection (UMAP).

Long-read data analysis

Generating CCS

Using SMRT-Link (version 9.0.0.92188), we performed circular consensus sequencing (CCS)
with the following modified parameters: maximum subread length 50,000 bp, minimum
subread length 10 bp, and minimum number of passes 3.

Unconcatenating long-reads

We used NCBI BLAST (version 2.5.0+) to map the 5' and 3' primers to CCS constructs, with
parameters: "-outfmt 7 -word_size 5" as described previously[24]. Sequences between two
successive primers were used as input for primer trimming using IsoSeq3 Lima (parameters:
--isoseq --dump-clips --min-passes 3). Cell barcodes and UMIs were then demultiplexed using
IsoSeq3 tag with parameter --design T-12U-16B. Finally, we used IsoSeq3 refine with option
--require-polya to remove concatemers and trim polyA tails. Only reads with a correct 5'-3'
primer pair, a barcode also found in the short-read data, a UMI, and a polyA tail were retained.

Isoform classification

Demultiplexing UMIs with IsoSeq3 dedup and calling isoforms on the cohort level with
collapse_isoforms_by_sam.py resulted in unfeasible runtimes. Therefore, we called isoforms
first on the cell level as a pre-filtering step. Long reads were split according to their cell
barcodes, and UMI deduplication was performed using IsoSeq3 dedup. Next, reads were
mapped and aligned to the reference genome (hg38) using minimap2 with parameters: -ax
splice -uf --secondary=no -C5. Identical isoforms were merged based on their aligned exonic
structure using collapse_isoforms_by_sam.py with parameters: -c 0.99 -i 0.95 --
gen_mol_count. We then classified isoforms using SQANTI3[32] with arguments: --skipORF --
fl_count --skip_report. We finally filtered artifacts including intrapriming (accidental priming of
pre-mRNA 'A's), reverse-transcriptase template switching artifacts, and mismapping to non-
canonical junctions. To create a unique isoform catalog for all our samples, we then retained
only reads associated with isoforms passing the SQANTI3 filter, and we ran
collapse_isoforms_by_sam.py, SQANTI3 classification and filtering again on all cells together.
The described pipeline is available here and was implemented in Snakemake, a reproducible
and scalable workflow management system[103].

3' and 5' isoform filtering

For SQANTI3-defined isoforms, incomplete splice match, novel in catalog, and novel not in
catalog, we only retained isoforms falling within 50 bp of a CAGE-validated transcription start

site (FANTOM5 CAGE database), and 50 bp of a polyA site form the PolyASite database[34].
The GENCODE database was used as a comparison, all protein-coding isoforms were
grouped under the GENCODE.full label, a subset including only full-length isoforms was
labeled as GENCOD.FL, and the Matched Annotation from NCBI and EMBL-EBI (MANE[35])
was named GENCODE.MANE.

Cell type-specific isoforms

717 Considering only the SQANTI3-defined 'full splice match', 'novel not in catalog' and 'novel in
718 catalog' isoforms with at least 3 reads, we established the following classification: "Cell-
719 specific" isoforms are present in only 1 cell and "cell type-specific" isoforms are present in >=3
720 cells of a unique cell type.
721

Cell type annotation
723 Cells were annotated with long-reads the same way as short-reads, using scROSHI. The
724 major cell types were modified according to gene expression in long-reads. Immune subtype
725 marker gene lists were unchanged.
726

Normalization and visualization
728
729 Long-read gene expression counts were normalized and visualized as described above for
730 short reads. Long-read gene expression counts were normalized using 10,000 features
731 instead of the default 3,000 in sctransform[96].

Mutation detection

Positions of mutations from Foundation Medicine's targeted NGS panel (Foundation One CDx)
mutations described in Supplementary Table 1 were used as reference. One mutation not
present in the list, TP53_P151H, was detected in IGV in Patient 1 and added to the list. If a
position was mutated at least in one cell belonging to a distal biopsy sample, the mutation was
classified as a germline variant. Cells with one mutated read in one of the positions were
considered mutated.

Alternative polyadenylation (APA) analysis

740 To analyze differences between 3'UTR lengths, we used a modified version of DaPars2[104]
741 (https://github.com/ArthurDondi/DaPars2_LR), with an APA site detection adapted to long-
742 read coverage.
743
Briefly, we identified the 3'UTR exon of each isoform, and overlapping 3'UTR exons with
different 5' start positions were discarded from analysis, as they create false positive APAs.
Then, for each remaining 3'UTR, we computed the coverage for each cell type. The distal site
position was defined as the most 3' position with coverage superior to 10 in any cell type:

$L^* = max(\{k : w_k^c > 10, 1 < k < L, 1 < c < m\})$

where L is the length of the annotated 3′ UTR region and $L*$ is the defined distal site position.

w_k^c is the coverage in cell type c at position k, and m is the total number of cell types.

We inferred the exact location of APA sites by maximizing the coverage gap between the 50

positions before and after the possible APA sites, based on the two-polyA-site model, the most

common model of APA regulation:

755
$$(C^*, P^*) = argmax_{1<c<m, 1<P<L^*} \left(\frac{1}{50} \sum_{i=1}^{50} w_{P+i}^c - \frac{1}{50} \sum_{i=1}^{50} w_{P-i}^c \right)^2$$

where P is the estimated length of alternative proximal 3′ UTR, and the optimal proximal site

$P*$ in cell type $C*$ is the one with the maximal objective function value. w_k^c is the coverage at

the position k and cell type c, and m is the total number of cell types.

Then distal and proximal site coverages W_d and W_p in condition C were defined as:

$W_d = \frac{1}{50} \sum_{i=1}^{50} w_{L^*-i}^c$, $W_p = \frac{1}{50} \sum_{i=1}^{50} w_{P^*-i}^c$

The fraction of distal polyA site usage is then defined as $F = \frac{w_d}{w_d + w_p}$

The degree of difference in APA usage in cell types C_1 and C_2 can be quantified as a

Fraction Change, which is capable of identifying 3′ UTR lengthening (positive index) or

shortening (negative index):

$Fraction\ Change = F_{C_1} - F_{C_2}.$

Differential isoform tests

Differential isoform testing was performed using a χ2 test as previously described in

Scisorseq[26]. Briefly, counts for each isoform ID were assigned to individual cell types, and

genes were discarded if they were mitochondrial, ribosomal, or if they did not reach sufficient

depth per condition (25 reads per condition per gene). P-values from a χ2 test for differential

isoform usage were computed per gene where a sufficient depth was reached, and we

corrected for multiple testing using Benjamini Hochberg correction with a 5% false discovery

rate. If the corrected p-value was ≤0.05 and the sum of change in the relative percent of

isoform (ΔΠ) of the top two isoforms in either positive or negative direction was more than

20%, then the gene was called differentially spliced. To classify the top differentially spliced

genes, we took the rank of genes by ΔΠ and corrected p-values, and took the square root of

the multiplication of those two ranks. The smallest ranks obtained this way were considered

as the top differentially expressed genes. Differentially used isoforms were visualized using
ScisorWiz[105].

Isoform biotypes

784   Novel isoforms biotypes were assessed internally by the GENCODE team with biotypes
785   matching those described by Frankish *et al.*[106]. Known isoform biotypes were retrieved from
786   the GENCODE v36 database.

Biotype change analysis

788   In genes with differential isoform usage between conditions, we compared the biotypes of the
789   most expressed isoform of each condition, and if they were not identical this was considered
790   a change in biotype.

Protein-coding gene expression

792   Considering only genes with protein-coding biotype (GENCODE) with a minimal expression
793   of 20 UMIs in a condition, we assessed the ratio of non-coding isoforms (NMD, intron retention,
794   etc.) expression to total gene expression (non-coding and coding).

Pathway enrichment analysis

We used GSVA to perform pathway enrichment analysis. When comparing TME and distal
cells, gene sets were obtained from MSigDB[107], except for the miR-29 targets that were
obtained from Cushing and colleagues[108]. When we searched for pathways enriched in
lengthened or shortened 3'UTRs in cancer cells, we used the investigate function of the GSEA
webpage (https://www.gsea-msigdb.org/gsea/msigdb/human/annotate.jsp), and compared
against all MSigDB gene sets[107].

Fusion Discovery

Mapped reads from isoform classification were pooled. We called reads mapping to two
separate genes at a distance of more than 100,000 bp or to different chromosomes using
fusion_finder.py (cDNA_Cupcake package, https://github.com/Magdoll/cDNA_Cupcake) with
parameters --min_locus_coverage_bp 200 -d 1000000. Fusion isoforms with sufficient depth
(min. 10 reads) were kept, and their breakpoint, expression per cell type, and number of cells
in which they are expressed were assessed.

Short-reads re-alignment to *IGF2BP2::TESPA1*

We designed a custom reference including *IGF2BP2::TESPA1* transcriptomic breakpoint as
well as the wild-type *IGF2BP2* and *TESPA1* exon junction covering the breakpoint. The
reference was composed of 6 sequences of 80 nucleotides (40 bases upstream and

downstream of the breakpoint), sequences XXX_1 and XXX_2 represent the breakpoints of

the two main isoforms seen in each gene:

>TESPA1_wt_1

TTCTGTCAGACCACATGCTGTTGTGGTGGTGGAGAAAGCAATTCTGGAGGCTGGCAAATCCAAG

GTCAAAAGCCTGCA

>TESPA1_wt_2

TTCTGTCAGACCACATGCTGTTGTGGTGGTGGAGAAAGCTTCACGAGTCTTTGCCAGCAAAAGTC

TGGTGGTGGTGGG

>IGF2BP2_wt_1

ATGTGACGTTGACAACGGCGGTTTCTGTGTCTGTGTTGACTTGTTCCACATTCTCCACTGTCCCA

TATTGAGCCAAAA

>IGF2BP2_wt_2

ATCACTGGATTGTGTGTTCTTCTGAATTACTTCTTTAGGCTTGTTCCACATTCTCCACTGTCCCAT

ATTGAGCCAAAA

>TESPA1_IGF2BP2_fusion_1

TTCTGTCAGACCACATGCTGTTGTGGTGGTGGAGAAAGCCTTGTTCCACATTCTCCACTGTCCCA

TATTGAGCCAAAA

>TESPA1_IGF2BP2_fusion_2

CAAATCCAAGGTCAAAAGCCTGCATCTGGTGAGGGCCTCCTTGTTCCACATTCTCCACTGTCCCA

TATTGAGCCAAAA

Patient 2 reads were aligned to this reference using minimap2 with parameters: -ax sr --

secondary=no. Reads mapping unambiguously to one of those reference sequences were

then attributed to the cell type to which their cell barcode belonged.

scDNA analysis

844 Cell Ranger DNA was used to demultiplex and align Chromium-prepared sequencing

845 samples. We used the cellranger-dna mkfastq command to generate FASTQ files from the

846 Illumina raw BCL files, and we ran the command cellranger-dna cnv to align FASTQ files to

847 the hg38 reference genome, call cells, and estimate copy numbers. We obtained the copy

848 number profiles and detected the main clonal structure of samples using SCICoNE[109].

DNA breakpoint validation

To validate in scDNA data breakpoints found in scRNA data, we used the putative scRNA

breakpoint reads as a reference to re-align scDNA reads using BWA with options: -pt8 -CH.

For the *IGF2BP2::TESPA1* fusion, the reference was composed of 3 sequences of 184

nucleotides (92 bases upstream and downstream of the breakpoint):

>IGF2BP2_WT

CAAACTTGTAGAAATGTGAATTTTTCTTGTTATTTTACAAGATTTGCAAAGGGACCTGAGACCCCG

AAAAGCTTAAGGACTACTGTTAAAAATACTGTTTGTTAAATAACTTTAAAGCAGCTGCAGCCTTTAT

GGGTTGCAGGGAGTTGTATGTAATGCTCAGAAAGAGCTGCCACTGAGAAT

>TESPA1_WT

TTCAATGATGTGGGCTGATTAGAACATAGCTGAAAGCAGGTGTTGGGATATTGATTTCCATGGCT

GGTCCTCACCTGTTACAAAACTTCTACTACAATGAGTTTCAAACTTCAATATGCAATCAATTATCTA

ACCTAAAGATCTTGGTAAAACTGTGATTCATTAGGTCTGGGGTGGGGGCTG

>IGF2BP2_TESPA1_Fusion

TTCAATGATGTGGGCTGATTAGAACATAGCTGAAAGCAGGTGTTGGGATATTGATTTCCATGGCT

GGTCCTCACCTGTTACAAAACTTCTACTACTGTTTGTTAAATAACTTTAAAGCAGCTGCAGCCTTT

ATGGGTTGCAGGGAGTTGTATGTAATGCTCAGAAAGAGCTGCCACTGAGAAT

Reads mapping unambiguously to one of those reference sequences were then attributed to

the clone to which their cell barcode belonged.

Data availability

The raw sequencing files reported in this study have been deposited in the European Genome-phenome Archive (EGA) under the accession number EGAS00001006807. Patient 2 scDNA reads mapping the *IGF2BP2::TESPA1* fusion breakpoint are available at https://eth-nexus.github.io/tu-pro_website/publications/dondi_et_al_2022/. TCGA data was obtained at https://portal.gdc.cancer.gov/repository, with the following specifications: Project=TCGA-OV, DataType=Gene Expression Quantification.

Code availability

The code used to pre-process and analyze the data of this study has been deposited in the GitHub repository: https://github.com/cbg-ethz/scIsoPrep. The code to analyze differential poly-adenylation site in long-reads has been deposited in the GitHub repository: https://github.com/ArthurDondi/DaPars2_LR.

Acknowledgements

We thank Ina Nissen (Genomics Facility Basel) for technical support with PacBio sequencing, Ching-Yeu Liang (University Hospital Basel) for assistance with sample dissociation, and Anne Bertolini for her help with the scAmpi R package. We also wish to thank Elisabeth Tseng (Pacific Bioscience) for her help with cDNA_Cupcake, SQANTI3, and long-read sequencing analysis, Anoushka Joglekar (Weill Cornell Medicine) for her help with ScisorSeq and ScisorWiz as well as Lucia Csepregi for her help with immune repertoire analysis. We thank Olivier Belli for his help with 3'UTR shortening mechanisms. Furthermore, we are grateful to Adam Frankish (Wellcome Sanger Institute) and the GENCODE Project team for their help with the annotation of isoforms. Graphical illustrations were created with BioRender.com. Illumina sequencing was carried out in the Genomics Facility Basel of the University of Basel and the Department of Biosystems Science and Engineering, ETH Zurich. PacBio sequencing was done in the Functional Genomics Center Zurich of the University of Zurich and ETH Zurich, and the Lausanne Genomic Technologies Facility, University of Lausanne.

Funding Information

Part of this work was funded by the SNSF SPARK grant #190413, the grants #2017-510 of the Strategic Focal Area "Personalized Health and Related Technologies (PHRT)" of the ETH Domain, the "Personalized Health and Related Technologies (PHRT)" for Pioneer Project #715 and the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie grant agreement #766030.

Author contributions

U. L, and C. B. acquired funding and conceived and designed the experiments. V. H. S provided patient material. U. L., F .J., A. D., and C. B. selected the clinical cohort. U. L. performed 10x Genomics sample processing as well as short-read and long-read sequencing library preparation. The Tumor Profiler Consortium provided scDNA-seq data and NGS panel results. A. D. designed the pre-processing pipeline and implemented it with the help of N. Borgsmueller. A. D. conducted all computational analyses. F. J. and R. C. performed experimental validation. F. S. assisted in short-read scRNA-seq analysis. A. D., U. L., F. J., N. Beerenwinkel, and C. B. interpreted the data. A. D. and U. L. wrote the manuscript with contributions from all authors. All authors read and approved the final manuscript.

Tumor Profiler Consortium authors list

Rudolf Aebersold[2], Melike Ak[28], Faisal S Al-Quaddoomi[9,17], Silvana I Albert[7], Jonas Albinus[7], Ilaria Alborelli[24], Sonali Andani[6,17,26,31], Per-Olof Attinger[11], Marina Bacac[16], Daniel Baumhoer[24], Beatrice Beck-Schimmer[39], Niko Beerenwinkel[4,17], Christian Beisel[4], Lara Bernasconi[27], Anne Bertolini[9,17], Bernd Bodenmiller[8,35], Ximena Bonilla[6,17,26], Lars Bosshard[9,17], Byron Calgua[24], Ruben Casanova[35], Stéphane Chevrier[35], Natalia Chicherova[9,17], Ricardo Coelho[18], Maya D'Costa[10], Esther Danenberg[37], Natalie Davidson[6,17,26], Monica-Andreea Drăgan[4], Reinhard Dummer[28], Stefanie Engler[35], Martin Erkens[14], Katja Eschbach[4], Cinzia Esposito[37], André Fedier[18], Pedro Ferreira[4], Joanna Ficek[6,17,26], Anja L Frei[31], Bruno Frey[13], Sandra Goetze[7], Linda Grob[9,17], Gabriele Gut[37], Detlef Günther[5], Martina Haberecker[31], Pirmin Haeuptle[1], Viola Heinzelmann-Schwarz[18,23], Sylvia Herter[16], Rene Holtackers[37], Tamara Huesser[16], Alexander Immer[6,12], Anja Irmisch[28], Francis Jacob[18], Andrea Jacobs[35], Tim M Jaeger[11], Katharina Jahn[4], Alva R James[6,17,26], Philip M Jermann[24], André Kahles[6,17,26], Abdullah Kahraman[17,31], Viktor H Koelzer[31], Werner Kuebler[25], Jack Kuipers[4,17], Christian P Kunze[22], Christian Kurzeder[21], Kjong-Van Lehmann[6,17,26], Mitchell Levesque[28], Ulrike Lischetti[18], Sebastian Lugert[10], Gerd Maass[13], Markus G Manz[30], Philipp Markolin[6,17,26], Martin Mehnert[7], Julien Mena[2], Julian M Metzler[29], Nicola Miglino[1], Emanuela S Milani[7], Holger Moch[31], Simone Muenst[24], Riccardo Murri[38], Charlotte KY Ng[24,34], Stefan Nicolet[24], Marta Nowak[31], Monica Nunez Lopez[18], Patrick GA Pedrioli[3], Lucas Pelkmans[37], Salvatore Piscuoglio[18,24], Michael Prummer[9,17], Natalie Rimmer[18], Mathilde Ritter[18], Christian Rommel[14], María L Rosano-González[9,17], Gunnar Rätsch[3,6,17,26], Natascha Santacroce[4], Jacobo Sarabia del Castillo[37], Ramona Schlenker[15], Petra C Schwalie[14], Severin Schwan[11], Tobias Schär[4], Gabriela Senti[27], Wenguang Shao[7], Franziska Singer[9,17], Sujana Sivapatham[35], Berend Snijder[2,17], Bettina Sobottka[31], Vipin T Sreedharan[9,17], Stefan Stark[6,17,26], Daniel J Stekhoven[9,17], Tanmay Tanna[4,6], Alexandre PA Theocharides[30], Tinu M Thomas[6,17,26], Markus Tolnay[24], Vinko Tosevski[16], Nora C Toussaint[9,17], Mustafa A Tuncel[4,17], Marina Tusup[28], Audrey Van Drogen[7], Marcus Vetter[20], Tatjana Vlajnic[24], Sandra Weber[27],

Walter P Weber[19], Rebekka Wegmann[2], Michael Weller[33], Fabian Wendt[7], Norbert Wey[31],

Andreas Wicki[30,36], Mattheus HE Wildschut[2,30], Bernd Wollscheid[7], Shuqing Yu[9,17], Johanna

Ziegler[28], Marc Zimmermann[6,17,26], Martin Zoche[31], Gregor Zuend[32]

[1]Cantonal Hospital Baselland, Medical University Clinic, Rheinstrasse 26, 4410 Liestal,

Switzerland, [2]ETH Zurich, Department of Biology, Institute of Molecular Systems Biology,

Otto-Stern-Weg 3, 8093 Zurich, Switzerland, [3]ETH Zurich, Department of Biology, Wolfgang-

Pauli-Strasse 27, 8093 Zurich, Switzerland, [4]ETH Zurich, Department of Biosystems Science

and Engineering, Mattenstrasse 26, 4058 Basel, Switzerland, [5]ETH Zurich, Department of

Chemistry and Applied Biosciences, Vladimir-Prelog-Weg 1-5/10, 8093 Zurich, Switzerland,

[6]ETH Zurich, Department of Computer Science, Institute of Machine Learning,

Universitätstrasse 6, 8092 Zurich, Switzerland, [7]ETH Zurich, Department of Health Sciences

and Technology, Otto-Stern-Weg 3, 8093 Zurich, Switzerland, [8]ETH Zurich, Institute of

Molecular Health Sciences, Otto-Stern-Weg 7, 8093 Zurich, Switzerland, [9]ETH Zurich,

NEXUS Personalized Health Technologies, John-von-Neumann-Weg 9, 8093 Zurich,

Switzerland, [10]F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, 4070 Basel, Switzerland,

[11]F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, 4070 Basel, Switzerland, [12]Max Planck

ETH Center for Learning Systems, [13]Roche Diagnostics GmbH, Nonnenwald 2, 82377

Penzberg, Germany, [14]Roche Pharmaceutical Research and Early Development, Roche

Innovation Center Basel, Grenzacherstrasse 124, 4070 Basel, Switzerland, [15]Roche

Pharmaceutical Research and Early Development, Roche Innovation Center Munich, Roche

Diagnostics GmbH, Nonnenwald 2, 82377 Penzberg, Germany, [16]Roche Pharmaceutical

Research and Early Development, Roche Innovation Center Zurich, Wagistrasse 10, 8952

Schlieren, Switzerland, [17]SIB Swiss Institute of Bioinformatics, Lausanne, Switzerland,

[18]University Hospital Basel and University of Basel, Department of Biomedicine, Hebelstrasse

20, 4031 Basel, Switzerland, [19]University Hospital Basel and University of Basel, Department

of Surgery, Brustzentrum, Spitalstrasse 21, 4031 Basel, Switzerland, [20]University Hospital

Basel, Brustzentrum & Tumorzentrum, Petersgraben 4, 4031 Basel, Switzerland, [21]University

Hospital Basel, Brustzentrum, Spitalstrasse 21, 4031 Basel, Switzerland, [22]University Hospital

Basel, Department of Information- and Communication Technology, Spitalstrasse 26, 4031

Basel, Switzerland, [23]University Hospital Basel, Gynecological Cancer Center, Spitalstrasse

21, 4031 Basel, Switzerland, [24]University Hospital Basel, Institute of Medical Genetics and

Pathology, Schönbeinstrasse 40, 4031 Basel, Switzerland, [25]University Hospital Basel,

Spitalstrasse 21/Petersgraben 4, 4031 Basel, Switzerland, [26]University Hospital Zurich,

Biomedical Informatics, Schmelzbergstrasse 26, 8006 Zurich, Switzerland, [27]University

Hospital Zurich, Clinical Trials Center, Rämistrasse 100, 8091 Zurich, Switzerland, [28]University

Hospital Zurich, Department of Dermatology, Gloriastrasse 31, 8091 Zurich, Switzerland,

[29]University Hospital Zurich, Department of Gynecology, Frauenklinikstrasse 10, 8091 Zurich,

Switzerland, [30]University Hospital Zurich, Department of Medical Oncology and Hematology,

Rämistrasse 100, 8091 Zurich, Switzerland, [31]University Hospital Zurich, Department of

Pathology and Molecular Pathology, Schmelzbergstrasse 12, 8091 Zurich, Switzerland,

[32]University Hospital Zurich, Rämistrasse 100, 8091 Zurich, Switzerland, [33]University Hospital

and University of Zurich, Department of Neurology, Frauenklinikstrasse 26, 8091 Zurich,

Switzerland, [34]University of Bern, Department of BioMedical Research, Murtenstrasse 35,

3008 Bern, Switzerland, [35]University of Zurich, Department of Quantitative Biomedicine,

Winterthurerstrasse 190, 8057 Zurich, Switzerland, [36]University of Zurich, Faculty of Medicine,

Zurich, Switzerland, [37]University of Zurich, Institute of Molecular Life Sciences,

Winterthurerstrasse 190, 8057 Zurich, Switzerland, [38]University of Zurich, Services and

Support for Science IT, Winterthurerstrasse 190, 8057 Zurich, Switzerland, [39]University of

Zurich, VP Medicine, Künstlergasse 15, 8001 Zurich, Switzerland

Conflict of interest

988   The authors declare no competing interests.

References

1. Garraway, L. A. & Lander, E. S. Lessons from the cancer genome. *Cell* **153**, 17–37 (2013).

2. Hanahan, D. Hallmarks of cancer: New Dimensions. *Cancer Discov.* **12**, 31–46 (2022).

3. Hertzman Johansson, C. & Egyhazi Brage, S. BRAF inhibitors in cancer therapy. *Pharmacol. Ther.* **142**, 176–182 (2014).

4. Li, J. *et al.* A functional genomic approach to actionable gene fusions for precision oncology. *Sci. Adv.* **8**, eabm2382 (2022).

5. Schumacher, T. N. & Schreiber, R. D. Neoantigens in cancer immunotherapy. *Science* **348**, 69–74 (2015).

6. Yu, Y.-P. *et al.* Identification of recurrent fusion genes across multiple cancer types. *Sci. Rep.* **9**, 1074 (2019).

7. Bower, H. *et al.* Life expectancy of patients with chronic myeloid leukemia approaches the life expectancy of the general population. *J. Clin. Oncol.* **34**, 2851–2857 (2016).

8. Khan, M. *et al.* ALK inhibitors in the treatment of ALK positive NSCLC. *Front. Oncol.* **8**, 557 (2018).

9. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–421 (2013).

10. Zack, T. I. *et al.* Pan-cancer patterns of somatic copy number alteration. *Nat. Genet.* **45**, 1134–1140 (2013).

11. Gao, Q. *et al.* Driver fusions and their implications in the development and treatment of human cancers. *Cell Rep.* **23**, 227-238.e3 (2018).

12. Wang, Y., Shi, T., Song, X., Liu, B. & Wei, J. Gene fusion neoantigens: Emerging targets for cancer immunotherapy. *Cancer Lett.* **506**, 45–54 (2021).

13. Wei, Z. *et al.* The Landscape of Tumor Fusion Neoantigens: A Pan-Cancer Analysis. *iScience* **21**, 249–260 (2019).

14. Blencowe, B. J. Alternative splicing: new insights from global analyses. *Cell* **126**, 37–47 (2006).

15. Bonnal, S. C., López-Oreja, I. & Valcárcel, J. Roles and mechanisms of alternative splicing in cancer - implications for care. *Nat. Rev. Clin. Oncol.* **17**, 457–474 (2020).

16. Kahles, A. *et al.* Comprehensive Analysis of Alternative Splicing Across Tumors from 8,705 Patients. *Cancer Cell* **34**, 211-224.e6 (2018).

17. Marusyk, A. & Polyak, K. Tumor heterogeneity: causes and consequences. *Biochim. Biophys. Acta* **1805**, 105–117 (2010).

18. Whiteside, T. L. The tumor microenvironment and its role in promoting tumor growth. *Oncogene* **27**, 5904–5912 (2008).

19. Hedlund, E. & Deng, Q. Single-cell RNA sequencing: Technical advancements and
biological applications. *Mol. Aspects Med.* **59**, 36–46 (2018).

20. Marx, V. Method of the year: long-read sequencing. *Nat. Methods* **20**, 6–11 (2023).

21. Glinos, D. A. *et al.* Transcriptome variation in human tissues revealed by long-read
sequencing. *Nature* **608**, 353–359 (2022).

22. Philpott, M. *et al.* Nanopore sequencing of single-cell transcriptomes with scCOLOR-
seq. *Nat. Biotechnol.* **39**, 1517–1520 (2021).

23. Gupta, I. *et al.* Single-cell isoform RNA sequencing characterizes isoforms in thousands
of cerebellar cells. *Nat. Biotechnol.* **36**, 1197–1202 (2018).

24. Shi, Z.-X. *et al.* High-throughput and high-accuracy single-cell RNA isoform analysis
using PacBio circular consensus sequencing. *Nat. Commun.* **14**, 2631 (2023).

25. Al'Khafaji, A. M. *et al.* High-throughput RNA isoform sequencing using programmed
cDNA concatenation. *Nat. Biotechnol.* (2023) doi:10.1038/s41587-023-01815-7.

26. Joglekar, A. *et al.* A spatially resolved brain region- and cell type-specific isoform atlas
of the postnatal mouse brain. *Nat. Commun.* **12**, 463 (2021).

27. Hardwick, S. A. *et al.* Single-nuclei isoform RNA sequencing unlocks barcoded exon
connectivity in frozen brain tissue. *Nat. Biotechnol.* **40**, 1082–1092 (2022).

28. Veiga, D. F. T. *et al.* A comprehensive long-read isoform analysis platform and
sequencing resource for breast cancer. *Sci. Adv.* **8**, eabg6711 (2022).

29. Namba, S. *et al.* Transcript-targeted analysis reveals isoform alterations and double-
hop fusions in breast cancer. *Commun. Biol.* **4**, 1320 (2021).

30. Macintyre, G. *et al.* Copy number signatures and mutational processes in ovarian
carcinoma. *Nat. Genet.* **50**, 1262–1270 (2018).

31. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian
carcinoma. *Nature* **474**, 609–615 (2011).

32. Tardaguila, M. *et al.* SQANTI: extensive characterization of long-read transcript
sequences for quality control in full-length transcriptome identification and quantification.
Genome Res. **28**, 396–411 (2018).

33. Abugessaisa, I. *et al.* FANTOM5 CAGE profiles of human and mouse reprocessed for
GRCh38 and GRCm38 genome assemblies. *Sci. Data* **4**, 170107 (2017).

34. Herrmann, C. J. *et al.* PolyASite 2.0: a consolidated atlas of polyadenylation sites from
3' end sequencing. *Nucleic Acids Res.* **48**, D174–D179 (2020).

35. Morales, J. *et al.* A joint NCBI and EMBL-EBI transcript set for clinical genomics and
research. *Nature* **604**, 310–315 (2022).

36. Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP.
Nat. Biotechnol. **37**, 38–44 (2018).

37. Mayr, C. & Bartel, D. P. Widespread shortening of 3'UTRs by alternative cleavage and

1062   polyadenylation activates oncogenes in cancer cells. *Cell* **138**, 673–684 (2009).

1063 38. Maurer, B. *et al.* MicroRNA-29, a key regulator of collagen expression in systemic
1064   sclerosis. *Arthritis Rheum.* **62**, 1733–1743 (2010).

1065 39. Chen, Y. & Wang, X. miRDB: an online database for prediction of functional microRNA
1066   targets. *Nucleic Acids Res.* **48**, D127–D131 (2020).

1067 40. Jin, Y., Song, X., Sun, X. & Ding, Y. Up-regulation of collagen type V alpha 2 (COL5A2)
1068   promotes malignant phenotypes in gastric cancer cell via inducing epithelial-
1069   mesenchymal transition (EMT). *Open Med (Wars)* **18**, 20220593 (2023).

1070 41. Shintani, Y., Maeda, M., Chaika, N., Johnson, K. R. & Wheelock, M. J. Collagen I
1071   promotes epithelial-to-mesenchymal transition in lung cancer cells via transforming
1072   growth factor-beta signaling. *Am. J. Respir. Cell Mol. Biol.* **38**, 95–104 (2008).

1073 42. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
1074   collection. *Cell Syst.* **1**, 417–425 (2015).

1075 43. Owusu-Ansah, K. G. *et al.* COL6A1 promotes metastasis and predicts poor prognosis
1076   in patients with pancreatic cancer. *Int. J. Oncol.* **55**, 391–404 (2019).

1077 44. Ramadoss, S., Chen, X. & Wang, C.-Y. Histone demethylase KDM6B promotes
1078   epithelial-mesenchymal transition. *J. Biol. Chem.* **287**, 44508–44517 (2012).

1079 45. Du, B. *et al.* The potential role of TNFAIP3 in malignant transformation of gastric
1080   carcinoma. *Pathol. Res. Pract.* **215**, 152471 (2019).

1081 46. D'Arrigo, P. *et al.* The splicing FK506-binding protein-51 isoform plays a role in
1082   glioblastoma resistance through programmed cell death ligand-1 expression regulation.
1083   *Cell Death Discov.* **5**, 137 (2019).

1084 47. Jie, W. *et al.* Pathophysiological functions of rnd3/rhoe. *Compr. Physiol.* **6**, 169–186
1085   (2015).

1086 48. Han, Q. *et al.* Omental cancer-associated fibroblast-derived exosomes with low
1087   microRNA-29c-3p promote ovarian cancer peritoneal metastasis. *Cancer Sci.* **114**,
1088   1929–1942 (2023).

1089 49. Chou, J. *et al.* GATA3 suppresses metastasis and modulates the tumour
1090   microenvironment by regulating microRNA-29b expression. *Nat. Cell Biol.* **15**, 201–213
1091   (2013).

1092 50. Smyth, A., Callaghan, B., Willoughby, C. E. & O'Brien, C. The Role of miR-29 Family in
1093   TGF-β Driven Fibrosis in Glaucomatous Optic Neuropathy. *Int. J. Mol. Sci.* **23**, (2022).

1094 51. Yu, C.-C., Liao, Y.-W., Hsieh, P.-L. & Chang, Y.-C. Targeting lncRNA H19/miR-
1095   29b/COL1A1 Axis Impedes Myofibroblast Activities of Precancerous Oral Submucous
1096   Fibrosis. *Int. J. Mol. Sci.* **22**, (2021).

1097 52. Tian, X., Zuo, X., Hou, M., Li, C. & Teng, Y. LncRNA-H19 regulates chemoresistance to
1098   carboplatin in epithelial ovarian cancer through microRNA-29b-3p and STAT3. *J.*

1099   *Cancer* **12**, 5712–5722 (2021).

53. Lv, M. *et al.* lncRNA H19 regulates epithelial-mesenchymal transition and metastasis of bladder cancer by miR-29b-3p as competing endogenous RNA. *Biochim. Biophys. Acta Mol. Cell Res.* **1864**, 1887–1899 (2017).

54. An, Q. *et al.* circKRT7-miR-29a-3p-COL1A1 Axis Promotes Ovarian Cancer Cell Progression. *Onco Targets Ther* **13**, 8963–8976 (2020).

55. Panda, A. C. Circular RNAs Act as miRNA Sponges. *Adv. Exp. Med. Biol.* **1087**, 67–79 (2018).

56. García-Bartolomé, A. *et al.* Altered Expression Ratio of Actin-Binding Gelsolin Isoforms Is a Novel Hallmark of Mitochondrial OXPHOS Dysfunction. *Cells* **9**, (2020).

57. Chen, Z.-Y., Wang, P.-W., Shieh, D.-B., Chiu, K.-Y. & Liou, Y.-M. Involvement of gelsolin in TGF-beta 1 induced epithelial to mesenchymal transition in breast cancer cells. *J. Biomed. Sci.* **22**, 90 (2015).

58. Philippou, A., Maridaki, M., Pneumaticos, S. & Koutsilieris, M. The complexity of the IGF1 gene splicing, posttranslational modification and bioactivity. *Mol. Med.* **20**, 202–214 (2014).

59. Tagami, S., Eguchi, Y., Kinoshita, M., Takeda, M. & Tsujimoto, Y. A novel protein, RTN-XS, interacts with both Bcl-XL and Bcl-2 on endoplasmic reticulum and reduces their anti-apoptotic activity. *Oncogene* **19**, 5736–5746 (2000).

60. Gong, L. *et al.* RTN1-C mediates cerebral ischemia/reperfusion injury via ER stress and mitochondria-associated apoptosis pathways. *Cell Death Dis.* **8**, e3080 (2017).

61. Parreno, J., Amadeo, M. B., Kwon, E. H. & Fowler, V. M. Tropomyosin 3.1 association with actin stress fibers is required for lens epithelial to mesenchymal transition. *Invest. Ophthalmol. Vis. Sci.* **61**, 2 (2020).

62. Nacu, S. *et al.* Deep RNA sequencing analysis of readthrough gene fusions in human prostate adenocarcinoma and reference samples. *BMC Med. Genomics* **4**, 11 (2011).

63. Banday, A. R. *et al.* Genetic regulation of OAS1 nonsense-mediated decay underlies association with COVID-19 hospitalization in patients of European and African ancestries. *Nat. Genet.* **54**, 1103–1116 (2022).

64. Choy, J. Y. H., Boon, P. L. S., Bertin, N. & Fullwood, M. J. A resource of ribosomal RNA-depleted RNA-Seq data from different normal adult and fetal human tissues. *Sci. Data* **2**, 150063 (2015).

65. Martínez-Jiménez, F. *et al.* A compendium of mutational cancer driver genes. *Nat. Rev. Cancer* **20**, 555–572 (2020).

66. Tate, J. G. *et al.* COSMIC: the catalogue of somatic mutations in cancer. *Nucleic Acids Res.* **47**, D941–D947 (2019).

67. Mertens, F., Johansson, B., Fioretos, T. & Mitelman, F. The emerging complexity of

gene fusions in cancer. *Nat. Rev. Cancer* **15**, 371–381 (2015).

68.   Yang, W. *et al.* Immunogenic neoantigens derived from gene fusions stimulate T cell
responses. *Nat. Med.* **25**, 767–775 (2019).

69.   Berger, M. F. & Mardis, E. R. The emerging clinical relevance of genomics in cancer
medicine. *Nat. Rev. Clin. Oncol.* **15**, 353–365 (2018).

70.   Rynne-Vidal, A. *et al.* Mesothelial-to-mesenchymal transition as a possible therapeutic
target in peritoneal metastasis of ovarian cancer. *J. Pathol.* **242**, 140–151 (2017).

71.   Cai, J. *et al.* Fibroblasts in omentum activated by tumor cells promote ovarian cancer
growth, adhesion and invasiveness. *Carcinogenesis* **33**, 20–29 (2012).

72.   Kenny, H. A. *et al.* Mesothelial cells promote early ovarian cancer metastasis through
fibronectin secretion. *J. Clin. Invest.* **124**, 4614–4628 (2014).

73.   Qin, W. *et al.* TGF-β/Smad3 signaling promotes renal fibrosis by inhibiting miR-29. *J.*
Am. Soc. Nephrol. **22**, 1462–1474 (2011).

74.   Yu, P.-N. *et al.* Downregulation of miR-29 contributes to cisplatin resistance of ovarian
cancer cells. *Int. J. Cancer* **134**, 542–551 (2014).

75.   Dasari, S., Fang, Y. & Mitra, A. K. Cancer associated fibroblasts: naughty neighbors that
drive ovarian cancer progression. *Cancers (Basel)* **10**, (2018).

76.   Li, H. *et al.* IGF-IR signaling in epithelial to mesenchymal transition and targeting IGF-
IR therapy: overview and new insights. *Mol. Cancer* **16**, 6 (2017).

77.   Liefers-Visser, J. A. L., Meijering, R. A. M., Reyners, A. K. L., van der Zee, A. G. J. & de
Jong, S. IGF system targeted therapy: Therapeutic opportunities for ovarian cancer.
Cancer Treat. Rev. **60**, 90–99 (2017).

78.   Yang, Y. *et al.* Tumor Suppressor microRNA-138 Suppresses Low-Grade Glioma
Development and Metastasis via Regulating IGF2BP2. *Onco Targets Ther* **13**, 2247–
2260 (2020).

79.   Wang, D. *et al.* Tespa1 is involved in late thymocyte development through the regulation
of TCR-mediated signaling. *Nat. Immunol.* **13**, 560–568 (2012).

80.   Dhamija, S. & Menon, M. B. Non-coding transcript variants of protein-coding genes -
what are they good for? *RNA Biol.* **15**, 1025–1031 (2018).

81.   Dhamija, S. & Diederichs, S. From junk to master regulators of invasion: lncRNA
functions in migration, EMT and metastasis. *Int. J. Cancer* **139**, 269–280 (2016).

82.   Joglekar, A., Foord, C., Jarroux, J., Pollard, S. & Tilgner, H. U. From words to complete
phrases: insight into single-cell isoforms using short and long reads. *Transcription* 1–13
(2023) doi:10.1080/21541264.2023.2213514.

83.   Lischetti, U. *et al.* Dynamic thresholding and tissue dissociation optimization for CITE-
seq identifies differential surface protein abundance in metastatic melanoma. *Commun.*
Biol. **6**, 830 (2023).

84. Fortelny, N., Overall, C. M., Pavlidis, P. & Freue, G. V. C. Can we predict protein from
mRNA levels? *Nature* **547**, E19–E20 (2017).

85. Vogel, C. & Marcotte, E. M. Insights into the regulation of protein abundance from
proteomic and transcriptomic analyses. *Nat. Rev. Genet.* **13**, 227–232 (2012).

86. Liu, Y., Beyer, A. & Aebersold, R. On the Dependency of Cellular Protein Levels on
mRNA Abundance. *Cell* **165**, 535–550 (2016).

87. Volden, R. & Vollmers, C. Single-cell isoform analysis in human immune cells. *Genome
Biol.* **23**, 47 (2022).

88. Byrne, A. *et al.* Single-cell long-read targeted sequencing reveals transcriptional
variation in ovarian cancer. *BioRxiv* (2023) doi:10.1101/2023.07.17.549422.

89. Dutton, G. CRISPR-Cas9 Technology Cuts Clutter from Sequencing Libraries. *Genetic
Engineering & Biotechnology News* **41**, 24–25 (2021).

90. Lang, F., Schrörs, B., Löwer, M., Türeci, Ö. & Sahin, U. Identification of neoantigens for
individualized therapeutic cancer vaccines. *Nat. Rev. Drug Discov.* **21**, 261–282 (2022).

91. Lin, M. J. *et al.* Cancer vaccines: the next immunotherapy frontier. *Nat. Cancer* **3**, 911–
926 (2022).

92. Hebelstrup, K. H. *et al.* UCE: A uracil excision (USER)-based toolbox for transformation
of cereals. *Plant Methods* **6**, 15 (2010).

93. Griffiths, J. A., Richard, A. C., Bach, K., Lun, A. T. L. & Marioni, J. C. Detection and
removal of barcode swapping in single-cell RNA-seq data. *Nat. Commun.* **9**, 2667
(2018).

94. Bertolini, A. *et al.* scAmpi-A versatile pipeline for single-cell RNA-seq analysis from
basics to clinics. *PLoS Comput. Biol.* **18**, e1010097 (2022).

95. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet
identification in single-cell sequencing data using scDblFinder. *F1000Res.* **10**, 979
(2021).

96. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biol.* **20**, 296 (2019).

97. Levine, J. H. *et al.* Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like
Cells that Correlate with Prognosis. *Cell* **162**, 184–197 (2015).

98. Prummer, M. *et al.* scROSHI: robust supervised hierarchical identification of single cells.
NAR Genom. Bioinform. **5**, lqad058 (2023).

99. Irmisch, A. *et al.* The Tumor Profiler Study: integrated, multi-omic, functional tumor
profiling for clinical decision support. *Cancer Cell* **39**, 288–293 (2021).

100. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.
(2021).

101. Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.

Nat. Methods **12**, 453–457 (2015).

102. Sade-Feldman, M. *et al.* Defining T Cell States Associated with Response to Checkpoint
Immunotherapy in Melanoma. *Cell* **175**, 998-1013.e20 (2018).

103. Köster, J. & Rahmann, S. Snakemake--a scalable bioinformatics workflow engine.
Bioinformatics **28**, 2520–2522 (2012).

104. Feng, X. *et al.* TC3A: The Cancer 3′ UTR Atlas, *Nucleic Acids Research*. **46**, 1027–1030
(2018).

105. Stein, A. N., Joglekar, A., Poon, C.-L. & Tilgner, H. U. ScisorWiz: visualizing differential
isoform expression in single-cell long-read data. *Bioinformatics* **38**, 3474–3476 (2022).

106. Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Res. **47**, D766–D773 (2019).

107. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,
1739–1740 (2011).

108. Cushing, L. *et al.* miR-29 is a major regulator of genes associated with pulmonary
fibrosis. *Am. J. Respir. Cell Mol. Biol.* **45**, 287–294 (2011).

109. Kuipers, J., Tuncel, M. A., Ferreira, P., Jahn, K. & Beerenwinkel, N. Single-cell copy
number   calling   and   event   history   reconstruction.   *BioRxiv*   (2020)
doi:10.1101/2020.04.28.065755.

1228

Figure 1: Study design and long-read data overview.
(a) Schematic of freshly processed HGSOC omentum metastases and patient-matched tumor-free distal omentum tissue biopsies, scRNA-seq. **(b)** Definition of SQANTI-defined isoform structural categories. **(c)** Proportions of isoform structural categories detected in merged metastasis and distal omentum samples. Percentage and total number of isoforms per category are indicated. **(d)** Proportions of unique reads attributed to isoforms detected in **(c)**. Percentage and total number of UMIs per category are indicated. **(e)** Percentage of isoforms for which transcription start site is supported by CAGE (FANTOM5) data and transcription termination site is supported by poly-adenylation (PolyASite) data, per isoform structural categories. 'GENCODE.all' indicates all protein-coding isoforms in the GENCODE database, 'GENCODE.FL' is a subset of 'GENCODE.all' containing only isoforms tagged as full-length, and 'GENCODE.MANE' is a hand-curated subset of canonical transcripts, one per human protein-coding locus. **(f)** GENCODE-defined biotype composition of novel isoforms. **(g)** Biotype composition of the GENCODE database.

Figure 2: Clustering and cell type-specific isoform distribution.

(a) Cohort UMAP embeddings by data types and automatic cell type annotation. Top and bottom rows:

cell type labels based on short- and long-read data, respectively. Left column: embedding on short-read

data - gene level, middle column: embedding on long-read data - gene level, right column: embedding

on long-read data - isoform level. **(b)** Jaccard distance of cell populations in different UMAP

embeddings: short-reads - gene level versus long-reads - gene level (left), short-reads - gene level
versus long-reads - isoform level (middle), long-reads - gene level versus long-reads - isoform level
(right). Long-reads - gene level UMAP cohort visualizations of cells with at least one germline **(c)** or
somatic **(d)** mutation also found in targeted NGS panel data of matched patient samples. Germline
variants are variants detected in healthy omentum distal samples. **(e)** SQANTI-defined structural
category normalized distribution of isoforms detected per cell type (number of isoforms displayed in
white).

1258 **Figure 3: Epithelial-to-mesenchymal transition in the tumor microenvironment.**

1259 **(a)** Zoom of UMAP embeddings of the cohorts' long-read - gene level data (Fig. 2a, middle column)

1260 highlighting tumor and stromal (mesothelial and fibroblast) cells, colored by biopsy tissue type (left) and

1261 EMT gene set signal (right). **(b)** Volcano plot of genes with APA in mesothelial cells. Genes have either

1262 a lengthened (red) or shortened (blue) 3'UTR in TME compared to distal mesothelial cells. Differentially

1263 lengthened or shortened genes targeted by miR-29 are colored in green. Genes with -log10(p-adjusted)

1264 > 10 and |Fraction Change| > 0.4 are annotated. **(c)** IGV view of 3'UTR raw coverage of *COL1A2*,

1265 *COL6A1*, *COL3A1*, and *COL5A2* in tissue cell types. On the top left between brackets, the coverage

range is displayed throughout each condition. In blue, Ensembl canonical 3'UTR, and for each gene,
distal (d) and proximal (p) APA sites are annotated. **(d)** Log fold-change expression between TME and
distal mesothelial cells of lengthened genes targeted (+, green) or not targeted (-, red) by miR-29, and
shortened genes (blue). **(e)** Cohort UMAP embedding long-read data - gene level, colored by gene set
signal of ECM-related genes targeted by miR-29. **(f)** ScisorWiz representation of *COL1A1* isoforms.
Colored areas are exons, whitespace areas are intronic space, not drawn to scale, and each horizontal
line represents a single read colored according to cell types. Dashed boxes highlight the use of the
canonical 3' UTR in TME fibroblasts and mesothelial cells, while distal mesothelial cells use an earlier
3' exon termination.

1307

Figure 4: Differential isoforms and 3'UTR lengths in cancer.

(a) Number of genes with change in isoform usage between HGSOC and all distal cells. In orange, genes with differentially expressed isoforms and a change in relative isoform abundance >20% (>50% in green). In blue, genes with no differentially expressed isoforms or change in relative isoform abundance <20%. **(b)** Alluvial plot of biotypes of most expressed isoforms in HGSOC and distal cells in genes containing an isoform change >20% (n=960). Each vein represents the conversion of one biotype to another. For example, in 7 genes, the most expressed isoform in HGSOC cells is protein-coding, while distal cells' one is non-protein-coding. **(c)** Alluvial plot of biotypes of most expressed isoforms in HGSOC and distal cells in genes containing an isoform switch (>50% change, n=39). **(d)** ScisorWiz representation of isoforms in *IGF1*, each horizontal line represents a single isoform colored according to cell types. Exons are numbered according to the Gencode reference, *Class I* and *II* isoforms are isoforms with starting exons 1 and 2, respectively. **(e)** Top: IGV view of *OAS1* expression in patients. Patient 3 has low p46 expression compared to others. Bottom: zoom on the last exon of isoform p46, where all patients have at least one mutated A allele in the splice acceptor site. **(f)** Volcano plot of genes with APA in cancer versus distal cells. Genes have either a lengthened (red) or shortened (blue) 3'UTR in cancer cells compared to all distal cells. Differentially lengthened or shortened genes targeted by miR-29 are colored in green. Genes with -log10(p-adjusted) > 10 and |Fraction Change| > 0.5 are annotated.

1326
1327
Figure 5: Tumor and patient-specific detection of novel *IGF2BP2::TESPA1* gene fusion.
(a) Overview of wt *IGF2BP2*, wt *TESPA1,* and *IGF2BP2::TESPA1* gene fusion with exon structure. **(b)**
Overview of wt IGF2BP2, wt TESPA1, and fusion protein with protein domains. RRM: RNA-recognition
motif, KH: heterogeneous nuclear ribonucleoprotein K-homology domain, KRAP_IP3R_bind: Ki-ras-
induced actin-interacting protein-IP3R- interacting domain. **(c)** Violin plot showing patient- and tumor-
specific *IGF2BP2::TESPA1* fusion transcript detection in Patient 2. **(d)** UMI count in fusion-containing
vs. -lacking Patient 2 tumor cells. **(e)** UMAP embeddings of the cohorts' short-read data. Cells are
colored if they express *IGF2BP2* (red), *TESPA1* (green), or both (yellow) in short- (left panel) or long-
reads (right panel). **(f)** Raw expression of *TESPA1* (left) and *IGF2BP2* (right) in short- (top) or long-
reads (bottom), by sample and cell type. **(g)** IGV view of short-reads (top), non-fusion long-reads
(middle), and fusion long-reads (bottom) mapping to the 3'UTR of *TESPA1*. Non-fusion reads are either
triple hSNP-mutated or non-mutated, while fusion and short-reads are only triple hSNP-mutated.

1340

Figure 6: *IGF2BP2::TESPA1* fusion breakpoint validation in bulk and scDNA.
(a) Genotyping PCR on genomic DNA isolated from matched patient samples using gene-specific primers for *IGF2BP2::TESPA1* genomic breakpoint (top), wt *TESPA1* (middle) and wt *IGF2BP2* (bottom). **(b)** Copy number values per subclone in Patient 2 scDNA-seq data. Sublone 0 has multiple copy number alterations, indicative of cancer, while Subclone 1 is copy-number neutral, presumably non-cancer. **(c)** IGV view of scDNA reads aligning unambiguously to the *IGF2BP2::TESPA1* genomic breakpoint (top), wt *TESPA1* (middle), or wt *IGF2BP2* (bottom). In red, reads from Subclone 0 cells (cancer); in blue, reads from Subclone 1 cells (non-cancer).