Title: One thousand soils for molecular understanding of belowground carbon cycling

Authors: Maggie M. Bowman[1] Alexis E. Heath[2], Tamas Varga[1], Anil K. Battu[1], Rosalie
K. Chu[1], Jason Toyoda[1], Tanya E. Cheeke[3], Stephanie S. Porter[4], Kevan Moffett[4],
Brittany Letendre[4], Odeta Qafoku[1], John R. Bargar[1], Douglas Mans[1], Nancy Hess[1],
Emily B. Graham[1,3*]

Affiliation:

1) Pacific Northwest National Laboratory, Richland, WA, 99354, USA
2) Bowling Green State University, Bowling Green, OH, 43403, USA
3) School of Biological Sciences, Washington State University, Richland, WA, USA
4) School of the Environment, Washington State University, Vancouver, WA 98686,
USA

Abstract:

While significant progress has been made in understanding global carbon (C) cycling,
the mechanisms regulating belowground C fluxes and storage are still uncertain. New
molecular technologies have the power to elucidate these processes, yet we have no
widespread standardized implementation of molecular techniques. To address this gap,
we introduce the Molecular Observation Network (MONet), a decadal vision from the
Environmental Molecular Sciences Laboratory (EMSL), to develop a national network
for understanding the molecular composition, physical structure, and hydraulic and
biological properties of soil and water. These data are essential for advancing the next
generation of multiscale Earth systems models. In this paper, we discuss the 1000 Soils
Pilot for MONet, including a description of standardized sampling materials and
protocols and a use case to highlight the utility of molecular-level and microstructural
measurements for assessing the impacts of wildfire on soil. While the 1000 Soils Pilot
generated a plethora of data, we focus on assessments of soil organic matter (SOM)
chemistry via Fourier-transform ion cyclotron resonance-mass spectrometry and
microstructural properties via X-ray Computed Tomography to highlight the effects of
recent fire history in forested ecosystems on belowground C cycling. We observed
decreases in soil respiration, microbial biomass, and potential enzyme activity in soils
with high frequency burns. Additionally, the nominal oxidation state of carbon in SOM
increased with burn frequency in surface soils. This results in a quantifiable shift in the
molecular signature of SOM and shows that wildfire may result in oxidation of SOM and
structural changes to soil pore networks that persist into deeper soils.

Introduction

Soil organic matter (SOM) is a critical part of the global carbon (C) cycle. Belowground ecosystems contain more C than stored in terrestrial vegetation and the atmosphere combined (Crowther et al., 2019; Crowther et al., 2016; Jobbágy and Jackson, 2000), and SOM is the largest and most biologically active portion of soil C. SOM decomposition is regulated by a complex and interacting set of factors including soil structure, moisture distribution, temperature, pH, and nutrient status; collectively, these factors determine accessibility, bioavailability, and rate kinetics of SOM (Graham and Hofmockel, 2022). Despite the importance of SOM in the global C cycle, the drivers of SOM decomposition from molecular to continental scales are not well understood.

Recent research suggests that standardized, spatially-resolved, and high-resolution data may be critical to reducing the uncertainty surrounding estimates of belowground C dynamics under future climate scenarios (Lehmann and Kleber, 2015; Viscarra Rossel et al., 2019). For example, pore-scale structural characterization could reveal spatial factors that inhibit SOM decomposition, and molecular chemistry may provide insight into SOM reactivity that cannot be observed from bulk measurements. As of yet, such information is lacking from soil databases (e.g., Web Soil Survey, International Soil Radiocarbon Database, and Soils Data Harmonization), and no widespread implementations of standardized high-resolution techniques simultaneously investigate the myriad factors influencing SOM decomposition (Bond-Lamberty and Thomson, 2010; Lawrence et al., 2020; NRCS; Wieder et al., 2021).

Consequently, soil C cycles are poorly represented in biogeochemical and Earth System Models (ESMs) that predict changes in the global climate (Bradford et al., 2016). For example, most ESMs simulate changes in belowground C concentration within coarsely defined pools (i.e., agnostic of SOM composition) rather than representing chemical and microbial reactions within SOM pools (Bond-Lamberty et al., 2020; Malhotra et al., 2019; Rasmussen et al., 2018; Waring et al., 2020). The depth of soil molecular and microphysical/chemical information presents a major limitation to improving models at regional and Continental United States (CONUS) scales, where impacts of climate change are manifested and can be understood.

Led by a team at the Environmental Molecular Science Laboratory (EMSL), a new decadal initiative is developing a CONUS-scale database of soil compositional, physical, and metagenomic measurements that are open and FAIR (Findable, Accessible, Interoperable and Reusable), termed the Molecular Observation Network (MONet) program (Todd-Brown et al., 2022; Wilkinson et al., 2016). The goal of MONet is to facilitate molecular data syntheses and modelling activities that can reduce uncertainty in soil C simulation at regional and CONUS scales. In this article, we describe the 1000 Soils Pilot for MONet, which focuses on understanding the drivers and fluxes of soil C cycles through standardized molecular measurements. It represents our initial goal to collect soil cores from 1000 locations.

To highlight the power of high-resolution measurements for understanding soil structure, molecular SOM composition, and SOM decomposition, we also present a use case from the 1000 Soils Pilot. Because wildfires are becoming increasingly prevalent worldwide

(Abatzoglou and Williams, 2016; Aponte et al., 2016) and have lasting impacts on
ecosystem structure and function (Abatzoglou and Williams, 2016; Certini, 2005; Doerr
et al., 2006; Lombao et al., 2021; Matosziuk et al., 2020; Neary et al., 2005; Verma and
Jayakumar, 2012; Wang et al., 2012), we focus our use case on a set of three locations
with varied burned histories. We show that wildfire may result in oxidation of SOM and
in structural changes to soil pore networks that persist into deeper soils, which are not
investigated by typical sampling schema. These are important considerations in the
management of forest ecosystem restoration post-wildfire and present an interesting
avenue for more robust investigations as the MONet database expands.

Approach

Through partnerships with individual researchers and ecological networks (Figure 1A),
the 1000 Soils Pilot facilitated the collection of 76 sets of soil cores using a standardized
sampling kit and field protocols. Collaborators collected two 30-cm (3-inch dia.) replicate
intact cores using a slide hammer soil corer (AMS, Inc., USA), as well as four 10-cm (2-
inch dia.) soil cores using a wooden block and mallet. Other field-based measurements
included soil temperature, moisture, and electrical conductivity, measured using a Teros
12 Probe and ZSC Bluetooth interface (Meter Group, Inc., USA). Soil infiltration rate
was measured using the single-ring infiltrometer method (Dane and Topp, 2020; Klute
and Dirksen, 1986). A step-by-step sampling protocol is available in the SI and at
protocols.io. After sample collection, all data was generated by EMSL or EMSL-
contracted vendors. All data from the 1000 Soils Pilot are available via Zenodo
(https://zenodo.org/communities/emsl-monet; Bowman et al., 2022).

Figure 1: (A) Sites included in the 1000 Soils Pilot overlain on National Ecological
Observatory Network (NEON) eco-domains. The color of the symbol indicates site
affiliation as Academic (light blue), Critical Zone Network (CZNet; dark blue),
Department of Energy Science Focus Area (DOE SFA; light green), and NEON (dark
green). (B) Soil core analysis workflow.

Data Generation Overview.

Collaborators shipped all cores to EMSL on ice within 24 hr of core collection for
immediate processing. One 30-cm core was dedicated to a biotic workflow for time
sensitive measurements, and the other 30-cm core was dedicated to an abiotic
workflow for structural and hydraulic characterization (Figure 1B). To account for
surface heterogeneity, three of the 10 cm cores were homogenized with the top 10 cm
of the larger biotic core. A full list of data types generated by the 1000 Soils Pilot is
available in SI Table 1.

Immediately on arrival (<48 hr), we processed the biotic core by separating it into 10 cm
depth intervals, which were then sieved at 4 mm. The middle section was archived, and
all analyses were conducted on homogenized top (0-10 cm) and bottom (deepest 10
cm) sections as described below. We measured microbial biomass C and N via
chloroform fumigation (Brookes et al., 1985; Witt et al., 2000; Zhao et al., 2022).
Aliquots of K_2SO_4 extractions from the initial (background) steps of microbial biomass
measurements were archived at - 20° C for later analysis of NO_3 via Cd-reduction
method and NH_4 via spectroscopic determination (Bremner, 1965; Dahnke and
Johnson, 1990). Inorganic P was extracted from each soil section via Bray or Olsen
extraction (depending on pH) and subsequent analysis via methods outlined in (Bray
and Kurtz, 1945; Olsen et al., 1982). We collected sub-samples of soil for DNA
extraction and subsequent metagenomic sequencing on the Illumina NovaSeq platform
(stored at -80° C). We measured soil pH using a 1:1 soil to water ratio using a calibrated
pH probe. Gravimetric water content was determined via oven drying at 60° C.
Remaining soil from each section was air dried prior to the analyses below. Beta-
glucosidase potential activity and respiration rates were measured using colorimetric
assays and the CO_2 burst method. Cation concentrations from 1:10 ammonium acetate
extraction (K, Ca, Mg, Na) and 1:2 soil to diethylenetriaminepentaacetic acid (DPTA)
extraction solution (Zn, Mg, Cu, Fe, B, SO_4^{2-} - S) extraction were measured with
Inductively coupled plasma mass spectrometry (ICP-MS), while SOM concentration was
determined by total organic carbon-total nitrogen (TOC-TN) analyzer and composition
from liquid chromatography mass spectrometry (LC-MS), and Fourier transform ion
cyclotron mass spectrometry (FTICR-MS) analysis. Details on SOM characterization are
provided in the SI.

The abiotic core was used to determine soil structure and hydraulic properties. We used
X-ray Computed Tomography (XCT) to measure soil pore size, connectivity, distribution,
and volume, as detailed below. In addition, soil hydraulic properties were measured
using a series of instruments by The Meter Group (KSat, Hyprop, WP4C, Meter Group,
Inc., USA).

Detailed methods for data collection and analysis for XCT and FTICR-MS can be found
in the SI.

Case Study: Impacts of Wildfire Regimes on Soil; Warm Springs, Oregon

Importance of fire on ecosystems.

As the frequency of wildfires continue to increase due to anthropogenic climate change
and land management practices, it is critical to understand how fire disturbances affect
belowground biogeochemistry (Abatzoglou and Williams, 2016; Huffman and Madritch,
2018). Wildfires alter belowground processes that generally result in decreased
structural stability, microbial respiration and biomass, and SOM concentration (Certini,
2005; Doerr et al., 2006; Fairbanks et al., 2020; González-Pérez et al., 2004; Matosziuk
et al., 2020; Neary et al., 2005; Wang et al., 2012). For instance, Certini et al. (Certini et
al., 2011) reported wildfire significantly decreased total soil C, which was associated
with increased lignin degradation. Others have reported the selective loss of SOM
oxygen containing functional groups and/or with less condensed structures in response
to wildfire (González-Pérez et al., 2004). Changes in SOM concentration and
composition are coincident with structural and hydrologic changes including decreased
water infiltration and increased soil erosion (Verma and Jayakumar, 2012). Because
belowground C storage and bioavailability are dramatically altered by wildfire, it is
important to understand the nature of wildfire effects on SOM chemistry and
accessibility. Below, we describe a case study to demonstrate the utility of high-
resolution data for addressing this need.

Dataset Description.

Our primary intent in this paper is to stimulate discussion and interest in large-scale
collaborative efforts by introducing MONet. To facilitate this discussion, we present
initial results from three of the first cores in the 1000 Soils Pilot, collected in November
2021 in partnership with the Confederate Tribes of Warm Springs in Warm Springs, OR,
USA. This region typically experiences cold, snowy winters followed by warm dry
summers with a mean annual precipitation of 1778 mm and a mean annual precipitation
of 8.3 °C. Soil samples were collected from plots dominated by warm and moist type
grand fir (*A. grandis*) and snowbrush (*C. velutinus*). The unburned (UB plots) had forb
and shrub ground cover, mature living conifers, and surface litter present (>1mm) and
had no recent burn history in the last 20 years. The moderate burn frequency (MB) plots
contained shrubs and dead-standing trees but very little surface litter (<1 mm) and had
moderate severity burns in 2014 and 2020. The high frequency burn (HB) plots
consisted of a low shrub ground cover, no live or dead standing trees, and very little
surface litter (<1 mm) and had moderate severity burns in 2003, 2014 and 2020.

This subset of data is intended as a proof of concept for the utility of high-resolution
data in understanding impacts of fire burn histories on soil physio-chemical properties
and microbial functions. Though the dataset is small and does not allow for robust
statistics (n = 3), it provides a wealth of information that is unfeasible with traditional
biogeochemical approaches. With these differences in mind, we compared FTICR-MS
and XCT across burn histories and soil depth to show differences in molecular SOM
composition and soil structure.

Soil Biogeochemistry and Microbial Activity.

We first compared soil biogeochemical properties including, β-glucosidase activity (BG),
microbial biomass C (MBC) and N (MBN), and respiration at 24 and 96 hr for
differences among soil sections and burn histories (Fig 2A and 2B). Elevated microbial

biomass concentrations, respiration rates, and potential enzyme activities in the
unburned site are consistent with increased biological availability when compared to the
burned sites (Fairbanks et al., 2020). We found the highest respiration rate at UB
followed by the MB, with the lowest respiration at HB (Figure 2B). This is consistent with
other literature, showing the sterilizing effects of high frequency fires on microbial
respiration (Ajwa et al., 1999; Lombao et al., 2021). Across the first 24 hours after re-
wet, the rate of CO_2 produced was greater in the top 10 cm vs. the lowest 10 cm across
all cores. However, after 96 hours the rate of CO_2 production in soils from the bottom 10
cm was greater than the top 10 cm for all sites (Fairbanks et al., 2020). Similarly, β-
glucosidase potential activity and microbial biomass C and N were highest at UB (SI
Figure 1A and Figure 2B respectively). Lower microbial activity at the burned sites may
be related to chemical recalcitrance or physical protection of SOM related to the fires
(Ajwa et al., 1999; Lombao et al., 2021). For all the sites, respiration increased between
24 and 96 hours in the bottom 10 cm, possibly reflecting less bioavailable SOM in
deeper soils than in surface soils (Fang and Moncrieff, 2005). Corresponding changes
in soil chemistry are shown in SI Figure 2.

X-ray Computed Tomography.

To examine pore space heterogeneity across soil sections and burn histories, we used
XCT to interrogate the top and bottom 10 cm of each soil core. We compared total
porosity, pore connectivity, and mean pore volume between the top and bottom sections
of cores from UB, MB, and HB. Wildfire can impact soil structure at least to 10 cm depth
depending on its duration, frequency, and intensity (Doerr et al., 2006). For UB and HB,
we observed higher porosity and pore connectivity in the top sections when compared
to the bottom sections. This is likely related to the higher abundance of root systems in
the top 10 cm. However, there was minimal difference between total porosity or
connectivity between the top and bottom section of the MB site, possibly due to a large
root section resulting in a larger mean pore size (SI Figure 5 and SI Table 2). Across all
cores, total porosity was more variable in the top than bottom section (8.82-20.08% vs
6.52-8.49%; SI Table 2). Pore connectivity was highest in the top section at UB
(96.41%) and lowest in the bottom section at UB (70.12%); potentially indicating a
homogenizing effect of wildfire on pore network structure at MB and HB. Overall, these
data suggest that wildfire may decrease pore connectivity and total porosity in surface
soils, likely contributing to increased hydrophobicity and soil erosion, while the deeper
soils are more insulated from wildfire effects.

Figure 2: A) Microbial biomass C (MBC) and N (MBN) concentrations, B) respiration (24 and 96 hr; µg CO_2-C/g dry soil/day), C) SOM composition, and D) NOSC of SOM profiles. A and B are boxplots where the median is represented by a line, the edges of the boxes reflect 25[th] and 75[th] percentile, and the end of the whiskers reflect the minimum and maximum. C shows a principal component analysis of SOM composition, and D displays a density curve of NOSC values for each sample. For all plots, UB, MB, and HB are denoted in yellow, orange, and red respectively. In A and B, data from top and bottom sections are denoted by grey and white boxes respectively. In C, section is denoted by shape, and in D, section is denoted by dashed vs. solid lines.

Soil Organic Matter Composition.

Soil organic matter is derived from chemical, physical, and biological processes that decompose plant and animal residues into microbial metabolites and other byproducts (Jobbágy and Jackson, 2000; Köchy et al., 2015; Lehmann and Kleber, 2015; Schmidt et al., 2011; Tarnocai et al., 2009). This generates a heterogenous mixture of compounds in SOM pools. We used FTICR-MS to distinguish differences in SOM chemistry along depth profiles and across the three wildfire histories. There was a clear separation in water-extractable SOM composition between the top and bottom sections of each core as well as across burn regimes, as indicated by principal component analysis (Figure 2C). Additionally, the bottom sections exhibited more similar SOM

pools (i.e., tighter clustering) than the top sections, possibly signifying stronger impacts
of wildfire on surficial soils.

Since fire results in the oxidation of SOM to carbon dioxide (CO_2), we hypothesized that
the extent of SOM oxidation, as indicated by the nominal oxidation state of carbon
(NOSC), would be related to burn regime. In the top section, the average NOSC of
SOM increased with stronger burn regimes (-0.34, -0.29, and -0.11 respectively for UB,
MB, and HB), indicating more oxidization, while there was a binary effect of burning on
NOSC in the bottom section. MB and HB had similar mean NOSC that was higher than
UB (-0.16 vs -0.35). The presence of more oxidized SOM in burned soils, is likely
related to the incomplete oxidation of SOM and the addition of charred materials from
wildfire (Schmidt and Noack, 2000; Velasco-Molina et al., 2016; Watson et al., 2005).
While limited to three sites, changes in NOSC could be an indicator of previous fire
exposure, with the NOSC becoming more oxidized with increased high frequency fire
exposure that persists even into deeper soil horizons.

Inferred Biochemical Transformations and Microbial Metabolism of SOM.

We also analyzed FTICR-MS spectra to understand potential biochemical
transformations and microbial pathways involved in SOM decomposition. Commonly
inferred biochemical transformations such as the addition of methyl chain ($-CH_2$) groups
were abundant at all sites and depths. We also note that one of the most abundant
transformations observed was the conversion of carboxyl groups to CO_2 (greater than
1400 occurrences at each site and depth). The transformation of carboxyl groups to
CO_2 likely results in combustion byproduct associated with incomplete oxidation of SOM
during fire exposure. Within each core, the oxidation and/or addition of CO_2 accounted
for a greater proportion of transformations at depth when compared to the top section at
the same site. This is potentially related to the translocation of water-soluble oxidized
OM to deeper soil sections (Hockaday et al., 2007).

Additionally, we mapped compounds detected by FTICR-MS to microbial reference
pathways in KEGG to infer mechanisms of SOM decomposition. Across all samples,
nearly 50% of all mapped metabolites were detected in a set of eight metabolic
pathways. These included 'biosynthesis of secondary metabolites' (map01110, 11.7%),
'metabolic pathways' (map01100, 8.4%), 'biosynthesis of antibiotics' (map01130, 7.5%),
and 'microbial metabolism in diverse environments' (map01120, 4.9%) among other
pathways (SI Table 4). Notably, the degradation of aromatic compounds (map012220)
and the polycyclic aromatic hydrocarbons (PAH) degradation (map00624) were also
among pathways with the highest number of mapped metabolites (16th and 19th most
metabolites). The degradation of aromatic compounds was the most common pathway
in the core with high frequency burns (HB). The degradation of PAHs was most
common in the MB site; 1.61% of metabolites mapped to metabolic pathways in the
moderate frequency burned core were associated with the degradation of PAHs in
comparison to <1% of mapped metabolites in the unburned core.

Case Study Conclusions.

Overall, our results were consistent with literature describing wildfire impacts on soils
and provided greater understanding of molecular changes in soil organic matter
composition and structure than is possible with typical measurements. In soils with
recent fire history, we observed decreases in soil respiration, microbial biomass, and
potential enzyme activity (Certini, 2005; Fairbanks et al., 2020; Guénon et al., 2013;
Lombao et al., 2021; Neary et al., 2005; Wang et al., 2012). This was associated with
shifts in SOM composition, with higher rates of oxidation apparent in surface soils.
Deeper soils also showed increased NOSC as a function of previous fire history. By
inferring biogeochemical transformations from SOM pool composition, we also show the
possible conversion of carboxyl groups to CO_2 that is likely related to the oxidizing
effects of wildfire. Using metabolic pathway mapping, we observed that the degradation
of aromatic compounds and polycyclic aromatic hydrocarbons may also be associated
with wildfire. Finally, wildfire appeared to induce structural changes including decreased
pore connectivity and total porosity in surface soils, possibly contributing to
hydrophobicity and decreased soil infiltration which can result in greater soil erosion
(Certini, 2005; Verma and Jayakumar, 2012). The combined molecular and
microstructural changes observed in this use case provide important insight and
considerations in the management and modelling of forest ecosystems post-wildfire. As
the MONet database continues to expand, we hope to use this information to model
wildfire impacts on belowground C cycling at the regional to CONUS scale.

Towards a CONUS-scale high resolution molecular SOM database

Earth system models currently rely on soil attributes such as soil texture, moisture, and
C concentration to parameterize C fluxes. This highly simplified approach results in high
levels of uncertainty at the pore to core scale. Molecular information has the potential to
reduce this uncertainty through rapidly improving multiscale models (Blankinship et al.,
2018; Feng et al., 2021; Kramer and Chadwick, 2018; Nave et al., 2021; Quesada et al.,
2020; Rasmussen et al., 2018; Viscarra Rossel et al., 2019; Von Fromm et al., 2021; Yu
et al., 2021).

To achieve these goals, MONet is developing a database of high-resolution soil organic
matter, metagenomic, and structural information from soil cores across the CONUS. We
collaborate with existing databases and networks, as well as with individual
researchers, using standardized sample collection and high throughput processing
pipelines, metadata documentation, and raw and processed open data publication. The
focus of our data generation is on variables that are key to the parameterization of
hierarchical multiscale models beginning with microbially explicit models and
culminating in ESMs.

In this article, we present the 1000 Soils Pilot for MONet including a use case to
highlight the utility of high-resolution measurements for understanding the impacts of
wildfires on soils. While the statistical analysis is limited by the small sample number
(n=3), we observed trends in soil biogeochemical values, soil structure, and SOM
composition related to fire history and soil section that are consistent with existing
literature. We are currently accepting requests for collaboration, and we anticipate that
all data from the first set of 76 cores will be available in 2023. With the support and
collaboration of the scientific community, we aim to facilitate the next generation of

fundamental knowledge and model representations of soil C cycling, which in turn
constrain uncertainties in the global climate.

Acknowledgments: Collection of soils from the Warm Springs Reservation was funded
by a Washington State University, Vancouver Mini-Grant to SSP and KM, and a
Washington State University College of Arts and Sciences Seed Grant to SSP, KM, TC,
and EG. In addition, we would like to thank the tribal leaders from the Confederate
Tribes of Warm Springs. This research was performed on a project award
(https://dx.doi.org/10.46936/intm.proj.2021.60141/60000423) from the Environmental
Molecular Sciences Laboratory, a DOE Office of Science User Facility sponsored by the
Biological and Environmental Research program under Contract No. DE-AC05-
76RL01830. This material is based upon work supported by the U.S. Department of
Energy (DOE), Office of Science (SC), Biological and Environmental Research (BER),
Research Development and Partnership Pilot (RDPP), under Award Number(s) DE-
SC0023150.
347
This report was prepared as an account of work sponsored by an agency of the United
States Government. Neither the United States Government nor any agency thereof, nor
any of their employees, makes any warranty, express or implied, or assumes any legal
liability or responsibility for the accuracy, completeness, or usefulness of any
information, apparatus, product, or process disclosed, or represents that its use would
not infringe privately owned rights. Reference herein to any specific commercial
product, process, or service by trade name, trademark, manufacturer, or otherwise does
not necessarily constitute or imply its endorsement, recommendation, or favoring by the
United States Government or any agency thereof. The views and opinions of authors
expressed herein do not necessarily state or reflect those of the United States
Government or any agency thereof.

References:

Abatzoglou, J.T. and Williams, A.P. 2016. Impact of anthropogenic climate change on wildfire
across western US forests. Proceedings of the National Academy of Sciences 113(42),
11770-11775.
Ajwa, H.A., Dell, C.J. and Rice, C.W. 1999. Changes in enzyme activities and microbial
biomass of tallgrass prairie soil as related to burning and nitrogen fertilization. Soil
Biology and Biochemistry 31(5), 769-777.
Aponte, C., De Groot, W.J. and Wotton, B.M. 2016. Forest fires and climate change: causes,
consequences and management options. International Journal of Wildland Fire 25(8), i.
Blankinship, J.C., Berhe, A.A., Crow, S.E., Druhan, J.L., Heckman, K.A., Keiluweit, M.,
Lawrence, C.R., Marín-Spiotta, E., Plante, A.F., Rasmussen, C., Schädel, C., Schimel,
J.P., Sierra, C.A., Thompson, A., Wagai, R. and Wieder, W.R. 2018. Improving
understanding of soil organic matter dynamics by triangulating theories, measurements,
and models. Biogeochemistry 140(1), 1-13.
Bond-Lamberty, B. and Thomson, A. 2010. Temperature-associated increases in the global
soil respiration record. Nature 464(7288), 579-582.
Bond-Lamberty, B., Christianson, D.S., Malhotra, A., Pennington, S.C., Sihi, D., Aghakouchak,
A., Anjileli, H., Altaf Arain, M., Armesto, J.J., Ashraf, S., Ataka, M., Baldocchi, D.,
Andrew Black, T., Buchmann, N., Carbone, M.S., Chang, S.C., Crill, P., Curtis, P.S.,

Davidson, E.A., Desai, A.R., Drake, J.E., El-Madany, T.S., Gavazzi, M., Görres, C.M.,
Gough, C.M., Goulden, M., Gregg, J., Gutiérrez Del Arroyo, O., He, J.S., Hirano, T.,
Hopple, A., Hughes, H., Järveoja, J., Jassal, R., Jian, J., Kan, H., Kaye, J., Kominami,
Y., Liang, N., Lipson, D., Macdonald, C.A., Maseyk, K., Mathes, K., Mauritz, M., Mayes,
M.A., McNulty, S., Miao, G., Migliavacca, M., Miller, S., Miniat, C.F., Nietz, J.G., Nilsson,
M.B., Noormets, A., Norouzi, H., O'Connell, C.S., Osborne, B., Oyonarte, C., Pang, Z.,
Peichl, M., Pendall, E., Perez-Quezada, J.F., Phillips, C.L., Phillips, R.P., Raich, J.W.,
Renchon, A.A., Ruehr, N.K., Sánchez-Cañete, E.P., Saunders, M., Savage, K.E.,
Schrumpf, M., Scott, R.L., Seibt, U., Silver, W.L., Sun, W., Szutu, D., Takagi, K., Takagi,
M., Teramoto, M., Tjoelker, M.G., Trumbore, S., Ueyama, M., Vargas, R., Varner, R.K.,
Verfaillie, J., Vogel, C., Wang, J., Winston, G., Wood, T.E., Wu, J., Wutzler, T., Zeng, J.,
Zha, T., Zhang, Q. and Zou, J. 2020. COSORE: A community database for continuous
soil respiration and other soil-atmosphere greenhouse gas flux data. Global Change
Biology 26(12), 7268-7283.
Bowman, M.M., Heath, A.E., Varga, T., Battu, A.K., Kew, W., Chu, R.K., Toyoda, J., Qafoku, O.,
Bargar, J., Hess, N., Leichty, S., Townsend, A., Sconzo, N., Miller, P., Anderson, J.,
Stohel, I., Rosenstock, M., Lawver, A., Hauchambe, R., Clendinen, C. and Graham, E.B.
2022 1000 Soils Pilot Dataset. Laboratory, E.M.S. (ed), Zenodo.
Bradford, M.A., Wieder, W.R., Bonan, G.B., Fierer, N., Raymond, P.A. and Crowther, T.W.
2016. Managing uncertainty in soil carbon feedbacks to climate change. Nature Climate
Change 6(8), 751-758.
Bray, R.H. and Kurtz, L.T. 1945. Determination of total, organic, and available forms of
phosphorus in soils. Soil science 59(1), 39-46.
Bremner, J.t. 1965. Inorganic forms of nitrogen. Methods of Soil Analysis: Part 2 Chemical and
Microbiological Properties 9, 1179-1237.
Brookes, P., Landman, A., Pruden, G. and Jenkinson, D. 1985. Chloroform fumigation and the
release of soil nitrogen: a rapid direct extraction method to measure microbial biomass
nitrogen in soil. Soil biology and biochemistry 17(6), 837-842.
Certini, G. 2005. Effects of fire on properties of forest soils: a review. Oecologia 143(1), 1-10.
Certini, G., Nocentini, C., Knicker, H., Arfaioli, P. and Rumpel, C. 2011. Wildfire effects on soil
organic matter quantity and quality in two fire-prone Mediterranean pine forests.
Geoderma 167-168, 148-155.
Crowther, T.W., Riggs, C., Lind, E.M., Borer, E.T., Seabloom, E.W., Hobbie, S.E., Wubs, J.,
Adler, P.B., Firn, J., Gherardi, L., Hagenah, N., Hofmockel, K.S., Knops, J.M.H.,
Mcculley, R.L., Macdougall, A.S., Peri, P.L., Prober, S.M., Stevens, C.J. and Routh, D.
2019. Sensitivity of global soil carbon stocks to combined nutrient enrichment. Ecology
Letters 22(6), 936-945.
Crowther, T.W., Todd-Brown, K.E.O., Rowe, C.W., Wieder, W.R., Carey, J.C., Machmuller,
M.B., Snoek, B.L., Fang, S., Zhou, G., Allison, S.D., Blair, J.M., Bridgham, S.D., Burton,
A.J., Carrillo, Y., Reich, P.B., Clark, J.S., Classen, A.T., Dijkstra, F.A., Elberling, B.,
Emmett, B.A., Estiarte, M., Frey, S.D., Guo, J., Harte, J., Jiang, L., Johnson, B.R., Kröel-
Dulay, G., Larsen, K.S., Laudon, H., Lavallee, J.M., Luo, Y., Lupascu, M., Ma, L.N.,
Marhan, S., Michelsen, A., Mohan, J., Niu, S., Pendall, E., Peñuelas, J., Pfeifer-Meister,
L., Poll, C., Reinsch, S., Reynolds, L.L., Schmidt, I.K., Sistla, S., Sokol, N.W., Templer,
P.H., Treseder, K.K., Welker, J.M. and Bradford, M.A. 2016. Quantifying global soil
carbon losses in response to warming. Nature 540(7631), 104-108.
Dahnke, W. and Johnson, G.V. 1990. Testing soils for available nitrogen. Soil testing and plant
analysis 3, 127-139.
Dane, J.H. and Topp, C.G. (2020) Methods of soil analysis, Part 4: Physical methods, John
Wiley & Sons.

Doerr, S.H., Shakesby, R.A., Blake, W.H., Chafer, C.J., Humphreys, G.S. and Wallbrink, P.J. 2006. Effects of differing wildfire severities on soil wettability and implications for hydrological response. Journal of Hydrology 319(1-4), 295-311.

Fairbanks, D., Shepard, C., Murphy, M., Rasmussen, C., Chorover, J., Rich, V. and Gallery, R. 2020. Depth and topographic controls on microbial activity in a recently burned sub-alpine catchment. Soil Biology and Biochemistry 148, 107844.

Fang, C. and Moncrieff, J.B. 2005. The variation of soil microbial respiration with depth in relation to soil carbon composition. Plant and Soil 268(1), 243-253.

Feng, S., Lauvaux, T., Williams, C.A., Davis, K.J., Zhou, Y., Baker, I., Barkley, Z.R. and Wesloh, D. 2021. Joint CO2 Mole Fraction and Flux Analysis Confirms Missing Processes in CASA Terrestrial Carbon Uptake Over North America. Global Biogeochemical Cycles 35(7).

González-Pérez, J.A., González-Vila, F.J., Almendros, G. and Knicker, H. 2004. The effect of fire on soil organic matter—a review. Environment international 30(6), 855-870.

Graham, E.B. and Hofmockel, K.S. 2022. Ecological stoichiometry as a foundation for omics-enabled biogeochemical models of soil organic matter decomposition. Biogeochemistry 157(1), 31-50.

Guénon, R., Vennetier, M., Dupuy, N., Roussos, S., Pailler, A. and Gros, R. 2013. TRENDS IN RECOVERY OF MEDITERRANEAN SOIL CHEMICAL PROPERTIES AND MICROBIAL ACTIVITIES AFTER INFREQUENT AND FREQUENT WILDFIRES. Land Degradation & Development 24(2), 115-128.

Hockaday, W.C., Grannas, A.M., Kim, S. and Hatcher, P.G. 2007. The transformation and mobility of charcoal in a fire-impacted watershed. Geochimica et Cosmochimica Acta 71(14), 3432-3445.

Huffman, M.S. and Madritch, M.D. 2018. Soil microbial response following wildfires in thermic oak-pine forests. Biology and Fertility of Soils 54(8), 985-997.

Jobbágy, E.G. and Jackson, R.B. 2000. The vertical distribution of soil organic carbon and its relation to climate and vegetation. Ecological Applications 10(2), 423-436.

Klute, A. and Dirksen, C. (1986) Methods of Soil Analysis, pp. 687-734.

Köchy, M., Hiederer, R. and Freibauer, A. 2015. Global distribution of soil organic carbon – Part 1: Masses and frequency distributions of SOC stocks for the tropics, permafrost regions, wetlands, and the world. SOIL 1(1), 351-365.

Kramer, M.G. and Chadwick, O.A. 2018. Climate-driven thresholds in reactive mineral retention of soil carbon at the global scale. Nature Climate Change 8(12), 1104-1108.

Lawrence, C.R., Beem-Miller, J., Hoyt, A.M., Monroe, G., Sierra, C.A., Stoner, S., Heckman, K., Blankinship, J.C., Crow, S.E., Mcnicol, G., Trumbore, S., Levine, P.A., Vindušková, O., Todd-Brown, K., Rasmussen, C., Hicks Pries, C.E., Schädel, C., Mcfarlane, K., Doetterl, S., Hatté, C., He, Y., Treat, C., Harden, J.W., Torn, M.S., Estop-Aragonés, C., Asefaw Berhe, A., Keiluweit, M., Della Rosa Kuhnen, Á., Marin-Spiotta, E., Plante, A.F., Thompson, A., Shi, Z., Schimel, J.P., Vaughn, L.J.S., Von Fromm, S.F. and Wagai, R. 2020. An open-source database for the synthesis of soil radiocarbon data: International Soil Radiocarbon Database (ISRaD) version 1.0. Earth System Science Data 12(1), 61-76.

Lehmann, J. and Kleber, M. 2015. The contentious nature of soil organic matter. Nature 528, 60-68.

Lombao, A., Barreiro, A., Fontúrbel, M.T., Martín, A., Carballas, T. and Díaz-Raviña, M. 2021. Effect of repeated soil heating at different temperatures on microbial activity in two burned soils. Science of The Total Environment 799, 149440.

Malhotra, A., Todd-Brown, K., Nave, L.E., Batjes, N.H., Holmquist, J.R., Hoyt, A.M., Iversen, C.M., Jackson, R.B., Lajtha, K., Lawrence, C., Vindušková, O., Wieder, W., Williams, M., Hugelius, G. and Harden, J. 2019. The landscape of soil carbon data: Emerging

questions, synergies and databases. Progress in Physical Geography: Earth and
Environment 43(5), 707-719.
Matosziuk, L.M., Gallo, A., Hatten, J., Bladon, K.D., Ruud, D., Bowman, M., Egan, J., Heckman,
K., SanClements, M., Strahm, B. and Weiglein, T. 2020. Short-Term Effects of Recent
Fire on the Production and Translocation of Pyrogenic Carbon in Great Smoky
Mountains National Park. Frontiers in Forests and Global Change 3.
Nave, L.E., Bowman, M., Gallo, A., Hatten, J.A., Heckman, K.A., Matosziuk, L., Possinger, A.R.,
SanClements, M., Sanderman, J., Strahm, B.D., Weiglein, T.L. and Swanston, C.W.
2021. Patterns and predictors of soil organic carbon storage across a continental-scale
network. Biogeochemistry, 1-22.
Neary, D.G., Ryan, K.C. and DeBano, L.F. 2005 Wildland fire in ecosystems: effects of fire on
soils and water, U.S. Department of Agriculture, Forest Service, Rocky Mountain
Research Station.
NRCS, U.N.U.   United States Department of Agriculture Natural Resource Conservation
Service (2019) Web Soil Survey. Map generated over River Highlands State Park in
Cromwell, Connecticut. Database website: https://websoilsurvey. sc. egov. usda.
gov/App/HomePage. htm.
Olsen, S., Sommers, L. and Page, A. 1982. Methods of soil analysis. Part 2(1982), 403-430.
Quesada, C.A., Paz, C., Oblitas Mendoza, E., Phillips, O.L., Saiz, G. and Lloyd, J. 2020.
Variations in soil chemical and physical properties explain basin-wide Amazon forest soil
carbon concentrations. Soil 6(1), 53-88.
Rasmussen, C., Heckman, K., Wieder, W.R., Keiluweit, M., Lawrence, C.R., Berhe, A.A.,
Blankinship, J.C., Crow, S.E., Druhan, J.L., Hicks Pries, C.E., Marin-Spiotta, E., Plante,
A.F., Schädel, C., Schimel, J.P., Sierra, C.A., Thompson, A. and Wagai, R. 2018.
Beyond clay: towards an improved set of variables for predicting soil organic matter
content. Biogeochemistry 137(3), 297-306.
Schmidt, M.W.I. and Noack, A.G. 2000. Black carbon in soils and sediments: analysis,
distribution, implications, and current challenges. Global biogeochemical cycles 14(3),
777-793.
Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber,
M., Kögel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P.,
Weiner, S. and Trumbore, S.E. 2011. Persistence of soil organic matter as an
ecosystem property. Nature 478, 49-56.
Tarnocai, C., Canadell, J.G., Schuur, E.A.G., Kuhry, P., Mazhitova, G. and Zimov, S. 2009.
Soil organic carbon pools in the northern circumpolar permafrost region. Global
Biogeochemical Cycles 23(2), n/a-n/a.
Todd-Brown, K.E.O., Abramoff, R.Z., Beem-Miller, J., Blair, H.K., Earl, S., Frederick, K.J., Fuka,
D.R., Guevara Santamaria, M., Harden, J.W., Heckman, K., Heran, L.J., Holmquist, J.R.,
Hoyt, A.M., Klinges, D.H., Lebauer, D.S., Malhotra, A., Mcclelland, S.C., Nave, L.E.,
Rocci, K.S., Schaeffer, S.M., Stoner, S., Van Gestel, N., Von Fromm, S.F. and Younger,
M.L. 2022. Reviews and syntheses: The promise of big diverse soil data, moving
current practices towards future potential. Biogeosciences 19(14), 3505-3522.
Velasco-Molina, M., Berns, A.E., Macías, F. and Knicker, H. 2016. Biochemically altered
charcoal residues as an important source of soil organic matter in subsoils of fire-
affected subtropical regions. Geoderma 262, 62-70.
Verma, S. and Jayakumar, S. 2012. Impact of forest fire on physical, chemical and biological
properties of soil: A review. proceedings of the International Academy of Ecology and
Environmental Sciences 2(3), 168.
Viscarra Rossel, R.A., Lee, J., Behrens, T., Luo, Z., Baldock, J. and Richards, A. 2019.
Continental-scale soil carbon composition and vulnerability modulated by regional
environmental controls. Nature Geoscience 12(7), 547-552.

Von Fromm, S.F., Hoyt, A.M., Lange, M., Acquah, G.E., Aynekulu, E., Berhe, A.A., Haefele,
S.M., McGrath, S.P., Shepherd, K.D., Sila, A.M., Six, J., Towett, E.K., Trumbore, S.E.,
Vågen, T.-G., Weullow, E., Winowiecki, L.A. and Doetterl, S. 2021. Continental-scale
controls on soil organic carbon across sub-Saharan Africa. SOIL 7(1), 305-332.
Wang, Q., Zhong, M. and Wang, S. 2012. A meta-analysis on the response of microbial
biomass, dissolved organic matter, respiration, and N mineralization in mineral soil to fire
in forest ecosystems. Forest Ecology and Management 271, 91-97.
Waring, B.G., Sulman, B.N., Reed, S., Smith, A.P., Averill, C., Creamer, C.A., Cusack, D.F.,
Hall, S.J., Jastrow, J.D., Jilling, A., Kemner, K.M., Kleber, M., Liu, X.J.A., Pett-Ridge, J.
and Schulz, M. 2020. From pools to flow: The PROMISE framework for new insights on
soil carbon cycling in a changing world. Global Change Biology 26(12), 6631-6643.
Watson, J.G., Chow, J.C. and Chen, L.W.A. 2005. Summary of Organic and Elemental
Carbon/Black Carbon Analysis Methods and Intercomparisons. Aerosol and Air Quality
Research 5(1), 65-102.
Wieder, W.R., Pierson, D., Earl, S., Lajtha, K., Baer, S.G., Ballantyne, F., Berhe, A.A., Billings,
S.A., Brigham, L.M., Chacon, S.S., Fraterrigo, J., Frey, S.D., Georgiou, K., De Graaff,
M.-A., Grandy, A.S., Hartman, M.D., Hobbie, S.E., Johnson, C., Kaye, J., Kyker-
Snowman, E., Litvak, M.E., Mack, M.C., Malhotra, A., Moore, J.A.M., Nadelhoffer, K.,
Rasmussen, C., Silver, W.L., Sulman, B.N., Walker, X. and Weintraub, S. 2021.
SoDaH: the SOils DAta Harmonization database, an open-source synthesis of soil data
from research networks, version 1.0. Earth System Science Data 13(5), 1843-1854.
Wilkinson, M.D., Dumontier, M., Aalbersberg, I.J., Appleton, G., Axton, M., Baak, A., Blomberg,
N., Boiten, J.-W., Da Silva Santos, L.B., Bourne, P.E., Bouwman, J., Brookes, A.J.,
Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C.T., Finkers, R.,
Gonzalez-Beltran, A., Gray, A.J.G., Groth, P., Goble, C., Grethe, J.S., Heringa, J., 'T
Hoen, P.A.C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S.J., Martone, M.E., Mons,
A., Packer, A.L., Persson, B., Rocca-Serra, P., Roos, M., Van Schaik, R., Sansone, S.-
A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M.A., Thompson, M., Van
Der Lei, J., Van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P.,
Wolstencroft, K., Zhao, J. and Mons, B. 2016. The FAIR Guiding Principles for scientific
data management and stewardship. Scientific Data 3(1), 160018.
Witt, C., Gaunt, J.L., Galicia, C.C., Ottow, J.C. and Neue, H.-U. 2000. A rapid chloroform-
fumigation extraction method for measuring soil microbial biomass carbon and nitrogen
in flooded rice soils. Biology and Fertility of Soils 30(5), 510-519.
Yu, W., Weintraub, S.R. and Hall, S.J. 2021. Climatic and geochemical controls on soil carbon
at the continental scale: interactions and thresholds. Global Biogeochemical Cycles
35(3), e2020GB006781.
Zhao, Q., Thompson, A.M., Callister, S.J., Tfaily, M.M., Bell, S.L., Hobbie, S.E. and Hofmockel,
K.S. 2022. Dynamics of organic matter molecular composition under aerobic
decomposition and their response to the nitrogen addition in grassland soils. Science of
the Total Environment 806, 150514.
572