Title page

Influence of the mutation load on the genomic composition of hybrids between outcrossing and self-fertilizing species

Fréderic Fyon[1, *], Waldir M. Berbel-Filho[2]

[1] Department of Biology, Royal Holloway University of London, Egham, TW20 0EX, UK.

[2] Department of Biology, University of Oklahoma, Norman, Oklahoma, USA.

* Corresponding author: frederic.fyon@gmail.com

Keywords: Hybridization **–** Mating systems – Mutational load – Recombination – Mutation rate – Self-fertilization

Abstract

Hybridization is a natural process whereby two diverging evolutionary lineages reproduce and create offspring of mixed ancestry. Differences in mating systems (e.g., self-fertilization and outcrossing) are expected to affect the direction and extent of hybridization and introgression in hybrid zones. Among other factors, selfers and outcrossers are expected to differ in their mutation loads. This has been studied both theoretically and empirically; however, conflicting predictions have been made on the effects mutation loads of parental species with different mating systems can have on the genomic composition of hybrids. Here we develop a multi-locus, selective model to study how the different mutation load built up in selfers and outcrossers as a result of selective interference and homozygosity impact the long-term genetic composition of hybrid populations. Notably, our results emphasize that genes from the parental population with lesser mutation load get rapidly over-represented in hybrid genomes, regardless of the hybrids own mating system. When recombination tends to be more important than mutation, outcrossers' genomes tend to be of higher quality and prevail. When recombination is small, however, selfers' genomes may reach higher quality than outcrossers' genomes and prevail. Taken together these results provide concrete insights into one of the multiple factors influencing hybrid genome composition and introgression patterns in hybrid zones with species containing species with different mating systems.

1. Introduction

Hybrid individuals are defined as progeny of mixed ancestry; they bear genes inherited from divergent evolutionary lineages. Introgressive hybridization (i.e., formation of hybrids with backcrossing) functions as bridges between divergent lineages, allowing gene flow and shared evolutionary history. Identifying the processes shaping the evolutionary fate of hybrid populations is a fundamental task for evolutionary biology, as well as for conservation, and wildlife management.

The possibility of hybridization happening, along with its ecological and genomic consequences, depends, among other factors, on the mating systems and reproductive biology of the parental species and their hybrids. Most plants and many animals are simultaneous hermaphrodites (JARNE AND AULD 2006; THE TREE OF SEX CONSORTIUM 2014). In these species, self-fertilization (selfing) – offspring produced by the fusion of gametes stemming from different meiosis in the same individual – is a possibility. Selfing contrasts with outcrossing as in the latter the progeny is produced by the fusion of gametes proceeding from different meiosis in different individuals (hermaphrodites or not). Mating systems are usually defined in three main categories according to the selfing rate: obligate outcrossing (selfing rate ≤ 20%), predominantly selfing (selfing rate > 80%), and mixed-mating (selfing rate > 20% and ≤ 80%) (SHIMIZU AND TSUCHIMATSU 2015). It is only recently that attention has been brought on the effects that different mating systems have on the occurrence of hybridization and on the genomic composition of hybrid populations (HU 2015; KIM *et al.* 2018; PICKUP *et al.* 2019).

Hybrid zones are regions of contact between divergent species/lineages that often produce individuals of admixed ancestry through interspecific mating. Hybrid zones between species with

different mating systems (e.g., predominantly selfing and obligate outcrossing) are commonly

reported in plants (RUHSAM *et al.* 2011; PICKUP *et al.* 2019; OSTEVIK *et al.* 2021), but examples

are scarce in animals (BERBEL-FILHO *et al.* 2021). Some predictions on how reproductive

systems (BRANDVAIN AND HAIG 2005) and reproductive timing (MARTIN AND WILLIS 2007;

BERBEL-FILHO *et al.* 2021), along with other ecological factors (BUSCH *et al.* 2022), can

influence the occurrence and outcome of hybridization in these cases have been extensively

provided. On the other hand, few studies have tried to address how mating systems differences

influence (i) the quality of each parental genomes inherited by first-generation hybrids (F1), and

(ii) how the genetic composition (defined in terms of genes' species-of-origin) of hybrids

evolves in the long-term as a result (HU 2015; KIM *et al.* 2018; PICKUP *et al.* 2019). Disparate,

sometimes contradictory predictions on these points have been made; the present work was

designed to disentangle these and provide a unified framework for the influence of mutation load

on hybrid ancestry.

Predictions concerning hybrids' genetic composition and evolution have focused on mutation

load. They rely on the idea that species with different mating systems are expected to harbor

different mutation loads (ARUNKUMAR *et al.* 2015; PICKUP *et al.* 2019). High homozygosity

caused by long-term selfing means that recombination is – most of the time – inefficient at

shuffling genetic combinations, which results in higher rates of hitch-hiking, background

selection, and selective interference (WRIGHT *et al.* 2013). That is, selection is expected to be

less efficient in selfing populations, which are thus predicted to have more slightly deleterious

alleles become fixed by random genetic drift (CHARLESWORTH *et al.* 1993). This relates to the

classic argument that the absence of effective recombination in asexual and self-fertilizing

species should lead them to extinction due to mutational meltdown (GABRIEL *et al.* 1993; LYNCH

et al. 1993; LYNCH *et al.* 1995). Altogether, this predicts that genomes of self-fertilizing species

tend to be of worse quality. On the other hand, however, highly deleterious recessive alleles are

expected to be purged by selection in selfers as they are "exposed" to selection by high

homozygosity, while in outcrossers they can be maintained at low frequencies as they are partly

hidden from selection in heterozygotes (WANG *et al.* 1999). In other words, here selection is

predicted to be more efficient in self-fertilizing species, resulting in genomes of better quality in

selfers. Overall, selfers' genomes are expected to harbor more fixed slightly deleterious,

codominant mutations and less strongly deleterious, recessive mutations (ARUNKUMAR *et al.*

2015). How are these two alternative mutation loads expected to interact and determine genetic

composition and evolution of hybrid populations?

The answer to this question is likely to depend on several parameters of the selection processes

in parental species, but also on the hybrid mating system. Highly deleterious, recessive mutations

coming from outcrossing ancestors may rapidly be purged by selection if the hybrids self-

fertilize (as these mutations rapidly become homozygous and visible to selection), taking

together any linked genes. This process is likely to limit the introgression of outcrossers'

genomes into selfers' (PICKUP *et al.* 2019). On the other hand, the many fixed slightly

deleterious, codominant mutations coming from the selfing ancestors should be purged out in an

outcrossing hybrid, this time restricting introgression of self-fertilizers' genomes into

outcrossers' (PICKUP *et al.* 2019). Under this argument, everything happens as if each parental

genome was "adapted" to the mating system it has evolved in and tends to fare worse in

alternative mating systems. This raises the question, does the mating system of the hybrid

population have a critical influence on the evolution of hybrid genome composition, or is this

simply determined by the respective quality of the parental populations' genomes?

Here, we aim at answering these questions by building a multi-locus model allowing for

selection, recombination, mutation, and drift to interplay and modify simple genomes. Running

numerical simulations for a diversity of scenarios, we were able to study the effects of selective

interference and homozygosity on (1) the relative mutation loads of self-fertilizing and

outcrossing populations, and (2) the long-term genetic composition of hybrid populations

deriving from parental populations with different mating systems.

First, our work allows to confirm some of the classic arguments on mutation load in self-

fertilizing species. As another theoretical study showed recently (SIANTA *et al.* 2022), how

mutation loads in self-fertilizing and outcrossing populations compare with one another strongly

depends on the recombination rate between selected loci. At high recombination, outcrossers

appear to be more efficient at limiting mutation load, as frequent reshuffling of genes

significantly reduce selective interference and fixation of deleterious mutations. However, at low

recombination regime, outcrossers suffer from selective interference just as much as selfers. The

only difference between the two then reside in the homozygosity of selfers that purge strongly

deleterious recessive mutations, resulting in them having a lesser mutation load relative to

outcrossers.

Second, we were able to determine the parameters that have a notable influence on the genetic

composition and evolution of the hybrid population. Interestingly, we show that the mating

system of the hybrid population does not largely influence the outcome of the simulations.

Hybrid genetic composition in the long term appears to be primarily dictated by the relative

mutation load of the parental species: genomes of worse quality get preferentially eliminated.

This result provides a new key to understand empirical data related to hybrid genome

composition and patterns of introgression in hybrid zones between selfer and outcrosser species.

Depending on genomic (recombination rates, mutation rates), ecological (intensity of selection),

and demographical (population size, bottlenecks, colonization events) parameters, one can now

predict how mutation load may have built in the parental species, and how much this could have

contributed to hybrid genome composition.

2. Methods

We used SLiM3 (HALLER AND MESSER 2019) software to code and run the model. SLiM3 has

been used in a diversity of population genetics studies as it provides a resource-effective and

flexible computing framework. Because the model is of a stochastic nature, each simulation is

repeated 100 times, and mean values as well as standard deviations are calculated from these 100

iterations. The code used here is available on the following GitHub repository:

https://github.com/FredericFyon/Mating-systems.

2.1. Theoretical populations

Our aim here is to follow the genetic alleles at different loci in populations with different mating

systems, along with their hybrids. To do so, we always model three populations: (1) an

exclusively or mostly self-incompatible (SI) population that reproduces via obligate outcrossing

(outcrossers); (2) a self-compatible (SC) population that reproduces exclusively or mostly by

means of self-fertilization (selfers); (3) a population born by hybridization of the two previous

ones. Populations do not interact with each other (except at the one generation where

hybridization happens) and do not compete. Each population size is set at N_{pop}. In the following,

we call σ_i the rate of self-fertilization of a given population *i*. We refer to the self-fertilization

rate of the outcrossing, self-fertilizing, and hybrid populations as $\sigma_o \leq 0.1$, $\sigma_s \geq 0.9$ and σ_h

respectively.

2.2. Genetic architecture

The genome of all individuals in the three populations consists of N_{loc} loci, equally distributed

along a chromosome. We note r the recombination rate between two adjacent loci. We consider

here an infinite-allele model: each mutation occurring is unique, all loci potentially have an

infinite number of alleles. In this model we only consider deleterious mutations and note s the

detrimental effect of a particular mutation on the fitness of the host individual.

2.3. Life Cycle

At each generation, selection is modelled by choosing the parents of every offspring born at this

generation. To do that, we first determine if an offspring was produced by self-fertilization or

not. With probability σ, it is produced by self-fertilization. In that case, we randomly draw one

parent from the distribution of individuals of the same population at the previous generation. In

contrast, with probability 1 - σ, the offspring is producing by outcrossing: we randomly draw two

parents. Drawn parents are then accepted with a probability equal to their fitness w_i. We assume

fitness to be purely multiplicative among loci (we assume no epistasis): $w_i = \prod_j w_{ij}$, with $w_{i,j}$

being the contribution of locus $j \in [1, N_{loc}]$ to the fitness of individual i. $w_{i,j}$ depends on (1) the

detrimental effect s_{ij}^1 and s_{ij}^2 of the alleles present on the two homologous chromosomes at locus

j of individual i, and on (2) the dominance coefficients h_{ij}^1 and h_{ij}^2 of these two alleles. Following

standard population genetics design, if the alleles are not identical-by-descent (that is, if the

individual is heterozygote), $w_{i,j}$ is calculated as:

$$w_{i,j} = \prod_{k=1}^{2} (1 - h_{ij}^k s_{ij}^k)$$

If they are identical by-descent (homozygote), then $w_{i,j}$ simply is: $w_{i,j} = 1 - s_{ij}$ (with $s_{ij}^1 =$

$s_{ij}^2 = s_{ij}$).

Following empirical data suggesting that the dominance of a deleterious mutation is

negatively correlated with its detrimental effect (SIMMONS AND CROW 1977; PHADNIS AND FRY

2005; AGRAWAL AND WHITLOCK 2011), we assume in this model a negative exponential

relationship between s and h:

$$h_{ij}^k = \frac{e^{-\alpha s_{ij}^k}}{2}$$

with α being an arbitrarily chosen parameter that allows to modulate this relationship. Here, we

will always assume that $\alpha = 7$ for simplicity. This allows to approximately match empirical data

of mean dominance of mildly deleterious mutations ($s = 0.1$) being around 0.25 (MANNA *et al.*

2011; however see SIMMONS AND CROW 1977; GARCÍA-DORADO & CABALLERO 2000; for

example, for other estimates). The exponential function allows for a long-tailed asymptotic

distribution of dominance effects, letting highly deleterious mutations to have a very low but

strictly positive dominance. Also, the function means that slightly deleterious mutations

approach codominance. This allows us to incorporate in the model deleterious mutations in the

whole range of dominance values from almost complete recessivity to almost codominance.

Overall, this allows the whole selective process to be determined by only one parameter: the

distribution of detrimental effects of deleterious mutations.

In addition to selection, the life cycle includes recombination and mutation.

Recombination happens at a rate r being any two adjacent loci. A recombination event

exchanges the composition of the two homologous chromosomes in an individual from the

recombination point to the downstream end on the chromosomes. Of course, multiple

recombination events can take place during any meiotic event, leading to potentially mosaic

inherited chromosomes.

Mutations happen with probability μ per locus per chromosome per individual. When a

mutation occurs, we draw its detrimental effect from a negative exponential distribution of mean

1/λ. This means that there is a vast majority of slightly deleterious, codominant mutations

arising, along with some mildly deleterious, recessive mutations and very few highly deleterious,

strongly recessive mutations.

2.4. Hybridization

At first, we consider only the parental populations of outcrossers and selfers. These reproduce

within themselves for a certain number of generations t_1. Then, a hybridization event happens.

Hybridization is modelled by created a third, hybrid population. This population is created by

drawing for each individual one parent from each parental population. As a result, at $t_1 + 1$,

every individual of the hybrid population bears one chromosome from one parental population,

and the other from the other parental population. After that initial hybridization event, we

consider that there is no additional gene flow from the parental populations (as backcrossing

would blur the sole effect of mutation load). We let the hybrid population reproduce within itself

and evolve in the absence of additional mutation. This allows us to determine the evolutionary

fate of the parental genomes within the hybrid population.

2.5. Parameter values

Unless specified otherwise, simulations throughout this work have been run with the following

parameter values: *Npop = 500*, *Nloc = 100*, $\lambda = 10$, $\sigma_o = 0$ (purely outcrossing population) and

$\sigma_s = 1$ (purely self-fertilizing population). We discuss later the influence of these parameters and

provide   examples   with   some   other   values   in   the   Supplementary   Material.

3. Results

3.1. Mutation Load in Outcrossers and Selfers

We present in the Supplementary Material some results of how the mutation load builds up in outcrossing and self-fertilizing populations. We only rapidly present these here as they are already known patterns (SIANTA *et al.* 2022).

Following classic population genetics arguments, our model shows that outcrossers' genomes present higher heterozygosity, more deleterious mutations, and lower dominance coefficients than selfers' genomes.

Our results also emphasize that the relative rates of fixation of deleterious mutations in outcrossers and selfers highly depend on mutation and recombination rates. Higher mutation rates mean more fixations. Interestingly, we see that when recombination rate is large, selfers accumulate more fixed deleterious mutations, as they suffer from selective interference: high rates of homozygosity mean recombination is mostly inefficient at reshuffling genetic associations. However, when recombination rates are small, outcrossers suffer from selective interference just as much as selfers. Consequently, outcrossers may accumulate more deleterious mutations than selfers, as they suffer from a reduction of selection efficacy associated with the detrimental effect of mutations being partially hidden in heterozygotes.

Overall, this allows us to determine areas of the parameter space where outcrossers' genomes are of better quality (high recombination rates), and areas of the parameter space where selfers' genomes are of better quality (low recombination rates). Parting from here, we can investigate how these genomes fare when put together in hybrids.

 3.2. Genetic Composition of Hybrids between Outcrossers and Selfers

In the following, we look at the proportions of genes in the hybrid population coming from each

of the parental populations. On Fig. 1, we illustrate the genomic composition of hybrid

populations, considering a case where $\mu = 10^{-5}$ and where the parental outcrossing and self-

fertilizing populations have diverged during 10,000 generations prior to hybridization. We

associate these patterns with the mutation load built-up in the parental populations.

Fig. 1*: Consequences of the accumulation of mutation load in the parental populations on the*

genomic composition of hybrids. Here, an outcrossing (purple) and a self-fertilizing (red)

populations evolve for 10,000 generations. First column shows the number of fixed deleterious

Figure 1 exemplifies three different scenarios. In scenario (i), the mutation rate and time left for

mutation accumulation are so small that mutation load barely builds up. Outcrossing and self-

fertilizing populations accumulate similar amounts of deleterious mutations. As a result, they do

not differ much in population mean fitness. The outcrossing population has lower fitness due to

higher frequencies of highly deleterious, strongly recessive mutations, but the difference is

marginal. As a result, in the hybrid population around half of the time the genes of the

outcrossing parents prevail, half the time the genes of the self-fertilizing parents prevail. We see

that there is no case of mixture of genes of the two populations. This is because the

recombination rate is too low here. Eventually, one haplotype goes to fixation, and because the

recombination rate is so small this haplotype is one chromosome of either one of the parental

populations. The two populations having similar mean fitnesses; in fact, they both have many

chromosomes without any deleterious mutations. Most probably, one of these "optimal"

chromosomes goes to fixation in the hybrid population, and there is almost equal probability that

the "optimal" haplotype that goes to fixation is from one or the other parental populations.

The results of scenario (i) strongly rely on the fact that deleterious mutations are rare, so that

many genomes are completely free of them or harbor very few of them. In the following

scenarios, we consider alternative cases where there are many detrimental mutations around,

either fixed or segregating. Here, we use mutation rate values that are not realistic but allow us to

simulate a scenario with many mutations emerge, which is arguably more realistic than the

scenario (i) where most genomes are "perfect".

In scenario (ii) there is significant accumulation of deleterious mutations. Because the

recombination rate is still low, both the outcrossing and the self-fertilizing populations suffer

from selective interference. Heterozygosity in the outcrossing populations helps deleterious

mutations go to fixation; as a result, we see in this scenario that the outcrossing population

accumulates more fixed deleterious mutations than the self-fertilizing population. This translates

in the outcrossing population having a much lower population mean fitness after 10,000

generations than the outcrossing population. In this case, we see that the genes from the self-

fertilizing population always prevail in hybrids. Again, because recombination rate is small, the

haplotype that goes to fixation is an entire chromosome of one of the parental populations, here

always the self-fertilizing population since its genomes are of much better quality.

In scenario (iii), high recombination rates mean that the outcrossing population suffer much less

from selective interference than the self-fertilizing population. Therefore, it accumulates less

fixed deleterious mutations, and have a somewhat better fitness after 10,000 generations than the

self-fertilizing population. In this case, we see that on average the genes from the outcrossing

population prevails. However, we see that hybrid genomes are a mixture of genes from the two

parental populations. This is because here high recombination rates means that a mosaic

haplotype goes to fixation. Because the genomes of the outcrossing population are of better

quality here, this mosaic tends to contain more genes from the outcrossing parents than genes

from the self-fertilizing parent.

3.3. Effects of hybrid mating system on hybrid genome composition

On Fig. 1, the hybrid population always outcross to reproduce. To gain insights into how the

mating system of the hybrid population impacts its genome composition, we show on Fig. 2

histograms of the part of the genome of the hybrid populations that stems from the self-fertilizing

parental populations for different values of r, μ, but also σ_h, the probability of hybrids to

reproduce via self-fertilization.

The mating system of the hybrid seems to have only a marginal effect on the genomic

composition of the hybrids. Regardless whether hybrids self-fertilize or outcross, it does not

change which genes tend to prevail. When the genomes of the two parental populations are of

similar quality ($\mu = 1.10^{-5}$), the probability for each type of genome to prevail remains around

one half whether the hybrids self-fertilize or outcross. When the genomes of the self-fertilizing

parents tend to prevail ($\mu = 1.10^{-4}$, $r = 10^{-6}$), they prevail also in both cases. Finally, when the

genomes of the outcrossing parents tend to prevail ($\mu = 1.10^{-4}$, $r = 10^{-1}$), hybrid genomes are

biased towards outcrossing parents' heredity again in both cases.

The effect of the mating system mostly appears when $r = 10^{-1}$. In that case, there is a

fundamental difference depending on how the hybrids reproduce. If the hybrids outcross, then

their genomes are frequently recombined. As evoked earlier, this means that hybrids genomes

become a mixture of the two parental populations, instead of being only composed of genes from

one parental populations. When $\mu = 5.10^{-4}$, this means that the genomes of the hybrids will be

composed of a mixture of genes biased towards outcrossing heredity instead of being composed

exclusively of genes from the outcrossing population.

Fig. 2: *Hybrid genome composition depends on* μ, r *and* σ$_h$. *We illustrate here histograms of the*

part of the hybrid genomes that were inherited from the self-fertilizing parental population.

Results are displayed for various recombination rates (first column: r = 10^{-6}, *second column:* 10^{-}

[1]), mutation rates (top four rows: $\mu = 10^{-5}$, bottom four rows: $\mu = 5.10^{-4}$), and rates at which the

hybrid population reproduces by self-fertilization (first and third rows: $\sigma_h = 0$, second and fourth

rows: $\sigma_h = 1$).

Overall, we see that when there are many mutations around, the genomic composition of hybrids

between species with different mating systems mostly depends on the recombination rate

between loci. If recombination is low, outcrossers accumulate more mutations than selfers and

their genome tend to disappear in hybrids. In contrast, if recombination rate is high, selfers'

genomes accumulate more deleterious mutations than outcrossers and their genes tend to

disappear in hybrids. Overall, the hybrid mating system seems to have a marginal impact: the

hybrid genomic composition seems primarily directed by parental populations' mutation load

and genome quality.

Fig. 2 was drawn for complete self-fertilization ($\sigma_o = 1$ and $\sigma_h = 1$). However, in nature, self-

fertilization is often associated with low rates of outcrossing (JARNE AND AULD 2006). To take

that into account, we show on Fig. 3 the effect of assuming self-fertilization rates of only 90%.

We see that most of the time, low rates of outcrossing do not change the average ancestry in

hybrids (around 50% of hybrid genome coming come the self-fertilizing parent when $\mu = 10^{-5}$,

only the self-fertilizing parents' genes transmitted to hybrids when $\mu = 10^{-5}$ and $r = 10^{-6}$).

However, patterns reverse in the case where $\mu = 10^{-5}$ and $r = 10^{-1}$. For these parameter values,

only outcrossing parents' genes are inherited in hybrids when $\sigma_o = \sigma_h = 1$, while a majority of

self-fertilizing parents' genes are transmitted when $\sigma_o = \sigma_h = 0.9$. We saw on Fig. 1 that for these

parameter values, outcrossing species' mean fitness just slightly overcome selfing species' mean

fitness. When the self-fertilization rate of the selfing species is increased, it accumulates less

deleterious mutations, enough for it to gain better fitness and its genes to be preferentially

transmitted in hybrids. Overall, we see again that ancestry patterns will depend on the

accumulated mutation load in the parental species, which will depend on parameters such as μ

and r, as seen before, but also the exact reproductive modes of each of the parental species.

_Fig. 3__: Hybrid genome composition also depends on the exact reproductive modes. Same values_

of r and µ are used as for Fig. 2. We now compare $\sigma_o = \sigma_h = 1$ _(first and third rows), with_ $\sigma_o = \sigma_h$

= 0.9. Again, histograms are shown for 100 iterations, with the x-axis _representing the part of_

 the hybrid genome that, 10,000 generations after hybridization, was inherited from the self-

fertilizing parents.

Another effect of introducing some low rates of outcrossing in self-fertilizing hybrids is that it

introduces recombination. This does not seem to have much of an effect when $r = 10^{-6}$, as

recombination rate is marginal. However, we see when $r = 10^{-1}$ that this recombination allows

for the hybrid genome to exhibit mixed ancestry, instead of being composed of genes from only

one or the other parental species.

3.4. Effects of divergence time on hybrid genome composition

The only case where hybrids' genomes become a mix of genes from the two populations is when

the hybrids outcross and recombination is frequent. In that case, we illustrate on Fig. 3 how the

hybrid genome composition changes depending on the time of divergence prior to hybridization

t_1. As the populations have gone through longer divergence, the selfers´ genome is of worst

quality compared with outcrossers´ genomes (see Fig. 1). Consequently, and as illustrated on

Fig. 3, the part of hybrid genome descending from self-fertilizing population ancestry tend to

decrease with time prior to hybridization. Also, we see that the standard deviation of the self-

fertilizing ancestry decreases: as differences between outcrossers and selfers amplify, results of

hybridization are more stable from one iteration to the next.

Fig. 4: Hybrid genomic mixture depends on how much time passed before hybridization. We illustrate the histograms of the part of the hybrid genomes that were inherited from the self-fertilizing parental population for different number of generations of divergence between the parental populations prior to hybridization. We illustrate results for divergence times $t_1 = 1,000$ (lighter pink), $t_1 = 10,000$ (pink) and $t_1 = 50,000$ (darker red).

4. Discussion

The transition from obligate outcrossing to predominantly selfing has happened multiple times in plants and animals (JARNE AND CHALESWORTH 1993; JARNE AND AULD 2006; THE TREE OF SEX CONSORTIUM 2014). This transition comes with a combination of phenotypic and genomic changes collectively known as the "selfing syndrome" (SHIMIZU AND TSUCHIMATSU 2015; CUTTER 2019). Among those changes, outcrossers and selfers are expected to differ in their mutation load. Theory predicts that higher levels of heterozygosity in outcrossing species allows to better hide mutations that are simultaneously highly deleterious and highly recessive from selection, resulting in these reaching higher frequencies in outcrossers when compared to selfers (WANG *et al.* 1999). In contrast, theory also predicts that higher levels of homozygosity in self-fertilizing species result in less efficient recombination events, and thus stronger selective interference (WRIGHT *et al.* 2013; HARTFIELD *et al.* 2017). As a result, slightly deleterious, co-

dominant alleles are more likely to escape purging out by selection and increase in frequency by

random genetic drift (CHARLESWORTH *et al.* 1993). Our results generally agree with these

predictions. As it turns out, there seems to be pros and cons for both outcrossing and self-

fertilization in terms of mutation load. An important question is, which of the two processes is

the strongest and most influences the genetic content of outcrossing and selfing species?

Our results, in line with other recent theoretical results (SIANTA *et al.* 2022) emphasize the

importance of the recombination rate in the build-up of the mutation load in outcrossers and

selfers. We show that when recombination between adjacent loci is relatively high, outcrossers

suffer little from selective interference. As a result, few deleterious mutations drift to fixation,

fewer than in the self-fertilizing species that do suffer from selective interference. In this case,

outcrossers evolve better genomes than their self-fertilizing counterparts (higher population

mean fitness). Importantly, the prediction changes when we assume a low recombination rate

between adjacent loci. In that case, outcrossers also suffer from selective interference since

genetic associations are rarely re-shuffled. In addition, higher heterozygosity in outcrossers

means higher frequency of deleterious, recessive mutations, as their detrimental effects are

partially hidden in heterozygotes. These higher frequencies means that it is easier for these

deleterious mutations to drift to fixation in outcrossers. As a result, we see that when

recombination rates are low, outcrossers tend to accumulate more fixed deleterious mutations,

which eventually translate into outcrossers evolving worse genomes (lower population mean

fitness).

An interesting consequence of this result is that relative fitnesses may vary within genomes.

Local rates of recombination are known to vary within genomes (NACHMAN AND CHURCHILL

1996; KONG *et al.* 2002; JENSEN-SEAMAN *et al.* 2004). It is possible, depending on how other

factors influencing selection and drift, that this could mean that areas of higher recombination

rate exhibit higher mutation accumulation in a self-fertilizing species than in its outcrossing

counterpart, while the reverse would hold true in areas of lower recombination rate.

Mutation accumulation by selective interference highly depends on genetic drift, and thus on the

effective population size N_e. On one hand, increasing population size limits the fixation of

deleterious mutations. In simulations shown in Supplementary Material Figure S1, with $\mu = 10^{-5}$

and $r = 10^{-6}$, increasing population size from 500 to 5000 decreases the deleterious mutation

fixation rate approximately by a factor 10. On the other hand, there are particular demographic

events that leads to N_e values much lower than the actual population size such as population

bottlenecks or colonization events. Importantly, self-fertilizing species seem to be prone to this

kind of events as they can in principle more easily colonize new environments and found new

populations (THEOLOGIDIS *et al.* 2014; NOEL *et al.* 2016; GROSSENBACHER *et al.* 2017). This

may explain empirical patterns of relaxed purifying selection observed in some self-fertilizing

species (SLOTTE *et al.* 2010; BURGARELLA *et al.* 2015; WANG *et al.* 2021). However, evidence

for purifying selection in selfers is often mixed (GLÉMIN *et al.* 2006; HAUDRY *et al.* 2008;

ESCOBAR *et al.* 2010). This mixed empirical evidence may be caused by the demographic history

of the species as mentioned above, but also due to other factors such as divergence time between

selfers and outcrossers, biased gene conversion, and stability of selfing as a reproductive strategy

(ESCOBAR *et al.* 2010). Overall, it is likely that self-fertilizing species suffer from greater

mutation accumulation than what we predict here due to their demographic and evolutionary

history.

How does mutation load in the parental species influence the ancestry of hybrids' genes?

Previous studies have hypothesized that hybrid individuals' mating system should strongly

impact the evolution of the genetic composition of hybrids and the levels of introgression to be

expected in a hybrid zone. PICKUP *et al.* (2019) predicts that in outcrossing hybrids, the many

slightly deleterious, co-dominant mutations coming from the self-fertilizing ancestry should be

selected against and purged while the highly deleterious, recessive mutations coming from the

outcrossing ancestry should stay hidden in heterozygosity and be maintained. That is, the authors

predict a purge of selfing ancestry in outcrossing hybrids. Similarly, they predict that in selfing

hybrids, the highly deleterious, recessive mutations coming from the outcrossing ancestry should

rapidly become homozygote and be selected against, while the many slightly deleterious, co-

dominant mutations should keep on evading selection and be maintained. This time, the

outcrossing ancestry is expected to be purged in selfing hybrids. Everything happens as if the

mutation load was adapted to a certain reproductive mode and should be maintained in a hybrid

that shares the same reproductive mode. This should result in limited introgression between

outcrossing and self-fertilizing species, in both directions.

Though appealing, this verbal argument does not appear to be verified in our simulations. Our

results emphasize that what matters is merely which parental species has the strongest mutation

load. The genome ancestry with the worst fitness will most likely be purged in hybrids,

independently of the hybrid mating system itself.

KIM *et al.* (2018) produced a similar model of introgression between outcrosser and selfer

parental populations. In their simulations, they obtained that genomes from outcrossing

populations are of better quality because selfer genomes accumulate fixed deleterious mutations

by selective interference, predicting limited introgression of selfer into outcrosser genomes and

greater introgression of outcrosser into selfer genomes. A few genomic parameters differ

between our work and this paper. First, the authors use a gamma distribution for the selective

coefficient of new deleterious mutations whose shape and scale parameters are such that overall

selection is much weaker in their study. This may alter the mutation load balance by increasing

genetic drift. However, it increases drift in both outcrossers and selfers, and simulations run with

$s = 0.01$ (which provides a selective coefficient distribution much closer to the one in KIM *et al.*

(2018)) show an acceleration of deleterious mutation accumulation similar in the two parental

species (see Supplementary Material Figure S2). KIM *et al.* (2018) also consider a genomic

structure based on chromosome 1 of *Arabidopsis thaliana*. This notably means that they consider

a greater number of loci under purifying selection, which is expected to increase the relative

importance of selective interference and mutation accumulation. However, simulations with

1,000 and 10,000 instead of 100 loci (Supplementary Material Figure S3) again emphasize that

mutation accumulation increases in both selfers and outcrossers; in both cases, outcrossers still

accumulate more deleterious mutations in our simulations.

Focusing on the case of *Arabidopsis thaliana* KIM *et al.* (2018) acknowledge that they look at a

scenario with relatively high recombination rates. As such, they may simply fall in a scenario

where selective interference may play a stronger role in selfers (high recombination rates). How

their result is generalizable to other self-fertilizing plant and animals remains to be assessed.

Mutation load is not the only factor that may influence introgression patterns between self-

fertilizing and outcrossing species. Behavioral and physiological constraints are also major

factors that are known to affect hybridization and introgression directions. In a hybrid zone

between species with different mating systems, introgression is expected to happen more

frequently from the selfing species to the outcrosser, than vice-versa (KIM *et al.* 2018; PICKUP *et*

al. 2019). This expectation relies on the premise that selfing (especially prior selfing (TIAN-BI *et*

al. 2008; BRYS *et al.* 2016; BERBEL-FILHO *et al.* 2021)) provides a very limited window of

459 opportunity for outcrossing, by conspecific, heterospecific, or potential male hybrids (PICKUP *et*

460 *al.* 2019; BERBEL-FILHO *et al.* 2021). This pattern of higher levels of introgression from selfers

461 into outcrossers has been commonly found in plant systems (RUHSAM *et al.* 2011; RUHSAM *et al.*

462 2013; BRANDVAIN *et al.* 2014). Our findings extend that expectation, but from a different

463 perspective. Our results indicate that under certain scenarios (here, lower recombination rates)

464 self-fertilizing species may exhibit lower mutation load relative to outcrossing species.

465 Following what we uncovered here, this should be expected to bias hybrid ancestry towards the

466 selfer parents, that is to result in higher levels of introgression from selfers to outcrossers.

467 However, under other scenarios (here, higher recombination rates), our model predicts the

468 contrary: that hybrid ancestry would be biased towards the outcrossing, fitter parents.

469 We argue that our model provides insights into one of the factor, mutation load, that may

470 influence the direction of introgression in hybrids zones between selfers and outcrossers. Of

471 course, many other factors also influence such introgression and should be taken into account

472 when comparing introgression patterns with parental species' mutation load. For instance, the

473 weak inbreeder/strong outbreeder hypothesis (WISO) (BRANDVAIN AND HAIG 2005) states that

474 due to the enhanced opportunity for genomic conflict, outcrossers' gametes are more competitive

475 than selfers' ones. Given equal fertilization opportunities, if hybrids mostly outcross, higher

476 introgression is expected from outcrossers to selfers under the WISO premises. Another factor

477 that can contribute for deviations from the expectations of our model is selfing timing. Selfing

478 can happen before (prior selfing), during (competitive), or after (delayed selfing) the opportunity

479 for outcrossing (LLOYD 1979). The parental selfing time may limit the window of opportunity

480 from an outcrossing hybrid to the selfing parental species, making the major direction of

481 hybridization more common from selfer to outcrosser, than vice-versa (BRYS *et al.* 2016;

BERBEL-FILHO *et al.* 2021). Finally, unilateral genetic incompatibilities may bias the direction of

introgression (TURELLI AND MOYLE 2007) in both directions, regardless of the hybrids mating

system and recombination rates. In our work, we focused solely on the effect of mating systems

via mutation loads and have shown that parental species' loads are good predictors of long-term

genomic composition of hybrid species. Altogether, the examples presented above represent

interesting avenues of research whenever deviations from this prediction are found.

5. Authors Contribution

FF and WMB-F designed the study. FF coded the model and ran the simulations. FF and WB-F

interpreted the results and wrote the manuscript.

6. Acknowledgement

FF and WMB-F are supported by a NSFDEB-NERC Research Grant NE/T009322/1. We thank

Christelle Fraisse, Benjamin C. Haller and Peter Ralph, along with editors and reviewers, for

comments and suggestions that helped us improve the quality of the manuscript.

References

Agrawal, A. F., and M. C. Whitlock, 2011 Inferences About the Distribution of Dominance Drawn From Yeast Gene Knockout Data. Genetics 187**:** 553-566.

Arunkumar, R., R. W. Ness, S. I. Wright and S. C. H. Barrett, 2015 The Evolution of Selfing Is Accompanied by Reduced Efficacy of Selection and Purging of Deleterious Mutations. Genetics 199**:** 817-829.

Berbel-Filho, W. M., A. Tatarenkov, G. Pacheco, H. M. V. Espírito-Santo, M. G. Lira *et al.*, 2021 Against the odds: hybrid zones between mangrove killifish species with different mating systems. Genes 12**:** 1486.

Brandvain, Y., and D. Haig, 2005 Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. The American Naturalist 166**:** 330-338.

Brandvain, Y., A. M. Kenney, L. Flagel, G. Coop and A. L. Sweigart, 2014 Speciation and introgression between *Mimulus nasutus* and *Mimulus guttatus*. PLoS Genetics 10**:** e1004410.

Brys, R., J. Van Cauwenberghe and H. Jacquemyn, 2016 The importance of autonomous selfing in preventing hybridization in three closely related plant species. Journal of Ecology 104**:** 601-610.

Burgarella, C., P. Gayral, M. Ballenghien, A. Bernard, P. David et al., 2015 Molecular evolution of freshwater snails with contrasting mating systems. Molecular Biology and Evolution 32: 2403-2416.

Busch, J. W., S. Bodbyl☐Roels, S. Tusuubira and J. K. Kelly, 2022 Pollinator loss causes rapid adaptive evolution of selfing and dramatically reduces genome☐wide genetic variability. Evolution.

Charlesworth, D., M. T. Morgan and B. Charlesworth, 1993 Mutation accumulation in finite
outbreeding and inbreeding populations. Genetical Research 61**:** 39-56.

Cutter, A. D., 2019 Reproductive transitions in plants and animals: selfing syndrome, sexual
selection and speciation. New Phytologist 224**:** 1080-1094.

Escobar, J. S., A. Cenci, J. Bolognini, A. Haudry, S. Laurent et al., 2010 An integrative test of
the dead-end hypothesis of selfing evolution in triticeae (Poaceae). Evolution: 2855-2872.

Frankham, R., 2005 Genetics and extinction. Biological Conservation 126**:** 131-140.

Gabriel, W., M. Lynch and R. Bürger, 1993 Muller's ratchet and mutational meltdowns.
Evolution 47**:** 1744-1757.

García-Dorado, A. and A. Caballero, 2000 On the average coefficient of dominance of
deleterious spontaneous mutations. Genetics 155:1991-2001.

Glémin, S., E. Bazin and D. Charlesworth, 2006 Impact of mating systems on patterns of
sequence polymorphism in flowering plants. Proceedings of the Royal Society B:
Biological Sciences 273: 3011-3019.

Grossenbacher, D. L., Y. Brandvain, J. R. Auld, M. Burd, P. O. Cheptou et al., 2017
Self☐compatibility is over☐represented on islands. New Phytologist 215: 469-478.

Haller, B. C., and P. W. Messer, 2019 SLiM 3: forward genetic simulations beyond the Wright–
Fisher model. Molecular Biology and Evolution 36**:** 632-637.

Hartfield, M., T. Bataillon and S. Glémin, 2017 The evolutionary interplay between adaptation
and self-fertilization. Trends in Genetics 33**:** 420-431.

Haudry, A., A. Cenci, C. Guilhaumon, E. Paux, S. Poirier et al., 2008 Mating system and
recombination affect molecular evolution in four *Triticeae* species. Genetics Research 90:
97-109.

Hu, X. S., 2015 Mating system as a barrier to gene flow. Evolution 69**:** 1158-1177.

Igic, B., and J. W. Busch, 2013 Is self☐fertilization an evolutionary dead end? New Phytologist

198**:** 386-397.

Jarne, P., and J. R. Auld, 2006 Animals mix it up too: the distribution of self-fertilization among

hermaphroditic animals. Evolution 60**:** 1816-1824.

Jarne, P., and D. Chalesworth, 1993 The evolution of the selfing rate in functionally

hermaphrodite plants and animals. Annual Review of Ecology, Evolution, and

Systematics 24**:** 441-466.

Jensen-Seaman, M. I., T. S. Furey, B. A. Payseur, Y. Lu, K. M. Roskin *et al.*, 2004 Comparative

recombination rates in the rat, mouse, and human genomes. Genome Research 14**:** 528-

538.

Kim, B. Y., C. D. Huber and K. E. Lohmueller, 2018 Deleterious variation shapes the genomic

landscape of introgression. PLos Genetics 14**:** e1007741.

Kong, A., D. F. Gudbjartsson, J. Sainz, G. M. Jonsdottir, S. A. Gudjonsson *et al.*, 2002 A high-

resolution recombination map of the human genome. Nature Genetics 31**:** 241-247.

Lloyd, D. G., 1979 Some reproductive factors affecting the selection of self-fertilization in

plants. The American Naturalist 113**:** 67-79.

Lynch, M., R. Bürger, D. Butcher and W. Gabriel, 1993 The Mutational Meltdown in Asexual

Populations. Journal of Heredity 84**:** 339-344.

Lynch, M., J. Conery and R. Bürger, 1995 Mutational meltdowns in sexual populations.

Evolution 49**:** 1067-1080.

Manna, F., G. Martin and T. Lenormand, 2011 Fitness Landscapes: An Alternative Theory for

the Dominance of Mutation. Genetics 189**:** 923-937.

Martin, N. H., and J. H. Willis, 2007 Divergence associated with mating system causes nearly

complete reproductive isolation between sympatric Mimulus species. Evolution 61**:** 68-

82.

Nachman, M. W., and G. A. Churchill, 1996 Heterogeneity in rates of recombination across the

mouse genome. Genetics 142**:** 537-548.

Noel, E., Y. Chemtob, T. Janicke, V. Sarda, B. Pelissie et al., 2016 Reduced mate availability

leads to evolution of self-fertilization and purging of inbreeding depression in a

hermaphrodite. Evolution 70: 625-640.

Ostevik, K. L., J. L. Rifkin, H. Xia and M. D. Rausher, 2021 Morning glory species

co☐occurrence is associated with asymmetrically decreased and cascading reproductive

isolation. Evolution Letters 5**:** 75-85.

Phadnis, N. and J. D. Fry, 2005 Widespread correlations between dominance and homozygous

effects of mutations: implications for theories of dominance. Genetics, 171: 385-392.

Pickup, M., Y. Brandvain, C. Fraisse, S. Yakimowski, N. H. Barton *et al.*, 2019 Mating system

variation in hybrid zones: facilitation, barriers and asymmetries to gene flow. New

Phytologist 224**:** 1035-1047.

Poon, A., and S. P. Otto, 2000 Compensating for our load of mutations: freezing the meltdown of

small populations. Evolution 54**:** 1467-1479.

Rand, D. M., 2008 Mitigating Mutational Meltdown in Mammalian Mitochondria. PLoS Biology

6**:** e35.

Ruhsam, M., P. Hollingsworth and R. Ennos, 2011 Early evolution in a hybrid swarm between

outcrossing and selfing lineages in Geum. Heredity 107**:** 246-255.

Ruhsam, M., P. M. Hollingsworth and R. A. Ennos, 2013 Patterns of mating, generation of diversity, and fitness of offspring in a Geum hybrid swarm. Evolution 67**:** 2728-2740.

Sianta, S. A., S. Peischl, D. A. Moeller and Y. Brandvain, 2022 The efficacy of selection may increase or decrease with selfing depending upon the recombination environment. Evolution.

Simmons, M. J., and J. F. Crow, 1977 Mutations affecting fitness in *Drosophila* populations. Annual Review of Genetics, 11:49-78.

Shimizu, K. K., and T. Tsuchimatsu, 2015 Evolution of selfing: recurrent patterns in molecular adaptation. Annual Review of Ecology, Evolution, and Systematics 46**:** 593-622.

Slotte, T., J. P. Foxe, K. M. Hazzouri and S. I. Wright, 2010 Genome-wide evidence for efficient positive and purifying selection in *Capsella grandiflora*, a plant species with a large effective population size. Molecular Biology and Evolution 27: 1813-1821.

Takebayashi, N., and P. L. Morrell, 2001 Is self-fertilization an evolutionary dead end? Revisiting an old hypothesis with genetic theories and a macroevolutionary approach. American Journal of Botany 88**:** 1143-1150.

The Tree of Sex Consortium, 2014 Tree of Sex: A database of sexual systems. Scientific Data 1.

Theologidis, I., I. M. Chelo, C. Goy and H. Teotónio, 2014 Reproductive assurance drives transitions to self-fertilization in experimental *Caenorhabditis elegans*. BMC Biology 12: 1-21.

Tian-Bi, Y.-N. T., E. K. N'goran, S.-P. N'guetta, B. Matthys, A. Sangare *et al.*, 2008 Prior selfing and the selfing syndrome in animals: an experimental approach in the freshwater snail Biomphalaria pfeifferi. Genetics research 90**:** 61-72.

Turelli, M., and L. C. Moyle, 2007 Asymmetric postmating isolation: Darwin's corollary to

610   Haldane's rule. Genetics 176**:** 1059-1088.

611   Wang, J., W. G. Hill, D. Charlesworth and B. Charlesworth, 1999 Dynamics of inbreeding

612   depression due to deleterious mutations in small populations: mutation parameters and

613   inbreeding rate. Genetical Research 74**:** 165-178.

614   Wang, X.-J., S. C. H. Barrett, L. Zhong, Z.-K. Wu, D.-Z. Li et al., 2021 The genomic selfing

615   syndrome accompanies the evolutionary breakdown of heterostyly. Molecular Biology

616   and Evolution 38: 168-180.

617   Wright, S. I., S. Kalisz and T. Slotte, 2013 Evolutionary consequences of self-fertilization in

618   plants. Proceedings of the Royal Society B: Biological Sciences 280**:** 20130133.