Deuterated buprenorphine retains pharmacodynamic properties of buprenorphine

Venumadhav Janganati[a,^], Paloma Salazar[b,^], Brian J. Parks[b], Gregory S. Gorman[c], Paul L. Prather[b], Peter A. Crooks[d], Lisa K. Brents[b,*]

a. TCG GreenChem Inc., Richmond, VA, USA

b. Department of Pharmacology and Toxicology, College of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR, USA

c. Pharmaceutical Sciences Research Institute, McWhorter School of Pharmacy, Samford University, Birmingham, AL, USA

d. Department of Pharmaceutical Sciences, College of Pharmacy, University of Arkansas for Medical Sciences, Little Rock, AR, USA

^co-first authors; authors contributed equally

Category: Neuropharmacology

*Corresponding author: Lisa Brents

Mailing address: 4301 West Markham Street, Slot 611, Little Rock, AR, 72205

Email: LBRENTS@uams.edu

ABSTRACT

Buprenorphine (**BUP**) is an opioid that is an effective treatment for opioid use disorder (**OUD**). However, evidence from our group and others suggests that when it is taken during pregnancy, its active metabolite norbuprenorphine (**NorBUP**) contributes to neonatal opioid withdrawal syndrome without significantly contributing to treatment. Therefore, reducing or eliminating metabolism of BUP to NorBUP without altering its therapeutic and side effect profile is a novel strategy that will likely lower total fetal exposure to opioids and thus improve offspring outcomes. Precision deuteration is a chemical modification that alters pharmacokinetics without altering pharmacodynamics. Here we report the first synthesis of deuterated buprenorphine (**BUP-D2**) and initial tests of its pharmacodynamics relative to buprenorphine. Using homogenates made from cells that were transfected with the human mu, delta, or kappa opioid receptors, we measured receptor-binding affinity of BUP-D2 (relative to BUP) using competition receptor binding assays. We also used these homogenates to measure potency and efficacy of BUP-D2 (relative to BUP) to activate G-proteins in the [^{35}S]GTPγS binding assay. Finally, we used the warm-water tail withdrawal assay to compare the antinociceptive effects of BUP-D2 and BUP. Our two-step synthesis method provided a 48% yield and the product was ≥99% deuterated. Like BUP, BUP-D2 had sub-nanomolar affinity for opioid receptors. BUP-D2 also activated opioid receptors and induced antinociception with equal potency and efficacy relative to BUP. These results indicate that BUP-D2 retains key pharmacodynamic properties of BUP and therefore holds promise as an alternative to BUP.

GRAPHICAL ABSTRACT

Deuterated buprenorphine (BUP-D2) retains pharmacodynamic properties of buprenorphine (BUP)

Deuteration is a strategy applied to drugs to alter pharmacokinetics without affecting pharmacodynamics.

We describe the synthesis and initial pharmacodynamic testing of deuterated buprenorphine (BUP-D2).

Buprenorphine | Deuterated buprenorphine

EQUAL OPIOID RECEPTOR AFFINITY

EQUAL OPIOID RECEPTOR-MEDIATED G-PROTEIN ACTIVATION

EQUAL ANTINOCICEPTIVE POTENCY AND EFFICACY

IMPACT: BUP-D2 LIKELY RETAINS THERAPEUTIC EFFECTS OF BUP AND THUS HAS POTENTIAL AS A BUP SUBSTITUTE.

SUPPORTED BY: NIH National Institute on Drug Abuse | NIH National Center for Advancing Translational Sciences | UAMS Translational Research Institute

Janganati, Salazar, et al., 2022

Keywords: buprenorphine, deuteration, opioid receptor pharmacodynamics, drug development, warm-water tail withdrawal, competition receptor binding, G-protein activation

Abbreviations: **BUP,** buprenorphine; **BUP-D2,** deuterated buprenorphine; **CHO,** Chinese Hamster Ovary cells; **DAMGO,** [D-Ala^2, *N*-MePhe^4, Gly-ol]-enkephalin; **DPDPE**, [D-Pen^2,D-Pen^5]enkephalin; **DPMs**, decays per minute; **GDP,** guanosine diphosphate; **GTPγS,** guanosine 5'-O-[gamma-thio]triphosphate; **hDOR,** human delta opioid receptors; **hMOR**, human mu opioid receptors; **hKOR,** human kappa opioid receptors; **NorBUP,** norbuprenorphine; **MPE,** maximum possible effect; **NOWS,** neonatal opioid withdrawal syndrome; **TLC,** thin layer chromatography; **OUD,** opioid use disorder;

1. <u>INTRODUCTION</u>

Chronic use of opioids during pregnancy negatively affects mothers and their children, and has increased more than five-fold in the United States since the beginning of the opioid addiction and overdose crisis[1,2]. Neonatal opioid withdrawal syndrome (**NOWS**) is a major adverse consequence of prenatal opioid exposure. NOWS is defined by multisystem dysregulation, resulting in poor feeding, vomiting, diarrhea, sweating, lacrimation, tremors, seizures, hyperalgesia, poor sleep, extreme irritability, and sensitivity to lights and sounds[3]. This complex syndrome interferes with growth and development of the neonate and often requires care in costly neonatal intensive care units for the first days, and even weeks, of life[4]. In addition to NOWS, prenatal opioid use is associated with other adverse birth outcomes, including maternal death[5], stillbirth[5,6], preterm labor[5]/birth[6-9], premature rupture of membranes[5], intrauterine growth restriction[5,8], placental abruption[5], low birth weight (< 2500 g)[9], being small for gestational age (weight < 2 standard deviations below sex- and gestational age-specific mean)[9], and having congenital malformations[9].

Maternal opioid use disorder (**OUD**) drives chronic opioid use, high stress, and poor self-care during pregnancy, but can be treated with the mu opioid receptor agonists methadone or buprenorphine (**BUP**) as part of medication-assisted treatment. Medication-assisted treatment stabilizes the maternal-fetal dyad by preventing opioid withdrawal and craving, thus diminishing maternal motivation to pursue and use potentially dangerous illicit opioids. Duration of prenatal medication-assisted treatment is associated with decreased risk of overdose[10], preterm birth[10], and low birth weight[10]. However, because methadone and BUP are opioids, risk of NOWS increased with medication-assisted treatment[10]. Prenatal treatment with BUP yields better neonatal outcomes than prenatal treatment with methadone, including shorter hospital stays and less medication needed to treat withdrawal[11]. However, because BUP is a partial agonist of mu opioid receptors (**MOR**), the receptor subtype that mediates most clinically sought effects of opioids, chronic use of BUP often causes dependence and leads to withdrawal upon cessation of exposure. As such, approximately half of children born to women treated with BUP develop moderate-to-severe NOWS that requires weaning with an opioid agonist such as morphine or methadone[11,12]. Although prenatal BUP treatment currently produces the best outcomes for neonates of women with OUD, improved treatments are clearly needed.

Multiple lines of evidence suggest that an active metabolite of BUP called norbuprenorphine (**NorBUP**) contributes to the adverse effects of BUP on the fetus. Following chronic prenatal BUP use, concentrations of

NorBUP in the human placenta, umbilical cord plasma, and meconium are at least 10-fold higher than BUP concentrations[13-15], suggesting that fetal exposure to NorBUP exceeds BUP. NOWS severity is independent of maternal BUP dose[13,16,17], suggesting that factors other than dosing (e.g., individual variability in BUP metabolism) determine fetal exposure to BUP and its active metabolites. For example, it is plausible that NOWS is more severe for neonates exposed *in utero* to higher levels of NorBUP due to dominance of the NorBUP metabolic pathway over alternative pathways, such as glucuronide conjugation. This possibility is supported by the results of a study in which investigators measured umbilical cord blood concentrations of BUP and its metabolites (which represents fetal blood concentrations at delivery) and determined that NorBUP, but not BUP, was positively correlated with NOWS severity[13]. Furthermore, the glucuronide conjugate of BUP was negatively correlated with NOWS severity[13], suggesting that bias for glucuronide conjugation (over *N*-dealkylation to form NorBUP) is protective, and bias for NorBUP formation (over glucuronide conjugation) is more harmful. Additionally, we previously showed with a rat model of NOWS that prenatal exposure to NorBUP induces dependence and neonatal withdrawal at levels comparable to morphine[18]. NorBUP likely contributes little to the therapeutic centrally mediated effects of BUP treatment due to its restriction to the periphery by p-glycoprotein in the blood-brain barrier[19,20]. Altogether, this evidence suggests that decreasing fetal exposure to NorBUP by reducing NorBUP formation can potentially improve short-term neonatal outcomes, such as NOWS, and the long-term effects of opioid exposure on neurodevelopment.

In this study, we applied precision deuteration to BUP and determined if deuteration affected its pharmacodynamic properties, including opioid receptor affinity, potency, and efficacy. Precision deuteration involves replacing hydrogen atoms with their heavy isotope deuterium, which exploits the primary kinetic isotope effect to slow bond-breaking by metabolic enzymes[21] (**Fig. 1**). The expected net result of this small change is altered

Buprenorphine   Norbuprenorphine   Deuterated buprenorphine

Figure 1. Norbuprenorphine formation. In humans, norbuprenorphine is the primary major metabolite of buprenorphine, as represented by the bold arrow. Norbuprenorphine is formed by oxidative *N*-dealkylation of buprenorphine's cyclopropyl methyl group by various cytochrome P450s. Precision deuteration is expected to make this group less vulnerable to metabolic cleavage and thus reduce its metabolism to norbuprenorphine, as represented by the thin broken arrow.

pharmacokinetics (*e.g.,* prevention of metabolite formation) with little or no change in pharmacodynamics[22,23]. In the present work, we synthesized a deuterated buprenorphine (**BUP-D2**) by replacing two methylene hydrogens with deuterium atoms adjacent to the tertiary nitrogen in the buprenorphine structure (**Fig. 2**) to inhibit oxidative dealkylation of the *N*-cyclopropylmethyl

Figure 2. Synthesis of BUP-D2. See Materials and Methods for description of synthesis.

moiety by liver cytochrome P450's. We hypothesized that BUP and BUP-D2 have equal affinity, potency, and efficacy for opioid receptors, and have equal potency and efficacy of antinociception. Here, we report the synthesis and initial testing of BUP-D2, compared to BUP, for opioid receptor affinity and G-protein activation using a radioligand competition receptor binding and [^{35}S]GTPγS binding assays, respectively. We then tested BUP-D2 and BUP in the warm water tail withdrawal assay, a rat model of antinociception, to determine whether precision deuteration alters *in vivo* buprenorphine effects. Because of virtually indistinguishable pharmacodynamic properties relative to BUP, the results of the present study suggest that BUP-D2 shows promise as an alternative to BUP for treating OUD.

2. **MATERIALS AND METHODS**

2.1 **Drugs**

The National Institute on Drug Abuse Drug Supply Program (Rockville, MD) provided norbuprenorphine (**NorBUP**) in free base form. Buprenorphine (**BUP**) was purchased from Tocris (Minneapolis, MN). Morphine sulfate salt pentahydrate was purchased from Sigma-Aldrich (St. Louis, MO). With the exception of NorBUP, solvents and chemicals used to synthesize deuterated buprenorphine were purchased from Sigma-Aldrich (St. Louis, MO). [^{3}H]diprenorphine (25.1 Ci/mmol) and [^{35}S]GTPγS (1250 Ci/mmol) were purchased from Perkin-Elmer (Waltham, MA). Guanosine-5-diphosphate (GDP) and unlabeled GTPγS were purchased from Fisher

Scientific (Waltham, MA) and Sigma-Aldrich (St. Louis, MO), respectively. [D-Ala^2, NMe-Phe^4, Gly-ol^5]-enkephalin (**DAMGO**) was purchased from Cayman Chemical (Ann Arbor, MI). [D-Pen^{2,5}]Enkephalin, [D-Pen^2,D-Pen^5]Enkephalin (**DPDPE**) and U-50,488 were purchased from Tocris (Minneapolis, MN). Naltrexone hydrochloride was purchased from Sigma Aldrich (St. Louis, MO). Drugs for *in vitro* assays were dissolved at concentrations of 10^{-2} M in 100% DMSO and stored at -20°C until used in experiments. Drugs for *in vivo* experiments were dissolved in a sterile saline solution (0.9% NaCl) containing 5% DMSO.

2.2 BUP-D2 synthesis

BUP-D2 was synthesized in a two-step process as illustrated in **Figure 2** and described here:

Step 1. *Cyclopropyl((4R,4aS,6R,7R,7aR,12bS)-9-hydroxy-6-(2-hydroxy-3,3-dimethylbutan-2-yl)-7-methoxy-1,2,5,6,7,7a-hexahydro-4a,7-ethano-4,12-methanobenzofuro[3,2-e]isoquinolin-3(4H)-yl)methanone:*

Triethylamine (44.96 mg, 0.444 mmol) and cyclopropanecarbonyl chloride (23.22 mg, 0.222 mmol) were added to norbuprenorphine hydrochloride (100 mg, 0.222 mmol) dissolved in dichloromethane (3 mL) at 0 °C. The reaction mixture was stirred for 1 h at 0 °C. After completion of the reaction, which was monitored by thin layer chromatography (**TLC**), water (5 mL) was added and the reaction mixture was extracted with dichloromethane (2x5 mL). The organic layer was washed with water (3x5 mL), dried over anhydrous Na$_2$SO$_4$ and concentrated under reduced pressure to afford crude product. The crude product was purified by column chromatography (silica gel, 30-40 % EtOAc in hexane) and obtained mixture of diastereomers (yield: 75 %).

Step 2. *(4R,4aS,6R,7R,7aR,12bS)-3-(cyclopropylmethyl-d2)-6-(2-hydroxy-3,3-dimethylbutan-2-yl)-7-methoxy-1,2,3,4,5,6,7,7a-octahydro-4a,7-ethano-4,12-methanobenzofuro[3,2-e]isoquinolin-9-ol:*

To the cyclopropylcarbonyl norbuprenorphine (diasteriomeric mixture) (80 mg, 0.17 mmol) in tetrahydrofuran (3 mL), lithium aluminum deuteride (27.92 mg, 0.68 mmol) was added at 0 °C. The reaction mixture was stirred at 50 °C for 3 h. After completion of the reaction, which was monitored by TLC, the reaction mixture was brought to 0 °C and quenched with 0.03 mL of water+0.03 mL of 15% aqueous sodium hydroxide+0.09 mL of water. The reaction mixture was stirred for 30 min at ambient temperature and filtered. The filtrate was concentrated under reduced pressure to afford crude product. The crude product was further purified by column chromatography (silica gel, 30 % EtOAc in hexane) to afford the pure product as a white solid. The solid was dissolved in diethyl ether and cooled to 0 °C. 2M HCl solution in diethyl ether (~ 0.1 mL) was added and stirred for 1 h. The reaction

mixture was filtered and washed with diethyl ether to afford white solid (yield: 64 %). The products were purified by flash column chromatography (silica gel; EtOAc/Hexane). ^{1}H and ^{13}C NMR spectra were recorded on a Varian 400 MHz spectrometer equipped with a Linux workstation running on vNMRj software. All the spectra were phased, baseline was corrected where necessary, and solvent signals (CDCl$_3$) were used as reference for both ^{1}H and ^{13}C spectra. TLC separations were carried out on pre-coated silica gel plates (F 254 Merck).

^{1}H NMR (300 MHz, CDCl$_3$; **Supporting Data Figure 1A**): δ 6.81 (d, J = 7.6 Hz, 1H), 6.50 (d, J = 7.6 Hz, 1H), 5.81 (s, 1H), 4.51 (s, 1H), 4.44 (s, 1H), 3.51 (s, 3H), 2.98-2.83 (m, 3H), 2.62-2.55 (m, 1H), 2.32-2.12 (m, 3H), 2.01-1.92 (m, 1H), 1.85-1.64 (m, 3H), 1.54 (s, 1H), 1.34 (s, 3H), 1.31-1.25 (m, 1H), 1.1-1.0 (m, 9H), 0.81-0.62 (m, 2H), 0.51-0.41 (m, 2H), 0.12-0.06 (m, 2H) *ppm*. ^{13}C NMR (CDCl$_3$, 100 MHz; **Supporting Data Figure 1B**): δ 145.4, 137.3, 132.5, 128.2, 119.5, 116.4, 96.9, 80.8, 79.6, 58.2, 52.5, 46.4, 43.6, 40.3, 35.9, 35.6, 33.4, 29.6, 26.3, 22.9, 20.1, 18.1, 9.2, 4.0, 3.1 *ppm*. MS (ESI), m/z: 470 (M+H)$^+$.

Enrichment of the product with deuterium was determined by LC/MS/MS to be >99% g/atom (**Supporting Data Figure 2**). The LC/MS/MS system used for this analysis consisted of Shimadzu system (Columbia MD) equipped with LC20-AD dual HLPC pumps, an SIL20-AC HT autosampler, and a DGU-20A2 in-line degasser. Detection was performed using an Applied BioSystems 4000 QTRAP (Applied Biosystems, Foster City, CA) triple quadrupole mass spectrometer operated in the positive ion mode. Mass calibration, data acquisition and quantitation were performed using Applied Biosystem Analyst 1.6.21 software (Applied Biosystems, Foster City, CA).

2.3 Cell Culture

Chinese hamster ovary (**CHO**) cells stably transfected with human mu opioid receptors (**CHO-hMOR**) were a gift from Dr. Dana Selley (Virginia Commonwealth University, Richmond, VA). CHO cells were stably transfected with human delta opioid receptors (**CHO-hDOR**) in our laboratory as described previously[24]. CHO cells stably transfected with human kappa opioid receptors (**CHO-hKOR**) were a gift from Dr. Lee-Yuan Liu-Chen (Temple University, Philadelphia, PA). CHO-hMOR and CHO-hDOR cells were cultured in DMEM Nutrient Mix F12 1:1 media containing 10% FetalPlex®, 1% penicillin-streptomycin, and 0.2 mg/mL hygromycin (Gibco, Waltham, MA). CHO-hKOR cells were cultured in DMEM Nutrient Mix F12 1:1 (Gibco, Waltham, MA) media containing 10% FetalPlex® (Gemini Bioproducts, Sacramento, CA), 1% penicillin-streptomycin (Corning,

Corning, NY), and 0.27 mg/mL G418 Sulfate (Fisher Scientific, Houston, TX). Cells were grown in a humidified chamber with 5% CO_2 at 37°C in sterile T175 flasks and were harvested at 100% confluence with 0.04% EDTA in phosphate buffered saline (pH 7.5). Harvested cells were centrifuged at 1040 x g to form cell pellets and supernatant was removed. Cell pellets were stored at -80°C until used to make membrane homogenates.

2.4 **Membrane Homogenate Preparation**

Up to five cell pellets, each made from up to four 100% confluent T175s, were thawed on ice, then transferred to a 40 mL Dounce glass homogenizer and suspended in 10 mL (or ~2 mL/cell pellet) of ice-cold homogenization buffer (50 mM HEPES, pH 7.4, 3 mM $MgCl_2$, and 1 mM EGTA). Ten strokes were applied to the contents of the homogenizer with a coarse-grade pestle, and the contents were then centrifuged at 40,000 x g for 10 minutes at +4°C. Supernatants were discarded, and the pellet was again transferred to the homogenizer, suspended in 10 mL of homogenization buffer, coarsely homogenized with 10 strokes, and centrifuged at 40,000 x g for 10 minutes at +4°C twice more, with the supernatant discarded each time. The resulting pellet was homogenized using 10 strokes of a fine-grade pestle in 10 mL (or ~2 mL/cell pellet) of ice-cold 50 mM HEPES. Homogenates were aliquoted in 0.5 or 1 mL aliquots and stored at -80°C until used in experiments. Protein concentrations of homogenates were determined using the BCA Protein Assay (Thermo Scientific, Walham, MA).

2.5 **Competition Receptor Binding**

Competition receptor binding assays were performed using 25, 50, or 100 µg of membrane homogenates made from CHO-hMOR, CHO-hKOR, or CHO-hDOR cells, respectively, per sample. Membranes were incubated with 1 nM of the non-selective opioid antagonist [³H]diprenorphine plus vehicle or naltrexone (10 µM, non-specific binding) or various concentrations of either non-radioactive morphine, buprenorphine, or deuterated buprenorphine in a buffer containing 5 mM $MgCl_2$, 50 mM Tris-HCl, and 0.05% bovine serum albumin. Total reaction volume was 1 mL. Samples were incubated for 90 minutes at room temperature to achieve equilibrium before reactions were rapidly terminated by vacuum filtration onto Whatman GF/B glass microfiber filters (Brandel, Gaithersburg, MD) with a Brandel cell harvester (Brandel, Gaithersburg, MD). Filters were washed three times with 5 mL of ice-cold assay buffer and samples were transferred to 7-mL scintillation vials. Four milliliters of ScintiVerse™ BD Cocktail scintillation fluid (Fisher Scientific, Waltham, MA) were added to scintillation vials, and allowed to incubate overnight. Radioactivity retained on filters was then quantified by liquid

scintillation spectrophotometry using the Perkin-Elmer Tri-Carb 2910TR Liquid Scintillation Analyzer (Walham, MA). Non-specific binding was determined in each experiment using a receptor-saturating concentration of non-radioactive naltrexone, a non-selective opioid antagonist. Specific binding for each sample was determined by subtracting radioactive counts (decays per minute, **DPMs**) measured in these non-specific binding samples from the total radioactive counts obtained in samples. Samples were conducted in duplicate within each experiment, and at least three independent experiments were combined to produce each receptor binding curve.

2.6 [^{35}S]GTPγS binding assay

Activation of MOR, DOR, and KOR by BUP and BUP-D2 was quantified using an [^{35}S]GTPγS, a radiolabeled non-hydrolyzable analogue of GTP that binds and labels activated G-proteins. In this assay, 50 µg of membrane homogenates were incubated with 0.1 nM [^{35}S]GTPγS in the presence of vehicle, unlabeled GTPγS (10 µM, non-specific binding) or opioids (BUP, BUP-D2, or controls) for 30 minutes at 30°C. The receptor-selective full agonists DAMGO, U50, 488, and DPDPE were used as positive controls to measure activation of hMOR, hKOR, and hDOR, respectively. Naltrexone, a non-selective opioid antagonist, was included as a negative control. Assay buffer contained 20 mM HEPES, 100 mM NaCl, 10 mM MgCl2, 10 uM GDP, and 0.05% bovine serum albumin. Membrane homogenates and GDP were pre-incubated together for 5 minutes at room temperature before being combined with the other components. Reactions were rapidly terminated using vacuum filtration onto Whatman GF/B glass microfiber filters (Brandel, Gaithersburg, MD) using a Brandel cell harvester, followed by three 5 mL washes of ice-cold 50 mM Tris-HCl (pH 7.4) buffer containing 0.05% bovine serum albumin. Filters were transferred to 7 mL scintillation vial and 4 mL of ScintiVerse™ BD Cocktail scintillation fluid was added to each vial. Following overnight incubation, bound radioactivity in samples was determined by liquid scintillation spectrophotometry using the Perkin-Elmer Tri-Carb 4910 TR Liquid Scintillation Analyzer (Walham, MA). Non-specific binding was determined in each experiment using a receptor-saturating concentration of non-radioactive GTPγS. Specific binding for each sample was determined by subtracting radioactive counts (DPMs) measured in these non-specific binding samples from the radioactive counts in the sample. Samples were conducted in triplicate within each experiment, and at least three independent experiments were combined to produce for each G-protein activation curve.

2.7 Animal Care and Use

Studies with rodents were approved by the University of Arkansas for Medical Sciences Institutional Animal Care and Use Committee (IACUC) prior to commencement of experiments. Female Long Evans rats with indwelling jugular catheters were purchased from Charles River Laboratories (Wilmington, MA). Rats weighed 200-250 grams upon arrival and throughout the study. All rats were singly housed in Plexiglass cages with enrichment. Rats were kept on a 14-hour light/10-hour dark cycles with free access to food in a temperature and humidity-controlled room.

2.8 Warm-water tail withdrawal assay

The analgesic activity of BUP and BUP-D2 were compared using the warm-water tail withdrawal procedure in rats. In this procedure, the latency of a rat to remove its tail from warm water (50°C) up to a cutoff of 20 seconds was measured as an endpoint of nociception. Control trials using a non-noxious water temperature (40°C) were conducted between test trials to confirm that tail withdrawals during test trials were specifically a nociceptive response (as opposed to a learned response). Prior to testing, animals were acclimated to the procedure with control trials until they had at least two successful control trials (i.e., trials in which animals did not remove tail from non-noxious water before cutoff). Then a test trial was conducted using 50°C water to determine baseline latency to withdraw the tail. Ten minutes later rats were given an intravenous injection via jugular catheter of either BUP or BUP-D2 (0.01, 0.03, 0.1, or 0.3 mg/kg, 2 ml/kg). Additional test trials were conducted 10 and 60 minutes after dosing. Control trials were conducted 5 minutes before dosing and 5 and 30 minutes after dosing. Investigators were blinded to treatments during the

2.9 Statistical analysis

Analyses were performed using GraphPad Prism (Version 8.0; San Diego, CA). After baseline and/or non-specific values were subtracted and normalization corrections were applied, data were fit using "three-parameter agonist (or inhibitor) vs. response" non-linear regression (for competition receptor binding and $[^{35}S]GTP\gamma S$ binding assays) or linear regression (for the warm-water tail withdrawal assay). In the competition receptor binding experiments, the log-transformed IC_{50} value of each replicate was obtained from the non-linear regression analyses and converted to a log-transformed K_i value (a normally distributed quantitative measure of receptor affinity) by applying the Cheng-Prusoff equation[25]. Statistically significant differences in affinity for each receptor subtype among BUP, BUP-D2, and morphine were determined by one-way ANOVA and Tukey's

Multiple Comparison *post-hoc* test (p < .05). In the concentration-effect [^{35}S]GTPγS binding experiments, log-transformed EC_{50} values and log-transformed 95% confidence intervals were derived from non-linear regression analyses applied to the aggregate data. Lack of overlap of log-transformed confidence intervals, within receptor subtype, indicated statistically significant group differences in receptor potency. "Top" values (and their 95% confidence intervals) calculated by GraphPad Prism from the agonist vs. response non-linear regression analyses are reported here as the maximum effect (E_{max}) values. For single-concentration [^{35}S]GTPγS binding experiments, data are reported as percent vehicle control, and a one-way ANOVA with Tukey's multiple comparisons test was conducted within receptor subtype. The ED_{50} of antinociception in the warm-water tail withdrawal assay was determined by first converting tail withdrawal latencies to percent maximum possible effect (**% MPE**) using the following formula: % MPE = (X – baseline) / cutoff – baseline) * 100, wherein "X" equals tail withdrawal latency following treatment (in seconds), "baseline" equals tail withdrawal latency prior to treatment (in seconds) and "cutoff" equals the investigator-imposed maximum trial time, which was 20 seconds in this experiment. Then, we applied simple linear regression to the ascending limb of the inverted-U-shaped dose-response curve (i.e., on data points up to, and including, the 0.1 mg/kg dose).

Figure 3. BUP-D2 and BUP have equal, high affinity for opioid receptors. Data points and error bars represent mean and standard error of mean, respectively, of ^{3}H-diprenorphine specific binding in homogenates containing hMOR (A), hDOR (B), or hKOR (C) as a percent of controls (y-axis) in the presence of varying concentrations (x-axis) of unlabeled morphine (circles), BUP (filled triangles), or BUP-D$_2$ (unfilled triangles). "Control" refers to ^{3}H-Diprenorphine binding in the presence of minimal unlabeled drug (10^{-14} M). n = 3-4 independent experiments performed in triplicate.

Finally, we interpolated the dose at which %MPE = 50 (i.e., the half-maximal effect) from the linear regression curve. We applied an F-test to the ascending limb data at each time point to test for differences in slope or intercept of the linear regressions. To test for differences between BUP and BUP-D2 at each dose and overall, we conducted a two-way ANOVA with Sidak's multiple comparisons test with data from all doses (i.e., ascending and descending limb) for each time point.

3. <u>RESULTS</u>

3.1 <u>Competition binding</u>

BUP-D2 binding to opioid receptors like BUP is crucial for BUP-D2 to retain the therapeutic effects of BUP. We used a competition receptor binding assay to determine and compare the affinities of BUP-D2 and BUP for human MOR, human DOR, and human KOR, and used morphine as a positive control.

Table 1. Affinity of BUP-D2 for opioid receptors						
	Log K_i values (M) Mean (95% CI)			K_i values (nM) Mean (SEM)		
	hMOR	**hDOR**	**hKOR**	**hMOR**	**hDOR**	**hKOR**
BUP-D2	-10.30 (-11.09, -9.51)	-9.54 (-10.82, -8.26)	-10.21 (-11.75, -8.66)	0.0771 (0.0383)	0.424 (0.229)	0.101 (0.494)
BUP	-10.00 (-10.23, -9.78)	-9.96 (-10.47, -9.45)	-9.55 (-11.08, -8.01)	0.103 (0.0156)	0.119 (0.0317)	0.511 (0.342)
Morphine	**-8.66 (-9.07, -8.25)**	**-6.59 (-7.11, -6.07)**	**-6.71 (-7.60, -5.83)**	2.50 (0.704)	279 (83.5)	239 (101)

K_i, inhibitor constant; **CI**, confidence interval; **SEM**, standard error of mean; **Bolding** indicates significant difference from BUP and BUP-D2 (within receptor subtype), as determined by Tukey's multiple comparisons test ($p < .005$). Statistical analyses were conducted with log-transformed values only. n = 3-4 independent experiments conducted in duplicate.

In these experiments, morphine and BUP exhibited affinity for each opioid receptor type similar to previously reported affinities[26,27] (**Fig. 3**; **Table 1**). BUP-D2 and BUP exhibited sub-nanomolar affinity at each opioid receptor type (**Table 1**). BUP-D2 and BUP had significantly greater affinity for hMOR than did morphine, but did not differ from each other (**Fig. 3A; Table 1**). Likewise, BUP-D2 and BUP affinities for hDOR did not differ from each other significantly, and both exhibited significantly greater affinity for hDOR than did morphine (**Fig. 3B; Table 1**). Finally, BUP-D2 and BUP affinities for hKOR did not differ from each other, and both had significantly greater affinity for hKOR than did morphine (**Fig. 3C; Table 1**).

3.2 <u>GTPγS Binding</u>

After determining that BUP-D2 retains the binding affinity of BUP for opioid receptors, we wanted to determine whether BUP-D2 also retained the potency and intrinsic efficacy of BUP at opioid receptors. To accomplish this goal, we first measured [^{35}S]GTPγS binding in homogenates containing each opioid receptor type in the presence or absence of a receptor-saturating concentration (1 µM) of BUP, BUP-D2, a positive control, or a negative control. The full agonists DAMGO, DPDPE, and U50, 488 were used as positive controls for G-protein activation by hMOR, hDOR, and hKOR, respectively, and naltrexone (a non-selective opioid antagonist) was used as a negative control for all receptor types. Positive controls significantly activated their respective receptor by at least 175% of vehicle control levels (**Fig. 4**). As expected from previous reports[26,27], BUP stimulated G-proteins in hMOR homogenates greater than baseline activity measured in vehicle controls, but less than positive control-stimulated activity (**Fig. 4A**), indicating that it acts as a partial agonist at hMOR. BUP-D2 likewise stimulated G-protein activation with partial agonist activity at hMOR (**Fig. 4A**). BUP and BUP-D2 did not stimulate G-proteins in hDOR homogenates and equally stimulated G-proteins in hKOR homogenates, suggesting that BUP and BUP-D2 act as neutral antagonists of hDOR and partial agonists of hKOR (**Fig. 4B and 4C**). The negative control naltrexone did not stimulate G-proteins in hMOR or

Figure 4. BUP-D2 and BUP activate opioid receptors with equal maximal efficacy. [^{35}S]GTPγS specific binding (y-axis) represents G-protein activation in homogenates containing hMOR (A), hDOR (B), or hKOR (C) in the presence of vehicle (0.01% DMSO) or a receptor-saturating concentration (1 µM) of positive control full agonists, negative control antagonist naltrexone, BUP, or BUP-D2 (x-axis). Bars represent group means, closed circles represent values obtained from independent experiments, and error bars represent 95% confidence intervals. Within each panel, columns sharing letters are not significantly different (One-way ANOVA, Tukey's multiple comparison test, $p < 0.05$, $n = 3\text{-}5$ independent experiments performed in triplicate).

hDOR homogenates; however, it curiously stimulated G-proteins in hKOR homogenates with partial agonist activity.

Figure 5. BUP-D2 and BUP activate hMOR and hKOR with equal, high potency. [^{35}S]GTPγS specific binding (y-axis) represents G-protein activation in homogenates containing hMOR (A) or hKOR (B) in the presence of increasing concentrations (x-axis) of DAMGO (open circles, A), U50,488 (open circles, B), BUP (closed triangles), or BUP-D2 (open triangles). Symbols and error bars represent means and standard error of mean from independent experiments (n = 3-8 independent experiments per concentrations, conducted in triplicate).

To determine potency to activate opioid receptors, we next determined the concentration-effect of BUP and BUP-D$_2$ to stimulate G-proteins in hMOR and hKOR homogenates (**Fig. 5**). Due to the lack of G-protein activation in screening experiments, concentration-effect curves of BUP and BUP-D2 were not generated using hDOR homogenates. BUP and BUP-D2 exhibited equal potency to activate G-proteins in hMOR homogenates and were both significantly more potent than the full agonist positive control DAMGO (**Fig. 5A; Table 2**). Likewise, in hKOR homogenates, BUP and BUP-D2 exhibited equal potency to activate G-proteins and were both significantly more potent than the full agonist positive control U50, 488 (**Fig. 5B; Table 2**). Greater potency of BUP to activate hMOR and hKOR relative to DAMGO and U50, 488, respectively, was expected based on previous reports[26,27]. Maximal efficacies for BUP and BUP-D2 in the concentration-effect experiment were consistent with partial agonist efficacies determined in the screening experiment (**Fig. 4** vs. **Fig. 5**; **Table 2**), providing more evidence that BUP and BUP-D2 are partial agonists of hMOR and hKOR receptors.

Table 2. Potency and efficacy of BUP-D2 for opioid receptors								
	Log EC$_{50}$ (M) Mean (95% CI)			EC$_{50}$ (nM) Mean (SEM)			Emax (% of vehicle) Mean (SEM)		
	hMOR	hDOR	hKOR	hMOR	hDOR	hKOR	hMOR	hDOR	hKOR
BUP-D2	-10.71 (-11.30, -9.74)	NA	-11.18 (-11.67, -10.72)	0.0197 (0.0115)	NA	0.00655 (0.00368)	191.5 (178.1, 208.6)	^106.94 (92.00, 121.90)	155.0 (147.5, 162.6)
BUP	-10.85 (-11.38, -10.14)	NA	-11.14 (-11.71, -10.60)	0.0140 (0.00770)	NA	0.00719 (0.00470)	194.5 (183.3, 206.4)	^114.32 (100.3, 128.33)	154.1 (146.4, 161.9)
DAMGO	**-7.38 (-7.75, -7.03)**	--	--	42.1 (14.8)	--	--	**307.8 (280.5, 339.4)**	--	--
DPDPE	--	--	--	--	--	--	--	**^172.00 (157.75, 186.19)**	--
U50-488	--	--	**-8.21 (-8.57, -7.89)**	--	--	6.16 (2.01)	--	--	**293.1 (275.2, 311.8)**

EC50, half-maximal effective concentration; **CI,** confidence interval; **SEM**, standard error of mean; **NA**, not applicable (neutral antagonist); **--**, not determined; **^**, determined using a single, receptor-saturating concentration (1 μM). **Bolding** indicates significant differences from BUP and BUP-D2, as determined by lack of overlap of 95% confidence intervals. No statistical testing was conducted with EC$_{50}$ values. n = 3-8 independent experiments per concentration, conducted in triplicate.

3.3 Warm-water tail withdrawal analgesia assay

The antinociceptive effects of BUP and BUP-D2 were compared using the warm water tail withdrawal assay in female Long-Evans rats. In this experiment, we measured the latencies of rats to remove their tails from 50°C water 10 minutes and 60 minutes after an intravenous injection of either BUP or BUP-D2, up to a maximum cutoff of 20 seconds. Baseline latencies were equal for the BUP and BUP-D2 groups (**Fig. 6**). At both time points, there were no differences in tail withdrawal latency between BUP and BUP-D2 at any dose (**Fig. 6**). Likewise, there was no difference between BUP-D2 and BUP in ED$_{50}$ values or percent maximum possible effect (**Table 3**). Maximal antinociception was achieved at 0.1 mg/kg at both time points for both drugs and declined at the higher dose of 0.3 mg/kg. This inverted U-shaped curve is atypical for most opioid agonists but is characteristic of BUP[28], thus providing more evidence that BUP-D2 retains the opioid effects of BUP.

4. DISCUSSION

The present study demonstrates that BUP-D2 retains BUP's high affinity for opioid receptors, as well as BUP's partial agonist efficacy (for hMOR and hKOR) or neutral antagonist efficacy (for hDOR), and high potency

(for hMOR and hKOR). This is important evidence that BUP-D2 has the potential to substitute for BUP for the treatment of opioid use disorder. Ongoing testing in our laboratory will determine whether BUP-D2 is an improvement over BUP by resisting metabolism to NorBUP, an active metabolite that has been implicated in NOWS when BUP is taken during pregnancy. The present study is the first to apply the innovative strategy of precision deuteration to BUP and investigate this compound as a potentially novel therapeutic agent.

The unique therapeutic and safety profile of BUP, relative to methadone and other opioids, is thought to emerge from its high affinity for MOR, DOR, and KOR[26], in combination with its high potency and partial agonist activity for MOR[26]. Determining the affinity of BUP-D2, relative to BUP for opioid receptors is the first step to determining whether BUP-D2 has potential as a therapeutic agent. Using a radioligand competition binding assay, we determined that BUP-D2 and BUP similarly displaced [3H]diprenorphine from each opioid receptor subtype, yielding equivalent sub-nanomolar affinities. Interestingly, [3H]diprenorphine binding in the presence of extremely low (femtomolar) concentrations of BUP and BUP-D2 was surprisingly lower than [3H]diprenorphine binding in the presence of vehicle controls, suggesting that BUP and BUP-D2 apparently displaces [3H]diprenorphine at these concentrations. Additionally, unlike competition binding curves for the positive controls and morphine, the binding curves for BUP and BUP-D2 did not have a Hill slope that approximated -1. BUP and

Figure 6. BUP-D2 fully retains the antinociceptive potency and efficacy of BUP in the warm water tail withdrawal assay. Data points and error bars represent the mean and standard error, respectively, of the latency for rats to remove their tails from 50°C water (y-axis) before (baseline, "BL") and after (10 minutes, A, and 60 minutes, B,) intravenous injection with BUP (closed triangles) or BUP-D2 (open triangles) at varying doses (x-axis). p > 0.05. two-way ANOVA with Sidak's multiple comparisons test, n = 3-4 independent animals per dose.

BUP-D2 instead displaced [³H]diprenorphine in a gradual linear manner throughout the entire range of concentrations used, with no clear minimum threshold for displacement. This behavior is possibly explained by differences in the dissociation rates of

Table 3. Potency and efficacy of BUP and BUP-D2 antinociception, mean (lower, upper 95% confidence interval)

	BUP	BUP-D2
10 minutes		
ED_{50} (mg/kg)	0.052 (0.0464, 0.0584)	0.056 (0.050, 0.065)
% MPE (at 0.1 mg/kg)	90.1 (68.8, 111.4)	84.4 (43.5, 125.2)
60 minutes		
ED_{50} (mg/kg)	0.049 (0.040, 0.061)	0.049 (0.045, 0.054)
% MPE (at 0.1 mg/kg)	96.8 (83.2, 110.5)	97.9 (89.1, 106.8)

ED50, half-maximal effective dose; **% MPE**, percent maximum possible effect; slopes and intercepts were equal, as tested by F-test (P > 0.45), and there is overlap of 95% confidence intervals between BUP and BUP-D2 for ED_{50} and % MPE values at both time points. n = 3-4 independent animals per dose

[³H]diprenorphine and BUP. Others have previously reported that after binding to opioid receptors, BUP dissociates from the receptor in ~40 minutes[29,30]. By comparison, diprenorphine and most other opioids dissociate from opioid receptors in 5-10 minutes[29-31]. The prolonged binding of BUP likely allows it to compete with [³H]diprenorphine at much lower concentrations than would be expected. Kinetic binding experiments specifically designed to measure dissociation rates are needed to definitively confirm that BUP-D2 retains the slow dissociation rate of BUP. In any case, BUP-D2 exhibiting atypical receptor binding similar to that of BUP at low concentrations is further evidence that precision deuteration does not alter the opioid receptor binding properties of BUP.

After confirming that receptor affinity for BUP-D2 was unchanged, relative to BUP, we next determined the potency and efficacy of BUP-D2 relative to BUP. It is important for BUP-D2 to retain the partial agonist efficacy of BUP, such that it stimulates MORs enough to prevent withdrawal (following proper induction) without the high risk of diversion, abuse, respiratory depression, and death observed with higher efficacy opioids. We used both *in vitro* and *in vivo* models to compare the potency and efficacy of BUP-D2 to BUP. Our *in vitro* approach measured BUP- and BUP-D2-induced G-protein stimulation in homogenates of CHO cells that were stably transfected with human MOR, DOR, and KOR, and determined that BUP-D2 and BUP stimulated G-proteins with equal potency and efficacy. More studies are needed to evaluate the potency and efficacy of BUP-D2 for other pathways downstream of G-protein activation that may affect its pharmacology at the whole-organism level, including β-arrestin-2 recruitment, and modulation of mitogen-activated protein kinase activity

and gating of G-protein coupled inward rectifying potassium channels, as well as adenylyl cyclase activity and cAMP-modulated calcium channel opening and intracellular calcium concentrations. However, our results of the warm water tail withdrawal assay in rats provides the first evidence that BUP-D2 retains potency and efficacy similar to that of BUP for antinociception at the *in vivo* level. Importantly, BUP-D2 exhibited an inverted U-shaped curve that is distinctive of BUP[28]. Increasing antinociception at doses on the ascending limb of the dose-response curve predominantly results from MOR activation[32]. However, at higher doses, activation of nociception/orphanin FQ receptors by BUP is thought to antagonize MOR-induced antinociception, causing the descending limb of the dose-response curve[33]. The close matching of both ascending and descending limbs for BUP and BUP-D2 is compelling evidence that precision deuteration of BUP does not alter these pharmacodynamic effects.

This innovative study takes the important first steps for developing a potential improved treatment for OUD during pregnancy. Future work will determine whether the pharmacokinetics of BUP-D2 differ from those of BUP, such that, as would be predicted, significantly less NorBUP is formed following administration of BUP-D2. Additional questions to be answered include whether such a reduction in maternal plasma concentrations can lower total fetal opioid exposure enough to reduce NOWS.

Acknowledgements and role of the funding sources: We acknowledge and thank the National Institute on Drug Abuse Drug Supply Program for providing norbuprenorphine for this study. This work was supported by the National Institutes of Health (NIH) National Center for Advancing Translational Sciences (NCATS) under award numbers U54TR001629 and KL2TR000063 and by the NIH National Institute on Drug Abuse (NIDA) under award numbers R21DA049585 and T32DA022981. Additional support was provided by the American Society for Pharmacology and Experimental Therapeutics through a Summer Undergraduate Research Fellowship. The funding sources had no involvement in the preparation of the manuscript or the decision to submit it for peer review and publication.

Author contributions: **Janganati**: conceptualization, methodology, validation, formal analysis, investigation, writing—review and editing; **Salazar**: formal analysis, investigation, writing—original draft; **Parks**: investigation, writing—review and editing; **Gorman:** investigation, writing—review and editing; **Prather:** conceptualization, resources, writing—review and editing, funding acquisition; **Crooks**: conceptualization, methodology, resources, writing—review and editing, supervision, funding acquisition; **Brents:**

conceptualization, formal analysis, investigation, writing—review and editing supervision, project administration, funding acquisition.

5. <u>REFERENCES</u>

1. Hirai, A. H.; Ko, J. Y.; Owens, P. L.; Stocks, C.; Patrick, S. W. *JAMA* **2021,** *325,* 146.

2. Haight, S. C.; Ko, J. Y.; Tong, V. T.; Bohm, M. K.; Callaghan, W. M. *MMWR Morb. Mortal. Wkly. Rep.* **2018,** *67,* 845.

3. Jansson, L. M.; Velez, M.; Harrow, C. *J. Opioid Manag.* **2009,** *5,* 47.

4. Strahan, A. E.; Guy, G. P.; Bohm, M.; Frey, M.; Ko, J. Y. *JAMA pediatrics* **2020,** *174,* 200.

5. Maeda, A.; Bateman, B. T.; Clancy, C. R.; Creanga, A. A.; Leffert, L. R. *Anesthesiology* **2014,** *121,* 1158.

6. Brogly, S. B.; Velez, M. P.; Werler, M. M.; Li, W.; Camden, A.; Guttmann, A. *Epidemiology* **2021,** *32,* 448.

7. Lemon, L. S.; Naimi, A.; Caritis, S. N.; Platt, R. W.; Venkataramanan, R.; Bodnar, L. M. *Paediatr. Perinat. Epidemiol.* **2018,** *32,* 213.

8. Azuine, R. E.; Ji, Y.; Chang, H.; Kim, Y.; Ji, H.; DiBari, J.; Hong, X.; Wang, G.; Singh, G. K.; Pearson, C. *JAMA network open* **2019,** *2,* e196405.

9. Nørgaard, M.; Nielsson, M. S.; Heide-Jørgensen, U. *Substance abuse: research and treatment* **2015,** *9,* SART. S23547.

10. Krans, E. E.; Kim, J. Y.; Chen, Q.; Rothenberger, S. D.; James, A. E.; Kelley, D.; Jarlenski, M. P. *Addiction* **2021,** *116,* 3504.

11. Jones, H. E.; Kaltenbach, K.; Heil, S. H.; Stine, S. M.; Coyle, M. G.; Arria, A. M.; O'Grady, K. E.; Selby, P.; Martin, P. R.; Fischer, G. *N. Engl. J. Med.* **2010,** *363,* 2320.

12. Bartu, A. E.; Ilett, K. F.; Hackett, L. P.; Doherty, D. A.; Hamilton, D. *Aust. N. Z. J. Obstet. Gynaecol.* **2012,** *52,* 342.

13. Shah, D.; Brown, S.; Hagemeier, N.; Zheng, S.; Kyle, A.; Pryor, J.; Dankhara, N.; Singh, P. *Springerplus* **2016,** *5,* 854.

14. Kacinko, S. L.; Jones, H. E.; Johnson, R. E.; Choo, R. E.; Huestis, M. A. *Clin. Pharmacol. Ther.* **2008,** *84,* 604.

15. Concheiro, M.; Jones, H. E.; Johnson, R. E.; Choo, R.; Shakleya, D. M.; Huestis, M. A. *Ther. Drug Monit.* **2010,** *32,* 206.

16. Jones, H. E.; Dengler, E.; Garrison, A.; O'Grady, K. E.; Seashore, C.; Horton, E.; Andringa, K.; Jansson, L. M.; Thorp, J. *Drug Alcohol Depend.* **2014,** *134,* 414.

17. O'Connor, A. B.; O'Brien, L.; Alto, W. A. *Eur. Addict. Res.* **2016,** *22,* 127.

18. Griffin, B. A.; Caperton, C. O.; Russell, L. N.; Cabalong, C. V.; Wilson, C. D.; Urquhart, K. R.; Martins, B. S.; Zita, M. D.; Patton, A. L.; Alund, A. W.; Owens, S. M.; Fantegrossi, W. E.; Moran, J. H.; Brents, L. K. *J. Pharmacol. Exp. Ther.* **2019,** .

19. Brown, S. M.; Campbell, S. D.; Crafford, A.; Regina, K. J.; Holtzman, M. J.; Kharasch, E. D. *J. Pharmacol. Exp. Ther.* **2012,** *343,* 53.

20. Alhaddad, H.; Cisternino, S.; Decleves, X.; Tournier, N.; Schlatter, J.; Chiadmi, F.; Risede, P.; Smirnova, M.; Besengez, C.; Scherrmann, J. M.; Baud, F. J.; Megarbane, B. *Crit. Care Med.* **2012,** *40,* 3215.

21. Miwa, G. T.; Walsh, J. S.; Kedderis, G. L.; Hollenberg, P. F. *J. Biol. Chem.* **1983,** *258,* 14445.

22. Harbeson, S. L.; Morgan, A. J.; Liu, J. F.; Aslanian, A. M.; Nguyen, S.; Bridson, G. W.; Brummel, C. L.; Wu, L.; Tung, R. D.; Pilja, L.; Braman, V.; Uttamsingh, V. *J. Pharmacol. Exp. Ther.* **2017,** *362,* 359.

23. Uttamsingh, V.; Gallegos, R.; Liu, J. F.; Harbeson, S. L.; Bridson, G. W.; Cheng, C.; Wells, D. S.; Graham, P. B.; Zelle, R.; Tung, R. *J. Pharmacol. Exp. Ther.* **2015,** *354,* 43.

24. Yadlapalli, J. S. K.; Ford, B. M.; Ketkar, A.; Wan, A.; Penthala, N. R.; Eoff, R. L.; Prather, P. L.; Dobretsov, M.; Crooks, P. A. *Pharmacological research* **2016,** *113,* 335.

25. Cheng, Y.; Prusoff, W. H. *Biochem. Pharmacol.* **1973,** *22,* 3099.

26. Huang, P.; Kehner, G. B.; Cowan, A.; Liu-Chen, L. Y. *J. Pharmacol. Exp. Ther.* **2001,** *297,* 688.

27. Olson, K. M.; Duron, D. I.; Womer, D.; Fell, R.; Streicher, J. M. *PloS one* **2019,** *14,* e0217371.

28. Christoph, T.; Kogel, B.; Schiene, K.; Meen, M.; De Vry, J.; Friderichs, E. *Eur. J. Pharmacol.* **2005,** *507,* 87.

29. Olofsen, E.; Algera, M. H.; Moss, L.; Dobbins, R. L.; Groeneveld, G. J.; van Velzen, M.; Niesters, M.; Dahan, A.; Laffont, C. M. *JCI insight* **2022,** *7,* .

30. Sadee, W.; Rosenbaum, J.; Herz, A. *J. Pharmacol. Exp. Ther.* **1982,** *223,* 157.

31. Cunningham, V. J.; Hume, S. P.; Price, G. R.; Ahier, R. G.; Cremer, J. E.; Jones, A. K. *Journal of Cerebral Blood Flow & Metabolism* **1991,** *11,* 1.

32. Yamamoto, T.; Shono, K.; Tanabe, S. *J. Pharmacol. Exp. Ther.* **2006,** *318,* 206.

33. Lutfy, K.; Eitan, S.; Bryant, C. D.; Yang, Y. C.; Saliminejad, N.; Walwyn, W.; Kieffer, B. L.; Takeshima, H.; Carroll, F. I.; Maidment, N. T. *Journal of Neuroscience* **2003,** *23,* 10331.