Leveraging a machine learning derived surrogate phenotype to improve power for genome-wide association studies of partially missing phenotypes in population biobanks

Zachary R. McCaw*, Jianhui Gao, Xihong Lin, Jessica Gronsbell*.

Correspondence: ZRM zmccaw@alumni.harvard.edu, JG j.gronsbell@utoronto.ca.

Abstract

While population biobanks have dramatically expanded opportunities for genome-wide association studies (GWAS), these large-scale analyses bring new statistical challenges. A key bottleneck is that phenotypes of interest are often partially missing. For example, phenotypes derived from specialized imaging modalities are often only measured for a subset of the cohort. Fortunately, biobanks contain surrogate phenotype information, in the form of routinely collected clinical data, that can often be leveraged to build machine learning (ML) models that accurately predict missing values of the target phenotype. However, simply imputing the missing values of the target phenotype can invalidate subsequent statistical inference. To address this significant barrier, we introduce SynSurr, an approach that jointly analyzes an incompletely observed target phenotype together with its predicted value from an ML model. As the ML model can combine or synthesize multiple sources of evidence to infer the missing phenotypic values, we refer to its prediction as a "synthetic surrogate" for the target phenotype. SynSurr estimates the same effect size as a standard GWAS of the target phenotype, but does so with increased power when the synthetic surrogate is correlated with the target phenotype. Unlike classical imputation, SynSurr does not require that the synthetic surrogate is obtained from a correctly specified generative model, only that it is correlated with the target outcome. SynSurr is also computationally feasible at biobank scale and has been implemented in the open source R package `SurrogateRegression`. In a genome-wide ablation analysis of 2 well-studied traits from the UK Biobank (UKBB), SynSurr consistently recovered more of the associations present in the full sample than standard GWAS using the observed target phenotypes. When applied to 6 incompletely measured body composition phenotypes from the UKBB, SynSurr identified 15.6 times as many genome-wide significant associations than standard GWAS, on average, and did so at 2.9 times the level of significance. These associations were highly enriched for biologically relevant gene sets and overlapped substantially with known body composition associations from the GWAS catalog.

Introduction

Over the last decade, there has been immense interest in conducting genomic research in population biobanks such as FinnGenn ($> 200{,}000$ individuals), the Million Veteran Program ($> 500{,}000$ veterans), and the UK Biobank ($> 400{,}000$ individuals)[1,2]. While these rich data sources have created an unprecedented opportunity to increase both the scope and precision of genome-wide association studies (GWAS), they have simultaneously introduced new statistical challenges[3,4]. Among these challenges, incompletely measured or partially missing phenotypes are a key issue[5]. Missingness often arises due to the difficulty, expense, or invasiveness of ascertaining the target phenotype[6]. Examples of partially missing phenotypes within the UK Biobank (UKBB)[7] include body composition phenotypes obtained from dual-energy X-ray absorptiometry (DEXA) scans[8], neurological[9] and cardiac[10] structural features extracted from functional magnetic resonance imaging, optic morphology parameters extracted from retinal fundus images[11], and sleep wake patterns extracted from wearable accelerometry trackers[12]. Each of these phenotypes was ascertained, at least initially, in only a subset of the full cohort. Restricting the analysis to only the subset individuals with observed target phenotypes would substantially diminish statistical power. To address this practical and statistical issue, we propose a novel method for conducting large-scale GWAS of a partially missing target phenotype that leverages a machine learning (ML) derived surrogate phenotype.

As biobanks are replete with demographic and clinical information from baseline assessments and electronic health records, an ML model can often be developed to accurately predict the partially missing phenotype, increasing power for subsequent GWAS[13–15]. In contrast to the target phenotype, which is only observed for a subset of the cohort, the ML-derived phenotype is available for the entire cohort, enabling GWAS with the full data set. There are now several approaches that can leverage ML-based predictions in performing a GWAS[16,17]. The simplest approach is to perform GWAS using the predicted phenotype as the outcome of interest for all subjects. This approach performs well if the prediction model is highly accurate. However, if the predictions are inaccurate, then genetic variants associated with the ML-derived phenotype may not in fact associate with the target phenotype, and the estimated effect sizes will not reflect the effect of genotype on the target phenotype. Moreover, these approaches may lose power by disregarding the fact that the target phenotype was actually observed for a subset of the cohort[18,19].

Another simple approach is to retain the observed values of the target phenotype, impute the missing values using the ML model, and perform standard GWAS analysis using the pooled observed and imputed target phenotypes. Although simple to implement, such single-

imputation based inference is generally invalid as it ignores imputation uncertainty by treating imputed values as if they were the actual values[17,20–22]. Specifically, single-imputation regards the observed and imputed values as originating from the same distribution, when in fact the imputed values originate from a different distribution and are less variable than their observed counterparts. In the context of linear regression, it is well known that failing to account for the reduced variability induced by imputation will lead to systematic underestimation of the residual variance and consequent inflation of the type I error[23]. In response, multiple imputation (MI)[24–26] has been proposed to overcome the limitations of single imputation. MI generates multiple imputations of missing values, performs the analysis with each set of imputed values, and then combines the resulting estimates, via Rubin's Rules, to correctly accounting for uncertainty in the imputation. The major limitation of MI is that the imputations must come from a correctly specified model[26–29]. When the imputation model is misspecified, as seems likely in practice, the estimates resulting from MI are biased.

To overcome the limitations of imputation-based inference, we propose synthetic surrogate analysis, or "SynSurr" for short, which improves power for GWAS of a partially missing (quantitative) target phenotype. Within a model-building data set, we first train an ML model for predicting the target phenotype on the basis of readily available surrogate data. The model is then transferred to the GWAS data set, where it is applied to generate a "synthetic surrogate" phenotype for all subjects. The partially observed target phenotype and fully observed synthetic surrogate are then *jointly* analyzed within a bivariate outcome framework. SynSurr falls within the previously described surrogate phenotype regression framework[30], but the estimation and inference procedures described here have been accelerated for application at biobank scale. Unlike performing GWAS solely with the ML-derived synthetic surrogate, SynSurr estimates the same effect size as a standard GWAS (using only the observed values of the target phenotype). SynSurr also boosts power to discover variants associated with the target phenotype through a joint analysis that directly leverages the correlation between the target phenotype and its synthetic surrogate. Finally, unlike imputation-based inference, SynSurr does not require the strong and often untenable assumption that the ML model which generates the synthetic surrogate is correctly specified. In fact, SynSurr remains valid and suffers negligible loss of power even when the synthetic surrogate is completely uncorrelated with the target outcome.

We illustrate the methodological advantages and practical implementation of SynSurr through extensive analyses of simulated and real data. Our simulation studies illustrate three key properties of SynSurr, which are backed by theoretical analyses provided in the Supplementary Methods. First, SynSurr is *robust* in the sense that it is no less powerful than standard

GWAS even when the synthetic surrogate is uncorrelated with the target phenotype. Second, SynSurr is more *powerful* than standard GWAS when the synthetic surrogate is correlated with the target phenotype. The power advantage increases with the target-surrogate correlation and the extent of missingness. Third, SynSurr always provides *valid inference*, meaning that the effect size estimates are unbiased and the type I error is properly controlled, even when the ML model that generates the synthetic surrogate is misspecified (i.e. differs from the generative model for the target phenotype).

Our real data analyses of phenotypes from the UK Biobank (UKBB) include an ablation analysis of two phenotypes with minimal missingness, height and forced expiratory volume in 1 second (FEV1), and an application of SynSurr to 6 body composition phenotypes, measured by DEXA, with substantial missingness. The ablation analysis demonstrates that SynSurr and standard GWAS identify all the same variants in the absence of missingness, but that SynSurr is uniformly more powerful in the presence of missingness. SynSurr achieves this power advantage not by inflating the false discovery rate or distorting the estimated effect size, but by leveraging the surrogate outcome to obtain more precise estimation. The application to DEXA phenotypes demonstrates the substantial opportunity for improved power with SynSurr. Compared to standard GWAS, SynSurr identified 15.6 times as many genome-wide significant variants, on average, and did so at 2.9 times the level of significance. Moreover, the variants identified by SynSurr are salient to the phenotypes under study, being significantly enriched for gene sets related to body composition, and overlapping substantially with GWAS catalog variants previously associated with body composition traits.

Results

Overview of Method

Figure 1 provides a graphical overview of a SynSurr GWAS. The overarching goal is to utilize surrogate information to improve power for GWAS of a partially missing target phenotype. The complete data set is first split into model-building and inference data sets. The model-building data set includes the target phenotype and a set of surrogates, but does not require genetics. The model-building data set must also be fully labeled, meaning the target phenotype is available for all subjects. The inference data set, in which GWAS is performed, requires genetics and surrogates, however, it need only be partially labeled, (i.e. the target phenotype is only partially observed).

Within the model-building data set, a ML model is trained to predict the target phenotype

on the basis of surrogate information. The trained model is then transferred to the inference data set, where it is used to predict the target phenotype for all subjects. Because the ML model synthesizes multiple surrogates to predict the target phenotype, we refer to its output as a *synthetic surrogate*. Finally, within the inference data set, the partially missing target phenotype and the fully observed synthetic surrogate are jointly regressed on genotype and covariates to identify genetic variants associated with the target outcome.

In contrast to standard GWAS, which can only incorporate subjects whose target phenotype is observed, SynSurr improves power by allowing subjects whose target phenotypes are missing to enter the analysis. Critically, unlike imputation-based inference, SynSurr does not require that the synthetic surrogate is generated from a correctly specified imputation model. SynSurr is robust, remaining valid and not foregoing power even if the synthetic surrogate is uncorrelated with the target phenotype. Moreover, SynSurr improves power to the extent that the target and synthetic surrogate outcomes are correlated. As we will show, SynSurr provides an unbiased estimate for the effect of genotype on the target outcome, increases estimation precision, and improves power for identifying variants specifically associated with the target phenotype.

The Methods section provides a mathematical description of the SynSurr model and an overview of the estimation procedure. The Supplementary Methods provide a detailed derivation of maximum likelihood estimates for all model parameters, their standard errors, and a Wald test for evaluating the association between genotype and phenotype.

Generation of synthetic surrogates

SynSurr depends on the availability of a synthetic surrogate, $\hat{Y}$, that is constructed for all subjects and ideally correlated with the phenotype of interest, Y. We focus on the setting where the surrogate is a prediction of Y from an ML model. This setting is particularly relevant in population biobanks where certain phenotypes are too invasive, expensive, or time-consuming to measure for the entire cohort, but can be reliably predicted from a model trained on readily available surrogate data (e.g. information from electronic health records, administrative data, or baseline assessments). The surrogate data is distinct from the genetic data G, but may contain covariates X that are also included in the GWAS. In training a prediction model to generate the synthetic surrogate, the goal is to obtain a prediction $\hat{Y}$ that is highly correlated with the target phenotype Y. More specifically, Y and $\hat{Y}$ should be highly correlated after adjusting for the covariates X included in the association model. As detailed in the Supplementary Methods section, a stronger residual correlation leads to

Figure 1: **Graphical overview of a SynSurr GWAS. A.** The complete data set is first split into a fully labeled model-building data set, including the target phenotype and surrogates, and a partially labeled inference data set, which also includes genetics. **B.** Within the fully labeled model-building data set, a ML model is trained to predicted the target phenotype on the basis of surrogates. **C.** The ML model is transferred to the partially labeled inference data set and applied to predict the target outcome for all subjects. The predicted value of the target outcome is referred to as the "synthetic surrogate". Importantly, the synthetic surrogate is maintained as a separate and distinct outcome from the partially missing target phenotype. **D.** Finally, within the inference data set, the partially missing target phenotype and the fully observed synthetic surrogate are jointly regressed on genotype and covariates to identify genetic variants specifically associated with the target outcome.

increased power. To capture the potentially complex relationship between a set of covariates and the phenotype of interest, we recommend that the synthetic surrogate be generated from a nonlinear ML model. For settings with well-defined covariates, tree-based models such as random forest[31] or Extreme Gradient Boosting (XGBoost)[32] generally perform well and are straightforward to train with standard software. Neural network models can also be used and are particularly advantageous for deriving synthetic surrogates from images or free-text data.

We recommend training the prediction model on a data set that is independent of that used for association testing. In practice, given a single large data set, a fraction of the data with observed values for Y would be split off at random to form the model-building data. The remainder of the data, likely including all subjects with unobserved values for Y, would be reserved as the inference data set (see **Figure 1**).

SynSurr overcomes the pitfalls of imputation-based inference

Recall that our goal is to perform inference on the association between Y and G when the outcome Y is partially observed, but a synthetic surrogate $\hat{Y}$ is available for all subjects. To proceed with the analysis, recall that the simplest approach is to focus on $\hat{Y}$, which is observed for all subjects, as the outcome of interest. This approach, however, changes the research question from performing inference on the association between Y and G (i.e. β_G in Equation 4) to performing inference on the association between $\hat{Y}$ and G (i.e. α_G in Equation 4). Moreover, analyzing $\hat{Y}$ instead of Y can reduce power and lead to spurious associations if $\hat{Y}$ is an imperfect proxy for Y. To preserve the original research question, an alternative approach is to generate a new outcome Y^* by imputing missing values of Y with $\hat{Y}$:

$$Y^* = \begin{cases} Y & \text{if } Y \text{ is observed,} \\ \hat{Y} & \text{if } Y \text{ is missing.} \end{cases} \tag{1}$$

Single imputation is generally invalid, as the analytical standard errors will be underestimated[23,26]. To perform valid inference after imputation, multiple imputations of the missing data can be generated, and the resulting point estimates and standard errors combined using Rubin's rules[23,26]. A major limitation of multiple imputation, as demonstrated below, is that its validity generally requires a correctly specified imputation model. A key contribution of the the present work is to provide a method for utilizing $\hat{Y}$ to improve inference on the association between Y and G *without* requiring that $\hat{Y}$ be generated from a correctly specified imputation model.

As an example of the pitfalls that can arise when performing imputation-based inference, we generated phenotypes from the following model:

$$Y = G\beta_G + X\beta_X + \epsilon. \tag{2}$$

Here Y represents the target phenotype, G the genotype, X a correlated covariate, and ϵ a residual. The true effect size is $\beta_G = 0.1$, corresponding to a variant with $h^2 = 1\%$. The total sample size is $n = 10^4$ and 25% of the values of Y are missing. We compare the performance of an *oracle estimator*, which has access to Y before the introduction of missingness, the *standard estimator*, which only has access to the observed values of Y, single and multiple imputation estimators, and the proposed SynSurr estimator. The imputation models are as follows:

1. A correctly specified model, which imputes Y based on G and X.

2. A misspecified model, which imputes Y using G only.

3. A misspecified model, which imputes Y using X only.

Each imputation model was fit on an independent model-building data set of size 10^3 then
applied to generate $\hat{Y}$ for all subjects in the inference data set. The single and multiple
imputation estimators used Y^* as their outcome, constructed via Equation (1), while the
corresponding SynSurr estimator regarded $\hat{Y}$ as the synthetic surrogate. All estimators uti-
lized Equation (2) as the association model, allowing for direct comparison of the resulting
estimates of β_G.

Figure 2 and **Supplementary Table 1** present the point estimates and analytic and em-
pirical standard errors of the imputation-based estimators compared to SynSurr. The oracle
estimator (green) is unbiased for β_G, and as a correctly specified maximum likelihood estima-
tor, its standard error (SE) represents the best that is attainable in the absence of missing
data[33]. The standard estimator (orange) is also unbiased, and because observations with
missing outcomes do not contribute to this estimator, its SE is larger than that of oracle.
The single imputation (SI; red) and multiple imputation (MI; blue) estimators based on the
correctly specified imputation model are unbiased. While the 95% confidence interval (CI)
for the MI estimator has proper coverage, as indicated by the agreement between the ana-
lytical CI (dotted) and the empirical CI (solid), the analytical CI for the SI estimator falls
short of the empirical CI. This occurs because the SE of the SI estimator is underestimated.
Consequently, inference based on the SI estimator would lead to an overstatement of signif-
icance (i.e. inflated type I error)[20]. Estimates based on an incorrectly specified imputation
model are biased, whether the the misspecification was due to omission of the variable of in-
terest (i.e. G) or simply a correlated covariate (i.e. X)[25]. In contrast, the synthetic surrogate
estimator (purple) is unbiased regardless of what collection of covariates is used to generate
$\hat{Y}$. Because $\hat{Y}$ was generated from a linear model, and the covariates used to generate $\hat{Y}$
are conditioned on the inference model, the SynSurr estimator is not notably more efficient
than the standard estimator in this simulation. In practice, to improve the surrogacy of $\hat{Y}$
and allow the SynSurr estimator to realize a statistical efficiency gain, $\hat{Y}$ should either be
generated from a non-linear model or incorporate covariates apart from those included in
the GWAS model (ideally both).

Figure 2: **Unlike imputation-based estimators, SynSurr is robust to misspecification of the imputation model.** The true value for the parameter of interest is $\beta_G = 0.1$, corresponding to a variant with $h^2 = 1\%$. For each estimator, the mean value across 10^3 simulations is shown by the point, and two 95% confidence intervals (CIs) are presented: the dotted CI is based on the root-mean-square analytical standard error (SE) while the solid CI is based on the empirical SE. The oracle estimator has access to the complete version of Y, before 25% of values were set to missing. The standard estimator has access to the observed values of Y only. The imputation-based estimators (SI: single imputation, MI: multiple imputation) impute the missing values of Y using an imputation-model fit on an independent data set. The set of covariates used to fit the imputation model are shown as a tuple: the imputation model based on G and X is correctly specified, whereas that based on G alone or X alone is misspecified. The SynSurr estimator (SS) jointly analyzes the partially missing Y with the synthetic surrogate $\hat{Y}$, where $\hat{Y}$ is generated for all subjects from the imputation model.

SynSurr is robust to uninformative surrogates and yields more precise inference with informative surrogates

Unlike imputation-based inference, which is sensitive to correct specification of the imputation model, SynSurr is robust to the choice of synthetic surrogate in that it (i) consistently estimates the effect of the genotype on the target phenotype regardless of how well the surrogate predicts the target phenotype and (ii) provides improved power over standard GWAS when the synthetic surrogate is correlated with the target phenotype. To demonstrate these points, we again simulated phenotypes from Equation (2) and varied the proportion of subjects with missing phenotypes and the correlation between Y and $\hat{Y}$ (see Supplementary Section 3.1). **Supplementary Figure 1** presents box plots of $\hat{\beta}_G - \beta_G$ for the standard GWAS estimator and the SynSurr estimator under different levels of missingness in Y. In

panel A, the synthetic surrogate $\hat{Y}$ is completely uncorrelated with, and in fact independent of, the phenotype of interest Y. Nevertheless, the synthetic surrogate estimator is unbiased and no less efficient than the standard GWAS estimator, as indicated by consistent widths of the box plots. This demonstrates that SynSurr is robust to the use of an uninformative surrogate. However, as demonstrated in panel B, when $\hat{Y}$ is correlated with Y, the precision of the SynSurr estimator increases with the number of subjects with missing phenotypes (also see **Supplemental Figure 2**). These qualitative observations are supported by a quantitative comparison of standard errors in **Supplementary Table 2** as well as a theoretical analysis in the Supplementary Methods.

SynSurr controls type I error and increases power relative to standard GWAS

Building on the findings of the previous section, we again simulated phenotypes from Equation (2) to evaluate the type I error and power of SynSurr across various missing rates, choices of synthetic surrogate, and levels of SNP heritability. **Figure 3** demonstrates that, across missingness rates and target-surrogate correlations, the SynSurr p-values are uniformly distributed under the null hypothesis of no association. **Supplemental Table 3** presents the type I error and power at a SNP heritability of $h^2 = 0.5\%$ ($\beta_G \approx 0.07$), together with the average χ^2 statistics. The type I error is consistently controlled across missing rates and choices of the synthetic surrogate. Consonant with the previous section and with the efficiency analysis in the Supplementary Methods, **Supplemental Table 3** demonstrates that the power of SynSurr increases with the target-synthetic surrogate correlation and level of target missingness. For instance, when the missingness rate is 90% and the correlation between the synthetic surrogate and the target phenotype is 0.75, there is a 27% increase in power relative to the standard analysis at the same missing rate. The relative efficiency analysis in **Supplemental Figure 2** further supports this finding as the SynSurr estimator exhibits lower standard error compared to standard across increasing missingness rate and target-surrogate correlation. Moreover, **Figure 4** illustrates the benefit of SynSurr with respect to power across SNP heritabilities ranging from 0.1% to 1.0%. As the proportion of subjects with missing target phenotypes increases, the benefit of SynSurr with a well-correlated surrogate phenotype is increasingly apparent. Interestingly, the relative efficiency of SynSurr does not depend on the SNP heritability (**Supplemental Figure 3**).

Figure 3: **Uniform quantile-quantile plots for the SynSurr p-values under the null hypothesis of no association across various missingness rates and target-surrogate correlations.** In all cases, the number of subjects with observed phenotypes was $n = 10^3$. The number of subjects with missing phenotypes was varied to achieve the indicated level of missingness. The synthetic surrogate has correlation $\rho \in \{0.00, 0.25, 0.50, 0.75\}$ with the target phenotype. The number of simulation replicates is 10^6.

Figure 4: **Power of SynSurr across various missing rates, target-surrogate correlations, and SNP heritabilities.** In all cases, the number of subjects with observed phenotypes was $n = 10^3$. The number of subjects with missing phenotypes was varied to achieve the indicated level of missingness. In each panel, the synthetic surrogate has correlation $\rho \in \{0.00, 0.25, 0.50, 0.75\}$ with the target phenotype and the SNP heritability was varied from 0.1% to 1%. When there is no missingness, SynSurr is equivalent to the standard analysis and shows no variation across values of ρ. The power of SynSurr increases with increasing missingness and target-surrogate correlation. The number of simulation replicates is 10^4.

Evaluation on UK Biobank (UKB) data

We next demonstrate the advantages of SynSurr over a standard GWAS with several analyses of the UKBB. We first consider GWAS of two well-studied traits, height and FEV1. These phenotypes are measured for a majority of the cohort and enable us to compare the performance of SynSurr with standard GWAS as the target phenotype is increasingly ablated. We then perform SynSurr analysis of a 6 incompletely observed body composition phenotypes measured by DEXA scan.

SynSurr recovers more oracle associations than standard GWAS as the target phenotype is increasingly ablated

The genotype and sample quality control steps are described in the Methods. Our analyses included 435,468 directly genotyped genetic variants. In total, 349,474 unrelated subjects were included in the height GWAS and 308,518 in the FEV1 GWAS. For each phenotype, sets of kin up to the third-degree were identified. One subject was randomly allocated to the GWAS data set and the remaining subjects to the model-building data set. The model-building data were used to construct a random forest model for predicting the target phenotype on the basis of age, sex, and various anthropomorphic measurements (see Supplementary Section 5.2). Note that having related subjects in the model-building and inference (GWAS) data sets is not problematic because (i) the GWAS data set is not being used to evaluate generalization performance and (ii) the GWAS data set itself contains no related subjects.

For each phenotype, we performed an oracle GWAS prior to the introduction of missingness. This GWAS establishes the number of genome-wide significant (GWS) associations that could be detected if the target phenotype were fully observed. Then, 25%, 50%, 75%, and 90% of the target phenotypes were ablated at random. Both standard and SynSurr GWAS were performed on the remaining data. Prior to the introduction of missingness, the correlation between the target and synthetic surrogate phenotypes in the GWAS data set was $R^2 = 0.67$ for height and $R^2 = 0.51$ for FEV1. Scatter plots of predicted vs. observed height and FEV1 in the model-building and GWAS data are shown in **Supplemental Figures 4-5**. **Table 1** presents the numbers of oracle associations recovered by both the standard and SynSurr GWAS. SynSurr consistently recovers a higher proportion of the oracle associations than standard GWAS. Importantly, as demonstrated in **Supplementary Table 5**, SynSurr does not achieve this higher recovery by having a higher false discovery rate. Rather, SynSurr is leveraging the correlated surrogate outcome to obtain more precise estimates of the standard error (**Supplementary Table 6**; also see **Supplementary Table 7**). For example,

at 90% missingness, the mean square standard error of the SynSurr estimator, across oracle associations, is 7.4×10^{-5} (height) and 8.3×10^{-5} (FEV1) compared with 9.8×10^{-5} (height) and 9.7×10^{-5} (FEV1) for the standard estimator (relative improvements of 33% and 17%, respectively). **Supplementary Figure 7** verifies that SynSurr is estimating the same effect size as the oracle GWAS (so too is standard GWAS, see **Supplementary Figure 8**). Even with 90% of target phenotypes ablated, the R^2 for the effect sizes between SynSurr and oracle is 0.90 for height and 0.87 for FEV1. With 50% of target phenotypes ablated, the R^2 rises to 0.99 and 0.98 respectively, and in the absence of missingness, $R^2 = 1.00$.

Table 1: **Number of genome-wide significant SNPs recovered by standard and SynSurr GWAS across increasing ablation of the target phenotype.** The oracle method establishes the number of genome-wide significant (GWS) variants ($p < 5 \times 10^{-8}$) that would be identified in the absence of missingness. Missingness was introduced by ablating 25%, 50%, 75%, and 90% of the target phenotypes. Standard and SynSurr GWAS were performed on each of the ablated data sets. The number of full-sample GWS variants recovered by each method is reported. Also see Supplementary Tables 5-7.

Missing Rate(%)	Height				FEV1			
	n_{obs}	Oracle	Standard	SynSurr	n_{obs}	Oracle	Standard	SynSurr
0	0	7,177	7,177(100%)	7,177(100%)	0	974	544(55.85%)	544(55.85%)
0.25	87,368	7,177	4,896(68.22%)	5,305(73.92%)	7,7129	974	335(34.39%)	357(36.65%)
0.5	174,737	7,177	2,742(38.21%)	3,421(47.67%)	154,259	974	178(18.28%)	208(21.36%)
0.75	262,105	7,177	834(11.62%)	1,243(17.32%)	231,388	974	42(4.31%)	49(5.03%)
0.9	314,526	7,177	192(2.68%)	329(4.58%)	277,666	974	0(0%)	0(0%)

SynSurr identifiers more associations with partially missing body composition phenotypes than standard GWAS

We next performed SynSurr analyses of 6 incompletely measured DEXA phenotypes: android, arm, gynoid, leg, trunk, and total mass. In each case, the model-building data set included 4,584 subjects with observed target outcomes, while the GWAS data set included 347,498 subjects, 29,577 (8.5%) of which have observed target phenotypes. Within the model-building data set, a random forest was trained to predict the DEXA phenotype on the basis of age, sex, height, body weight, and body mass index. The fitted models were then transferred to the GWAS data set, where a synthetic surrogate outcome was generated for all subjects. Among subjects in the inference data set with an observed DEXA phenotype, the average R^2 between the target phenotype and the synthetic surrogate was 0.78.

Figure 5 presents the number of GWS associations ($p < 5 \times 10^{-8}$) for standard GWAS, based on the subjects with observed DEXA measurements only, and SynSurr GWAS, as well as the average χ^2 statistic at the union of variants that reached significance in either GWAS.

Standard GWAS identified 7 GWS variants, while SynSurr GWAS identified between 46 and 172 GWS variants, for an average of 109 (15.6-fold improvement). The average χ^2 statistic at GWS variants was 46.5 for SynSurr GWAS, compared with 15.8 for standard GWAS, a 2.9-fold improvement. To check control of the type I error, the SynSurr analysis was repeated using permuted phenotypes. The uniform quantile-quantile plots in **Supplemental Figure 9** show no evidence of type I error inflation under the null.

The mirrored Manhattan plots in **Supplementary Figure 10** indicate that SynSurr did not merely inflate the significance of standard GWAS results, but elevated some signals while suppressing others. To further validate our putative associations, we overlapped our GWS variants with publicly available associations from the GWAS catalog[34] for the following traits: body fat distribution, body fat percentage, fat body mass, lean body mass (**Supplemental Figure 11**). On average, 74.5% of the DEXA associations identified by SynSurr overlapped with a previously reported body composition association from the GWAS catalog. To investigate the biological function of the GWS variants, we performed gene set enrichment analysis using FUMA[35]. On average, 392 gene sets were enriched among the SynSurr results, while no significant enrichment were identified with the standard GWAS results. For all phenotypes, gene sets related to body fat distribution were among the most significant, and numerous enrichments related to anthropometric traits were identified (**Supplementary Data 1**).

Discussion

We introduced SynSurr, a simple and computationally tractable procedure for leveraging a surrogate phenotype generated by a machine learning (ML) model to improve inference on partially missing target outcome. In the course of a typical SynSurr analysis, the data are split into disjoint model-building and GWAS data sets. A ML model trained in the former is applied to generate the synthetic surrogate in the latter. GWAS is performed by jointly regressing the partially missing target phenotype and its synthetic surrogate on genotype and covariates. Unlike imputation, SynSurr is robust to the choice of surrogate, not requiring the often untenable assumption that the synthetic surrogate is generated from a correctly specified imputation model. We demonstrated that a SynSurr GWAS is robust, powerful, and valid: providing power at least equaling, but typically exceeding standard GWAS, while controlling the type I error and estimating the proper effect size. The power advantage afforded by SynSurr increases with the extent of target missingness and the target-surrogate correlation. In real data ablation analyses, SynSurr consistently recovered more of the full-sample associations than standard GWAS, and did so without either inflating the false discovery rate or distorting the estimated effect sizes. Instead, SynSurr leverages cor-

Figure 5: **Comparing SynSurr and standard GWAS with respect to the number and significance of genome-wide significant associations for body composition traits. A.** Number of genome-wide significant associations ($p < 5 \times 10^{-8}$) with DEXA body composition traits for standard and synthetic surrogate (SynSurr) GWAS. **B.** Average χ^2 statistic at the union of variants that reached genome-wide significance under either method.

relation between the target and synthetic surrogate to obtain more precise estimates of the parameter of interest (i.e. to reduce the standard error). Finally, when applied to GWAS of 6 partially missing body composition phenotypes from the UKB, SynSurr identified 15.6 times more genome-wide significant associations than standard GWAS. These associations were significantly enriched for gene sets related to body composition and overlapped considerably with body composition associations from the GWAS catalog.

Although performing valid inference by means of imputation is possible, it typically requires

a correctly specified imputation model and the use of multiple imputations to correctly estimate standard errors. Here we propose jointly modeling a partially missing outcome together with its predicted value, rather than replacing missing values of the target outcome by their predicted values. In doing so, we avoid the often unverifiable assumption that the imputation model is correctly specified, and obtain valid statistical inference in a single shot. This robust approach to handling missing data has broad potential for applications outside of GWAS.

Our current approach to SynSurr involves splitting the data into independent model-building and GWAS data sets. For our analyses in UKB, the model-building set was composed of those individuals with observed target outcomes who could not be included in the GWAS set because a relative was already present. For the relatively simple class of random forest models utilized for generating synthetic surrogates, our model-building data sets were sufficiently sized to obtain reasonable target-surrogate correlations. For more complex models (e.g. deep neural networks), larger model-building sets may be required. A trade-off arises, in terms of power for the final GWAS, between allocating more subjects to the model-building data set and retaining more subjects in the GWAS data set. By allocating more subjects to the former, we potentially obtain better target-surrogate correlations, but at the cost of having fewer subjects available for statistical inference in the GWAS. Studying this power trade-off and principled approaches to determining the size of the model-building set is an avenue for future research.

Lastly, SynSurr is not without limitations. First, the method was derived under joint normality of the target phenotype and the synthetic surrogate. As phenotypes and surrogates may follow non-normal distributions, we suggest applying the rank-based inverse normal transformation (INT) to both the target and the surrogate prior to analysis[36], and did so for all analyses presented in this paper. While INT only guarantees marginal normality of each phenotype, it is commonly used in GWAS and mitigates the extent to which the joint distribution can depart from bivariate normality[30]. Our current work focuses on extending SynSurr to a broader class of distributions by dropping the bivariate normality assumption. This can be achieved by replacing the score equations derived from maximum likelihood theory with a set of weighted estimating equations[37]. Moreover, we plan to extend SynSurr to generalized linear models with outcomes from the exponential family using a zero-mean augmentation approach[19,38]. Second, the validity of SynSurr assumes that the missing phenotypes are missing at random (see Supplemental Methods). It is of future research interest to extend SynSurr to the setting when data are missing nor at random. Third, SynSurr does not currently accommodate related individuals. Future work will extend SynSurr to

allow for related individuals by adding a subject-specific random effect with cross-subject covariance proportional to the sample kinship matrix[39].

Methods

Standard GWAS

For the standard GWAS, the target phenotype Y is regressed on genotype G and covariates X among those subjects whose target phenotypes are observed:

$$Y = G\beta_G + X\beta_X + \epsilon.$$

The null hypothesis $H_0 : \beta_G = 0$ is evaluated using the standard Wald test[40]:

$$T_W = \frac{\hat{\beta}_G^2}{\mathbb{SE}^2(\hat{\beta}_G)}, \tag{3}$$

where $\hat{\beta}_G$ is the ordinary least squares (OLS) estimate of the genetic effect, and $\mathbb{SE}(\hat{\beta}_G)$ is the corresponding standard error.

SynSurr GWAS

Within the inference data set, suppose the target phenotype Y is observed for n_{obs} subjects and missing for n_{miss}. Let $n = n_{\text{obs}} + n_{\text{miss}}$ denote the total sample size. Define R as an indicator that Y was observed:

$$R = \begin{cases} 1 & \text{if } Y \text{ is observed,} \\ 0 & \text{if } Y \text{ is missing.} \end{cases}$$

Denote by $\hat{Y}$ a synthetic surrogate for Y that is available for all n subjects. We recommend that $\hat{Y}$ be obtained from a nonlinear ML model, such as a random forest or neural network, trained on an independent model-building data set. Let G denote the genotype and X a vector of covariates, such as age, sex, and genetic principal components. To make use of both Y and $\hat{Y}$ in evaluating the association between Y and G, SynSurr utilizes the joint association model:

$$\begin{bmatrix} Y \\ \hat{Y} \end{bmatrix} \mid Z = \begin{bmatrix} Z & 0 \\ 0 & Z \end{bmatrix} \begin{bmatrix} \beta \\ \alpha \end{bmatrix} + \begin{bmatrix} \epsilon_T \\ \epsilon_S \end{bmatrix} \tag{4}$$

where $Z = (G, X)^T$, $\beta = (\beta_G, \beta_X)^T$, $\alpha = (\alpha_G, \alpha_X)^T$, and the residuals $(\epsilon_T, \epsilon_S)^T$ follow a bivariate normal distribution:

$$\begin{bmatrix} \epsilon_T \\ \epsilon_S \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \Sigma_{TT} & \Sigma_{TS} \\ \Sigma_{ST} & \Sigma_{SS} \end{bmatrix} \right).$$

(Notationally, T denotes the target phenotype, namely Y, and S denotes the synthetic surrogate, namely $\hat{Y}$.) In model (4), β_G is the parameter of interest, which quantifies the association between the genotype and the target phenotype. This is the same parameter estimated by standard GWAS of Y on G among the n_{obs} subjects with observed target phenotypes (see robustness simulations). Studies that regress the synthetic phenotype $\hat{Y}$ on G estimate α_G instead of β_G, and these effect sizes can differ. SynSurr enables inference on β_G while making use of all n subjects. Moreover, SynSurr is computationally tractable at biobank scale, requiring only two ordinary least squares regressions:

1. First, among all n subjects, regress $\hat{Y}$ on Z to obtain an estimate of α.

2. Second, among the n_{obs} subjects with observed phenotypes ($R = 1$), regress Y on $(\hat{Y}, Z)^T$ to obtain an estimate of the associated regression coefficient, denoted $(\delta, \gamma)^T$.

The validity of this two-step approach is demonstrated through a reparameterization of the log-likelihood function which demonstrates that the association parameter can be recovered as $\beta = \gamma + \delta\alpha$. Details of this equivalence, as well as derivation of the Wald test for SynSurr, are provided in the Supplementary Methods.

UKBB genotype and sample quality control

Our UKBB data release contains genotypes for 488,377 subjects and 784,256 directly geno-typed variants. Prior to the analysis, we performed the following common quality control steps[41]:

1. Excluded individuals with $> 10\%$ missing genotypes.

2. Excluded SNPs with a genotyping rate $< 90\%$.

3. Excluded SNPs with a Hardy-Weinberg Equilibrium $p < 10^{-5}$.

4. Excluded SNPs with MAF $< 1\%$.

5. Included only who self-identified as 'White British' and have very similar genetic ancestry based on a principal components analysis of the genotypes (Data-Field 22006).

6. Selected one member of each set of subjects with a kinship coefficient greater than 0.0625 (the threshold for third-degree relatives) for the GWAS data set, and allocated the remaining subjects to the model-building data set.

The results from our genotype QC are summarized in **Supplementary Table 4**. Note that we allowed the model-building data set to contain related individuals.

GWAS catalog overlap analysis

Summary statistics for `body fat distribution`, `body fat percentage`, `fat body mass`, `lean body mass` were downloaded from the NHGRI-EBI GWAS catalog[34]. After concatenating and reducing to 1 record per unique combination of chromosome and base pair, this set contained 984 associated variants. Overlap of study variants with GWAS catalog variants was assessed using the `GenomicRanges`[42] package in `R`[43]. A study variant was considered overlapped if fell within 250 kb of GWAS catalog variant for one of the aforementioned traits.

Data availability

This work used genotypes and phenotypes from the UK Biobank study (https://www.ukbiobank.ac.uk) and our access was approved under application 64875.

Code availability

`SurrogateRegression` is available as an `R`[43] package on the Comprehensive R Archive Network: https://CRAN.R-project.org/package=SurrogateRegression.

Replication code for the analyses presented in this paper is available on GitHub at: https://github.com/jianhuig/SyntheticSurrogateAnalysis.

References

1. Gaziano, J. M. *et al.* Million Veteran Program: A mega-biobank to study genetic influences on health and disease. *Journal of clinical epidemiology* **70,** 214–223 (2016).

2. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562,** 203–209 (2018).

3. Beesley, L. J. *et al.* The emerging landscape of health research based on biobanks linked to electronic health records: Existing resources, statistical challenges, and potential opportunities. *Statistics in medicine* **39,** 773–800 (2020).

4. Tan, V. Y. & Timpson, N. J. The UK Biobank: A Shining Example of Genome-Wide Association Study Science with the Power to Detect the Murky Complications of Real-World Epidemiology. *Annual Review of Genomics and Human Genetics* **23** (2022).

5. Wei, W.-Q. & Denny, J. C. Extracting research-quality phenotypes from electronic health records to support precision medicine. *Genome medicine* **7,** 1–14 (2015).

6. Banda, J. M., Seneviratne, M., Hernandez-Boussard, T. & Shah, N. H. Advances in electronic phenotyping: from rule-based definitions to machine learning models. *Annual review of biomedical data science* **1,** 53 (2018).

7. Allen, N. E., Sudlow, C., Peakman, T., Collins, R., *et al. UK biobank data: come and get it.* 2014.

8. Littlejohns, T. J. *et al.* The UK Biobank imaging enhancement of 100,000 participants: rationale, data collection, management and future directions. *Nature communications* **11,** 1–12 (2020).

9. Elliott, L. *et al.* Genome-wide association studies of brain imaging phenotypes in UK Biobank. *Nature* **562,** 210–216 (2018).

10. Pirruccello, J. *et al.* Analysis of cardiac magnetic resonance imaging in 36,000 individuals yields genetic insights into dilated cardiomyopathy. *Nat Commun* **11,** 2254 (2020).

11. Alipanahi, B. *et al. Large-scale machine learning-based phenotyping significantly improves genomic discovery for optic nerve head morphology* 2020. arXiv: 2011.13012 [q-bio.GN].

12. Li, X. & Zhao, H. Automated feature extraction from population wearable device data identified novel loci associated with sleep and circadian rhythms. *PLoS Genet* **16,** e1009089 (2020).

13. Zhang, Y. *et al.* High-throughput phenotyping with electronic medical record data using a common semi-supervised approach (PheCAP). *Nature protocols* **14,** 3426–3444 (2019).

14. Liao, K. P. *et al.* High-throughput multimodal automated phenotyping (MAP) with application to PheWAS. *Journal of the American Medical Informatics Association* **26,** 1255–1262 (2019).

15. Yang, S., Varghese, P., Stephenson, E., Tu, K. & Gronsbell, J. Machine learning approaches for electronic health records phenotyping: A methodical review. *medRxiv* (2022).

16. Hormozdiari, F. *et al.* Imputing phenotypes for genome-wide association studies. *The American Journal of Human Genetics* **99,** 89–103 (2016).

17. Wang, S., McCormick, T. H. & Leek, J. T. Methods for correcting inference based on outcomes predicted by machine learning. *Proceedings of the National Academy of Sciences* **117,** 30266–30275 (2020).

18. Huang, J. *et al.* PIE: A prior knowledge guided integrated likelihood estimation method for bias reduction in association studies using electronic health records data. *Journal of the American Medical Informatics Association* **25,** 345–352 (2018).

19. Tong, J. *et al.* An augmented estimation procedure for EHR-based association studies accounting for differential misclassification. *Journal of the American Medical Informatics Association* **27,** 244–253 (2020).

20. Rubin, D. B. Inference and missing data. *Biometrika* **63,** 581–592 (1976).

21. Hubbard, R. A., Tong, J., Duan, R. & Chen, Y. Reducing bias due to outcome misclassification for epidemiologic studies using EHR-derived probabilistic phenotypes. *Epidemiology* **31,** 542–550 (2020).

22. Hong, C., Liao, K. P. & Cai, T. Semi-supervised validation of multiple surrogate outcomes with application to electronic medical records phenotyping. *Biometrics* **75,** 78–89 (2019).

23. Little, R. J. & Rubin, D. B. *Statistical Analysis with Missing Data* 2nd (John Wiley & Sons, 2002).

24. Rubin, D. *Multiple Imputation for Nonresponse in Surveys* (John Wiley & Sons, 1987).

25. Rubin, D. B. Multiple imputation after 18+ years. *Journal of the American statistical Association* **91,** 473–489 (1996).

26. Van Buuren, S. *Flexible Imputation of Missing Data* 2nd (Chapman and Hall/CRC, 2018).

27. Bartlett, J. W. & Hughes, R. A. Bootstrap inference for multiple imputation under uncongeniality and misspecification. *Statistical methods in medical research* **29,** 3533–3546 (2020).

28. Austin, P. C., White, I. R., Lee, D. S. & van Buuren, S. Missing data in clinical research: a tutorial on multiple imputation. *Canadian Journal of Cardiology* **37,** 1322–1331 (2021).

29. Murray, J. S. Multiple imputation: a review of practical and theoretical findings. *Statistical Science* **33,** 142–159 (2018).

30. McCaw, Z. R., Gaynor, S. M., Sun, R. & Lin, X. Leveraging a surrogate outcome to improve inference on a partially missing target outcome. *Biometrics* **Online ahead of print** (2022).

31. Breiman, L. Random forests. *Machine learning* **45,** 5–32 (2001).

32. Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. *CoRR* **abs/1603.02754.** arXiv: 1603.02754. http://arxiv.org/abs/1603.02754 (2016).

33. Casella, B. & Berger, R. *Statistical Inference.* 2nd ed. (Duxbury/Thomson Learning, Pacific Grove, CA, 2002).

34. Buniello, A., MacArthur, J., Cerezo, M., *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* **47,** D1005–D1012 (2019).

35. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* **8,** 1826 (2017).

36. McCaw, Z., Lane, J., Saxena, R., Redline, S. & Lin, X. Operating characteristics of the rank-based inverse normal transformation for quantitative trait analysis in genome-wide association studies. *Biometrics* **76,** 1262–1272 (2020).

37. Robins, J. & Rotnitzky, A. Semiparametric Efficiency in Multivariate Regression Models with Missing Data. *Journal of the American Statistical Association* **90,** 122–129 (1995).

38. Wang, X. & Wang, Q. Semiparametric linear transformation model with differential measurement error and validation sampling. *Journal of Multivariate Analysis* **141,** 67–80 (2015).

39. Po-Ru, L., Tucker, G., Bulik-Sullivan, B., Vilhjalmsson, B., Finucane, H., *et al.* Efficient Bayesian mixed model analysis increases association power in large cohorts. *Nature Genetics* **47,** 284–290 (2015).

40. Seber, G. *The Linear Model and Hypothesis. A General Unifying Theory* 1st ed. (Springer Cham, 2015).

41. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *The American Journal of Human Genetics* **81,** 559–575 (2007).

42. Lawrence, M., Huber, W., Pages, H., *et al.* Software for Computing and Annotating Genomic Ranges. *PLoS Comput Biol* **9,** e1003118 (2013).

43. R Core Team. *R: A Language and Environment for Statistical Computing* R Foundation for Statistical Computing (Vienna, Austria, 2022). `https://www.R-project.org/`.

Acknowledgments

This work was supported by the National Institutes of Health grants R35 CA197449 and F31 HL140822 (to ZRM); and R35-CA197449, U19-CA203654, R01-HL163560, U01-HG012064, and U01-HG009088 (to XL); and the Natural Sciences and Engineering Research Council of Canada grant RGPIN-2021-03734 and a Connaught New Researcher Award (to JeG).

Author information

Authors and Affiliations

These authors jointly supervised this work: Xihong Lin and Jessica Gronsbell.

Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, US.
Zachary R. McCaw, Xihong Lin

Department of Statistical Sciences, University of Toronto, Toronto, Ontario, CA.
Jianhui Gao, Jessica Gronsbell

Department of Statistics, Harvard University, Cambridge, MA, US.
Xihong Lin

Department of Computer Science, University of Toronto, Toronto, Ontario, CA.
Jessica Gronsbell

Department of Family & Community Medicine, University of Toronto, Toronto, Ontario, CA.
Jessica Gronsbell

Contributions

ZRM, XL, and JeG designed the study and the experiments. ZRM implemented the software with input from XL. ZRM and JiG performed the simulation experiments. JiG conducted analyses of the UK Biobank data. ZRM performed the overlap analysis. ZRM and JeG wrote the first draft of the manuscript and all co-authors provided intellectual revisions.