Title: Shifts in competitive structures can drive variation in species' phenology

Authors: Patricia Kaye T. Dumandan[1], Glenda M. Yenni[2], and S. K. Morgan Ernest[2]

Institutional Affiliations:

1. School of Natural Resources and Environment, University of Florida, Gainesville, FL, USA

2. Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL, USA

Corresponding author: Patricia Kaye T. Dumandan (pdumandan@ufl.edu)

ABSTRACT

For many species, a well-documented response to anthropogenic climate change is a shift in various aspects of its life-history, including its timing or phenology. Often, these phenological shifts are associated with changes in abiotic factors used as proxies for resource availability or other suitable conditions. Resource availability, however, can also be impacted by competition, but the impact of competition on phenology is less studied than abiotic drivers. We fit generalized additive models (GAMs) to a long-term experimental dataset on small mammals monitored in the southwestern United States and show that altered competitive landscapes can drive shifts in breeding timing and prevalence, and that, relative to a dominant competitor, other species exhibit less specific responses to environmental factors. These results suggest that plasticity of phenological responses, that is often described in the context of annual variation in abiotic factors, can occur in response to biotic context as well. Variation in phenological responses under different biotic conditions shown here further demonstrates that a more nuanced understanding of shifting biotic interactions is useful to better understand and predict biodiversity patterns in a changing world.

INTRODUCTION

Characterizing species' responses to various environmental conditions is useful in determining their sensitivity and adaptive capacity to the consequences of ongoing global environmental change (Glick et al., 2011). Among the known responses to climate change, shifts in phenology or the timing of key life-history events of organisms, are the most well-documented (Cohen et al., 2018; Roslin et al., 2021; Visser & Both, 2005; Walther et al., 2002). Often, the timing of important phenological events (e.g., reproduction) match with periods of high resource

availability to support high fitness and survival costs (Bradshaw & Holzapfel, 2007; Buss et al.,

2021; Forchhammer et al., 1998; Helm et al., 2013; Reed et al., 2013; Walther et al., 2002).

Fitness costs of reproduction are often attributed to the increased energetic demands of

maintaining and developing reproductive organs and behaviors (Audzijonyte & Richards, 2018)

and survival costs are incurred from increased predation risk (Kelt et al., 1995). To avoid

unfavorable conditions during the reproductive period, species may use abiotic (i.e., temperature,

day length, precipitation) and biotic cues (i.e., flowering, prey availability) to regulate and shape

reproductive phenology (Denny et al., 2014; Kenagy & Bartholomew, 1981; Wolkovich et al.,

2014).

Timing reproductive activities to overlap with periods of high resource availability is

important for consumer species because decoupled resource and consumer phenology can

negatively impact production and survival of offspring (Thackeray et al., 2010; Wann et al.,

2019). Thus, many studies assess phenological drivers related to either reliable environmental

cues of resource availability or biotic cues related directly to the emergence or abundance of prey

(i.e., resource) species (DeLucia et al., 2012; Gordo & Sanz, 2005; Pearce-Higgins & Yalden,

2004). However, resource availability is determined by more than just the intrinsic population

dynamics of prey. Competition from other individuals can deplete important resources and create

resource limitation even when resource supply is initially high. Recent theoretical models (Carter

& Rudolf 2022) have demonstrated that resource depletion from competition (intra- or inter-

specific) favors early initiators of reproduction over latter initiators and drives high phenological

synchrony as a result.

Understanding how competition impacts phenology is important because, as species enter

and leave communities in response to environmental changes, the timing and magnitude of

resource availability perceived by resident species may also change (Carter & Rudolf 2022) and

promote shifts in both timing and the apparent response to abiotic cues. If phenology is fixed to

specific abiotic cues or if competition is a process of lesser importance, then we would expect

removing competitors to have little impact on the timing or magnitude of reproductive events.

We would also see little difference in the important drivers associated with reproductive activity

under different competitive scenarios. However, changes to reproductive phenology in response

to competition are possible, given its potential to alter resource availability. The study by Carter

and Rudolf (2022) posits that competitive impacts can differ depending on whether a focal

species initiates their phenology concurrent to or before their competitor. Late initiation allows a

dominant competitor to draw down valuable resources, reducing reproductive success for the

focal species. Moreover, assuming plasticity in an individual's ability to decide to enter a

reproductive state, the presence of a competitor may drive decreases in the prevalence of

reproductive individuals. If reproductive timing is a fixed trait, only individuals with certain

phenological timings that reduce the competitive impact may enter breeding condition. However,

while species' interactions (i.e., competition) can theoretically impact reproductive timing and

investment, empirical assessment of these impacts are limited.

Long-term natural field experiments, where competitive networks are explicitly

manipulated, are well-suited for investigating the role of intra- and inter- specific competition in

driving population processes, including reproduction (Ernest et al., 2016; Kelt et al., 1995,

Mysterud et al., 2008; Ostfeld, 1985). In long-term field experiments where dominant

competitors are removed, the effect of their loss on the reproductive phenology of other species

in the system can be evaluated by comparing the reproductive parameters of a focal species in

plots with and without a dominant competitor. Here, we used long-term data on small mammals

monitored in an experimental field site near Portal, Arizona to assess the effects of competition

on reproductive phenology. This long-term experiment (1977 to present) in the southwestern

USA generates fine-scale (monthly) data on desert rodents under different biotic contexts by

experimentally removing a dominant competitor, the kangaroo rats (*Dipodomys* spp.) in a subset

of the experimental plots. In this study, we compared the breeding phenology of desert pocket

mice (*Chaetodipus penicillatus*) and Bailey's pocket mice (*Chaetodipus baileyi*) in *Dipodomys*

accessible (or Dipo+) and *Dipodomys* inaccessible (or Dipo-) plots to determine the effects of

interspecific competition on the phenology of natural populations. Specifically, we determined

periods of significant change in the monthly proportions of breeding male and female individuals

in both plot types and assessed the sex-specific responses to environmental conditions, and intra-

and inter-specific competition of both species in each plot type (Dipo+ and Dipo-).

METHODS

Study System

To assess the impacts of competition on reproductive phenology, we used data from a

long-term field experiment located in the Chihuahuan Desert near Portal, Arizona

(31°56'20.29"N 109° 4'47.44"W). Initiated in 1977, this study site consists of 24 experimental

plots (50m x 50 m) divided up among treatments that manipulate the desert rodent community.

Each plot is fenced, and we manipulate access to plots using different sized openings through the

fencing to generate some plots containing an unmanipulated rodent community (Dipo+), a

community where *Dipodomys* spp. (three species of behaviorally dominant competitors of which

Dipodomys merriami is the most common) are only occasionally present in low numbers (Dipo-

), and plots where all rodents are excluded (total rodent exclosure). In this study, we only used

data from the Dipo+ and Dipo- plots. Because of the spatial extent of the experiment (the plots

are contained in a 20-ha area), the rodents share the same habitat and experience the same

environment, allowing us to assess the impacts of competition without correlated changes in

abiotic conditions or habitat structure. All plots were sampled monthly, and the sex, species, and

reproductive condition of each individual were recorded affording us a monthly record of

reproductive activity for every species in the community. Treatments on some plots have

changed over time, so for this study we restricted our analysis to data collected from 1988 to

2014 (Appendix S1: Fig. 1), the longest period when the experimental treatments were consistent

for the most plots. More detailed description of the experimental plots is discussed elsewhere

(Ernest et al. 2016).

Rodent data

Desert rodents are an ideal model system for asking questions about the role of

competition on phenology because male and female reproductive investments are largely known

in rodents (Mysterud et al., 2008) and competition is a strong structuring process in desert rodent

communities (Brown and Munger 1985, Bledsoe and Ernest 2019). We focused our analyses on

two species, *C. baileyi* and *C. penicillatus*, because they are numerous on both target plot types

between 1988-2014 and because previous work has demonstrated strong competitive interactions

between each other and with *Dipodomys* spp. (Bledsoe and Ernest 2019). Because the timing of

phenology initiation relative to the dominant competitor is important in Clark and Rudolf's

experiment (2022), we also analyzed the temporal dynamics of *Dipodomys*' breeding for

comparison. We used the 'portalr' package (Christensen et al., 2019) to extract a monthly time

series by species (*Dipodomys* spp., *C. baileyi*, *C. penicillatus*), treatment (Dipo+, Dipo-), and sex

(male, female) of the abundance (i.e., all individuals regardless of reproductive status) and

counts of only reproductive individuals. Males were considered in reproductive state if they were

recorded as scrotal (descended), recently scrotal, or had minor signs of scrotal testes. Females

were considered reproductive if recorded as pregnant (after palpating the belly), and/or exhibited

either red and/or enlarged nipples, and/or those with swollen and/or plugged vaginas. To limit

the introduction of biases in our dataset through the inclusion of juveniles, we identified a

minimum threshold of the body mass for the breeding individuals (McLean and Guralnick, 2020)

for each species and sex (i.e., *C. baileyi* male= 16 g, *C. baileyi* female= 21 g, *C. penicillatus*

male= 10 g, *C. penicillatus* female= 12 g).

Environmental data

We used monthly data on NDVI (an index of plant greenness), mean temperature (°C),

cumulative precipitation (millimeters; sliding window calculating rainfall accumulations) that

fell during warm or cool months (calculated as the sum of precipitation on days when minimum

temperature is > or < 4 °C, a biologically relevant cutoff for the transition between warm and

cool rainy seasons which generate different plant assemblages), and biomass data of *C. baileyi*,

C. penicillatus, and all *Dipodomys* spp. We also examined lags of 0 and 1 month, and cumulative

values (i.e., precipitation) for environmental data because weather affects vegetation and

population dynamics dynamically through temporal processes (Ding et al., 2020; Wen et al.,

2019) and the importance of lags and seasonal accumulations have repeatedly been demonstrated

in explaining rodent population dynamics at this site (Thibault et al 2010). We used biomass data

of *C. baileyi*, *C. penicillatus*, and all *Dipodomys* spp. as proxies for competitive effects

(henceforth referred to as biotic factors).

 Data Analysis

We built generalized additive models (GAMs) to characterize the reproductive phenology

of male and female *C. baileyi* and *C. penicillatus* in different treatments, and to determine the

relationship between the proportion of breeding individuals with environmental (i.e., weather and

resource availability) and biotic (i.e., the intraspecific and interspecific competitive context)

factors. We also built GAMs to characterize the reproductive phenology of *Dipodomys* spp. to

compare any shifts in the breeding phenology of *Chaetodipus* spp. with that of the dominant

competitor. We pooled all data on *Dipodomys* spp. monitored at the site (*D. ordii*, *D. merriami*,

D. spectabilis) and focused only on their timing in Dipo+ plots, where we have adequate

observation data on the group. We used GAMs because they are appropriate for fitting data that

may exhibit non-linear trends and associations (Pedersen et al. 2019), a dynamic typical of

species' phenology that wax and wane through time.

Quantifying between-treatment differences in phenology- To assess the difference in the breeding

phenology of rodents on Dipo+ and Dipo- plots, we estimated a separate smoother for month,

and for each plot type, which were treated as ordered factors (Rose et al. 2012). By treating the

treatments as ordered factors (P_0= Dipo+, P_1= Dipo-), we were able to determine difference in

seasonal trends (i.e., smooths) between the Dipo + (i.e., treated as the reference level) and Dipo-

plots. We built species- and sex- specific models that had the general form:

153
$$Y_{i,j} = \alpha_0 + \alpha_{1j} P_{i,j} + f_j(T_1) + f_j(T_2) \quad \text{(Eqn. 1)}$$

where Y is the response variable (i.e., proportion of breeding individuals), α_0 is the model

intercept (i.e., the mean value of *Y* in the reference treatment, P_0 ; i.e., Dipo+ plot), α_1 is the

difference in the mean response between the Dipo+ (P_0) and Dipo- plot (P_1), $f()$ are smooth

functions of month (T_1), representing the breeding phenology of rodents, that are parameterized

using cubic cyclic regression splines (Pederson et al. 2019), and of year (T_2), representing long

term trends, that are parameterized using Gaussian process smooths (Wood 2017) for the *j*th

treatment.

Fitting the drivers of breeding phenology- To assess how the competitive environment may

impact model fits and the relative importance of drivers, we built species-, sex- and treatment-

specific models that included terms for environmental factors (terms for NDVI, warm and cool

precipitation, and mean monthly temperature with lag of zero and one), a proxy for intraspecific

competition (i.e., population biomass), and for interspecific competition (i.e., biomass of

competitors) as biotic (i.e., competitive context) factors. Pairwise correlation tests showed low

correlation among environmental covariates used to fit all models ($r < |0.5|$; see Appendix for

results). We built separate models for each treatment, species, and sex combination. These

models followed the general form:

$$Y \sim Binomial\ (n\,,p)\ \ (\text{Eqn. 2})$$

$$logit\ (p) = \alpha\ +\ \sum_{m=1}^{2} f_m(T_m)\ +\ \sum_{k=1}^{10}(\beta_k X_k)\ \ (\text{Eqn. 3})$$

In Eqn. 2, Y is the number of breeding individuals, which follows a binomial distribution with

the parameters n (number of independent trials, e.g., total number of males) and p (probability of

success, e.g., proportion of breeding males). In Eqn. 3, p is modeled as a function of the intercept

(α), the smooth function for month ($f_1(T_1)$), the smooth function for year ($f_2(T_2)$), and linear

terms for the effects of environmental and biotic predictors (β_k). Similar parameterizations of the

smooth terms done in Eqn. 1 were followed. We standardized all predictors by centering each

value on the mean and dividing them by two standard deviations (Gelman, 2008) to facilitate

interpretation of results. All models were implemented in a restricted maximum likelihood

(REML) framework using the 'mgcv' package ver 1.8-31 (Wood 2017) in R ver 4.1.1 (R Core

Team, 2021). Data (10.5281/zenodo.7301687) and code (10.5281/zenodo.8121230) used to

conduct data analyses are archived on Zenodo.

Model Interpretation

Reproductive phenology- From Eqn. 1, we determined significant treatment effects and

visualized differences between treatments by plotting the difference smooths, which reflect the

estimated differences in seasonality between Dipo+ and Dipo- plots. Then, we determined the

beginning and end of each breeding season for each species-, sex-, treatment- combination by

identifying periods of significant change in the proportion of breeding individuals for each

category using the output from models built using Eqn. 3. Here, the first derivatives of the

seasonal spline ($f_1(T_1)$) were estimated using finite differences to find the time periods when the

derivatives are significantly increasing or decreasing (i.e., does not contain zero in the

confidence interval). This and the 95% pointwise confidence interval were determined using the

derivatives function in the 'gratia' package (Simpson, 2023). We interpreted periods when the

rate of change in the seasonal spline ($f_1(T_1)$) were increasing as the onset of breeding season

(i.e., when more individuals are exhibiting reproductive characteristics), and decreasing rates of

change as the termination of breeding season (i.e., when fewer individuals are exhibiting

reproductive characteristics). We also performed exploratory analyses on the reproductive

phenology of *C. penicillatus* at different time periods of *C. baileyi* establishment in the site. *C.*

baileyi colonized the site around 1995, causing major shifts in resource use and the competitive

landscape, especially for *C. penicillatus* (Bledsoe and Ernest 2019, Diaz and Ernest 2022). To

assess the effect of the shift in the competitive landscape caused by this natural experiment, we

also built simpler models (without terms for competition) for all species-, sex-, and treatment-

specific datasets to determine the relative importance of these terms in explaining variation in

breeding proportions. Methods and results of these exploratory works are in the Appendix.

Association between covariates and proportion of breeders- We assessed whether individuals of

both pocket mice species exhibited different responses to drivers with and without *Dipodomys*

spp. present. We interpreted the coefficient values based on their signs (positive or negative) to

assess whether the effect of a given factor on the proportion of breeding individuals was positive

or negative. We also used the distribution curves of the estimated slopes from the models to

better understand how *Chaetodipus* spp. phenology relates to its dominant competitor.

RESULTS

Both *Chaetodipus* spp. exhibited changes in their breeding patterns when *Dipodomys* spp.

were removed from the rodent community. Changes included shifts in breeding onset (i.e., *C.*

baileyi females and to some extent, males), and prevalence of breeding individuals (*C.*

penicillatus females). Some relationships between covariates and the proportion of breeders did

not differ between *Dipodomys* accessible and *Dipodomys* inaccessible plots. However, the

responses of both *Chaetodipus* spp. to most covariates were broader than that of *Dipodomys* spp.

in Dipo+ plots and were narrower in Dipo- plots.

Female Bailey's pocket mouse (C. baileyi)

Female *C. baileyi* on Dipo+ plots went into breeding condition earlier than those on Dipo

- plots (s(difference)= 1.55, p= 0.05, deviance explained= 46.7%; January to March vs. February

to April, respectively; Fig.1, Appendix S1: Fig. S2), but declines in reproductive activity

occurred during similar timeframes on both Dipo+ and Dipo- plots (June-August vs. July-Sept,

respectively; Fig. 1). On both plots, the onset of the breeding season for female *C. baileyi*

occurred later than that for females of the dominant competitor, *Dipodomys* spp. (Fig. 1). Thus,

in the absence of *Dipodomys* spp., female *C. baileyi* delayed reproduction by ~1 month and in

the presence of *Dipodomys* spp., entered reproductive condition earlier and stayed in

reproductive condition longer.

The best fit models for female *C. baileyi* on Dipo+ and Dipo- inaccessible plots

overlapped in their important drivers, but also contained important differences. On Dipo+ plots,

female reproductive condition was positively associated with warm precipitation (lag 0; amount

of warm precipitation on corresponding month) and cool precipitation (lag 1; amount of cool

precipitation a month prior; variation explained= 66.4%; R^2= 0.57; Fig. 2) while on Dipo- plots,

warm precipitation (lag 0) was still significant but cool precipitation was not. Additional

important associations that emerged in the Dipo- model that did not appear in the Dipo+ model

included a negative association with NDVI (lag 0), a positive association with NDVI (lag 1), and

a positive association with *C. penicillatus* biomass (variation explained= 69.3%; R^2= 0.66; Fig.

3). Distributions of the covariate slope estimates (Fig. 4 and Appendix S1: Fig. S5) exhibited

broader distributions on Dipo+ plots for some environmental and biotic factors than either

Dipodomys spp. or *C. baileyi* on Dipo- plots.

Male Bailey's pocket mouse (C. baileyi)

The breeding timing of male *C. baileyi* did not significantly differ between Dipo+ and

Dipo- plots (*s*(difference= 1.35, *p*= 0.11, deviance explained= 20.7%; Appendix S1: Fig. S2).

However, despite a lack of a statistically significant treatment effect, the treatments exhibit a

slight divergence in their seasonal smooths that peak around June-July. The environmental

GAMs fit separately to Dipo+ and Dipo- plots were also suggestive of a difference in the

seasonal signal – with the Dipo+ model exhibiting no seasonal signal whereas a weak seasonal

signal was detected on Dipo- plots. Together these results may indicate that *C. baileyi* males

exhibit a weak phenological shift when *Dipodomys* spp. are present, but additional data and

analyses are required. The breeding onset of *C. baileyi* occurred about a month earlier than

female *C.baileyi* on the same plots (Fig.1).

Models show that male *C. baileyi* on Dipo+ plots were positively associated with *C.*

penicillatus biomass, and negatively associated with mean temperature (lag 0) and *C. baileyi*

biomass (variation explained= 21.8%, R2= 0.17; Fig. 2). Male *C. baileyi* on Dipo- plots did not

share a similar sensitivity to mean temperature but showed a similar relationship to the other

variables: a positive association with *C. penicillatus* biomass and a negative association with *C.*

baileyi biomass (variation explained= 44.3%; R^2= 0.42; Fig. 2). Moreover, male *C. baileyi* on

both Dipo+ and Dipo- plots exhibited similarly broad distributions in slope estimates to some

environmental variables (Fig. 4), and tighter ones to some biotic factors (Appendix S1: Fig. S5).

While the breadth of these distributions appears to vary little between treatments, these

distributions are much broader than those exhibited by *Dipodomys* spp.

Female desert pocket mouse (C. penicillatus)

The breeding timing of female *C. penicillatus* on Dipo+ and Dipo- plots did not

significantly vary (*s*(difference=0.00, *p*= 0.73; Appendix S1: Fig. S2). However, breeding

intensity was significantly higher on Dipo- than on Dipo+ plots (0.15, *p*= 0.02, deviance

explained= 26.6%; Fig. 3). Thus, in the absence of *Dipodomys* spp., female *C. penicillatus* are

more likely to go into reproductive mode than their counterparts on Dipo+ plots.

Female *C. penicillatus* on Dipo+ plots were positively associated with mean temperature

(lag 0), warm precipitation (lag 0), cool precipitation (lag 1), and *C. penicillatus* biomass, and

negatively associated with cool precipitation (lag 0) and *C. baileyi* biomass (variation explained=

55.5%; R^2= 0.53; Fig. 2). Female *C. penicillatus* on Dipo- plots responded similarly to mean

temperature (lag 0), warm precipitation (lag 0), and *C. baileyi* biomass on Dipo+ plots, but,

additionally, were positively associated with NDVI (lag 1; Fig. 2). Females on both plot types

exhibited relatively broad distributions in their slope estimates for environmental factors (Fig. 4).

Additionally, there was only marginal overlap between *Dipodomys* and *C. penicillatus* responses

to temperature and precipitation, which may suggest differences in how these environmental

cues are being viewed by both species.

Male desert pocket mouse (C. penicillatus)

Like females, the breeding timing of male *C. penicillatus* on Dipo+ and Dipo- plots did

not significantly vary (s(difference)= 0.00, p= 0.44, deviance explained= 52.1%; Appendix S1:

Fig. S2). Moreover, the proportion of breeding males on Dipo+ and Dipo- plots also did not

significantly differ. However, despite the lack of difference in seasonal signals and breeding

intensities, we noted subtle differences in the duration of breeding activity on both plots. The

estimated periods of increased reproductive activity were much longer on Dipo+ than Dipo- plots

(Fig. 3), which may suggest that male *C. penicillatus* prolong their breeding activities in the

presence of a dominant competitor. However, without a significant treatment effect, this result

should be treated cautiously.

The proportions of breeding male *C. penicillatus* on *Dipodomys* inaccessible plots were

negatively associated with warm precipitation (lag 1; variation explained= 52.1%; R2= 0.48; Fig.

3) but in general males exhibited little relationship with most drivers (Dipo+ model variation

explained= 63.6%; R^2= 0.68; Fig. 2). Like male *C. baileyi*, male *C. penicillatus* on both plots

also exhibited broader distributions of slope estimates to environmental variables and a tighter

response to biotic factors (Fig. 4 and Appendix S1: Fig S2). Finally, male *C. penicillatus*

responses to most factors did not differ from the responses of *Dipodomys* spp. (Fig. 4).

DISCUSSION

These results show that the presence or absence of a dominant competitor can alter the

reproductive phenology of small mammals by shifting the timing of the breeding season or

influencing the proportion of individuals entering reproductive condition. The earlier breeding

onset of *C. baileyi* females is consistent with Carter & Rudolf's results (2022), where

competitors that time their activities before or coincident with their main competitor fare better

in their reproductive activities. The precise mechanism for how *Dipodomys* is influencing the

timing of *C. baileyi* female phenology is unclear and we cannot distinguish between resource

depletion, behavioral interference, or other *Dipodomys* spp. mediated impacts, nor can we assess

whether shifts in phenology are due to individual plasticity (i.e., individuals changing the timing

of their breeding activities) or selection (i.e., the presence of *Dipodomys* spp. favors individuals

that would always breed early). While phenological shifts are one possible response to changes

in the competitive landscape, our results demonstrate that it is not necessarily the only way

species may respond. Female *C. penicillatus* exhibited no significant differences in their

phenology between treatments, but a higher proportion of females entered reproductive condition

in Dipo- plots. One possible explanation for *C. penicillatus*' lower breeding intensity in Dipo+

plots could be decreased body condition when *Dipodomys* spp. are present, with lower body

condition potentially preventing reproduction. We do see a small, but significant (t-test p= 0.005)

difference in mass between reproductive *C. penicillatus* females on Dipo+ (μ= 19.07 g) and

Dipo- plots (μ= 19.75 g). Iteroparous species- i.e., with multiple opportunities to reproduce- can

skip reproductive events or terminate in-process reproduction under extremely stressful

environments (Stearns 1992). Thus, *C. penicillatus* females may be opting to skip a particular

breeding season to wait for better conditions or be incapable of entering reproductive condition

due to their own nutritional state. While Carter & Rudolf's model (2022) does incorporate

intraspecific variation, it provides limited insights for the impacts of individual-level flexibility

in entering a phenological event. The results from both *C. baileyi* and *C. penicillatus* females

suggest that further assessment of individual-level plasticity - in addition to intraspecific trait

variation - may be useful.

Within-species phenological synchrony, which occurs when there is low variation in the

timing of a biological event for a given species, may arise from consistent responses among

individuals to external cues (i.e., shared responses to environmental factors at a given time

scale). Carter and Rudolf (2022) posit that the benefits of low and high phenological synchrony

are dependent on competitive context - specifically on the relative timing and synchrony in

competing species. However, given the nature of our data, it is not immediately clear how to

quantify the strength of intraspecific synchrony for our populations. If intraspecific synchrony is

driven by individuals exhibiting strong and consistent responses to drivers, then tight

distributions of environmental covariate estimates may indicate higher synchrony, and broader

distributions may imply more coordinated responses to the environment. Consistent with the

impression of higher synchrony, female *Dipodomys* spp. exhibit strong responses to

environmental variables with narrower distribution curves for parameter estimates, and thus, low

within-species variability in their environmental responses (Fig 4). In contrast, *Chaetodipus* spp.

exhibited a broader range of responses to external stressors. This larger variation in the

parameter estimates was generally consistent on both treatments and across sexes, suggesting

that high intraspecific variability in these species may not be driven solely by the direct impacts

of competition with *Dipodomys* spp. This may mean that these *Chaetodipus* spp. exploit

environmental conditions that are underutilized by *Dipodomys* spp. as an adaptive strategy.

While shifts in biotic context appear to drive changes in the breeding patterns of the

females for both *Chaetodipus* spp., it did not appear to be the case for males. The sex-related

differences in response to environment and competitive landscape are not necessarily surprising

since, in mammals, female reproduction is often limited by resource availability for gestation and

lactation and male reproduction by access to mates (Mysterud et al 2008). Because female

mammals are energetically and nutritionally providing for their own increased resource needs

and those of their offspring, females may be particularly sensitive to the effects of resource

depletion caused by interspecific competition. This may explain the observed phenological shifts

of female *C. baileyi* and increased breeding intensity of *C. penicillatus* in Dipo- plots. In

contrast, males for many mammal species - including those at our field site - are frequently

uninvolved in offspring rearing. This may explain the weak and lack of seasonal signal for

breeding detected in male *C. baileyi* and *C. penicillatus*, respectively. If male energetic costs are

primarily due to acquiring and defending females from other conspecifics (Ostfeld 1985), then

their phenology may be less sensitive to interspecific competition and more driven by female

reproductive changes and intraspecific competition for mates.

Our results also show that the effects of competition on phenology may be species

specific. While both *Chaetodipus* spp. have repeatedly demonstrated that *Dipodomys* have strong

negative impacts on their population sizes (Brown and Munger 1984, Ernest and Brown 2001,

Bledsoe and Ernest 2019), they showed different phenological responses to the removal of

Dipodomys. Clearly, more empirical and theoretical research is required to better understand how

and when competition will cause shifts in timing as opposed to prevalence. The differences in

model fits between treatments also suggest that continual reassessment of environmental drivers

of phenology may be warranted as biodiversity change alters the competitive environments many

species are experiencing. Future studies that can tease apart the direct and indirect impacts of

competition may be useful in developing a robust understanding of the effects of competition in

species' phenology. By recognizing that competitive landscapes can influence the responses of

breeding individuals, we gain a more nuanced understanding of one of the key biological

phenomena that can alter ecosystem dynamics.

ACKNOWLEDGMENTS

We thank the numerous volunteers, staff, and research assistants, who collected the Portal

Project data over the years. This work was supported by research grants awarded to SKME by

the US National Science Foundation (NSF DEB-1929730) and USDA National Institute of Food

and Agriculture, Hatch Project FLA-WEC-005983.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

REFERENCES

Audzijonyte, Asta, and Shane A. Richards. 2018. "The Energetic Cost of Reproduction and Its

Effect on Optimal Life-History Strategies." *The American Naturalist*, June.

https://doi.org/10.1086/698655.

Bledsoe, Ellen K., and S. K. Morgan Ernest. 2019. "Temporal Changes in Species Composition

Affect a Ubiquitous Species' Use of Habitat Patches." *Ecology* 100 (11): e02869.

https://doi.org/10.1002/ecy.2869.

Bradshaw, William E., and Christina M. Holzapfel. 2007. "Evolution of Animal

Photoperiodism." *Annual Review of Ecology, Evolution, and Systematics* 38 (1): 1–25.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110115.

Brown, James H., and James C. Munger. 1985. "Experimental Manipulation of a Desert Rodent

Community: Food Addition and Species Removal." *Ecology* 66 (5): 1545-1563.

Buss, Nicholas, Lindsey Swierk, and Jessica Hua. 2021. "Amphibian Breeding Phenology

Influences Offspring Size and Response to a Common Wetland Contaminant." *Frontiers in*

Zoology 18 (1): 31. https://doi.org/10.1186/s12983-021-00413-0.

Carter, Shannon K., and Volker H. W. Rudolf. 2022. "Exploring Conditions That Strengthen or

Weaken the Ecological and Evolutionary Consequences of Phenological Synchrony." *The*

American Naturalist 200 (5): E189–206. https://doi.org/10.1086/720899.

Christensen, Erica, Glenda Yenni, Hao Ye, Juniper Simonis, Ellen Bledsoe, Renata Diaz, Shawn

Taylor, Ethan White, and S. Ernest. 2019. "Portalr: An R Package for Summarizing and

Using the Portal Project Data." *Journal of Open Source Software* 4 (33): 1098.

https://doi.org/10.21105/joss.01098.

Cohen, Jeremy M., Marc J. Lajeunesse, and Jason R. Rohr. 2018. "A Global Synthesis of Animal

Phenological Responses to Climate Change." *Nature Climate Change* 8 (3): 224–28.

https://doi.org/10.1038/s41558-018-0067-3.

DeLucia, Evan H., Paul D. Nabity, Jorge A. Zavala, and May R. Berenbaum. 2012. "Climate

Change: Resetting Plant-Insect Interactions1." *Plant Physiology* 160 (4): 1677–85.

https://doi.org/10.1104/pp.112.204750.

Denny, Ellen G., Katharine L. Gerst, Abraham J. Miller-Rushing, Geraldine L. Tierney, Theresa

M. Crimmins, Carolyn A. F. Enquist, Patricia Guertin, et al. 2014. "Standardized Phenology

Monitoring Methods to Track Plant and Animal Activity for Science and Resource

Management Applications." *International Journal of Biometeorology* 58 (4): 591–601.

https://doi.org/10.1007/s00484-014-0789-5.

Dumandan, Patricia Kaye. 2022, November 16. patdumandan/ReproPhenology: v1.0.0. Zenodo.

Diaz, Renata M., and S. K. Morgan Ernest. 2022. "Maintenance of Community Function through

Compensation Breaks down over Time in a Desert Rodent Community." *Ecology* 103 (7):

e3709. https://doi.org/10.1002/ecy.3709.d

Ding, Yongxia, Zhi Li, and Shouzhang Peng. 2020. "Global analysis of time-lag and-

accumulation effects of climate on vegetation growth." *International Journal of Applied*

Earth Observation and Geoinformation 92:102179.

https://doi.org/10.1016/j.jag.2020.102179

Ernest, S. K. M., Glenda M. Yenni, Ginger Allington, Erica M. Christensen, Keith Geluso, Jacob

R. Goheen, Michele R. Schutzenhofer, et al. 2016. "Long□term Monitoring and

Experimental Manipulation of a Chihuahuan Desert Ecosystem near Portal, Arizona (1977–

2013) - Morgan Ernest - 2016 - Ecology - Wiley Online Library." 2016.

https://esajournals.onlinelibrary.wiley.com/doi/full/10.1890/15-2115.1.

Ernest, S. K. M., Glenda M. Yenni, Ginger Allington, Ellen K. Bledsoe, Erica M. Christensen,

Renata Diaz, Keith Geluso, Jacob R. Goheen, Qinfeng Guo, Edward Heske, Douglas Kelt,

Joan M. Meiners, Jim Munger, Carla Restrepo, Douglas A. Samson, Michele R.

Schutzenhofer, Marian Skupski, Sarah R. Supp, Katherine M. Thibault, Shawn D. Taylor,

Ethan P. White, Diane W. Davidson, James H. Brown, and Thomas J. Valone. 2022.

weecology/PortalData: 3.139.0. Zenodo.

Forchhammer, Mads C., Eric Post, and Nils Chr Stenseth. 1998. "Breeding Phenology and

Climate¨." *Nature* 391 (6662): 29–30. https://doi.org/10.1038/34070.

French, Alan R. 1977. "Circannual Rhythmicity and Entrainment of Surface Activity in the

Hibernator, Perognathus Longimembris." *Journal of Mammalogy* 58 (1): 37–43.

https://doi.org/10.2307/1379725.

Gelman, Andrew. 2008. "Scaling regression inputs by dividing by two standard

deviations." *Statistics in medicine* 27(15): 2865-2873.

Glick, Patricia, Bruce A Stein, Naomi A Edelson, and National Wildlife Federation. 2011.

Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessment.

Washington, D.C.: National Wildlife Federation.

Gordo, Oscar, and Juan José Sanz. 2005. "Phenology and Climate Change: A Long-Term Study

in a Mediterranean Locality." *Oecologia* 146 (3): 484–95. https://doi.org/10.1007/s00442-

005-0240-z.

Helm, Barbara, Rachel Ben-Shlomo, Michael J. Sheriff, Roelof A. Hut, Russell Foster, Brian M.

Barnes, and Davide Dominoni. 2013. "Annual Rhythms That Underlie Phenology:

Biological Time-Keeping Meets Environmental Change." *Proceedings of the Royal Society*

B: Biological Sciences 280 (1765): 20130016. https://doi.org/10.1098/rspb.2013.0016.

Kelt, Douglas, Mark Taper, and Peter L. Meserve. 1995. "Assessing the Impact of Competition

on Community Assembly: A Case Study Using Small Mammals - Kelt - 1995 - Ecology -

Wiley Online Library." 1995.

https://esajournals.onlinelibrary.wiley.com/doi/abs/10.2307/1940935.

Kenagy, G. J., and George A. Bartholomew. 1981. "Effects of Day Length, Temperature, and

Green Food on Testicular Development in a Desert Pocket Mouse Perognathus Formosus."

Physiological Zoology 54 (1): 62–73.

McLean, Bryan S., and Robert P. Guralnick. 2021. "Digital Biodiversity Data Sets Reveal

Breeding Phenology and Its Drivers in a Widespread North American Mammal." *Ecology*

102 (3): e03258. https://doi.org/10.1002/ecy.3258.

Mysterud, Atle, Christophe Bonenfant, Leif Egil Loe, Rolf Langvatn, Nigel G. Yoccoz, and Nils

C. Stenseth. 2008. "The Timing of Male Reproductive Effort Relative to Female Ovulation

in a Capital Breeder." *Journal of Animal Ecology* 77 (3): 469–77.

https://doi.org/10.1111/j.1365-2656.2008.01365.x.

Ostfeld, Richard S. 1985. "Limiting Resources and Territoriality in Microtine Rodents." *The*

American Naturalist 126 (1): 1–15.

Pearce-Higgins, J. W., and D. W. Yalden. 2004. "Habitat Selection, Diet, Arthropod Availability

and Growth of a Moorland Wader: The Ecology of European Golden Plover Pluvialis

Apricaria Chicks." *Ibis* 146 (2): 335–46. https://doi.org/10.1111/j.1474-919X.2004.00278.x.

Pedersen, Eric J., David L. Miller, Gavin L. Simpson, and Noam Ross. 2019. "Hierarchical

generalized additive models in ecology: an introduction with mgcv." *PeerJ* 7: e6876.

https://doi.org/10.7717/peerj.6876

R Core Team. 2021. "R: A language and environment for statistical computing". *R Foundation*

for Statistical Computing. Vienna, Austria. http://www.R-project.org

Reed, Thomas E., Stephanie Jenouvrier, and Marcel E. Visser. 2013. "Phenological Mismatch

Strongly Affects Individual Fitness but Not Population Demography in a Woodland

Passerine." *Journal of Animal Ecology* 82 (1): 131–44. https://doi.org/10.1111/j.1365-

2656.2012.02020.x.

Rose, Neil L., Handong Yang, Simon D. Turner, and Gavin L. Simpson. 2012. "An assessment

of the mechanisms for the transfer of lead and mercury from atmospherically contaminated

organic soils to lake sediments with particular reference to Scotland, UK." *Geochimica et*

Cosmochimica Acta 82: 113-135.

Roslin, Tomas, Laura Antão, Maria Hällfors, Evgeniy Meyke, Coong Lo, Gleb Tikhonov, Maria

del Mar Delgado, et al. 2021. "Phenological Shifts of Abiotic Events, Producers and

Consumers across a Continent." *Nature Climate Change* 11 (3): 241–48.

https://doi.org/10.1038/s41558-020-00967-7.

Simpson, Gavin L. 2018. "Modelling Palaeoecological Time Series Using Generalised Additive

Models." *Frontiers in Ecology and Evolution* 6.

https://www.frontiersin.org/article/10.3389/fevo.2018.00149.

Simpson, Gavin L. 2023. Gratia: Graceful ggplot-Based Graphics and Other Function for GAMs

Fitted using mgcv. R package version 0.8.1. https://gavinsimpson.github.io/gratia/.

Stearns, Stephen. 1992. The Evolution of Life Histories. Oxford University Press.

Thackeray, Stephen J., Timothy H. Sparks, Morten Frederiksen, Sarah Burthe, Philip J. Bacon,

James R. Bell, Marc S. Botham, et al. 2010. "Trophic Level Asynchrony in Rates of

Phenological Change for Marine, Freshwater and Terrestrial Environments." *Global*

Change Biology 16 (12): 3304–13. https://doi.org/10.1111/j.1365-2486.2010.02165.x.

Thibault, Katherine M., S. K. Morgan Ernest, Ethan P. White, James H. Brown, and Jacob R.

Goheen. 2010. "Long-Term Insights into the Influence of Precipitation on Community

Dynamics in Desert Rodents." *Journal of Mammalogy* 91(4): 787–97.

https://doi.org/10.1644/09-MAMM-S-142.1.

Visser, Marcel E, and Christiaan Both. 2005. "Shifts in Phenology Due to Global Climate

Change: The Need for a Yardstick." *Proceedings of the Royal Society B: Biological*

Sciences 272 (1581): 2561–69. https://doi.org/10.1098/rspb.2005.3356.

Walther, Gian-Reto, Eric Post, Peter Convey, Annette Menzel, Camille Parmesan, Trevor J. C.

Beebee, Jean-Marc Fromentin, Ove Hoegh-Guldberg, and Franz Bairlein. 2002. "Ecological

Responses to Recent Climate Change." *Nature* 416 (6879): 389–95.

https://doi.org/10.1038/416389a.

Wann, Gregory T., Cameron L. Aldridge, Amy E. Seglund, Sara J. Oyler-McCance, Boris C.

Kondratieff, and Clait E. Braun. 2019. "Mismatches between Breeding Phenology and

Resource Abundance of Resident Alpine Ptarmigan Negatively Affect Chick Survival."

Ecology and Evolution 9 (12): 7200–7212. https://doi.org/10.1002/ece3.5290.

Wen, Youyue, Xiaoping Liu, Qinchuan Xin, Jin Wu, Xiaocong Xu, Fengsong Pei, Xia Li et al.

2019. "Cumulative effects of climatic factors on terrestrial vegetation growth." *Journal of*

Geophysical Research: Biogeosciences 124 (4): 789-806.

https://doi.org/10.1029/2018JG004751

Wolkovich, Elizabeth M., Benjamin I. Cook, and T. Jonathan Davies. 2014. "Progress towards

an Interdisciplinary Science of Plant Phenology: Building Predictions across Space, Time

and Species Diversity." *New Phytologist* 201 (4): 1156–62.

https://doi.org/10.1111/nph.12599.

Wood, Simon. 2017. "Generalized Additive Models: An Introduction with R (2nd edition).

Chapman and Hall/CRC.

FIGURE LEGENDS

Figure 1. Reproductive phenology of female (left panel) and male (right panel) *Dipodomys* spp.

    in *Dipodomys* accessible plots (top panel), and *C. baileyi* in *Dipodomys* accessible

    (middle panel) and inaccessible (bottom panel) plots in an experimental field site near

    Portal, Arizona. Plots show estimated smooths of the month term from species-, sex-, and

    treatment-specific generalized additive models (GAMs). Blue and red shading indicates

    periods of significant increase and decrease in breeding proportions, respectively. In

    cases where seasonal patterns or periods of significant increase or decrease could not be

    estimated from the model, raw data on proportion of breeding individuals were plotted

    (male *Dipodomys* spp. and *C. baileyi*;).

Figure 2. Association between select environmental and biotic factors with the proportion of

    reproductive female (left panel) and male (right panel) *C. baileyi* (top panel) and desert

    pocket mouse (bottom panel) in *Dipodomys* accessible and inaccessible plots.

Figure 3. Reproductive phenology of female (left panel) and male (right panel) *Dipodomys* spp.

    in *Dipodomys* accessible plots (top panel), and *C. penicillatus* in *Dipodomys* accessible

    (middle panel) and inaccessible (bottom panel) plots in an experimental field site near

    Portal, Arizona. Details on the configuration of these plots are like Figure 1.

Figure 4. Distribution curves of the abiotic covariate slope estimates indicating the response of

female (left panel) and male (right panel) *C. baileyi* (top panel) and *C. penicillatus*

(bottom panel) in *Dipodomys* acessible and inaccessible plots, compared to *Dipodomys*

spp. estimates. Curves indicate the normal distribution with mean equal to the point

estimate, and standard deviation equal to the standard error of the coefficients.

FIGURES

539
540   Figure 1.
541

543
544

Figure 2.

547
548   Figure 3.

550
551
552
553   Figure 4.