Manuscript type: Article

Title: Shifts in competitive structures can drive variation in species' phenology

Authors: Patricia Kaye T. Dumandan[1], Glenda M. Yenni[2], and S. K. Morgan Ernest[2]

Institutional Affiliations:

1. School of Natural Resources and Environment, University of Florida, Gainesville, FL, USA

2. Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL, USA

Corresponding author: Patricia Kaye T. Dumandan (pdumandan@ufl.edu)

ABSTRACT

Characterizing patterns and drivers of species' responses to various environmental

contexts is useful in determining their adaptive capacity to the consequences of ongoing global

environmental change. For many species, a well-documented response to anthropogenic climate

change is a shift in the timing of their key life-history events or phenology. Often, these

phenological shifts are associated with changes in abiotic factors used as proxies for resource

availability or other suitable conditions. However, less is known about how species' interactions

that can lead to resource limitation influence phenological responses. Using long-term

experimental data on small mammals monitored in the southwestern United States, we show that

altered competitive landscapes can drive shifts in breeding timing and duration, and that, relative

to a dominant competitor, other species exhibit less specific responses to environmental factors.

These results suggest that plasticity of phenological responses, that is often described in the

context of annual variation in abiotic factors, can occur in response to biotic context as well.

Additionally, in response to competitive pressures, species may employ adaptive strategies, such

as low phenological synchrony, to compensate for the adverse demographic impacts of

mismatched resource and species' phenology. Variation in phenological responses under

different biotic conditions shown here further demonstrate that shifting biotic interactions are

rampantly occurring in many natural systems and a more nuanced understanding of its impacts

on population dynamics is useful to better understand and predict biodiversity patterns in a

changing world.

INTRODUCTION

Characterizing species' responses to various environmental conditions is useful in

determining their sensitivity and adaptive capacity to the consequences of ongoing global

environmental change (Glick et al., 2011). Among the known responses to climate change, shifts

in phenology or the timing of key life-history events of organisms, are the most well-documented

(Cohen et al., 2018; Roslin et al., 2021; Walther et al., 2002). Often, the timing of important

phenological events (e.g., reproduction) match with periods of high resource availability to

support high fitness and survival costs (Bradshaw & Holzapfel, 2007; Buss et al., 2021;

Forchhammer et al., 1998; Helm et al., 2013; Reed et al., 2013; Walther et al., 2002). Fitness

costs of reproduction are often attributed to the increased energetic demands of maintaining and

developing reproductive organs and behaviors (Visser & Both, 2005) and survival costs are

incurred from increased predation risk (Audzijonyte & Richards, 2018; Kelt et al., 1995). To

avoid unfavorable conditions during the reproductive period, species may use abiotic (i.e.,

temperature, day length, precipitation) and biotic cues (i.e., flowering, prey availability) to

regulate and shape reproductive phenology (Denny et al., 2014; Kenagy & Bartholomew, 1981;

Wolkovich et al., 2014).

Timing reproductive activities to overlap with periods of high resource availability is

important for consumer species. Thus, many studies assess phenological drivers related to either

reliable environmental cues of resource availability or biotic cues related directly to the

emergence or abundance of prey (i.e., resource) species (DeLucia et al., 2012; Gordo & Sanz,

2005; Pearce-Higgins & Yalden, 2004). In fact, decoupled resource and consumer phenology

often has negative implications for successful production and survival of offspring (Thackeray et

al., 2010; Wann et al., 2019). However, resource availability is determined by more than just the

intrinsic population dynamics of predator and prey. Competition from other species can deplete

important resources and create resource limitation even when resource supply is initially high

and can negatively impact individuals and populations. In fact, demographic rates were found to

be lower for those that initiate their biological activities (i.e., breeding) after their competitors

(Carter & Rudolf 2022). Thus, plasticity of species' phenology under different biotic contexts

could have important implications for interpreting observed phenological changes as ecosystems

increasingly undergo turnover in their component species. However, it is unclear whether

plasticity exists to allow shifts in a species' phenology when a competitor is absent.

While competition has the potential to impact reproductive timing and investment, it may

not similarly impact all individuals within a species, especially for species where variation in the

degree and type of offspring investment depends on the sex of the parent (Mysterud et al., 2008).

In general, female animals play a primary role in provisioning offspring from conception until

birth or hatching. In mammals, this period of primary responsibility for offspring provisioning is

lengthened through lactation. Because female mammals are energetically and nutritionally

providing for their own increased resource needs and those of their offspring, females may be

particularly sensitive to the effects of resource depletion caused by interspecific competition. In

contrast, males for many mammal species are frequently uninvolved in offspring rearing. Their

enhanced energetic demands during this period are due to acquiring and defending females from

other conspecifics (Ostfeld 1985), and thus, may be largely influenced by intraspecific

competition for mates. These differences in reproductive constraints and pressures may also be

reflected in their use and sensitivity to environmental cues. With logical support for sex-specific

phenological responses, it is therefore important that such sources of variation are considered

when assessing the effects of competition on reproductive phenology. However, data that allow

sex-specific assessment of competitive effects on phenology in dynamic environments are often

limited.

Long-term field experiments, where sex-specific differences in reproductive investments

of species are largely known and competitive networks are explicitly manipulated, are well-

suited for investigating the role of competition in driving reproductive phenology (Ernest et al.,

2016; Kelt et al., 1995, Mysterud et al., 2008; Ostfeld, 1985). In long-term field experiments

where dominant competitors are removed, the effect of their loss on the reproductive phenology

of other species in the system can be evaluated by comparing the reproductive parameters of a

focal species in control and treatment set-ups during different temporal contexts. Here, we used

long-term data on small mammals monitored in an experimental field site near Portal, Arizona to

assess the effects of competition on reproductive phenology. This long-term experiment (1977 to

present) in the southwestern USA generates fine-scale (monthly) data on desert rodents under

different biotic contexts. The experiment involves the removal of a dominant competitor, the

kangaroo rats (*Dipodomys* spp.), in a subset of the experimental plots (hereafter referred to as

exclosure plots). In this study, we compared the breeding phenology of desert pocket mice

(*Chaetodipus penicillatus*) and Bailey's pocket mice (*C. baileyi*) in control and exclosure plots.

Specifically, we determined periods of significant change in the monthly proportions of breeding

male and female individuals in both plot types. We also assessed the sex-specific responses to

abiotic conditions, and intra- and inter-specific competition of both species in each plot type

(control and exclosure). In doing so, we can gain a more robust understanding of phenological

responses to environmental factors, which may vary in future landscape and climate change

scenarios.

METHODS

 Data Curation and Manipulation

We used time-series data on rodents and environmental variables collected at an

experimental field site near Portal, Arizona (31°56'20.29"N 109° 4'47.44"W) to determine the

effects of biotic context (presence of kangaroo rats) and abiotic factors (temperature, NDVI,

precipitation) on the reproductive phenology of two pocket mice species (*C. penicillatus* and *C.*

baileyi). We used rodents as model species in this study because male and female reproductive

investments of rodents are largely known (Mysterud et al., 2008). Moreover, desert rodents are

especially ideal for answering questions about phenological plasticity given the resource-limited

environment they inhabit, which may drive variation in their responses to both biotic and abiotic

factors.

In this study, we used data collected from 1988 to 2014, because this is the longest period

when the experimental treatments were consistent for the most plots. We used data obtained

from control plots, where all rodents including kangaroo rats have access, and exclosure plots,

where all other rodents were allowed access through smaller holes in the fences surrounding the

plots. More detailed description of the experimental plots is discussed elsewhere (Ernest et al.

2016). All plots were sampled monthly, and the sex, species, and reproductive condition of

individuals were recorded.

Rodent data- We assembled a dataset on the observed total abundance and counts of

reproductive *C. baileyi* and *C. penicillatus* in Portal, Arizona using the 'portalr' package

(Christensen et al., 2019). We created subsets of the data for each type of plot (control and

exclosure) and sex (male and female). In calculating the proportion of reproductive individuals,

we considered those that had testes described to be scrotal (descended), recently scrotal, or had

minor signs of scrotal testes as reproductive males. We considered individuals that were

described to be pregnant (after palpating the belly), and/or those with either red and/or enlarged

nipples, and/or those with swollen and/or plugged vaginas as reproductive females. To limit the

introduction of biases in our dataset through the inclusion of juveniles, we identified a minimum

threshold of the body mass for the breeding individuals (McLean and Guralnick, 2020) for each

species and sex (i.e., *C. baileyi* male= 16 g, *C. baileyi* female= 21 g, *C. penicillatus* male= 10 g,

C. penicillatus female= 12 g). We filtered out individuals that had a body mass below the

threshold for each group.

Environmental data- We used monthly data on NDVI (an index of plant greenness),

mean temperature (°C), cumulative precipitation (millimeters) that fell during warm or cool

months (calculated as the sum of precipitation on days > or < 4 °C), and biomass data of *C.*

baileyi, *C. penicillatus*, and all species of *Dipodomys*. We manipulated data to obtain lagged

values (lag =1 month) of all abiotic variables. We explored the effects of time lagged variables

and time accumulation effects of precipitation because climate affects vegetation dynamically

through temporal processes (Ding et al., 2020; Wen et al., 2019). We assumed that the

cumulative effect of precipitation is three months (90 days) given the aridity of the system (Ma et

al., 2021). We used biomass data of *C. baileyi*, *C. penicillatus*, and all species of *Dipodomys* as

proxies for competitive effects.

Data Analysis

We built generalized additive models (GAMs) to characterize the reproductive phenology

of male and female *C. baileyi* and *C. penicillatus* in different biotic contexts, and to determine

the relationship between the proportion of breeding individuals with abiotic and biotic factors.

134 We also built GAMs to characterize the reproductive phenology of kangaroo rats so that we

135 could compare any shifts in the breeding phenology of *Chaetodipus* spp. with that of the

136 dominant competitor. We pooled all data on *Dipodomys* spp. monitored in the site (*D. ordii*, *D.*

137 *merriami*, *D. spectabilis*) and focused only on their timing in control plots, where we have

138 adequate observation data on the group.

139 We used GAMs because they are appropriate for fitting data that may exhibit non-linear

140 trends and associations (Pedersen et al. 2019), a dynamic typical of species' phenology that wax

141 and wane through time. Here, we fitted species-, sex-, and treatment-specific models to different

142 subsets of the dataset. For example, we built a separate model for data on male *C. penicillatus* in

143 control plots. We built models that included terms for abiotic factors (terms for NDVI, warm and

144 cool precipitation, and mean monthly temperature with lag of zero and one), a proxy for

145 intraspecific competition (i.e., population biomass), and for interspecific competition (i.e.,

146 biomass of competitors). We built separate models for all combinations of data. These models

147 followed the general form:

148$$Y \sim Binomial\ (n\,,p)\ \text{(Eqn. 2)}$$

149$$logit\ (p) = \alpha + \sum_{m=1}^{2} f_m(T_m) + \sum_{k=1}^{10} (\beta_k X_k)\ \text{(Eqn. 3)}$$

150 In Eqn. 2, Y is the number of breeding individuals, which follows a binomial distribution with

151 the parameters n (number of independent trials, e.g., total number of males) and p (probability of

152 success, e.g., proportion of breeding males). In Eqn. 3, p is modeled as a function of the intercept

153 (α), the smooth function for month ($f_1(T_1)$), which was parameterized using cubic cyclic

154 regression splines (Pederson et al. 2019), the smooth function for year ($f_2(T_2)$), which was

155 parameterized using Gaussian process smooths (Wood 2006), and linear terms for the effects of

abiotic and biotic predictors (β_k). We used Pearson's correlation coefficient (r) to test for

collinearity among the predictors included in the models ($|r| < 0.5$). We standardized all

predictors by centering each value on the mean and dividing them by two standard deviations

(Gelman, 2008) to facilitate interpretation of results. All models were implemented in a restricted

maximum likelihood (REML) framework using the 'mgcv' package ver 1.8-31 (Wood and

Wood 2015) in R ver 4.1.1 (R Core Team, 2021). Data (10.5281/zenodo.7301687) and code

(10.5281/zenodo.7327503) used to conduct data analyses are archived in Zenodo.

Model Interpretation

Reproductive phenology- We characterized rodent breeding phenology by identifying

periods of significant change in the proportion of breeding individuals (i.e., when the rates of

change were non-zero). Here, we approximated the derivatives of the spline function for month

through finite differences to determine the direction of the function (i.e., increasing or

decreasing; Simpson, 2018). In this case, we estimated the rate of change in the fitted smoother

(i.e., month) to identify periods when the proportion of breeding individuals increased or

decreased. We interpreted periods when the rate of change in the fitted smoother for month were

increasing as the onset of breeding season (i.e., when more individuals are exhibiting

reproductive characteristics), and when the rate of change were decreasing as termination of

breeding season (i.e., when fewer individuals are exhibiting reproductive characteristics). We

also performed exploratory work on the reproductive phenology of *C. penicillatus* at different

time periods of *C. baileyi* establishment in the site. *C. baileyi* colonized the site in 1995, causing

major shifts in resource use and the competitive landscape, especially for *C. penicillatus*

(Bledsoe and Ernest 2019, Diaz and Ernest 2022). To assess the effect of the shift in the

competitive landscape caused by this natural experiment, we also built simpler models (without

terms for competition) for all species-, sex-, and treatment- specific datasets to determine the

relative importance of these terms in explaining variation in breeding proportions. Details of the

methods and results of these exploratory works are found in the Appendix.

Association between covariates and proportion of breeders- We examined relationships

between covariates and the proportion of breeders in two ways. First, we assessed whether

individuals of both pocket mice species exhibited different environmental responses under

different biotic contexts. We interpreted the coefficient values based on their signs (positive or

negative) to assess whether the effect of a given factor on the proportion of breeding individuals

was positive or negative. Then, we compared species-specific responses in each plot to the

response of the kangaroo rats to infer species-specific strategies for competition relative to the

dominant competitor in the system. We used the distribution curves of the slopes estimated from

models of both pocket mice species and related it to the distribution curves of the slopes from

models of kangaroo rats (i.e., *Dipodomys* spp. and *C. baileyi* in control and exclosure plots,

Dipodomys spp. and *C. penicillatus* in control and exclosure plots).

RESULTS

Both species of pocket mouse exhibited changes in their phenology when kangaroo rats

are removed from the rodent community (Figs. 1 and 2). Changes included shifts in onset and

duration, depending on species and sex (details below; Figs. 1 and 2). Some relationships

between covariates and the proportion of breeders did not differ between control and kangaroo

rat exclosure plots (Fig. 3), however, the responses of both pocket mice species to most

covariates were relatively broader than that of kangaroo rats in control plots (Fig 4a and 4b) and

were narrower in exclosure plots (Fig. 4c and 4d).

Reproductive phenology of pocket mice under different biotic conditions

Female Bailey's pocket mouse (C. baileyi)- Females on control plots went into breeding

condition earlier than those on exclosure plots (January to March vs. February to April,

respectively; Fig 1c, e), but declines in reproductive activity occurred during similar timeframes

on both exclosures and controls (June-August vs. July-Sept, respectively; Figure 1c, e). On both

plots, the onset of the breeding season for female *C. baileyi* occurred later than that for females

of the dominant competitor, kangaroo rats (Fig. 1a, 1c, 1e). Thus, in the absence of kangaroo

rats, female *C. baileyi* delayed reproduction by ~1 month and in the presence of kangaroo rats,

entered reproductive condition earlier and stayed in reproductive condition longer.

The best fit models for female *C. baileyi* on controls and exclosures overlapped in their

important drivers, but also contained important differences. On control plots, female

reproductive condition was positively associated with warm precipitation (lag 0) and cool

precipitation (lag 1; variation explained= 66.4%; $R^2 = 0.57$; Fig. 2a). The best fit model for *C.*

baileyi on exclosures also found a positive role for warm precipitation (lag 0), but not for cool

precipitation. Additional important associations that emerged in the exclosure model that did not

appear in the control model included a negative association with NDVI (lag 0), a positive

association with NDVI (lag 1), and desert pocket mouse biomass (variation explained= 69.3%;

$R^2 = 0.66$; Fig. 2a). Moreover, female *C. baileyi* in control plots exhibited a wider range of

responses to some abiotic and biotic factors than kangaroo rats, and those in exclosure plots (Fig.

4a and Appendix S1 Fig. S2).

Male Bailey's pocket mouse (*C. baileyi*)- The main difference between males on controls

and exclosures was a lack of a seasonal signal in the proportion of reproductive males on

controls, in contrast to a weak seasonal signal on the exclosure. This suggests that in the presence

of kangaroo rats, male *C. baileyi* do not follow a particular schedule when they are more likely to

be reproductively active. Without kangaroo rats present, male *C. baileyi* exhibited weak seasonal

reproductive behavior, with their breeding onset occurring about a month earlier than female

C.baileyi on the same plots (Fig.1e).

Models show that male *C. baileyi* on control plots were positively associated with *C.*

penicillatus biomass, and negatively associated with mean temperature (lag 0) and *C. baileyi*

biomass (variation explained= 21.8%, $R2 = 0.17$; Fig. 2b). Male *C. baileyi* on exclosure plots did

not share a similar sensitivity to mean temperature but showed a similar relationship to the other

variables: a positive association with *C. penicillatus* biomass and a negative association with *C.*

baileyi biomass (variation explained= 44.3%; $R2 = 0.42$; Fig. 2b). Moreover, male *C. baileyi* in

both control and exclosure plots exhibited similar generalized responses to some abiotic factors

(Fig. 4b), and more specialized responses to some biotic factors (Appendix S1: Figure S2).

However, in general, they exhibited more generalized responses to both abiotic and biotic factors

compared to kangaroo rats.

Female desert pocket mouse (C. penicillatus)- Females on control plots delayed their

reproductive onset by nearly a month in comparison to females on exclosure plots (November to

January (Fig. 3c) on control plots while those on exclosure plots were increasing around October

to December (Fig. 3e)). Many breeding females left breeding condition from May to August,

which was the same regardless of plot type. Thus, in the absence of kangaroo rats, female *C.*

penicillatus entered reproductive condition earlier and stayed in reproductive condition longer

than females observed in exclosure plots (Fig. 3c and 3e).

Female *C. penicillatus* on control plots were positively associated with mean temperature

(lag 0), warm precipitation (0), cool precipitation (lag 1), and *C. penicillatus* biomass, and

negatively associated with cool precipitation (lag 0) and *C. baileyi* biomass (variation explained=

55.5%; $R2 = 0.53$; Fig. 2c). Female *C. penicillatus* on exclosure plots responded similarly to

mean temperature (lag 0), warm precipitation (lag 0), and Bailey's pocket mouse biomass on

control plots, but, additionally, was positively associated with NDVI (lag 1; Fig. 2c). Female *C.*

penicillatus in both controls and exclosures exhibited relatively generalized responses to abiotic

factors (Fig. 4c). Additionally, there is marginal overlap between kangaroo rat and *C.*

penicillatus responses to temperature and precipitation, which may suggest differences in how

these environmental cues are being viewed by both species. In contrast, their responses to biotic

factors were generally similar with that of their dominant competitor, kangaroo rats (Appendix

S1: Fig S2).

Male desert pocket mouse (C. penicillatus)- For male *C. penicillatus*, the end to the

breeding season occurred sooner on exclosure plots than on controls. The proportions of

breeding male *C. penicillatus* began increasing in January on both controls and exclosures but

finished decreasing on control plots in September in comparison to August on exclosure plots

(Fig. 3d). Thus, the presence of kangaroo rats resulted in a longer breeding season for male *C.*

penicillatus.

Male *C. penicillatus* on both control (variation explained= 63.6%; $R2 = 0.68$) and

exclosure (variation explained= 52.1%; $R2 = 0.48$) plots were generally not associated with any

predictors (Fig. 2d). However, the proportions of breeding male *C. penicillatus* on exclosure

plots were negatively associated with warm precipitation (lag 1; Fig. 2d). Like male *C. baileyi*,

male *C. penicillatus* in both control and exclosure plots also exhibited a relatively more

generalized response to abiotic and less generalized response to biotic factors (Fig. 4d and

Appendix S1: Fig S2). Finally, male *C. penicillatus* responses to most factors did not differ from

the responses of their dominant competitors, *Dipodomys* spp.

DISCUSSION

Our results show that small mammals can shift the timing of their breeding phenology in

response to the presence or absence of an important competitor. Differences in breeding

phenology were observed between control and kangaroo rat exclosure plots for females of both

species and - to some extent - males. Shifts in breeding phenology occurred in a variety of ways -

changes in the timing of initiation and/or cessation of breeding season, as well as changes in the

duration and degree of seasonality (or synchrony) exhibited. Our results add to our understanding

of animal phenology by providing evidence that observed shifts in timing can be driven not only

by shifts in abiotic cues, but also in a species' competitive environment.

While we detected shifts in breeding phenology in response to our experimental

manipulation of kangaroo rats, it is not immediately clear how changes in the competitive milieu

are driving changes in the breeding phenology of the two pocket mice. Models examining

interactions between competition and phenology timing show that competitors that time their

activities before or coincident with their main competitor exhibit higher survival rates whereas

those that come afterwards tend to suffer from poorer condition and outcomes (Carter and Rudolf

2022). A similar process might explain why *C. baileyi* on controls enter reproductive condition

sooner than those on kangaroo rat exclosures since reproducing later may mean lower resource

availability when kangaroo rats are present. We also see that *C. penicillatus* females initiate their

reproductive season before either *C. baileyi* or the kangaroo rats initiate theirs, regardless of

whether kangaroo rats are present or not. Initiating their reproductive season before these two

dominant competitors is also consistent with Carter and Rudolf's (2022) simulation. However,

the relatively delayed breeding onset of female *C. penicillatus* in controls, where both kangaroo

rats and Bailey's pocket mouse are present (Fig 2) is not consistent with Carter & Rudolf (2022),

as there should be little advantage to delaying reproduction in a more competitive environment,

where you would otherwise avoid competition by initiating reproduction early.

The counterintuitive response of *C. penicillatus* may emerge from a unique aspect of its

biology. Unlike the other two species, *C. penicillatus* is known to enter intermittent torpor during

the winter depending upon energetic state and ambient temperature (French 1977). In this

system, we note slight increases in the proportion of breeding female *C. penicillatus* in exclosure

plots during winter months, suggesting that many of the individuals that do not enter torpor are

in good enough condition to initiate reproductive condition. A possible explanation for *C.*

penicillatus' delay in breeding in the controls could be related to decreased body condition when

kangaroo rats are present - diverting more individuals into torpor and fewer into reproductive

pathways. While Carter & Rudolf (2022) incorporate intraspecific variation into their model, it

provides limited implications for temporal variation in the condition of individuals of species

that are capable of multiple breeding seasons. Our results suggest that expanding Carter &

Rudolf (2022) to incorporate individual decision-making on if and when to initiate reproductive

events based on energetic condition could provide important insights into how species may shift

phenology in response to changes in their competitive environment.

Males of both pocket mice species also exhibited changes in reproductive timing in

response to altered competitive environments. Male *C. penicillatus* in control plots were going

into and out of the breeding season longer than those on exclosures while *C. baileyi* males

exhibited a weak seasonal signal when kangaroo rats were absent. Decreased phenological

synchrony, in the form of longer periods of increase and decrease for *C. penicillatus* (Fig 3) and

a lack of seasonal signal for *C. baileyi* (Fig. 1) in control plots, may have been a competitive

strategy of these pocket mice to offset costs of delayed breeding relative to kangaroo rats (Carter

and Rudolf 2022). However, these changes may also be consistent with previous knowledge on

the advantages obtained by males of various species when they are always reproductively ready,

or are close to being ready, when opportunities to breed arise (Laaksonen et al., 2002; Mysterud

et al., 2008). This may especially be important in highly seasonal environments where it may be

energetically impossible for females to breed year-round, and so males may enter the breeding

season earlier and leave it later than females (Bronson, 1985). Alternatively, a shortened

breeding season may also be an adaptive strategy by males under stressful conditions. In this

system, male *C. penicillatus* on exclosure plots may opt to not increase energy allocation for

breeding when environmental conditions are not optimal for females since the metabolic

consequences of breeding under adverse conditions can adversely impact future demographic

success (Descamps et al. 2007, Lane et al. 2010)

Within-species phenological synchrony, which occurs when there is low variation in the

timing of a biological event for a given species, may arise from consistent responses among

individuals to external cues (i.e., shared responses to environmental factors at a given time

scale). Rudolf and Carter (2022) posit that the implications of low and high phenological

synchrony are dependent on competitive context. In this case, there is potentially a benefit to

high synchrony (i.e., shorter breeding period and/or specific responses to environmental cues) in

some groups (e.g., female kangaroo rats) and low synchrony (i.e., longer breeding period and/or

less specific responses to environmental cues) in others (e.g., male kangaroo rats and *C. baileyi*

males on controls). However, given the nature of our data, it is not immediately clear how to

quantify the strength of intraspecific synchrony. If intraspecific synchrony is driven by

individuals exhibiting strong and consistent responses to drivers, then the model results may

provide some insights into the degree of intraspecific synchrony. Consistent with the impression

of higher synchrony, female kangaroo rats exhibit strong responses to environmental variables

with narrower distribution curves for parameter estimates, and thus, low within-species

variability in their environmental responses (Fig 4). In contrast, both pocket mice species

exhibited a broader range of responses to external stressors (Fig 4). This larger variation in the

parameter estimates was generally consistent on both exclosures and controls and across males

and females, suggesting that high intraspecific variability in these species may not be driven

specifically by the direct impacts of competition with kangaroo rats. This may mean that these

pocket mice exploit environmental conditions that are underutilized by kangaroo rats as an

adaptive strategy. Consequently, this may facilitate population-level shifts in phenological timing

in response to changes in competitive context. While shifts in biotic context appear to drive

changes in the breeding phenology of both pocket mice, these had a subtle impact on the

structure of the best fit model on controls versus exclosures. Important environmental drivers on

control plots were generally also important on exclosure plots but the strength and significance

of some drivers changed depending on the biotic context. What this may mean for the

interoperability of models fit under one biotic context for predicting phenology in other biotic

contexts is currently unclear but bears further investigation.

Overall, the variation in the phenological responses of desert rodents under different

biotic conditions that we have documented here highlight the important role of species

interactions in driving phenology. While these results cannot distinguish whether the impacts of

competition occur through resource depletion, behavioral interference, or other impacts of the

kangaroo rats, these results show that removing a competitor can influence a species' phenology.

Future studies that can tease apart these direct and indirect impacts may be useful in developing a

robust understanding of the effects of competition in species' phenology. By recognizing that

competitive landscapes can influence the responses of breeding individuals, we gain a more

nuanced understanding of one of the key biological phenomena that can alter ecosystem

dynamics.

ACKNOWLEDGMENTS

We thank the numerous volunteers, staff, and research assistants, who collected the Portal

Project data over the years. This work was supported by research grants awarded to SKME by

the US National Science Foundation (NSF DEB-1929730) and USDA National Institute of Food

and Agriculture, Hatch Project FLA-WEC-005983.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

REFERENCES

Audzijonyte, Asta, and Shane A. Richards. 2018. "The Energetic Cost of Reproduction and Its

Effect on Optimal Life-History Strategies." *The American Naturalist*, June.

https://doi.org/10.1086/698655.

Behrends, Philip, Martin Daly, and Margo I. Wilson. 1986. "Aboveground Activity of Merriam's

Kangaroo Rats (Dipodomys Merriami) in Relation to Sex and Reproduction." *Behaviour* 96

(3/4): 210–26.

Bledsoe, Ellen K., and S. K. Morgan Ernest. 2019. "Temporal Changes in Species Composition

Affect a Ubiquitous Species' Use of Habitat Patches." *Ecology* 100 (11): e02869.

https://doi.org/10.1002/ecy.2869.

Bradshaw, William E., and Christina M. Holzapfel. 2007. "Evolution of Animal

Photoperiodism." *Annual Review of Ecology, Evolution, and Systematics* 38 (1): 1–25.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110115.

Bronson, Franklin H.1985. "Mammalian reproduction: an ecological perspective." *Biology of

reproduction* 32 (1):1-26.

Buss, Nicholas, Lindsey Swierk, and Jessica Hua. 2021. "Amphibian Breeding Phenology

Influences Offspring Size and Response to a Common Wetland Contaminant." *Frontiers in

Zoology* 18 (1): 31. https://doi.org/10.1186/s12983-021-00413-0.

Carter, Shannon K., and Volker H. W. Rudolf. 2022. "Exploring Conditions That Strengthen or

Weaken the Ecological and Evolutionary Consequences of Phenological Synchrony." *The

American Naturalist* 200 (5): E189–206. https://doi.org/10.1086/720899.

Christensen, Erica, Glenda Yenni, Hao Ye, Juniper Simonis, Ellen Bledsoe, Renata Diaz, Shawn

Taylor, Ethan White, and S. Ernest. 2019. "Portalr: An R Package for Summarizing and

Using the Portal Project Data." *Journal of Open Source Software* 4 (33): 1098.

https://doi.org/10.21105/joss.01098.

Cohen, Jeremy M., Marc J. Lajeunesse, and Jason R. Rohr. 2018. "A Global Synthesis of Animal

Phenological Responses to Climate Change." *Nature Climate Change* 8 (3): 224–28.

https://doi.org/10.1038/s41558-018-0067-3.

DeLucia, Evan H., Paul D. Nabity, Jorge A. Zavala, and May R. Berenbaum. 2012. "Climate

Change: Resetting Plant-Insect Interactions1." *Plant Physiology* 160 (4): 1677–85.

https://doi.org/10.1104/pp.112.204750.

Denny, Ellen G., Katharine L. Gerst, Abraham J. Miller-Rushing, Geraldine L. Tierney, Theresa

M. Crimmins, Carolyn A. F. Enquist, Patricia Guertin, et al. 2014. "Standardized Phenology

Monitoring Methods to Track Plant and Animal Activity for Science and Resource

Management Applications." *International Journal of Biometeorology* 58 (4): 591–601.

https://doi.org/10.1007/s00484-014-0789-5.

Descamps, Sébastien, Stan Boutin, Dominique Berteaux, and Jean-Michel Gaillard. 2007.

"Female red squirrels fit Williams' hypothesis of increasing reproductive effort with

increasing age." *Journal of Animal Ecology*. 1192-1201.

Diaz, Renata M., and S. K. Morgan Ernest. 2022. "Maintenance of Community Function through

Compensation Breaks down over Time in a Desert Rodent Community." *Ecology* 103 (7):

e3709. https://doi.org/10.1002/ecy.3709.

Ding, Yongxia, Zhi Li, and Shouzhang Peng. 2020. "Global analysis of time-lag and-

accumulation effects of climate on vegetation growth." *International Journal of Applied*

Earth Observation and Geoinformation 92:102179.

https://doi.org/10.1016/j.jag.2020.102179

Ernest, S. K. M., Glenda M. Yenni, Ginger Allington, Erica M. Christensen, Keith Geluso, Jacob

R. Goheen, Michele R. Schutzenhofer, et al. 2016. "Long□term Monitoring and

Experimental Manipulation of a Chihuahuan Desert Ecosystem near Portal, Arizona (1977–2013) - Morgan Ernest - 2016 - Ecology - Wiley Online Library." 2016. https://esajournals.onlinelibrary.wiley.com/doi/full/10.1890/15-2115.1.

Forchhammer, Mads C., Eric Post, and Nils Chr Stenseth. 1998. "Breeding Phenology and Climate¨." *Nature* 391 (6662): 29–30. https://doi.org/10.1038/34070.

French, Alan R. 1977. "Circannual Rhythmicity and Entrainment of Surface Activity in the Hibernator, Perognathus Longimembris." *Journal of Mammalogy* 58 (1): 37–43. https://doi.org/10.2307/1379725.

Gelman, Andrew. 2008. "Scaling regression inputs by dividing by two standard deviations." *Statistics in medicine* 27(15): 2865-2873.

Glick, Patricia, Bruce A Stein, Naomi A Edelson, and National Wildlife Federation. 2011. *Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessment*. Washington, D.C.: National Wildlife Federation.

Gordo, Oscar, and Juan José Sanz. 2005. "Phenology and Climate Change: A Long-Term Study in a Mediterranean Locality." *Oecologia* 146 (3): 484–95. https://doi.org/10.1007/s00442-005-0240-z.

Helm, Barbara, Rachel Ben-Shlomo, Michael J. Sheriff, Roelof A. Hut, Russell Foster, Brian M. Barnes, and Davide Dominoni. 2013. "Annual Rhythms That Underlie Phenology: Biological Time-Keeping Meets Environmental Change." *Proceedings of the Royal Society B: Biological Sciences* 280 (1765): 20130016. https://doi.org/10.1098/rspb.2013.0016.

Kelt, Douglas, Mark Taper, and Peter L. Meserve. 1995. "Assessing the Impact of Competition on Community Assembly: A Case Study Using Small Mammals - Kelt - 1995 - Ecology -

Wiley Online Library." 1995.

https://esajournals.onlinelibrary.wiley.com/doi/abs/10.2307/1940935.

Kenagy, G. J., and George A. Bartholomew. 1981. "Effects of Day Length, Temperature, and

Green Food on Testicular Development in a Desert Pocket Mouse Perognathus Formosus."

Physiological Zoology 54 (1): 62–73.

Laaksonen, Toni, Erkki Korpimäki, and Harri Hakkarainen. 2002. "Interactive Effects of

Parental Age and Environmental Variation on the Breeding Performance of Tengmalm's

Owls." *Journal of Animal Ecology* 71 (1): 23–31. https://doi.org/10.1046/j.0021-

8790.2001.00570.x.

Lane, Jeffrey E., Stan Boutin, John R. Speakman, and Murray M. Humphries. 2010. "Energetic

Costs of Male Reproduction in a Scramble Competition Mating System." *Journal of Animal*

Ecology 79 (1): 27–34. https://doi.org/10.1111/j.1365-2656.2009.01592.x.

Ma, Yu-Jun, Fang-Zhong Shi, Xia Hu, and Xiao-Yan Li. 2021. "Climatic Constraints to Monthly

Vegetation Dynamics in Desert Areas Over the Silk Road Economic Belt." *Remote*

Sensing 13 (5): 995. https://doi.org/10.3390/rs13050995

McLean, Bryan S., and Robert P. Guralnick. 2021. "Digital Biodiversity Data Sets Reveal

Breeding Phenology and Its Drivers in a Widespread North American Mammal." *Ecology*

102 (3): e03258. https://doi.org/10.1002/ecy.3258.

Mysterud, Atle, Christophe Bonenfant, Leif Egil Loe, Rolf Langvatn, Nigel G. Yoccoz, and Nils

C. Stenseth. 2008. "The Timing of Male Reproductive Effort Relative to Female Ovulation

in a Capital Breeder." *Journal of Animal Ecology* 77 (3): 469–77.

https://doi.org/10.1111/j.1365-2656.2008.01365.x.

Ostfeld, Richard S. 1985. "Limiting Resources and Territoriality in Microtine Rodents." *The*

American Naturalist 126 (1): 1–15.

Pearce-Higgins, J. W., and D. W. Yalden. 2004. "Habitat Selection, Diet, Arthropod Availability

and Growth of a Moorland Wader: The Ecology of European Golden Plover Pluvialis

Apricaria Chicks." *Ibis* 146 (2): 335–46. https://doi.org/10.1111/j.1474-919X.2004.00278.x.

Pedersen, Eric J., David L. Miller, Gavin L. Simpson, and Noam Ross. 2019. "Hierarchical

generalized additive models in ecology: an introduction with mgcv." *PeerJ* 7: e6876.

https://doi.org/10.7717/peerj.6876

R Core Team. 2021. "R: A language and environment for statistical computing". *R Foundation*

for Statistical Computing. Vienna, Austria. http://www.R-project.org

Reed, Thomas E., Stephanie Jenouvrier, and Marcel E. Visser. 2013. "Phenological Mismatch

Strongly Affects Individual Fitness but Not Population Demography in a Woodland

Passerine." *Journal of Animal Ecology* 82 (1): 131–44. https://doi.org/10.1111/j.1365-

2656.2012.02020.x.

Roslin, Tomas, Laura Antão, Maria Hällfors, Evgeniy Meyke, Coong Lo, Gleb Tikhonov, Maria

del Mar Delgado, et al. 2021. "Phenological Shifts of Abiotic Events, Producers and

Consumers across a Continent." *Nature Climate Change* 11 (3): 241–48.

https://doi.org/10.1038/s41558-020-00967-7.

Simpson, Gavin L. 2018. "Modelling Palaeoecological Time Series Using Generalised Additive

Models." *Frontiers in Ecology and Evolution* 6.

https://www.frontiersin.org/article/10.3389/fevo.2018.00149.

Thackeray, Stephen J., Timothy H. Sparks, Morten Frederiksen, Sarah Burthe, Philip J. Bacon,

James R. Bell, Marc S. Botham, et al. 2010. "Trophic Level Asynchrony in Rates of

Phenological Change for Marine, Freshwater and Terrestrial Environments." *Global*

Change Biology 16 (12): 3304–13. https://doi.org/10.1111/j.1365-2486.2010.02165.x.

Visser, Marcel E, and Christiaan Both. 2005. "Shifts in Phenology Due to Global Climate

Change: The Need for a Yardstick." *Proceedings of the Royal Society B: Biological*

Sciences 272 (1581): 2561–69. https://doi.org/10.1098/rspb.2005.3356.

Walther, Gian-Reto, Eric Post, Peter Convey, Annette Menzel, Camille Parmesan, Trevor J. C.

Beebee, Jean-Marc Fromentin, Ove Hoegh-Guldberg, and Franz Bairlein. 2002. "Ecological

Responses to Recent Climate Change." *Nature* 416 (6879): 389–95.

https://doi.org/10.1038/416389a.

Wann, Gregory T., Cameron L. Aldridge, Amy E. Seglund, Sara J. Oyler-McCance, Boris C.

Kondratieff, and Clait E. Braun. 2019. "Mismatches between Breeding Phenology and

Resource Abundance of Resident Alpine Ptarmigan Negatively Affect Chick Survival."

Ecology and Evolution 9 (12): 7200–7212. https://doi.org/10.1002/ece3.5290.

Wen, Youyue, Xiaoping Liu, Qinchuan Xin, Jin Wu, Xiaocong Xu, Fengsong Pei, Xia Li et al.

2019. "Cumulative effects of climatic factors on terrestrial vegetation growth." *Journal of*

Geophysical Research: Biogeosciences 124 (4): 789-806.

https://doi.org/10.1029/2018JG004751

Wolkovich, Elizabeth M., Benjamin I. Cook, and T. Jonathan Davies. 2014. "Progress towards

an Interdisciplinary Science of Plant Phenology: Building Predictions across Space, Time

and Species Diversity." *New Phytologist* 201 (4): 1156–62.

https://doi.org/10.1111/nph.12599.

Wood, Simon, and Maintainer Simon Wood.2015. "Package 'mgcv'." *R package version* 1, no.

29: 729.

FIGURE LEGENDS

Figure 1. Reproductive phenology of female (left panel) and male (right panel) kangaroo rats in

        control plots (top panel), and Bailey's pocket mice in control (middle panel) and

        exclosure (bottom panel) plots in an experimental field site near Portal, Arizona. Plots

        show estimated smooths of the month term from species-, sex-, and treatment-specific

        generalized additive models (GAMs). Blue and red shading indicates periods of

        significant increase and decrease in breeding proportions, respectively. In cases where

        seasonal patterns or periods of significant increase or decrease could not be estimated

        from the model, raw data on proportion of breeding individuals were plotted (male

        Dipodomys spp. and *C. baileyi*; B, D).

Figure 2. Association between select abiotic and biotic factors with the proportion of

        reproductive female (left panel) and male (right panel) Bailey's pocket mouse (top panel)

        and desert pocket mouse (bottom panel) in control and kangaroo rat exclosure plots.

Figure 3. Reproductive phenology of female (left panel) and male (right panel) kangaroo rats in

        control plots (top panel), and desert pocket mice in control (middle panel) and exclosure

        (bottom panel) plots in an experimental field site near Portal, Arizona. Details on the

        configuration of these plots are like Figure 1.

Figure 4. Distribution curves of the abiotic covariate slope estimates indicating the response of

        female (left panel) and male (right panel) Bailey's pocket mouse (top panel) and desert

        pocket mouse (bottom panel) in control and kangaroo rat exclosure plots, compared to

        kangaroo rat estimates. Curves indicate the normal distribution with mean equal to the

        point estimate, and standard deviation equal to the standard error of the coefficients.

FIGURES

Figure 1.

Figure 2.

Figure 3.

Figure 4.