Flexible parental care ensures robustness of post-embryonic development

Ahva L. Potticary*, Christopher B. Cunningham, and Allen J. Moore

Department of Entomology, University of Georgia, Athens, Georgia, 30602 USA

*Corresponding Author:

Ahva L. Potticary

120 Cedar Street

Biological Sciences West #360

Athens, GA 30602

ORCID identifier:

Ahva L. Potticary: 0000-0002-1157-5315

Christopher B. Cunningham: 0000-0003-3965-2076

Allen J. Moore: 0000-0002-1498-3322

Preprint Servers:

BioRxiv: DOI forthcoming

Classification:

Evolution

Keywords: maternal effect, transgenerational effect

This PDF file includes:
Main Text
Figures 1 - 3
25

Abstract

Parental care is expected to buffer offspring against poor environments, yet such environments are

common and parental care is both rare and incredibly variable. Moreover, parental care is complex,

composed of multiple acts that together influence offspring fitness. We tested the hypothesis that

parental care can act as a buffer because it is composed of multiple component behaviors that allow

for robustness of post-embryonic development in response to changing environments. We test this

hypothesis in the burying beetle *Nicrophorus orbicollis*, a species with complex parental care, by

determining how this species responds to a poor developmental environment due to reduced parental

care. We manipulated the duration of parental care received and measured both offspring and

grandoffspring traits of males and females. As expected, reduced care affected key traits that influence

fitness including offspring development time and mass. However, offspring that received reduced care

responded by increasing uniparental care when they became parents and, in addition, shifted the type

of parenting behavior used to transfer nutrients to offspring. Parents that received reduced care spent

more time directly feeding offspring, while parents that received full care invested more time

maintaining the food resource. As a result, both the number and mass of grandoffspring were

unaffected by the developmental experience of their parent. Our results show that flexibility in the

behaviors composing parental care can overcome poor developmental environments and limit negative

parental effects to a single generation, suggesting that parental care can confer robustness both within

and across generations.

Significance Statement

Parental care is thought to buffer offspring against poor environments during their development, but what components of parenting shift to achieve this buffering? Here we manipulated the amount of care provided by burying beetles, an insect that directly feeds its offspring. Reduced care affected offspring development and mass. However, when these offspring reached adulthood, they modified their own care so that the effects were limited to a single generation. We suggest that these result show that parental care not only serves as a buffer within-generation but can also confer robustness against poor natal environments across generations.

Introduction

Parental care is thought to evolve to buffer developing young from harsh environments (1). Yet despite their importance for offspring development, parental care strategies show a bewildering amount of variation within and across taxa (2-7) and have arisen across a wide diversity of ecological conditions (1, 8, 9). Moreover, while parental care is often treated as a single trait, it is a category encompassing multiple behaviors, such as offspring feeding, maintenance of resources, nest defense, preparation and preservation of food, among others (10, 11). Thus, it is unclear how a composite trait like parental care buffers offspring development when the types of environmental hazards that an organism may experience are expected to be incredibly variable over space and time.

Implicit to the idea that parental care acts as a buffer is that it ensures appropriate offspring development despite variation in environmental insults. The maintenance of function despite changing environments – or robustness - is a ubiquitous feature of complex systems from artificially evolved electronic circuits to gene networks to ecological communities (12-16). Research across systems indicates that robustness in different environments is achieved though changes in the lower-level units that compose the system. For example, if a single gene stops functioning in a gene network, functionality may be maintained at the level of the network because multiple genes can produce the same product/outcome (17). As parental care is a composite trait, a robustness perspective predicts that buffering of offspring development across environments may be accomplished based on changes in different components of the system – that is, the behaviors composing parental care (18, 19). Selection for robustness itself can produce complex and unpredictable structures in lower-level units (16), as mechanisms that increase robustness against perturbation often trade-off with other types of functions (e.g., replication rate; 20). Since parental care is expected to evolve to buffer offspring, one solution for the diversity and complexity of parental care strategies is that there is selection for behavioral

robustness to protect offspring against a range of threats, rather than for a particular behavior in a

specific environment. Yet, whether behavior follows similar principles of robustness as other complex

systems is poorly understood.

Here we test the hypothesis that parental care can act as a buffer because it confers robustness

to post-embryonic development for the subsocial burying beetle *Nicrophorus orbicollis* through

alteration of the components of parental care. If parental care confers robustness, robustness theory

predicts that, across environments, the form and frequency of specific parental care behaviors may

change to maintain offspring developmental outcomes across a range of conditions. Burying beetles

require small animal carcasses to breed, and compete with other burying beetles and species that feed

on carrion for this essential resource (21). Upon securing a carcass, parents bury it partially or fully

underground, depending on species (20), remove the exterior (e.g., fur), and coat it with compounds

that slow decomposition through removal of bacteria and fungi and form it into a ball with an opening

for larvae to occupy and within which parents predigest the carcass to form a "soup" (4, 22). Females

lay eggs in the soil and upon hatching, larvae crawl to the carcass and enter the cavity. Parents begin

feeding larvae immediately upon detection, and larvae of *N. orbicollis* cannot survive without at least

several hours of direct regurgitation of predigested food from the parent (23). Parents enable nutrient

transfer from carcasses to offspring using multiple behaviors, including burying and defending

carcasses (which in part acts to ensure that the carcass is available for longer; 21, 24), destruction of

bacterial and fungal consumers (25, 26), creation of larval cavities to support offspring self-feeding

(23, 27, 28), and direct transfer of food to larvae through regurgitation (29). Larvae do not feed after

they leave the carcass, and for this reason, adult body size is largely determined by the efficiency of

resource acquisition during the larval period on carcass. Body size is a key fitness trait for *Nicrophorus*,

as larger beetles are more effective competitors for carcasses (21, 30). Thus, parental care creates a

direct link between a larva's developmental experience and competitive ability in later life.

We experimentally induced perturbation of the parental care environment by altering the

duration of parental care that mixed foster broods received to produce larvae that experience reduced

(six hours) or full care (approximately seven days). Poor developmental environments, including those

arising from poor parental care, often produce cascading multigenerational effects (reviewed in 31)

and thus represent an environmental perturbation. We then tracked these individuals into adulthood

and determined how removal of parental care influenced developmental times, behavior, and

reproductive success. Beetles that received full care were larger and developed faster during the

vulnerable period on the breeding carcass than beetles that received reduced care. However, this effect

did not carry through to the next generation (i.e., to the grandoffspring). Parents that received reduced

care as larvae spent more time regurgitating food to their larvae than parents that had received full care.

As a result, parents from full and reduced care treatments produced larvae of similar numbers and

weights. Together, these results indicate that parental care behavior may not only buffer offspring

development within generation but may act as a developmental buffer that mitigates the legacy of

environments past. This robustness was achieved through flexible changes of parental behaviors that

transfer nutrients to offspring and the ability of offspring to accept these differences.

Results

Effect of Perturbation of Care Environment on Development.

We set up 36 families, 18 in each treatment, of mixed-parentage broods with a single, unrelated female

caretaker in the full and reduced care treatments. Overall larval survival was high in both treatments;

from an initial 10 fostered larvae placed on carcass, 3-10 survived to dispersal from the mouse with a

mean 8.3 ± 0.4 surviving larvae for full care families and 8.7 ± 0.4 for reduced care ($F_{1,34} = 0.430$, P =

0.516). Three (2%) larvae died after dispersal from the full care treatment, and 3 (1.9%) died from the

reduced care treatment. A further 14 (9.6%) died in the pupal stage in the full care treatment, and 16

(10.4%) in the reduced care.

Larvae in the full care treatment developed faster on the carcass than larvae that received

reduced care (χ^2 = 18.86, P < 0.0001, N = 305). Larvae that received full care completed development

from hatch to dispersal in 7 - 8 days (mean = 7.4) and reduced care larvae dispersed in 8 - 9 days (mean

= 8.3) in reduced care families; **Fig. 1A)**. The largest effect of reduced care was on larval weight at

dispersal from the carcass (**Fig. 1B**). There was significant variation between care treatments ($F_{1,269}$ =

433.253, P < 0.0001), as well as caretakers nested within treatments ($F_{34,269}$ = 10.155, P < 0.0001).

Controlling for variation amongst families, larvae that received full care weighed 500.4 ± 4.4 mg,

whilst larvae that received reduced care weighed 373.4 ± 4.2 mg. Larval weight at dispersal was highly

correlated with adult size (r = 0.91, N = 248).

However, even though larvae in the full care treatment developed faster on the carcass, duration

of total post-embryonic development (hatch to eclosion) was influenced primarily by larval weight at

dispersal ($F_{1,232}$ = 77.283, P < 0.0001) and not on care treatment ($F_{1,232}$ = 1.311, P = 0.2535; **Fig. 1C**).

There was significant variation attributable to caretaker ($F_{34,261}$ = 9.301, P < 0.0001), larval weight

($F_{1,261}$ = 34.501, P < 0.0001) and duration of parental care received ($F_{1,261}$ = 13.47, P = 0.0003) on time

spent after dispersal to pupation. The same factors influenced duration of pupal development (pupae

to adult eclosion) with significant variation arising from larval weight ($F_{1,231}$ = 55.586, P < 0.0001),

caretaker ($F_{34,231}$ = 3.173, P < 0.0001), but not duration of parental care received ($F_{1,231}$ = 0.883, P =

0.348). Thus, the duration of care received influenced the speed with which the larvae developed on

the carcass, but this only shifted development time.

Robustness of Parental Care Against Perturbation.

We next established 89 uniparental broods using the resulting offspring from the reduced and full care

treatments (26 full care males, 20 full care females, 24 reduced care males, and 17 reduced care

females). There is no evidence that the type of parenting received influenced the likelihood that

individuals would provide parental care. Focal male caretakers were more likely to abandon before

eggs hatched (reduced care males: 11/24 breeding attempts; full care males: 11/26 breeding attempts)

than female caretakers (reduced care females: 2/17 breeding attempts; full care females 2/20 breeding

attempts). Once larvae hatched, most focal parents stayed for the entire larval development on the

carcass. When caretakers did abandon larvae, it was almost exclusively males (10/29 males, 1/33

females; $\chi^2 = 11.640$, df $= 1$, P $= 0.0012$). The one female that abandoned early had received full care;

for males, four had received full care and six received reduced care.

Parental care behavior of both males and females was affected by the length of care they

received as larvae. There are strong differences between uniparental females and males in this species

(**Fig. 2**). Females spent more time on the carcass during the observation period than did males ($F_{1,55} =$

17.610, P < 0.001; **Fig. 2***A*) regardless of the duration of parental care they received ($F_{1,55} = 1.878$, P

$= 0.176$) and there was not a significant interaction between the duration of care received and sex of

the caregiver on amount of time spent on the carcass ($F_{1,55} = 1.332$, P $= 0.253$). Direct care was strongly

influenced by the duration of parental care received ($F_{1,55} = 6.761$, P $= 0.012$); individuals that received

reduced care provided more direct care themselves and this pattern was consistent for both males and

females (**Fig. 2***B*); there was no interaction between the sex of the caregiver and the effect of parenting

received ($F_{1,55} = 0.111$, P $= 0.741$). Females provided more direct care than males under both treatments

($F_{1,55} = 12.699$, P $= 0.0008$). In contrast, indirect care was not affected by the duration of parental care

received ($F_{1,55} = 1.181$, P $= 0.282$; **Fig. 2***C*) or sex of the caregiver ($F_{1,55} = 3.072$, P $= 0.085$), and there

was no statistically significant interaction between these treatments ($F_{1,55} = 0.705$, P $= 0.405$).

Overall, the care a parent received as a larva did not result in negative effects on grandoffspring.

There was no statistically significant effect of caregiver sex ($F_{1,56}$ = 3.583, P = 0.064) or duration of

care received by parent ($F_{1,56}$ = 0.057, P = 0.812) on the length of time larvae spent on the carcass, but

there was a significant interaction ($F_{1,56}$ = 5.796, P = 0.019; **Fig. 3A**). Offspring from females that had

received reduced care took longer to develop on the carcass than offspring from females that received

full care, but the opposite was true for offspring reared by males. We again included carcass mass as a

covariate ($F_{1,56}$ = 1.564, P = 0.216). There was no effect of caregiver sex on average larval mass ($F_{1,56}$

= 1.855, P = 0.179), care received by parent ($F_{1,56}$ = 1.340, P = 0.252) or their interaction ($F_{1,56}$ = 0.477,

P = 0.493), controlling for carcass mass as a covariate ($F_{1,56}$ = 0.047, P = 0.829; **Fig. 3B**). Females

reared more offspring than did males in general ($F_{1,56}$ = 15.521, P = 0.0002), but there was no effect of

parental treatment on the number of offspring reared ($F_{1,56}$ = 0.095, P = 0.759; **Fig. 3C**) and no

interaction between caregiver sex and the duration of parental care received ($F_{1,56}$ = 0.429, P = 0.515).

We included carcass mass as a covariate ($F_{1,56}$ = 1.562, P = 0.217), but given the restricted range of

masses used we see no effect here or in any of our analyses.

Discussion

We tested the hypothesis that parental care can act as a buffer because it allows for robustness of post-

embryonic development by altering components of care in response to perturbation. We show that the

architecture of parental care behaviors of burying beetles translates developmental perturbation into

consistent phenotypic outcomes across generations. Experimentally minimizing parental care greatly

impacted ecologically relevant offspring traits that are aligned with fitness, including duration of the

vulnerable stage on the carcass (**Fig. 1A**; 21, 29) and body size (**Fig. 1B**; 30). Parents that received

reduced parental care themselves spent more time directly transferring nutrients to offspring, as

opposed to parents that received full care, who invested more in indirect mechanisms to enable nutrient

acquisition of larvae (i.e., maintenance of a food resource). This reflects that parents differentially

express functionally redundant behaviors – i.e., providing self-feeding opportunities to offspring versus

direct provisioning – in response to perturbation. This resulted in the negative effects of a poor

developmental environment being constrained to a single generation. Together, these data may reflect

robustness of parental care to perturbation during development, even when that perturbation arises

from variation in the care environment itself, which is not always the case (31-33).

Parental care strategies show a bewildering amount of variation across taxa (2-7) and have

arisen across a wide diversity of ecological conditions (1, 8, 9). Our results may provide an explanation

for why there is so much variation in parental care (e.g., 34) given its importance to offspring

development (10). Theory and empirical work indicates that selection for robustness itself can produce

unpredictable and sometimes counterintuitive patterns in lower-level units (16), in this case the

behaviors composing a complex behavior, as mechanisms that increase robustness against perturbation

often trade-off with other types of functions (e.g., replication rate; 20). A "byzantine" complexity of

biological systems may be what enables their continued functionality across a wide array of

environmental perturbations (35, 36). Moreover, once a robust system has evolved, it may be able to

accommodate environmental variation and dampen changes in natural selection (12, 19). Theoretical

work on parental care found that stochastic environments are expected to be associated with a range of

care strategies (37), which indirectly supports the idea that selection for robustness may increase the

complexity and diversity of parental care strategies. *Nicrophorus orbicollis* live in highly heterogenous

and variable environments where they are generally competitively superior to other species due to

larger body size (21, 38, 39), and parental care is important for body size across *Nicrophorus* spp. (40).

This may indicate that the great variation observed both in how individual burying beetles show care

(34, 41) and who shows care (e.g., uniparental to biparental to communal; 4, 5, 42) may emerge from

selection for consistency of offspring phenotypes relative to the prevailing environmental context (38),

which may require variation in the form and frequency of components of parental care (**Fig. 2**).

Selection experiments for larger offspring across a diversity of perturbations may shed light on whether

selection on the robustness of offspring post-embryonic development influences the structure of

parental care.

Robustness is often misunderstood to reflect a lack of change in response to perturbation, but

robustness refers to consistency of function, which often requires flexibility or change at the level of

interacting units (13). Systems spend more time in robust states because, over time, fragile systems do

not last (12). Biology is hierarchical by nature. For this reason, robustness at one level of level of

organization, e.g., the outcome of parental care, does not imply a lack of change at other levels, e.g.,

the expression of component parental care behaviors. That parental care has evolved independently

multiple times across taxa (e.g., 7, 43) may indicate that it appears because parental care strategies

confer robustness, or, diverse taxa end up repeatedly evolving parental care strategies because these

are the most stable mechanisms for ensuring appropriate offspring development. A robustness

perspective further reconciles why there is such a great diversity of life history, ecological, and social

contexts associated with the evolution of parenting (1, 7, 37, 44). Future work could disentangle

whether robustness of parental care in burying beetles can also ensure functional offspring outcomes

under a suite of both historical and novel perturbations.

Organisms both construct and evolve in response to their experience of the environment (45).

Robust systems are expected to emerge over evolutionary time, and the architecture of such systems

are often enabled by modularity and functional redundancy (12, 13). We found that removing parental

care induced strong changes in ecologically relevant traits. Offspring that received poor parenting

responded by increasing uniparental care when they became parents and shifting the type of nutrient-

transfer behavior they showed, resulting in the amelioration of perturbation during the development of

parents. In conclusion, parenting may be an effective evolutionary mechanism for buffering poor past

and present environments, including those imposed during the parents' development. This robustness

may be enabled by dynamic changes in the units of parental care – the component behaviors

themselves.

Materials and Methods

Insect colony and husbandry.

We used *Nicrophorus orbicollis* from an outbred colony maintained at the University of Georgia that

is regularly supplemented with wild-caught beetles from Whitehall Forest in Athens, GA, USA. All

beetles were maintained in an incubator (Percival, model no. I41VLC8, Perry, IA) at 22 ± 0.1°C on a

14:10 hour light:dark cycle that simulated summer breeding conditions. We bred all individuals used

here in the laboratory to ensure that parentage, age, and to standardize rearing conditions. We housed

beetles individually at larval dispersal in 9-cm-diameter by 4-cm-deep circular compostable deli

containers (EP-RDP5, Eco-Products) filled with ~2.5 cm of potting soil (Performance Organics All

Purpose In-Ground Soil, Miracle-Gro). Larvae had no further social interactions with other burying

beetles until they were randomly allocated into treatments. We defined individuals as age Day 1 on the

day of eclosion. Beetles were fed *ad libitum* organic ground beef twice a week following adult eclosion.

Effect of Perturbation of Care Environment on Development.

We manipulated the duration of parental care to determine the effect of parental care on offspring traits

related to fitness and their expression of care once they attained adulthood. We randomly mated focal

virgins at Day 13-19 by placing an unrelated male and female in a small plastic container and observed

them to ensure that mating occurred. The pair was then moved to a plastic breeding box filled with

approximately two cm of potting soil moistened with distilled water and containing a freshly thawed

mouse 20.8 ± 0.6 g SD (max = 22.2, min = 20.0; Rodent Pro, Inglefield, IN, U.S.A.). There was no

difference in the size of the mouse provided between treatments (20.9 g in full care treatment, 20.8 g

in reduced care treatment; $F_{1,34} = 0.153$, P = 0.698). We then returned breeding boxes to the incubator

and kept them on a darkened shelf to simulate underground breeding conditions. We checked breeding

boxes for eggs every twenty-four hours. Once eggs we observed eggs, we moved the female and brood

ball to a new breeding box filled with approximately two cm of moistened potting soil, while the male

was removed. We left eggs in the original breeding box with a small piece of ground beef. Once 90

hours had elapsed since egg laying, we checked the breeding box containing the eggs every hour to

determine when larvae had hatched and crawled to the beef. After we observed the focal females'

larvae hatch, we provided the focal female with an unrelated, mixed brood of 10 larvae. Burying beetles

identify their offspring temporally and eat any larvae that arrive too early on the carcass (46), and thus

we only provided females with larvae after her own larvae had hatched. For the full duration parental

care treatment, we allowed females to associate with larvae until at least 2 of the larvae had dispersed

from the breeding carcass (~7 - 8 days). For the reduced duration parental care treatment, we allowed

females to associate with larvae for 6 hours and then the female was removed. We checked

experimental treatments daily to determine larval dispersal date, subsequently, larval survival.

We measured several offspring traits related to fitness (47), including mass at dispersal,

duration of developmental periods, and adult size. We measured larval mass at dispersal to the nearest

0.1 mg. We also measured the duration of development on the carcass (from hatch until dispersal),

which when parental care and all larval growth occurs; the number of days of development from larval

dispersal to pupation; the number of days as pupae (from pupation to adult eclosion), and total duration

of development (from larval hatch until adult eclosion). When the larvae left the carcass (dispersal),

we placed larvae into individual containers with ~2 cm of moistened soil. We checked treatments daily

to determine the number of surviving larvae, length of larval period, pupation date, and adult ecolsion

date. We measured the pronotum length of full and reduced care offspring on Day 14 – 16 post-eclosion

with calipers to the nearest 0.01 mm.

Robustness of Care Against Perturbation.

At Day 14 – 16 following eclosion, adult beetles that received full or reduced care as offspring were

provided with an unrelated, virgin mate from our stock colony and a 19.6 - 21.2 g thawed mouse carcass

(mean = 20.6 g). These were placed into breeding container with ~2 cm of moistened soil. As variation

of parental care may manifest as variation of investment across the breeding cycle, breeding boxes

were fitted with an escape chamber to allow focal beetles to desert the breeding attempt. Breeding

boxes were checked daily for eggs. Beetles that escaped into the escape chamber prior to egg laying

were replaced in the breeding box. Beetles that escaped after egg-laying and remained in the escape

chamber for over 15 minutes were counted as deserters and removed. Once larvae were observed, the

non-focal partner was removed, and the focal male or female was left in a uniparental context to care

for the offspring. If the focal beetle deserted the breeding attempt prior to larval hatch, no further

monitoring was conducted on that breeding attempt.

As rates of parental provisioning may be affected by the number of larvae (48), we measured

parental care using both behavioral measures and larval traits associated with fitness. Behavioral

observations were conducted twenty-four hours after larva were first observed on the carcass for each

focal individual to quantify parental care. We moved breeding boxes into an observation room, and we

gave the focal beetle at least 30 minutes to acclimate prior to observation. All observations were

conducted under red lights to ensure no light effects on individual care behavior, as *N. orbicollis* are

nocturnal (49). In each behavioral observation, we recorded both direct and indirect parental care

behaviors (50) for 15 minutes. Direct care is defined as when the male or female fed larvae. Indirect

care is defined as activities related to carcass cleaning and maintenance, which allows for the

preservation of the carcass for future offspring provisioning and thus represents a type of parental care

without direct interactions with offspring. We also recorded time spent showing "no care", defined as

when the parent was on the carcass but not engaged in direct or indirect care, or the individual was off

the carcass. Behaviors were recorded continuously as both a duration and frequency (e.g., number of

times an adult fed larvae). This gave each individual three scores (direct care, indirect care, and no

care) which together summed to 15 minutes.

We measured offspring traits related to fitness reared by males or females that had received full

or reduced care as larvae themselves. We did not manipulate the number of offspring but instead

allowed parents to cull the number without interference. Of the remaining 61 uniparental broods, we

measured the number of offspring reared, the mass of the offspring at dispersal, and the amount of time

larvae spent on the carcass from hatching to dispersal. Measurements of development times and larval

mass were the same as used in the experimental manipulation of the duration of parental care (see

previous section).

Statistical Analyses.

We performed all statistical analyses using JMP Pro (v. 16.0.0, http://jmp.com) and produced figures

in SigmaPlot (v. 14.5, http://www.sigmaplot.co.uk). Results are means ± SE unless otherwise noted.

Effects on continuous offspring traits were analyzed with nested ANOVA, with family nested within

treatment. We analyzed categorical data with Pearson's Chi-square goodness-of-fit test. We analyzed

parental care behaviors and larval traits of grand offspring with two-way ANOVA with caretaker sex

and parenting experienced as factors.

Acknowledgements and Funding Sources

We would also like to thank Kathryn Kollars for assistance constructing escape chambers and for

developing the beetle graphics used in this study. We would also like to thank Patricia Moore and

Hans Otto for discussions and/or feedback on the manuscript, and Emily Shelby for assistance with

figures. ALP and AJM conceived of study and developed methods. ALP, AJM, and CBC collected

life history data. ALP conducted behavioural observations. AJM performed statistical analyses. ALP

and AJM created figures. ALP, AJM, and CBC all contributed to writing and editing the manuscript.

ALP was funded by a USDA cooperative agreement to AJM.

References

1.   E. O. Wilson, *Sociobiology: the New Synthesis* (Harvard University Press, Cambridge, 1975).
2.   A. L. Potticary, R. A. Duckworth, A neuroendocrine perspective on the origin and evolution
of cooperative breeding. *Ornithology* **138** (2021).
3.   H. J. Brockmann, "Cooperative breeding in wasps and vertebrates: the role of ecological
constraints." in The Evolution of Social Behavior in Insects and Arachnids*,* J. C. Choe, B. J.
Crespi, Eds. (Cambridge University Press, Cambridge, 1997), pp. 347-371.
4.   M. P. Scott, The ecology and behavior of burying beetles. *Ann. Rev. Entomol.* **43**, 595-618
(1998).
5.   K. M. Benowitz, E. C. McKinney, A. J. Moore, Difference in parenting in two species of
burying beetle, Nicrophorus orbicollis and Nicrophorus vespilloides. *J. Ethol.* **34**, 315-319
(2016).
6.   A. Cockburn, Prevalence of different modes of parental care in birds. *Proc. Biol. Sci.* **273**,
1375-1383 (2006).
7.   J. E. Mank, D. E. L. Promislow, J. C. Avise, Phylogenetic perspectives in the evolution of
parental care in ray-finned fishes *Evolution* **59**, 1570-1578 (2005).
8.   W. Jetz, D. R. Rubenstein, Environmental uncertainty and the global biogeography of
cooperative breeding in birds. *Curr. Biol.* **21**, 72-78 (2011).
9.   S. T. Emlen, The evolution of helping. 1. An ecological constraints model *Am. Nat.* **119**, 29-
39 (1982).
10.   N. J. Royle, S. H. Alonzo, A. J. Moore, Co-evolution, conflict and complexity: what have we
learned about the evolution of parental care behaviours? *Curr. Op. Behav. Sci.* **12**, 30-36
(2016).
11.   T. Székely, V. Remeš, R. P. Freckleton, A. Liker, Why care? Inferring the evolution of
complex social behaviour. *J. Evol. Biol.* **26**, 1381-1391 (2013).
12.   A. Wagner, Robustness, evolvability, and neutrality. *FEBS Lett.* **579**, 1772-1778 (2005).
13.   H. Kitano, Biological robustness. *Nature Rev. Gen.* **5**, 826-837 (2004).
14.   A. R. Ives, S. R. Carpenter, Stability and diversity of ecosystems. *Science* **317**, 58-62 (2007).

15. K. H. ten Tusscher, P. Hogeweg, Evolution of networks for body plan patterning; interplay of
modularity, robustness and evolvability. *PLoS Comp. Biol.* **7**, e1002208 (2011).
16. A. Thompson, P. Layzell, R. S. Zebulum, Explorations in design space: unconventional
electronics design through artificial evolution. *IEEE Trans. Evol. Comput.* **3**, 167-196 (1999).
17. P. Holme, Metabolic robustness and network modularity: a model study. *PloS One* **6**, e16605
(2011).
18. K. Chenard, R. Duckworth, "The special case of behavioral plasticity?" in *Phenotypic
Plasticity & Evolution: Causes, Consequences, and Controversies,* D. W. Pfennig, Ed. (CRC
Press, Boca Raton, 2021), 10.1201/9780429343001-16, pp. 301-325.
19. A. L. Potticary, R. A. Duckworth, Multiple environmental stressors induce an adaptive
maternal effect. *Am. Nat.* **196**, 487-500 (2020).
20. C. O. Wilke, J. L. Wang, C. Ofria, R. E. Lenski, C. Adami, Evolution of digital organisms at
high mutation rates leads to survival of the flattest. *Nature* **412**, 331-333 (2001).
21. I. C. Robertson, Nest intrusions, infanticide, and parental care in the burying beetle,
Nicrophorus orbicollis (Coleoptera: Silphidae). *J. Zool.* **231**, 583-593 (1993).
22. S. P. Shukla *et al.*, Microbiome-assisted carrion preservation aids larval development in a
burying beetle. *Proc. Natl. Acad. Sci.* **115**, 11274-11279 (2018).
23. A. Capodeanu-Nägler *et al.*, Offspring dependence on parental care and the role of parental
transfer of oral fluids in burying beetles. *Front. Zool.* **15**, 33 (2018).
24. S. T. Trumbo, D. S. Sikes, Resource concealment and the evolution of parental care
in burying beetles. *J. Zool.* **315**, 175-182 (2021).
25. W. Haberer, S. Steiger, J. K. Müller, Dynamic changes in volatile emissions of breeding
burying beetles. *Physiol. Entomol.* **39**, 153-164 (2014).
26. A. N. Arce, P. R. Johnston, P. T. Smiseth, D. E. Rozen, Mechanisms and fitness effects of
antibacterial defences in a carrion beetle. *J. Evol. Biol.* **25**, 930-937 (2012).
27. A. Duarte, D. Rebar, A. C. Hallett, B. J. M. Jarrett, R. M. Kilner, Evolutionary change in the
construction of the nursery environment when parents are prevented from caring for their
young directly. *Proc. Natl. Acad. Sci.* **118**, e2102450118 (2021).
28. A.-K. Eggert, J. K. Müller, "Biparental care and social evolution in burying beetles: lessons
from the larder" in *The Evolution of Social Behaviour in Insects and Arachnids,* J. C. Choe,
B. J. Crespi, Eds. (Cambridge University Press, Cambridge, 1997), pp. 216.
29. A.-K. Eggert, M. Reinking, J. K. Müller, Parental care improves offspring survival and
growth in burying beetles. *Anim. Behav.* **55**, 97-107 (1998).
30. M. Otronen, The effect of body size on the outcome of fights in burying beetles
(*Nicrophorus*). *Ann. Zool. Fenn.* **25**, 191-201 (1988).
31. F. A. Champagne, Epigenetic mechanisms and the transgenerational effects of maternal care.
Front. Neuroendocrinol. **29**, 386-397 (2008).
32. M. J. Meaney, Maternal care, gene expression, and the transmission of Individual differences
in stress reactivity across generations. *Ann. Rev. Neurosci.* **24**, 1161-1192 (2001).
33. R. M. Kilner *et al.*, Parental effects alter the adaptive value of an adult behavioural trait. *eLife*
4 (2015).
34. I. Meierhofer, H. H. Schwarz, J. K. Müller, Seasonal variation in parental care, offspring
development, and reproductive success in the burying beetle, *Nicrophorus vespillo. Ecol.
Entomol.* **24**, 73-79 (1999).
35. A. Wagner, *Robustness and Evolvability in Living Systems* (Princeton University Press,
Princeton, 2013), vol. 24.
36. J. M. Carlson, J. Doyle, Complexity and robustness. *Proc. Natl. Acad. Sci.* **99 Suppl 1**, 2538-
2545 (2002).

37. M. B. Bonsall, H. Klug, The evolution of parental care in stochastic environments. *J. Evol.*
Biol. **24**, 645-655 (2011).
38. J. C. Creighton, Population density, body size, and phenotypic plasticity of brood size in a
burying beetle. *Behav. Ecol.* **16**, 1031-1036 (2005).
39. S. D. Schrempf, K. W. Burke, J. D. Wettlaufer, P. R. Martin, Behavioral dominance
interactions between *Nicrophorus orbicollis* and *N. tomentosus* burying beetles (Coleoptera:
Silphidae). *PeerJ* **9**, e10797 (2021).
40. B. J. M. Jarrett, M. Schrader, D. Rebar, T. M. Houslay, R. M. Kilner, Cooperative
interactions within the family enhance the capacity for evolutionary change in body size.
Nature Ecol. Evol. **1** (2017).
41. M. P. Scott, Facultative adjustment of the duration of parental care by burying beetles. *J.*
Insect Behav. **11**, 597-603 (1998).
42. S. Trumbo, Monogamy to communal breeding: exploitation of a broad resource base by
burying beetles (*Nicrophorus*). *Ecol. Entomol.* **17**, 289-298 (1992).
43. N. Royle, P. Smiseth, M. Kölliker, *The Evolution of Parental Care* (Oxford University Press,
Oxford, 2012), 10.1093/acprof:oso/9780199692576.003.0001.
44. J. W. Y. Wong, J. Meunier, M. Kollicker, The evolution of parental care in insects: the roles
of ecology, life history and the social environment. *Ecol. Entomol.* **38**, 123-137 (2013).
45. R. C. Lewontin, "Gene, organism, and environment" in *Evolution from Molecules to Men,* D.
S. Bendall, Ed. (Cambridge University Press, Cambridge, 1983).
46. J. A. Oldekop, P. T. Smiseth, H. D. Piggins, A. J. Moore, Adaptive switch from infanticide to
parental care: How do beetles time their behaviour? *J. Evol. Biol.* **20**, 1998-2004 (2007).
47. J. E. Lock, P. T. Smiseth, A. J. Moore, Selection, inheritance, and the evolution of parent-
offspring interactions. *Am. Nat.* **164**, 13-24 (2004).
48. C. M. Rauter, A. J. Moore, Time constraints and trade-offs among parental care behaviours:
effects of brood size, sex and loss of mate. *Anim. Behav.* **68**, 695-702 (2004).
49. D. S. Wilson, W. Knollenberg, J. Fudge, Species packing and temperature dependent
competition among burying beetles (Silphidae, *Nicrophorus*). *Ecol. Entomol.* **9**, 205-216
(1984).
50. C. A. Walling, C. E. Stamper, P. T. Smiseth, A. J. Moore, The quantitative genetics of sex
differences in parenting. *Proc. Natl. Acad. Sci.*, 18430-18435 (2008).

Figure legends

Figure 1. Care influences larval developmental time and size. *(A)* Mosaic plot indicating the percent

of larvae dispersing from the carcass after 7, 8, or 9 days (white, light grey, or dark grey, respectively).

Larvae that received full care developed faster on the carcass than larvae receiving reduced care.

Sample sizes noted within mosaic plot. *(B)* Violin plot of distribution of final larval weight between

individuals that received the full or reduced care treatment (Full care: N = 149, Reduced care: N =

156). Asterisks indicate statistically significant results of contrasts (with P < 0.05). *(C)* Duration of

larval period (in days) from larval hatch until ecolsion, or emergence of adult beetles, was statistically

dependent on larval weight at dispersal and not on care treatment (N = 233). Grey circles and solid line

refer to larvae that received reduced care, while open circles and the dashed line refer to individuals

that received full care.

Figure 2. Parental behavior of females and males affected by the care they received as larvae. *(A)*

Females spent more time on the carcass during the observation period than did males regardless of how

much care they received. In addition, *(B)* both females and males that experienced reduced care

increased the amount of direct care given to their offspring but *(C)* not indirect care. Error bars are

mean ± SEM. N = 15 observations for all treatments except Full care males, where N = 14 observations.

Asterisks indicate statistically significant results of contrasts (with P < 0.05).

Figure 3. Sex of parent influences offspring outcomes but not developmental perturbation. (*A*)

There was no statistically significant effect of caregiver sex or duration of care received by parent on

the length of time larvae spent on the carcass, but there was a significant interaction. *(B)* There was no

statistically significant effect of caregiver sex, the care they received as a larva, or their interaction, on

larval mass. *(C)* Females reared more offspring than did males, but there was no effect of parental

treatment on the number of offspring reared. Namely, the care a parent received as a larva did not result

in negative effects on grandoffspring. Full care males: N = 13 breeding attempts, Full care females: N

= 18, and N = 15 for both Reduced care males and females. Female data are indicated with open circles

or violin plots, while male data are closed circles or violin plots. Error bars are ± SEM. Asterisks

indicate statistically significant results of contrasts (with P < 0.05).

A

B

C

F1

Full *Reduced* *Full* *Reduced*
Female   Male

Parental care treatment