Offspring overcome poor parenting by being better parents

Ahva L. Potticary*, Christopher B. Cunningham, and Allen J. Moore

Department of Entomology, University of Georgia, Athens, Georgia, 30602 USA

*Corresponding Author:

Ahva L. Potticary

120 Cedar Street

Biological Sciences West #360

Athens, GA 30602

ORCID identifier:

Ahva L. Potticary: 0000-0002-1157-5315

Christopher B. Cunningham: 0000-0003-3965-2076

Allen J. Moore: 0000-0002-1498-3322

Article Type:

Letter

Keywords: maternal effect, paternal effect, transgenerational effect

ABSTRACT

The evolutionary repercussions of parental effects are often discussed as fixed effects on offspring

phenotype that can even have negative influences on offspring fitness if parental and offspring

environments do not match. Yet individuals are not passive recipients and may mitigate the persistence

of parental effects through their own behavior. Here we test how the burying beetle *Nicrophorus*

orbicollis, a species with highly developed parental care, responds to poor parenting. We manipulated

the duration of parental care received and measured both offspring and grandoffspring traits of males

and females. As expected, reducing parental care negatively impacted key traits that influence fitness,

including offspring development time and body size. However, offspring that received reduced care

increased uniparental care when they became parents and, in addition, shifted the type of parenting

behavior to more direct feeding of offspring. Parents that had received full care as larvae invested more

time maintaining the food resource, a form of indirect parental care. As a result, both the number and

mass of offspring were unaffected by the different developmental experience of their parent. Our

results show that flexible parental care can overcome poor parental environments and limit negative

parental effects to a single generation.

IMPACT SUMMARY

Parental care is one of the most visible means by which parents influence the development of their

offspring and is expected to evolve to buffer offspring from challenging environments. Yet, variation

in parental care itself can have negative effects on offspring that cascades across generations. How can

offspring overcome these poor developmental environments? Here we test the hypothesis that

organisms can compensate for their own poor developmental environments by changing their parental

care behavior in the burying beetle *Nicrophorus orbicollis*. Specifically, we asked whether receiving

poor parenting begets poor parenting. We manipulated the amount of parental care provided by burying

beetles, an insect that feeds its offspring regurgitated food. Reduced parental care resulted in smaller

offspring that developed more slowly in early developmental stages. However, when these offspring

reached adulthood, they modified their own parental care so that the grandoffspring were unaffected

by the negative conditions experienced by the previous generation. We suggest that these result show

that parental care not only serves as a buffer within-generation but can also compensate for poor natal

environments across generations.

Introduction

Organisms inhabit challenging environments that require accommodation of or buffering against

environmental variation. Which response prevails depends on both an organism's experience of the

current environment and the environments experienced by ancestors. Parental effects are particularly

clear examples of this, as all parents influence the development of their offspring (Roach and Wulff

1987; Mousseau and Fox 1998; Molinier et al. 2006; Marshall and Uller 2007; Royle et al. 2012),

ranging from short-term impacts on offspring to effects that persist for multiple generations (Cheverud

and Moore 1994; Molinier et al. 2006; Yin et al. 2019). As environments can change rapidly or

unpredictably, parental effects on offspring development can range from positive (Badyaev et al. 2006;

Galloway and Etterson 2007; Dantzer et al. 2013; Duckworth et al. 2015) to negative (Gluckman et al.

2005; Sheriff and Love 2013; Potticary and Duckworth 2018) depending on how well offspring

phenotypes match prevailing conditions (Bernardo 1996; Bateson et al. 2014). Consequently, the

evolutionary repercussions of parental effects are often viewed as fixed effects on offspring phenotype,

but this is an assumption that should be tested (Badyaev and Uller 2009).

Individuals are not passive recipients of the environment, and the persistence of a parental effect

depends on the interactions between offspring and their environment in later life (Cheverud and Moore

1994; Bernardo 1996; Francis et al. 2002), which can be directly mediated by individual behavior. One

of the most recognizable parental effects that can impact offspring development is parental care

behavior, and variation in parental care behavior can produce transgenerational effects (Rossiter 1996;

Champagne 2008). Adaptive parental effects are broadly expected to evolve to help offspring overcome

challenging environments through modification of offspring phenotypes or buffering (Wilson 1975;

Mousseau and Fox 1998; Badyaev and Uller 2009). In the former, parents prepare offspring to face a

future environment (Mousseau and Dingle 1991; Dantzer et al. 2013; Potticary and Duckworth 2020)

and are thought to evolve when parental experience is predictive of the environment that will be

experienced by offspring (Bateson et al. 2014; Burgess and Marshall 2014). Alternatively, parents may

buffer offspring from the environment they experienced by changing how they provision offspring

with resources, showing experience-dependent parental care.

Here we test the hypothesis that organisms can compensate for their own poor developmental

environments by changing their parental care behavior in the burying beetle *Nicrophorus orbicollis*.

Specifically, we examine how *N. orbicollis* as parents overcome poor parenting received as larvae. We

investigated this by manipulating the duration of parental care that mixed foster broods received to

produce larvae that experience reduced (six hours) or full parental care (approximately seven days).

Burying beetles require small animal carcasses to breed and show complex parental care behavior

(Robertson 1993; Eggert and Müller 1997; Scott 1998a). Parental care is not a single behavior, but a

category describing a suite of behaviors (Royle et al. 2012). In *Nicrophorus*, parents bury and defend

carcasses, which in part acts to ensure that the carcass is available for longer (Robertson 1993; Trumbo

and Sikes 2021), eliminate bacterial and fungal consumers to avoid putrefaction (Arce et al. 2012;

Haberer et al. 2014), create cavities with predigested carrion "soup" to support offspring self-feeding

(Eggert and Müller 1997; Capodeanu-Nägler et al. 2018; Duarte et al. 2021), and directly transfer food

to larvae through regurgitation of pre-digested carrion (Eggert et al. 1998). Larvae do not feed after

they leave the carcass until they metamorphose into adults, and thus adult body size is largely

determined by the efficiency of resource acquisition during the larval period on carcass. Body size is a

key fitness trait for *Nicrophorus*, as larger beetles are more effective competitors for carcasses (Otronen

1988; Robertson 1993). Thus, parental care creates a direct link between a larva's developmental

experience and competitive ability upon reaching adulthood, and the outcome of this effect will depend

on the ecological context offspring experience in later life.

Our study was designed to ask if poor parenting begets poor parenting. We found that, in *N.*

orbicollis, it does not; parents behaviorally compensated for their own poor rearing environment,

showing experience-dependent parental care and an adaptive parental effect ensuring that the influence

of poor parenting on development does not persist. This suggests that parental care behavior may not

only buffer offspring development within a generation but may also act as a buffer that mitigates the

legacy of environments past.

Materials and Methods

INSECT COLONY AND HUSBANDRY

We used *Nicrophorus orbicollis* from an outbred colony maintained at the University of Georgia that

is regularly supplemented with wild-caught beetles from Whitehall Forest in Athens, GA, USA. All

beetles were maintained in an incubator (Percival, model no. I4-66VL, Perry, IA) at $22 \pm 0.1°C$ on a

14:10 hour light:dark cycle that simulated summer breeding conditions. We bred all individuals used

here in the laboratory to ensure that parentage, age, and to standardize rearing conditions. We housed

beetles individually at larval dispersal in 9-cm-diameter by 4-cm-deep circular compostable deli

containers (EP-RDP5, Eco-Products) filled with ~2.5 cm of organic potting soil. Larvae had no further

social interactions with other burying beetles until they were randomly allocated into treatments. We

defined individuals as age Day 1 on the day of eclosion to adulthood. Beetles were fed *ad libitum*

organic ground beef twice a week following adult eclosion.

EFFECT OF REMOVING PARENTAL CARE ON OFFSPRING DEVELOPMENT

We manipulated the duration of parental care to determine the effect of parental care on offspring traits

related to fitness and their expression of care once they attained adulthood. We randomly mated focal

virgins at Day 13-19 by placing an unrelated male and female in a small plastic container and observed

them to ensure that mating occurred. The pair was then moved to a plastic breeding box filled with

approximately two cm of potting soil moistened with distilled water and containing a freshly thawed

mouse 20.8 ± 0.6 g SD (max = 22.2, min = 20.0; Rodent Pro, Inglefield, IN, U.S.A.). There was no

difference in the size of the mouse provided between treatments (20.9 g in full care treatment, 20.8 g

in reduced care treatment; $F_{1,34} = 0.153$, P = 0.698). We then returned breeding boxes to the incubator

and kept them on a darkened shelf to simulate underground breeding conditions. We checked breeding

boxes for eggs every twenty-four hours. Once eggs we observed eggs, we moved the female and brood

ball to a new breeding box filled with approximately two cm of moistened potting soil, while the male

was removed. We left eggs in the original breeding box with a small piece of ground beef. Once 90

hours had elapsed since egg laying, we checked the breeding box containing the eggs every hour to

determine when larvae had hatched and crawled to the beef. After we observed the focal females'

larvae hatch, we provided the focal female with an unrelated, mixed brood of 10 larvae. Burying beetles

identify their offspring temporally and eat any larvae that arrive too early on the carcass (Oldekop et

al. 2007), and thus we only provided females with larvae after her own larvae had hatched. For the full

duration parental care treatment, we allowed females to associate with larvae until at least 2 of the

larvae had dispersed from the breeding carcass (~7 - 8 days). For the reduced duration parental care

treatment, we allowed females to associate with larvae for 6 hours and then the female was removed.

We checked experimental treatments daily to determine larval dispersal date, subsequently, larval

survival.

We measured several offspring traits related to fitness (Lock et al. 2004), including mass at

dispersal, duration of developmental periods, and adult size. We measured larval mass at dispersal to

the nearest 0.1 mg. We also measured the duration of development on the carcass (from hatch until

dispersal), which is when parental care and all larval growth occurs; the number of days of development

from larval dispersal to pupation; the number of days as pupae (from pupation to adult eclosion), and

total duration of development (from larval hatch until adult eclosion). When the larvae left the carcass

(dispersal), we placed larvae into individual containers with ~2 cm of moistened soil. We checked

treatments daily to determine the number of surviving larvae, length of larval period, pupation date,

and adult ecolsion date. We measured the pronotum length of full and reduced care offspring on Day

14 – 16 post-eclosion with calipers to the nearest 0.01 mm.

EFFECT OF VARIATION IN PARENTAL CARE ENVIRONMENT ON SUBSEQUENT GENERATIONS

At Day 14 – 16 following eclosion, adult beetles that received full or reduced care as offspring were

provided with an unrelated, virgin mate from our stock colony and a 19.6 - 21.2 g thawed mouse carcass

(mean = 20.6 g). These were placed into breeding container with ~2 cm of moistened soil. As variation

of parental care may manifest as variation in investment across the breeding cycle, breeding boxes

were fitted with an escape chamber to allow focal beetles to desert the breeding attempt. Breeding

boxes were checked daily for eggs. Beetles that escaped into the escape chamber prior to egg laying

were replaced in the breeding box. Beetles that escaped after egg-laying and remained in the escape

chamber for over 15 minutes were counted as deserters and removed. Once larvae were observed, the

non-focal partner was removed, and the focal male or female was left in a uniparental context to care

for the offspring. If the focal beetle deserted the breeding attempt prior to larval hatch, no further

monitoring was conducted on that breeding attempt.

As rates of parental provisioning may be affected by the number of larvae (Rauter and Moore

2004), we measured parental care using both behavioral measures and larval traits associated with

fitness. Behavioral observations were conducted twenty-four hours after larva were first observed on

the carcass for each focal individual to quantify parental care. We moved breeding boxes into an

observation room, and we gave the focal beetle at least 30 minutes to acclimate prior to observation.

All observations were conducted under red lights to ensure no light effects on individual care behavior,

as *N. orbicollis* are nocturnal (Wilson et al. 1984). In each behavioral observation, we recorded both

direct and indirect parental care behaviors for 15 minutes (Walling et al. 2008). Direct care is defined

as when the male or female fed larvae. Indirect care is defined as activities related to carcass cleaning

and maintenance, which allows for the preservation of the carcass for future offspring provisioning and

thus represents a type of parental care without direct interactions with offspring. We also recorded time

spent showing "no care", defined as when the parent was on the carcass but not engaged in direct or

indirect care, or the individual was off the carcass. Behaviors were recorded continuously as both a

duration and frequency (e.g., number of times an adult fed larvae). This gave each individual three

scores (direct care, indirect care, and no care) which together summed to 15 minutes.

We measured offspring traits related to fitness reared by males or females that had received full

or reduced care as larvae themselves. We did not manipulate the number of offspring but instead

allowed parents to cull the number without interference. Of the remaining 61 uniparental broods, we

measured the number of offspring reared, the mass of the offspring at dispersal, and the amount of time

larvae spent on the carcass from hatching to dispersal. Measurements of development times and larval

mass were the same as used in the experimental manipulation of the duration of parental care (see

previous section).

STATISTICAL ANALYSES

We performed all statistical analyses using JMP Pro (v. 16.0.0, http://jmp.com) and produced figures

in SigmaPlot (v. 14.5, http://www.sigmaplot.co.uk). Results are means ± SE unless otherwise noted.

Effects on continuous offspring traits were analyzed with nested ANOVA, with family nested within

treatment. We analyzed categorical data with Pearson's Chi-square goodness-of-fit test. We analyzed

parental care behaviors and continuous larval traits of grand offspring with two-way ANOVA or, a

GLM with a Poisson distribution for categorical data, with caretaker sex and parenting experienced as

factors.

Results

EFFECT OF REDUCING PARENTAL CARE ON OFFSPRING DEVELOPMENT

We set up 36 families, 18 in each treatment, of mixed-parentage broods with a single, unrelated female

caretaker in the full and reduced parental care treatments. Overall larval survival was high for both

treatments; from an initial 10 fostered larvae placed on carcass, 3-10 survived to dispersal from the

mouse with a mean 8.3 ± 0.4 surviving larvae for full care families and 8.7 ± 0.4 for reduced care ($F_{1,34}$

$= 0.430$, P $= 0.516$). Three (2%) larvae died after dispersal from the full care treatment, and three

(1.9%) died from the reduced care treatment. A further 14 (9.6%) died in the pupal stage in the full

care treatment, and 16 (10.4%) in the reduced care.

Larvae in the full care treatment developed faster on the carcass than larvae that received

reduced care ($\chi^2 = 18.86$, P < 0.0001, N $= 305$). Larvae that received full care completed development

from hatch to dispersal in 7 - 8 days (mean $= 7.4$) and reduced care larvae dispersed in 8 - 9 days (mean

$= 8.3$ in reduced care families; **Fig. 1*A*)**. The largest effect of reduced care was on larval weight at

dispersal from the carcass (**Fig. 1*B***). There was statistically significant variation between care

treatments ($F_{1,269} = 433.253$, P < 0.0001), as well as caretakers nested within treatments ($F_{34,269} =$

10.155, P < 0.0001). Controlling for variation amongst families, larvae that received full care weighed

500.4 ± 4.4 mg, whilst larvae that received reduced care weighed 373.4 ± 4.2 mg. Larval weight at

dispersal was highly and positively correlated with adult size ($r = 0.91$, N $= 248$).

234  Even though larvae in the full care treatment developed faster on the carcass, duration of total

235 post-embryonic development (hatch to eclosion) was influenced primarily by larval mass at dispersal

236 ($F_{1,232} = 77.283$, P < 0.0001) and not on care treatment ($F_{1,232} = 1.311$, P = 0.2535; **Fig. 1C**). There was

237 statistically significant variation attributable to caretaker ($F_{34,261} = 9.301$, P < 0.0001), larval weight

238 ($F_{1,261} = 34.501$, P < 0.0001) and duration of parental care received ($F_{1,261} = 13.47$, P = 0.0003) on time

239 spent after dispersal to pupation. The same factors influenced duration of pupal development (pupae

240 to adult eclosion) with statistically significant variation arising from larval weight ($F_{1,231} = 55.586$, P

241 < 0.0001), caretaker ($F_{34,231} = 3.173$, P < 0.0001), but not duration of parental care received ($F_{1,231} =$

242 0.883, P = 0.348). Thus, the duration of care received influenced the speed with which the larvae

243 developed on the carcass, but this only shifted development time.

245 **EFFECT OF VARIATION IN PARENTAL CARE ENVIRONMENT ON SUBSEQUENT GENERATIONS**

247 We next established 89 uniparental broods using the resulting offspring from the reduced and full care

248 treatments (26 full care males, 20 full care females, 24 reduced care males, and 17 reduced care

249 females). There is no evidence that the type of parenting received influenced whether parents

250 abandoned prior to showing parental care to larvae. Focal male caretakers were more likely to abandon

251 before eggs hatched (reduced care males: 11/24 breeding attempts; full care males: 11/26 breeding

252 attempts) than female caretakers (reduced care females: 2/17 breeding attempts; full care females 2/20

253 breeding attempts). Once larvae hatched, most focal parents stayed for the entire larval development

254 on the carcass. When caretakers did abandon larvae, it was almost exclusively males (10/29 males,

255 1/33 females; $\chi^2 = 11.640$, df = 1, P = 0.0012). The one female that abandoned early had received full

256 care; for males, four had received full care and six received reduced care.

257  Parental care behavior of both males and females was affected by the length of care they

258 received as larvae. There are strong differences between uniparental females and males in this species

(**Fig. 2**). Females spent more time on the carcass during the observation period than did males ($F_{1,55} =$

17.610, P < 0.001; **Fig. 2A**) regardless of the duration of parental care they received ($F_{1,55} = 1.878$, P

= 0.176) and there was not a statistically significant interaction between the duration of care received

and sex of the caregiver on amount of time spent on the carcass ($F_{1,55} = 1.332$, P = 0.253). Direct care,

or the time parents spent regurgitating to their larvae, was strongly influenced by the duration of

parental care received ($F_{1,55} = 6.761$, P = 0.012); individuals that received reduced care provided more

direct care themselves and this pattern was consistent for both males and females (**Fig. 2B**); there was

no interaction between the sex of the caregiver and the effect of parenting received ($F_{1,55} = 0.111$, P =

0.741). Females provided more direct care than males under both treatments ($F_{1,55} = 12.699$, P =

0.0008). In contrast, indirect care, or time spent maintaining the carcass, was not affected by the

duration of parental care received ($F_{1,55} = 1.181$, P = 0.282; **Fig. 2C**) or sex of the caregiver ($F_{1,55} =$

3.072, P = 0.085), and there was no interaction between these treatments ($F_{1,55} = 0.705$, P = 0.405).

Overall, the care a parent received as a larva did not result in negative effects on grandoffspring.

Using GLM with Poisson distribution, there was no statistically significant effect of caregiver sex (χ^2

= 0.269, df = 1, P = 0.604), care received by the parent ($\chi^2 = 0.007$, P = 0.932) or their interaction (χ^2

= 0.439, P = 0.508). Carcass mass did not have a statistically significant influence ($\chi^2 = 0.118$, P =

0.731). There was no effect of caregiver sex on average larval mass ($F_{1,56} = 1.855$, P = 0.179), care

received by parent ($F_{1,56} = 1.340$, P = 0.252) or their interaction ($F_{1,56} = 0.477$, P = 0.493), controlling

for carcass mass as a covariate ($F_{1,56} = 0.047$, P = 0.829; **Fig. 3A**). Females reared more offspring than

did males in general ($F_{1,56} = 15.521$, P = 0.0002), but there was no effect of parental treatment on the

number of offspring reared ($F_{1,56} = 0.095$, P = 0.759; **Fig. 3B**) and no interaction between caregiver

sex and the duration of parental care received ($F_{1,56} = 0.429$, P = 0.515). Carcass mass as a covariate

was also not statistically significant ($F_{1,56} = 1.562$, P = 0.217).

Discussion

Adaptive parental effects are expected to evolve to help offspring overcome challenging environments

through modification of offspring phenotypes or buffering (Wilson 1975; Mousseau and Fox 1998;

West-Eberhard 2003; Badyaev and Uller 2009), yet parents can have negative impacts on offspring

(Rauter and Moore 2002; Royle et al. 2012) especially if parental and offspring environments do not

match (Marshall and Uller 2007; Sheriff and Love 2013; Bateson et al. 2014). We tested the hypothesis

that parental care can act as a buffer because it allows parents to compensate for their own poor

developmental environment when they become parents themselves. Experimentally reducing parental

care greatly influenced offspring traits that are aligned with fitness, including duration of the vulnerable

stage on the carcass (**Fig. 1***A*; Robertson 1993; Eggert & Müller 1997) and body size (**Fig. 1***B*; Otronen

1988). However, there was no difference in the size or number of offspring when those same offspring

themselves were parents. This consistency of offspring outcomes was enabled by a shift in the

expression of the components of parental care behavior. Parents that received reduced parental care

themselves spent more time directly transferring nutrients to offspring, as opposed to parents that

received full care, who invested more in indirect mechanisms to enable nutrient acquisition of larvae

(i.e., maintenance of a food resource). As a result, parents that received reduced parental care

constrained the negative effects of their own poor developmental environment to a single generation.

Together, these data suggest that parents behaviorally compensate for their own poor rearing

environment, showing experience-dependent parental care and an adaptive parental effect ensuring that

poor developmental environments do not persist.

Beetles that received full parental care were larger and developed faster on the carcass, although

larger beetles overall took longer to develop into adult beetles. Why might development on the carcass

be different relative to other developmental periods? On a local level, developmental rate can be

directly influenced by variation in parental care. Parents regurgitate to larvae, and direct provisioning has been shown to increase offspring size or growth rate across taxa (Naef-Daenzer and Keller 1999; Lock et al. 2004; Lock 2012). The carcass is the only source of nutrition for both larvae and parents during early larval development (Keppner et al. 2023). In other *Nicrophorus* species, multiple parental behaviors increase larval growth and survival, such as reducing exploitation competition by removing bacteria (Cotter and Kilner 2010; Arce et al. 2012), selective culling of the brood (Bartlett 1987; Bartlett and Ashworth 1988), and maintenance of a larval cavity in the carcass (Capodeanu-Nägler et al. 2018; Duarte et al. 2021). An ultimate explanation for rapid development on the carcass may be that there is selection arising from competition or predation for offspring to leave the carcass quickly. Studies in the wild find that *N. orbicollis* nests are frequently discovered by scavengers or taken over by conspecifics, which generally eat larvae and eggs during successful takeovers (Robertson 1993). High predation risk has been associated with more rapid development rates in the nest across bird species (Martin et al. 2011). Thus, we suggest that local provisioning rates serve to enhance offspring growth rates while on the carcass, reducing the risk at this vulnerable stage, while still allowing larvae to attain large adult size.

Parental effects range from short-term effects that dissipate when offspring pass a certain developmental stage (Gluckman et al. 2005), to effects that persist for multiple generations, even for the same traits (e.g., stress reactivity; Meaney 2001; Mustoe et al. 2014; Sheriff et al. 2017). Previous work in other *Nicrophorus* species and other taxa have documented such transgenerational effects of poor parental care environments (e.g., Meaney 2001; Champagne 2008; Lock 2012; Kilner et al. 2015). An unanswered question is why the effects of poor developmental environment are propagated across generations under some circumstances but not in others. The persistence of parental effects likely depends both abiotic and biotic factors, such as the durability of physiological mechanisms that transmit them (Weaver et al. 2004; Horsthemke 2018) and concordance between the parent and offspring environments (Gluckman et al. 2005; Marshall and Uller 2007; Bateson et al. 2014).

Moreover, the persistence of parental effects likely depends on whether they arise from an evolved interaction between parents and offspring development, or are merely a disruption of normal development (Gluckman et al. 2005). Adaptive parental effects are predicted to induce variation in offspring traits that are functionally related to the environment that triggered them (e.g., Tollrian 1995; Duckworth et al. 2015; Sheriff et al. 2017; Sharda et al. 2021). In our study, *N. orbicollis* that received poor parenting changed their own parental care behavior to reduce the influence of a poor developmental environment, which also requires that offspring are receptive to changes in parental care behavior (Smiseth et al. 2003). Together, these data imply that parental care acts a developmental buffer across generations and may indicate evolution for this purpose in *N. orbicollis,* particularly given how important body size is for competitive interactions in this species (Robertson 1993; Creighton 2005; Schrempf et al. 2021). While there are many processes that may alter the persistence of parental effects, the results of this study suggest that changes in parental care enables the reversal of poor developmental environments.

Parental care, and behavior more broadly, is often hypothesized to act as a buffer on both ecological and evolutionary timescales (Huey et al. 2003; Duckworth 2009; Muñoz and Losos 2018). Parental care strategies show tremendous variation across taxa (Brockmann 1997; Scott 1998a; Mank et al. 2005; Cockburn 2006; Benowitz et al. 2016; Potticary and Duckworth 2021) and have arisen across a wide diversity of ecological conditions (Wilson 1975; Jetz and Rubenstein 2011). Our results may provide one explanation for why there can be so much variation in parental care (e.g., Meierhofer et al. 1999) given its importance to offspring development (Royle et al. 2016). It has been broadly suggested that variable environments are expected to produce systems that have greater flexibility, in order to maintain consistent function across changing environments (Kitano 2004; Kashtan and Alon 2005; Kashtan et al. 2007; Chenard and Duckworth 2021). That is, environments that are challenging in many ways may require flexible parental care to maintain consistent function, such as large offspring body size, across environments. *N. orbicollis* live in highly heterogenous and variable environments

(Collard et al. 2021; Wettlaufer et al. 2023; Potticary et al. in press), larger beetles win access to

carcasses (Robertson 1993; Creighton 2005; Schrempf et al. 2021), and larger *Nicrophorus* species are

more likely to show obligate care (Jarrett et al. 2017). Together, this may indicate that the great

variation observed both in how individual burying beetles show care (Scott 1998b; Meierhofer et al.

1999) and who shows care (e.g., uniparental to biparental to communal; Trumbo 1992; Scott 1998a;

Benowitz et al. 2016) may emerge from selection for consistency of offspring phenotypes relative to

the prevailing environmental context (Creighton 2005).

Organisms are not passive recipients of the environment, but rather influence their environment

in the short term and evolve in response to social and ecological conditions in the long term (Lewontin

1983). We found that removing parental care induced strong changes in traits associated with fitness.

Offspring that received poor parenting responded by increasing uniparental care when they became

parents and shifting the type of nutrient-transfer behavior they showed, resulting in the amelioration of

poor developmental environments experienced by parents. Parenting may be an effective evolutionary

mechanism for buffering poor past and present environments, including those imposed during the

parents' development. This buffering may be enabled by dynamic changes in the units of parental care

– the component behaviors themselves.

ACKNOWLEDGEMENTS AND FUNDING SOURCES

We thank Kathryn Kollars for assistance constructing escape chambers, insect colony care, and for

developing the beetle graphics. We also thank Trish Moore and Hans Otto for discussions and/or

feedback on the manuscript, and Emily Shelby for assistance with figures produced by R. The authors

declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

ALP and AJM conceived of study and developed methods. ALP, AJM, and CBC collected life history

data. ALP conducted behavioural observations. AJM performed statistical analyses. ALP and AJM

created figures. ALP, AJM, and CBC all contributed to writing and editing the manuscript. ALP was

funded by a USDA cooperative agreement to AJM.

DATA ACCESSIBILITY

The data underlying this article are available in the Dryad Digital Repository (doi: pending).

References

Arce, A. N., P. R. Johnston, P. T. Smiseth, and D. E. Rozen. 2012. Mechanisms and fitness effects of
antibacterial defences in a carrion beetle. J. Evol. Biol. 25:930-937.
Badyaev, A. V., T. L. Hamstra, K. P. Oh, and D. A. Acevedo Seaman. 2006. Sex-biased maternal
effects reduce ectoparasite-induced mortality in a passerine bird. Proc. Natl. Acad. Sci.
103:14406-14411.
Badyaev, A. V. and T. Uller. 2009. Parental effects in ecology and evolution: mechanisms, processes
and implications. Phil. Trans. Roy. Soc. B. 364:1169-1177.
Bartlett, J. 1987. Filial cannibalism in burying beetles. Behav. Ecol. Sociobiol. 21:179-183.
Bartlett, J. and C. M. Ashworth. 1988. Brood size and fitness in *Nicrophorus vespilloides*
(Coleoptera:Silphidae). Behav. Ecol. Sociobiol. 22:429-434.
Bateson, P., P. Gluckman, and M. Hanson. 2014. The biology of developmental plasticity and the
Predictive Adaptive Response hypothesis. J. Physiol. 592:2357-2368.
Benowitz, K. M., E. C. McKinney, and A. J. Moore. 2016. Difference in parenting in two species of
burying beetle, *Nicrophorus orbicollis* and *Nicrophorus vespilloides*. J. Ethol. 34:315-319.
Bernardo, J. 1996. Maternal effects in animal ecology. Am. Zool. 36:83-105.
Brockmann, H. J. 1997. Cooperative breeding in wasps and vertebrates: the role of ecological
constraints. Pp. 347-371 *in* J. C. Choe, and B. J. Crespi, eds. The Evolution of Social
Behavior in Insects and Arachnids. Cambridge University Press, Cambridge.
Burgess, S. C. and D. J. Marshall. 2014. Adaptive parental effects: the importance of estimating
environmental predictability and offspring fitness appropriately. Oikos 123:769-776.
Capodeanu-Nägler, A., M. A. Prang, S. T. Trumbo, H. Vogel, A.-K. Eggert, S. K. Sakaluk, and S.
Steiger. 2018. Offspring dependence on parental care and the role of parental transfer of oral
fluids in burying beetles. Front. Zool. 15:33.

Champagne, F. A. 2008. Epigenetic mechanisms and the transgenerational effects of maternal care. Front. Neuroendocrinol. 29:386-397.

Chenard, K. and R. Duckworth. 2021. The special case of behavioral plasticity? Pp. 301-325 *in* D. W. Pfennig, ed. *Phenotypic Plasticity & Evolution: Causes, Consequences, and Controversies*. CRC Press, Boca Raton.

Cheverud, J. M. and A. Moore. 1994. Quantitative genetics and the role of the environment provided by relatives in behavioral evolution. Pp. 68-100 *in* C. Boake, ed. Quantitative genetic studies of behavioral evolution. University of Chicago Press, Chicago.

Cockburn, A. 2006. Prevalence of different modes of parental care in birds. Proc. Biol. Sci. 273:1375-1383.

Collard, A. E., J. D. Wettlaufer, K. W. Burke, D. V. Beresford, and P. R. Martin. 2021. Body size variation in a guild of carrion beetles. Can. J. Zool. 99:117-129.

Cotter, S. C. and R. M. Kilner. 2010. Sexual division of antibacterial resource defence in breeding burying beetles, *Nicrophorus vespilloides*. J. Anim. Ecol. 79:35-43.

Creighton, J. C. 2005. Population density, body size, and phenotypic plasticity of brood size in a burying beetle. Behav. Ecol. 16:1031-1036.

Dantzer, B., A. E. M. Newman, R. Boonstra, R. Palme, S. Boutin, M. M. Humphries, and A. G. McAdam. 2013. Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal. Science 340:1215-1217.

Duarte, A., D. Rebar, A. C. Hallett, B. J. M. Jarrett, and R. M. Kilner. 2021. Evolutionary change in the construction of the nursery environment when parents are prevented from caring for their young directly. Proc. Natl. Acad. Sci. 118:e2102450118.

Duckworth, R. A. 2009. The role of behavior in evolution: a search for mechanism. Evol. Ecol. 23:513-531.

Duckworth, R. A., V. Belloni, and S. R. Anderson. 2015. Cycles of species replacement emerge from locally induced maternal effects on offspring behavior in a passerine bird. Science 347:875-877.

Eggert, A.-K. and J. K. Müller. 1997. Biparental care and social evolution in burying beetles: lessons from the larder. Pp. 216 *in* J. C. Choe, and B. J. Crespi, eds. *The Evolution of Social Behaviour in Insects and Arachnids*. Cambridge University Press, Cambridge.

Eggert, A.-K., M. Reinking, and J. K. Müller. 1998. Parental care improves offspring survival and growth in burying beetles. Anim. Behav. 55:97-107.

Francis, D. D., J. Diorio, P. M. Plotsky, and M. J. Meaney. 2002. Environmental enrichment reverses the effects of maternal separation on stress reactivity. J. Neurosci. 22:7840-7843.

Galloway, L. F. and J. R. Etterson. 2007. Transgenerational plasticity is adaptive in the wild. Science 318:1134-1136.

Gluckman, P. D., M. A. Hanson, H. G. Spencer, and P. Bateson. 2005. Environmental influences during development and their later consequences for health and disease: implications for the interpretation of empirical studies. Proc. Roy. Soc. B. 272:671-677.

Haberer, W., S. Steiger, and J. K. Müller. 2014. Dynamic changes in volatile emissions of breeding burying beetles. Physiol. Entomol. 39:153-164.

Horsthemke, B. 2018. A critical view on transgenerational epigenetic inheritance in humans. Nat. Comm. 9:2973.

Huey, R. B., P. E. Hertz, and B. Sinervo. 2003. Behavioral drive versus behavioral inertia in evolution: a null model approach. Am. Nat. 161:357-366.

Jarrett, B. J. M., M. Schrader, D. Rebar, T. M. Houslay, and R. M. Kilner. 2017. Cooperative interactions within the family enhance the capacity for evolutionary change in body size. Nature Ecol. Evol. 1:0178.

Jetz, W. and D. R. Rubenstein. 2011. Environmental uncertainty and the global biogeography of cooperative breeding in birds. Curr. Biol. 21:72-78.

Kashtan, N. and U. Alon. 2005. Spontaneous evolution of modularity and network motifs. Proc. Natl. Acad. Sci. 102:13773-13778.

Kashtan, N., E. Noor, and U. Alon. 2007. Varying environments can speed up evolution. Proc. Natl. Acad. Sci. 104:13711-13716.

Keppner, E. M., M. Laubenthal, M. A. Prang, T. Conrad, and S. Steiger. 2023. Harsh nutritional environment has positive and negative consequences for family living in a burying beetle. Ecol. Evol. 13:e9699.

Kilner, R. M., G. Boncoraglio, J. M. Henshaw, B. J. M. Jarrett, O. De Gasperin, A. Attisano, and H. Kokko. 2015. Parental effects alter the adaptive value of an adult behavioural trait. eLife 4.

Kitano, H. 2004. Biological robustness. Nature Rev. Gen. 5:826-837.

Lewontin, R. C. 1983. Gene, organism, and environment *in* D. S. Bendall, ed. *Evolution from Molecules to Men*. Cambridge University Press, Cambridge.

Lock, J. E. 2012. Transgenerational effects of parent and grandparent gender on offspring development in a biparental beetle species. Biol. Lett. 8:408 - 411.

Lock, J. E., P. T. Smiseth, and A. J. Moore. 2004. Selection, inheritance, and the evolution of parent-offspring interactions. Am. Nat. 164:13-24.

Mank, J. E., D. E. L. Promislow, and J. C. Avise. 2005. Phylogenetic perspectives in the evolution of parental care in ray-finned fishes Evolution 59:1570-1578.

Marshall, D. J. and T. Uller. 2007. When is a maternal effect adaptive? Oikos 116:1957-1963.

Martin, T. E., P. Lloyd, C. Bosque, D. C. Barton, A. L. Biancucci, Y.-R. Cheng, and R. Ton. 2011. Growth rate variation among passerine species in tropical and temperate sites: an antagonistic interaction between parental food provisioning and nest predation risk Evolution 65:1607-1622.

Meaney, M. J. 2001. Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations. Ann. Rev. Neurosci. 24:1161-1192.

Meierhofer, I., H. H. Schwarz, and J. K. Müller. 1999. Seasonal variation in parental care, offspring development, and reproductive success in the burying beetle, *Nicrophorus vespillo*. Ecol. Entomol. 24:73-79.

Molinier, J., G. Ries, C. Zipfel, and B. Hohn. 2006. Transgeneration memory of stress in plants. Nature 442:1046-1049.

Mousseau, T. A. and H. Dingle. 1991. Maternal effects in insect life histories. Ann. Rev. Entomol. 36:511-534.

Mousseau, T. A. and C. W. Fox. 1998. Maternal effects as adaptations. Oxford University Press, Oxford.

Muñoz, M. M. and J. B. Losos. 2018. Thermoregulatory behavior simultaneously promotes and forestalls evolution in a tropical lizard. Am. Nat. 191:E15-E26.

Mustoe, A. C., J. H. Taylor, A. K. Birnie, M. C. Huffman, and J. A. French. 2014. Gestational cortisol and social play shape development of marmosets' HPA functioning and behavioral responses to stressors. Dev. Psychobiol. 56:1229-1243.

Naef-Daenzer, B. and L. F. Keller. 1999. The foraging performance of great and blue tits (*Parus major* and *P. caeruleus*) in relation to caterpillar development, and its consequences for nestling growth and fledging weight. J. Anim. Ecol. 68:708-718.

Oldekop, J. A., P. T. Smiseth, H. D. Piggins, and A. J. Moore. 2007. Adaptive switch from infanticide to parental care: how do beetles time their behaviour? J. Evol. Biol. 20:1998-2004.

Otronen, M. 1988. The effect of body size on the outcome of fights in burying beetles (*Nicrophorus*). Ann. Zool. Fenn. 25:191-201.

Potticary, A. L. and R. A. Duckworth. 2018. Environmental mismatch results in emergence of cooperative behavior in a passerine bird. Evol. Ecol. 32:215-229.

Potticary, A. L. and R. A. Duckworth. 2020. Multiple environmental stressors induce an adaptive maternal effect. Am. Nat. 196:487-500.

Potticary, A. L. and R. A. Duckworth. 2021. A neuroendocrine perspective on the origin and evolution of cooperative breeding. Ornithol. 138.

Potticary, A. L., H. W. Otto, J. V. McHugh, and A. J. Moore. in press. Spatiotemporal variation in the competitive environment, with implications for how climate change may affect a species with parental care. Ecol. Evol.

Rauter, C. M. and A. J. Moore. 2002. Quantitative genetics of growth and development time in the burying beetle *Nicrophorus pustulatus* in the presence and absence of post-hatching parental care. Evolution 56:96-110.

Rauter, C. M. and A. J. Moore. 2004. Time constraints and trade-offs among parental care behaviours: effects of brood size, sex and loss of mate. Anim. Behav. 68:695-702.

Roach, D. A. and R. D. Wulff. 1987. Maternal effects in plants. Ann. Rev. Ecol. Syst. 18:209-235.

Robertson, I. C. 1993. Nest intrusions, infanticide, and parental care in the burying beetle, *Nicrophorus orbicollis* (Coleoptera: Silphidae). J. Zool. 231:583-593.

Rossiter, M. 1996. Incidence and consequences of inherited environmental effects. Ann. Rev. Ecol. Syst. 27:451-476.

Royle, N., P. Smiseth, and M. Kölliker. 2012. *The Evolution of Parental Care*. Oxford University Press, Oxford.

Royle, N. J., S. H. Alonzo, and A. J. Moore. 2016. Co-evolution, conflict and complexity: what have we learned about the evolution of parental care behaviours? Curr. Op. Behav. Sci. 12:30-36.

Schrempf, S. D., K. W. Burke, J. D. Wettlaufer, and P. R. Martin. 2021. Behavioral dominance interactions between *Nicrophorus orbicollis* and *N. tomentosus* burying beetles (Coleoptera: Silphidae). PeerJ 9:e10797.

Scott, M. 1998a. The ecology and behavior of burying beetles. Ann. Rev. Entomol. 43:595-618.

Scott, M. P. 1998b. Facultative adjustment of the duration of parental care by burying beetles. J. Insect Behav. 11:597-603.

Sharda, S., T. Zuest, M. Erb, and B. Taborsky. 2021. Predator-induced maternal effects determine adaptive antipredator behaviors via egg composition. Proc. Natl. Acad. Sci. 118.

Sheriff, M. J., A. Bell, R. Boonstra, B. Dantzer, S. G. Lavergne, K. E. McGhee, K. J. MacLeod, L. Winandy, C. Zimmer, and O. P. Love. 2017. Integrating ecological and evolutionary context in the study of maternal stress. Integ. Comp. Biol. 57:437-449.

Sheriff, M. J. and O. P. Love. 2013. Determining the adaptive potential of maternal stress. Ecol. Lett. 16:271-280.

Smiseth, P. T., C. T. Darwell, and A. J. Moore. 2003. Partial begging: an empirical model for the early evolution of offspring signalling. Proc. Royal Soc. B. 270:1773-1777.

Tollrian, R. 1995. Predator-induced morphological defenses: costs, life history shifts, and maternal effects in *Daphnia pulex*. Ecology 76:1691-1705.

Trumbo, S. 1992. Monogamy to communal breeding: exploitation of a broad resource base by burying beetles (*Nicrophorus*). Ecol. Entomol. 17:289-298.

Trumbo, S. T. and D. S. Sikes. 2021. Resource concealment and the evolution of parental care in burying beetles. J. Zool. 315:175-182.

Walling, C. A., C. E. Stamper, P. T. Smiseth, and A. J. Moore. 2008. The quantitative genetics of sex differences in parenting. Proc. Natl. Acad. Sci. 105:18430-18435.

Weaver, I. C. G., N. Cervoni, F. A. Champagne, A. C. D'Alessio, S. Sharma, Seckl, Jr., S. Dymov, M. Szyf, and M. J. Meaney. 2004. Epigenetic programming by maternal behavior. Nature Neurosci. 7:847-854.

West-Eberhard, M. 2003. Developmental plasticity and evolution. Oxford University Press, New
York.
Wettlaufer, J. D., A. Ye, H. A. MacMillan, and P. R. Martin. 2023. A test of the competitive ability–
cold tolerance trade-off hypothesis in seasonally breeding beetles. Ecol. Entomol. 48:55-68.
Wilson, D. S., W. G. Knollenberg, and J. Fudge. 1984. Species packing and temperature dependent
competition among burying beetles (Silphidae, *Nicrophorus*). Ecol. Entomol. 9:205-216.
Wilson, E. O. 1975. *Sociobiology: the new synthesis*. Harvard University Press, Cambridge.
Yin, J., M. Zhou, Z. Lin, Q. Q. Li, and Y.-Y. Zhang. 2019. Transgenerational effects benefit
offspring across diverse environments: a meta-analysis in plants and animals. Ecol. Lett.
22:1976-1986.

Figure legends

Figure 1. Care influences larval developmental time and size. *(A)* Mosaic plot indicating the percent

of larvae dispersing from the carcass after 7, 8, or 9 days (white, light grey, or dark grey, respectively).

Larvae that received full care developed faster on the carcass than larvae receiving reduced care.

Sample sizes noted within mosaic plot. *(B)* Violin plot of distribution of final larval weight between

individuals that received the full or reduced care treatment (Full care: N = 149, Reduced care: N =

156). Asterisks indicate statistically significant contrasts (P < 0.05). *(C)* Duration of larval period (in

days) from larval hatch until ecolsion, or emergence of adult beetles, was statistically dependent on

larval weight at dispersal and not on care treatment (N = 233). Grey circles and solid line refer to larvae

that received reduced care, while open circles and the dashed line refer to individuals that received full

care.

Figure 2. Parental behavior of females and males affected by the care they received as larvae. *(A)*

Females spent more time on the carcass during the observation period than did males regardless of how

much care they received. In addition, *(B)* both females and males that experienced reduced care

increased the amount of direct care given to their offspring but *(C)* not indirect care. Error bars are

mean ± SEM. N = 15 observations for all treatments except Full care males, where N = 14 observations.

Asterisks indicate statistically significant contrasts (P < 0.05).

Figure 3. Sex of parent influences offspring outcomes but not the parental developmental

environment. *(A)* Violin plot of average larval mass for female (open violin) or male parents (closed

violin) that received full or reduced care. There was no statistically significant effect of caregiver sex,

the care they received as a larva, or their interaction, on larval mass. (***B***) Females reared more offspring

than did males, but there was no effect of parental treatment on the number of offspring reared. Namely,

the care a parent received as a larva did not result in negative effects on grandoffspring. Full care males:

N = 13 breeding attempts, full care females: N = 18, and N = 15 for both reduced care males and

females. Female data are indicated with open circles, while male data are closed circles. Error bars are

± SEM. Asterisks indicate statistically significant contrasts (P < 0.05).

Parental care environment experienced by parent