Highly defended nudibranchs do not 'escape' to visually diverse background habitats

Cedric P. van den Berg[1†], John A. Endler[2], Karen L. Cheney[1]

[1]School of Biological Sciences, The University of Queensland, 4072 Brisbane, Australia

[2]Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Geelong,
Victoria 3216, Australia

Corresponding author: Cedric P. van den Berg

Email: c.vandenberg@uq.edu.au

ORCID ID: 0000-0001-6422-7237

[†]Current address: School of Biological Sciences, The University of Bristol, BS81TQ, United Kingdom

Lay Summary

According to the 'escape and radiate' hypothesis, potent chemical defences are thought to enable
animals to be less dependent on matching visual backgrounds for camouflage. We investigated this
idea in 13 species of nudibranch molluscs and found strong evidence against this hypothesis. In fact,
species with strong chemical defences were found on up to 56.9% more uniform backgrounds. This
suggests that warning colouration in nudibranchs might be specialised for specific visual backgrounds.

Abstract

The 'escape and radiate' hypothesis predicts that once species have evolved aposematism, defended
species can utilise more visually diverse background habitats as they 'escape' the need to be well
camouflaged. This enables species to explore new ecological niches, resulting in increased
diversification rates. To test the 'escape' component of this hypothesis, we examined whether the
background habitats of 13 nudibranch mollusc species were more variable for highly defended species
(in terms of unpalatability), compared to undefended (palatable) species. We obtained 157 colour
pattern statistics from the Quantitative Colour Pattern Analysis (QCPA) framework to analyse
backgrounds viewed through the eyes of a potential predator (triggerfish, *Rhinecanthus aculeatus*).
This was done at viewing distances simulating an escalating predation sequence. Contrary to the
hypothesis, we found no significant positive correlations between the diversity of visual backgrounds
and the strength of chemical defences, but instead found numerous significant negative correlations.
Highly defended species were viewed against backgrounds up to 56.9% less variable than non-
defended species. Furthermore, the strongest correlations were found at viewing distances coinciding

with likely prey detection and identification. Our results provide empirical evidence against

background generalisation in aposematic nudibranch species. Instead, we suggest that less variable

visual backgrounds may strengthen aposematic signalling.

Keywords

Aposematism, camouflage, defensive animal colouration, escape and radiate hypothesis, signal

honesty, visual ecology, visual modelling, warning signals

Introduction

Animals and plants use aposematic colour patterns for advertising defences to potential

predators (Poulton, 1890). One of the hypothesised benefits of an organism evolving aposematism is

the potential to 'escape' the costs of crypsis, including limited movement and thus access to resources

due to the need to match a specific background habitat or preventing detection through motion

contrast for primary defence (Endler, 1984; Merilaita & Tullberg, 2005; Regan & Beverley, 1984).

Therefore, Merilaita & Tullberg (2005) suggested that alternative visual defence strategies, such as

aposematism, are more likely to evolve in highly variable visual environments, where crypsis is difficult

to achieve. The evolution of aposematic signalling then provides increased access to resources across

ecological niches as individuals are not restricted to particular habitats to avoid predator detection.

This reduced dependency on crypsis may then drive increased rates of speciation and diversification

(Arbuckle & Speed, 2015; Ehrlich & Raven, 1964). This process was coined 'escape and radiate' by

Thompson (1989) and has since been adopted as a core concept underpinning the importance of

chemical defences for adaptive radiations (e.g. Arbuckle & Harris, 2021).

Theoretical modelling supports the conclusion that aposematic species are less constrained

by the visual properties of their habitat (habitat generalists) than undefended background-matching

species (habitat specialists) (Merilaita et al., 1999; Speed et al., 2010). However, empirical evidence

for this in natural systems across multiple species is limited. Here, we tested the hypothesis that more

defended species inhabit more variable habitats using 13 species of co-occurring Eastern Australian

nudibranch species (see Fig. S1 for images of all species). Nudibranchs are an intriguing system for the

study of defensive animal colouration due to the presence of potent chemical defences in many

species (Avila, 1995; Winters et al., 2018, 2022), and colour patterns that range from bold aposematic

displays to near-perfect camouflage (Debelius & Kuiter, 2007). Indeed, the evolution of specialised

glands (mantle dermal formations) for the storage and secretion of defensive chemicals (Carbone et

al., 2013; Wägele et al., 2006), in combination with the evolution of aposematic signalling, may have
led to an adaptive radiation in nudibranchs (Gosliner, 2001).

Until recently, studies quantifying visual backgrounds in the context of defensive animal
colouration have relied on the independent consideration of spatial and spectral (colour and
luminance) properties (e.g. Cortesi & Cheney, 2010; Nokelainen et al., 2021). In this study, we
considered these pattern components in combination (van den Berg, Troscianko, et al., 2020) using
calibrated digital photography, the multispectral image calibration and analysis toolbox (MICA)
(Troscianko & Stevens, 2015) and its' integrated Quantitative Colour Pattern Analysis (QCPA)
framework (van den Berg, Troscianko, et al., 2020). This enabled us to assess the spatiochromatic
information of natural backgrounds upon which nudibranch individuals (n = 226) were found. QCPA
provides a descriptive array of image statistics capturing the spatiochromatic properties of each
background. We analysed backgrounds using the intraspecific interquartile range (IQR) of colour
pattern parameters, including: 1) colour adjacency analysis (CAA), which describes pattern geometry;
2) visual contrast analysis (VCA), which describes pattern boldness based on pattern element
properties; and 3) boundary strength analysis (BSA), which describes the contrast of boundaries
between pattern elements. This was performed according to the physiological limitations of an
ecologically relevant observer, a triggerfish (*Rhinecanthus aculeatus*). We analysed different viewing
distances to model an escalating predation sequence to account for changes in background
appearance as a predator approaches.

We then compared these colour pattern parameters to measures of the strength of chemical
defence for each species, which ranged from highly unpalatable to palatable using previously
published data from antifeedant assays with rockpool shrimp (*Palaemon serenus*) (Winters et al.,
2022). We hypothesised that highly defended (unpalatable) species would be found on more variable
visual backgrounds, and weakly defended species (palatable) would be found on backgrounds that
were less variable. Furthermore, we predicted that we would find a stronger correlation between the
level of defence and habitat variability at distances where prey detection and identification are most
likely (Endler, 1986, 1991; Ruxton et al., 2018), rather than their subjugation or consumption.

Materials and Methods

Study species. We took calibrated digital photographs of 226 Dorid nudibranchs belonging to 13
species (8-27 individuals per species, median = 16) (Supplementary Information, Fig. S1) from four
locations on the east coast of Australia: Sunshine Coast (SE Queensland, QLD), Gold Coast (SE QLD),

Cook Island (New South Wales, NSW) and Nelson Bay (NSW) between March 2016 and February 2021.

Species were identified visually using various nudibranch ID guides (Coleman et al., 2015; Debelius &

Kuiter, 2007; Gosliner et al., 2018): *Phyllidia varicosa* (n = 8); *Phyllidiella pustulosa* (n = 18); *Phyllidia*

elegans (n = 8); *Phyllidia ocellata* (n = 23); *Goniobranchus collingwoodi* (n = 15); *Goniobranchus*

splendidus (n = 25); *Glossodoris vespa* (n = 15); *Doriprismatica atromarginata* (n = 27), *Hypselodoris*

bennetti (n = 10); *Chromodoris kuiteri* (n = 17); *Chromodoris elisabethina* (N = 21); *Discodoris sp.* (n =

15); *Aphelodoris varia* (n = 24). *Discodoris sp*. in our study may have constituted a mixture of closely

related species, including *Sebadoris fragilis*, *Jorunna pantheris, Tayuva lilacina* and undescribed

species; however, these species are visually indiscriminable. Therefore, they are named *Discodoris sp.*

as they were found in the same locations, have no known chemical defences and are predominantly

nocturnal (Larkin et al., 2018). Only three out of the 13 species (*Doriprismatica atromarginata,*

Goniobranchus splendidus, Hypselodoris obscura) were found in comparable numbers across sites in

NSW and SE QLD, with most species found either in NSW or SE QLD (see Supplementary Information,

Table S1). Nudibranchs were photographed underwater against their natural habitat using a calibrated

Olympus EPL-5 with a 60mm macro lens in an Olympus PT-EP10 underwater housing using white LED

illumination from a combination of VK6r and PV62 Scubalamp video lights (van den Berg, Troscianko,

et al., 2020; see Supplementary Information Fig. S2 for spectral sensitivities of camera, spectral output

of LED lights, and irradiance spectrum of light environment at 5 m). All pictures were taken at roughly

a 90-degree angle relative to each animal and its background. Examples of images are provided in Fig.

1.

Figure 1. Representative nudibranch and background images for 4 out of the 13 species in this study. From top to bottom: *Discodoris sp., Phyllidia ocellata, Goniobranchus splendidus, Doriprismatica atromarginata.*

Image analysis. Using ImageJ (Schneider et al., 2012) and the MICA toolbox (Troscianko & Stevens, 2015), the images were manually segmented into regions of interest (ROI), selecting the animal from its background and defining a size standard using a custom-made colour and grey standard placed in each image (Fig. 1). All images were aligned, so the nudibranch was aligned vertically before analysis using QCPA (van den Berg, Troscianko, et al., 2020). To analyse the images, we used the visual system parameters of a trichromatic triggerfish, *Rhinecanthus aculeatus* (with spectral sensitivities of 413 nm, 480 nm and 528 nm λ_{max}; Cheney et al. 2022). This species (adult total length, TL ~ 15 cm) is a common shallow reef inhabitant found throughout the Indo-Pacific and feeds on invertebrates, algae, and detritus (Randall et al., 1997) and is deemed a potential predator of nudibranchs. We analysed for viewing distances of 2 cm, 5 cm, 10 cm, 30 cm and 50 cm assuming a triggerfish spatial acuity of 3 cycles per degree (Champ et al., 2014). The viewing distances reflect a range covering the maximal distance at which the largest specimen and coarse background detail would be detectable to the closest possible distance reflecting ultimate contact between predator and prey or visual background.

Following acuity modelling, the images were processed with a Receptor Noise Limited (RNL) ranked

filter (falloff: 3, radius: 5, repetition: 5) and clustered using RNL clustering with a colour discrimination

threshold of 2 ΔS (Cheney et al., 2019) and a luminance contrast threshold of 4 ΔS (van den Berg,

Hollenkamp, et al., 2020). We used Weber fractions based on a relative photoreceptor abundance of

1:2:2:2 (sw:mw:lw:dbl) and photoreceptor noise of 0.05, resulting in 0.07:0.05:0.05:0.05.

QCPA analysis was achieved using a custom batch script (currently available on request). This

resulted in a highly descriptive array of 157 colour pattern statistics for each animal, named with their

respective prefixes. Wherever pattern statistics could not be calculated, Infinite or NA values were

treated as 0. All pattern analyses used a segmented image, and measured transitions between pixels

along horizontal and vertical sampling transects in a transition matrix (as per (Endler, 2012). Statistics

ending with 'hrz' or 'vrt' are the horizontal (across body axis) and vertical (along body axis) versions

(analysing the respective transition matrix only) of their respective statistic (analysing the full

transition matrix). For example, a background with algae or seagrass will likely have an elongated

aspect ratio, whereas a uniform sandy background will not. A detailed description of each pattern

statistic is found in van den Berg, Troscianko, et al. (2020).

Level of chemical defence. As a measure of chemical defence for each species, we used our previously

published data from antifeedant assays with rockpool shrimp (*Palaemon serenus*) (Winters et al.,

2022). Assay data for *Glossodoris vespa* is presented in Winters *et al.* (2018). In summary, the data

were obtained from extracting secondary metabolites from nudibranchs and added to food pellets

made from squid mantle at increasing concentrations. Effective dose (ED_{50}) values were calculated

based on the concentration that elicited a rejection response in at least 50% of the shrimp. For this

study, the resulting ED_{50} values were normalised and calculated as 1-ED_{50} to range from 0 (most

palatable) to 1 (least palatable). Where multiple estimates of ED_{50} existed for a species due to multiple

extracts/assays being performed, the average value was used.

Statistical analysis. All statistical analyses were conducted using R Software (R Core Team, 2021)

(v4.1.2). To quantify background variability, we calculated the interquartile range (IQR) for each

pattern statistic of each species. All statistics were then normalised to range from 0 (least variable) to

1 (most variable), with all trait data subjected to an ordered quantile normalisation (Peterson &

Cavanaugh, 2020) using the bestNormalize package (v1.8.2). We first assessed whether there were

colour pattern differences in background between populations for those species that were found

across all four locations (*G. splendidus and D. atromarginata*); however, we found no significant

differences between Queensland and New South Wales locations in the mean IQR across all 157

parameters, using species and viewing distance as a random factor (lmer, $F_{1,17}$ = 1.56, p = 0.22). To

investigate whether there was a phylogenetic correlation between our pattern statistics and
palatability, we pruned the phylogeny from Cheney et al. (2014) using the 'ape' package (Paradis &
Schliep, 2019) (v5.6-1) with *Aphelodoris varia* added manually according to its taxonomic classification
in the World Register of Marine Species (WoRMS) (Appeltans et al., 2012). However, the small number
of species made it impossible to estimate the phylogenetic signal reliably using phylogenetic least
square regression (pgls). As a result, we estimated a phylogenetic signal close to 0 across all viewing
distances when analysing the median IQR across all 157 pattern statistics. This finding is in line with
previous studies (Cheney et al., 2014; Cortesi & Cheney, 2010), which have determined a very small
phylogenetic signal across various image statistics and Dorid nudibranch species, meaning that closely
related species can look considerably different. Therefore, we analysed the data assuming a negligible
phylogenetic signal using a Pearson correlation test in the 'caret' package (Kuhn, 2021) (v6.0-90). As
all pattern statistics are defined by the same three core properties (colour, luminance and pattern),
we applied a conservative Bonferroni correction of n = 3 to avoid an excessive type II error (Moran,
2003), resulting in an adjusted significance threshold of p = 0.017.

Results

The intraspecific Interquartile-Range (IQR) of many visual background colour pattern statistics
was negatively correlated with the level of unpalatability at each viewing distance (Table 1), indicating
that the visual backgrounds of unpalatable (defended) species were less variable than the background
of palatable (undefended) species. We did not find any significant positive correlations among all 785
comparisons of colour pattern statistics with chemical defences, nor did we find any moderately
strong positive correlations (r > 0.4, p = < 0.18). Below, we report both significant values of negative
correlations (r < -0.65, p < 0.017), but also non-significant moderate correlations (r < -0.4 or r > 0.4, p
= < 0.18). To reflect the behavioural context of predator-prey interactions, we present the results from
the largest viewing distance to the smallest to replicate an escalating predation sequence (Endler,
1991).

At 50 cm, only coarse background details could be resolved by the visual system of our
triggerfish predator. We found no significant negative correlations passing the Bonferroni correction
(Table 1, Table S2); however, there were moderate negative correlations for 25 pattern statistics with
unpalatability (Pearson correlation, r < -0.4, p = < 0.18). We did not find any significant positive
correlations, and there were no moderate positive correlations (all r < 0.34, p > 0.29) (Table S2).

At 30 cm, four out of 157 pattern statistics were significantly negatively correlated with
unpalatability (Pearson correlation: $r < -0.5$, $p < 0.017$; Table 1). All significantly correlated pattern
statistics at 30 cm indicated more unpalatable species to be found on backgrounds with lower
variability in their geometric complexity along the horizontal axis relative to an animal (CAA.C.Hrz) as
well as reduced variability in the coefficient of variation of the luminance contrast (ΔS) relative to the
patch sizes in each background (VCA.CVSL) (Table 1). There were moderate negative correlations for
47 pattern statistics with unpalatability (Pearson correlation, $r < -0.4$, $p = < 0.18$). We did not find any
significant positive correlations, and there were no moderate positive correlations (all $r < 0.18$, $p >$
0.55) (Table S2).

At 10 cm, six out of 157 pattern statistics were significantly negatively correlated with
unpalatability (Pearson correlation $r < -0.65$, $p < 0.017$; Table 1, Table S2). More unpalatable species
were found on less variable visual backgrounds with lower variability of the coefficient of variation of
chromaticity contrast between colour pattern elements (VCA.CVS) as well as backgrounds less variable
concerning the regularity of the transitions between colour pattern elements overall (CAA.Jt) and in
the vertical axis of a background relative to the anterior-posterior axis of an animal (CAA.Jt.Vrt). More
unpalatable animals were also found on backgrounds with reduced variability in their Shannon colour
diversity (CAA.Qc) (Table 1). There were moderate negative correlations for 47 pattern statistics with
unpalatability (Pearson correlation, $r < -0.4$, $p = < 0.18$). We did not find any significant positive
correlations, and there were no moderate positive correlations (all $r < 0.26$, $p > 0.40$) (Table S2).

At 5 cm, three out of 157 pattern statistics were significantly negatively correlated with
unpalatability ($r < -0.65$; $p < 0.017$; Table 1, Table S2). The strongest correlations were observed at
this distance, with the coefficient of variation luminance contrast between colour pattern elements of
a background (VCA.CVSL) being negatively correlated with unpalatability (Pearson correlation: $r = -$
0.75, $p = 0.003$; Fig. 2). In other words, for VCA.CVSL, highly defended species were viewed against
backgrounds that were 56.99% less variable than non-defended species. There were moderate
negative correlations for 26 pattern statistics with unpalatability (Pearson correlation, $r < -0.4$, $p = <$
0.18). We did not find any significant positive correlations, and there were no moderate positive
correlations (all $r < 0.24$, $p > 0.44$) (Table S2).

At 2 cm, no significant correlations passing the Bonferroni correction could be found (Table 1,
Table S2). There were moderate negative correlations for 23 pattern statistics with unpalatability
(Pearson correlation, $r < -0.4$, $p = < 0.18$). We did not find any significant positive correlations, and
there were no moderate positive correlations (all $r < 0.39$, $p > 0.19$) (Table S2).

Although we found individual pattern parameters were negatively correlated with
unpalatability, we did not find that the mean IQR of all 157 pattern parameters for each species was
significantly correlated with unpalatability at any viewing distance (50 cm: Pearson correlation $r = -0.24$, $p = 0.436$; 30 cm: Pearson correlation $r = -0.536$, $p = 0.059$; 10 cm: Pearson correlation $r = -0.51$, $p = 0.078$; 5 cm: Pearson correlation $r = -0.09$, $p = 0.780$; 2 cm: Pearson correlation $r = -0.33$, $p = 0.273$).

Table 1. All significantly correlated pattern statistics for each viewing distance with a Bonferroni adjusted Pearson correlation significance of *p* < 0.017. Statistics are ranked according to their significance at each viewing distance. Pearson correlations (r) are also expressed as a percentage.

Viewing distance	Stage in Predation Sequence	Pattern statistic	Pearson Correlation [r] / % [r²*100]	P value
50cm	Encounter / Detection	None	None	None
30cm	Detection / Identification	CAA.C.Hrz	-0.68 / 46.11	0.011
		VCA.CVSL	-0.66 / 43.88	0.014
		VCA.CVSL.Hrz	-0.66 / 43.88	0.014
		VCA.CVSL.Vrt	-0.66 / 43.88	0.014
10cm	Identification / Approach	VCA.CVS	-0.69 / 48.00	0.009
		VCA.CVS.Hrz	-0.69 / 48.00	0.009
		VCA.CVS.Vrt	-0.69 / 48.00	0.009
		CAA.Jt.Vrt	-0.68 / 46.58	0.010
		CAA.Qc	-0.67 / 45.74	0.011
		CAA.Jt	-0.66 / 42.90	0.015
5cm	Approach / Subjugation	VCA.CVSL	-0.75 / 56.99	0.003
		VCA.CVSL.Hrz	-0.75 / 56.99	0.003
		VCA.CVSL.Vrt	-0.75 / 56.99	0.003
2cm	Subjugation / Consumption	None	None	None

Figure 2. An example colour pattern statistic of the background (IQR of luminance contrast; VCA.CVSL) plotted

against the normalised unpalatability score of each species, at a 5cm viewing distance, (Pearson correlation: r =

-0.75, p = 0.003)

Discussion

We found several significant and moderately negative correlations between the strength of chemical defences and the variability of individual background statistics at any viewing distance, indicating that specific image statistics of the visual backgrounds inhabited by 13 Dorid nudibranch species are up to 56.99% less variable in species with higher levels of chemical defences (unpalatability). Importantly, we found no significant or moderately positive correlations at any viewing distance. This indicates that, contrary to the predictions derived from the 'escape and radiate' hypothesis (Merilaita & Tullberg, 2005; Thompson, 1989), well-defended nudibranch species utilised a smaller range of background habitats, and were less likely to be background generalists than cryptic species. Our study provides the first spatiochromatic quantification of natural background variability for aposematic species beyond the previously separated modalities of colour and spatial frequency. These findings are among the strongest quantitative correlations between primary and secondary defences known to date (White & Umbers, 2021).

Background variability did not vary consistently across all 157 colour pattern statistics and at all distances. This indicates that a distinct set of visual features (rather than all possible colour pattern characteristics) may be ecologically relevant at a given viewing distance. Our data also shows that some of these statistics are relevant for both along and perpendicular to the body long axis of the animal found on each background, whereas others do not (Table 1). This is interesting regarding the potential importance of feature selectivity in the visual perception of prey and their environment by potential predators (Howe & Purves, 2005; Hubel & Wiesel, 1962). Specifically, our results show that the variability of visual contrast (VCA) of colour saturation (VCA.CVS) and luminance (VCA.SL), in addition to the regularity of the background patterning (CAA.Qc & CAA.Jt) was reduced in species with strong chemical defences. Colour, luminance and pattern contrast are crucial for the detectability of aposematic species (Mappes et al., 2005) and a reduction in the variability of these properties in visual backgrounds is of likely consequence to the signalling ability of aposematic species.

As predicted, the strongest and most correlations between background variability and chemical defences occurred at distances corresponding to prey detection, identification, and approach, with correlations occurring at distances between 5 and 30 cm. From our data, prey detection at distances beyond 30 cm may only be relevant for a few very large species of nudibranchs. In contrast, aspects of object identity and quality are mediated by the intrinsic properties of the animal itself and processed by the predator post-detection and likely independent of distractor features present in the visual background (excluding examples of masquerade) (Endler, 1991; Guilford & Dawkins, 1993; Ruxton et al., 2018). Indeed, the lack of significant correlations at very close distances

(2 cm) might suggest that visual backgrounds are more important in object detection rather than
discrimination.

However, our modelling choice of 3 cycles per degree (cpd) is likely conservative (Cheney et
al., 2022), and feature detection in the visual system of triggerfish could happen at multiple spatial
scales simultaneously (Hubel & Wiesel, 1962; van den Berg et al., 2022). Furthermore, nudibranch
species are highly variable in size, and spatial acuity within a predator community is also likely variable.
Therefore, if and how specific properties of colour patterns in Dorid nudibranchs correlate with
predator cognition will need to be validated with behavioural studies.

Nudibranchs mainly use chemotaxis to move around their environment to find food and
mates, with visual input only relevant for basic phototaxis, such as determining daytime or detecting
shelter (Eakin et al., 1967). Therefore, unlike many other aposematic species, nudibranchs will unlikely
choose resting and foraging microhabitats based on visual cues. Furthermore, to our knowledge, no
study has quantified the mobility of different nudibranch species, and the diets of many nudibranch
species (however, see Rudman & Bergquist, 2007), and the spatiotemporal distribution of food
sources remain poorly known. Some of the most unpalatable nudibranchs in this study are known to
forage on sponges containing highly potent secondary metabolites (e.g. latrunculin a) at least at some
stage during their lifetime (e.g. *Chromodoris elisabethina*) (Cheney et al., 2016). In contrast, others
can synthesise compounds *de novo* (Cimino & Ghiselin, 1999). Nudibranchs with more specialist diets
may need to increase mobility to find uncommon food sources. These then allow the animals to
maintain functional levels of secondary metabolites or pigments required to maintain salient
aposematic colour patterns. Background specialisation might therefore be increased in species
dependent on such supplemental food sources. Specifically, colour patterns in aposematic species
may be designed to be particularly efficient on certain backgrounds but not others. This might help
explain our results while aligning with the abundance of chemically defended but likely cryptic Dorid
nudibranch species (Cortesi & Cheney, 2010). Intriguingly, the same mechanism could potentially also
lead to reduction in observed background variability. However, our study does not consider species
relying on masquerade (Ruxton et al., 2018), which is commonly seen in extreme background
specialists (living on one species of prey exclusively) where nudibranchs mimic their food source, such
as various species of the genus *Phyllodesmium* (e.g. *Phyllodesmium lizardensis*).

Our study also only considered the degree of unpalatability rather than toxicity (or both) to
quantify the level of the chemical defence (Winters et al., 2022). These two mechanisms of defence
are not always related (Mappes et al., 2005), but predators are more likely to learn to avoid
aposematic prey that is initially aversive through taste, rather than toxicity (although we are not aware

of conclusive empirical evidence for this). Future studies investigating correlations between
background variability and additional modalities of prey profitability (such as toxicity or nutritional
value) would be of great interest.

When discussing correlations between visual properties of backgrounds and the evolution of
aposematic colouration, Merilaita & Tullberg (2005) specifically focused on diurnal lepidopterans as
day-active and nocturnal species are expected to be under very different sets of constraints from
visual predation than nudibranchs. Indeed, our study uses species found during both day and (to a
much lesser extent) night (Supplementary Information, Table S1); further studies are needed to
investigate the correlation between daytime activity and habitat use in nudibranchs (Larkin et al.,
2018).

Finally, it is possible that, for some colour pattern statistics, there could be a strong
phylogenetic signal, which we have not been able to pick up in this study as we only sampled 13 out
of many co-occurring sea slug species at either sampling site (e.g. Schubert & Smith, 2020). To test
this, we would need to increase our sample sizes and more systematic sampling across geographical
and seasonal ranges. This would allow applying sophisticated comparative phylogenetic analyses
necessary for properly investigating coevolutionary processes in Dorid nudibranchs. However,
obtaining calibrated imaging data suitable for visual modelling is tedious and resource intensive. As a
result, our study is the first to analyse spatiochromatic signalling properties of aposematic species
across more than a handful of species.

In conclusion, while not confirming the escape and radiate hypothesis, our findings are not at
odds with existing literature. Furthermore, the mechanisms underlying the 'escape and radiate'
hypothesis are often unclear, with little empirical evidence of the process (Suchan & Alvarez, 2015).
Nevertheless, our study provides the most thorough investigation of correlations between chemical
defences and visual background diversity in aposematic animals to date.

Acknowledgements

We thank Dr. Nicholas Condon for his assistance with the creation of automated QCPA scripts for
analysing the image, several volunteers for assistance with fieldwork and image analysis. We also
thank the staff running the High-Performance Computing infrastructure at The University of
Queensland (Wiener & Awoonga) and Dr. Simone Blomberg, who provided infrastructure that
contributed to the computing of image statistics.

Conflict of Interest

The authors have no conflict of interest to declare.

Data availability

The data is available on UQ's digital repository eSpace: https://doi.org/10.48610/8f7b70e

Author contributions

CPvdB conceived the original study with KLC & JAE providing critical input; CPvdB collected and

analysed the data; CPvdB and KLC led the writing of the manuscript. All authors contributed critically

to the drafts and gave final approval for publication.

Funding

This work was funded by the Australian Research Council (FT190199313 and DP180102363 awarded

to KLC and JAE), two Holsworth Wildlife Research Endowment grants awarded to CPvdB, and a

student research grant from the Australasian Society for the Study of Animal Behaviour awarded to

CPvdB.

References

Appeltans, W., Bouchet, P., Boxshall, G. A., De Broyer, C., de Voogd, N. J., Gordon, D. P., Hoeksema,

B. W., Horton, T., Kennedy, M., Mees, J., Poore, G. C. B., Read, G., Stöhr, S., Walter, T. C., &

Costello, M. J. (eds). (2012). *World Register of Marine Species*. http://www.marinespecies.org

Arbuckle, K., & Harris, R. J. (2021). Radiating pain: venom has contributed to the diversification of

the largest radiations of vertebrate and invertebrate animals. *BMC Ecology and Evolution*,

21(1), 1–12. https://doi.org/10.1186/s12862-021-01880-z

Arbuckle, K., & Speed, M. P. (2015). Antipredator defenses predict diversification rates. *Proceedings*

of the National Academy of Sciences of the United States of America, *112*(44), 13597–13602.

https://doi.org/10.1073/pnas.1509811112

Avila, C. (1995). Natural products of opisthobranch molluscs: a biological review. *Oceanography and*

Marine Biology: An Annual Review, *33*, 487–559.

Carbone, M., Gavagnin, M., Haber, M., Guo, Y.-W., Fontana, A., Manzo, E., Genta-Jouve, G.,

Tsoukatou, M., Rudman, W. B., Cimino, G., Ghiselin, M. T., & Mollo, E. (2013). Packaging and

Delivery of Chemical Weapons: A Defensive Trojan Horse Stratagem in Chromodorid

Nudibranchs. *PLoS ONE*, *8*(4), e62075. https://doi.org/10.1371/journal.pone.0062075

Champ, C. M., Wallis, G., Vorobyev, M., Siebeck, U., & Marshall, J. (2014). Visual acuity in a species of coral reef fish: *Rhinecanthus aculeatus*. *Brain, Behavior and Evolution*, *83*(1), 31–42. https://doi.org/10.1159/000356977

Cheney, K. L., Cortesi, F., How, M. J., Wilson, N. G., Blomberg, S. P., Winters, A. E., Umanzör, S., & Marshall, N. J. (2014). Conspicuous visual signals do not coevolve with increased body size in marine sea slugs. *Journal of Evolutionary Biology*, *27*(4), 676–687. https://doi.org/10.1111/jeb.12348

Cheney, K. L., Green, N. F., Vibert, A. P., Vorobyev, M., Marshall, N. J., Osorio, D. C., & Endler, J. A. (2019). An Ishihara-style test of animal colour vision. *Journal of Experimental Biology*, *222*(1), jeb189787. https://doi.org/10.1242/jeb.189787

Cheney, K. L., Hudson, J., de Busserolles, F., Luehrmann, M., Shaughnessy, A., van den Berg, C., Green, N. F., Marshall, N. J., & Cortesi, F. (2022). Seeing Picasso: an investigation into the visual system of the triggerfish *Rhinecanthus aculeatus*. *Journal of Experimental Biology*, *225*(7), jeb243907. https://doi.org/10.1242/jeb.243907

Cheney, K. L., White, A., Mudianta, I. W., Winters, A. E., Quezada, M., Capon, R. J., Mollo, E., & Garson, M. J. (2016). Choose your weaponry: Selective storage of a single toxic compound, latrunculin a, by closely related nudibranch molluscs. *PLoS ONE*, *11*(1), 1–16. https://doi.org/10.1371/journal.pone.0145134

Cimino, G., & Ghiselin, M. T. (1999). Chemical defense and evolutionary trends in biosynthetic capacity among dorid nudibranchs (Mollusca: Gastropoda: Opisthobranchia). *Chemoecology*, *9*(4), 187–207. https://doi.org/10.1007/s000490050052

Coleman, N., Cobb, G., & Mullins, D. (2015). *Nudibranchs Encyclopedia*. Thomas Slone, Masalai Press and Tim Hochgrebe.

Cortesi, F., & Cheney, K. L. (2010). Conspicuousness is correlated with toxicity in marine opisthobranchs. *Journal of Evolutionary Biology*, *23*(7), 1509–1518. https://doi.org/10.1111/j.1420-9101.2010.02018.x

Debelius, H., & Kuiter, R. H. (2007). *Nudibranchs of the world*. IKAN-Unterwasserarchiv.

Eakin, R. M., Westfall, J. A., & Dennis, M. J. (1967). Fine Structure of the Eye of a Nudibranch Mollusc, *Hermissenda Crassicornis *. *Journal of Cell Science*, *2*(3), 349–358. https://doi.org/10.1242/jcs.2.3.349

Ehrlich, P. R., & Raven, P. H. (1964). Butterflies and Plants: A Study in Coevolution. *Evolution*, *18*(4),
586. https://doi.org/10.2307/2406212

Endler, J. A. (1984). Progressive background in moths, and a quantitative measure of crypsis.
Biological Journal of the Linnean Society, *22*(3), 187–231. https://doi.org/10.1111/j.1095-
8312.1984.tb01677.x

Endler, J. A. (1986). Defense Against Predators. In *Predator-prey relationships, Perspectives and*
Approaches from the Study of Lower Vertebrates (pp. 109–134). University of Chicago Press.

Endler, J. A. (1991). Interactions between predators and prey. In J. R. Krebs & N. B. Davies (Eds.),
Behavioural Ecology (pp. 169–196). Blackwell Scientific.

Endler, J. A. (2012). A framework for analysing colour pattern geometry: Adjacent colours. *Biological*
Journal of the Linnean Society, *107*(2), 233–253. https://doi.org/10.1111/j.1095-
8312.2012.01937.x

Gosliner, T. M. (2001). Aposematic coloration and mimicry in opisthobranch mollusks: new
phylogenetic and experimental data. *Bollettino Malacologico*, *37*(37), 163–170.

Gosliner, T. M., Valdes, A., & Behrens, D. W. (2018). *Nudibranch & Sea Slug Identification - Indo-*
Pacific (2nd ed.). New World Publications.

Guilford, T., & Dawkins, M. S. (1993). Receiver psychology and the design of animal signals. *Trends in*
Neurosciences, *16*(August 1990), 430–436. https://doi.org/10.1016/S0003-3472(05)80600-1

Howe, C. Q., & Purves, D. (2005). Natural-scene geometry predicts the perception of angles and line
orientation. *Proceedings of the National Academy of Sciences of the United States of America*,
102(4), 1228–1233. https://doi.org/10.1073/pnas.0409311102

Hubel, D. H., & Wiesel, T. N. (1962). Receptive fields, binocular interaction and functional
architecture in the cat's visual cortex. *The Journal of Physiology*, *160*(1), 106–154.
https://doi.org/10.1113/jphysiol.1962.sp006837

Kuhn, M. (2021). *caret: Classification and Regression Training. R package version 6.0-90.*
https://CRAN.R-project.org/package=caret (6.0-88). https://cran.r-project.org/package=caret

Larkin, M. F., Smith, S. D. A., Willan, R. C., & Davis, T. R. (2018). Diel and seasonal variation in
heterobranch sea slug assemblages within an embayment in temperate eastern Australia.
Marine Biodiversity, *48*(3), 1541–1550. https://doi.org/10.1007/s12526-017-0700-9

Mappes, J., Marples, N., & Endler, J. A. (2005). The complex business of survival by aposematism.

Trends in Ecology and Evolution, *20*(11), 598–603. https://doi.org/10.1016/j.tree.2005.07.011

Merilaita, S., & Tullberg, B. S. (2005). Constrained camouflage facilitates the evolution of

conspicuous warning coloration. *Evolution*, *59*(1), 38–45. https://doi.org/10.1111/j.0014-

3820.2005.tb00892.x

Merilaita, S., Tuomi, J., & Jormalainen, V. (1999). Optimization of cryptic coloration in heterogenous

habitats. *Biological Journal of the Linnean Society*, *67*(2), 151–161.

https://doi.org/10.1006/bijl.1998.0298

Moran, M. D. (2003). Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos*,

100(2), 403–405. https://doi.org/10.1034/j.1600-0706.2003.12010.x

Nokelainen, O., Scott-Samuel, N. E., Nie, Y., Wei, F., & Caro, T. (2021). The giant panda is cryptic.

Scientific Reports, *11*(1), 1–10. https://doi.org/10.1038/s41598-021-00742-4

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics*, *35*(3), 526–528. https://doi.org/10.1093/bioinformatics/bty633

Peterson, R. A., & Cavanaugh, J. E. (2020). Ordered quantile normalization: a semiparametric

transformation built for the cross-validation era. *Journal of Applied Statistics*, *47*(13–15), 2312–

2327. https://doi.org/10.1080/02664763.2019.1630372

Poulton, E. B. (1890). *The Colours of Animals*. Kegan Paul, Trench & Trubner.

R Core Team. (2021). *R: A Language and Environment for Statistical Computing* (4.1.). R Foundation

for Statistical Computing. http://www.r-project.org/

Randall, J. E., Allen, G. R., & Steene, R. C. (1997). *Fishes of the Great Barrier Reef and Coral Sea*.

Crawford House Publishing.

Regan, D., & Beverley, K. I. (1984). Figure–ground segregation by motion contrast and by luminance

contrast. *Journal of the Optical Society of America A*, *1*(5), 433.

https://doi.org/10.1364/josaa.1.000433

Rudman, W. B., & Bergquist, P. R. (2007). A review of feeding specificity in the sponge-feeding

Chromodorididae (Nudibranchia: Mollusca). *Molluscan Research*, *27*(2), 60–88.

Ruxton, G. D., Allen, W. L., Sherratt, T. N., & Speed, M. P. (2018). *Avoiding Attack* (Vol. 1). Oxford

University Press. https://doi.org/10.1093/oso/9780199688678.001.0001

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image

analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Schubert, J., & Smith, S. D. A. (2020). Sea slugs—"rare in space and time"—but not always. *Diversity*,
12(11), 1–14. https://doi.org/10.3390/d12110423

Speed, M. P., Brockhurst, M. A., & Ruxton, G. D. (2010). The dual benefits of aposematism: Predator
avoidance and enhanced resource collection. *Evolution*, *64*(6), 1622–1633.
https://doi.org/10.1111/j.1558-5646.2009.00931.x

Suchan, T., & Alvarez, N. (2015). Fifty years after Ehrlich and Raven, is there support for plant-insect
coevolution as a major driver of species diversification? *Entomologia Experimentalis et*
Applicata, *157*(1), 98–112. https://doi.org/10.1111/eea.12348

Thompson, J. N. (1989). Concepts of coevolution. *Trends in Ecology and Evolution*, *4*(6), 179–183.
https://doi.org/10.1016/0169-5347(89)90125-0

Troscianko, J., & Stevens, M. (2015). Image calibration and analysis toolbox – a free software suite
for objectively measuring reflectance, colour and pattern. *Methods in Ecology and Evolution*,
6(11), 1320–1331. https://doi.org/10.1111/2041-210X.12439

van den Berg, C. P., Endler, J. A., Papinczak, D. E. J., & Cheney, K. L. (2022). Using colour pattern
edge contrast statistics to predict detection speed and success in triggerfish (Rhinecanthus
aculeatus) . *Journal of Experimental Biology*. https://doi.org/10.1242/jeb.244677

van den Berg, C. P., Hollenkamp, M., Mitchell, L. J., Watson, E. J., Green, N. F., Marshall, N. J., &
Cheney, K. L. (2020). More than noise: Context-dependant luminance contrast discrimination in
a coral reef fish (Rhinecanthus aculeatus). *Journal of Experimental Biology*, *223*(21),
jeb.232090. https://doi.org/10.1242/jeb.232090

van den Berg, C. P., Troscianko, J., Endler, J. A., Marshall, N. J., & Cheney, K. L. (2020). Quantitative
Colour Pattern Analysis (QCPA): A comprehensive framework for the analysis of colour patterns
in nature. *Methods in Ecology and Evolution*, *11*(2), 316–332. https://doi.org/10.1111/2041-
210X.13328

Wägele, H., Ballesteros, M., & Avila, C. (2006). Defensive Glandular Structures In Opisthobranch
Molluscs — From Histology To Ecology. In *Oceanography and Marine Biology* (Vol. 44, Issue
June, pp. 197–276). https://doi.org/10.1201/9781420006391.ch5

White, T. E., & Umbers, K. D. L. (2021). Meta-analytic evidence for quantitative honesty in
aposematic signals. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1949).
https://doi.org/10.1098/rspb.2021.0679

Winters, A. E., Chan, W., White, A. M., Berg, C. P., Garson, M. J., & Cheney, K. L. (2022). Weapons or

deterrents? Nudibranch molluscs use distinct ecological modes of chemical defence against

predators. *Journal of Animal Ecology*, *91*(4), 831–844. https://doi.org/10.1111/1365-

2656.13643

Winters, A. E., White, A. M., Dewi, A. S., Mudianta, I. W., Wilson, N. G., Forster, L. C., Garson, M. J., &

Cheney, K. L. (2018). Distribution of Defensive Metabolites in Nudibranch Molluscs. *Journal of*

Chemical Ecology, *44*(4), 384–396. https://doi.org/10.1007/s10886-018-0941-5

498